







E. Rogers
1835.

Presented by J. E. Placerville

Received of
1891
J. J. [unclear]
[unclear]

PHARMACOPŒIA UNIVERSALIS.

ELIASHVILI UNIVERSITY

THE
PHARMACOPŒIA UNIVERSALIS ;
OR, COMPLETE
ENCYCLOPÆDIA
OF THE
MATERIA MEDICA,

CONTAINED IN

THE PHARMACOPŒIAS OF LONDON, EDINBURGH, AND DUBLIN, AS
WELL AS OF ALL THOSE OF EUROPE AND AMERICA ;

AND OF

THE DISPENSATORIES, FORMULARIES, AND CHEMICAL WORKS

OF

AINSLIE, BIGELOW, BRANDE, BRERA, BRUGNATELLI, BUCHNER, CHEVALIER,
CHEVREUIL, COKE, CULLEN, DAVY, DE LENS, DUNCAN, GRAY, GUIBOUT,
HENRY, HUFELAND, MAGENDIE, MURRAY, ORFILA, PARIS, PHILIPS,
PIDERIT, POIRET, RATIER, RICHARD, ROBINET, ROSE, SPIELMANN,
THOMSON, VAN MONS, AND NUMEROUS OTHERS :

GIVING

THE OFFICINAL AND OTHER SYNONIMES AND SCIENTIFIC CHARACTERS OF
SIMPLE SUBSTANCES, THE PREPARATIONS OF THE COMPOUNDS, ACCORDING
TO THE VARIOUS FORMULÆ, AND THE USUAL DOSES PRESCRIBED.

By A. J. L. JOURDAN, M. D. OF PARIS.

EDITED

By J. RENNIE, A. M., A. L. S.

VOL. I.

LONDON

PUBLISHED BY JOSEPH ROBINS

BRIDE COURT, FLEET STREET.

MDCCCXXXIII.



PHARMACOPOEIA UNIVERSALIS.



THE
PHARMACOPŒIA UNIVERSALIS ;

OR, COMPLETE

ENCYCLOPÆDIA

OF THE

MATERIA MEDICA,

CONTAINED IN

THE PHARMACOPŒIAS OF LONDON, EDINBURGH, AND DUBLIN, AS
WELL AS OF ALL THOSE OF EUROPE AND AMERICA ;

AND OF

THE DISPENSATORIES, FORMULARIES, AND CHEMICAL WORKS

OF

AINSLIE, BIGELOW, BRANDE, BRERA, BRUGNATELLI, BUCHNER, CHEVALIER,
CHEVREUIL, COXE, CULLEN, DAVY, DE LENS, DUNCAN, GRAY, GUIBOURT,
HENRY, HUFELAND, MAGENDIE, MURRAY, ORFILA, PARIS, PHILIPS,
PIDERIT, POIRET, RATIER, RICHARD, ROBINET, ROSE, SPIELMANN,
THOMSON, VAN MONS, AND NUMEROUS OTHERS :

GIVING

THE OFFICINAL AND OTHER SYNONIMES AND SCIENTIFIC CHARACTERS OF
SIMPLE SUBSTANCES, THE PREPARATIONS OF THE COMPOUNDS, ACCORDING
TO THE VARIOUS FORMULÆ, AND THE USUAL DOSES PRESCRIBED.

By A. J. L. JOURDAN, M. D. OF PARIS.

EDITED

By J. RENNIE, A. M., A. L. S.

VOL. II.

LONDON :

PUBLISHED BY JOSEPH ROBINS,

BRIDE COURT, FLEET STREET.

MDCCCXXXIII.

THE PHARMACOPŒIA UNIVERSALIS.

HAMAMELIS VIRGINIANA, LINN.—WHITE HAZEL.

COXE.

A shrub of North America. (*Tetrandria Digynia*, LINN.; *Berberideæ*, JUSS.; *Fig.* Cat. Carol. III. tab. 2.)

The bark is employed. It is bitter and astringent, leaving a disagreeable impression on the tongue.

HEDERA HELIX, LINN.—IVY.

SYNONIMES:—Lierre grim pant, FR.; Ephed, GERM.; Brectan, BOHEM.; Vintergront, DAN.; Hiedra, SPAN.; Klyf, DUT.; Edera, ITAL.; Bluszez, POL.; Hera, PORT.; Margræn, SWED.

BRUNS.; HISP.; GALL.; FERR.; GENEV.; HAMB.; WIRTEM.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

Europe. (*Pentandria Monogynia*, LINN.; *Caprifoliaceæ*, JUSS.; *Fig.* Lab. Icon. 614.)

The leaves and fruit are employed.

The leaves (*Folia Hederæ arboreæ*) are glabrous, shining, of a deep green colour, oval or three or five-lobed, of a variable form.

The berries are not very succulent, of a blackish-green colour, having from three to four monospermal cells.

This shrub is found in the greater part of

HELIANTHUS TUBEROSUS, LINN.—JERUSALEM ARTICHOKE.

SYNONIMES:—Topinambour, FR.; Erdartischocke, Jerusalemsartischocke, GERM.; Jærdæble, DAN.; Jirasol tuberoso, SPAN.; Aardpeeren, DUT.; Girasol tuberoso, ITAL.; Girasolo Brasil, PORT.; Jærdæple, SWED.

This plant is originally from Brazil; it is cultivated in Europe. (*Syngenesia Polygamia Superflua*, LINN.; *Synanthereæ*, CASS.; *Fig.* Jacq. Hort. Vindob. II. t. 16.)

The root is employed; it is polymorphous, reddish without, fleshy white within, inodorous, and almost insipid.

GALL.; BERGIUS.

HELIOTROPIMUM EUROPÆUM, LINN.—TURNSOL.

SYNONIMES:—Heliotrope d'Europe, Tournesol, Herbe aux Verrues, FR.; Sonnenwende, GERM.; Hierba verruguerra,

SPAN.; Eliotropio maggiore, ITAL.; Tornesol, PORT.

GALL.

B

A plant of Europe. (*Pentandria Monogynia*, LINN.; *Boraginææ*, JUSS.; *Fig. Jacq. Fl. Austr. t. 207.*)

The herb (*Herba Verrucariæ seu Cancri*) is employed. It consists of branching stems, furnished with short hairs and with

oval, petiolated leaves, somewhat hairy, wrinkled, and of a whitish-green colour. It has a bitter taste.

Its juice was formerly esteemed for occasioning the separation of warts.

HELMINTHOCORTON.—CORSICAN MOSS.

SYNONIMES:—*Helminthocortos*, *Leimithocorton*, *Eleminthocorton*, *Melithocorton*, *Corallina Corsica seu rubra*, *Muscus Helminthocortos seu corallinus seu marinus* seu *Helminthocortus*.—Mousse de Corse, Mousse de mer, Coralline de Corse, Fr.

AMST.; ANT.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

This marine production is obtained chiefly from the coasts of Corsica. It has a disagreeable odour, and a saline and nauseous taste.

It is a mixture of more than twenty, as well marine plants as of what M. LAMOUROUX calls *polypiers flexibles*, as the *Fucus Helminthocorton*, LATOUR., *F. Cricoides*, GOOD., *F. barbatus*, GOOD., *T. Sodoïdes*, DESF., *T. fascicla*, RO., *Desmarestia aculeata*, LMX., *Gigartina plicata*, LMX., *Ceramium catenatum*, *agagrophilum albinum*, *incurvum*, *forcipatum*, *scoparium*, *gracile*, *cancellatum*, CAND., *Ulva lactuca*, *squamaria*, *pavonia*, CAND., *Corallina rubens*, *officinarum*, fragments of *Zostera marina*, *Sertularia*, *Gorgonia*, *Tubularia*, the *Acetabularia mediterranea*, LMX., sand, and shells. But, usually, in this informal mixture the *Gigartina Helminthocorton*, LMX. predominates, an alga (*Fig. Journ. de Phys. t. XX. tab. 1*) which we have seen to constitute from an eighth to a third of it. This plant is about two inches long, of a pale brownish, or reddish colour, forming close tufts, composed of many slender stems, having three or four horned and again straightened, almost simple branches, with pointed extremities, sensibly articulated, having hemispherical, lateral, loose, and sessile fructiferous tubercles.

It is slightly stimulant, and is reputed to be a powerful anthelmintic, particularly for children.—Dose, from ten grains to two drams, in honey.

VERMIFUGE POWDER. (GALL.; FORM. HÔP. FR.; CADET DE GASSICOURT)

R. Corsican Moss,	} of each, an equal part.
Seeds of Artemisia,	
Tops of Wormwood,	
Tansey,	
Leaves of Scordium,	
Senna,	
Rhubarb,	
Mix.	

VERMIFUGE BOLUSES.

Boli anthelmintico-tonici. (BRERA)

R. Corsican Moss,	} of each, one scruple.
Extract of Artemisia	
Seeds,	
Make six boluses.—Dose, one, every two hours.	
R. Corsican Moss,	} of each, one scruple.
Extract of Artemisia	
Seeds,	
Make six boluses.—Dose, one, every two hours.	

INFUSION OF CORSICAN MOSS.

Infusum anthelminticum. (SWEDIAUR)

R. Corsican Moss, from two to eight drams.
Boiling Water, one pound.
Macerate for an hour, and strain.—Dose, one pound, every morning, to be taken by glassfuls.

DECOCTION OF CORSICAN MOSS.

Decoctum Helminthocorti. (FORM. HÔP. FR.; DISP. CHA. PARIS; NIEMANN; RATIER)

R. Corsican Moss, two drams.
Water, a sufficient quantity
to obtain half a pound of strained decoction after half an hour's boiling, express, strain, and add
Simple Syrup, two drams.
Mix. (FORM. HÔP. FR.)

NIEMANN prescribes two drams of moss, enough of water to obtain four ounces of

decoction, and half an ounce of honey.—
RATIER—one ounce of moss, and two pounds of water.—**DISP. CHA. PARIS**—one ounce of moss, two pounds and a half of water, and four drams of liquorice.

VERMIFUGE DECOCTION.

R. Corsican Moss, half an ounce.
 Herb of Gratiola, one dram.
 Water, six ounces.

Boil down to five ounces, and add to the strained decoction

Sulphate of Iron, half a dram.
 Honey, half an ounce.

Mix.—Dose, a spoonful, every two hours, for an adult.

JELLY OF CORSICAN MOSS. (**AMST.** ; **GALL.** ; **CADET DE GASSICOURT** ; **NIE-MANN** ; **VAN MONS**)

R. Corsican Moss, one ounce.
 Water, two pounds.

Boil slowly in a covered vessel till reduced to eight ounces. Strain, and add

Sugar, one ounce and a half.

Boil down to five ounces, and set aside in a cool place. (**AMST.**)

R. Corsican Moss, four ounces.
 Spring Water, four pounds.

Boil, and add

White Wine, one pound.
 — Sugar, one pound and a half.
 Isinglass, two drams.

Clarify, strain, and evaporate. (**GALL.**)

NIE-MANN gives the same formula, suppressing the wine.

R. Corsican Moss, two pounds four ounces.
 Red Wine, twelve pounds.

Infuse for twelve hours in a water-bath ; then boil, strain, and put it on the fire again with

White Sugar, twelve pounds.

Clarify, strain, and reduce the whole to about fifteen pounds. (**CADET DE GASSICOURT**)

R. Corsican Moss, one part.
 Water, seven parts.

Infuse for two hours in a covered vessel at almost the boiling heat, and strain with expression : infuse the residuum, in a water-bath, in

Rhenish Wine, one part and a half.

Express at the end of twelve hours. Evaporate the former infusion to one part and a half, mix it with the former, and dissolve

White Sugar, six parts.

Skim, and strain through a close flannel. (**VAN MONS**)

Dose, from two drams to one ounce or more.

SYRUP OF CORSICAN MOSS. (**GENEV.** ; **PIERQUIN**)

R. Corsican Moss, twelve ounces.
 Water, eight pounds.

Boil for two hours, and strain : boil the residue in

Water, five pounds.

Strain. Mix the decoctions, and add

Sugar, three pounds,

and boil to the consistence of syrup. (**PIERQUIN**)

GENEV. prescribes four ounces of moss, a sufficient quantity of water, and one pound of sugar.

To be taken by spoonfuls.

VERMIFUGE SYRUP. (**CADET DE GASSICOURT**)

R. Corsican Moss, twelve ounces.
 Water, a sufficient quantity
 to exhaust it by boiling. Strain, and add to the liquor

White Wine, twelve ounces.
 White of Egg, one.
 Sugar, four pounds.
 Cochineal, thirty-six grains.
 Alum, twenty-four grains.

Boil, filter, and boil again to the proper consistence.—To be taken by spoonfuls alone, or diluted with ordinary drink.

VERMIFUGE POTION. (**CADET DE GASSICOURT** ; **RATIER** ; **VAN MONS**)

R. Corsican Moss, one dram.
 Boiling Water, four ounces.
 Simple Syrup, one ounce.

Infuse, and mix. (**RATIER**)

RATIER elsewhere prescribes from three to six drams of moss, six ounces of water, and one ounce of syrup of honey.

R. Corsican Moss, one dram.
 Boiling Water, six ounces.

Boil for eight minutes, and to the strained liquor when cold, add

Orange-flower Water, half an ounce.
 Lemon Juice, one ounce.

Mix. (**CADET DE GASSICOURT** ; **SAINTE-MARIE**)

To be taken at one dose, in the morning, fasting, for three or four days.

VERMIFUGE LOHOCH. (**BORIES** ; **PIERQUIN** ; **RATIER**)

R. Mucilaginous Lohoch, four ounces.

Powdered Corsican Moss, half a dram.
 Oil of Almonds, } of each, one ounce.
 Syrup of Lemon, }

Mix. (RATIER)

R. Strong Infusion of Corsican Moss, four ounces.
 Castor Oil, two ounces.
 Syrup of Peach Flowers, one ounce.
 Proto-chloruret of Mercury, four grains.
 Rhubarb, in powder, twelve grains.

Mix. (PIERQUIN)

R. Oil of Almonds, four ounces.
 Powdered Corsican Moss, one dram.
 Syrup of Lemon, one ounce.
 Camphor, twenty grains.
 HOFFMANN'S Liquor, twenty-five drops.
 Orange-flower Water, } of each,
 Purslain Water, } two ounces.
 Mother-wort Water, }

Mix. (BORIES; PIERQUIN)

HELYCHRYSUM.

Two species of this genus of plants are employed :

1. *Helychrysum arenarium*, GÆRTN.

SYNONIME :—*Stæchas citrin*, FR.

BRUNS.; GALL.; WIRTEM.; HERBIP.; BERGIUS.; MURRAY and GMELIN.; SPIELMANN.

A plant of Europe. (*Syngenesia Polygamia Superflua*, LINN.; *Synantherææ*, CASS.; *Fig. Zorn, Ic. pl. t. 352.*)

The flowers (*Flores Stæchadis citrinæ seu Germanicæ seu Amaranthæ lutei seu Helychrysi*) are employed. They are radiated, oval, cup-shaped, of a citrine-yellow or

golden colour. They have very little odour, and a slightly styptic taste.

2. *Helychrysum Stæchas*, CAND.—Everlasting Flower.

SYNONIMES :—*Stæchas de Naples*, FR.; *Gemeines Ruhrkraut*, GERM.

GALL.

A plant of the middle of Europe. (*Fig. Zaun. Ist. p. 92. t. 279.*)

The flowers (*Flores Stæchadis Neapolitanæ*) are employed. They are sub-cylindrical radiated cup-shaped, about two lines in length, of a yellow colour, and inodorous.

HENCHERA AMERICANA, LINN.—AMERICAN SANICLE, ALUM ROOT.

AMER.; COXE.

A plant of North America. (*Pentandria Digynia*, LINN.; *Saxifragææ*, JUSS.; *Fig. Herm. Parad. t. 131.*)

The root is employed; it is very astringent.

HEPATICA TRILOBA, CAND.—HEPATICA, LIVERWORT.

SYNONIMES :—*Hépatique des jardins*, FR.; *Leberkraut*, GERM.; *Gaternjk*, BOHEM.; *Liverurt*, DAN.; *Fegatella*, SPAN.; *Le-verkruid*, DUT.; *Watrobnik Ziele*, POL.; *Hepatica*, PORT.; *Blasippa*, SWED.

BRUNS.; HISP.; GALL.; GENEV.; WIRTEM.; HERBIP.; BERGIUS.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of Europe. (*Polyandria Polygamia*, LINN.; *Ranunculacææ*, JUSS.; *Fig. Zorn, Ic. pl. t. 5.*)

The herb (*Herba Hepaticæ nobilis seu Trifolliis aurei*) consists of coriaceous leaves, crenated and heart shaped at the base, half

divided into three lobes, glabrous above, and slightly pubescent underneath.

The flowers are of a grey, red, or violet colour.

It has an astringent taste, but is without odour.

It is slightly astringent.

HERACLEUM SPONDILIUM, LINN.—COW PARSNIP.

SYNONIMES:—Berce, Fausse acanthe, FR.; Unæchte Bärenklau, GERM.; Esfondilio, SPAN.; Beerenklaw, DUT.; Spondilio, ITAL.; Parszoz porstinaski, POL.; Canabraz, PORT.

AMER.; GALL.; WIRTEM.; COXE; MURRAY and GMELIN.

A plant of Europe and North America. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Blackw. Herb. t. 540.*)

The root and herb are employed.

The root (*Radix Brancæ ursinæ* seu *Ursi*

seu *Spondylii* seu *Germanicæ*) is long, thick, and branching. It is white, and has an acrid and caustic taste.

The herb consists of a grooved stem, furnished with winged, rough leaves, whitish underneath, having crenulated, lobed leaflets. It has an agreeable and sweetish taste.

ORNE recommends the powdered root against epilepsy, in the dose of two drams, prescribing at the same time an infusion of the leaves. MANN has administered a decoction of it in dyspepsia, with success.

HERMODACTYLI,—HERMODACTYLES.

SYNONIMES:—Hermodactes, Hermodattes, FR.; Fremde Zeitlosenwurzel, GERM.

ANT.; BRUNS.; HISP.; GALL.; WIRTEM.; BERGIUS; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

This name is given to certain tuberosities which grow on the collar (*au collet*) of an undetermined liliaceous plant (*Fig. Lob. Ic. 146*), growing in the East, and which GROVONIUS improperly designates *Colchicum Illyricum*. [Sir H. HALFORD has stated that the hermodactyles he had from the East are a species of *colchicum*.—EDIT.]

They are compressed, almost triangular, concave and grooved on one side, swollen on the other, the circumference rounded and

blunt, yellowish without, and white within, tolerably dense, fragile, and of a grained texture. Their taste is very feeble, and their odour strong and nauseous.

They should not be confounded with the white tuberosities of the size of a nut, which are disposed in the form of fingers round the root of the *Iris tuberosa*, LINN.; a plant of Europe and of the Levant; and which are called *Spurious Hermodactyles*.

Hermodactyles are reputed, without reason, to be slightly laxative. LECANU has found in them neither emetine nor veratrine, but a fatty matter, much starch, a yellow colouring principle, gum, and various salts.

HERNIARIA.—RUPTURE-WORT.

Two species of this genus are used in medicine:

1. *Herniaria glabra*, LINN.

SYNONIMES:—Herniole glabre, Turquette, FR.; Bruchkraut, GERM.; Bridurt, DAN.; Yerba turca, SPAN.; Klein duitend knoof, DUT.; Sporyz trzeci, POL.; Herniaria, PORT.; Brackært, SWED.

BRUNS.; HISP.; GALL.; GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of Europe (*Pentandria Digynia*, LINN.; *Paronychieæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 382.*)

The herb (*Herba Herniariæ* seu *Milligranæ* seu *Empetri*) is employed. It con-

sists of a branching, slender, glabrous stem, furnished with very small oval, oblong, green leaves. It has a feeble and herbaceous odour, and a bitter saline, and styptic taste.

It is slightly astringent.

2. *Herniaria hirsuta*, LINN.

GALL.; SPIELMANN.

A common plant of Europe. (*Fig. Zaur. Ist. p. 138. t. 254.*)

The herb differs from the preceding only by its hairy leaves.

HIERACIUM.

Two species of this genus are used in medicine:

1. *Hieracium Murorum*, LINN.

SYNONIMES:—Epervière des murailles, Pulmonaire des Français, FR.; Mauershabichtskraut, GERM.

GALL.

A plant common throughout Europe. (*Syngenesia Polygamia Æqualis*, LINN.; *Synantherææ*, CASS.; *Fig. CEd. Fl. Dan. t. 1513.*)

The herb (*Herba Pulmonariæ Gallicæ* seu *Auriculæ Muris majoris*) is employed. It consists of a rosette of oval, radical leaves, slightly toothed, rather angular towards the base, slightly crenated at the insertion of the foot-stalk, thin, and very hairy underneath, on the margin, and on the foot-stalks. Its taste is feebly bitter and saline.

It is tonic.

2. *Hieracium Pilosella*, LINN.—Mouse-ear.

SYNONIMES:—Oreille de Souris, Piloselle,

FR.; Nagelkraut, Habichtskraut, Mausährchen, GERM.; Mauseære, DAN.; Pilosella, PORT.; and SPAN.; Nagelkruid, DUT.; Pelosella, ITAL.; Nidoszialec, POL.; Musæan, SWED.

BRUNS.; HISP.; GALL.; WIRTEM.; MURRAY and GMELIN; SPIELMANN.

This plant is common throughout Europe. (*Fig. Zorn, Ic. pl. t. 174.*)

The herb and flowers are employed.

The herb (*Herba Pilosellæ* seu *Auriculæ Muris*) consists of a slender, pubescent, whitish stem, and of oval, oblong, obtuse, entire leaves, narrowed at the base into foot-stalks, green above, and bristled, particularly towards the margin, with long, white, scattered hairs; beneath it is whitish and cottony.

The flowers are yellow semi-flosculous cups.

The taste of this plant is bitterish. It was formerly prescribed in diarrhœa and dysentery.

HIRUDO OFFICINALIS, LINN.—MEDICINAL LEACH.

SYNONIMES:—Sangsue, FR.; Blutigel, GERM.; Blodigle, DAN.; Sanguisuela, SPAN.; Mignatta, ITAL.; Blodiglar, SWED.

DUBL.; ANT.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FORM. HÔP. FR.; GENEV.; HAMB.; HASS.; OLDEN.; CAST. BORUS.; BORUS.; SAX.;

SUEC.; WIRTEM.; BRUGNATELLI; COXE; GUIBOUT; SPIELMANN.

An aquatic, invertebrate, red-blooded animal (*Annelida Enterobranchia hirudineæ*, LATR.). Its body is long and blackish, with lines of various colours, spotted underneath with yellow.

HORDEUM VULGARE, LINN.—BARLEY.

SYNONIMES :—Orge, FR.; Gerste, GERM.; Geemen, BOHEM.; Byg, DAN.; Cebada, SPAN.; Gerst, DUT.; Orzo, ITAL.; Jecznicz, POL.; Cevada, PORT.; Korn, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

This plant is cultivated in every part of Europe. (*Triandria Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 509.*)

The seed (*Hordeum crudum seu integrum*) is employed. It is oval, oblong, bellied, angular, pointed at the two ends, yellow, and grooved longitudinally. When deprived of its yellow pellicle it is called *Decorticated Barley* (*H. mundatum seu decortiatum*:—Orge mondé, FR.; Geschalte gerste, Gerstengraupe, GERM.; Orzo scorticato, ITAL.; Korngryn, SWED.) which, again, after being reduced to small white spheres, marked with a black longitudinal line, takes the name of Pearl Barley (*H. perlatum*:—Orge perlé, FR.; Perlgraupe, GERM.; Orzo perlato, ITAL.)

Barley contains hordeine, starch, sugar, gum, gluten, and resin. It is analeptic, and is also reputed emollient.

PREPARED BARLEY MEAL.

Farina Hordei præparata, Amylum Hordei, Hordeum præparatum. (BAVAR.; HANN.; SAX.)

R. Barley Meal, any quantity.

Enclose it in a linen bag, leaving a space between it and the ligature; put the bag into a tinned copper pan filled with water, and boil for twelve hours, replacing the water in proportion as it becomes evaporated: then dry the meal, after having removed the thick paste which envelopes it. When dried, take off the brown crust and horny mass which is formed, and pulverize the rest.

It is analeptic; and is recommended in pulmonary consumption, atrophy, and the wasting of the flesh attending lactation.—One or two spoonfuls are to be boiled with sixteen ounces of cow's milk, and sweetened. This quantity is for an adult; one half is sufficient for a child under twelve years.

BARLEY SUGAR.

Saccharum hordeatum seu hordaceum. (BRUNS.; NIEMANN.; SPIELMANN)

R. Barley, eight ounces.
Spring Water, one hundred ounces.
Boil for an hour, strain, and add
Sugar, sixteen ounces.

Boil again till the thread which detaches itself from the spatula, breaks of itself; then pour it upon an oiled slab, and cut it into convenient pieces. (SPIELMANN)

NIEMANN prescribes one pound of sugar, and a sufficient quantity of decoction of barley.—BRUNS.—one pound and a half of decoction, two pounds of sugar, and four grains of saffron.

DECOCTION OF BARLEY; BARLEY WATER.

Decoctum Hordei integri seu decorticati, Aqua Hordei, Ptisana Hippocratica. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BELG.; GALL.; FERR.; FORM. HÔP. FR.; HASS.; LIPP.; LUSIT.; ROSS.; SUEC.; HERBIP.; WURCEB.; BRERA.; BRUGNATELLI.; COXE.; CADET DE GASSICOURT.; DISP. CHA. PARIS.; PIDERIT.; RATIER.; SPIELMANN.; VAN MONS)

Decoctum Hordei. (LOND.; DUBL.)

Decoctum Hordei Distichi. (EDINB.)

R. Pearl Barley, two ounces.
Water, four pints and a half.

First wash off extraneous matters with cold water, and then boil for a short time in half a pint of water. This being rejected, pour on the remainder of the water, boiling; then boil down to one half, and strain. (LOND.)

DUBL.; and EDINB. require five pounds of water for the second boiling.

The same formula, with unimportant variations, is given by AMER.; ANT.; BATAV.; BELG.; HASS.; HERBIP.; COXE.; and PIDERIT.

R. Pearl Barley, two ounces.
Water, twelve ounces.

Boil for a short time and reject the liquor; then add more water, boiling, in such quantity that after a short boiling, and when the barley has burst, there remains about two pounds of strained decoction. (BATAV.)

R. Pearl Barley, washed in hot water,
one part.
Water, fifty parts.

Boil down to thirty-four parts; strain the decoction through close flannel, set it aside, and pour off thirty-two parts. (VAN MONS)

FERR.; and BATAV. prescribe two ounces of washed pearl barley and four pounds of water, to be reduced to two.—ROSS.—six ounces of barley and ten pounds of water, to be reduced to eight.—LIPP.; and BRUGNATELLI—two ounces of barley and five pounds of water, to be reduced to two and a half.—AMST.—two ounces of barley and six pounds of water, to be reduced to three.—LUSIT.—three ounces of barley and four pounds of water, to be reduced to two.—CADET DE GASSICOURT—half an ounce of barley and one pound and a half of water, to be reduced to one pound.

R. Decorticated Barley, rubbed with
a coarse cloth, and washed,
half an ounce.

Water, two pounds and a half.

Boil down to two pounds ; strain and add to the decoction,

Syrup of Marshmallows, one ounce.

Mix. (GALL.)

R. Barley, washed several times
with hot water, three ounces.
Root of Dog-grass, one ounce.
Spring Water, ten pounds.

Boil down to eight pounds, and strain.
(SUEC.; SPIELMANN)

R. Decorticated Barley, coarsely
ground, four ounces.
Spring Water, thirty-two pounds.

Boil for a quarter of an hour, and strain
Liquorice Root. two ounces.

Boil down slowly to two thirds. (WURCEB.)

SPIELMANN directs two ounces of washed barley and one ounce of dog-grass to be boiled in one hundred and twenty-eight ounces of water till reduced to ninety-six and half an ounce of liquorice to be added towards the end.—**FORM. HÔP. FR.**—half an ounce of barley to be washed in boiling water, and then to be boiled in a sufficient quantity of water to obtain two pounds of strained decoction, adding towards the end, one dram of liquorice.—**RATIER**—one ounce of barley to be boiled for five minutes in two pounds of water; this liquor to be rejected, three pounds of water added, and the boiling continued till reduced to two pounds, adding towards the end one dram of liquorice.—**VAN MONS**—to boil one part of dog-grass with one and a half of barley, in ninety-six parts of water, till reduced to sixty-six, adding towards the end, one part of liquorice; the liquor to be strained before cold, and sixty-four parts decanted.—

SWEDIAUR—to boil one ounce and a half of pearl barley and the same of dog-grass, in sixteen pounds of water, to be reduced to twelve, adding one ounce of liquorice.—**SPIELMANN**—to boil one ounce of barley and half an ounce of dog-grass, in sixty-four ounces of water, till reduced to one half, adding two drams of liquorice.—**PIERQUIN**—to boil one pound of barley and half a pound of dog-grass, in an arbitrary quantity of water, till reduced to forty-four pounds, adding towards the end, four ounces of liquorice.—**SPIELMANN**—to boil one ounce of barley in five pounds of water, till reduced to four, adding towards the end, four drams of liquorice.

This is one of the most ancient medicinal drinks we know of; and it is from this that others have been called *Ptisans*, which is derived from the Greek *πιττανή*, *decorticated barley*. It is necessary first to wash the barley, and then to submit it to a short boiling, in order to take up an astrigent principle which might communicate to the water other than the demulcent and calming virtues which are imparted by the boiled grain.

DECOCTION OF BARLEY WITH GUM.

Aqua hordeata cum gummi. (GENEV.;
AUGUSTIN)

R. Barley, three ounces.

Boil it with about one half of water, and reject this. Then pour upon it

Boiling water, four pounds.

Boil down to one half, adding towards the end.

Liquorice, half an ounce.

Dissolve finally in the liquor.

Gum Arabic, one ounce and a half.

To be taken by glassfuls. (GENEV.)

R. Pearl Barley,	six or seven drams.
Gum Arabic,	one dram.
Spring Water,	two pounds.

Boil, strain, and add

White Sugar, a sufficient quantity
to sweeten. (AUGUSTIN)

ACIDULATED DECOCTION OF BARLEY.

Decoctum Hordei acidulatum seu *acidulum*
seu *citratum*. (SUEC.; SAINTE-MARIE;
SPIELMANN; SWEDIAUR)

R. Pearl Barley, two ounces.

After having washed it, boil it for a short time with half a pound of water: reject this and boil it again with

Water, five pounds.

till one half of it is consumed. Add to it when strained,

Oxymel, two ounces.

Mix. (SAINTE-MARIE; SWEDIAUR)

R. Barley, washed several times with hot water, three ounces.

Root of Dog-grass, one ounce.

Spring Water, ten pounds.

Boil down to eight pounds, and add when strained,

Oxymel, four ounces.

Mix. (SUEC.; SPIELMANN)

R. Lemons, sliced, five.

Decoction of Barley, six pounds.

Boil till reduced to two thirds, strain, and add

Sugar, four ounces.

Mix.

The acidulated decoction of barley is more grateful to patients than the simply sweetened decoction. In general, from half an ounce to one ounce of oxymel is sufficient for one pound of decoction. The juice of pomegranate or of gooseberries, strawberry vinegar, citric or tartaric acid, supertartrate of potass, &c., may also be employed. It is a drink which it is well to know how to vary almost to infinity, in order to accommodate it to all the tastes and caprices of patients, it being one of those most employed, and one of the easiest to be procured.

ANTI-SCORBUTIC DECOCTION. (NIEMANN)

R. Decorticated Barley, eight ounces.
Water, eight pounds.

Boil down to six pounds, and add

Expressed Juice of Cranberries, four ounces.

Syrup, any quantity.

Mix.

COMPOUND DECOCTION OF BARLEY.

Decoctum Hordei compositum. (LOND.; DUBL.; HISP.; HAMB.; HERBIP.; SWEDIAUR)

R. Decoction of Barley, two pints.
Figs, sliced, two ounces.

Liquorice Root, cut and bruised, half an ounce.

Raisins, stoned, two ounces.
Water, one pint.

Boil down to two pints, and strain. (LOND.)

R. Decoction of Barley, four pints.
Raisins, stoned, } of each,

Figs, stoned, } two ounces.

Liquorice Root, cut and bruised, half an ounce.

VOL. II.

To the boiling decoction first add the raisins, then the figs, and, lastly, the liquorice towards the end of the boiling, which is to be continued, until two pints of strained decoction remain. (DUBL.; and COXE)

R. Decorticated Barley, half an ounce.
Water, three pounds.

Boil down to two pounds, adding towards the end,

Dried Leaves of Ground Ivy, two drams.

Liquorice, grated, } of each,
Flowers of Red Pop- } one dram and
pies, } a half.

Strain. (HISP.)

R. Decorticated Barley, two ounces.
Hartshorn Shavings, one ounce.

Spring Water, thirty ounces.

Boil down to two thirds, and add when strained,

Despumated Honey, } of each,
Vinegar, } two ounces.

Mix. (HANN.)

R. Barley, } of each, two ounces.
Figs, }
Raisins, }

Water, four pounds.

Boil down to two pounds, and strain. (SWEDIAUR)

R. Decorticated Barley, one ounce.
Root of Seneka, half an ounce.

Nitrate of Potass, two drams.

Spring Water, a sufficient quantity,

to obtain, after boiling, two pounds of strained decoction. Add to this

Rob of Gooseberries, } of each,
Elder, } six drams.

Oxymel, two ounces.

Mix. (HERBIP.)

PECTORAL DECOCTION. (HISP.; CADET DE GASSICOURT; VAN MONS)

R. Pearl Barley, two ounces.
Water, six pounds.

Boil, and on removing it from the fire, add

Canadian Maidenhair, } of each,
Flowers of Coltsfoot, } half a handful.

Infuse for eight minutes, strain through a sieve, and put into the liquor

Raisins, stoned, two ounces.
Figs, six.

When the liquor is nearly cold, strain without expression, set aside, and pour off four pounds. (VAN MONS)

R. Decorticated Barley, one ounce.
Jujubes, twelve.

Figs, six.

C

Raisins, } of each, half an ounce.
 Liquorice, }
 Maidenhair, }
 Water, three pounds.

Boil down to two pounds, and strain.
 (HISP)

R. Pearl Barley, two ounces.
 Water, six pounds.

Boil, and add

Raisins, } of each, two ounces.
 Figs, }
 Liquorice, }

Boil down then to one half, and add, if necessary, two drams of nitrate of potass.
 (CADET-DE GASSICOURT)

It is demulcent, emollient, and pectoral.
 —Dose, six ounces.

FOREST'S DECOCTION.

Decoctum Petri Foresti. (ANT.; VAN MONS)

R. Red Chick Peas, } of each, one ounce.
 Pearl Barley, }
 Water, a sufficient quantity

to obtain thirty-two ounces of strained decoction. On removing it from the fire, add

Figs, half an ounce.
 Liquorice, two drams.

When cold, strain, and make an emulsion with

Melon Seeds, one ounce.
 Strain. (VAN MONS)

ANT. requires sweet almonds to form the emulsion with.

LIEUTAUD'S REFRESHING APOZEM. (PIERQUIN; SAINTE-MARIE)

R. Barley, four drams.
 Root of Cichory, one ounce.
 Cold Seeds, three drams.
 Leaves of Sorrel, one handful.
 Water, two pounds.

Boil till reduced to a pint and a half. To be taken in the morning, fasting, at four glassfuls, adding to each, a spoonful of syrup of gooseberries.

SOOTHING INJECTION. (PIERQUIN)

R. Ears of Barley, half a handful.
 Liquorice, half an ounce.
 Flowers of Mallow, two handfuls.
 Water, a sufficient quantity.

Boil down to one pound, strain, and add
 Sugar Candy, one ounce.

HUMULUS LUPULUS, LINN.—HOP.

SYNONIMES: — Houblon, FR.; Hopfen, GERM.; Humle, DAN.; and SUEC.; Lupulo, ITAL.; and SPAN.; Hoppe, DUT.; Chmiel, POL.; Hombrecillo, PORT.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BAVAR.; BRUNS.; HISP.; GALL.; FENN.; FORM. HOP. FR.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY.; and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.

This plant is cultivated in all parts of Europe. (*Diacia Pentandria*, LINN.; *Urticeae*, JUSS.; *Fig. Zorn, Ic. pl. v. 522, 541.*)

The cones of the female flowers, called *Hops* (*Strobili seu Coni seu Flores Lupuli*) are employed. They are ovoid, formed of membranous, oval, concave, imbricated yellowish scales. They have a strong and aromatic odour, and a bitter taste.

They contain, according to YVES, a peculiar principle *Lupuline*, which is the source of their activity.

They are tonic and sub-narcotic; and are principally prescribed in scrofulous disorders, rachitis, and tabes mesenterica, and are also extolled as febrifuge and sedative.

Dose, from half a dram to two drams, of the powder; and from two to four ounces of the expressed juice.

AROMATIC SPECIES. (POL.; BORUS.; SARD.; NIEMANN)

R. Hops, one pound.
 Chamomile Flowers, }
 Lavender Flowers, } of each,
 Herb of Rosemary, } three ounces.
 — Wild Thyme,
 — Tansey,

Cut, and mix.

AQUEOUS EXTRACT OF HOPS.

Extractum Humuli seu Lupuli. (LOND.; ANT.; HANN.; LIPP.; SUEC.; COXE.; VAN MONS)

R. Hops, four ounces.
 Water, one gallon.

Boil down to four pints, strain while hot, and evaporate. (LOND.; COXE)

R. Hops, two pounds.
 Water, ten pounds.

Boil for a quarter of an hour, and ex-

press; set aside and decant the liquor, then evaporate to the proper consistence. (ANT.)

R. Hops, } of each, one part.
Cold Water, }

Macerate for twelve hours and add
Boiling Water, four parts.

Strain when cold, clarify the liquor with white of egg, put it on the fire, and when it begins to thicken, strain it through a sieve, and evaporate, constantly stirring. (VAN MONS)

R. Hops, one part.
Boiling Water, eight parts.

Boil for a quarter of an hour, and express: boil the residue with four parts of water, and strain again. Mix the decoctions, decant after sufficient repose, and evaporate slowly to the required consistence. (HANN.)

SUC. directs one part of hops to be boiled with two parts of water, and then expressed; the residuum to be boiled in fresh water, and the united liquors evaporated in a water-bath. — LIPP. — three pounds of hops to be macerated for twelve hours in nine of boiling water; then to be boiled, expressed, and evaporated.

Dose, from half a dram to one dram.

ALCOHOLIC EXTRACT OF HOPS. (AUSTR.)

R. Hops, two parts.
Alcohol (0.910), three parts.
Water, nine parts.

Digest in a water-bath, express, distil off the alcohol, and evaporate the rest in a water-bath, to the proper consistence.

The same dose for this as for the preceding.

TINCTURE OF HOPS.

Tinctura Humuli seu Lupuli seu Humuli Lupuli seu Lupuli Strobilorum. (LOND.; EDINB.; AMER.; ANT.; HANN.; SUC.; COXE; NIEMANN; SWEDIAUR; VAN MONS)

Tinctura Humuli. (LOND.)

R. Hops, five ounces.
Proof Spirit, two pints.

Macerate for fourteen days, and strain.

Tinctura Humuli Lupuli. (EDINB.; SUC.)

R. Hops, five ounces.
Dilute Alcohol, five pounds and a half.

Digest for seven days, express and filter.

NIEMANN; and SWEDIAUR prescribe half an ounce of hops, and half a pound of proof spirit. — ANT. — one part of hops and eight parts of alcohol (20 degrees). — HANN. — one of hops and six of alcohol. — AMER.; and COXE. — five ounces of hops and one pound of alcohol.

It is narcotic. — Dose, from half a dram to two drams.

INFUSION OF HOPS. (ELLIS; RATIER)

R. Hops, one ounce.
Boiling Water, two pounds.

Infuse, and strain.

It is aromatic and bitter; and is recommended in scrofula, scurvy, and asthenic disorders.

A more active preparation may be obtained by infusing hops in wine, after having sprinkled it with a sufficient quantity of proof spirit.

DISTILLED WATER OF HOPS.

Aqua Lupuli. (HANN.; NIEMANN; VAN MONS)

R. Hops, eight parts.
Water, a sufficient quantity.

Distil twenty-four parts. (HANN.; NIEMANN)

R. Hops, four parts.
Proof Spirit, one fourth part.
Water, thirty parts.

Distil six parts. (VAN MONS)

It is stomachic and subnarcotic. — Dose, a spoonful.

STOMACHIC MIXTURE.

Mixtura e Lupulo. (NIEMANN)

R. Aqueous Extract of Hops, two or three drams.

Water of Hops, seven ounces.

Tincture of Hops, half an ounce, or one ounce.

Syrup of Orange Peel, one ounce.

Mix. — Dose, a spoonful, every three hours.

OINTMENT OF HOPS.

Unguentum Lupuli. (VAN MONS)

R. Dried Hops, two ounces.
Lard, ten ounces.

Digest in a moderate heat for several hours, and express.

It is recommended in cancer to appease pain.

POWDER OF LUPULINE. (FEE; HENRI; MAGENDIE)

R. Lupuline, one part.
White Sugar, two parts.

Triturate.

OINTMENT OF LUPULINE. (HENRI)

- R. Lupuline, one dram.
Fat, three drams.

Digest in a water-bath, and strain through a close cloth.

TINCTURE OF LUPULINE. (FERR.; FEE; MAGENDIE)

- R. Lupuline, bruised, one ounce.
Alcohol (36 degrees), two ounces.

Digest for six days in a covered vessel, express strongly, and add

Alcohol (36 degrees),

a sufficient quantity

to obtain three ounces of tincture.

It is aromatic, bitter, and tonic.—YVES says besides that this tincture is narcotic.—The dose is not yet well determined.

SYRUP OF LUPULINE.

- R. Tincture of Lupuline, one part.
Syrup, seven parts.

Mix.

HYPOCISTIS.—HYPOCIST.

SYNONIMES:—*Succus Hypocistidis*.—Hypociste, FR.; Hypocistensaft, Zistensaft, GERM.; Hypocistis, BOHEM.

BRUNS.; HISP.; GALL.; WIRTEM.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

This substance is in dry, heavy, black masses, opaque, but shewing a bright fracture. It is inodorous, and possesses an astringent and slightly acid taste.

It is obtained by evaporating to dryness in the sun, the expressed juice, either of the fruit or of the whole plant of the *Cytinus Hypocistis*, LINN., a small parasitical plant of the middle of Europe. (*Gynandria Dodecandria*, LINN.; *Aristolochiæ*, JUSS.; Fig. Cav. Ic. 2. p. 55. t. 171.)

It is astringent.—Dose, from half a scruple to a dram.

HYDRARGYRUM.—MERCURY, QUICKSILVER.

SYNONIMES:—*Hydrargyrum*, *Mercurius*, *Argentum vivum*.—Mercure, Vif-Argent, FR.; Quecksilber, GERM.; Abuk, Zibakk, ARAB.; Chuy-yin, CHIN.; Quecksolv, DAN.; Azogue, Mercurio, SPAN.; and PORT.; Para, Parah, HIND.; and DUK.; Kwekzilva, DUT.; Argento vivo, Mercurio, ITAL.; Rassa, MAL.; Seemab, PERS.; Parada, Rasa, Sutam, SANS.; Quicksilfver, SWED.; Rasam, TAM.; and TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; COXE.; PARIS.

The mercury furnished by commerce ought not to be applied to the purposes of medicine till after it has been freed from the foreign substances which are mixed with it. It is then called *Purified Mercury*; *Mercurius vivus depuratus seu rectificatus*, *Hydrargyrum purificatus*. All pharmaco-

pœias do not indicate a process for this purification.

Hydrargyrum Purificatum.

Pour mercury into an iron retort; then apply heat, and distil the pure mercury. (LOND.)

- R. Mercury, six parts.
Distil slowly four parts. (DUBL.)

Hydrargyrum Purificatus. (EDINB.)

- R. Mercury, six parts.
Iron Filings, one part.

Triturate together, and distil from an iron retort.

AMER.; AMST.; SARD.; and COXE give the same process as EDINB.

BAVAR.; FERR.; GENEV.; HANN.; HISP.; LUSIT.; OLDEN.; ROSS.; SUEC.; WIRTEM.; BRUGNATELLI.; and PIDERIT direct mercury to be distilled from a retort communicating with a globular receiver half filled with water.—FENN.; and HASS.

—the same apparatus to be employed, but a mixture of three pounds of the metal and one ounce of sulphur to be distilled.

BRUNS. requires the metal to be distilled with subcarbonate of potass or with quick lime, and afterwards to be strained through chamois leather.

LIPP. directs it to be distilled into a receiver half filled with water, then to digest it for twelve hours in nitric acid diluted with three parts of water, and lastly to wash it well.

SAX. directs a mixture of equal parts of black sulphuret of mercury and of quick lime, to be distilled.—FULD.—six ounces of roll-sulphur to be melted in an iron mortar, and two pounds of mercury to be added to it by portions; the black mass which results from their intimate union to be powdered and mixed with eighteen ounces of powdered quick lime: this mixture to be put into a retort and the product received in a globular receiver half filled with water.—HISP.; GALL.; FERR.; POL.; and VAN MONS—red sulphuret of mercury to be distilled with an equal weight of iron filings. This process gives the *Mercury revived from Cinnabar*.

The mode of purification recommended by LUT and CHEVALLIER is this. Fill a metal retort with mercury to two-thirds, and place it in a covered furnace; adapt to the neck of the retort an iron tube, to the other extremity of which attach a wetted linen bag which is to be partly suspended in an earthen vessel full of water: the operation being finished, pour off the water, dry the metal in bibulous paper, and strain it through chamois leather.

Simple metallic mercury is employed only in a very few preparations, themselves but seldom used. In this form, it does not exercise any action on the animal economy, except that which may result from its weight, though many erroneously believe it to be poisonous. But when it is in a state of intimate division, it becomes one of the most powerful stimulants known, and may prove poisonous, either by its quantity or by the continuity of its action, although, in the latter case, its most ordinary effect is that of producing a series of pathological phenomena, which terminate in death very gradually.

Although it is well demonstrated that mercury is introduced into the circulation through the medium of the absorbents, yet we are still ignorant, whether its effects depend on a direct action exercised by it on organs, into the tissue of which it may be carried by the blood; or whether they result from the sympathetic action of tissues associated with each other by nervous connexions, or by a similarity of structure with

that, at the surface to which it is applied: but this last hypothesis seems, at present, the more probable. It has in its favour the infinite variability of the effects of mercury, according to external circumstances, the constitution or condition of the subject, the nature of the preparation, the dose, &c. This variability has made mercury to be classed at the same time among the antispasmodics, diuretics, emmenagogues, stimulants, purgatives, sialagogues, and sudorifics; according as it may have been found to augment more particularly one or other of the secretions or excretions.

METALLIC MERCURY.

I. *Preparations in which the action of Mercury is not modified, at least in a notable manner, by other substances.*

A.

Mercury in mass.

POTION OF MERCURY.

Potio Mercurii vivi. (FULD.; BRUGNATELLI)

R. Purified Mercury,

from two to four ounces.

Divide into eight doses, one to be taken every eight hours, in intromusception.

BRERA has administered in this manner twenty-six ounces of mercury to a patient who was not, he says, cured, but remarkably relieved (*soulagé notablement*).

SIMPLE MERCURIAL WATER.

Aqua Mercurii vivi cocta. (FULD.; BRUGNATELLI)

R. Mercury, two pounds.
Water, four pounds.

Boil for two hours in a matrass, and decant.

It is anthelmintic.—It is administered, alone or sweetened, to children; to free them from ascarides or lumbrici.

Dose, from half an ounce to three ounces, for subjects of from one to six years of age.

DECOCTION OF MERCURY.

Decoctum Hydrargyri. (HERBIP.)

R. Purified Mercury, two ounces.
Distilled Water of Tansey, one pound.

Boil in a covered earthen pot, constantly moving the metal with a rod; decant the water, and add to it,

Syrup of Peach Flowers, one ounce.

Mix.

It is anthelmintic.

ROSENSTEIN'S ANTHELMINTIC DECOCTION.
(AUGUSTIN; PIDERIT)

- R. Root of Dog-grass, half an ounce.
Orange Peel, half a dram.
Cinnamon, half a scruple.
Spring Water, eight pounds.

Boil to the consumption of three-fourths;
then strain and add.

Mercury, two ounces.

Boil again in a covered vessel, stirring
frequently, and decant slowly.

It is recommended in verminal disorders
of children, in the dose from eight ounces
to one pound, alone or with milk.

RESOLVENT PLASTER. (BORIES)

- R. Emplastrum Diabotani, two ounces.
Juice of Nightshade, two drams.

Triturate with yolk of egg, and add
Amalgam of equal parts of Mercury
and of Lead, two drams.
Spermaceti, one dram and a half.

Mix.

B.

Mercury divided by means of Mucilage.

MERCURIAL MUCILAGE OF GUM ARABIC.

Mucilago mercurialis Plenki. (SAX.; WIRTEM.; BRUGNATELLI)

- R. Purified Mercury, one dram.
Gum Arabic, two drams.
Water, two spoonfuls.

Triturate the mixture in a glass mortar,
with a wooden pestle, till the metal shall
have disappeared. (BRUGNATELLI)

SAX.; and WIRTEM. prescribe one dram
of mercury, half an ounce of gum and a
sufficient quantity of water.

It is stimulant. The revulsive action
which it sometimes exerts has made it to be
regarded as debilitant, for it has been re-
commended in inflammatory dysentery and
in some other internal phlegmasiæ.

It has been employed externally, in ure-
thritis and in syphilitic ophthalmia and ul-
ceration of the throat. It has been also ad-
ministered against lumbricoid ascariæ; for
which purpose it is mixed with eight
ounces of fumitory water, and half an
ounce of syrup: of which mixture one or
two spoonfuls are to be taken morning and
evening.

SYRUP OF MERCURY AND OF GUM.

*Syrupus Hydrargyri, Mercurius gummosus
Plenki.* (GALL.; FERR.; FULD.; HASS.;
LIPP.; NIEMANN; SAUNDERS; SPIEL-
MANN; VAN MONS)

- R. Pure Mercury, one dram.
Gum Arabic, in powder, three drams.

Triturate, till the mercury is extinguish-
ed; then add

Syrup of Poppies, half an ounce.

Triturate again to form a mucilage.
(GALL.; FERR.; HASS.; LIPP.; NIE-
MANN; SPIELMANN)

- R. Mucilage of Gum Arabic, six drams.
Mercury, one dram.

Triturate sufficiently, and add

Simple Syrup, four ounces.

Mix well. (FULD.)

- R. Mercury, half a dram.
Gum Arabic, one dram and a half.
Water, three drams.

Triturate till the metal is extinguished;
then add,

Syrup of Violets, two ounces.
Elder-flower Water, one ounce.

Mix. (SAUNDERS)

- R. Gum Arabic, three drams.
Water, seven drams and a half.

Make a mucilage, and add

Mercurial Ointment, made with
Cocoa-nut Oil, one dram and a half.

Triturate together, and incorporate gra-
dually,

Simple Syrup, six ounces.

Preserve. (VAN MONS)

In administering this syrup, it is to be so
dosed that from three to six or eight grains
be taken daily. The last formula scarcely
differs from the succeeding.

MERCURIAL MARMALADE.

Conditum hydrargyro-unguinosum. (SWE-
DIAUR; VAN MONS)

- R. Mercurial Ointment, made with
Cocoa-nut Oil, three drams.
Mucilage of Gum Arabic, six drams.

Mix together and make an emulsion;
then add gradually,

Honey, four ounces.

This preparation is recommended in acute
urethritis suppressed by the abuse or un-
timely use of purgatives.—Dose, from two
to four tea-spoonfuls.

LAGNEAU'S GUMMY SYRUP OF MERCURY.
(CADET DE GASSICOURT)

- R. Mercury, one scruple.
Gum Arabic, one ounce.
Compound Syrup of Cichory,
a sufficient quantity.

Triturate in a glass mortar to the perfect
extinction of the metal, and add

Compound Syrup of Cichory,
one ounce and a half.

Mix.—To be administered in the same mode as the preceding.

MERCURIAL MIXTURE.

Mixtura Mercurii gummosa. (BRUNS.; LUSIT.; SARD.)

R. Mercury, one dram.
Mucilage of Tragacanth, half an ounce.

Triturate to extinction, and add

Simple Syrup, one ounce.
Distilled Water, eight ounces.

Mix. (LUSIT.)

R. Mercury, one dram.
Gum Arabic, two drams.
Simple Syrup, half an ounce.

Triturate the metal to extinction, and add,
Mint Water, six ounces.
Compound Juniper Water, one ounce.

Mix. (SAUNDERS)

R. Gum Arabic, three drams.

Sprinkle it, in a marble mortar, with a little water, that it may take the consistence of mucilage; then add

Mercury, one dram.

Triturate it to perfect extinction, and add gradually;

Kermesine Syrup, half an ounce.
Fumitory Water, eight ounces.

Mix. (BRUNS.)

The mode of administration has been indicated above, in the article of mercurial mucilage.

SCHLESINGER has prescribed a preparation analogous to this in its mode of action, which he designates *Solutio Hydrargyri gummosi*; the following is the formula:

R. Mercury, ten grains.
Spermaceti, half a dram.
Yolk of Egg, one.
Syrup of Marshmallows, one ounce.
Distilled Water, two ounces.

Triturate and mix. (AUGUSTIN)

BALSAMIC MERCURIAL MIXTURE. (BRUNS.)

R. Copaiba, }
Gum Arabic, } of each, half a dram.
Mercury, one dram.

Triturate to extinction, and add

Syrup of Kermes, } of each,
Fumitory Water, } two ounces.

Mix.

It is recommended particularly in urethritis.

PLENK'S MERCURIAL MILK. (NIEMANN; SAINTE-MARIE)

R. Mercury, one dram.
Gum Arabic, four drams.
Syrup of Poppies, a sufficient quantity

to make a mucilage. After having triturated it for a long time in a mortar, add gradually to it,

Boiling Cow's Milk, eight ounces.

It is recommended in lotions, in gonorrhœic ophthalmia; in baths, in phymosis caused by ulceration of the glans; and in gargles, in syphilitic ulceration of the pharynx, &c.

SIMPLE MERCURIAL INJECTION.

Injectio hydrargyrica simplex. (SWEDIAUR)

R. Mercurial Ointment, two drams.
Mucilage of Gum Arabic, half an ounce.

Triturate together to form an emulsion; add gradually to it,

Water, three ounces.

It is employed in chronic urethritis.—Two drams are to be injected four times a day.

PLENK'S MERCURIAL INJECTION.

Injectio mercurialis Plenktii. (SPIELMANN)

R. Mercury, one dram.
Gum Arabic, three drams.
Syrup of Poppies, half an ounce.

Triturate to extinction, and add

Spring Water, twelve ounces.
Rue Water, two ounces.

It is recommended in gonorrhœa and leucorrhœa.

HUFELAND has proposed the following similar but more simple injection.

R. Mercury, one dram.
Gum Arabic, half an ounce.
Syrup of Opium, a sufficient quantity.

Triturate the mercury with the gum; adding the syrup gradually till the whole is reduced to a mucilage. (AUGUSTIN)

PLENK'S MERCURIAL GARGLE.

Gargarisma mercuriale Plenktii. (SPIELMANN)

R. Mercury, half a dram.
Gum Arabic, three drams.
Syrup of Poppies, half an ounce.
Proto-chloruret of Mercury, six grains.

Triturate to extinction, and add

Decoction of half an ounce of
the Herb of Upright Clematis
in water, twenty-six ounces.
Honey of Roses, one ounce.
Tincture of Myrrh, one dram.

It is extolled in syphilitic pharyngitis and
ozæna.

LAGNEAU'S PASTILLES. (CADET DE GAS-
SICOURT)

R. Sugar, ten ounces.
Mercury, two ounces.
Gum Arabic, one ounce.
Vanilla, one dram.

Make, according to art, one hundred and
seventy six pastilles, each of which contains
two grains of mercury.

TROCHISCS OF MERCURY. (PH. SYPH.)

R. Mercury, one ounce.
Sugar Candy, two ounces.

Triturate with a few drops of water to
the perfect extinction of the metal; then add

Mucilage of Gum Arabic,
a sufficient quantity,
and make trochiscs of six grains each.

PLENK'S MERCURIAL PILLS.

*Pilulæ mercuriales Plenkii seu Hydrargyri
gummosæ.* (ANT.; BELG.; PH. SYPH.;
AUGUSTIN; VAN MONS)

R. Mercurial Mucilage, one part.
Crumb of Bread, one part and a half.
Honey, a sufficient quantity
to make a pilular mass. (VAN MONS)

R. Mercury, one dram.
Gum Arabic, } of each, two drams.
Honey, }
Liquorice Powder, half an ounce.

Make pills of three grains each. (AU-
GUSTIN)

R. Mercury, one scruple.
Starch, one dram.
Mucilage of Gum Arabic,
a sufficient quantity.

Triturate to perfect extinction, and add

Liquorice Powder, a sufficient quantity,
and divide the mass into twenty pills. (PH.
SYPH.)

R. Mercury, one dram.
Gum Arabic, three drams.
Syrup of Rhubarb, a sufficient quantity.

Triturate in a marble mortar to the per-
fect extinction of the metal, and add

Crumb of Bread, half an ounce.

Make a pilular mass. (ANT.; BELG.)

PLENK'S COMPOUND MERCURIAL PILLS.
(GALL.; FERR.; BRUGNATELLI; CADET
DE GASSICOURT; PIERQUIN; VAN
MONS)

R. Mercurial Mucilage, three parts.
Extract of Hemlock, one part.
Liquorice Powder, a sufficient quantity

to make a pilular mass. (VAN MONS)

R. Mercury, one dram.
Gum Arabic, six drams.

Triturate, with a little water, to perfect
extinction, and add

Extract of Hemlock, one dram.

Make pills of two grains each. (GALL.;
FERR.; BRUGNATELLI; CADET DE GAS-
SICOURT; PIERQUIN)

C.

*Mercury divided by means of Saccharine
Substances.*

SACCHARATED OR SACCHARINE MERCURY.

*Æthiops saccharatus, Hydrargyrum saccha-
ratum, Mercurius saccharatus, Oxydum
Hydrargyri saccharatum, Saccharum
Mercurii vermifugum.* (HISP.; FERR.;
FULD.; HANN.; LIPP.; SARD.; HERBIP.;
CADET DE GASSICOURT; SPIELMANN;
VAN MONS)

R. Mercury, one part.
White Sugar, two parts.

Triturate together to the perfect extinc-
tion of the metal. (HANN.; SARD.; SPI-
ELMANN; VAN MONS)

FULD.; LIPP.; and CADET DE GASSI-
COURT prescribe yellow sugar-candy.

R. Mercury, one part.
White Sugar, two parts.
Water, a sufficient quantity

to make a soft paste, to be triturated to the
complete extinction of the metal. (HISP.)

R. Mercury, } of each,
White Sugar, } half an ounce.
Essential Oil of Tansey, }
twenty-six drops.

Triturate. (HERBIP.)

R. Mercury, } of each,
White Sugar, } half an ounce.
Supertartrate of Potass, } two scruples.

Triturate. (FERR.)

This preparation is recommended for
children and delicate subjects affected with
syphilitic affections.—It is given in choco-
late or coffee.

MERCURIAL HONEY.

Mel hydrargyrum. (PH. SYPH.; NIEMANN; SWEDIAUR)

R. Mercury, } of each,
Despumated Honey, } one ounce.
Triturate to extinction.

ANDRY'S ANODYNE POTION. (CADET DE GASSICOURT)

R. Orange-flower Water, two ounces.
Rose Water, } of each,
Syrup of Poppies, } one ounce.
Mercury, six drams.

Extinguish the mercury in the syrup, and then add the water.

It is recommended in cardialgia, gastrodynia, and spasmodic emesis.—Dose, a spoonful, morning and evening.

MERCURIAL BOLUSES.

Boli cærulei. (AUGUSTIN; ELLIS; SAUNDERS)

R. Mercury, ten grains.
Conserve of Roses, one scruple.
Triturate to extinction.

GLYCYRRHIZED MERCURIAL BOLUSES.
(PH. SYPH.; AUGUSTIN)

R. Mercury, ten grains.
Soft Extract of Liquorice, one scruple.
Triturate.

PILLS OF MERCURY.

Pilule Hydrargyri. (LOND.; EDINB.; DUBL.; AMER.; GENEV.; SUEC.; BRUGNATELLI; CADET DE GASSICOURT; COXE; ELLIS; NIEMANN; RATIER; VAN MONS)

R. Purified Mercury, two drams.
Confection of Red Roses, three drams.
Liquorice Root, powdered, one dram.

Triturate the mercury and the confection together till the former disappears, then add the powder, and mix. (LOND.; DUBL.)

AMER.; BRUGNATELLI; COXE; and NIEMANN give the same formula.

R. Purified Mercury, } of each,
Conserve of Red } one
Roses, } ounce.
Starch, two ounces.

Triturate the mercury with the conserve till the globules have entirely disappeared, adding, if necessary, a

VOL. II.

little mucilage of gum arabic; then add the starch, and with a little water beat the whole into a mass, to be directly divided into four hundred and eighty equal-sized pills. (EDINB.)

The same formula is given by BRUGNATELLI; CADET DE GASSICOURT; ELLIS; and VAN MONS.

R. Mercury, } of each,
Conserve of Dog Roses, } one dram.
Triturate to extinction, and make, with liquorice powder, thirty-six equal pills. (RATIER)

GENEV. prescribes one part of mercury, and two parts of conserve of roses.

R. Mercury, } of each,
Conserve of Roses, } two drams.
White Sugar, one dram.

After the extinction of the metal, add

Starch, } of each,
Liquorice Powder, } half a dram.
Water, a small quantity.

Make two hundred and eighty-eight pills. (SUEC.)

MELLATED PILLS OF MERCURY. (HISP.)

R. Mercury, } of each, one ounce.
Honey, }
After the extinction of the metal, add Crumb of New Bread, two ounces, and make a pilular mass.

R. Mercury, two ounces.
Honey, two ounces and a half.
Triturate to extinction; add Liquorice Powder, one ounce and a half, and make a pilular mass.

GLYCYRRHIZED PILLS OF MERCURY.
(LUSIT.; PH. SYPH.)

R. Mercury, } of each, one ounce.
Honey, }
Extract of Liquorice, two ounces.
Make a pilular mass. (PH. SYPH.)

R. Mercury, } of each,
Extract of Liquorice, } eight parts.
Powdered Liquorice, one part.

Make a pilular mass. (PH. SYPH.)

LUSIT. prescribes two of mercury, two of extract, and one of powder.

VERMIFUGE POWDER.

Pulvis anthelminticus. (HERBIP.)

R. Root of Jalap, half an ounce.
Valerian, } of each,
Male Fern, } two
Saccharated Mercury, } drams.

D

Dried Sulphate of Iron, two scruples.
Oleosaccharum of Tansey,
half an ounce.

Make a fine powder.

D.

Mercury divided by means of earthy Substances.

MERCURY WITH CHALK.

Hydrargyrum cum Creta, Mercurius seu Ethiops alcalisatus. (LOND.; DUBL.; FULD.; HERBIP.; WIRTEM.; PIDERIT)

R. Purified Mercury, by weight,
three ounces.
Prepared Chalk, five ounces.

Triturate till globules are no longer to be seen. (LOND.)

R. Mercury, } of each, two parts.
Manna, }
Prepared Chalk, one part.

Triturate the mercury with the manna in an earthen mortar, adding so much water as to give to the mixture the consistence of syrup; continue the trituration till the globules have entirely disappeared: then add, still triturating, one eighth of the chalk, and when it is well mixed, add sixteen parts of hot water, and agitate the mixture. Set aside the mixture for some time, and then decant the liquid. Wash the powder a second and a third time, in order to remove the whole of the manna: finally, add the remainder of the chalk to the moist powder, and dry in bibulous paper. (DUBL.)

FULD.; HERBIP.; WIRTEM.; and PIDERIT give the same process as LOND., but require prepared crab's eyes instead of chalk.—WIRTEM. prescribes the same proportions.—FULD.; HERBIP.; and PIDERIT—one part of mercury, and two parts of crab's eyes.

MERCURY WITH MAGNESIA.

Hydrargyrum cum Magnesia, Mercurius alcalisatus. (DUBL.; LIPP.)

DUBL. directs this to be prepared in the same manner as the mercury with chalk, employing carbonate of magnesia instead of chalk.

R. Carbonate of Magnesia, one ounce.
Mercury, half an ounce.

Triturate to the perfect extinction of the metal. (LIPP.)

This preparation and the preceding are the best that can be employed, when metallic mercury is required to be administered.

MERCURIAL BOLUSES. (SAUNDERS)

R. Alkalized Mercury, half a dram.
Conserve of Roses, one scruple.
Simple Syrup, a sufficient quantity.
Mix.

DETERGENT APPLICATION.

Topique deterensif. (BORIES)

R. Mercury, } of each,
Prepared Chalk, } half an ounce.
After the extinction of the metal, add
Honey of Roses, two ounces.

FOUQUET employed this preparation in the dressing of syphilitic ulcers.

RESOLVENT LIQUOR. (SAINTE-MARIE)

R. Mercury, half a dram.
Quick Lime, two drams.

Triturate to perfect extinction, and put the mixture into a bottle full of water holding in solution

Soap, four drams.

This liquor is reputed a powerful resolvent in cold and indolent tumours, and particularly in engorgements of the thyroid gland.—It is to be warmed, and compresses are to be saturated with it.

E.

Mercury divided by means of fat and resinous Bodies.

INJECTION OF MERCURY AND OF TURPENTINE.

Injectio hydrargyro-terebinthinata. (VAN MONS)

R. Mercury, } of each, two drams.
Turpentine, }

Triturate for an hour, and add

Lard, half a dram.

Then make an emulsion with

Mucilage of Gum Arabic, two drams.
Water, three ounces.

It is recommended in obstinate discharges of the urethra.—Two drams to be injected thrice a day.

PILLS OF MERCURY AND OF TURPENTINE. (PH. SYPH.; VAN MONS)

R. Mercury, four parts.
Turpentine, one part.

After the extinction of the metal, add
Thick Mucilage of Gum Arabic,
two parts.

Powdered Biscuit, a sufficient quantity,
and divide the mass into pills of five grains
each. (VAN MONS)

R. Mercury, one ounce.
Venice Turpentine,
one dram and a half.

Extinguish the metal, adding, if required,
a few drops of spirit of turpentine, and with

Liquorice Powder, a sufficient quantity,
make a mass to be divided into eighty pills.
(PH. SYPH.)

VAN MONS prescribes also two drams
each, of mercury and of turpentine, and
after extinction, a sufficient quantity of a
mixture of an equal part each of liquorice
and marshmallows root.

MERCURIAL PILLS. (FERR.; PIERQUIN)

R. Mercury, one ounce.
Turpentine, a sufficient quantity

to extinguish the metal; add
Powdered Amber, } of each,
Liquorice, } two drams.
Syrup of Marshmallows,
a sufficient quantity,

and make pills of four grains each. (PIER-
QUIN)

R. Mercury, one ounce.
Balsam of Mecca, one ounce and a half.

Triturate to the extinction of the metal,
and incorporate

Mass of Opium Pills, one dram.
Make a pilular mass. (FERR.)

Dose, from eight to twelve grains.

MERCURIAL OINTMENT.

Unguentum Hydrargyri seu mercuriale seu Neapolitanum. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; CADET DE GASSICOURT; COXE; PIDERIT; RATIER; SPIELMANN; SWEDIAUR; PH. SYPH.; VAN MONS)

The extinguishing of mercury being a long and tedious operation, various means have been proposed to render it easier: these may be reduced to seven, namely:

1. The addition of turpentine.

2. The addition of a little old mercurial ointment.

3. The addition of a little nitrous ointment.

4. *DUFILHO'S process.* Put the mercury into a phial having a convex and pointed bottom; fill this to one half with pure water, and shake it for some time, keep the mouth of the phial covered with the thumb; then, having let the globules deposit themselves, decant the water, and pour the mercury upon the fat; twenty minutes are then sufficient to extinguish it.—BOMPOIS had recourse, according to FEE, to a similar process; it is this: put into a quart bottle eight ounces of mercury, and six ounces of lard liquified with two ounces of olive oil; to stop the bottle, and agitate the mixture till it acquires the consistence of treacle; to pour it then into a mortar, and to triturate it strongly for half an hour.

5. *DUMESNIL'S process.* Put into a mortar four ounces of fat, one pound of mercury, and one ounce of oil of almonds; a quarter of an hour's agitation is sufficient to extinguish the metal entirely.

6. *PLANCHE'S process.* Triturate mercury with oil of eggs, and then add fat as usual.

7. *HERNANDES' process.* Warm the mortar so as to melt the fat; then add the mercury, and triturate till perfectly cold.—HASS. prescribes a similar method: melt the fat, then remove it from the fire; add the mercury when it begins to cool, and triturate.

In a medicinal point of view, it matters very little which of these processes be followed. But it should be observed that old ointment, turpentine, and nitrous ointment give a disagreeable smell to the ointment, and that into which turpentine enters, does not always form a proper application. It should also be remarked that rancid ointment is liable to occasion erysipelatous inflammation of the skin, which is sometimes very troublesome and disagreeable; but the general effects of the preparation are not diminished by that quality, for it has been remarked that the older the ointment the more efficacious it is.

But what chiefly interests the physician is the knowledge of the respective proportions of the mercury and of its excipient. But these proportions vary almost to infinity. Many pharmacopœias give only one, while others prescribe two, and even three.

The researches made by modern chemists, as to the most efficacious means for attenuating mercury and incorporating it with fat, having rendered useless all that has been formerly done on this subject, we shall pass

in silence the operatory processes indicated in the various pharmacopœias, and confine ourselves to mentioning the composition of the ointment or ointments in question, required by each. In doing this we shall divide them according to the proportion of the mercury to the fat body serving as its excipient. Of these preparations some are,

1. In the proportion of one to one.

One ounce each of mercury and of prepared lard. (DUBL.;* GALL.; FORM. HÔP. FR.; GENEV.; HASS.; BERGIUS; BRUGNATELLI; CADET DE GASSICOURT; RATIER; SPIELMANN)

Two ounces of mercury, and one ounce each, of lard and of goat's suet. (FRNN.; SUEC.)

Two pounds of mercury, twenty-three ounces of lard, and one ounce of mutton suet. (*Ung. Hydr. fortius*, LOND.*)—one pound of mercury, nine ounces of lard, and three of suet. (FERR.)

Four ounces of mercury, three and a half of lard, and half an ounce of mercurial ointment. (SWEDIAUR)

Two ounces of mercury, half an ounce of mutton suet, and one ounce and a half of cocoa nut oil. (PH. SYPH.)

This ointment is stronger than the preceding, as also is that prescribed by VAN MONS:—Mercury, sixteen parts, mercurial ointment, one part. †

The following, on the contrary, are weaker:

Mercury, one part, turpentine, an indeterminate but sufficient quantity for the extinction of the metal, lard one part. (SARD.)

Mercury, three parts, suet, one part, lard, three parts. (AMER.; COXE)

Mercury, one ounce, fresh oil of eggs, twenty drops, cocoa nut oil liquified with a little oil of almonds, one ounce. (HENRI; CADET DE GASSICOURT)

This process gives an ointment of a very agreeable odour, and which, although of a firm inconsistency, is of as easy application. The mercury and the oil of eggs are put into a small deep stone mortar and triturated together for a quarter of an hour; a porcelaine mortar, with its pestle, is also to be heated, and the cocoa-nut oil to be put into it as soon as it is melted, the mercury divided by the oil of eggs is to be added, and the mixture triturated for half

* The mode of preparation prescribed by LOND.; DUBL.; and EDINB. is simply trituration to the perfect extinction of the metal.—EDIT.

† Here the author has obviously made some mistake.—EDIT.

an hour, keeping the mortar tolerably warm that the mixture may preserve a certain liquidity: it is then to be suffered to cool gradually, and the trituration to be continued for a quarter of an hour longer.

2. In the proportion of one to two.

One part of mercury and two parts of lard. (*Ung. Hyd. mitius*, DUBL.*; HISP.; ROSS.; SAX.; SWEDIAUR)

One pound of mercury, eight ounces of mutton suet, sixteen ounces of lard, and one dram of oil of lemon. (HANN.; POL.; BORUS.)—one part each, of mercury, of mutton suet, and of lard. (EDINB.)*

EDINB. also prescribes two other ointments into which two parts and three parts of mercury enter.—Twelve parts of mercury, eight of mutton suet, and sixteen of lard. (OLDEN; CAST. BORUS.)—half an ounce of mercury, two drams of mutton suet, and two ounces of lard. (FULD.)—three parts of mercury, two of mutton suet, and four of lard. (BAVAR.)—half a pound each, of mercury and of suet, and one pound and a half of lard. (AMST.)

Four ounces of mercury, one ounce and a half of turpentine, ten ounces of lard, and one dram of essential oil of lavender. (BRUNS.; WIRTEM.)

3. In the proportion of two to five.

Four parts of mercury, a sufficient quantity of turpentine for its extinction, and ten parts of lard. (SAX.)

4. In the proportion of one to three.

Two ounces of mercury, five ounces of lard, five drams of turpentine, and half a dram of olive oil. (PIDERIT)—one ounce of mercury, half an ounce of turpentine, and three ounces of lard. (HERBIP.)

5. In the proportion of one to four.

One ounce of mercury, one of suet, and three ounces of lard. (DAN.; LUSIT.)—three ounces of mercury, half an ounce of turpentine, and one pound of wax ointment. (AUSTR.)—two parts of mercury, one part of nitrous ointment, six parts of lard, and one part of oil. (BATAV.; BELG.)

6. In the proportion of one to five.

One pound of stronger mercurial ointment and two pounds of lard. (*Ung. Hydr. mitius*, LOND.)

7. In the proportion of two to nine.

Four ounces of mercury, two of turpentine, and sixteen of lard. (HANN.; LIPP.)

8. *In the proportion of one to six.*

One ounce of mercury, and six ounces of lard. (GENEV.; RATIER)—two parts of mercury, one part of turpentine, and twelve parts of lard. (VAN MONS)

9. *In the proportion of one to eight.*

One ounce of mercurial ointment and three ounces of lard. (GALL.; FORM. HÔP. FR.)

10. *In the proportion of one to ten.*

Three ounces of mercurial ointment and one pound of lard. (ANT.)

If to these numerous varieties, in use at this time, of one preparation, we add all those which are now thrown aside, it will be seen that we ought to be very circumspect in reading treatises, ancient or modern, on syphilitic diseases, when we would wish to draw correct deductions respecting the mercury exhibited as a curative means in these diseases. At the present day, when the ointment prepared with equal parts of mercury and of fat is employed, from half a dram to one dram is generally used for each rubbing-in, when this operation is performed every day; but when it is performed every other day, then from one dram to two drams may be used. It is very seldom that this quantity is exceeded, and almost always dangerous to do so.

MERCURIAL CERATE. (RATIER)

R. Mercurial Ointment, four ounces.
Simple Cerate, ten ounces.

Mix.—It is employed especially in the dressing of syphilitic ulcers.

NITROUS MERCURIAL OINTMENT. (VAN MONS)

R. Mercurial Ointment, one part.
Nitrous Ointment, two parts.

Melt the latter with a moderate heat and at the moment of its cooling, incorporate together.

DIGESTIVE MERCURIAL OINTMENT. (RATIER)

R. Mercurial Ointment, } of each,
Digestive Ointment, } four ounces.

Mix.—It is recommended to be applied to atonic syphilitic ulcers.

PLENK'S MERCURIAL BALSAM.

Balsamum mercuriale Plenkii. (SPIEL-MANN)

R. Mercury, two drams.
Turpentine, one dram.

Triturate to extinction, and add

Lard, six drams.
ARCEUS' Balsam, nine drams.
Proto-chloruret of Mercury, twenty-three grains.

Mix.—This was much extolled by PLENK as an application to syphilitic ulcers.

ANTI-ARTHRITIC OINTMENT. (VAN MONS)

R. Mercurial Ointment, one dram.
Marshmallows Ointment, three drams.

Mix.—The parts affected with gout are recommended to be rubbed in the evening with a dram of this ointment. It is also said to be serviceable in chronic hepatitis, in frictions to the region of the liver, after having added half a dram of camphor to it.

SANCHEZ' MERCURIAL OINTMENT. (SAINTE-MARIE)

R. Mercury, four ounces.
Honey, four drams.

Extinguish the metal, and add

Powdered Camphor, four drams.
Cocoa Nut Oil, eight ounces.

Triturate for seventy hours [by steam power?—EDIT.].

In frictions.—Dose, from two drams to one ounce.

ELLIS prescribes also an ointment prepared with one ounce of simple mercurial ointment and one dram of camphor.

PILLS OF MERCURIAL OINTMENT. (HENRI; RATIER)

R. Mercurial Ointment, one ounce.
Starch or Powdered Marshmallows Root, a sufficient quantity
to make a mass. (HENRI)

RATIER directs a mass to be made of two drams of the ointment and a sufficient quantity of liquorice powder, to be divided into one hundred and forty-four pills.

SEDILLOT'S PILLS. (BORIES; CADET DE GASSICOURT; PIERQUIN; RATIER)

R. Mercurial Ointment, three ounces.
Medicinal Soap, two ounces.
Starch or Liquorice Powder, one ounce.

Make pills of from four to six grains each.

LAGNEAU has modified this formula, thus:

R. Mercurial Ointment prepared with
Cocoa Nut Oil, half an ounce.
Sugar, in powder, two drams.
Mucilage of Gum Arabic, a sufficient quantity

to make a mass, to be divided into one hundred and forty-four pills; to the mass two or three drams of the soap may be added.

Great eulogiums have been lately be-

stowed on these and the preceding pills: this happens, in medicine, to every thing bearing the stamp of novelty. But they are disgusting preparations: they should be rejected, were it but for the variability of the mercurial ointment, for it is absurd, when so powerful a medicament as mercury is prescribed, not to apportion it with sufficient precision so as to be certain of the actual quantity of it taken by the patient. LAGNEAU'S pastilles or PLENK'S pills before-mentioned, are infinitely preferable both on account of cleanliness and of safety: if necessary, the second may be rendered aperient by the addition of soap.

OINTMENT AGAINST LICE.

Unguentum Pediculorum seu Sabadillæ cum Mercurio seu de Staphisagria. (BRUNS.; HANN.; OLDEN.; PALAT.; SARD.; WIRTEM.; SPIELMANN)

R. Mercury, one ounce and a half.
Lard, one pound and a half.

After the extinction of the metal, add
Powdered Stavesacre Seeds,
four ounces.

Mix. (SARD.)

R. Mercury, two ounces.
Turpentine, one ounce.

Triturate, and add
Lard, three pounds.
Oil of Bay, half a pound.
Powdered Sabadilla, two ounces.
—— Root of Black Hellebore,
one ounce.

Mix. (OLDEN.)

R. Mercurial Ointment, eighteen ounces.
Powdered Sabadilla, } of each,
—— Stavesacre, } two ounces.
—— White Hellebore, one ounce.
Oil of Bay, nine ounces.
—— Lavender, one ounce.

Mix. (SPIELMANN)

R. Venice Soap, three ounces.

Dissolve it in water, and boil the solution with

Lard, three pounds
to the consumption of the humidity, and add

Oil of Bay, six ounces.
Powdered Stavesacre Seeds,
two ounces.
—— Root of White Hellebore,
one ounce.

Mercury, triturated to extinction
with one ounce of Turpentine,
two ounces.

Mix. (BRUNS.; HANN.; PALAT.; WIRTEM.)

JUCH'S OINTMENT OR PLASTER. (VAN MONS)

R. Mercury, three parts.
Turpentine, one part.
Triturate to extinction, and add gradually,
Mutton Suet, one part.
and then a mixture of
Turpentine, one part.
Spermaceti, one part and a half.
Yellow Wax, five parts.
melted together and half cold.

MERCURIAL PLASTER.

Emplastrum Hydrargyri seu ex Hydrargyro seu hydrargyricum seu mercuriale seu mercuriale resolvens.

1. *Without Lead.* (BAVAR.; FULD.; LIPP.; LUSIT.; SAX.)

R. Mercury, three parts.
Turpentine, one part.

Having triturated to extinction, add, triturating till perfectly cold, a liquified and half-cold mixture of

Mutton Suet, two parts and a half.
Yellow Wax, five parts and a half.

Preserve. (BAVAR.)

R. Mercury, extinguished with one
third of Turpentine, six parts.

Triturate it with a melted and nearly cold mixture of

Yellow Wax, ten parts.
Spermaceti, three parts.
Mutton Suet, two parts.

Mix. (SAX.)

R. Wax Plaster, one pound.
Mercury extinguished in one ounce
of Turpentine, three ounces.

Triturate together. (LIPP.)

FULD.; and LUSIT. prescribe one part of mercury extinguished with turpentine, and sixteen parts of hemlock or other plaster.

2. *With Lead.* (LOND.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BELG.; DAN.; MIL. COPEN.; HISP.; FENN.; FERR.; HASS.; PALAT.; ROSS.; SUEC.; WIRTEM.; HERBIP.; BRERA; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Purified Mercury, three ounces.
Sulphuretted Oil,
one fluid dram.

Lead Plaster, one pound

Triturate the two former till the mercury is extinguished; then add

gradually the melted plaster, and mix. (LOND.; BRERA; COXE)

R. Olive Oil, }
Resin, } of each, one part.
Mercury, three parts.
Plaster of Semi-vitreous
Oxide of Lead, six parts.

Extinguish the mercury in the oil and resin, previously melted together and half cooled; then add the melted plaster, and mix intimately. (EDINB.; AMER.; BATAV.; COXE)

R. Lead Plaster, one pound.
Melt slowly, and add
Mercurial Ointment, three ounces.
Mix. (FERR.)

R. Mercury, four ounces.
Venice Turpentine, half an ounce.
Lead Plaster, melted, twelve ounces.

Extinguish the mercury in the turpentine, add the plaster slowly and mix. (MIL. COPEN.)

HASS.; and PIDERIT prescribe three ounces each, of mercury and of turpentine, and one pound of plaster.—ROSS.—three ounces of mercury, one ounce and a half of turpentine, and one pound of plaster.—SWEDIAUR—eight ounces of mercury, three drams of turpentine, and one pound of plaster.—DAN.; and HERBIP.—one ounce and a half of mercury extinguished in turpentine, and one pound of plaster.

R. Mercury, three ounces.
Lard, one ounce and a half.
Mutton Suet, half an ounce.

Triturate till the metal is extinguished, and add slowly, a melted mixture of

Lead Plaster, one pound.
Yellow Wax, three ounces.

Mix. (AMST.)

R. Mercury, four ounces.
Mercurial Ointment, } of each,
Venice Turpentine, } one ounce.

Extinguish the metal and add slowly, a melted mixture of

Compound Galbanum Plaster, one pound.
Yellow Wax, one ounce.

Mix in a warm mortar. (ANT.; BELG.)

R. Mercury, one ounce.
Turpentine, three drams.

Having extinguished the metal, add,

Lead Plaster, two ounces.
Yellow Wax, half an ounce,

melted together. (FENN.; SUEC.)

HANN.; OLDEN.; POL.; and BORUS. prescribe eight ounces of mercury, four of

turpentine, two pounds of plaster, and half a pound of wax.

R. Mercury, twelve parts.
Venice Turpentine, three parts.
Liquid Storax, two parts.
Oil of Bay, one part and a half.

Extinguish the metal in the turpentine, add the storax, and then the oil: then put the mortar into warm water, and add gradually,

Lead Plaster, forty-eight parts.
Yellow Wax, three parts,

melted and mixed together. (VAN MONS)

VAN MONS also prescribes twenty ounces of mercury, five of turpentine, three of storax, two and a half of oil of bay, seventy of plaster, and one and a half of wax.

R. Mercury, one pound.
Turpentine, four ounces.
Vigo's Plaster, melted,
six pounds eight ounces

Mix. (HISP.)

R. Mercury, three ounces.
Turpentine, one ounce and a half.
Diabotanium Plaster, one pound.

Mix. (HISP.)

This stimulant plaster is reputed resolvent and discutient. It is applied to indolent tumours of the lymphatic glands. The numerous variations presented by it, in the proportional quantity of mercury which it contains, are of less importance than those which were noticed in the ointment; but which, however, cannot be without some influence in its effects. The last formula but one gives the celebrated *Emplastrum de Vigo cum Mercurio*, which only differs from the others in being more complicated, and consequently less active, the mercury in it being, if we may use the expression, choked by an absurd assemblage of dissimilar drugs.

MERCURIAL BOUGIES. (VAN MONS)

R. Mercurial Plaster, three parts.
Yellow Wax, one part.

Melt, mix, and make bougies.

AMMONIACAL MERCURIAL PLASTER.

Emplastrum Ammoniaci seu de Ammoniaco cum Mercurio.

1. *Without Lead.* (LOND.; DUBL.; BRUGNATELLI; GADET DE GASSICOURT; COXE; SPIELMANN; VAN MONS)

Emplastrum Ammoniaci cum Hydrargyro.

R. Gum Ammoniac, one pound.
Purified Mercury, three ounces.

Sulphuretted Oil, one fluid dram.

Extinguish the mercury in the oil, and add the melted plaster. (LOND.; BRUGNATELLI; CADET DE GASSICOURT; COXE)

R. Mercury, three ounces.
Turpentine, two drams.

After the extinction of the metal, add slowly

Gum Ammoniac, melted,
one pound.

Mix well. (DUBL.)

VAN MONS prescribes nine parts of mercury, one part of oil, and thirty-six parts of gum.

R. Mercury, three ounces.
Sulphuretted Oil, one dram.

After the extinction of the metal, add

Gum Ammoniac, melted, twelve ounces.
Turpentine, three ounces.

Mix. (SPIELMANN)

2. With Lead. (BRUNS.; GALL.; FORM. HOP. FR.; SARD; WIRTEM.)

R. Mercury eight ounces.
Venice Turpentine, one ounce.
Liquid Storax, half an ounce.

Triturate and add

Lead Plaster, one pound,
previously melted and mixed with
Gum Ammoniac, half a pound.

Mix. (BRUNS.; SARD.; WIRTEM.)

R. Mercury, three hundred and eighty parts.
Turpentine, sixty-four parts.
Liquid Storax, one hundred and ninety-two parts.

Triturate in an iron mortar, to the extinction of the metal; then

Lead Plaster, two hundred and fifty parts.
Yellow Wax, } of each,
Resin, } eighty-four parts.

Melt together, and when the mixture is half-cold, add to it,

Powdered Gum Ammoniac, } of each,
Bdellium, } twenty
Olibanum, } parts,
Myrrh, }

then

Saffron, twelve parts,

and finally the extinguished mercury. Mix intimately, and before the mass is perfectly cold, add

Oil of Lavender, eight parts.

Preserve. (GALL.)

R. Mercury, ninety-five parts.
Liquid Storax, forty-eight parts.

Triturate to extinction. Melt separately and together,

Lead Plaster, three hundred and twelve parts.
Yellow Wax, sixteen parts.

Mix, and add,

Turpentine, } of each, sixteen parts.
Resin, }
Gum Ammoniac, } of each, five parts.
Olibanum, }
Myrrh, }

Then incorporate

Saffron, in powder, three parts,
and finally the mercurial mass. (FORM. HOP. FR.)

The observations made on the mercurial plaster, apply equally to this.

PLASTER AGAINST BUBOES.

Emplastrum ad Bubones venereos.

R. Mercury, two drams.
Sublimed Sulphur, two ounces.

Extinguish the metal, and add,

Super-acetate of Lead, three drams.

Mix, and incorporate with a melted and half cold mixture of

Plaster of Gum Ammoniac, }
Melilot, } of each,
Cumin, } half
Galbanum, } an ounce.
Sagapenum, }

PLASTER OF THE FOUR RESOLVENTS.
(GALL.; BRERA; CADET DE GASSICOURT)

R. Mercurial Plaster, }
Compound Galbanum } of each,
Plaster, } an equal part.
Plaster of Hemlock, }
Soap, }

Melt, and mix intimately. (GALL.; CADET DE GASSICOURT)

BRERA, who omits the plaster of soap, gives this formula under the name of *Emplastrum resolvens compositum*.

It is stimulant, resolvent, and maturative.

PLASTER OF MERCURY WITH HEMLOCK.

Emplastrum hydrargyri cum Conio seu resolvens cum Mercurio. (MIL. COPEN.)

R. Mercurial Plaster, } of each,
Hemlock Plaster, } half an ounce.

Knead together.

It is recommended in periostosis and enlargements of the epididymis.

D.

By Iron.

GRAPHITIC ÆTHIOPS.

Æthiops graphiticus, Graphites hydrargyratus. (NIEMANN; SAUNDERS; VAN MONS)

R. Mercury, one part.
Carburet of Iron, two parts.
Triturate to the perfect extinction of the metal.

This preparation has been celebrated in herpes.—Dose, ten grains.

ANTI-HERPETIC ELECTUARY.

Electuarium resolvens anti-herpeticum.
(BRERA)

R. Graphitic Æthiops, two drams.
Rhubarb, half a dram.
Extract of Cichory, half an ounce.
Despumated Honey, a sufficient quantity.

Mix.

E.

By Opium.

OPIATE MERCURIAL LINIMENT. (SAINTE-MARIE)

R. Mercurial Ointment (at equal parts), four scruples.
Oil of Almonds, two ounces.
SYDENHAM'S Laudanum, one dram.

Mix.

It is recommended in inflammatory ulcerations of the gland, in baths, or in injections, according to the state of the prepuce, twice or thrice a day.

OPIATE MERCURIAL OINTMENT. (BRERA; RATIER)

R. Mercurial Ointment, ten grains.
Opium, one grain.

Mix. (BRERA)

R. Mercurial Ointment, } of each,
Opium Cerate, } an equal part.

Mix. (RATIER)

CHAUSSIER frequently prescribes two or three frictions daily, to the abdomen, with a dram of the last ointment, towards the end of peritonitis, as a proper means of promoting the termination of the inflammation.

R. Mercurial Ointment (at equal parts), four ounces.
Powdered Opium, two ounces.

Mix.—To be applied to painful syphilitic ulcers. (RATIER)

F.

By Purgative Substances.

COMPOUND MERCURIAL SUGAR.

Saccharum mercuriale compositum.
(BRUNS.)

R. Mercury, half an ounce.
White Sugar, two ounces.
After the perfect extinction of the metal, add,
Powdered Jalap, half an ounce.
Mix.

PURGATIVE PILLS. (ELLIS)

R. Mass of Mercurial Pills, one scruple.
Powdered Aloes, twenty-five grains.
Make fifteen pills.—Dose, one every two hours.

BELLOSTE'S PILLS.

R. Mercury, ninety-six parts.
Honey, two hundred and fifty parts.
Triturate to the extinction of the metal, and add
Powdered White } of each,
Agaric, } sixteen parts.
Aloes, }
Scammony, sixty-four parts.
White Pepper, twelve parts.

Roll the pills in a powder made with equal parts of mechoacan and of jalap exhausted by alcohol. (HENRI; FEE)

This is the true formula of BELLOSTE'S pills, celebrated at the same time as anti-syphilitic and as purgative. They have been modified in a great many different ways, from which an infinity of pills have been formed called *Neapolitan, mercurial*, and *anti-venereal*; these we shall bring together in this article, because they all have mercury combined with purgative substances, and only differ by the choice or the dose of the latter.

R. Mercury, four parts.
Honey, forty parts.

Triturate to extinction, and add

Socotrine Aloes, } of each,
White Agaric, } eight parts.
Mace, } of each, one part.
Cinnamon, }

Make pills of three grains each. (FORM. HÔP. FR.)

This formula approaches that of BELLOSTE.

The following is a modification, by LEMERY, of BARBEROUSSE'S celebrated pills; from which it is known that BELLOSTE contrived his:

R. Mercury, six drams.

Turpentine, a sufficient quantity
to extinguish the metal. Add
Aloes, six drams.
White Agaric, half an ounce.
Rhubarb, one dram.

Make a pilular mass. (SPIELMANN)

R. Mercury, } of each,
Venice Turpentine, } one ounce.

Triturate to the extinction of the metal,
and add

Scammony, }
Rhubarb, } of each, one ounce.
Jalap, }
Despumated Honey,

a sufficient quantity
to make a pilular mass. (FERR.)

R. Mercury, } of each, one ounce.
Scammony, }
Supertartrate of Potass, } of each,
Resin of Jalap, } half
Rhubarb, } an ounce.
Simple Syrup, a sufficient quantity.

Extinguish the mercury with the super-
tartrate and a little syrup, add the rest, and
make pills of four grains each. (GENEV.)

The following also resemble BELLOSTE's
pills, and are called *Purgative Mercurial*
Pills:

R. Mercury, four parts.
Honey, a sufficient quantity

to extinguish the metal. Add

Powdered Cinnamon, } of each,
Mace, } one part.
Aloes, } of each,
Scammony, } eight parts.
Honey, a sufficient quantity.

Divide the mass into pills of four grains
each. (GALL; VAN MONS)

The *Neapolitan Pills*, of which the fol-
lowing is the formula, are also very similar:

R. Mercury, six drams.
Honey, a sufficient quantity

to extinguish the metal. Add

Powdered Aloes, four drams.
Rhubarb, three drams.
Scammony, two drams.
White Agaric, one dram.
Sassafras, } of each,
Cinnamon, } one scruple.
Mace, }

Mix. (CADET DE GASSICOURT; PIER-
QUIN)

BAUME' has corrected BELLOSTE's for-
mula in the following manner:

R. Mercury, } of each,
Supertartrate of Potass, } one ounce.

Triturate with a little honey to the ex-
tinction of the metal, and add

Powdered Scammony, } of each,
Jalap, } one ounce.

Honey, a sufficient quantity

to make a pilular mass. (ANT.; SPIEL-
MANN)

RATIER prescribes three ounces of mer-
cury, ten grains of supertartrate of potass,
three drams of scammony, three drams of
jalap, and a sufficient quantity of syrup.—
VAN MONS—four parts of mercury, two of
cream of tartar, a sufficient quantity of
syrup to extinguish the metal, one part of
gum arabic, four parts of jalap, four of
scammony, and a sufficient quantity of sy-
rup to form a pilular mass.

R. Mercury, } of each, one ounce.
Scammony, }
Sugar, two drams.

Triturate to the extinction of the metal,
moistening with a little white wine, and
add

Powdered Jalap, one ounce.

Mix. (CADET DE GASSICOURT)

R. Mercury, two drams.
Venice Turpentine, a sufficient quantity
to extinguish the metal. Add

Scammony, } of each, two drams.
Jalap, }
Gamboge, }
Syrup of Buckthorn, a sufficient quantity.

Divide the mass into pills of four grains
each. (SAUNDERS)

R. Mercury, two ounces.
Manna, one ounce.
Jalap, four ounces.
Turpentine, } of each,
Sublimed Sulphur, } two drams.
Gamboge, }
Syrup of Buckthorn, a sufficient quantity.

Divide the mass into pills of four grains
each. (CADET DE GASSICOURT)

R. Mercury, half an ounce.
Mucilage of Gum Arabic, one ounce.

Triturate to perfect extinction, and add

Extract of Guaiac Wood, } of each,
Aloes, } half
Jalap, } an ounce.
Syrup of Rhubarb, a sufficient quantity.

Mix. (SARD.)

R. Mercury, one ounce.
White Sugar, half an ounce.

Triturate to the perfect extinction of the
metal, sprinkling with a sufficient quantity
of oxymel. Then add gradually

Scammony, } of each, one ounce.
Jalap, }
Oxymel, a sufficient quantity.

And add towards the end

Nitrate of Potass,	one dram.
Camphor,	six grains.

Make pills of ten grains each. (PIERQUIN)

It will be readily seen that the dose ought not to be the same for all these pills. It must be varied according to the strength of the purgatives entering into their composition.

That of LEMERY's pills is one dram, and that of BAUME' half a dram.

PILLS AGAINST TËNIA. (BORIES.)

R. Mercury,	one ounce and a half.
Lard,	one ounce.

After the extinction of the metal, add

Powdered Aloes, a sufficient quantity to make a mass to be divided into pills of four grains each.

G.

By Cinchona.

HEISTER'S ANTHELMINTIC ELECTUARY. (BRUNS.; CADET DE GASSICOURT; SPIELMANN)

R. Mercury,	half an ounce.
Gum Arabic,	one ounce.
Water,	a sufficient quantity.

Triturate to extinction, and add

Cinchona, in fine powder, one ounce.
Syrup of Mint, a sufficient quantity to give to the mixture the consistence of a soft electuary.

The dose varies according to the age. A tea spoonful is usually given morning and evening.

H.

By Sulphur

MERCURY WITH SULPHUR; BLACK SULPHURET OF MERCURY.

Sulphuretum Hydrargyri nigrum seu simplex, Hydrargyrus sulphuratus niger seu cum Sulphure, Hydrargyrum sulphuratum nigrum, Mercurius cum Sulphure, Pulvis hypnoticus, Æthiops mineralis seu narcoticus.—Æthiops Mineral. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; HANN.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PARIS; SPIELMANN; VAN MONS)

R. Purified Mercury,	} of each, an equal weight.
Sublimed Sulphur,	

Triturate till the metal is extinguished. (LOND.; DUBL.; EDINB.)

EDINB. says that it may also be made with double the quantity of mercury.

AMER.; AMST.; ANT.; BATAV.; BELG.; DAN.; FERR.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; COXE; and SWEDIAUR also prescribe equal parts.

ANT. directs a little water to be added to facilitate the trituration.

DESTOUCHES has proposed to sprinkle the mixture during the trituration with hydrosulphate of potass; and GEIGER, with hydrosulphate of ammonia.

R. Mercury,	two parts.
Sublimed Sulphur,	one part.

Triturate till the metal is extinguished. (AUSTR.; HISP.; GALL.; ROSS.; PLANCHE; SPIELMANN)

There are other variations with respect to the proportions. Thus, BRUNS.; PALAT.; and WIRTEM. prescribe three parts of mercury and two of sulphur.—SARD.—two of mercury and one and a half of sulphur.

VAN MONS admits four sorts of æthiops, to which he gives particular denominations, namely,

1. *Æthiops mineral*, or *Sulphur triturated with Mercury*, which is prepared by triturating together, in a marble mortar with a large pestle, for a quarter of an hour, one part and a half of roll sulphur, and nineteen parts of mercury;
2. *Æthiops Mineral of Harris*, for which, having taken mercury and sulphur, of each an arbitrary quantity, the former is to be triturated with a twelfth of the latter, to perfect extinction, after which another twelfth is to be added, and then gradually the remainder of the sulphur;
3. *Oxydulated Æthiops Mineral*, obtained by triturating together, one part and a half of roll sulphur, nine and a half of mercury, and ten and a fourth of red oxide of mercury;
4. *Double Æthiops Mineral* or *Super-sulphuret of Mercury*, which requires nine parts of mercury to be triturated with one part and a half of roll sulphur.

Æthiops mineral called black sulphuret of mercury, is not a combination of mercury and of sulphur, as has been for a long time believed; but is formed of a simple mixture of the two substances. MARTIUS has prepared it by submitting a bottle containing a mixture of mercury and of sulphur to the motion of a saw-mill, making it follow the movements of the machine for fifteen days.

It is stimulant, anti-syphilitic, and anthelmintic. It is never administered alone.

ANTI-GONORRHOIC PILLS. (BORIES ; PIERQUIN)

- R. Black Sulphuret of Mercury, one ounce and a half.
 Rhubarb, } of each, two drams.
 Myrrh, }
 Compound Syrup of Cichory, a sufficient quantity.

Make pills of six grains each.

BARBEYRUE prescribed them to be taken in the morning twice a week for two or three months.

MERCURIAL VERMIFUGE POWDER. (GALL.; HAMB.)

- R. Black Sulphuret of Mercury, } of each, one dram.
 Jalap, }
 Root of Valerian, two drams.

Mix. (HAMB.)

- R. Black Sulphuret of Mercury, } of each,
 Cornachine Powder, } an equal part.
 Triturate together. (GALL.)

ANTI-STRUMIC PILLS. (GALL.)

- R. Black Sulphuret of Mercury, } of each,
 Scammony, } two ounces.
 Deutoxide of Antimony, } of each,
 Prepared Wood-lice, } three drams.
 Soap, }
 Compound Syrup of Smallage, a sufficient quantity.

Make pills of four grains each.

PURGATIVE AND ALTERNATIVE ELECTUARY. (BORIES)

- R. Black Sulphuret of Antimony, half an ounce.
 Confection of Senna, one ounce.
 Powdered Senna, } of each,
 Rhubarb, } two drams.
 Jalap, }
 Wood-lice, }
 Scammony, one dram.
 Sulphate of Magnesia, half an ounce.
 Syrup of Smallage, a sufficient quantity.

Mix.—It is recommended in abdominal obstructions, and in dropsy.—Dose, one dram in the morning, fasting.

ANTI-STRUMOUS POWDER. (SAINTE-MARIE)

- R. Black Sulphuret of Mercury, four drams.

Prepared Crab's Eyes, two ounces.
 Powdered Amber, one ounce.

Mix, and divide into sixty-four equal doses.—First, one is to be taken, and then two, daily.

VERMIFUGE POWDER.

Pulvis contra Vermes. (SPIELMANN)

- R. Black Sulphuret of Mercury, two drams.
 Corsican Moss, half an ounce.
 Essential Oil of Wormwood, half a scruple.

Mix.—FULLER prescribed this powder for children, and ordered as much as might be taken upon the point of a knife to be administered night and morning. This is a singular manner of indicating the dose, which should be such that the child may take from two to four grains at most of the sulphuret, at each dose.

VERMIFUGE PASTILLES.

Rotula contra Vermes. (FERR.; WIRTEM.)

- R. Black Sulphuret of Mercury, } of each,
 Seeds of Artemisia, } two scruples.
 Corsican Moss }
 Cinnamon, }
 White Sugar, boiled in Rose Water to the proper consistence, four ounces.

Make pastilles. (WIRTEM.)

- R. Chocolate, fifteen ounces.
 Black Sulphuret of Mercury, six drams.
 Seeds of Artemisia, one ounce.
 Make seventy-two pastilles. (FERR.)

VERMIFUGE ELECTUARY. (PIERQUIN)

- R. Black Sulphuret of Mercury, } of each,
 Seeds of St. John's-wort, } half an ounce.
 Rue, }
 Wormwood, }
 Tansey, }
 Syrup of Wormwood, a sufficient quantity.

Mix.—CHIRAC gave half a dram of it in the morning, and the same in the evening; afterwards giving a decoction of male fern.

ALTERNATIVE ELECTUARY. (SAUNDERS; SWEDIAUR)

- R. Black Sulphuret of Mercury, } of each,
 Guaiac Resin, } half an ounce.
 Pulp of Prunes, one ounce.
 Simple Syrup, a sufficient quantity.

Mix.—It is recommended in itch, and other cutaneous diseases.—Dose, one dram, twice a day.

DETERGENT OINTMENT. (HERBIP.)

R. Black Sulphuret of Mercury, one ounce.
Lard, three ounces.
Triturate.

SULPHURETTED MERCURIAL OINTMENT.

Unguentum e Sulphure hydrargyrato.
(VAN MONS)

R. Sublimed Sulphur, one part.
Mercurial Ointment, two parts.
Triturate for a long time, and add gradually
Lard, fifteen parts.
Mix.

VAN MONS also prescribes one of sulphur, and two each, of the ointment and of lard.

It is useful when rubbed upon the skin, in itch.

ANTI-PSORIC OINTMENT. (VAN MONS)

R. Black Sulphuret of Mercury, one ounce.
Soap, half an ounce.
Mix, and add
Ointment of Camphor, one ounce and a half.

Mix.—It is recommended in obstinate itch. Like the two preceding, it requires to be used with reserve and circumspection.

ANTI-PSORIC CERATE. (VAN MONS)

R. Black Sulphuret of Mercury, four parts.
Prepared Sulphuret of Lead, one part.
Mix together in a mortar, and add
Cerate, softened with heat, sixteen parts.
Triturate.

RESOLVENT PILLS. (SAINTE-MARIE)

R. Black Sulphuret of Mercury, one scruple.
Extract of Dandelion, three drams.
Gum Ammoniac, two drams.
Squill, in powder, half a dram.
Syrup, a sufficient quantity.
Make pills of three grains each.

FINKE extolled these in ascites occasioned by engorgements of the abdominal viscera.—Dose, five, thrice a day.

ANTI-HERPETIC PILLS. (BORIES)

R. Mercury, two ounces.
Sublimed Sulphur, } of each,
Sulphate of Potass, } half an ounce.
Camphor, half a dram.
Triturate to the extinction of the metal, and add
Guaiac Resin, one ounce.
Peruvian Balsam, two drams.
Syrup of Clove Pink, a sufficient quantity.

Make pills of four grains each.

They are recommended by BARTHEZ, as resolvent and diaphoretic, in herpetic and other affections of the skin.—Dose, three, in the morning, fasting.

I.

By Sulphur and Antimony.

MERCURY WITH SULPHUR AND SULPHURET OF ANTIMONY; SULPHURET OF MERCURY AND OF ANTIMONY.

Sulphuretum Hydrargyri et Stibii seu stibiolum, Hydrargyrum stibiato-sulphuratum, Æthiops antimonialis.—Antimonial Æthiops. (AMST.; AUST.; BATAV.; BAVAR.; BELG.; BRUNS.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; BORUS.; ROSS.; SAX.; SARD.; WIRTEM.; HERBIP.; CADET DE GASSICOURT; PIDERIT; PLANCHE; SAUNDERS; SPIELMANN; SWEDIAUR; VAN MONS)

R. Mercury, four ounces.
Sulphuret of Antimony, three ounces.
Sulphur, two ounces.

Triturate together to the perfect extinction of the metal. (AMST.; BATAV.; BELG.; FENN.; HASS.; CADET DE GASSICOURT; PIDERIT)

GENEV.; HASS.; LIPP.; and PIDERIT recommend a few drops of water to be added.

LIPP. prescribes one ounce of mercury, one of sulphuret of antimony, and half an ounce of sulphur.—HANN.; POL.; BORUS.; SAX.; and SAUNDERS—one ounce each, of mercury and of sulphur, and two ounces of sulphuret.—VAN MONS—nine parts and a half of mercury, five and a half of sulphuret, and one and a half of sulphur.

R. Mercury, } of each,
Sulphuret of Antimony, } an equal part.
Triturate together. (AUST.; BAVAR.; ROSS.)

BRUNS.; DAN.; OLDEN.; SARD.; WIRTEM.; PLANCHE; SPIELMANN; and SWEDIAUR prescribe two parts of sulphuret, and one part of mercury.

- R. Mercury, three parts and a half.
Proto-hydro-sulphate of Antimony, one part.

Triturate to extinction. (VAN MONS)

VAN MONS elsewhere prescribes two parts of mercury and three of sulphuretted hydro-sulphate of antimony.—LUSIT.—one and a half of mercury and one of sulphuretted hydro-sulphate.—FERR.—one and a half of mercury and one of proto-hydro-sulphate of antimony.

As with the preceding, this is but a simple mixture, the mercury being merely much subdivided.—Dose, from six to twenty grains.

POWDER OF SULPHURET OF MERCURY AND OF ANTIMONY.

Pulvis ex Æthiope antimoniali seu ex Hydrargyro stibiato-sulphurato. (CAST. BORUS.; WURCEB.; AUGUSTIN)

- R. Sulphuret of Mercury and of
Antimony, six grains.
Liquorice Root, ten grains.

Mix. (WURCEB.)

CAST. BORUS.; and AUGUSTIN prescribe five grains of æthiops and fifteen of liquorice to be taken every four hours.

ANTI-HERPETIC POWDER.

Pulvis ad Herpetem. (HAMB.)

- R. Sulphuret of Mercury and of } of each,
Antimony, half
Sulphuret of Mercury, } a dram.
Guaiaec Resin, }
Liquorice, } of each, one dram.
Magnesia, }

Mix intimately.

ALTERATIVE POWDER.

Pulvis purificans. (HUFELAND)

- R. Guaiaec Resin, half a dram.
Sulphuret of Mercury and of } of each,
Antimony, half
Subcarbonate of Magnesia, } scruple.
Oleosaccharum of Fennel, half a dram.

Mix.—Dose, for an adult, in twenty-four hours. One half of it to children of seven years, and one third to those of four years.

ALTERATIVE POWDER.

Pulvis Hydrargyri Stibiato-sulphurato cum Cicuta. (AUGUSTIN; VAN MONS)

- R. Sulphuret of Mercury and of } of each,
Antimony, an
Dried Fecula of Hemlock, equal
Subcarbonate of Magnesia part.
White Sugar, }

Triturate.—Dose, from four to sixteen grains. It is recommended by HUFELAND

in the chronic exanthemata of children. The use of it should be continued for some time.

ALTERATIVE PILLS. (PH. SYPH.)

- R. Mercury, four ounces.
Sulphuret of Antimony, three ounces.
Sulphur, two ounces.

Triturate to the perfect extinction of the metal, and add

Honey, a sufficient quantity.

Divide the mass into pills of five grains.

COMPOUND PILLS OF SULPHURET OF MERCURY AND OF ANTIMONY. (DAN.; HAMB.; LUSIT.; WIRTEM.; HERBIP.; AUGUSTIN; SPIELMANN; VAN MONS)

- R. Sulphuret of Mercury and of
Antimony, one ounce.
Powdered Extract of Guaiaec, half an ounce.

Honey, a sufficient quantity,

to make a pilular mass. (LUSIT.)

- R. Sulphuret of Mercury and of } of each,
Antimony, one
Guaiaec Resin, } dram.
Extract of Bitter-sweet, two drams.
Simple Syrup, a sufficient quantity.

Make pills of two grains each. (AUGUSTIN)

- R. Sulphuret of Mercury and of } of each,
Antimony, half an
Medicinal Regulus of Anti- } ounce.
mony, }
Guaiaec Resin, one ounce.
Extract of Sarsaparilla, }

Make a pilular mass. (WIRTEM.; SPIELMANN)

- R. Mercury, six drams.
Honey, half an ounce.

Triturate to the perfect extinction of the metal, and add

Sulphuretted Hydro-sulphate } of each,
of Antimony, half an
Guaiaec Resin, } ounce.
Mucilage of Gum Arabic, a sufficient quantity,

to make a homogeneous mass. (HERBIP.)

- R. Mercury, three parts.
Sulphuretted Hydro-sulphate } of each,
of Antimony, two
Honey, } parts.

After the extinction of the metal, add

Gum Arabic, one part.
Guaiaec Resin, two parts.
Simple Syrup, a sufficient quantity,

to make a pilular mass. (VAN MONS)

- R. Mercury, six drams.
Sulphuretted Hydro-sulphate of
Antimony, half an ounce.

Extinguish the metal, and add

Soap, } of each,
 Guaiac Resin, } half an ounce.
 Simple Syrup, a sufficient quantity.

to make a pilular mass. (DAN.)

R. Sulphuret of Mercury and of } of each,
 Antimony, } one
 Sulphuret of Mercury, } dram.
 Extract of Hemlock }
 one dram and a half.
 Powdered Hemlock, half a dram.

Make pills of two grains each. (HAMB.)

These and the preceding pills are celebrated as alterative and resolvent. They are chiefly recommended in itch, hydropic affections, and in cases of ill-conditioned ulcers. —The dose of it may be carried to half a dram.

BALDINGER'S PILLS. (SAINTE-MARIE)

R. Mercury, one dram.
 Sulphuret of Antimony, four scruples.
 Sublimed Sulphur, two scruples.
 Opium, one scruple.

Triturate intimately with

Syrup of Marshmallows,
 a sufficient quantity.

Make a hundred and fifty pills.

HUXHAM recommended these pills, since modified by BALDINGER, in cases of painful affections of the bones and of leucorrhœa considered syphilitic; in tinea, scrofula, and itch.—Dose, five or six, daily

DIAPHORETIC PILLS. (PIERQUIN)

R. Sulphuret of Mercury } of each,
 and of Antimony, } half an ounce.
 Guaiac Resin, }
 Soap, } of each,
 Extract of Bitter-sweet, } one ounce.
 Conserve of Fumitory, } of each,
 Compound Syrup of } a sufficient
 Smalage, } quantity.

Mix.

ANTI-SYPHILITIC BOLUSES.

Boli in Necrosi syphilitica. (BRERA)

R. Nitric Oxide of Mercury, one grain.
 Sulphuret of Mercury and of Antimony,
 forty-eight grains.
 Rob of Elder, } of each, a
 Liquorice Powder, } sufficient quantity.

Make eight boluses.—Dose, one, morning and evening.

ALTERATIVE ELECTUARY. (SAINTE-MARIE)

R. Mercury, } of each,
 Sulphate of Antimony, } half an ounce.
 Guaiac Resin, one ounce.

After the extinction of the metal, add a sufficient quantity of rob of elder diluted with one third of water to make an electuary.

This preparation was vaunted by HERRENSCHVAND in syphilitic cutaneous diseases. —Dose, one dram, twice a day, beginning with a scruple, and augmenting gradually.

ANTI-HERPETIC ELECTUARY. (BORIES)

R. Mercury, one scruple.
 Sublimed Sulphur, } of each,
 Regulus of Antimony, } half an ounce.
 Extract of Bitter-sweet, } of each,
 Meadow Anemone, } two
 Powdered Wild Violet, } drams.
 Camphor, one dram.
 Syrup of Fumitory,
 a sufficient quantity.

Mix.—It is extolled in obstinate tetter, particularly those called scrofulous and syphilitic.—Dose, one dram, morning and evening.

OXIDE OF MERCURY.

I. *Mixed with Metallic Mercury.*

GREY OXIDE OF MERCURY.

Mercurius niger Moschati, Pulvis cinereus Hydrargyri, Oxydum Hgdrargyri nigrum seu Cinereum, Oxydum hydrargyrosum, Oxydum hydrargyricum præcipitatum, Hydrargyrum oxydulatum nigrum. (LOND.; EDINB.; DUBL.; AMER.; AUSTR.; BATAV.; FENN.; FERR.; BORUS.; SEEC.; BORIES; COXE; PALAT.; SWEDIAUR)

Hydrargyrum Oxydum cinereum.

R. Submuriate (Proto-chloruret) of Mercury, one ounce.
 Lime Water, one gallon.

Boil the submuriate in the lime water, continually stirring, till the grey oxide subsides, wash this with distilled water, and dry it. (LOND.; and COXE)

R. Submuriate of Mercury, half an ounce.
 Lime Water, five pounds.

Boil the submuriate in the lime water for a quarter of an hour, in a covered vessel. Pour off the supernatant liquor, wash the oxide in distilled water, and dry it. (EDINB.)

Hydrargyri Oxydum nigrum. (DUBL.)

R. Sublimed Calomel (Proto-chloruret of Mercury), one part.

Hot Solution of Caustic

Potass, four parts.

Rub them together till the black oxide forms. Wash it well with water, and dry it with a moderate heat, on bibulous paper.

R. Proto-chloruret of Mercury, one part.
Solution of Potass, three parts.
Distilled Water, one part.

Boil, occasionally stirring, for an hour; wash well the powder which is formed, in cold water, and dry it with a moderate heat. (AUST. ; FERR. ; FENN.)

BORIES prescribes the caustic soda, and VAN MONS the caustic barytes.

BORUS. requires the proto-chloruret to be taken while moist from its preparation, and to macerate it in a cool and dark place, with a sufficient quantity of solution of caustic potass; the powder which is formed, to be washed with several waters, and then dried.

R. Proto-chloruret of Mercury,
in powder, one part.
Lime Water twenty parts.

Boil for an hour, constantly stirring; then decant the liquor, pour an equal quantity of lime water upon the residuum, and boil it again: wash the powder with hot water, and dry it slowly. (BATAV.)

AMER. prescribes one ounce of proto-chloruret and eight pints of lime water; to be boiled once.

It is stimulant, sialagogue, and laxative.—Dose, from the eighth of a grain to two grains.

POWDER OF GREY OXIDE OF MERCURY.
(AUGUSTIN)

R. Grey Oxide of Mercury, three grains.
Sugar, one dram and a half.
Opium, two grains.

Divide into six papers.—Dose, one, morning and evening.

RESOLVENT BOLUSES.

Boli resolventes. (BRERA)

R. Subcarbonate of Soda, two scruples.
Proto-chloruret of Mercury, six grains.
Rob of Elder, } of each, a
Liquorice Powder, } sufficient quantity.

Make six boluses.—Dose, one, every two hours.

R. Grey Oxide of Mercury, one grain.
Extract of Hemlock, three grains.
—— Bitter-sweet,
a sufficient quantity.

Make three boluses, to be taken in the course of the day.

VOL. II.

The former of these are recommended in slow inflammations, those particularly of the peritoneum; the latter, in cases of cold tumours, scrofulous or arthritic affections; and engorgements of the viscera after inflammation.

ANTI-SYPHILITIC PILLS. (AUGUSTIN)

R. Grey Oxide of Mercury, one dram.
Crumb of Bread, a sufficient quantity
to make sixty pills.—Dose, one daily.

R. Grey Oxide of Mercury, six grains.
Camphor, twenty-four grains.
Mucilage of Gum Arabic,
a sufficient quantity.

Make twelve pills.—Dose, one or two daily.

STOMACHIC ANTI-SYPHILITIC BOLUSES.

Boli anti-syphilitici stomachici. (BRERA)

R. Grey Oxide of Mercury,
from one to four grains.
Extract of Gentian, one dram.

Make three boluses, to be taken in the course of the day.

RESOLVENT ANTI-SYPHILITIC BOLUSES.

Boli resolventes et anti-syphilitici. (BRERA)

R. Grey Oxide of Mercury,
one grain and a half.
Extract of Aconite, twelve grains.

Make four boluses.—Dose, one every three hours.

ANTI-ARTHRITIC PILLS. (AUGUSTIN)

R. Grey Oxide of Mercury, ten grains.
Opium, } of each,
Extract of Hemlock, } five grains.
Sulphuretted Hydrosulphate of
Antimony, one scruple.
Guaiac Resin, one dram.
Copaiba, a sufficient quantity.

Dose, from two to four grains of the mass, every two hours.

RESOLVENT PILLS. (AUGUSTIN)

R. Grey Oxide of Mercury, one scruple.
Extract of Milfoil, two drams.
—— Henbane, two scruples.
Powdered Rhubarb, one scruple.

Make pills of two grains each.—Dose, fifteen, at thrice, daily; in glandular engorgements.

MOSCATI'S MERCURIAL PILLS. (BORIES;
CADET DE GASSICOURT)

R. Grey Oxide of Mercury, two scruples.
Aqueous Extract of Opium,
twelve grains.

F

Aqueous Extract of Cinchona,
two drams.

Mix, and divide into twenty pills.

JEROMEL has proposed to replace
PLENK'S pills with the following.

R. Grey Oxide of Mercury, one dram.
Powdered Marshmallows Root,
four drams.

Extract of Hemlock, one dram.

Make pills of two grains each.

BLACK MERCURIAL LOTION; BLACK WASH;
BLACK PHAGEDENIC LIQUOR.

*Aqua Hydrargyri muriatici mitis nigra seu
mercurialis nigra, Lotio Hydrargyri.*
(MIL. COPEN.; PH. SYPH.; AUGUSTIN;
ELLIS; NIEMANN; SWEDIAUR; VAN
MONS)

R. Proto-chloruret of Mercury,
sixteen grains.

Lime Water, four ounces.

Triturate. (MIL. COPEN.)

AUGUSTIN and NIEMANN prescribe two
drams of proto-chloruret and one pound of
lime water.—VAN MONS—two scruples of
proto-chloruret and twenty-eight ounces of
lime water.—PH. SYPH.; ELLIS; and SWE-
DIAUR—one dram of proto-chloruret and
four ounces of lime water.

It is employed, particularly in England,
as a lotion in syphilitic eruptions, invete-
rate gonorrhœa in women, and also to sy-
philitic ulcers.

MERCURIAL INJECTION.

Injectio vaginalis cum Mercurio. (AUGUS-
TIN; BRERA)

R. Proto-chloruret of Mercury,
half a dram.

Gum Arabic, one dram.

Lime Water, one ounce and a half.

Plantain Water, three ounces.

This injection is to be used thrice a day,
in blennorrhœa accompanied with local en-
gorgement. (BRERA)

AUGUSTIN prescribes one dram of pro-
to-chloruret, one dram of gum arabic, two
ounces of water, and two ounces of lime
water.

MORETTI'S GREY OXIDE OF MERCURY.
(GALL.; CADET DE GASSICOURT.; SWE-
DIAUR)

R. Mercury, five ounces.
Concentrated Sulphuric Acid,
six ounces.

Put them into a glass retort placed in a
sand-bath, and adapt a receiver; distil with
a gradually augmented heat, till no more

vapours are disengaged; then let the appa-
ratus cool, remove the salt from the retort,
and triturate it in a glass mortar with

Mercury, five ounces,
to perfect combination. Then boil the mass
with

Solution of Caustic Potass,
a sufficient quantity.

Pour the whole upon a filter. Boil again
with potass till the precipitate dissolves en-
tirely in acetic acid. Then wash it well with
pure water, and dry it in the shade. (CADET
DE GASSICOURT)

VAN MONS indicates nearly the same
process; only prescribing caustic soda.

The following process somewhat resem-
bles this.

R. Proto-nitrate of Mercury, one part.
Distilled Water, sixteen parts.

Dissolve, and add gradually

Solution of Potass,
a sufficient quantity,

or till no more precipitate is formed; wash
this with water, and dry it. (GALL.)

There is no real difference between MO-
RETTI'S oxide and the preceding one. They
were both formerly considered protoxides
of mercury: but GUIBOUT has shown them
to be merely a mixture of mercury in a state
of intimate division and of the red oxide.

ANTI-PSORIC OINTMENT.

Unguentum ad Scabiem. (VAN MONS)

R. Nitric Oxide of Mercury, } of each,
Mercury, } one part.
Precipitated Sulphur, sixteen parts.

Triturate to the extinction of the metal,
and add

Lard, thirty-two parts.

Mix.—Although this ointment is not
much to be recommended in the treatment
of itch, yet it deserves to be preferred to
that of DOUBLE and PLANCHE, which will
hereafter be spoken of.—From two drams
to one ounce, according to the age of the
patient, is to be rubbed in at night.

GREY MERCURIAL OINTMENT.

Unguentum Hydrargyri griseum. (PH.
SYPH.; SWEDIAUR)

R. Nitric Oxide of Mercury, one part.
Mercury, three parts.

Triturate to the extinction of the metal
and add

Lard, eight parts.

Mix. (SWEDIAUR)

PH. SYPH. prescribes one part of oxide,
forty-eight parts of mercury, and forty-

eight of lard. It was supposed, by this mixture, to accelerate the passing of the mercury into the state of protoxide; but this effect does not take place, though the preparation is rendered rather more active than the common mercurial ointment.

CAMPHORATED GREY MERCURIAL OINTMENT. (AUGUSTIN)

R. Grey Mercurial Ointment, one dram.
Camphorated Ointment, three drams.

Mix.—LENTIN has recommended this in chronic rheumatism.

II. In a state of Purity.

This is the red oxide, called deutoxide or peroxide of mercury when the metal was thought susceptible of two degrees of oxidation.

LIPP. has given it a place in the materia medica, and also gives a process for its preparation.

A.

In the state of Hydrate.

R. Aqueous Solution of Deuto-chloruret of Mercury, any quantity.
Solution of Caustic Potass, a sufficient quantity.

Pour this into the former, till no more precipitate is formed; wash and dry this. (SUEC.)

YELLOW MERCURIAL LOTION; YELLOW WASH; YELLOW PHAGEDENIC LIQUOR.

Aqua muriatico-mercurialis rubra, Liquor Muriatis Deutoxido-hydrargyro-calcarei, Lotio Hydrargyri lutea, Solutio Muriatis Calcis hydrargyrata. (AMST.; ANT.; BAVAR.; MIL. COPEN.; HISP.; FERR.; FORM. HÔP. FR.; HANN.; ROSS.; SARD.; WIRTEM.; HERBIP.; PH. SYPH.; BRUGNATELLI; ELLIS; HENRI; SPIELMANN; SWEDIAUR; VAN MONS)

R. Deuto-chloruret of Mercury, ten grains.
Lime Water, four ounces.

Triturate, gradually adding the water. (MIL. COPEN.; ANT.)

HANN.; ROSS.; and WIRTEM. prescribe twenty-four grains of the salt, and sixteen ounces of the water.—FORM. HÔP. FR.—nineteen grains of the salt, and one pound of the water.—SPIELMANN—half a dram of the salt, and twelve ounces of the water.—AMST.; and HERBIP.—half a dram of the salt, and one pound of the water.—VAN MONS—one dram of the salt, and thirty-two ounces of the water.—BAVAR.—two drams of the salt, and twelve ounces of the water.—HISP.—from one to three grains of the salt, and one ounce of the water.—FERR.—twenty grains of the salt, and one pound of the water.

R. Deuto-chloruret of Mercury, one part and a fourth.

Water, three hundred and seventy-five parts.

Dissolve, and add

Lime Water, a thousand parts.

Mix. (HENRI)

BRUGNATELLI directs sixteen ounces of lime water to be added to a solution of one scruple of the salt in a sufficient quantity of water.

R. Deuto-chloruret of Mercury, one dram.
Lime Water, one pound.

Dissolve, and add

Proof Spirit, four ounces.

Mix. (WIRTEM.)

PH. SYPH.; and SWEDIAUR prescribe from thirty to sixty grains of the salt, two pounds of the water, and half a pound of proof spirit.

This preparation consists of a very dilute solution of hydrochlorate of lime, and of a precipitate of the hydrate of deutoxide of mercury, when it has a yellow colour, and a mixture of deuto-chloruret and of deutoxide, when it is red or rather brick-coloured.

R. Deuto-chloruret of Mercury, one dram.
First Lime Water, two pounds.

Triturate. (SARD.)

First lime water containing a little potass, this solution contains hydrochlorate of potass.

R. Deuto-chloruret of Mercury, one scruple.

Sub-carbonate of Potass, one dram.

Triturate in a glass mortar, adding gradually

Distilled Water, one pound.

Mix. (HISP.)

This preparation differs from the preceding in that hydrochlorate of potass alone is in solution, the precipitate being subcarbonate of mercury.

This lotion is applied to syphilitic, scrofulous, and other spreading and obstinate ulcers. Before using, it should be shaken so as to mix the sediment with the liquor, for to that it owes, if not all, the greater share of its activity.

ALOETIC MERCURIAL LIQUOR.

Aqua aloetica mercurialis. (FULD.; SPIELMANN)

R. Deuto-chloruret of Mercury, half a dram.

Lime Water, twelve ounces.

Dissolve, and add

Aloes, in powder, six drams.

Honey of Roses, one ounce and a half.

Mix. (SPIELMANN)

Hydrargyri Oxydum nitricum.
(DUBL.)

R. Purified Mercury, two parts.
Dilute Nitrous Acid, three parts.

Dissolve the mercury, and apply heat till the matter is converted into red scales.

R. Mercury, any quantity.
Nitric Acid (32 degrees), a sufficient quantity,

to dissolve all the metal; pour the metal into a flat bottomed matrass placed in a sand-bath; then pour in the acid, and when the solution is effected, augment the heat to evaporate the liquor to dryness: then increase it till no more nitrous fumes are disengaged, and the mass becomes of a fine red colour. Preserve it when cold. (AMST.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; WIRTEM.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

PAISSE' has made known the following process, framed from that of the Dutch manufacturers, and which is considered a better one:

R. Mercury, one hundred pounds.
Nitric Acid (1, 3 to 1.37), one hundred and forty pounds.

Dissolve the mercury with the heat of a sand-bath; evaporate by distillation; and when the formation of nitrous gas indicates the decomposition of the nitrate, remove the receiver, and apply a constant and moderate heat for about eight hours, till a candle-wick recently extinguished inflames on being put into the matrass.

HENRI recommends the fire not to be pushed till oxygen is disengaged, as BAVAR. and PAISSE' direct, because it announces the commencement of the decomposition of the oxide.

AMST.; HISP.; HANN.; OLDEN.; POL.; BORUS.; ROSS.; HERBIP.; and COXE direct the operation to be effected at twice, and after having evaporated the solution to dryness, the residuum to be pulverized and calcined in another vessel.

BATAV. recommends the product to be digested for several hours with ten parts of hot water and a little solution of potass; the powder to be then washed several times with hot water. By this means the oxide is deprived of a small quantity of nitric acid, or rather of sub-deuto-nitrate of mercury, which might have remained adherent to it.

R. Salt, obtained by the pre- } of each,
ceding process, } one part.
Mercury, }
Distilled Water, a sufficient quantity.

Triturate together to the perfect extinction of the metal; then calcine, as above, in a crucible. (ANT.; SWEDIAUR)

BAVAR. and FENN. prescribe two parts of mercury.—SUEC.—two-thirds of the metal.

It is excitant, irritant, and poisonous in a high degree. It was formerly, however, much employed, and is still prescribed in syphilitic diseases, either internally, in the dose from half a grain to one grain, or externally, in frictions, mixed with saliva, in the quantity of ten grains, each friction.

POWDER OF NITRIC OXIDE OF MERCURY.

Pulvis Hydrargyri oxydati rubri. (AUGUSTIN; NIEMANN)

R. Red Oxide of Mercury, one grain.
Sulphuret of Antimony, two drams.

Triturate together, and divide into eight equal parts.

It is recommended in syphilitic disorders.—Dose, one part, at twice, morning and evening, in water. The dose may be a little augmented.

PLENK'S CAUSTIC POWDER.

Pulvis ex Hydrargyro rubro cum Alumine, Pulvis causticus seu aluminoso-mercurialis. (FERR.; FULD.; CAST. BORUS.; PH. SYPH.; AUGUSTIN; SWEDIAUR)

R. Nitric Oxide of Mercury, one part.
Alum, two parts.

Mix by trituration. (FERR.; FULD.; CAST. BORUS.; AUGUSTIN; NIEMANN; SWEDIAUR)

PH. SYPH. prescribes an equal part of each.

R. Nitric Oxide of Mercury, } of each,
Calcined Alum, } one dram.
Powdered Savine, half an ounce.

Mix intimately. (NIEMANN)

It is recommended to remove warts and luxuriant granulations of ulcers.

ESCHAROTIC POWDER. (NIEMANN)

R. Nitric Oxide of Mercury, }
Acetate of Copper, } of each,
Fused Potass, } one dram.
Alum, }
Powdered Savine, } of each,
Charcoal, } two drams.

Triturate, and make a fine powder.

This is a much more active preparation than the preceding.

DRY COLLYRIUM.

Collyre sec. (RATIER)

- R. Red Oxide of Mercury, ten grains.
Tutty, twenty grains.
White Sugar, two drams.
Triturate.

This powder is blown into the eyes to remove spots on the cornea which occur after ophthalmia.

GREEN PRECIPITATE.

Mercurius præcipitatus viridis. (WIRTEM.)

- R. Mercury, four ounces.
Copper, half an ounce.

Dissolve each, separately, in
Nitric Acid, a sufficient quantity.

Mix together the two solutions, and evaporate in a phial placed in a sand-bath. Pour upon the residuum

Distilled Vinegar, one pound,
and leave it to digest till perfectly dissolved; then distil off the vinegar, and pulverize the green mass which remains in the retort.

It is violently emetic and drastic. It is extolled as a specific in syphilitic diseases, and is employed also, externally, as a detergent to ulcers. It is a formidable poison.

BOLUS OF NITRIC OXIDE OF MERCURY.

Bolus ex Hydrargyro oxydato rubro. (PH. SYPH.; AUGUSTIN)

- R. Nitric Oxide of Mercury, } of each,
Opium, } one grain.
Extract of Liquorice, a sufficient quantity

to unite them and make a bolus.

PILLS OF NITRIC OXIDE OF MERCURY.

Pilule anti-venereæ seu syphiliticæ seu polychrestæ. (FULD.; HERBIP.; ELLIS; SAUNDERS; PH. SYPH.)

- R. Nitric Oxide of Mercury, eight grains.
Extract of Liquorice, a sufficient quantity

to make a mass, to be divided into twelve pills. (PH. SYPH.)

- R. Nitric Oxide of Mercury, one grain.
Prepared Red Coral, two grains.
Conserve of Roses, a sufficient quantity

to make one pill. (SAUNDERS)

- R. Nitric Oxide of Mercury, three grains.
Opium, one grain.
Oil of Cloves, two drops.

Make three pills.

- R. Nitric Oxide of Mercury, } of each,
Opium, } one dram.
Balsam of Peru, a sufficient quantity

to make pills of one grain each. (HERBIP.)

- R. Nitric Oxide of Mercury, ten grains.
Sulphuretted Hydrsulphate } of each,
of Antimony, } twenty
Assafœtida } grains.
Opium, five grains.
Syrup of Poppies, a sufficient quantity

to make a pilular mass. (FULD.)

The dose is one pill at most, of the three first formulæ. As to the latter, from two to eight grains of the mass may be given. They are all reputed resolvent and anti-spasmodic.

MERCURIAL HONEY. (VAN MONS)

- R. Nitric Oxide of Mercury, } of each,
White Sugar, } one scruple.
Triturate together, and add gradually
Thick Honey, three ounces.

Mix.—It is recommended as an application to certain syphilitic ulcers.

OINTMENT OF NITRIC OXIDE OF MERCURY.

Balsamum ophthalmicum rubrum, Unguentum Deutoxidi Hydrargyri nitrati seu Hydrargyri Nitrico-oxydi seu Hydrargyri rubri seu ex Hydrargyro oxydato rubro seu Mercurii rubrum seu ophthalmicum rubrum seu Oxydi Hydrargyri rubri seu Oxydi hydrargyrici nitrati seu Subnitratis hydrargyrici.—Golden Ointment, Anti-psoric Ointment, LYON'S Ointment, PELLETAN'S Ophthalmic Ointment. (LOND.; EDINB.; DUBL.; AMER.; AMST.; BATÁV.; BELG.; DAN.; HAMB.; OLDEN.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUÉC.; HERBIP.; AUGUSTIN; BORIES; BRUGNATELLI; CADET DE GASSICOURT; COXE; HUFELAND; RATIER; SPIELMANN; VAN MONS)

*Unguentum Hydrargyri Nitrico-Oxydi. (LOND.)**Unguentum Hydrargyri Oxydi nitrati. (DUBL.)*

- R. Nitric Oxide of Mercury, one ounce.
White Wax, two ounces.
Prepared Lard, six ounces.

Melt together the wax and lard, then add the nitric oxide in very fine powder, and mix.

Unguentum Oxydi Hydrargyri rubri. (EDINB.)

- R. Nitric Oxide of Mercury, in fine powder, one part.
Lard, eight parts.

Mix well.

AMER.; and COXE give the same formula as EDINB.—BATAV.; BELG.; and AUGUSTIN prescribe one part of the oxide and thirty-two parts of lard.—HANN.—one of oxide and ten of lard.—AMST.—one of oxide and forty-eight of lard.—HUFELAND—one of oxide and thirty-six of lard.—AUGUSTIN—from one to four grains of oxide and two drams of walnut oil.

R. Nitric Oxide of Mercury, one part.
Fresh Butter melted slowly,
eight parts.
Triturate together. (DAN.; SUEC.)

R. Nitric Oxide of Mercury, eight parts.
Lard, eighty parts.
Oil of Lemon, one part.
Mix. (BORUS.)

R. Nitric Oxide of Mercury, one part.
Fresh Butter, four parts.
Yellow Wax, two parts.

Melt together the two latter, and add the oxide in fine powder. (CAST. BORUS.)

VAN MONS prescribes one part of oxide, twelve parts of butter, and six of wax; and calls the product *Yellow Ophthalmic Balsam*.—RATIER—one and a half of oxide, four of white wax, and twenty-four of butter.—ROSS.—seventeen grains of oxide, three drams of butter, and forty-five grains of white wax.—SAX.—one part of oxide, nine parts of butter, and three of white wax.—SPIELMANN—two drams fifteen grains of oxide, three ounces of butter, and six drams of white wax.—HANN.; and OLDEN.—one part of oxide, two parts of butter, and two of white wax.

R. Nitric Oxide of Mercury, one part.
White Wax, three parts.
Olive Oil, eight parts.

Melt the two latter, and add the former. (BRUGNATELLI)

PLANCHE gives this formula: one part of oxide, four parts of white wax, and eight of oil; which he calls *Anti-psoric Ointment*.—WURCEB. prescribes one of oxide and six of common cerate.—POL.—one of oxide and ten of cerate.

R. Red Oxide of Mercury, one part.
Ointment of Roses, eight parts.

Mix. (RATIER)

BORIES; and CADET DE GASSICOURT prescribe one part of oxide and two parts of ointment.

R. Nitric Oxide of Mercury, ten grains.
Camphor, six grains.
Fresh Butter, two drams.

Triturate together. (AUGUSTIN)

This ointment, the strength of which varies so much, since the oxide is found with the fat in the proportion of from one to two to one to forty-eight, is employed as

a stimulant in chronic affections of the eyelids and of the exterior membrane of the eye, when there is no inflammation or scarcely any. One or two grains of it are introduced into the inner angle of the eye twice a day. DOUBLE recommends PLANCHE's ointment in itch; the quantity prescribed in the formula is to be divided into six for as many frictions, to be made in the evening before the fire. But an ointment, into which the nitric oxide of mercury enters in the proportion of one in thirteen, is certainly dangerous, and should not be recommended in such an affection.

RICHTER'S OPHTHALMIC OINTMENT.

Unguentum ophthalmicum Richteri. (HANN)

R. Nitric Oxide of Mercury, one part.
Cocoa Nut Oil, } of each, three parts.
Prepared Lard, }

Melt together the two latter, and when nearly cold, mix intimately the oxide.

ESCHAROTIC OINTMENT. (ELLIS)

R. Nitric Oxide of Mercury, one dram.
Venice Turpentine, one ounce.
Triturate together.

ANTI-SYPHILITIC OINTMENT. (AUGUSTIN)

R. Nitric Oxide of Mercury, five grains.
Grey Mercurial Ointment, one dram.
Mix.—For one friction.

The use of this ointment, continued to sufficient salivation, remaining in bed in a warm chamber, and at the same time being subjected to a severe and watery regimen for several weeks, constitutes the method extolled by LOUVRIER and RUST for the radical cure of syphilis.

BÖCKING'S MERCURIAL OINTMENT.

(FORM. HÔP. FR.; PH. SYPH.; RATIER; SAUNDERS; SAINTE-MARIE)

R. Nitric Oxide of Mercury, one part.
Resinous Ointment, six parts.

Mix. (SAUNDERS)

PH. SYPH. prescribes one part of oxide and ten parts of ointment.—SAINTE-MARIE—one of oxide and twelve of ointment.—FORM. HÔP. FR.—one of oxide and sixteen of ointment.—RATIER—one of oxide and forty-eight of ointment.

It is applied as a stimulant to indolent and particularly syphilitic ulcers.

In using this ointment, it should be commenced with very weak; the active ingredient might be augmented or diminished according to the effects produced by it, which should be carefully watched.

MERCURIAL OINTMENT. (SAINTE-MARIE)

- R. Nitric Oxide of Mercury, } of each,
 Camphor, } one dram.
 Fresh Butter, } four ounces.
 White Wax, } three drams.

Melt together the wax and butter, then add the oxide, and finally the camphor.

It is recommended in inflammation of the glans penis (*balanite*), to be rubbed upon the glans, every night on going to bed, for a fortnight. It is a means of keeping up this insignificant affection, which yields in a few days to cleanliness alone.

OPHTHALMIC BALSAM. (VAN MONS)

- R. Red Oxide of Mercury, four parts.
 — Sulphuret of Mercury, two parts.
 Camphor, one part.
 White Cerate, ninety-four parts.

Triturate.

STIMULANT OINTMENT.

Unguentum Elemi cum Alumine. (AUGUSTIN)

- R. Ointment of Elemi, half an ounce.
 — Storax, two drams.
 Balsam of Peru, half a dram.
 Nitric Oxide of Mercury, fifteen grains.
 Calcined Alum, five grains.
 Oil of St. John's-wort, a sufficient quantity.

Mix.—MICHAELIS has praised it as an application to wounds and ulcers of a bad aspect.

OINTMENT FOR SCROFULOUS ULCERS.

Unguentum ad Ulcera Lymphatica.
 (VAN MONS)

- R. Nitric Oxide of Mercury, one part.
 Plaster of Hemlock, } of each,
 Ointment of Resin, } forty parts.
 Mix.

OINTMENT FOR SYPHILITIC ULCERS.

Unguentum ad Ulcera venerea. (VAN MONS.)

- R. Nitric Oxide of Mercury, one part.
 Mercurial Ointment, four parts.
 Ointment of Resin, thirty-two parts.
 Mix.

SUPPURATIVE OINTMENT.

Unguentum suppurativum. (VAN MONS)

- R. Nitric Oxide of Mercury, one part.
 Triturate it with a few drops of camphorated oil, and then add to it

ARCÆUS' Balsam, twelve parts.
 Ointment of Resin, forty-eight parts.
 Mix.

SAINT-YVES OPHTHALMIC BALSAM.

Balsamum ophthalmicum Yveanum, Unguentum Hydrargyri rubrum. (BRUNS.; FULD.; HANN.; HASS.; LIPP.; ROSS.; SAX.; HERBIP.; AUGUSTIN.; CADET DE GASSICOURT.; PIDERIT.; PIERQUIN.; SPIELMANN.; VAN MONS.)

- R. Nitric Oxide of Mercury, five parts.
 Oxide of Zinc, two parts.

Mix, and throw it into

Fresh Butter, twenty parts.
 Yellow Wax, ten parts.

melted together. Triturate. (LIPP.)

FULD. prescribes fifteen parts of oxide, six of oxide of zinc, sixty of butter, and thirty of wax.

- R. Nitric Oxide of Mercury, thirty parts.
 Oxide of Zinc, twelve parts.
 Camphor, dissolved in Oil of Egg, nine parts.

White Wax, forty-eight parts.
 Fresh Butter, two hundred and forty-eight parts.

Melt together the two latter, add the others, and triturate till cold. (BRUNS.; HANN.; HASS.; ROSS.; HERBIP.; CADET DE GASSICOURT.; PIDERIT.; PIERQUIN.; SPIELMANN.)

VAN MONS prescribes three parts of oxide of mercury, three of oxide of zinc, one of camphor triturated with oil of almonds, sixteen of wax, and ninety-six of butter.

- R. Ointment of Nitric Oxide of Mercury, sixty parts.
 Oxide of Zinc, two parts.
 Camphor, one part.
 Mix. (SAX.)

SPIELMANN prescribes ninety-six parts of ointment, three of oxide of zinc, and two of camphor.—AUGUSTIN—five grains of nitric oxide, four of tutty, two of camphor, and two drams of fresh butter; or, four grains of nitric oxide, ten of oxide of zinc, as much of extract of opium, and two drams of lard.

This ointment is scarcely different from that of the simple ointment of the nitric oxide, but it is said to be preferable when there is inflammation. It is considered detergent, resolvent, refrigerant, and corroborant. Applied too freely, it is irritant.

DUPUYTREN'S OPHTHALMIC OINTMENT.
 (RATIER)

- R. Red Oxide of Mercury, one part.
 Sulphate of Zinc, two parts.
 Lard, ninety-six parts.

Mix.—It is recommended in chronic and obstinate scrofulous ophthalmias. The margins of the eye-lids are lightly rubbed with it.

RED OINTMENT. (VAN MONS)

- R. Red Oxide of Mercury, } of each,
 Lead, } one part.
 Lard, } eight parts.
 Triturate the oxides gradually with the lard.

JOERDEN'S OINTMENT. (AUGUSTIN ; NIEMANN)

- R. Red Oxide of Mercury, one part.
 Fresh Butter, sixteen parts.
 Mix well together, and add
 Solution of Subacetate of Lead, six parts,
 and triturate till the liquid shall have totally disappeared.

RUST'S OINTMENT. (AUGUSTIN)

- R. Nitric Oxide of Mercury, from four to six grains.
 Fresh Butter, two drams.
 Solution of Subacetate of Lead, } of each,
 SYDENHAM'S Laudanum, } half a dram.
 Mix.

REGEN'T'S OINTMENT. (GALL. ; FORM. HÔP. FR. ; PIERQUIN)

- R. Fresh Butter, washed with
 Rose Water, 4,05.
 Camphor, }
 Nitric Oxide of Mercury, } of each,
 Acetate of Lead, } 0.25.

Triturate for a long time. (GALL.)

FORM. HÔP. FR. prescribes one part each, of nitric oxide, of camphor, and of acetate, and eighteen parts of lard.—PIERQUIN—twenty of nitric oxide, twenty of acetate, one of camphor, and seventy of butter.

DESAULT'S OINTMENT. (CADET DE GASSICOURT ; PIERQUIN ; RATIER)

- R. Nitric Oxide of Mercury, } of each,
 Acetate of Lead, } half
 Calcined Alum, } an ounce.
 Tutty, }
 Deuto-chloruret of Mercury, one grain.
 Lard, three ounces.

Mix. (RATIER)

PIERQUIN prescribes four drams each, of nitric oxide, of acetate, of alum, and of tutty, forty-eight grains of deuto-chloruret, and three ounces of lard.—CADET DE GASSICOURT—one dram each, of nitric oxide, of litharge, of tutty, and of calcined alum,
 VOL. II.

twelve grains of deuto-chloruret, and a sufficient quantity of cerate.

This ointment is celebrated against herpetic eruptions and chronic inflammations of the eye-lids. It differs from the preceding only in being more irritant.

BELL'S OINTMENT. (BORIES)

- R. Nitric Oxide of Mercury, } of each,
 Prepared Calamine, } one dram
 Litharge, } and a half.
 Tutty, } one dram.
 Red Sulphuret of Mercury, half a dram.
 Lard, one scruple.
 two ounces.

Triturate, and add
 Balsam of Peru, fifteen drops.

OPHTHALMIC OINTMENT. (AUGUSTIN)

- R. Nitric Oxide of Mercury, from ten to twenty grains.
 Fresh Butter, three drams.
 Sub-acetate of Copper, from six to eight grains.
 White Bole, } of each, one scruple.
 Opium, }
 Triturate.—BENEDICT recommends it in pterigium.

- R. Nitric Oxide of Mercury, six grains.
 Sub-acetate of Copper, ten grains.
 Tutty, fourteen grains.
 Fresh Butter, half an ounce.
 Triturate.

FALCK'S MERCURIAL CERATE.

Ceratum detergens seu mercuriale Falckii.
 (HERBIP. ; CADET DE GASSICOURT ; PIDERIT ; VAN MONS)

- R. Nitric Oxide of Mercury, half an ounce.
 Proto-chloruret of Mercury, two ounces.
 Acetate of Lead, one ounce.

Levigate and incorporate them with a melted mixture of

White Wax, four ounces.
 Olive Oil, six ounces.

Then add

Oil of Lavender, half a dram.

Mix well. (PIDERIT)

HERBIP. gives the same formula, but omits the oil of lavender.—CADET DE GASSICOURT substitutes walnut oil for the olive oil, and reduces the wax to one ounce.

- R. Red Oxide of Mercury, one half part.
 Acetate of Lead, one part.
 Olive Oil, six parts.
 Yellow Wax, one part.

Triturate the oxide and the salt separately with a little oil ; add the rest of the oil to

the wax melted over a slow fire, and incorporate in the mixture first the salt, and then the oxide. (VAN MONS)

It is stimulant, and detergent, and is applied to syphilitic ulcers, either in frictions or on lint.

FALCK'S OINTMENT. (BORIES)

- R. Nitric Oxide of Mercury, two drams.
 Amalgam of Tin and Mercury,
 at equal parts, four drams.
 Rose Ointment, one ounce.
 Triturate together, and add
 Oil of Peppermint, twenty drops.

Mix.—It is recommended against hæmorrhoidal tumours.

FALCK'S BOUGIES. (HASS.; PIDERIT)

- R. Venice Turpentine, one dram.
 Gum Lac, in powder, half a dram.
 Melt slowly, and add
 Mercurial Plaster, two ounces.
 Proto-chloruret of Mercury, two drams.
 Nitric Oxide of Mercury, levigated,
 two scruples.

Mix, and make bougies.

SULPHURET OF MERCURY.

I. Black Sulphuret.

BAVAR.; FERR.; LIPP.; SUEC.; VAN MONS.

This preparation has been for a long time confounded, under the name of *Æthiops mineral*, with the mechanical mixture of sulphur and mercury; but it differs from the latter in its proportions being constant, and also, according to SEGUIN, in changing when heated sufficiently in a tube, to red sulphuret without losing in weight.

A.

In a state of purity.

1. Prepared by the action of fire.

- R. Sublimed Sulphur, one part.
 Melt them over a slow fire, in an earthen vessel, and add gradually
 Mercury, heated in an iron spoon,
 one part.

Stir incessantly till the mass shall have acquired a uniform black colour, and levigate when cold. (BAVAR.; SUEC.; VAN MONS)

FERR. prescribes three parts of sulphur, and two of mercury.

2. Prepared by Precipitation.—Hypo-hydro-sulphuret of Mercury.

- R. Deuto-chloruret of Mercury,
 twelve parts and a half.

Hydro-sulphate of Iron, } of each,
 Concentrated Sulphuric } five parts.
 Acid,

Dissolve the deuto-chloruret in water, and introduce into the solution the gas disengaged from the mixture of the two other substances. Then add

Water, twenty-five parts.

Collect, wash, and dry the precipitate. (VAN MONS)

- R. Subcarbonate of Potass, two ounces.
 Sublimed Sulphur, four ounces.
 Lime Water, two pounds.

Boil in a long-necked glass retort placed in a sand-bath, till the sulphur is dissolved. Set aside the liquor to cool, filter, and add drop by drop a solution made without heat with

Mercury, two ounces.
 Nitric Acid, four ounces.

When the effervescence has ceased, set apart the liquor, pour off the clear part, wash the precipitate with distilled water, and dry it with a gentle heat. (FERR.; LIPP.; VAN MONS)

VAN MONS also prescribes solution of potass instead of subcarbonate.

B.

Mixed with Sulphuret of Antimony.

This preparation ought to be distinguished from the *Mercury with Sulphur and Sulphuret of Antimony*.

- R. Sulphur, one part and a half.
 Fused Potass, } of each,
 Sulphuret of } five parts and a half.
 Antimony,

Boil these in water till nothing more is dissolved, and filter. Boil the strained liquor, and instil into it

Solution of Nitrate of Mercury,
 a sufficient quantity,
 till no more precipitate is formed. Wash and dry this. (VAN MONS)

VAN MONS also proposes,

1. To substitute the solution of deuto-chloruret of mercury for that of the nitrate, and to pour suddenly twelve parts of it into the liquor.

2. To melt together six parts of sulphuret of potass, and five and a half of sulphuret of antimony; to pulverize the mixture when cold, and to triturate it with a little water; to add twelve parts of deuto-chloruret of mercury dissolved in water, or twenty of proto-chloruret: the precipitate to be washed in a great quantity of water, and dried.

Sugar Candy, three ounces.
Oil of Cinnamon, from half a dram to one dram.

Triturate well together. (BRUNS.; OLDEN.; NIEMANN; SWEDIAUR)

WIRTEM.; and SPIELMANN prescribe six drams of sulphuret, two ounces of sugar, and two scruples of oil of cinnamon.

It is stimulant, and is rather celebrated as cordial, stomachic, and analeptic.—Dose, a scruple.

CAMERARIUS' CEPHALIC POWDER. (WIRTEM.)

- R. Red Sulphuret of Mercury, half a dram.
Rock Crystal, } of each, one dram.
Prepared Amber, }
Elk's Hoof, }
—Horn, prepared } of each,
without Fire, } three drams.
Hart's Horn prepared }
without Fire, }
Prepared Human Skull, six drams.
White Sugar, one ounce and a half.

Make a powder.

The preceding powder should be substituted for this absurd composition.—Dose, from half a scruple to three scruples.

ANTI-EPILEPTIC POWDER.

Pulvis epilepticus ruber seu ad Guttetam.
(SARD.; WIRTEM.; BORIES; PIERQUIN)

- R. Red Sulphuret of Mercury, } of each,
Prepared Amber, } one
Sea Unicorn's Horn, } ounce.
Root of Peony, } of each,
Red Coral, } two ounces.

Make a powder. (WIRTEM.)

Dose, from half a scruple to half a dram.

- R. Root of Peony, } of each,
Hartshorn, } half an ounce.
Prepared Amber, }
Castor, one dram.
Red Sulphuret of Mercury, two drams.

Mix. (SARD.)

- R. Root of Peony, } of each, one ounce.
—Valerian, }
Seeds of Rue, } of each,
Myrrh, } half an ounce.
Castor, }
Amber, }
Red Sulphuret of Mercury, two ounces.

Mix. (BORIES; PIERQUIN)

- R. Mistletoe, } of each,
Red Sulphuret of Mercury, } two
Elk's Hoof, } drams.
Prepared Human Skull, half an ounce.

Mix. (WIRTEM.)

Dose, half a scruple to children, and from a scruple to half a dram to adults.

STAHL'S TEMPERANT POWDER.

Pulvis anti-spasmodicus seu nitrosus cinnabarinus seu salinus compositus seu temperans ruber. (AMST.; BATAV.; BELG.; BRUNS.; GALL.; LIPP.; OLDEN.; PALAT.; WIRTEM.; CADET DE GASSICOURT; PIERQUIN; SAUNDERS; SWEDIAUR; VAN MONS)

- R. Red Sulphuret of Mercury, one part.
Nitrate of Potass, four parts.
Sulphate of Potass, one part.

Triturate together. (BATAV.; BELG.)

AMST.; and VAN MONS prescribe one part of sulphuret and four parts each, of the two salts.—SWEDIAUR—one of sulphuret, sixteen of nitrate, and eight of sulphate.—VAN MONS—one of sulphuret, and sixteen each, of the two salts.—OLDEN.—one of sulphuret, and twelve each, of the two salts.—GALL.; CADET DE GASSICOURT; and PIERQUIN—two of sulphuret, and nine each, of the two salts.—LIPP.; and WIRTEM.—three of sulphuret, and sixteen each, of the two salts.

- R. Red Sulphuret of Mercury, one part.
Prepared Oyster Shells, } of each,
Nitrate of Potass, } two parts.
Sulphate of Potass, }

Mix. (VAN MONS)

PALAT. prescribes six drams each, of nitrate, of sulphate, and of oyster shells, and three of sulphate.

- R. Red Sulphuret of Mercury, one part.
Prepared Crab's Eyes, six parts.

Triturate together. (SAUNDERS)

- R. Red Sulphuret of Mercury, five parts.
Crab's Eyes, } of each,
Prepared Red Coral, } eight parts.
—Mother of }
Pearl, } of each,
Sulphate of Potass, } sixteen parts.
Nitrate of Potass, thirty-two parts.

Mix. (BRUNS.; WIRTEM.)

Although evidently stimulant, this powder has for a long time enjoyed great celebrity as a refrigerant and alterative.—The dose of that furnished by the last formula is from one to two scruples.

TEMPERANT POTION. (VAN MONS)

- R. STAHL'S Temperant Powder, fifteen grains.
Almond Emulsion, six ounces.
Syrup of Marshmallows, one ounce.
Orange-flower Water, four drams.

Mix.—This is recommended at the commencement of acute diseases. It can then only do harm by the excitement which it occasions, in spite of the name given to it, and which it does not deserve.

DIAPHORETIC BOLUSES. (SAINTE-MARIE)

- R. Red Sulphuret of Mercury, one scruple.
 Camphor, two scruples.
 Theriaca, one ounce.
 Syrup, a sufficient quantity

to make a mass to be divided into twenty-four boluses.

Three to be taken in the course of the day, with some warm drink.

RESOLVENT PILLS.

Pilule Conii Cinnabarinæ. (AUGUSTIN)

- R. Red Sulphuret of Mercury, } of each,
 Extract of Milfoil, } one
 Hemlock, } dram.

Mix.—Fourteen grains of the mass to be taken morning and evening; in syphilitic glandular engorgements.

MERCURIAL PASTILLES. (SPIELMANN)

- R. Levigated Red Sulphuret of
 Mercury, two scruples.
 Prepared Amber, one scruple.
 Mucilage of Tragacanth,
 a sufficient quantity
 to make pastilles.—For fumigation.

RED CERATE. (SWEDIAUR)

- R. Red Sulphuret of Mercury, fifteen grains.
 Yellow Resin, half an ounce.
 Wax, } of each, half a pound.
 Lard, }

Melt the three latter, and add the sulphuret.

It is stimulant and desiccative, applied to ulcers.

BLACK OINTMENT. (VAN MONS)

- R. Red Sulphuret of Mercury, three parts.
 Hydrosulphate of Lead, one part.
 Beef Marrow, twelve parts.
 Triturate.

OINTMENT OF RED SULPHURET OF MERCURY. (ELLIS)

- R. Red Sulphuret of Mercury, two ounces.
 Semi-vitreous Oxide of Lead, four ounces.
 Subcarbonate of Lead, one ounce and a half.
 Proto-chloruret of Mercury, two drams.
 Levigate each substance separately, then mix and incorporate the powders in a melted mixture of
 Yellow Wax, six ounces.
 Oil of Roses, two pounds.

ANTI-HERPETIC OINTMENT. (SAINTE-MARIE; RATIER)

- R. Red Sulphuret of Mercury, one dram.
 Camphor, one scruple.
 Common Cerate, one ounce.

Triturate. (SAINTE-MARIE)

This is frequently applied with success, by ALIBERT, to tetters; the ointment is to be renewed at least twice a day.

- R. Red Sulphuret of Mercury, one dram and a half.
 Hydrochlorate of Ammonia, half a dram.
 Lard, two ounces.

Triturate together, and add

- Rose Water, one dram.

Mix.—It is especially recommended in the morbus pedicularis, to moderate the itching, and diminish the production of the lice.

ANTI-ARTHRITIC ELECTUARY. (CADET DE GASSICOURT)

- R. Red Sulphuret of Mercury, one ounce and a half.
 Conserve of Roses, two ounces.
 Guaiac Resin, one ounce.
 Syrup of Ginger, a sufficient quantity.

Mix.—It is recommended in gouty-rheumatic affections unattended with fever.—Dose, a tea spoonful twice a day.

PURGATIVE ELECTUARY. (RATIER)

- R. Red Sulphuret of Mercury, } of each,
 Jalap, } two drams.
 Scammony, } one dram.
 Squill, } of each,
 Resin of Jalap, } one dram and a half.
 Syrup of Buckthorn, a sufficient quantity.

Mix.—Dose, from half a dram to two drams.—It is employed by FOUQUIER in dropsy and painter's colic.

RED SPARADRAP. (VAN MONS)

- R. Red Sulphuret of Mercury, one part.
 Turpentine, six parts.
 White Wax, twelve parts.

Melt, and incorporate; then dip strips of linen in the mass, and polish them.

CINNABARINE EXTRACT OF OPIUM.

Extractum Opii cinnabarinum, Laudanum cinnabarinum. (HISP.)

- R. Aqueous Extract of Opium, six drams.
 Saffron, in powder, } of each,
 Red Sulphuret of Mercury, } three drams.

Make them into a mass with a little water.

PROTO-CHLORURET OF MERCURY.

Antiquarium, Aquila alba seu mitigata, Calomelas, Draco mitigatus, Hydrargyrum muriaticum dulcis seu oxydulatus seu muriatus dulcis, Hydrargyrum muriaticum mite, Mercurius dulcis seu muriaticus seu præcipitatus albus seu sublimatus dulcis, Murias Hydrargyri oxydulati seu mitis seu suboxygenatus seu hydrargyrosus mitis seu Oxyduli Hydrargyri mitis, Manna Metallorum, Panchymagogum minerale, Panchymagogus Quercetanus Proto-chloruretum Hydrargyri, Submurias Hydrargyri seu Hydrargyri mitis seu præcipitatus seu sublimatus.—Calomel, Mild Muriate of Mercury. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPE.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PARIS; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

1. The ordinary process.

Submurias Hydrargyri mitis sive Calomelas. (EDINB.)

R. Muriate (Deuto-chloruret) of Mercury, four parts.
Purified Mercury, three parts.
Triturate the muriate in a glass mortar with a little water to prevent the acrid powder from rising; then add the mercury and triturate till it is extinguished. Put the mass, dried, into an oblong flask, which it fills to one third, and sublime it in a sand-bath. Triturate the sublimed powder, and sublime again: finally reduce it to fine powder, and wash it with boiling distilled water.

This formula presents numerous varieties. HANN.; OLDEN.; BORUS.; and SARD. give the same as EDINB.—SAX. gives the same, but requires but one sublimation.

R. Deuto-chloruret of Mercury, four parts.
Mercury, three parts.

Triturate in a glass mortar, adding a little alcohol, till the metal is extinguished; put the powder into a phial which it fills to one third only; stop the phial, placed in a sand-bath, first with paper, and afterwards with a stone stopper: gradually increase the fire, and keep it up till nothing remains at the bottom of the vessel. Then reduce the sublimed mass to powder, and boil with double

its weight of distilled water and one twentieth of hydrochlorate of ammonia. Decant the liquid, wash the powder twice or thrice with distilled water, and then dry it with a gentle heat. (ANT.; BATAV.; BELG.; FERR.; ROSS.)

BRUNS.; PALAT.; and WIRTEM. prescribe an equal part each, of deuto-chloruret and of mercury, trituration with alcohol, two successive sublimations, and washing with a boiling solution of one dram of hydrochlorate of ammonia to every pound of water.—HERBIP.—an equal part each, of deuto-chloruret and of mercury, trituration with alcohol, one sublimation, and washing with boiling water alone.—AUSTR.—five ounces of deuto-chloruret and four of mercury, one sublimation, and washing with cold water.—DAN.—one pound of deuto-chloruret and nine ounces of mercury, two sublimations, and washing.—AMST.—one pound of deuto-chloruret and ten ounces and a half of mercury, trituration with a little water, three sublimations, and no washing.—FORM. HÔP. FR.—twenty-four parts of deuto-chloruret and fifteen of mercury, trituration with water, two sublimations, and washing with boiling water.—LUSIT.—one pound of deuto-chloruret and nine ounces of mercury, trituration with water, two sublimations, and washing with boiling water.—GALL.—430 of deuto-chloruret and 320 of mercury, trituration with water, several successive sublimations, and washing with boiling water.—SWEDIAUR—fifteen ounces of deuto-chloruret and twelve of mercury, trituration with alcohol, two sublimations, and washing with boiling water.—BAVAR.—four parts of deuto-chloruret and two of mercury, two sublimations, and washing with four parts of boiling alcohol.—VAN MONS—12,7 of deuto-chloruret and 9,5 of mercury, one sublimation, and no washing.—BRUGNATELLI—an equal part of each, trituration with water, and no washing.—HISP.—eight of deuto-chloruret and six of mercury, several successive sublimations, and washing with boiling water.—SPIELMANN—thirty-two ounces of deuto-chloruret and twenty-four of mercury, and two sublimations.—AMER.; and COXE—one pound of deuto-chloruret and nine ounces of mercury, and two successive ebullitions.

HASS.; and PIDERIT prescribe one pound of deuto-chloruret and ten ounces of mercury, and trituration with water; but direct the product after the first sublimation to be carefully triturated with an ounce of mercury, and the mixture to be sublimed afresh.

The washing, omitted by several pharmacopœias, is an important precaution for the freeing of the preparation from a little deuto-chloruret which always remains in it. (HENRI)

2. *Scheele's process.*

Calomelas præcipitatum. (DUBL., COXE)

R. Mercury, seventeen parts.
Dilute Nitrous Acid, fifteen parts.

Pour the acid upon the metal in a glass vessel, and at the end of the effervescence, digest in a moderate heat, shaking the vessel from time to time, for six hours. Then augment the heat till the liquor boils a little; pour it off then from the undissolved mercury, and immediately mix it with a boiling solution, previously prepared, of

Muriate of Soda, seven parts.
Water, forty parts.

Wash the precipitate which is formed with hot distilled water, till this is no longer made turbid with subcarbonate of potass; then dry it.

Submuriæ Hydrargyri præcipitatus. (EDINB.)

R. Dilute Nitrous Acid, } of each,
Purified Mercury, } eight
Muriate of Soda, } ounces.
four ounces and a half.
Boiling Water, eight pounds.

Mix the mercury with the acid, and towards the end of the effervescence, digest with a gentle heat, frequently shaking the vessel. It is necessary, however, that more mercury be mixed with the acid than it can dissolve, that a completely saturated solution may be obtained.

At the same time let the muriate of soda be dissolved in the boiling water; add to this solution the other while yet warm, and mix quickly together. When the precipitate shall have subsided, pour off the liquor, and wash the precipitated powder frequently with warm water, till this comes off tasteless.

The product is one of those commonly designated *White Precipitate*:

AMER.; AMST.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FENN.; FULD.; LUSIT.; PALAT.; ROSS.; SARD.; SUEC.; WIRTEM.; and SWEDIAUR give nearly the same formula. The following is a modification which deserves notice:

R. Crystallized Nitrate of Mercury, } of each,
Mercury, } an equal part.
Muriate of Soda,

Triturate in a glass mortar to the extinction of the metal, and sublime. (BRUGNATELLI)

3. *Hernbstaedt's process.*

R. Deuto-sulphate of Mercury, unwashed, obtained by distilling to dryness, in a retort, six ounces and a half of mercury and four of sulphuric acid,

Mercury, three ounces and a half.
Triturate to the extinction of the metal, and add

Muriate of Soda, five ounces and a half.

Mix by trituration, and sublime. (BRUGNATELLI; COXE; HENRI)

FERR. prescribes twelve parts of mercury, fourteen of sulphuric acid, pulverization of the salt with twelve parts of mercury, the mixture of eight parts of the mass with three of decrepitated muriate of soda, and sublimation.

Hydrargyri Submuriæ. (LOND.)

R. Purified Mercury, by weight, four pounds.
Sulphuric Acid, by weight, thirty ounces.

Muriate of Soda, one pound and a half.
Ammonia, eight ounces.

Boil two pounds of the mercury in a glass vessel with sulphuric acid, till dry sulphate of mercury is formed; when this is cold triturate it with the other two pounds of mercury in an earthen mortar till perfectly mixed: then add the muriate of soda, triturate again till all the metallic globules have disappeared, and then sublime. Reduce the sublimed mass to a very fine powder, pass it through a sieve, and mix it carefully with the muriate of ammonia previously dissolved in a gallon of boiling distilled

water. Set it apart that the powder may subside; then pour off the liquor, and wash the powder repeatedly with boiling distilled water, till solution of ammonia dropped into it effect no precipitate. Finally reduce it to a very fine powder in the manner directed for prepared chalk.

Calomelas Sublimatum. (DUBL.)

R. Persulphate of Mercury,
twenty-five parts.
Purified Mercury,
seventeen parts.
Dried Muriate of Soda,
ten parts.

Triturate together till the metal is extinguished, and sublime with a sufficient degree of heat. Reduce the sublimed mass to powder and sublime again. Then pulverize it and wash it frequently with distilled water till the decanted solution ceases to throw down a precipitate on the addition of a little carbonate of potass.

LIPP. directs four ounces each, of mercury and of sulphuric acid, to be mixed gradually together in a glass retort, and exposed to the heat of a sand-bath, till the mixture becomes converted into a white mass. The salt to be pulverized when cold, and triturated with two ounces and a half of mercury till the metal is extinguished; then four ounces and a half of decrepitated muriate of soda to be added, and the mixture sublimed; the product to be powdered and again sublimed. The mass to be again powdered and boiled for an hour in a solution of one ounce and a half of muriate of ammonia in five pounds of distilled water. The liquor to be then filtered, the powder to be washed in distilled water, and finally dried in bibulous paper.

The proto-chloruret of mercury is a preparation in very great use, particularly in England, where it seems to be considered as a sort of panacea. It is stimulant, alterative, anthelmintic, anti-spasmodic, depurative, diaphoretic, diuretic, errhine, purgative, resolvent, or sialagogue, according to the dose, the mode of application, the circumstances in which it is given, the constitution of the patient, and the substances with which it is combined. It has been extolled in syphilitic or pseudo-syphilitic diseases, scrofulous affections, chronic hepatitis, convulsive diseases, rheumatic affections, cutaneous diseases, and obstinate constipation.

In general, it only acts as an alterative, that is to say, as a revulsive stimulant or irritant, in the dose of from one to five grains; but in that of from five to fifteen grains, it acts as a purgative.

It is incompatible with chlorine, nitric acid, alkalis and their carbonates and sulphurets, lime water, copper, iron, and lead.

1. *Preparations in which the action of the Proto-chloruret of Mercury is not modified, at least sensibly, by any other substance.*

POWDER OF PROTO-CHLORURET OF MERCURY.

Pulvis e Mercurio dulci. (MIL. COPEN.; PH. SYPH.; SWEDIAUR)

R. Proto-chloruret of Mercury,
two grains.
Starch,
one scruple.

Mix. (MIL. COPEN.)

PH. SYPH. prescribes one or two grains of proto-chloruret, and fifteen grains of sugar.—SWEDIAUR—six grains of proto-chloruret, twelve of starch, and half a dram of white sugar.

All these powders may be administered at one dose. The following has been recommended in the syphilitic disorders of children:

R. Proto-chloruret of Mercury, one grain.
Sugar,
one scruple.

Mix, and divide into three doses; to be taken in the course of the day.

A mixture of two drams of proto-chloruret with half an ounce of sugar and the same quantity of olibanum may be proper in certain cases of syphilis, in fumigations, for each of which, from one to two drams of the powder may be used. Another mixture of four grains of proto-chloruret and of half a scruple of Armenian bole, divided into four doses for one day, has been proposed to be rubbed into the gums, internal surface of the lips, tongue, and palate, according to the method of CLARE and BRACHET.

VERMIFUGE POWDER.

Remedium anti-taniosum. (AUGUSTIN)

R. Proto-chloruret of Mercury,
one scruple.
Calcedined Hartshorn, } of each,
Red Sulphuret of Mercury, } ten grains.

Mix.—To be taken every four or five hours in the afternoon; in the evening, two ounces of oil of almonds are to be taken; on the following morning, fasting, in syrup, one third of a powder prepared with one dram of fern, half a dram of jalap, the same of gamboge, and the same of animal char-

coal. If the tænia is not expelled, the second third is to be taken at the end of two hours, and the last third two hours after.

This remedy has been proposed by BECK.

INJECTION OF PROTO-CHLORURET OF MERCURY.

Injectio ex Muriate Hydrargyri.
(PH. SYPH.)

- R. Proto-chloruret of Mercury, half an ounce.
Distilled Water, eight ounces.

Mix.—It is recommended in acute urethritis, but is most useful in certain cases of chronic urethritis. LAGNEAU gives a preferable formula:

- R. Proto-chloruret of Mercury, levigated, half an ounce.
Solution of Gum Arabic, twelve ounces.

Mix.

ANTI-SYPHILITIC GARGLE. (PH. SYPH.)

- R. Gum Arabic, one dram.
Decoction of Barley, five ounces.

Sweeten the solution with
Syrup of Marshmallows,
one ounce and a half,

and add

Proto-chloruret of Mercury,
fifteen grains.

MERCURIAL LAVEMENT.

Enema mercuriale. (AUGUSTIN)

- R. Proto-chloruret of Mercury, one scruple.
Starch, } of each,
Gum Arabic, } half an ounce.
Water, } eight ounces.

Mix by trituration.

ANTI-PSORIC LOTION.

Lotio Veratri mercurialis seu anti-psorica.
(AUGUSTIN)

- R. Decoction of White Hellebore, eight ounces.
Proto-chloruret of Mercury, one dram.
Nitrate of Potass, two drams.
Muriate of Ammonia, two ounces.

This is a bad preparation; but WILSON has extolled it as an excellent anti-psoric.

MIXTURE WITH PROTO-CHLORURET OF MERCURY. (SAUNDERS)

- R. Proto-chloruret of Mercury, half a scruple.
Syrup of Fumitory, two ounces.
Fumitory Water, one pound.

VOL. II.

Mix.—To be taken by half ounces or ounces.

HONEY WITH PROTO-CHLORURET OF MERCURY.

Mel e Muriate Hydrargyri seu Hydrargyri.
(PH. SYPH.; SWEDIAUR)

- R. Proto-chloruret of Mercury, one or two drams.
Honey, one ounce.

Triturate.—It is recommended in syphilitic ulceration of the throat; and is frequently preferable to ointment, for ulcers of the genital organs.

OINTMENT OF PROTO-CHLORURET OF MERCURY. (PH. SYPH.)

- R. Proto-chloruret of Mercury, one or two drams.
White Cerate, or Beef Marrow, one ounce.

Triturate for a quarter of an hour.

It is recommended in syphilitic diseases, in frictions to the skin; or in applications to diseased parts.

NIEMANN prescribes, under this title, an ointment composed of two scruples of proto-chloruret, with one ounce of resinous ointment; which MARRIANUS extolled in leprosy and in various pustular diseases of the skin.

ANTI-HERPETIC OINTMENT. (RATIER)

- R. Proto-chloruret of Mercury, one dram.
Rose Ointment, three drams.

Mix.—This ointment is prescribed by DUPUYTREN. Under the title of *English anti-herpetic Ointment*, PIERQUIN prescribes one dram of proto-chloruret, and four ounces of lard or of unguentum populeum.

- R. Proto-chloruret of Mercury, one dram.
Sublimed Sulphur, two drams.
Prepared Lard, one ounce.

It is recommended by FOUQUIER.

INJECTION AGAINST CARNOSSITIES OF THE URETHRA. (SAINTÉ-MARIE)

- R. Proto-chloruret of Mercury, one dram.
Oil of Scorpions, one ounce.
Balsamum tranquillum, two ounces.

Mix.—Having voided the urine, the volume of a large die of this mixture is to be injected into the urethra, and there retained during five minutes at least; and the operation is to be repeated thrice a day, for a month. Though this strange injection is useless against such imaginary things as carnosities of the urethra, yet it might be of service in some cases of obstinate chronic urethritis.

H

BOLUS OF PROTO-CHLORURET OF MERCURY.
(SWEDIAUR)

- R. Proto-chloruret of Mercury,
from five to ten grains.
Simple Syrup, a sufficient quantity
to make with powdered lycopodium a bolus.
It is said to be useful in hydrocephalus.

TONIC ANTHELMINTIC BOLUSES. (BRERA)

- R. Proto-chloruret of Mercury,
four grains.
Extract of Wormwood, one dram.
Make four boluses, to be taken at one dose.
- R. Proto-chloruret of Mercury, four grains.
Extract of Artemisia Seeds, one dram.
Make four boluses, one to be taken every two hours.
- They are recommended in cases of worms, where there is slight atony of the intestinal canal.

NERVINE ANTHELMINTIC BOLUSES.
(BRERA)

- R. Proto-chloruret of Mercury,
from four to eight grains.
Extract of Valerian, two scruples.
Make four boluses.—Dose, one every three hours, in verminal disorders accompanied with nervous symptoms.

PILLS OF PROTO-CHLORURET OF MERCURY.

Pilule ex Hydrargyro muriatico miti seu ex Mercurio dulci. (MIL. COPEN.: WURCEB.; AUGUSTIN; ELLIS; RATIER)

- R. Proto-chloruret of Mercury,
twelve grains.
Conserve of Roses, a sufficient quantity
to make twelve pills. (ELLIS)

- R. Proto-chloruret of Mer- } of each,
cury, } one dram.
Extract of Aconite, }
Simple Syrup, a sufficient quantity.

Make one hundred and twenty pills. (AUGUSTIN)

- R. Proto-chloruret of Mer- } of each,
cury, } half a dram.
Crumb of Bread, }
Water, a sufficient quantity

to make thirty pills. (MIL. COPEN.)

- R. Proto-chloruret of Mer- } of each,
cury, } an
Powdered Marshmallows, } equal part.
Syrup, a sufficient quantity.

Make pills of two grains each. (RATIER)

- R. Proto-chloruret of Mercury,
fifteen grains.

Extract of Liquorice,
one dram and a half.
Water, a sufficient quantity

to form a mass to be divided into sixty pills.
(WURCEB.)

The dose of these pills must be varied according to the effect desired to be produced, the constitution of the patient, and their own composition.

ANTI-ICTERIC PILLS. (CADET DE GAS-SICOURT)

- R. Proto-chloruret of Mercury, one dram.
Extract of Soap-wort, two drams.
Make a mass to be divided into seventy-two pills; two to be taken three or four times a day.

RESOLVENT PILLS. (BRERA)

- R. Proto-chloruret of Mercury,
four grains.
Extract of Dandelion, eighteen grains.
Make four pills, to be taken within twenty-four hours, in abdominal obstructions.

ANTI-SYPHILITIC PILLS.

Pilule anti-syphiliticæ et resolventes.
(BRERA)

- R. Proto-chloruret of Mercury,
four grains.
Extract of Bitter-sweet,
a sufficient quantity.
Make four pills, to be taken in the course of the day.

EMMENAGOGUE BOLUSES.

Bolus ex Sabina. (AUGUSTIN)

- R. Proto-chloruret of Mercury, two grains.
Extract of Savine, four grains.
Syrup of Red Poppies,
a sufficient quantity.

Make one bolus.—PERFECT has found it serviceable in the mania caused by amenorrhœa.

ALTERATIVE PILLS. (SAUNDERS)

- R. Proto-chloruret of Mercury, one dram.
Electuary of Scordium, two drams.
Liquorice Powder, a sufficient quantity
to make seventy-two pills.

PURGATIVE PILLS. (AMER.; COXE)

- R. Proto-chloruret of Mercury,
half a dram.
Soap, one scruple.
Water, a sufficient quantity.

Make forty pills.

This preparation should be rejected on account of the soap being incompatible with the proto-chloruret.

VIGAROUS' PILLS. (BORIES; PIERQUIN; SAINTE-MARIE)

- R. Mercury, four drams.
Deuto-chloruret of Mercury, two drams

Triturate together, adding a few drops of lemon juice, till the metal is extinguished; then add

Wheat Flower, five ounces.

Make, with lemon juice, a mass to be divided into pills of two grains each.

This is a bad preparation; and the uncertainty of the decomposition which takes place must render it a very hazardous one.—VIGAROUS gave first one pill a day, gradually augmenting the dose till the patient took three morning and evening.

BOLUSES AGAINST HOOPING-COUGH.

Boli in Pertussi. (BRERA)

R. Proto-chloruret of Mercury, four grains.
 Extract of Belladonna, two grains.
 Turpentine, six grains.
 Honey, } of each,
 Liquorice Powder, } a sufficient quantity

to make eight boluses.—Dose, one every two hours, for a child of from eight to ten years.

ANTI-SYPHILITIC PILLS. (PIERQUIN)

R. Mercury, } of each,
 Proto-chloruret of } half an ounce.
 Mercury, }
 Root of Pyrethrum, } of each,
 Senna, in powder, } one ounce.
 White Agaric, }
 Honey, a sufficient quantity

to make pills of six grains each. To these may also be applied the remarks which we made on VIGAROUS' Pills.

VERMIFUGE PASTILLES. (SPIELMANN)

R. Proto-chloruret of Mercury, two ounces.
 Sugar, boiled to a proper consistence with the Juice of Barberries, four ounces.
 Mix, and make pastilles.

BARTHEZ' VERMIFUGE PASTILLES. (BORRIES; CADET DE GASSICOURT; PIERQUIN)

R. Proto-chloruret of Mercury, three drams.
 White Sugar, one pound.
 Mucilage of Gum Arabic, a sufficient quantity

to make pastilles of the size of a shilling.—Dose, one or two to children, and six or eight to adults.

CHING'S WORM LOZENGES.

Trochisci anthelmintici Doctoris Ching.
 (AUGUSTIN)

R. Proto-chloruret of Mercury, two drams.
 White Sugar, ten drams.
 Tincture of Saffron, twenty drops.
 Mucilage of Tragacanth, a sufficient quantity.
 Make a hundred and twenty lozenges.—Dose, two or three.

VERMIFUGE COMFITS. (PIERQUIN)

R. Proto-chloruret of Mercury, half an ounce.
 Sugar, one ounce.
 Starch, half an ounce.
 Mucilage of Tragacanth, } of each,
 Essence of Bergamotte, } a sufficient quantity.
 Make one hundred and forty-four comfits.

II. *Preparations in which the action of Proto-chloruret of Mercury is more or less modified by foreign substances.*

A.

By Aloes.

PURGATIVE ELECTUARY. (PIERQUIN)

R. Proto-chloruret of Mercury, } of each,
 Extract of Aloes, } half a scruple.
 Subcarbonate of Potass, two grains.
 Syrup of Fumitory, a sufficient quantity.

Mix.—This preparation is bad, from the alkaline salt being incompatible with the proto-chloruret.

ANTHELMINTIC PILLS. (FULD.; HAMB.; AUGUSTIN)

R. Proto-chloruret of Mercury, } of each,
 Aqueous Extract of Aloes, } one scruple.
 Extract of Walnut, } of each,
 Assafœtida, } four scruples.
 Tincture of Tansey, a sufficient quantity

to make a pilular mass. (FULD.; HAMB.)

R. Proto-chloruret of Mercury, } of each,
 Socotrine Aloes, } one dram.
 Resin of Jalap, }
 Tincture of Rhubarb, a sufficient quantity

to make a pilular mass. (AUGUSTIN)

They are anthelmintic, anti-spasmodic, resolvent, and laxative.—Dose, from six to twenty grains.

ANTHELMINTIC PILLS.

Pilule anthelminticæ purgantes. (FULD.)

- R. Proto-chloruret of Mercury, three grains.
 Aqueous Extract of Aloes, } of each,
 Extract of Rhubarb, } six grains.
 ————— Tansey,
 Tincture of Tansey, a sufficient quantity.
- Mix.—To be taken at once; for an adult.

DRASTIC BOLUSES. (BORIES; PIERQUIN)

- R. Proto-chloruret of Mercury, fifteen grains
 Socotrine Aloes, } of each, six grains.
 Scammony, } fifteen grains.
 Rhubarb, } a sufficient quantity.
 Pulp of Cassia,

Mix and divide into proper doses.

These are recommended in rheumatic affections, on account of the salutary derivation which they sometimes produce by determining alvine evacuations and provoking hæmorrhoidal discharges.

PROPHYLACTIC PILLS FOR SMALL-POX.

Pilule ad Variolam. (PIERQUIN; SPIELMANN)

- R. Proto-chloruret of Mercury, } of each,
 Extract of Aloes, } twelve grains.
 Camphor, } eight grains.
 Guaiac Resin, } sixteen grains.

Make a mass to be divided into pills of two grains each.

ROSENSTEIN recommends them as a preservative against the small pox, and also as a preparative to inoculation.—The dose varies according to the age.

B.

By Antimonial preparations.

SIMPLE ALTERATIVE OR PLUMMER'S POWDER.

Pulvis alterans Plummeri seu Edinburgensis seu ex Hydrargyro muriatico miti cum Sulphure Stibii aurantiaco seu Hydrargyri stibiatus. (BAVAR.; BRUNS.; DAN.; MIL. COPEN.; FERR.; FORM. HÔP. FR.; FULD.; HAMB.; HANN.; LIPP.; CAST. BORUS.; ROSS.; WIRTEM.; AUGUSTIN; CADET DE GASSICOURT; HUFELAND; NIEMANN; RATIER; SPIELMANN)

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } an equal
 of Antimony, } part.

Triturate for a long time to obtain a very fine powder. (BAVAR.; BRUNS.; DAN.;

FERR.; HAMB.; HANN.; LIPP.; ROSS.; WIRTEM.; CADET DE GASSICOURT; SPIELMANN)

- R. Proto-chloruret of Mercury, one part.
 Sulphuretted Hydrosulphate of Antimony, two parts.
 Mix by trituration. (FULD.)

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } one
 of Antimony, } part.
 White Sugar, nine parts.

Triturate together. (RATIER)

HUFELAND prescribes half a scruple each, of proto-chloruret and of sulphuretted hydrosulphate and three drams of sugar.

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } one
 of Antimony, } part.
 Gum Arabic, ten parts.

Triturate together. (FORM. HÔP. FR.)

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } one
 of Antimony, } part.
 White Sugar, } of each, ten parts.
 Starch,

Triturate together. (MIL. COPEN.)

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } one
 of Antimony, } part.
 Gum Arabic, five parts.
 White Sugar, eight parts.

Triturate together. (NIEMANN)

AUGUSTIN prescribes one part of proto-chloruret, one of sulphuretted hydrosulphate, five parts of gum, and ten of sugar.

Whatever celebrity this preparation may have formerly possessed as resolvent and alterative, in scrofulous and syphilitic diseases, very little confidence deserves to be placed in it. The decomposition which results between two principal components renders it a very uncertain one, as well as every other into which it enters.—Dose, from six to ten grains daily, of the simple mixture at equal parts.

SIMPLE ALTERATIVE, OR PLUMMER'S PILLS.

Pilule Plummeri seu ex Stibio et Hydrargyro seu ex Hydrargyro muriatico miti cum Sulphure stibiato aurantiaco. (FULD.; GENEV.; HAMB.; LUSIT.; WURCEB.; AUGUSTIN; CADET DE GASSICOURT; NIEMANN; RATIER; SPIELMANN; SWEDIAUR)

- R. Proto-chloruret of Mercury, } of each,
 Sulphuretted Hydrosulphate } an equal
 of Antimony, } part.
 Extract of Liquorice, a sufficient quantity

to make a mass to be divided into pills of three grains each. (SWEDIAUR)

CADET DE GASSICOURT prescribes three ounces each, of proto-chloruret and sulphuretted hydrosulphate, two drams of extract of liquorice, and a sufficient quantity of mucilage of gum arabic to make a mass to be divided into pills of six grains each.—AUGUSTIN—fifteen grains each, of proto-chloruret and of sulphuretted hydrosulphate and one dram of extract of liquorice, to make pills of two grains each.—NIEMANN—two drams each, of proto-chloruret and of sulphuretted hydrosulphate, four scruples of extract of liquorice, and a sufficient quantity of mucilage of gum arabic.—HAMB.—half a dram of sulphuretted hydrosulphate fifteen grains of proto-chloruret of mercury, one dram of extract of wild mustard, and a sufficient quantity of liquorice powder to form a mass to be divided into forty-five pills.—SPIELMANN—six drams each, of proto-chloruret and of sulphuretted hydrosulphate, and four drams of extract of gentian.—LUSIT.—the same proportions but the extract of liquorice instead of that of gentian.—WURCEB.—fifteen grains each, of proto-chloruret and of sulphuretted hydrosulphate, and one dram and a half of extract of hemlock, to make sixty pills.—RATIER—half a dram each, of proto-chloruret and of sulphuretted hydrosulphate, half an ounce of extract of hemlock, and a sufficient quantity of powdered hemlock to make a mass to be divided into thirty-six pills.—GENEV.—one scruple each, of proto-chloruret and of sulphuretted hydrosulphate, and a sufficient quantity of rob of elder, to make twenty-four pills.

R. Proto-chloruret of Mercury, fifteen grains.
Sulphuretted Hydrosulphate of Antimony, half a dram.
Extract of Hemlock, two drams.
Copaiba, a sufficient quantity
to make a pilular mass. (FULD.)

To these pills may be compared those which SAINTE-MARIE gives under the title of *Diaphoretic Pills*, and which are prepared with two scruples each, of proto-chloruret of mercury and of proto-hydrosulphate of antimony, and a sufficient quantity of syrup; the mass being divided into thirty pills.

FINKE'S RESOLVENT PILLS. (SAINTE-MARIE)

R. Simple PLUMMER'S Powder, twelve grains.
Extract of Dandelion, three grains.
Gum Ammoniac, two grains.
Squill, in powder, half a dram.
Make pills of three grains each.

Five to be taken three or four times a day, taking after each dose, an infusion of cichory. These pills have appeared useful

after intermittent fevers, in engorgements of the abdominal viscera accompanied with swelling of the legs.

COMPOUND ALTERNATIVE, OR PLUMMER'S POWDER. (HASS.; HERBIP.; PIDERIT)

R. Proto-chloruret of Mercury, two drams.
Sulphuretted Hydrosulphate of Antimony, one dram.
Guaiaic Resin, one dram and a half.

Make a powder. (HASS.; PIDERIT)

HERBIP. directs an equal part to be taken of each of the three substances, and the resin to be triturated with a quantity of sugar equal to its own.

BASILIC POWDER.

Pulvis basilicus. (SAUNDERS)

R. Proto-chloruret of Mercury, } of each,
Deutoxide of Antimony, } five
Scammony, } grains.
Supertartrate of Potass, }

Mix.

COMPOUND ALTERNATIVE, OR PLUMMER'S PILLS.

Pilulæ Plummeri seu Submuriatis Hydrargyri compositæ seu resolventes cum Mercurio seu depurantes. (LOND.; EDINB.; DUBL.; AMER.; HASS.; CAST. BORUS.; ROSS.; SARD.; WIRTEM.; HERBIP.; AUGUSTIN.; NIEMANN.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

Pilulæ Submuriatis Hydrargyri compositæ. (LOND.; EDINB.)

Pilulæ Calomelanos compositæ. (DUBL.)

R. Submuriate (Proto-chloruret) of Mercury, } of
Precipitated Sulphuret } each,
(Sulphuretted Hydro- } one
sulphate) of Antimony, } dram.
Guaiaic Resin, two drams.
Mucilage of Gum Arabic, a sufficient quantity.

Triturate the two first together, add the resin, and finally the mucilage, till a pilular mass is formed.

AMER.; and COXE give the same formula.

R. Simple PLUMMER'S Powder, half an ounce.
Guaiaic Resin, four drams.
Copaiba, a sufficient quantity
to make a mass, one dram of which is to be divided into twelve pills. (WIRTEM.)

- R. Copaiba, one part.
Guaiac Resin, twelve parts.

Triturate together, adding gradually,

- | | | |
|-----------------------------|-----------------------|------------|
| Proto-chloruret of Mercury, | } of each, | six parts. |
| Sulphuretted Hydrosulphate | | |
| of Antimony, | | |
| Alcohol, | a sufficient quantity | |

to make a pilular mass. (VAN MONS)

- R. Proto-chloruret of Mercury, two grains.
Sulphuretted Hydrosulphate of Antimony, one dram.
Guaiac Resin, } of each, one dram
Extract of Gentian, } and a half.

Mix. (ROSS.)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } two
of Antimony, } drams.
Guaiac Resin, } one dram.
Myrrh, }
Balsam of Tolu, a sufficient quantity.

Make pills of five grains each. (SWEDIAUR)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } six
of Antimony, } drams.

Triturate together, and add

- | | | |
|-------------------------|-----------------------|--------------|
| Guaiac Resin, | } of each, | three drams. |
| Extract of Liquorice, | | |
| Mucilage of Gum Arabic, | a sufficient quantity | |

to make a pilular mass. (HERBIP.)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } two
of Antimony, } scruples.
Extract of Bitter-sweet, } of each,
Guaiacine Soap, } half an ounce.

Mix, and make pills of two grains each. (CAST. BORUS.)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } an
of Antimony, } equal
Guaiac Resin, } part.
Extract of Fumitory, } of each,
Copaiba, } a sufficient quantity.

Make pills of four grains each. (AUGUSTIN)

- R. Proto-chloruret of Mercury, two drams.
Sulphuretted Hydrosulphate of Antimony, one dram.
Guaiac Resin, one dram and a half.
Alcohol, a sufficient quantity.

Make pills of two grains each.—Dose, from one to four. (AUGUSTIN)

- R. Guaiacine Soap, } of each,
Extract of Bitter-sweet, } half an ounce.
Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } two
of Antimony, } scruples.

Mix.—Dose, from fourteen to thirty grains, morning and evening. (AUGUSTIN)

- R. Guaiacine Soap, one dram and a half.
Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } fifteen
of Antimony, } grains.
Root of Seneka,
Camphor,
Vinegar of Squill, a sufficient quantity.

Mix.—To be taken in the same dose as the preceding. (AUGUSTIN)

ANTHELMINTIC PILLS. (AUGUSTIN)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate of Antimony, } one
Compound Extract of Rhu- } dram.
barb,
Assafœtida,

Mix.—Dose, six grains, in the morning, augmenting it by three grains every day.

RESOLVENT PILLS. (SWEDIAUR)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } one
of Antimony, } dram.
Sulphate of Iron,
Myrrh,
Simple Syrup, a sufficient quantity.

Mix, and divide into pills of three grains each.

They are recommended in engorgements of the abdominal viscera supervening after quartan fever.—Dose, four, morning and evening.

- R. Proto-chloruret of Mercury, half a dram.
Sulphuretted Hydrosulphate } of each,
of Antimony, } one
Camphor, } scruple.
Guaiac Resin, half an ounce.
Extract of Burnet Saxifrage, } of each,
Hemlock, } one dram.

Make pills of two grains each. (SAUNDERS)

ANTI-HERPETIC PILLS. (BORIES)

- R. Proto-chloruret of Mercury, ten grains.
Sulphuretted Hydrosulphate of Antimony, five grains.
Black Sulphuret of Mercury, sixty grains.
Extract of Scabious, thirty grains.
Syrup of Fumitory, a sufficient quantity.

Make pills of four grains each.

Dose, three in the morning fasting, to be gradually augmented till the patient takes six daily at two doses.

KEMPF'S ANTI-SYPHILITIC PILLS.
(SAINTE-MARIE)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } two
of Antimony, } scruples.

Triturate together for a long time, and add

Extract of Soap-wort, } of each,
Guaiac Resin, } four scruples.
Camphor, } half a dram.

Incorporate the whole with
Theriac, a sufficient quantity.

Dose, twelve grains thrice a day with a strong decoction of china, burdock, or sarsaparilla root, augmenting gradually till the patient takes doses of twenty-four grains.

ANTI-SPASMODIC POWDER.

Pulvis Hydrargyri stibiatus cum Hyosciamo.
(AUGUSTIN)

- R. Proto-chloruret of Mer- } of each,
cury, } one grain.
Extract of Henbane, }
Sulphuretted Hydrosulphate of An- }
timony, } three grains.
Sugar, } half a dram.

Mix.—To be taken morning and evening, in inflammatory, asthenic, and spasmodic diseases.—It has been particularly useful in nyctalopia.

STHENIC PILLS. (CADET DE GASSICOURT)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } ten
of Antimony, } grains.
Opium, }
Proto-hydrosulphate of Antimony, } fifteen grains.
Balsam of Peru, a sufficient quantity.

Make pills of one grain each.

They are recommended in phthisis pulmonalis, and in dropsical and spasmodic affections.

COMPOUND ANTIMONIAL PILLS. (BRUGNATELLI)

- R. Proto-chloruret of Mercury, } of each,
Prepared Sulphuret of Anti- } three
mony, } drams.
Extract of Gentian, } of each,
Soap, } one dram.

Triturate the two first together, then add the extract, and finally the soap previously softened.

It is a bad preparation.

ANTI-SYPHILITIC PILLS. (SAINTE-MARIE)

- R. Proto-chloruret of Mercury, } of each,
Proto-hydrosulphate of An- } two
timony, } drams.

Guaiac Resin, four drams.
Peruvian Balsam, a sufficient quantity
to make a mass, each dram of which is to be divided into twelve pills.

They are extolled in chronic diseases of the skin reputed syphilitic.—Dose, one thrice a day, gradually augmenting one daily, till six are taken in twenty-four hours.

RESOLVENT PILLS. (BRERA)

- R. Proto-chloruret of Mercury, six grains.
Proto-hydrosulphate of Antimony, twelve grains.
Rob of Elder, a sufficient quantity.

Make six pills.—Dose, one every two hours, towards the end of pneumonia, and in the symptomatic cough from worms in children.

SWEDISH PILLS. (RATIER)

- R. Proto-chloruret of Mercury, one dram and a half.
Proto-hydrosulphate of Antimony, } of each,
Black Sulphuret of Mer- } one dram.
cury, }
Crumb of Bread, a sufficient quantity
to make one hundred and forty-four pills.

They are anti-syphilitic.

LAXATIVE MERCURIAL POWDER.

Pulvis e Mercurio. (GENEV.)

- R. Proto-chloruret of Mercury, two ounces.
Supertartrate of Potass, } of each,
Tartarized Antimony, } two scruples.
Sugar, } fifteen drams.
Subcarbonate of Magnesia, five ounces.

Mix.

DIAPHORETIC POWDER. (ELLIS)

- R. Proto-chloruret of Mercury, } of each,
Sulphuretted Hydrosulphate } six
of Antimony, } grains.
Guaiac Resin, } one dram.

Make six parcels.—One to be taken morning and evening in disorders of the skin.

STIMULANT RESOLVENT BOLUSES.

Boli resolventes et incitantes. (BRERA)

- R. Proto-chloruret of Mercury, sixteen grains.
Proto-hydrosulphate of Antimony, eight grains.
Camphor, thirty grains.
Rob of Elder, a sufficient quantity.

Make ten boluses.—Dose, one every two

hours in visceral inflammations with tendency to asthenia.

COMPOUND PILLS OF PROTO-CHLORURET OF MERCURY AND OF OPIUM. (AMER.; PH. SYPH.; COXE)

- R. Proto-chloruret of Mercury, half a dram.
 Opium, fifteen grains.
 Tartarized Antimony, four grains.
 Conserve of Dog Rose, a sufficient quantity
 to make fifteen pills. (PH. SYPH.)

AMER.; and COXE prescribe two drams of proto-chloruret, one dram of opium, one scruple of tartarized antimony, and a sufficient quantity of simple syrup, to make sixty pills,

HYDRAGOGUE PILLS. (SWEDIAUR)

- R. Proto-chloruret of Mercury, } of each,
 Deutoxide of Antimony, } five
 Scammony, } grains.
 Make three pills, for one dose.

EXPECTORANT PILLS. (ELLIS)

- R. Proto-chloruret of Mercury, twenty-four grains.
 Squill, half a dram.
 Tartarized Antimony, six grains.
 Opium, eighteen grains.
 Make twenty-four pills.—Dose, one every evening, in catarrhal affections.

JANIN'S HYDRAGOGUE PILLS. (CADET DE GASSICOURT; HUFELAND)

- R. Water, six pounds.
 Jalap, }
 Aloes, } of each, one pound.
 Senna, }
 Subcarbonate of Iron, } of each,
 Nitrate of Potass, } eight ounces
 and a half.
 Agaric, }
 Mechoacan, } of each,
 Rhubarb, } six ounces
 Scammony, } two drams.
 Root of Heath, (*Calluna*
vulgaris, PERSOÏN) }
 Hermodactyls, }
 Black Sulphuret of Mercury, five ounces.
 Supertartrate of Potass, two ounces one dram and a half.
 Gamboge, }
 Alhandal Trochiscs, } of each,
 Proto-chloruret of Mercury, } two
 Tartarized Antimony, } ounces.

Boil the senna and supertartrate of potass in the water for a quarter of an hour, and express strongly; to the liquor heated in an

iron vessel add the other substances finely powdered and intermixed, stirring continually over a slow fire till a pilular mass is obtained; divide this into pills of four grains each, roll them in powdered jalap, and dry them.

This is a monstrous assemblage; and it is rather strange that HUFELAND should have given it a place in his *Pharmacopœia for the Poor*, where he speaks of it in terms of praise.—Dose, four pills at first, augmenting it according to the effect produced.

C.

By *Assafoetida*.

VERMIFUGE POWDER.

Pulvis Gratiolæ cum Hgdrargyro. (AUGUSTIN)

- R. Herb of Gratiola, two scruples.
 Proto-chloruret of Mercury, five grains.
 Assafoetida, half a dram.
 Oil of Peppermint, three drops.

Divide into ten parcels.—Dose, one every three hours. It is extolled by HARGENS against tænia.

ANTHELMINTIC PILLS. (AUGUSTIN; SPIELMANN; VAN MONS)

- R. Proto-chloruret of Mercury, one part.
 Assafoetida, three parts.
 Sulphate of Iron, two parts.
 Simple Syrup, a sufficient quantity.
 Mix. (VAN MONS)

- R. Proto-chloruret of Mercury, } of each,
 Assafoetida, } one
 Rhubarb, } dram.
 Sagapenum, }
 Tincture of Gentian, a sufficient quantity.

Make pills of one grain each. (SPIELMANN)

They are recommended against ascarides and tænia.

- R. Extract of Rhubarb, } of each,
 Walnut, } half
 Assafoetida, } a dram.
 Proto-chloruret of Mercury, }

Make pills of one grain each.—Dose, from four daily. (AUGUSTIN)

ANTHELMINTIC AND ANTI-SPASMODIC BOLUSES. (BRERA)

- R. Proto-chloruret of Mercury, six grains.
 Assafoetida, one scruple.
 Extract of Valerian, two scruples.

Make eight boluses.—Dose, one every two hours, in verminal disorders accom-

panied with epilepsy, chorea, or spasmodic cholera.

RESOLVENT PILLS (SAUNDERS)

- R. Assafœtida, one dram.
 Powdered Acorns, } of each,
 Black Sulphuret of Mer- } half
 cury, } a dram.
 Proto-chloruret of Mercury, five grains.
 Syrup of Poppies, } of each, a
 Liquorice Powder, } sufficient quantity.

Make pills of two grains each.

EMMENAGOGUE PILLS. (PIERQUIN)

- R. Extract of Dandelion, } of each,
 Soapwort, } one dram.
 Sub-borate of Soda, } of each,
 Assafœtida, } half a dram.
 Black Sulphuret of } of each,
 Mercury, } twenty
 Proto-chloruret of Mercury, } grains.
 Syrup of Ground Ivy, a sufficient quantity.

D.

By Copaiba.

PILLS TO TERMINATE GONORRHEA.

(CADET DE GASSICOURT; PIERQUIN;
 RATIER)

- R. Proto-chloruret of Mercury, one dram.
 Conserve of Red Roses, four ounces.
 Dragon's Blood, half an ounce.
 Copaiba, one ounce.

Make pills of six grains each.

These have been extolled in chronic gonorrhœa which remains from atony.—Dose, one, then two and three, morning and evening, with ferruginous water for drink.

E.

By Catechu.

ASTRINGENT PILLS. (SAINTE-MARIE)

- R. Proto-chloruret of Mercury, one scruple.
 Catechu, } of each, five drams.
 Copaiba, }
 Syrup of Comfrey, a sufficient quantity

to make one hundred and fifty pills.—Dose, four, three times a day in suspected leucorrhœa, and in atonic blennorrhœa.

- R. Proto-chloruret of Mercury, eight grains.
 Catechu, one scruple.
 Crab's Eyes, two scruples.
 Copaiba, half a scruple.
 Syrup of Cinchona, a sufficient quantity

to make pills of four grains each. MORAND
 Vol. II.

prescribed them in atonic mucous discharges.—Dose, one or two, thrice a day.

- R. Proto-chloruret of Mercury, one scruple.
 Catechu, two drams.
 Gum Arabic, four scruples.
 Turpentine, a sufficient quantity

to make one hundred and fifty pills. QUARIN prescribed these pills in leucorrhœa and blennorrhœa, in the dose of four, thrice a day.

F.

By Camphor.

STIMULANT POWDER.

Pulvis Hydrargyri cum Camphora. (AUGUSTIN)

- R. Proto-chloruret of Mercury, twenty-four grains.
 Camphor, } of each,
 Prepared Oyster Shells, } two scruples.

Divide into eight parcels.—Dose, three or four daily, to excite the activity of the absorbent system, in dropsies.

PILLS OF PROTO-CHLORURET OF MERCURY
 AND OF CAMPHOR.

Pilule Muriatis Hydrargyri camphoratae.
 (SUEC.; RATIER; VAN MONS)

- R. Proto-chloruret of Mercury, one grain.
 Camphor, two grains.
 Simple Syrup, a sufficient quantity
 to make a pill with liquorice powder. (CADET DE GASSICOURT)

- R. Proto-chloruret of Mercury, two drams.
 Camphor, triturated with a few
 drops of Alcohol, one dram.
 White Sugar, three drams.
 Mucilage of Gum Arabic, a sufficient quantity

to make two hundred and forty pills. (SUEC.)

VAN MONS prescribes eight parts of proto-chloruret, four of camphor, twelve of sugar, one of tragacanth, and a sufficient quantity of water to make a mass.

G.

By Cinnamon.

PURGATIVE AND TONIC BOLUSES. (BO-
 RIES; PIERQUIN)

- R. Proto-chloruret of Mer- } of each,
 cury, } ten grains.
 Cinnamon, }
 Hermodactyls, one dram.
 Syrup of Peach Flowers, a sufficient quantity

to make ten boluses

H.

By Colocynth.

MERCURIAL PILLS. (BRUNS.; SARD.)

- R. Proto-chloruret of Mercury, half an ounce.
Compound Extract of Colocynth, one ounce and a half.
Mix. (BRUNS.)
- R. Proto-chloruret of Mercury, half an ounce.
Pills of Colocynth, with Aloes, one ounce.
Syrup of Buckthorn, a sufficient quantity.
Mix. (SAUNDERS)

I.

By Black Sulphuret of Mercury.

PURGATIVE BOLUSES. (PIERQUIN)

- R. Proto-chloruret of Mercury, fifteen grains.
Black Sulphuret of Mercury, ten grains.
Pulp of Cassia, a sufficient quantity.
To be taken in the morning, fasting.

ANTI-SYPHILITIC PILLS. (SAINTE-MARIE)

- R. Proto-chloruret of Mercury, } of each,
Black Sulphuret of Mer- } one
cury, } dram.
Linseed Meal, } of each, a
Mucilage of Gum } sufficient quantity
Arabic, }

to make thirty-six pills.

They are prescribed in the second stage of urethritis.—Dose, two daily, and a purgative every five days.

ANTI-LACTEAL PILLS. (BORIES)

- R. Proto-chloruret of Mercury, half a dram.
Black Sulphuret of Mercury, one dram.
Sulphate of Potass, half an ounce.
Nitrate of Potass, one dram.
Camphor, half a dram.
Resin of Jalap, one scruple.
Gum Arabic, one dram.
Syrup, a sufficient quantity

to make pills of four grains each.

Two to be taken morning and evening, several hours before and after taking food.

ANTI-SYPHILITIC ELECTUARY. (PIERQUIN)

- R. Proto-chloruret of Mercury, two ounces and a half.
Black Sulphuret of Mercury, two ounces.
Soap, eight ounces.
Mucilage of Tragacanth, a sufficient quantity.

Make a homogeneous mass, and add,

Extract of Borage, }	of each,
Camphor, Bugloss, }	three ounces.
Simple Syrup,	half a dram.
	a sufficient quantity.

Mix carefully.

It is recommended in syphilitic disorders complicated with scrofula.—Dose, from half a dram to one dram, in pills or in an appropriate liquid.

J.

By Iron.

FERRATED PROTO-CHLORURET OF MERCURY.

Murias Hydrargyri ferratus, Mercurius dulcis martiatus. (CADET DE GASSI-COURT; SWEDIAUR)

- R. Proto-chloruret of Mercury, six parts.
Black Oxide of Iron, one part.
Triturate together, and sublime. (SWE-DIAUR)

- R. Proto-chloruret of Mercury, three parts.
Hydrochlorate of Ammonia and of Iron, one part.

Triturate together. (CADET DE GASSI-COURT; SWEDIAUR)

This preparation is recommended by HARTMANN in scrofulous disorders, lymphatic engorgements, and vermical affections.—Dose, from twenty to thirty grains.

PURGATIVE AND TONIC ELECTUARY. (PIERQUIN)

- R. Proto-chloruret of Mer- } of each,
cury, } two drams.
Powdered Wood-lice, }
Sub-trito-carbonate of } of each,
Iron, } half an ounce.
Powdered Cinchona, }
Extract of Juniper, a sufficient quantity.

Mix.

K.

*By Guaiac Resin.**Errhine Powder.* (AUGUSTIN)

- R. Proto-chloruret of Mercury, ten grains.
Camphor, } of each, five grains.
Guaiac Resin, }
Sugar Candy, half a dram.
Cinchona, one scruple.

Mix, and divide into twenty parcels.

- R. Proto-chloruret of Mercury, one scruple.
Camphor, } of each, ten grains.
Guaiac Resin, }
Sugar Candy, two scruples.

Cinchona, one scruple.
Oil of Cloves, two drops.

Divide into twenty parcels.

- R. Proto-chloruret of Mercury, two scruples.
Camphor, } of each,
Guaiac Resin, } ten grains.
Sugar Candy, two scruples.
Cinchona, } of each,
Root of White Hellebore, } ten grains.
Oil of Cajeput, five drops.

Mix, and divide into twenty parcels.

These powders have been extolled by KLEBER in amaurosis.

- R. Proto-chloruret of Mercury, one grain.
Sugar, ten grains.
Ammoniacal Tincture of Guaiac, one drop.

Mix.

PILLS OF PROTO-CHLORURET OF MERCURY AND OF GUAIAIC. (CAST. BORUS.)

- R. Proto-chloruret of Mercury, one scruple.
Extract of Liquorice, } of each,
Guaiac Resin, } four scruples.
Make twenty-four pills.

ANTI-HERPETIC PILLS. (PIERQUIN)

- R. Proto-chloruret of Mercury, one dram.
Guaiac Resin, two drams.
Syrup of Water Lily, a sufficient quantity.

Make twenty-two pills.

Two to be taken morning and evening, with a strong infusion of hops.

ANTI-SYPHILITIC PILLS. (CADET DE GASSICOURT)

- R. Proto-chloruret of Mercury, } of each,
Guaiac Resin, } two drams.
Marshmallows Root, four ounces.
Syrup of Quince, a sufficient quantity
to make pills of four grains each.

ALIBERT directs five or six to be taken daily.

ANTI-RHEUMATIC PILLS. (PIERQUIN)

- R. Proto-chloruret of Mercury, one grain.
Guaiac Resin, } of each,
Flowers of Leopard's Bane, } four grains.

Make two pills for one dose. VIGAROUS recommended them.

ANTI-ARTHRITIC PILLS. (PIERQUIN; SAUNDERS)

- R. Proto-chloruret of Mercury, } of each,
Guaiac Resin, } one dram.

Soap, half an ounce.
Extract of Ox-gall, two drams.
Powdered Guaiac Wood, a sufficient quantity

to make pills of four grains each.

An unchemical preparation.—Dose, from two to four pills, twice a day.

ANTI-HERPETIC PILLS. (BATAV.)

- R. Proto-chloruret of Mercury, } of each,
Extract of Poisonous Sulphuret of Lime, } one dram.
Guaiac Resin, } of each,
Tartarized Antimony, } two drams.
Aqueous Extract of Opium, twelve grains.
Extract of Soap-wort, fifteen grains.
a sufficient quantity.

Mix.—The proto-chloruret is also decomposed in this preparation.

MERCURIAL PILLS. (FERR.)

- R. Proto-chloruret of Mercury, } of each,
Extract of Guaiac, } one ounce.
Camphor, one dram and a half.
Copaiba, a sufficient quantity

to make a pilular mass.—Dose, from twelve grains to half a dram.

L.

By Gum Ammoniac.

RESOLVENT PILLS. (PIERQUIN)

- R. Proto-chloruret of Mercury, one dram.
Gum Ammoniac, dissolved in
Vinegar of Squill, half an ounce.
Rhubarb, two drams.
Soap, } of each,
Extract of Soap-wort, } one ounce.
Conserve of Elicampagne, } of each,
Syrup, } a sufficient quantity

to make pills of four grains each.

This should also be rejected on the same account as the preceding.

M.

By Gamboge.

VERMIFUGE POWDER. (HISP.)

- R. Proto-chloruret of Mercury, five grains.
Gamboge, from six to ten grains.

Mix.—This is for an adult; to be taken in the morning, in syrup.

N.

By *Ipecacuanha*.MERCURIAL POWDER OF IPECACUANHA.
(SUEC.)

- R. Proto-chloruret of Mercury, one part.
Root of Ipecacuanha, four parts.
Triturate together.

MERCURIAL LAVEMENT.

Enema mercuriale seu mannatum. (AUGUSTIN)

- R. Manna, one or two ounces.
Water, six ounces.
Castor Oil, triturated with
Yolk of Egg, one or two ounces.
Proto-chloruret of Mercury, one scruple.

Ipecacuanha, in powder, half a dram or one dram.

The oil may be replaced by infusion of jalap.

- R. Proto-chloruret of Mercury, } of each,
Powdered Ipecacuanha, } one scruple.
Linseed Oil, one ounce.
Infusion of Chamomile, eight ounces.

Mix carefully.

O.

By *Jalap*.

VERMIFUGE POWDER.

Pulvis ad Vermes seu ad Lumbricos. (MIL. COPEN.; SUEC.; ELLIS; PIERQUIN; RATIER; SAUNDERS; SWEDIAUR; VAN MONS; PH. SYPH.)

- R. Jalap, twenty grains.
Proto-chloruret of Mercury, ten grains.

Mix. (PH. SYPH.; SWEDIAUR)

SUEC. prescribes three parts of jalap, and one part of proto-chloruret.—MIL. COPEN.—six of jalap, and one of the other.—HISP.—ten of jalap, and five or ten of the other.

- R. Proto-chloruret of Mercury, two grains.
Rhubarb, six grains.
Jalap, thirty grains.

Mix. (RATIER)

- R. Proto-chloruret of Mercury, eight grains.
Jalap, thirty grains.
Scammony, seven grains.
Sugar, half an ounce.

Mix. (PIERQUIN)

SAUNDERS prescribes one ounce of proto-chloruret, two drams of resin of jalap, and one ounce and a half each, of scammony and of coral.

- R. Jalap, two parts.

Rhubarb, } of each, six parts.
Santolina, }
Sulphate of Potass, one part.
Proto-chloruret of Mercury, one half part.

Mix. (VAN MONS)

- R. Proto-chloruret of Mercury, } of each,
Jalap, } five grains.
Rhubarb, }
Oil of Cinnamon, one drop.

Mix. (HISP.)

Dose, from fifteen to twenty grains, in syrup.

LAXATIVE POWDER FOR CHILDREN.

Pulvis evacuator Puerorum. (BRUNS.; NIEMANN; SPIELMANN)

- R. Proto-chloruret of Mercury, half an ounce.
Resin of Jalap, } of each,
Blanched Sweet Almonds, } two drams.
White Sugar, one ounce.
Deutoxide of Antimony, two scruples.
Oil of Lemon, two drops.

Mix. (BRUNS.; NIEMANN)

SPIELMANN prescribes one ounce of proto-chloruret, three drams of resin of jalap, one dram and a half of almonds, one dram of red sulphuret of mercury, and one ounce and a half of sugar.

Dose, from ten to twenty or thirty grains.

SIMPLE PILLS OF PROTO-CHLORURET OF
MERCURY AND OF JALAP.

Pilule ex Hydrargyro cathartice seu purgantes cum Hydrargyro muriatico miti. (MIL. COPEN.; SWEDIAUR)

- R. Proto-chloruret of Mercury, five grains.
Jalap, half a dram.
Mucilage of Gum Arabic, a sufficient quantity.

Make fifteen pills, for two doses.

VERMIFUGE PASTILLES. (HENRI)

- R. Proto-chloruret of Mercury, } of each,
Jalap, } twenty-four
White Sugar, } parts.
Tragacanth, five hundred parts.
Orange-flower Water, eight parts.
a sufficient quantity.

VERMIFUGE TROCHISCS.

Trochisci contra Vermes seu Hermetici. (BRUNS.; SPIELMANN; VAN MONS)

- R. Proto-chloruret of Mercury, four parts.
Resin of Jalap, one part.
White Sugar, sixteen parts.

Mucilage of Gum Arabic,
a sufficient quantity.

Make masses the weight of which should be augmented by five grains for each year of age; roll these masses into balls, envelope them in starch powder, and stamp upon each the number of the age to which it corresponds. (VAN MONS)

R. Proto-chloruret of Mercury, five drams.
Cinnamon, forty grains.
Resin of Jalap, seventy-five grains.
White Sugar, one ounce and a half.
Mucilage of Tragacanth,
a sufficient quantity

to make trochiscs for children, to be dosed according to age. (SPIELMANN)

R. Proto-chloruret of Mercury, } of each,
Jalap, } two scruples.
Artemisia Seeds, two drams.
White Sugar, four ounces.
Mucilage of Tragacanth,
a sufficient quantity.

Make a mass, and divide each dram of it into ten trochiscs. (BRUNS.)

SAPONACEOUS PILLS OF PROTO-CHLORURET OF MERCURY AND OF JALAP.

Pilulæ Hydrargyri jalapinæ seu mercuriales seu purgantes seu ex Resina Jalapæ seu laxantes. (BATAV.; BRUNS.; HANN.; HASS.; OLDEN.; POL.; CAST. BORUS.; BORUS.; WURCEB.; HUFELAND; RATIER)

R. Proto-chloruret of Mercury, } of each,
Resin of Jalap, } one part.
Soap, two parts.

Make pills of two grains each.—Dose, one or two. (BRUNS.; RATIER)

CAST. BORUS.; and WURCEB. prescribe one ounce of proto-chloruret, and one ounce and a half each, of resin, of jalap, and of soap.—HASS.—one ounce and a half of proto-chloruret, and six ounces each, of resin, and of soap; from sixteen to thirty grains of the mass for one dose.

R. Proto-chloruret of Mercury, one part.
Soap of Jalap, five parts.

Mix. (OLDEN.)

BATAV.; HANN.; POL.; BORUS.; and HUFELAND prescribe one part of proto-chloruret, and three of the soap.

This is a bad preparation, the soap decomposing the proto-chloruret, as in some of the following:

COMPOUND SAPONACEOUS PILLS OF PROTO-CHLORURET OF MERCURY AND OF JALAP. (AMER.; COXE)

R. Proto-chloruret of Mercury,
six drams two scruples.

Jalap, }
Rhubarb, } of each, one ounce.
Soap, }
Tartarized Antimony, }
Water, } twenty-eight grains.
a sufficient quantity

to form a pilular mass.

HYDRAGOGUE PILLS. (SWEDIAUR)

R. Proto-chloruret of Mercury, } of each,
Soap of Jalap, } an equal part.
Gamboge, }

Make pills of four grains each.—Dose, from three to six.

COMPOUND PILLS OF PROTO-CHLORURET OF MERCURY AND OF JALAP.

Pilulæ catarrhales seu mercuriales seu purgantes seu laxantes cum Mercurio. (DAN.; PALAT.; WIRTEM.; SAUNDERS.; SPIELMANN.; PH. SYPH.)

R. Proto-chloruret of Mercury, half an ounce.
Resin of Jalap, one dram.
Compound Extract of Colocynth, one ounce.
Oil of Orange, thirty drops.
Make a pilular mass. (DAN.; SPIELMANN.)

SAUNDERS prescribes two grains of proto-chloruret, two of resin of jalap, and six of compound extract of colocynth.—PH. SYPH.—four grains of proto-chloruret, four of resin of jalap, and half a dram of compound pills of myrrh.

R. Proto-chloruret of Mercury, }
Scammony, } of each,
Resin of Jalap, } one dram.
Compound Extract of Colocynth, }
Rhubarb, }
Tincture of Sassafras, }
a sufficient quantity

to make a pilular mass. (PALAT.; WIRTEM.)

It is anthelmintic, anti-syphilitic, and purgative.—Dose, twelve or fifteen grains.

CATHARTIC PILLS. (ELLIS; CADET DE GASSICOURT)

R. Proto-chloruret of Mercury,
twenty-five grains.

Jalap, }
 Rhubarb, } of each, half a dram.
 Soap, }
 Tartarized Antimony, }

Water, one dram and a half.
 a sufficient quantity

to make seventy-five pills.—Dose, two, to be repeated at the end of two hours, if necessary. (ELLIS)

R. Proto-chloruret of Mercury, half a dram.

Jalap, half an ounce.
 Sub-trito-carbonate of Iron, two drams.
 Wood-lice, two scruples.
 Syrup, a sufficient quantity

to make pills of four grains each. (CADET DE GASSICOURT)

The latter are recommended by BARTHEZ, in scrofula.

PURGATIVE BOLUSES. (PIERQUIN)

R. Proto-chloruret of Mercury, ten grains.
 Resin of Jalap, } of each,
 Sulphate of Potass, } six grains.
 Compound Syrup of Cichory, a sufficient quantity.

Mix.—CHIRAC frequently prescribed these.

PURGATIVE CHOCOLATE. (BORIES; PIERQUIN)

R. Proto-chloruret of Mercury, one ounce.
 Jalap, one ounce and a half.
 Chocolate of Health, one pound.
 Make pastilles of one dram each. (PIERQUIN)

BORIES prescribes one dram of proto-chloruret, six drams of jalap, and one pound of chocolate.

NEAPOLITAN ELECTUARY. (PIERQUIN)

R. Proto-chloruret of Mercury, } of each,
 Jalap, } one dram
 Scammony, } and a half.
 Rhubarb, } of each, two drams.
 Senna, }
 Guaiac, }
 Sarsaparilla, } of each, half an ounce.
 China Root, }
 Syrup of Roses, a sufficient quantity.

Pulverize and mix.

It is recommended in chronic discharges from the urethra.—Dose, one or two drams, in the morning, fasting, every other day for three weeks.

PURGATIVE AND VERMIFUGE BISCUITS. (PIERQUIN)

R. Proto-chloruret of Mercury, one ounce and a half.
 Jalap, three ounces.

Coriander, one dram.
 Wheat Flour, one pound.
 Sugar, one pound and a half.
 Eggs, twenty-four.
 Peel of Lemon, one.
 Essence of Bergamotte, twenty-four drops.

Orange-flower Water, one ounce.

Make biscuits of one ounce each, each of which contains fifteen grains of proto-chloruret.

O.

By Sulphate of Magnesia.

CHAUSSEIER'S LAXATIVE BOLUS. (RATIER)

R. Proto-chloruret of Mercury, three grains.
 Sulphate of Magnesia, four grains.
 Syrup of Buckthorn, a sufficient quantity

to make a bolus.

P.

By Opium.

POWDER OF PROTO-CHLORURET OF MERCURY AND OF OPIUM.

Pulvis muriatis Hydrargyri thebaicus seu Hydrargyri cum Opio. (SUEC.; MIL. COPEN.; AUGUSTIN)

R. Proto-chloruret of Mercury, two grains.
 Opium, one grain.
 Starch, one scruple.

Mix. (MIL. COPEN.; SUEC.)

AUGUSTIN prescribes two grains of opium, six of proto-chloruret, and one dram of sugar, to be divided into six parcels, as an excellent remedy in hepatitis.

DRY COLLYRIUM. (RATIER)

R. Proto-chloruret of Mercury, } of each,
 Jalap, } one scruple.
 Sugar, }
 Opium, } four grains.

Triturate together.

It is recommended in obstinate ophthalmia, particularly in that reputed syphilitic.

DIAPHORETIC POWDER. (ELLIS)

R. Proto-chloruret of Mercury, sixteen grains.
 Opium, four grains.
 Ipecacuanha, eight grains.

Make eight parcels.—Dose, one every hour or two.—It is recommended by CHAPMAN in dysentery.

R. Opium, three grains.
 Ipecacuanha, six grains.

Proto-chloruret of Mercury,
one grain and a half.
Nitrate of Potass, half a dram.

Make six parcels.—Dose, one every three or four hours.

EXPECTORANT POWDER. (ELLIS)

R. Proto-chloruret of Mercury, six grains.
Opium, three grains.
Ipecacuanha, six grains.
Nitrate of Potass, one dram.

Make six parcels.—Dose, one every two or three hours, in syrup.

PLENK'S BALSAMIC MERCURIAL LIQUOR. (SAINTE-MARIE)

R. Proto-chloruret of Mercury,
two scruples.
Tincture of Myrrh, four drams.
SYDENHAM'S Laudanum, } of each,
Concentrated Decoction of } one
Cinchona, } ounce.

Mix.—This liquor, applied on lint, frequently heals indolent syphilitic ulcers.

ANTI-SYPHILITIC MIXTURE. (SAINTE-MARIE)

R. Proto-chloruret of Mercury,
two drams.
Opium, twelve grains.
Conserve of Roses, four ounces.

Mix, and incorporate.

It is said that this quantity is sufficient for the treatment of an attack of syphilis.—It is taken by portions, in syrup of almonds, with a concentrated decoction of guaiac.

ANTI-SYPHILITIC OINTMENT.

R. Proto-chloruret of Mercury, one dram.
SYDENHAM'S Laudanum, two drams.
GALEN'S Cerate, one ounce.

Triturate for a long time, and add

Essence of Roses, one drop.

This is presented as a preventive against syphilitic diseases, being rubbed upon the glans, an hour before coition, and also as an aphrodisiac. In the former case it only serves to inspire a false security, and in the latter only vain hopes.

PILLS OF PROTO-CHLORURET OF MERCURY AND OF OPIUM. (MIL. COPEN.; VAN MONS)

R. Proto-chloruret of Mercury,
twelve parts.
Opium, one part.
Crumb of Bread, seventeen parts.
Despumated Honey,
a sufficient quantity

to make pills of two grains and a half each. (VAN MONS)

MIL. COPEN. prescribes four parts of proto-chloruret, one part of opium, and four parts of crumb of bread, to make thirty pills.

These pills are recommended in syphilitic diseases, particularly those characterized by ulcerations.

ANTI-HYDROPIC PILLS. (AUGUSTIN)

R. Proto-chloruret of Mercury,
half a dram.
Opium, one scruple
Rhubarb, one dram.
Extract of Quassia, three drams.

Make pills of two grains each.—Dose, nine or ten, thrice a day.

LESLIE'S PILLS. (PIERQUIN)

R. Proto-chloruret of Mercury,
two grains.
Opium, half a grain.

Mix.—To be taken thrice a day, in tic doloieux.

PURGATIVE PILLS. (ELLIS)

R. Proto-chloruret of Mercury,
twelve grains.
Rhubarb, twenty grains.
Opium, one grain.

Make eight pills.—Dose, four, and the remainder at the end of an hour if necessary.

PRINGLE'S PILLS. (GENEV.)

R. Compound Extract of Colocynth,
one scruple.
Proto-chloruret of Mercury,
five grains.
Opium, one grain and a half.
Make twelve pills.

ALTERATIVE PILLS. (RATIER)

R. Proto-chloruret of Mercury,
half a dram.
Extract of Opium, eighteen grains.
Hemlock, four drams.
Syrup, a sufficient quantity
to make thirty-two pills.

These are employed by DUBOIS in cases of engorgement, to appease pain.—Dose, two at first, to be gradually augmented.

SIALAGOGUE PILLS. (ELLIS)

R. Proto-chloruret of Mer-
cury, } of each,
Camphor, } one scruple.
Opium, twelve grains.
Simple Syrup, a sufficient quantity.

Make twenty pills.—Dose, one or two thrice a day.

Q.

By Lead.

ANTI-GONORRŔŔIC INJECTION.

Injectio in Gonorrhœa. (SAUNDERS)

- R. Gum Arabic, half an ounce.
Boiling Water, twelve ounces.
Dissolve, strain, and add
Proto-chloruret of Mercury, three drams.
Superacetate of Lead, four grains.
Mix.—A dram of this to be injected into the urethra thrice a day.

DESICCATIVE CERATE. (VAN MONS)

- R. Yellow Wax, forty parts.
Melt, and add
Plaster of Semi-vitreous } of each,
Oxide of Lead, } five parts.
Turpentine, }
When these are melted, add
Subcarbonate of Lead, two parts.
Proto-chloruret of Mercury, one part.
Mix.

MERCURIAL BOUGIES.

Candela seu Cereoli mercuriales. (FULD.; SPIELMANN; SWEDIAUR)

- R. Yellow Wax, six ounces.
Melt, and add
Solution of Acetate of Lead, half an ounce.
Proto-chloruret of Mercury, two drams.
Make bougies.

PLENK recommended them to restore suppressed gonorrhœa. They are no better in such a case than ordinary bougies.

OINTMENT AGAINST TINEA. (ELLIS)

- R. Proto-chloruret of Mercury, two drams.
Calcined Alum, } of each,
Subcarbonate of Lead, } half an ounce.
Oil of Turpentine, two drams.
Simple Cerate, one ounce and a half.
Mix.—This ointment to be applied in the evening and removed in the morning.

R.

By Rhubarb.

CATHARTIC POWDER. (GENEV.; ELLIS; SAUNDERS; SWEDIAUR)

- R. Proto-chloruret of Mercury, one part.
Rhubarb, two parts.
Mix. (SAUNDERS; SWEDIAUR)

ELLIS prescribes ten grains each, of the two, and one drop of oil of cinnamon; dose, twenty grains.—GENEV.—one dram of proto-chloruret, four scruples of rhubarb, and one scruple of relaxing powder.

VERMIFUGE BOLUSES. (SWEDIAUR)

- R. Proto-chloruret of Mercury, two grains.
Rhubarb, from half a scruple to two scruples.
Syrup, a sufficient quantity.
Mix.

WERLHOF'S PILLS. (SAINTE-MARIE)

- R. Proto-chloruret of Mercury, one scruple.
Extract of Rhubarb, two drams.
Syrup, a sufficient quantity
to make seventy-two pills.

They are recommended in leucorrhœa which is of a suspected character.—Dose, two, morning and evening. Their use should be continued for some time.

ANTHELMINTIC MIXTURE. (BRERA)

- R. Proto-chloruret of Mercury, six grains.
Infusion of Rhubarb, eight ounces.
Compound Syrup of Cichory, two ounces.
Mix.—To be taken by spoonfuls, in febrile affections from worms.

S.

By Scammony.

HYDRAGOGUE POWDER.

Pulvis purgans anthelmintica. (BRERA; SWEDIAUR)

- R. Proto-chloruret of Mercury, one part.
Scammony, two parts.
Mix. (SWEDIAUR)
Dose, from five to thirty grains, every four or five days.
R. Proto-chloruret of Mer- } of each,
cury, } three grains.
Sulphuretted Scammony, }

Mix. (BRERA)

This is for a child of six years, the dose is to be augmented or diminished by so many grains as the child may be more or less old by a year.

RESOLVENT OR DIGESTIVE POWDER. (SAINTE-MARIE)

- R. Proto-chloruret of Mercury, three parts.
Sulphuretted Scammony, eight parts.
Mix.

This is recommended by FINKE in obstructions of the abdominal viscera.—Dose, twelve grains, every morning, fasting, for several days in succession.

VERMIFUGE POWDER. (HERBIP.; CADET DE GASSICOURT; SPIELMANN)

R. Proto-chloruret of Mercury, five grains.
Scammony, six grains.
Rhubarb, four grains.

Mix. (SPIELMANN)

CADET DE GASSICOURT prescribes one dram each, of proto-chloruret, of rhubarb, and of scammony, and three drams of sugar.

R. Proto-chloruret of Mercury, } of each,
Scammony, } twelve
Gamboge, } grains.
five grains.

Mix.

These powders may be formed into boluses with conserve of wormwood or syrup.

ANTHELMINTIC PURGATIVE BOLUSES.

Boli anthelmintico-laxantes. (BRERA; SPIELMANN)

R. Proto-chloruret of Mercury, } of each,
Scammony, } ten grains.
Gamboge, } six grains.
Confection of Hyacinth, a sufficient quantity

to make two or three boluses. (SPIELMANN)

These boluses make the basis of NOUFER's celebrated remedy against tænia.

They are to be taken at intervals of a quarter of an hour, after having taken three drams of powdered fern root in six ounces of infusion of lime flowers, and drinking after them a cup or two of tea.

R. Proto-chloruret of Mercury, six grains.
Sulphuretted Scammony, eight grains.
Elder Rob, a sufficient quantity.

Make four boluses, one to be taken every two hours. (BRERA)

ANTHELMINTIC TROCHISCS.

Trochisci anthelmintici seu hermetici Screte, Rotule anthelminticae. (PALAT.; WIRTEM.)

R. Proto-chloruret of Mercury, one ounce and a half.
Prepared Hartshorn, two scruples.
Cinnamon, } of each,
Sulphuretted Scammony, } one scruple.
White Sugar, one ounce.
Conserve of Red Roses, two drams.

Make sixty-four trochiscs with mucilage of tragacanth. (WIRTEM.)

VOL. II.

PALAT. prescribes two drams of proto-chloruret, one dram each, of herb of carline thistle, of artemisia seeds, and of burnt hartshorn, half a dram of sulphuretted scammony, half a scruple of cinnamon, one ounce of sugar, and a sufficient quantity of mucilage.

T.

By Squill.

DIURETIC PILLS. (ELLIS; PARIS)

R. Proto-chloruret of Mercury, } of each,
Squill, } twelve grains.
Conserve of Roses, a sufficient quantity.
Make twelve pills.—Dose, one, morning and evening. (ELLIS)

R. Proto-chloruret of Mercury, six grains.
Squill, four grains.
Leaf of Digitalis, ten grains.
Myrrh, one scruple.
Assafoetida, half a dram.
Extract of Gentian, a sufficient quantity.

Make fifteen pills.—Dose, one, morning and evening. (ELLIS; PARIS)

ANTI-HYDROPIC PILLS. (PIERQUIN)

R. Proto-chloruret of Mercury, two drams and a half.
Squill, four drams.
Oxymel of Squill, a sufficient quantity.

Make one hundred pills.

They are purgative, resolvent, and diuretic.—Dose, at first two, morning and evening, to be gradually augmented.

RESOLVENT OINTMENT. (RATIER)

R. Proto-chloruret of Mercury, } of each,
Squill, } half a dram.
Prepared Lard, two drams.
Oil of Roses, four drops.

Mix.—This is employed by DUPUYTREN, in frictions, to chronic engorgements of the articulations.

U.

By Artemisia Seeds.

VERMIFUGE POWDER. (AUGUSTIN; SWEDIAUR)

R. Proto-chloruret of Mercury, half a dram.
Rhubarb, two ounces.
Artemisia Seeds, half an ounce.

Make a fine powder. (SWEDIAUR)

AUGUSTIN prescribes six grains each, of proto-chloruret, of rhubarb, and of artemisia seeds, and two drops of oil of cinnamon.

K

VERMIFUGE BOLUS. (RATIER)

- R. Proto-chloruret of Mercury, two grains.
 Artemisia Seeds, eight grains.
 Camphor, six grains.
 Simple Syrup, a sufficient quantity.

VERMIFUGE ELECTUARY. (VAN MONS)

- R. Proto-chloruret of Mercury, half a scruple.
 Rhubarb, }
 Artemisia Seeds, } of each. two drams.
 Valerian Root, }
 Conserve of Wormwood, one ounce and a half.

Oxymel of Squill, a sufficient quantity.

Mix.—It is recommended in epilepsy complicated with worms.—Dose, from two drams to half an ounce, according to the age and strength of the patient.

VOGLER'S ANTHELMINTIC ELECTUARY.
(BRUNS.; CADET DE GASSICOURT;
SWEDIAUR)

- R. Proto-chloruret of Mercury, from six to twelve grains.
 Artemisia Seeds, two drams.
 Jalap, one dram.
 Cinnamon, half a dram.
 Syrup of Peach Flowers, a sufficient quantity.

Mix.—To be taken in the morning in softened wafer.—Dose, one twelfth of this mass to children of from two to four years, one eighth for those of from five to eight, and one fourth and above for adults.

V.

By Turpentine.

FERNANDEZ' COLLYRIUM.

Injunctio Hydrargyri Muriatis camphorata.
(HISP.)

- R. Proto-chloruret of Mercury, half a scruple.
 Alum, }
 Alcoholized Sulphuric } of each,
 Ether, } one scruple.
 Camphor, }
 Turpentine, half an ounce.
 Yolks of Eggs, two.

Triturate in a glass mortar, and strain with expression.

ASTRINGENT ELECTUARY. (SAINTE-MARIE)

- R. Venice Turpentine, four drams.
 Rhubarb, three drams.
 Proto-chloruret of Mercury, half a scruple.
 Syrup, a sufficient quantity.

Mix.—Dose, one scruple, thrice a day, towards the end of gonorrhœa.

DEUTO-CHLORURET OF MERCURY.

Hydrargyrum muriatus seu muriatus corrosivus, Hydrargyrum muriatum corrosivum, Hydrargyrum corrosivum album, Hydrargyri Oxymurias seu Perurias, Mercurius muriatus seu sublimatus corrosivus, Murias Hydrargyri corrosivus seu oxygenatus sublimatione seu præcipitatione paratus, Murias hydrargyricum corrosivum, Murias oxygenatus Hydrargyri sublimatus, Superurias Hydrargyri, Murias Hydrargyri basi Oxydi imperfecti.—Corrosive Sublimate, Perchloruret, Bichloruret, or Bichloride of Mercury, Superoxygenated or Oxymuriate of Mercury. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PARIS; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

1. The Old Process.

- R. Mercury, four parts.
 Nitric Acid, a sufficient quantity
 to dissolve the metal. The solution being completed, evaporate the liquor to dryness, pulverize the mass, and add to it

Decrepitated Muriate of Soda, }
 Sulphate of Iron, calcined to whiteness, } of each,
 five parts.

Triturate well together; put the mixture into a phial which it may fill to one half; heat it slowly in a sand-bath, stopping the phial only when acid and watery vapours have ceased to ascend: having then closed it, augment the heat slowly. When the evaporation has ended, break the phial, and separate the white mass from the scoriæ. (AMST.; BATAV.; BELG.; BRUNS.; GENEV.; SARD.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR)

2. Kunkel's process.

Hydrargyri Oxymurias. (LOND.)

- R. Purified Mercury, by weight, two pounds.
 Sulphuric Acid, by weight, thirty ounces.
 Dried Muriate of Soda, four pounds.

Boil the mercury with the acid in a glass vessel, till the product becomes dry; when it is cold triturate it in an earthen mortar with the muriate. Finally, sublime it in a glass cucurbit, with a gradually raised heat.

Murias Hydrargyri corrosivus.
(EDINB.)

- R. Purified Mercury, two pounds.
Sulphuric Acid, two pounds and a half.
Dried Muriate of Soda, four pounds.

The operatory process is the same as that directed by LOND.; but the scorix are required to be separated.

Hydrargyri Murias corrosivum.
(DUBL.)

- R. Persulphate of Mercury, five parts.
Dried Muriate of Soda, two parts.

Rub these together to fine powder in an earthenware mortar; then sublime the mixture in a proper vessel with a gradually raised heat.

- R. Mercury, twelve ounces.
Concentrated Sulphuric Acid, twenty ounces.

Distil to dryness; pulverize the residuum in a glass mortar, and add to it

Muriate of Ammonia, in powder,
a weight equal to that of the mass.

Triturate well together; put the mixture into a phial which it may fill to about one half, and place it in a slowly heated sand-bath; leave it open while it disengages watery and acid vapours; then stop it with paper, and gradually augment the heat. The operation being finished, break the phial, and separate the sublimed white mass from all impurities which may be seen at its inferior surface. (AMER.; AUSTR.; BAVAR.; DAN.; MIL. COPEN.; HISP.; FENN.; HAMB.; LIPP.; OLDEN.; LUSIT.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; COXE; SWEDIAUR)

- R. Mercury, four parts.
Sulphuric Acid (66 degrees), five parts.

Boil till the whole is reduced to five parts. Then take of this

Deuto-sulphate of Mercury,	} of each,
Pulverized Muriate of Soda,	
	one part.

Black Oxide of Manganese, one part.

Mix together. At the end of several days, put the mixture into one or more flat-bottomed matrasses which should be filled to one third, and plunged into a sand-bath to the neck. Cover the orifice with a little pot turned upside down, and apply a gradual heat till the mass becomes totally sublimed. (FORM. HÔP. FR.)

GALL. prescribes four hundred and eighty parts of sulphate, the same of muriate, and four hundred and fifty of oxide. But HENRI observes that one fourth of the last would suffice, the purpose of which is only to convert into a deuto-sulphate the proto-sulphate which may exist in the salt first obtained.

FERR. directs six ounces of mercury to be dissolved in an equal quantity of sulphuric acid, the solution to be evaporated to dryness, the residuum to be triturated with its weight of muriate of soda, and the mixture to be sublimed.

- R. Red Oxide of Mercury, ten parts and a fourth.
Nitric Acid, a sufficient quantity
to dissolve the oxide. Evaporate to dryness, and triturate the residuum with
Muriate of Soda, four parts.
Sublime. (VAN MONS)

- R. Mercury, one pound.
Nitric Acid, a sufficient quantity
to dissolve the metal with heat. Pour into the liquor a hot solution made with

Muriate of Soda, one pound and a half.
Boiling Water, a sufficient quantity,
avoiding the red vapours which are disengaged; the abundant precipitate at first formed becomes soon dissolved. Set aside the liquor, and separate the prismatic crystals which form. Evaporate till a pellicle appears, and set apart to crystallize again. (FULD.)

- R. Red Oxide of Mercury, any quantity.
Hydrochloric Acid, a sufficient quantity

to dissolve the oxide. Evaporate in a glass vessel, and crystallize. (ANT.; VAN MONS)

The deuto-chloruret of mercury or corrosive sublimate is one of the most virulent poisons we know. It is employed internally as an irritant, and externally as an escharotic, and also in some arts, to preserve animal substances which it renders impusible, giving to them the hardness of wood. But it is principally as an anti-syphilitic that it has enjoyed and still enjoys a high celebrity. The English, however, have decided that it should not be depended on in syphilis, so that they have abandoned the use of it again almost entirely to empirics, from

whose hands it had passed, towards the end of the last century, into those of the regular practitioners. But it neither deserves this unqualified condemnation, nor such exaggerated encomiums as VAN SWIETEN bestowed on it on the faith of one of those men so common in all times, who make nothing of distorting facts and of uttering deliberate falsehoods to flatter the weaknesses of men above them. When used with judgment, it is a most valuable medicine, and is highly useful in many cases; but, abandoned to quackery or to blind routine, it becomes the fruitful source of an infinity of evils.

It is incompatible with the alkalis, alkaline subcarbonates and sulphurets, tartarized antimony, acetate of lead, nitrate of silver, white of egg, gluten, metallic mercury, iron, copper, lead, volatile oils, and with infusions and decoctions of vegetable astringents. BOULLAY has even asserted that all vegetable and animal substances have the property of separating its constituent principles, in a space of time which varies with each of them, and which indeed it would be useful to determine, so as to know precisely the value of certain medicaments of very extended use, in which the inventors have intended the sublimate to be the most active ingredient.

It is administered in the dose of one sixteenth, one eighth, one fourth, or one half, of a grain.

I. *Preparations in which the action of the Deuto-chloruret of Mercury is not modified, at least perceptibly, by any foreign substance.*

AQUEOUS SOLUTION OF DEUTO-CHLORURET
OF MERCURY.

*Aqua divina Fernellii seu phagedenica, Li-
quor mercurialis seu Muriatis Hydrar-
gyri seu Hydrargyri muriatici corrosivi.*
(AMST.; BATAV.; BAVAR.; MIL. CO-
PEN.; FERR.; HAMB.; HASS.; LIPP.;
CAST. BORUS.; SAX.; WURCEB.; CADET
DE GASSICOURT.; PIDERIT.; SAUNDERS;
SWEDIAUR)

R. Deuto-chloruret of Mercury, one dram.
Distilled Water, one pound.

Dissolve with trituration in a stone mortar. (AMST.)

BATAV.; and FERR. prescribe one dram to two pounds.—MIL. COPEN.; CAST. BORUS.; SAX.; WURCEB.; and SAUNDERS—one grain to one ounce.—SWEDIAUR—from twenty to thirty grains to two pounds.—HASS.—one scruple to ten ounces.—LIPP.—one scruple to one pound and a half.

HASS. ; and PIDERIT.—forty-four grains to one pound.—BAVAR.—one part to twenty-four parts.—WURCEB. elsewhere prescribes ten grains to four ounces.—CADET DE GASSICOURT—from twelve to eighteen grains to two pounds.

The greater part of these solutions are employed externally, in lotions, principally in itch; some of them, as that of BAVAR., are noticed as tests.

One part of the deuto-chloruret of mercury is soluble in about twenty parts of water at the ordinary temperature, and in thrice its weight of boiling water.

MERCURIAL OPHTHALMIC LIQUOR.

Collyrium ex Muriate Hydrargyri oxygenata.
(FULD. ; SWEDIAUR)

R. Dento-chloruret of Mercury, one grain.
Distilled Water, six ounces.

Dissolve. (FULP.)

SWEDIAUR prescribes from half a grain to one grain to four ounces of water.

MERCURIAL LOTION.

Lotio Hydrargyri. (GENEV.; AUGUSTIN)

R. Deuto-chloruret of Mercury, one scruple.
Distilled Water, twelve ounces.
Hydrochloric Acid, twenty-four drops.

Mix. (GENEV.)

AUGUSTIN prescribes a solution of three grains of deuto-chloruret in two pounds of water or rose water.

R. Deuto-chloruret of Mercury, two grains.
Subacetate of Copper, six grains.
Water, two pounds.

Mix.—It is recommended in obstinate tinea. (AUGUSTIN)

R. Deuto-chloruret of Mercury,	two drams.
Nitric Oxide of Mercury,	ten grains.
Spirit of Angelica,	of each,
Rosemary,	two drams.
Sulphate of Zinc,	four drams and a half.
Water.	four pounds.

Mix.—It is recommended in itch. (AUGUSTIN)

MERCURIAL LAVEMENT. (AUGUSTIN)

R. Deuto-chloruret of Mercury, six grains.
 Infusion of Linseed, eight ounces.

Mix.—It is useful against ascarides.

MERCURIAL INJECTION. (RATIER)

- R. Deuto-chloruret of Mercury, one grain.
Boiling Water, one ounce.

Mix.—For leucorrhœa.

AUGUSTIN recommends, after MONTAGGIA, in chronic gonorrhœa, one grain dissolved in four ounces of water.

ALIBERT'S MERCURIAL LOTION.

Eau rouge. (RATIER)

- R. Deuto-chloruret of Mercury, one dram.
Distilled Water, one pound.
Alkanet, a sufficient quantity,
to colour the solution.

Linen or sponge is to be moistened with this liquor and parts affected with inflammatory herpes bathed with it; it is recommended particularly when the tetters are supposed to have a syphilitic origin.

MERCURIAL COLLYRIUM.

Aqua ophthalmica mercurialis. (AUGUSTIN; HUFELAND)

- R. Deuto-chloruret of Mercury, half a grain.
Rose Water, three ounces.
Dissolve, and add
Mucilage of Quince Seeds, one dram.
Cherry-laurel Water, thirty-drops.

Mix. (HUFELAND)

AUGUSTIN prescribes a mixture of one drop of solution of deuto-chloruret and four ounces of distilled water.

ANTI-SYPHILITIC BATH. (RATIER)

- R. Deuto-chloruret of Mercury, from six to twelve grains.
Distilled Water, eight pounds.
Pure Water (at 30 degrees), two hundred pounds.

This bath should be taken in a wooden bathing-tub. It is highly dangerous in spite of the pompous eulogiums which have been bestowed on it.

ANTI-SYPHILITIC POTION. (FORM. HÔP. FR.; GENEV.: ELLIS)

- R. Solution of Deuto-chloruret of Mercury, ten parts.
Gum Arabic, one part.
Distilled Water, one hundred parts.

Dissolve the gum in the water, and add the solution. (FORM. HÔP. FR.)

- R. Deuto-chloruret of Mercury, two grains.
Pure Water, six ounces.

Spirit of Cinnamon, } of each,
Simple Syrup, } one ounce.

Mix. (ELLIS)

- R. Deuto-chloruret of Mercury, eight grains.
Distilled Water, fifteen ounces.
Hydrochloric Acid, twelve drops.
Syrup of Poppies, one ounce.

Mix. (GENEV.)

Dose, one or two spoonfuls, twice or thrice a day.

MERCURIAL GARGLE.

Gargarisma in Ulceribus venereis. (AUGUSTIN; BRERA; RATIER; PH. SYPH.)

- R. Deuto-chloruret of Mercury, six grains.
Decoction of Barley, one pound.
Honey of Roses, two ounces.

Mix. (PH. SYPH.)

RATIER prescribes from one to two grains of deuto-chloruret in six ounces of decoction of marshmallows, or two grains of deuto-chloruret, four ounces of distilled water, and half an ounce of syrup of honey. —BRERA—one grain of deuto-chloruret, one pound of decoction of barley, and half an ounce of honey of roses.

ANTI-SYPHILITIC LAVEMENT. (RATIER)

- R. Deuto-chloruret of Mercury, two grains.
Distilled Water, two ounces.
Decoction of Linseed, one pound.

Mix.—It is recommended in syphilitic affections of the rectum, when the patient cannot support the deuto-chloruret by the mouth.

DETERSIVE INJECTION.

Injectio in Nicrosi syphilitica. (BRERA)

- R. Deuto-chloruret of Mercury, two grains.
Decoction of Barley, one pound.
Tincture of Myrrh, one dram.

Mix.

COMPOUND PHAGEDENIC LIQUOR. (VAN MONS)

- R. Phagedenic Liquor, six ounces.
Extract of Myrrh, one dram.
Mix, and add
Tincture of Myrrh, half an ounce.

PIDERIT'S PHAGEDENIC LIQUOR. (PIDERIT)

- R. Phagedenic Liquor, six ounces.
Tincture of Saffron, half an ounce.
Solution of Myrrh, one dram.

RHAZES' Compound Powder of Sub-carbonate of Lead, one ounce.

Mix.—This is unchemical, and should be rejected.

ALCOHOLIC SOLUTION OF DEUTO-CHLORURET OF MERCURY.

Liquor Hydrargyri Oxymuriati seu syphiliticus Turneri, Murias Hydrargyri spirituosus liquidus, Solutio Mercurii Corrosivi seu Muriatis Hydrargyri oxygenati. (LOND.; AMER.; ANT.; MIL. COPEN.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; SAX.; HERBIP.; CADET DE GASSICOURT; COXE; ELLIS; SPIELMANN; SWEDIAUR; VAN MONS)

Liquor Hydrargyri Oxymuriati. (LOND.)

R. Oxymuriate (Deuto-chloruret) of Mercury, eight grains.
Distilled Water, fifteen fluid ounces.
Alcohol, one fluid ounce.

Dissolve the salt in the water, and add the alcohol.

AMER.; COXE; and SWEDIAUR give the same formula.—RATIER—prescribes sixteen grains of deuto-chloruret, one pound of water, and half a dram of alcohol.—FORM. HÔP. FR.—one part of the salt, nine hundred parts of water, and one hundred of alcohol (33 degrees).—GALL.—eight grains of the salt, fourteen ounces and a half of water, one ounce and a half of alcohol (36 degrees).—FERR.—six grains of the salt, four ounces of water, and eight ounces of alcohol.—CADET DE GASSICOURT—twenty grains of the salt, twenty-five drams of alcohol (36 degrees), and two pounds of water.

R. Deuto-chloruret of Mercury, twelve grains.
Spirit from Grain rectified twice, two pounds.

Dissolve with trituration. (ANT.; HERBIP.; SPIELMANN; SWEDIAUR; VAN MONS)

MIL. COPEN. prescribes one grain of the salt, and four ounces of proof spirit.—SAX. eight grains of the salt and one pound of proof spirit.—ELLIS—two grains of the salt and four ounces of proof spirit.

R. Deuto-chloruret of Mercury, one scruple.
White Sugar, half an ounce.
Triturate perfectly, and add
Proof Spirit, ten ounces.
Dissolve, and filter. (VAN MONS)

R. Deuto-chloruret of Mercury, six grains.
Alcohol, twelve ounces.

Dissolve, and add

Syrup of Tolu, one ounce and a half.

Mix. (FULD.; NIEMANN)

R. Deuto-chloruret of Mercury, eight grains.
Distilled Water, fifteen ounces.
Hydrochloric Acid, twelve drops.
Syrup of Poppies, one ounce.

Mix. (GENEV.)

After many trials VAN SWIETEN stopped at the proportion of twelve grains of deuto-chloruret to two pounds of spirit. He gave, morning and evening, half an ounce of this liquor, or a quarter of a grain of the salt, making the patient take afterwards a pound of emollient decoction mixed with a third or fourth of milk. The formula of GALL. gives half a grain to the ounce: but the ordinary mode in France is to dissolve eight grains of the salt in one dram of alcohol, and to dilute the solution with thirty-two ounces of distilled water. Thus, suppressing the alcohol, which serves no purpose whatever, and preserving the proportion of the two other components, we obtain a solution containing exactly a quarter of a grain to the ounce, which is a very convenient proportion for easily dosing and administering it. In the morning, fasting, half an ounce of the liquor, containing an eighth of a grain of the salt, is given on the first day; three quarters of an ounce, containing three-sixteenths of a grain, on the second; and on the third, one ounce, containing one fourth of a grain: and so on till half a grain is arrived at, which quantity should never be exceeded. It is frequently of advantage to divide the quantity into two takings, one for the morning and the other for the evening. It should be given in several ounces of decoction of barley, of linseed, or of marshmallows, of milk, of gum water, or syrup of gum; it may also be mixed with two pounds of ptisan, which the patient may drink by glassfuls in the course of the day.

CAMPHORATED MERCURIAL LIQUOR.

Liquor Hydrargyri camphorati seu ad Condylomata. (AUGUSTIN)

R. Deuto-chloruret of Mercury, half a dram.
Camphor, one dram.
Alcohol, one ounce.

RUST recommends it to dry and destroy condylomata.

ANTI-SYPHILITIC LIQUOR. (BRUNS.)

R. Deuto-chloruret of Mercury, twelve grains.

Fennel Water, } of each, one pound.
 Alcohol, }
 Rose Julep, } two ounces.

Mix.

ANTI-SYPHILITIC GARGLE. (DISP. CHA. PARIS)

R. VAN SWIETEN'S LIQUOR, two ounces.
 Honey of Roses, one ounce.
 Sudorific Decoction, six ounces.

Mix.

ANTI-SYPHILITIC LIQUOR. (PIERQUIN)

R. Deuto-chloruret of Mercury, twelve grains.
 Camphor, four grains.

Triturate together in a marble mortar, and dissolve in

Proof Spirit, two pounds.

Add towards the end

Syrup of Red Poppies, one ounce.

Mix.—Dose, from one to two spoonfuls, morning and evening, in half a pound of pectoral ptisan or of milk.

MARRYAT'S SOLUTION. (NIEMANN)

R. Deuto-chloruret of Mer- } of each,
 cury, } ten grains.
 Hydrochloric Acid, }
 Compound Spirit of Lavender, one ounce.

Mix.—Dose, twenty drops, morning and evening, in six spoonfuls of white wine, or decoction of marshmallows.

MINERAL MIXTURE. (NIEMANN)

R. Deuto-chloruret of Mercury, six grains.
 Distilled Water, nine ounces.

Dissolve, and add

Vinous Cinnamon Water, two ounces.
 Rectified Alcohol, } of each,
 Syrup of Marshmallows, } one ounce.

Mix.

BOURQUENOD'S POTION.

R. Extract of Liquorice, one pound and a half.
 Water, three pounds and a half.

Dissolve, and add

Deuto-chloruret of Mercury, twelve grains.

Dissolved in

Rectified Alcohol, two ounces.

Mix.—Dose, two spoonfuls in one pound of water.

SAINT-ILDEPHONT'S ANTI-SYPHILITIC SYRUP. (SAINTE-MARIE)

R. Deuto-chloruret of Mercury, fifteen grains.
 Alcohol, two drams.

Triturate together till no more bubbles appear and no deposit is formed. Then add

Syrup of Maidenhair, twenty-four ounces.

Place the whole in a water-bath till the alcohol is evaporated.—Dose, one spoonful at first, and two or three, in two pounds of strong decoction of marshmallows, to be taken in the course of the day. This remedy, continued for three weeks or a month, with milk for the patient's nourishment, frequently cures syphilitic disorders.

ANTI-SYPHILITIC GARGLE. (CADET DE GASSICOURT)

R. VAN SWIETEN'S LIQUOR, one dram.
 Decoction of Barley, one pound.
 Syrup, two ounces.

Mix.

GRINDEL'S PHAGEDENIC LIQUOR.

Aqua caustica ad Condylomata. (NIEMANN)

R. Deuto-chloruret of Mercury, half a dram.
 Camphor, one dram.
 Rectified Alcohol, one ounce.

Mix.

WHATELY'S INJECTION. (ELLIS; SAINTE-MARIE)

R. Deuto-chloruret of Mercury, one dram.
 Alcohol, one ounce.

Dissolve, and mix from two to twenty drops of this solution with

Distilled Water, four ounces.
 Sulphate of Zinc, from four to ten grains.

Mix.—It is recommended in blennorrhœa; one dram to be injected thrice a day.—The deuto-chloruret is decomposed by the sulphate.

ELLIS prescribes ten grains of sulphate, two of deuto-chloruret, and eight ounces of rose water.

ANTI-SYPHILITIC TINCTURE. (AUGUSTIN)

R. Deuto-chloruret of Mercury, one scruple.
 Alcohol, four ounces.

Dissolve, and add

Canadian Balsam, } of each,
 Guaiac Resin, } two ounces.

Dissolved in

Alcohol, four ounces.
Oil of Sassafras, two drams.

Mix.—Dose, ten drops every hour in chronic gonorrhœa.

ÆTHEREAL SOLUTION OF DEUTO-CHLORURET OF MERCURY.

Æther Mercurialis. (NIEMANN)

R. Deuto-chloruret of Mercury, sixteen grains.
Sulphuric Æther, one ounce.

Mix.—CHERON has given this formula.—Dose, from six to twelve drops, in milk, gum water, or sugar and water; and from two drams to half an ounce, externally, in frictions in the neighbourhood of the genital organs.

A long while before CHERON, LAFONTAINE had recommended to be taken, morning and evening, in a cupful of mucilaginous drink, from ten to twenty drops of a solution of four grains of deuto-chloruret in half an ounce of sulphuric æther.

The volatility of the excipient renders this preparation objectionable.

EMULSION OF DEUTO-CHLORURET OF MERCURY. (NIEMANN)

R. Deuto-chloruret of Mercury, two grains.
Sulphuric Æther, one dram.

Incorporate the solution with

Yolks of Eggs, two.

And add gradually

Peppermint Water, five ounces.
Syrup of Marshmallows, one ounce.

Mix.

CHERON'S MERCURIAL SYRUP.

R. White Sugar, one pound and a half.
Distilled Water, one pound.

Make a syrup, and when cold add

Æthereal Solution of Deuto-chloruret of Mercury, one ounce.

Mix.

CYRILLO'S OINTMENT.

Unguentum Cyrilli seu mercuriale corrosivum seu mercuriale aplyalizomenon seu Muriatis Hydrargyri oxygenati. (GALL.; FERR.; FORM. HÔP. FR.; SAX.; BORIES; BRUGNATELLI; CADET DE GASSICOURT; HUFELAND; NIEMANN)

R. Deuto-chloruret of Mercury, one part.
Prepared Lard, eight parts.

Triturate together for a long time, in a glass mortar. (GALL.; FERR.; FORM. HÔP. FR.; BRUGNATELLI)

R. Deuto-chloruret of Mercury, } of each,
Hydrochlorate of Ammonia, } one
Lard, } part.
eight parts.

Triturate for twenty-four hours. (SAX.; HUFELAND; NIEMANN)

BORIES; and CADET DE GASSICOURT prescribe one dram of deuto-chloruret, ten grains of hydrochlorate, and one ounce of lard.

CYRILLO directed one dram of it to be rubbed into the sole of the foot; CULLERIER thought half a dram was enough to begin with.

CORROSIVE PLASTER.

Emplastrum corrosivum. (GENEV.)

R. Deuto-chloruret of Mercury, half a dram.
Lard, half an ounce.
White Wax, two drams.

Mix.

CYRILLO'S OINTMENT WITH OPIUM. (PLANCHE)

R. Deuto-chloruret of Mercury, one part.
Prepared Lard, eight parts.

Triturate together, and add

Extract of Opium, ten parts.

Mix.—It is recommended in obstinate gonorrhœa, in slight frictions to the perineum.

ANTI-PSORIC OINTMENT.

Unguentum Muriatis oxygenatis Hydrargyri. (HISP.; BORIES)

R. Deuto-chloruret of Mercury, one part.
Subcarbonate of Lead, fifteen parts.
Lard, thirty parts.

Triturate together. (BORIES)

R. Fresh Butter, four ounces.
Turpentine, } of each, one ounce.
Oil of Roses, }

Melt over a very slow fire, and add

Subcarbonate of Lead, two ounces and a half.

Deuto-chloruret of Mercury, } of each,
Calcined Alum, } two drams.
Yolks of Eggs, two.
Lemon Juice, two ounces.

Mix.—This is a bad preparation, and should be rejected.

HOFFMANN'S MAJOR PILLS. (BRUNS.; FERR.; HAMB.; HANN.; WIRTEM.; AUGUSTIN; BRERA; PIERQUIN; RATIER; SAINTE-MARIE; SPIELMANN)

R. Deuto-chloruret of Mercury, fifteen grains.
Distilled Water, six drams.

Add to the solution
Crumb of Bread, two drams and a half.

Mix, and make one hundred and twenty pills, containing each one eighth of a grain of deuto-chloruret; they may be taken reduced to powder with sugar. (BRUNS.; HANN.; WIRTEM.)

SPIELMANN prescribes one dram of deuto-chloruret, a sufficient quantity of water to dissolve it, and nine drams of crumb of bread, to make pills of one grain each, which will be six hundred, each containing one tenth of a grain of deuto-chloruret. By some inconceivable mistake SPIELMANN directs four or five of them to be taken thrice a day.

R. Deuto-chloruret of Mercury, six grains.

Dissolve it in a sufficient quantity of water, and add

Crumb of Bread, } of each, a
White Sugar, } sufficient quantity,
to make one hundred and eighty pills, each of which contains one thirtieth of a grain of deuto-chloruret. (HAMB.; AUGUSTIN)

R. Deuto-chloruret of Mercury, one scruple.

Alcohol, two drams.
Wheat Flour, three drams.
Distilled Water, a sufficient quantity,
to make one hundred and sixty pills, each containing one eighth of a grain of deuto-chloruret. (RATIER)

FERR.; and BRERA prescribe half a grain of proto-chloruret, a sufficient quantity of alcohol to dissolve it, and a sufficient quantity of crumb of bread to make a mass of four pills.

R. Deuto-chloruret of Mercury, ten grains.

Sugar Candy, } of each,
Potato Starch, } one dram.
Water, a sufficient quantity,

to make a mass to be divided into eighty pills, containing each, one eighth of a grain of deuto-chloruret. (PIERQUIN)

R. Deuto-chloruret of Mercury, twelve grains.

Distilled Water, one dram.

Dissolve and add
Manna, two drams.

Liquorice Powder, a sufficient quantity,
to make a mass to be divided into ninety-six pills. (SAINTE-MARIE)

These last pills are the best of all. One is given at first, and then two, morning and evening.

VOGLER'S PILLS. (NIEMANN)

R. Extract of Liquorice, one dram.
VOL. II.

Gum Tragacanth, two scruples.
Deuto-chloruret of Mercury, half a scruple.

Mix the extract and gum by trituration, then add the salt dissolved in a little water and make a mass to be divided into one hundred pills; each contains one tenth of a grain of the deuto-chloruret.

SIALAGOGUE PILLS. (ELLIS)

R Deuto-chloruret of Mercury, } of each,
Hydrochlorate of Ammonia, } five
Distilled Water, thirty drops,
Conserve of Roses, one scruple.
Liquorice Powder,

a sufficient quantity,
to make forty pills, each of which contains one eighth of a grain of deuto-chloruret.

ESCHAROTIC TROCHISCS.

Trochisci Muriatii Oxygenati Hydrargyri.
(HISP.; GALL.; FORM. HÔP. FR.)

R. Deuto-chloruret of Mercury, one part.
Starch, two parts.
Mucilage of Gum Arabic,

a sufficient quantity,
to make trochiscs. (GALL.; FORM. HÔP. FR.)

HISP. directs four parts of deuto-chloruret, seven of starch, and one part of gum tragacanth, to be pulverized, and then reduced to a paste with water.

ANTI-SYPHILITIC CHOCOLATE.

R. Deuto-chloruret of Mercury, sixteen grains.

Alcohol, a sufficient quantity,
to dissolve it. Add.
Peruvian Balsam, four drams.
Prepared Cocoa Paste,

fourteen ounces.
Sugar, two ounces.

Each pound should be divided into thirty-two takings. It is a bad preparation and should be thrown aside.

II. *Preparations in which the action of Deuto-chloruret of Mercury is more or less modified by foreign substances.*

A.

By Aconite.

MERCURIAL PILLS OF ACONITE. (PIERQUIN; PLANCHE)

R. Deuto-chloruret of Mercury, two grains.
Extract of Aconite, one scruple.

L

Triturate for a long time in a glass mortar, and divide into twenty pills.

They are recommended in chronic disorders of the lymphatic system, in tetters, particularly those complicated with itch, scrofula, and syphilis.—Dose, one pill, morning and evening, augmenting it by one every ten days.

B.

By Alum.

ALUMINOUS MERCURIAL LIQUOR.

Aqua Mercurii vitriolati. (FULD.; AUGUSTIN)

R. Deuto-chloruret of Mercury, } of each,
Alum, } two drams.
Water, } two pounds.

Boil down to one half, and filter.

No trace of the deuto-chloruret is left in this preparation.

C.

By Antimony.

ANTI-HERPETIC ELECTUARY. (BORIES; PIERQUIN)

R. Deuto-chloruret of Mercury, three grains.
Sulphuret of Antimony, half an ounce.
Extract of Fumitory, one ounce.
Guaic Resin, half an ounce.
Syrup of Orange Peel, a sufficient quantity.

Mix.—FOUQUET prescribes from half a dram to one dram, twice a day. It is a bad preparation.

D.

By Hemlock.

ANTI-SYPHILITIC PILLS. (FULD.; ELLIS)

R. Deuto-chloruret of Mercury, six grains.
Distilled Water, a sufficient quantity, to dissolve it. Add
Extract of Hemlock, one dram.
Powdered Hemlock, a sufficient quantity, to make forty-eight pills.

ANTI-HERPETIC PILLS. (PIERQUIN)

R. Deuto-chloruret of Mercury, eight grains.
Rhubarb, one dram.

Soap, } of each,
Extract of Hemlock, } two drams.

Make sixty-four pills.

It is a bad preparation.

MERCURIAL GARGLE.

Gargarisma Belladonnæ cum Hydrargyro. (AUGUSTIN)

R. Herb of Belladonna, } of each,
Hemlock, } one handful.
Linseed, } two ounces.
Deuto-chloruret of Mercury, from six to ten grains.
Water, } two pounds.

Boil for a few minutes.

SANTER recommends it in affections of the throat which succeed scarlatina.

E.

By Scurvy-grass.

PORTAL'S ANTI-SCORBUTIC SYRUP. (BRUGNATELLI; CADET DE GASSICOURT; HENRI)

R. Root of Gentian, one ounce.
Madder, } of each, four drams.
Cinchona, }
Water, } four pounds.

Boil till reduced to two pounds; strain the decoction, and add to it

White Sugar, three pounds.

Clarify with white of egg, boil to the consistence of syrup, and strain.

On the other hand.

R. Leaves of Water Cress, } of each,
Scurvy-Grass, } four ounces.
Root of Horse-radish, } one ounce.

Bruise in a mortar, express the juice, filter this when cold, and dissolve in it, in a water-bath,

White Sugar, twenty-two ounces.

Strain, mix this with the preceding syrup, and add

Deuto-chloruret of Mercury, four grains,

previously dissolved in

Alcohol, two drams.

Mix.—It is recommended in diseases of the skin, scrofula, scurvy, and syphilitic affections of old standing.—Dose, from one to two ounces, in proper drink.

F.

By Copaiba.

PAJOT LAFOREST'S ELECTUARY.

R. Deuto-chloruret of Mercury, } of each,
Kino, } half an
Copaiba, } ounce.

Gum Arabic, one ounce and a half.
 White Sugar, five ounces.
 Mint Water, a sufficient quantity.

Make a mucilage with the gum and mint water, and triturate with it the kino, the deuto-chloruret, and the copaiba; add to the mixture a little rose water, then add the sugar, and a sufficient quantity of rose water to give a proper consistence.

It is recommended in chronic discharges from the urethra and at the commencement of acute urethritis. — Dose, two drams, morning and evening, alone or with any proper liquid.

G.

By Copper.

KRUGER'S CAUSTIC POWDER. (CADET DE GASSICOURT)

R. Deuto-chloruret of Mercury, } of each,
 Sulphate of Copper, } ten grains.
 Nitrate of Silver, } six grains.

Mix.—This is a bad preparation; it has been recommended to be applied in syphilitic excrescences, after having moistened them with a brush.

DRY COLLYRIUM. (VAN MONS)

R. Deuto-chloruret of Mercury, } of each,
 Muriate of Copper, } four grains.
 Florentine Iris, } half a dram.
 Fine Sugar, } two drams.

Mix.

H.

By Guaiac.

FALK'S ANTI-SYPHILITIC TINCTURE.

Elixir balsamicum purificans, Essentia Balsami Canadensis mundificans. (HERBIP.; PIDERIT; BORIES; PIERQUIN; SPIELMANN; VAN MONS)

R. Deuto-chloruret of Mercury, one scruple.
 Alcohol, one ounce.

Dissolve and add

Guaiac Resin, } of each,
 Canadian Balsam, } one ounce,
 dissolved in
 Alcohol, seven ounces.

Filter. Add further,

Oil of Sassafras, two drams.
 Mix, shaking the bottle. (BORIES; PIERQUIN; SPIELMANN)

R. Deuto-chloruret of Mercury, one scruple.

Rectified Alcohol, four ounces.

Dissolve, and add

Venice Turpentine, } of each,
 Guaiac Resin, } one ounce,
 dissolved in

Alcohol, four ounces.

Mix, and add

Oil of Sassafras, two drams.

Preserve. (HERBIP.)

PIDERIT gives the same formula, but substitutes turpentine for the Canadian balsam.

VAN MONS has properly simplified this formula in the following manner:

R. Deuto-chloruret of Mercury, one scruple.
 Oil of Sassafras, two drams.
 Tincture of Guaiac Resin, eight ounces.
 Proof Spirit, twelve ounces.

Dissolve and filter.

It is a powerful stimulant, and is recommended in affections attributed to the presence of the syphilitic virus, and various other acrimonies.—Dose, from ten to twenty drops, morning and evening, in an appropriate vehicle.

I.

By Gluten.

TADDEI'S PILLS.

R. Wheat Flour, one pound.

Make a paste with distilled water, and knead it under a stream of water; leave the gluten to itself for twenty-four hours in winter, and for eight or ten in summer; after which press it and agitate it with a solution made with

Amygdaline Soap, half an ounce.
 Water, one pound,

till a homogeneous liquor is produced, and strain it through a sieve. Then dissolve

Deuto-chloruret of Mercury, one dram
 in a sufficient quantity of water; pour into it an excess of glutinous liquor, till flocules cease to be formed. After several hours' repose, collect these on a cloth and dry them; then reduce them to a paste with

Extract of Guaiac, or of Gentian,
 a sufficient quantity,
 and divide the mass into one hundred and forty-four pills.

TADDEI extols these, in the treatment of syphilitic disorders, in the dose of from two or three, to six, eight, and even more daily. But they should be proscribed; for their intimate composition is not yet known. The deuto-chloruret is decomposed, or eight of them could never be taken at once, which

would be equivalent to four grains of this substance.

J.

By Gratiola.

ANTI-SYPHILITIC ELECTUARY. (AUGUSTIN; PH. SYPH.)

R. Deuto-chloruret of Mercury, three grains.
 Extract of Gratiola three drams.
 Rob of Elder, three ounces.

Mix.—HOLL prescribed it in painful tumours attributed to syphilis.

DEHAEN'S ANTI-SYPHILITIC MIXTURE.
 (SAINTE-MARIE)

R. Deuto-chloruret of Mercury, three grains.
 Extract of Gratiola, two drams.
 Aconite, one dram.
 Rob of Elder, six ounces.

Mix.—It is recommended in chronic syphilis.—Dose, a tea-spoonful, twice or thrice a day, gradually increasing the quantity of the gratiola to half an ounce.

K.

By Opium.

PILLS OF DEUTO-CHLORURET OF MERCURY AND OF OPIUM. (CAST. BORUS.; AUGUSTIN)

R. Deuto-chloruret of Mercury, ten grains.
 Opium, five grains.
 Hot Water, a sufficient quantity
 to dissolve them. Add

Grated Crumb of Bread, six scruples.
 Extract of Liquorice, four scruples.

Mix, and make one hundred pills. Dose, one or two, morning and evening. (CAST. BORUS.)

R. Deuto-chloruret of Mercury, two grains.
 Water, a sufficient quantity.

Dissolve, and add

Opium, four grains.
 Camphor, eight grains.
 Sugar, half a dram.
 Crumb of Bread, a sufficient quantity.

Make forty pills.—Dose, from three to five, morning and evening. (AUGUSTIN)

ANTI-SYPHILITIC PILLS. (RATIER; SAINTE-MARIE)

R. Deuto-chloruret of Mercury, four grains.
 Water, a sufficient quantity.
 Aqueous Extract of Opium, twelve grains.

Extract of Cinchona, one scruple.
 Simple Syrup, a sufficient quantity.

Make sixteen pills, each of which contains one-fourth of a grain of deuto-chloruret. (SAINTE-MARIE)

DUPUYTREN has thus modified these pills proposed by FRANK :

R. Deuto-chloruret of Mercury, } of each,
 Extract of Opium, } half a grain.
 Cinchona, ten grains.
 Powdered Cinchona, a sufficient quantity

to make two pills. (RATIER)

OPIATE ALCOHOLIC SOLUTION OF DEUTO-CHLORURET OF MERCURY. (MIL. COPEN.; AUGUSTIN; NIEMANN)

R. Deuto-chloruret of Mercury, one grain.
 Alcohol, four ounces.
 Tincture of Opium, one dram.

Mix. (MIL. COPEN.)

R. Deuto-chloruret of Mercury, two grains.
 Vinous Cinnamon Water, one ounce.
 SYDENHAM'S Laudanum, one scruple.

Mix. (NIEMANN)

AUGUSTIN prescribes four grains of the salt, three ounces of vinous cinnamon water, and two scruples of tincture of opium. The first formula deserves the preference.

SPANGENBERG'S LIQUOR. (PIERQUIN; SAINTE-MARIE)

R. Deuto-chloruret of Mercury, two grains.
 Tincture of Opium, one dram.
 Rose Water, four ounces.

Mix.—It is recommended, as a preventive of syphilis, to be applied as a lotion after a suspicious coitus.

OPIATE MERCURIAL INJECTION. (RATIER)

R. Deuto-chloruret of Mercury, twelve grains.
 Distilled Water, two pounds.
 SYDENHAM'S Laudanum, one ounce.

Mix.—To be applied to the syphilitic ulcers of women.

CONRAD'S OPHTHALMIC LIQUOR.

Aqua ophthalmica Conradi seu mercurialis seu Hydrargyri muriatici corrosivi cum Opio, Collyrium ex Hydrargyro muriatico corrosivo. (MIL. COPEN.; HANN.; WURCEB.; AUGUSTIN; RATIER; VAN MONS)

R. Deuto-chloruret of Mercury, one grain.
 Infusion of Elder Flowers, four ounces.
 Tincture of Opium, one dram.

Mix. (MIL. COPEN.)

- R. Deuto-chloruret of Mercury, one grain.
 Elder Flower Water, ten drams.
 Tincture of Opium, ten drops.
 Extract of Liquorice, one dram.

Mix.—Twenty drops of this solution to be taken every three hours. (AUGUSTIN)

- R. Deuto-chloruret of Mercury, one grain.
 Rose Water, four ounces.
 Extract of Opium, two grains.

Mix. (WURCEB.)

HANN.; and VAN MONS prescribe one grain of deuto-chloruret, six ounces of rose water, and two scruples of SYDENHAM'S laudanum.—RATIER—one grain of the salt, one scruple of tragacanth, four ounces of rose water, and eighteen drops of laudanum.—VAN MONS—two grains of the salt, half a scruple of extract of opium, four ounces of rose water, and half an ounce of mucilage of quince seeds.—AUGUSTIN—one ounce of chamomile water, and twenty grains of quince seeds, to be made into mucilage, and one grain of deuto-chloruret added.

MERCURIAL GARGLE. (RATIER)

- R. Dried Leaves of Hemlock, half an ounce.
 Water, one pound and a half.

Boil, strain, and add

Deuto-chloruret of Mercury, three grains.
 SYDENHAM'S Laudanum, one dram.
 Honey, two ounces.

Mix.

WRIGHT'S ELIXIR. (VAN MONS)

- R. Opium, one dram.
 Pimento, two drams.
 Seneka Root, three drams.
 Proof Spirit, eighty-two ounces.

Infuse without heat; and after sufficient extraction, express, and add to the liquor

Guaiac Resin, ten drams,
 and when dissolved,

Deuto-chloruret of Mercury, half a dram.

Dissolve, and filter.

SOLUBLE MERCURIAL BOUGIES. (AUGUSTIN)

- R. Deuto-chloruret of Mercury, four grains.
 Extract of Opium, one dram.
 Water, two ounces.
 Gum Arabic, a sufficient quantity

to make a thick liquid into which threads of cotton are to be plunged several times, letting them dry after each dipping.—

HECKER recommends them in chronic gonorrhoea.

L.

By Lead.

ESCHAROTIC TROCHISCS.

Trochisci de Minio. (GALL.; SARD.; WIRTEM.; CADET DE GASSICOURT; SPIELMANN; VAN MONS)

- R. Deuto-chloruret of Mercury, two parts.
 Deutoxide of Lead, one part.
 Grated Crumb of Bread, four parts.
 Rose Water, a sufficient quantity.

Make trochiscs of the volume and figure of oats. (GALL.; SPIELMANN)

WIRTEM. prescribes one part of deuto-chloruret, one of deutoxide, two parts of starch, two of crumb of bread, and a sufficient quantity of mucilage of tragacanth.—VAN MONS—two of deuto-chloruret, one of deutoxide, eight of crumb of bread, and a sufficient quantity of water.—SARD.—one dram of deuto-chloruret, one dram of deutoxide, one ounce of crumb of bread, and water.—CADET DE GASSICOURT—one ounce of deuto-chloruret, half an ounce of deutoxide, six ounces of crumb of bread, and water.

MERCURIO-SATURNINE INJECTION.

Injectio hydrargyro-saturnina seu de Plumbo et Mercurio, Solutio hydrargyrata. (PH. SYPH.; SWEDIAUR)

- R. Crystallized Protoxide of Lead, one ounce.
 Deuto-chloruret of Mercury, half a dram.
 Vinegar, five ounces.

Digest for twelve hours in a warm place, frequently shaking the vessel; pour off the clear part of the liquor, and to every two or four drams of it, add

Distilled Water, four ounces.

Mix. (PH. SYPH.)

- R. Deuto-chloruret of Mercury, } of each,
 Crystallized Acetate of Lead, } half a dram.
 Rose Water, one pound.

Dissolve. (PH. SYPH.)

- R. Deuto-chloruret of Mercury, two grains.
 Water, sixteen ounces.
 Dissolve, and add
 Solution of Sub-acetate of Lead, thirty drops.

Mix. (PH. SYPH.)

SWEDIAUR prescribes half a grain of

deuto-chloruret, four ounces of water, and eight drops of solution of sub-acetate; or, half a scruple of deuto-chloruret, fifteen ounces of water, and two drams of the solution; or, half a dram of deuto-chloruret, five ounces of solution, and the addition of one pound of water to every dram of the mixture of these two triturated together.

It is a bad preparation.—It has been recommended in both chronic and acute urethritis.

DESICCATIVE LOTION.

Lotio Plumbi composita. (SWEDIAUR)

R: Solution of Sub-acetate } of each,
of Lead, } half
Sulphate of Zinc, } an ounce.
Deuto-chloruret of Mercury, }
twenty grains.
Water, } two pounds.

Mix.—It is a bad preparation. It has been recommended in certain pruriginous eruptions of the genital organs.

CARDINAL DE LUYNES' ANTI-HERPETIC LIQUOR. (BORIES; CADET DE GASSICOURT; PIERQUIN)

R. Deuto-chloruret of Mercury, } one dram and a half.
Sub-carbonate of Lead, } four drams.
Alum, } three drams.
White of Egg, } one.
Rose Water, } eight ounces.

Mix. (PIERQUIN)

BORIES; and CADET DE GASSICOURT prescribe half an ounce of deuto-chloruret, two ounces of sub-carbonate, one ounce of alum, and two pounds of water.—BORIES very amusingly gives this formula as his own.

Compresses are to be moistened with it and applied to herpetic eruptions. It is a bad preparation, and should be used with caution.

PLENK'S CAUSTIC LIQUOR.

Liquor ad Condylomata. (HERBIP.; PH. SYPH.; AUGUSTIN; CADET DE GASSICOURT; SWEDIAUR; VAN MONS)

R. Deuto-chloruret of Mercury, } of each,
Alum, } one ounce.
Sub-carbonate of Lead, } of each,
Camphor, } one dram.
Alcohol, } of each,
Vinegar, } twelve ounces.

Mix, and preserve without filtering. (HERBIP.; VAN MONS)

CADET DE GASSICOURT prescribes two drams each, of proto-chloruret, of alum, of sub-carbonate of lead, and of camphor, and

one pound and a half each, of alcohol and of vinegar.—PH. SYPH.—one dram of deuto-chloruret, half a dram each, of alum, of camphor, and of crystallized acetate of lead, and half an ounce each, of alcohol and of vinegar.—AUGUSTIN; and SWEDIAUR suppress the preparation of lead, and prescribe one dram of deuto-chloruret, half a dram of alum, half a dram of camphor, half an ounce of vinegar, and half an ounce of alcohol.

Bits of lint are wetted, and syphilitic excrescences covered with them. It is a bad preparation, and the following should always be preferred:

STETTIN'S LIQUOR.

Aqua mercurialis seu Stettinensis ad usus externos. (BRUNS.; WIRTEM.)

R. Pomegranate Flowers, } half an ounce.
Super-acetate of Lead, } of each,
Super-tartrate of Potass, } one dram.
Root of Tormentil, } two drams.
Saccharated Alum, } one ounce.
Deuto-chloruret of Mercury, } of each,
Red Sanders Wood, } one dram.
Rectified Alcohol, } sixteen ounces.

Digest, and express. (BRUNS.)

R. Sub-carbonate of Lead, } one ounce.
Alum, } six drams.
Crystallized Protoxide of Lead, } half an ounce.

Deuto-chloruret of Mercury, } of each,
Hydrochlorate of Ammonia, } one dram.
Water of Garden Nightshade, } of each,
Water Lily, } four ounces.
Rose Water, }
Plantain Water, }
Vinegar, } half a pound.

Boil slightly, and preserve without straining. (WIRTEM.)

DESICCATIVE OINTMENT.

Unguentum siccativum (SWEDIAUR)

R. Prepared Tutty, } of each,
Deuto-chloruret of Mercury, } one scruple.
Red Oxide of Mercury, } of each,
Acetate of Lead, } two drams.
Calcined Alum, } four ounces.
Lard, }

Triturate.

M.

By Rhubarb.

MERCURIAL PILLS. (FEE; RATIER)

R. Deuto-chloruret of Mercury, } four grains.

Alcohol, a few drops.
 Rhubarb, } of each,
 Guaiac Resin, } one dram.
 Extract of Liquorice, a sufficient quantity

to make pills of five grains each, (FEE)

RATIER prescribes one scruple of deuto-chloruret, two drams of rhubarb, half an ounce of soap, and a sufficient quantity of liquorice powder and of syrup, to make one hundred and forty-four pills.

The latter is unchemical.

N.

By Scammony.

MERCURIAL PILLS OF SCAMMONY.

Pilule Scammonii hydrargyrosæ seu e Scammonio composiæ. (SUEC.)

R. Deuto-chloruret of } of each,
 Mercury, } twelve grains.
 Scammony, }
 Gamboge, five grains.
 Simple Syrup, a sufficient quantity
 to make six pills. (!)

By adding one dram of extract of rhubarb, and making twenty-four pills, a more commodious and less dangerous preparation would be obtained.

O.

By Senna.

LASSÔNE'S ANTI-SYPHILITIC PILLS. (SAINTE-MARIE)

R. Deuto-chloruret of Mercury, eighteen grains.
 Proto-chloruret of Mercury, fifty-four grains.

Triturate for a long time, and add

Guaiac Resin, half a dram.
 Senna, in powder, one dram.
 Syrup, a sufficient quantity

to make pills of two grains each.

Dose, three daily, to be gradually augmented.—It is a bad preparation.

IODURETS OF MERCURY.

I. *Proto-Ioduret.*

FERR.; HENRI; MAGENDIE.

R. Crystallized Proto-nitrate of Mercury, one hundred parts.
 Water, four hundred parts.

Dissolve, filter, and add

Ioduret of Potassium, dissolved in water acidulated with Nitric Acid, a sufficient quantity,

till a precipitate ceases to be formed.

Collect this on a filter, wash it with distilled water, till a black precipitate ceases to be produced by the addition of potass to the water, then dry it, and preserve it in a dark place.

II. *Deuto-Ioduret.*

FERR.; MAGENDIE.

R. Deuto-chloruret of Mercury, seventy parts.
 Ioduret of Potassium, one hundred parts.

Dissolve each separately in a sufficient quantity of distilled water; filter the two liquors, and mix them slowly together: collect the precipitate on a filter, wash it with distilled water till this comes off tasteless; then dry and pulverize it, and preserve in a dark place. (MAGENDIE)

R. Solution of Hydriodate of Potass, any quantity.

Add to it drop by drop

Solution of Deuto-chloruret of Mercury, a sufficient quantity, or, till it ceases to produce a precipitate. Wash and dry this. (FERR.)

The iodurets of mercury are powerful stimulants; the deuto-ioduret is even more energetic than the deuto-chloruret, and is employed in the treatment of syphilis complicated with scrofula. When left on the skin for a few moments, the deuto-ioduret has sometimes brought on an intense erysipelatous inflammation. The commencing dose is one-sixteenth of a grain, to be carried gradually to one-fourth.

ALCOHOLIC SOLUTION OF DEUTO-IODURET OF MERCURY. (FERR.; FEE; HENRI; MAGENDIE)

R. Deuto-ioduret of Mercury, twenty grains.

Alcohol (36 degrees), one ounce and a half.

Dissolve. (FEE; HENRI; MAGENDIE)

Twenty-six drops contain nearly one eighth of a grain of the ioduret.

FERR. prescribes only one ounce of alcohol, in which solution there are in every ten drops, a sixteenth of a grain.—Dose, ten, fifteen, and twenty drops in a glassful of distilled water; common water decomposes it.

ÆTHEREAL SOLUTION OF DEUTO-IODURET OF MERCURY. (FEE; MAGENDIE)

R. Deuto-ioduret of Mercury, twenty grains.

Sulphuric Æther, one ounce and a half.

Mix.—The dose of this should be less than that of the preceding.

PILLS OF IODURET OF MERCURY. (FERR.; BRERA; FEE; HENRI; MAGENDIE)

R. Proto or Deuto-ioduret of Mercury, one grain.
Extract of Liquorice, twelve grains.
Liquorice Powder, a sufficient quantity,
to make eight pills. (MAGENDIE)

FEE; and HENRI substitute extract of of juniper for that of liquorice.

R. Deuto or Proto-ioduret of Mercury, one grain.
Liquorice Powder, six grains.
Rob of Elder, a sufficient quantity.
Make eight pills. (FERR.)

R. Proto or Deuto-ioduret of Mercury, one grain.
Extract of Juniper, sixteen grains.
Liquorice Powder, a sufficient quantity,
to make eight boluses. (BRERA)

Dose, two, morning and evening, increasing the dose gradually till four are taken each time.

OINTMENT OF IODURET OF MERCURY. (BRERA; HENRI; MAGENDIE; PIERQUIN; RATIER)

R. Proto or Deuto-ioduret of Mercury, twenty grains.
Prepared Lard, one ounce and a half.
Triturate. (MAGENDIE)

R. Deuto-ioduret of Mercury, fifteen grains.
Lard, two ounces.
Essence of Bergamotte, twenty drops.
Mix. (RATIER)

R. Proto-ioduret of Mercury, half a dram.
Lard, one ounce and a half.
Essence of Bergamotte, fifteen drops.
Mix. (RATIER)

PIERQUIN prescribes fifteen grains of ioduret, one ounce of lard, and twenty drops of essence of cloves.

It is recommended to hasten the cicatrization of indolent ulcers. That with the deuto-ioduret is more stimulant than the other, and should be employed in less quantity.

CYANURET OF MERCURY.

Borussias seu *Cyanuretum Hydrargyri*. (BELG.; GALL.; FERR.; NIEMANN; PLANCHE; VAN MONS)

R. Cyanuret of Iron, six parts.
Nitric Oxide of Mercury, five parts.

Distilled Water, forty parts.

Mix the two former together, and add the latter heated. Boil for half an hour, constantly stirring, and then filter through paper. Wash the residuum with hot distilled water. Lastly, evaporate the liquor, filter, and set apart that crystals may form. (DUBL.)

R. Prussian Blue, two parts.
Red Oxide of Mercury, one part.

Pulverize these two substances, mix them together, and put them into a glass capsule. Pour upon them

Pure Water, eight parts.

Boil, constantly stirring with an iron spatula, till the mixture has acquired a yellow tint; then filter the liquor and wash the residuum with

Water, four parts.

Evaporate the mixed liquors till a drop being placed on a cold body becomes solidified; then set it aside in a cool place to crystallize, and purify with fresh crystallization. (BELG.; FERR.; GALL.)

HENRI recommends the Prussian blue to be first washed with diluted sulphuric or hydrochloric acid, in order to take from it the alumina, without which one would be acting on an uncertain and variable quantity of ferro-cyanate of iron.

R. Red Oxide of Iron, ten parts.
Prussian Blue, seven parts.

Pulverize each separately, adding a little alcohol to the second; put the powders into a matrass with

Water, eighty parts.

Boil in a sand-bath till the matter has assumed a yellowish-green tint; filter twice with the same paper; wash the residuum with cold water; put it again in the bath, and remove the fire; set it aside to crystallize slowly, and purify with a second crystallization. (NIEMANN; VAN MONS)

R. Red Oxide of Mercury, twelve drams.
Prussian Blue, eight ounces.
Distilled Water, twenty-seven ounces.

Levigate each separately, and put them into a porcelain capsule with nine ounces of the water; boil for half an hour, constantly stirring, then decant and filter: pour upon the residuum the remainder of the water, boiling; boil again for half an hour, and filter. Mix the two liquors, evaporate to a pellicle, and set aside to crystallize in a cool place. (PLANCHE)

This is a formidable poison; it has been tried in syphilitic cases, but its deleterious activity, which is still greater than that of the deuto-chloruret, should make it be

shunned. It is administered in the same manner but in smaller doses than the preceding.

CHAUSSEIER'S ANTI-SYPHILITIC LIQUOR.
(RATIER)

- R. Cyanuret of Mercury, eight grains.
Distilled Water, one pound.

Mix.—Each ounce contains half a grain, of cyanuret.

OINTMENT OF CYANURET OF MERCURY.
(RATIER)

- R. Cyanuret of Mercury, sixteen grains.
Lard, one ounce.
Essence of Lemon, fifteen drops.

Mix.—It is recommended in moist scaly tetter, accompanied with inflammation and violent itching.

AMMONIURET OF MERCURY.

Ammoniuretum Hydrargyri. (NIEMANN;
PLANCHE; VAN MONS)

- R. Red Oxide of Mercury, two parts.
Concentrated Solution of Ammonia, one part.

Shake quickly together, and then set aside for several hours; shake again, and continue thus till the matter has taken a crystalline appearance; then decant, wash, dry, pulverize, wash again, and finally dry and preserve it. (NIEMANN; VAN MONS)

- R. White Precipitate, any quantity.
Solution of Ammonia, a sufficient quantity

to dissolve it. (PLANCHE)

The ammoniuret of mercury has been proposed to be applied externally for the dispersion of exostoses.

LINIMENT OF AMMONIURET OF MERCURY.
(PLANCHE)

- R. Olive Oil, nine drams.
Ammoniuret of Mercury, half an ounce.
Triturate together in a glass mortar.

BORATE OF MERCURY.

Sal sedativum mercuriale, Boras Hydrargyri Mercurius boraxitus. (NIEMANN;
SPIELMANN; VAN MONS)

- R. Mercury, } of each,
Concentrated Nitric Acid, } two ounces.

The solution being effected, dilute the liquor with four parts of water, filter it, and add, gradually,

Solution of Sub-borate of Soda, a sufficient quantity,

or, till no more precipitate is formed; wash and dry this. (SPIELMANN; VAN MONS)

VOL. II.

- R. Proto-chloruret of Mercury, 22,2 parts.
Neutral Borate of Soda, 26,5 parts.

Triturate together in a glass mortar; at the end of a quarter of an hour add a little water, then more, and finally a large quantity: decant the liquor, wash the precipitate till the water comes off tasteless, and dry it. (NIEMANN; VAN MONS)

This salt may be employed instead of the proto-chloruret of mercury. Dose, two grains daily, to be gradually augmented.

CARBONATE OF MERCURY. (FERR.;
VAN MONS)

- R. Saturated Solution of Deuto-chloruret of Mercury, four parts.
Water, charged with Carbonic Acid, sixteen parts.

Pour these into a flask, and add gradually, shaking briskly, a solution of

Neutral Carbonate of Potass, one part.

Leave the flask in a cool place for fifteen days or more; then wash slightly the crystals which form at the surface of the liquor, and dry them. (VAN MONS)

- R. Acetate of Mercury, any quantity.
Boiling Water, a sufficient quantity
to dissolve the salt; filter the liquor while hot, and add to it,

Solution of Subcarbonate of Ammonia, a sufficient quantity, or, till no more precipitate is formed; wash and dry this. (FERR.)

WURTZ' PRECIPITATE.

Mercurius præcipitatus fuscus Wurtzii, Turpethum rubrum. (LIPP.)

- R. Solution of Proto-nitrate of Mercury, two pounds.
———— Subcarbonate of Potass, a sufficient quantity.

Add the latter to the former, till precipitate ceases to be formed; collect this, wash it with hot water, and dry it.

PHOSPHATE OF MERCURY.

Hydrargyrum phosphoratum seu phosphoreum, Phosphas Hydrargyri. (OLDEN;
BORUS.; COXE; NIEMANN; SWEDIAUR;
VAN MONS)

- R. Solution of Nitrate of Mercury, any quantity.
———— Phosphate of Soda, a sufficient quantity.

Add the latter to the former till no more precipitate is formed; wash, and dry this. (OLDEN.; BORUS.; NIEMANN; SWEDIAUR; VAN MONS)

- R. Deuto-chloruret of Mercury, any quantity.

M

Dissolve it in hot water, and add gradually,

Hot Solution of Sub-phosphate of
Soda, a sufficient quantity,

or, till no more precipitate is formed; wash and dry this. (VAN MONS)

R. Proto-chloruret of Mercury, one part.
Sub-phosphate of Soda, two parts.

Mix, and add, while trituring,

Hot Water, four parts.

At the end of an hour, pour off the liquid, wash the precipitate and dry it on a filter. (VAN MONS)

COXE requires the solution of nitrate of mercury to be decomposed by that of sub-phosphate of soda.

This is rarely used, but it once was in vogue in Germany and England, for diseases called degenerated and obstinate syphilis. — Dose, from half a grain to one grain, with sugar or cinnamon; it is also applied externally, in the form of an ointment. BOUDER thinks that we might substitute for this and many analogous preparations, the *Ammoníaco-mercurial Phosphate*, which he has discovered, and which is possessed of very great solubility. It enters, with proto-chloruret of mercury, and probably other mercurial salts, into the composition of the *Rose Precipitate* or *Mineral Rose*, obtained by pouring urine into a solution of nitrate of mercury, and which was formerly prescribed as purgative, in the dose of from six to ten grains.

PERSULPHATE OF MERCURY.

Hydrargyri Persulphas. (DUBL.)

R. Purified Mercury, } of each,
Sulphuric Acid, } six parts.
Nitric Acid, one part.

Expose the mixture in a glass vessel, to a gradually raised heat, till the matter, when nearly dry, assumes a white colour.

SUB-DEUTO-SULPHATE OF MERCURY.

Calx Mercurii vitriolata, Hydrargyrus vitriolatus seu flavus seu corrosivus flavus seu emeticus flavus seu præcipitatus flavus seu causticus flavus seu luteus, Oxydum Hydrargyri sulphuricum, Sulphas Hydrargyri luteum seu cum excessu Oxydi, Subsulphas Hydrargyri flavus, Turpethum minerale. — Yellow Precipitate, Turbith mineral. (DUBL.; EDINB.; AMER.; AMST.; BRUNS.; GALL.; FERR.; GENEV.; LIPP.; LUSIT.; SARD.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; NIEMANN; SPIELMANN; SWEDIAUR; VAN MONS)

Subsulphas Hydrargyri flavus.
(EDINB.)

R. Purified Mercury, two parts.
Sulphuric Acid, three parts.

Put them into a glass cucurbit placed in a sand-bath, and boil to dryness. Reduce the white residuum to powder, and throw it into boiling water; it will then be immediately converted into a yellow powder, which is to be frequently washed with warm water.

Sulphuricum Oxydum Hydrargyri.
(DUBL.)

R. Persulphate of Mercury, one part.
Boiling Water, twenty parts.

Triturate these together in a stone mortar, and pour off the supernatant liquor: wash the yellow powder which results with hot distilled water as long as the addition of solution of caustic potass to the effused water effects a precipitate, and finally dry it.

R. Mercury, one part.
Sulphuric Acid, two parts.

Heat gradually till a white mass is formed; dry this perfectly by augmenting the heat; then sprinkle it with distilled water. Triturate it well in a glass mortar, set it aside, decant the liquid, and wash the powder at the bottom of the vessel with distilled water, till this comes off tasteless. (AMST.; GENEV.; LUSIT.; SARD.; HERBIP.)

NIEMANN prescribes two parts of mercury one and a half of acid.—BRUGNATELLI.—three of mercury and four and a half of acid.—GALL.—two of mercury and two and a half of acid (66 degrees)—AMER.; COXE; and SPIELMANN.—two of mercury and three of acid.—BRUNS.; FERR.; and WIRTEM.—an equal part of each.—VAN MONS.—190 of mercury and 200 of acid.

LIPP. directs five ounces of mercury to be dissolved in six ounces of concentrated sulphuric acid, and having dried and pulverized the salt, the powder to be introduced into a flask containing two pounds of boiling water and one ounce of deliquesced subcarbonate of potass, and the precipitate to be carefully washed.

R. Sulphuric Acid diluted with its own weight of water, any quantity.
Red Oxide of Mercury, a sufficient quantity.

to saturate the acid. Evaporate to dryness, pulverize the mass, pour upon it ten times its weight of boiling water, agitate, set aside, decant, and wash and dry the precipitate. (VAN MONS)

It is violently emetic and purgative, and was formerly represented as a preservative from the small pox; but is scarcely employed at the present day, except in obstinate exanthemata, in which cases it is still reputed to have more efficacy than any other mercurial preparation. — Dose, from one fourth of a grain repeated twice or thrice a day, to one, two, or three grains, and this should not be exceeded.

EMETIC BOLUSES. (PIERQUIN; SAUNDERS)

R. Sub-deuto-sulphate of Mercury, seven grains.
Conserve of Roses, one scruple.

Make one bolus. (SAUNDERS)

PIERQUIN prescribes five grains of the salt.

These are both too strong; three grains would be sufficient.—It would be still better if such an emetic were renounced altogether.

OINTMENT OF SUB-DEUTO-SULPHATE OF MERCURY.

Unguentum Sulphatis Hydrargyri seu Turpethi mineralis. (BRUGNATELLI; CADET DE GASSICOURT; NIEMANN; RATIER; SAINTE-MARIE; SWEDIAUR)

R. Sub-deuto-sulphate of Mercury, one part.
Lard, eight parts.

Triturate the salt in a glass mortar, incorporate it in the lard melted slowly, and triturate till cold. (SAINTE-MARIE; SWEDIAUR)

BRUGNATELLI prescribes one part to four.—CADET DE GASSICOURT; and RATIER—one to sixteen.

R. Sub-deuto-sulphate of Mercury, one part.
Simple Cerate, ten parts.

Mix. (NIEMANN)

ALBERT frequently employs this ointment in the treatment of indolent tetter.

CULLERIER'S ANTI-HERPETIC OINTMENT. (PIERQUIN; RATIER)

R. Sub-deuto-sulphate of Mercury, } of each,
SYDENHAM'S Laudanum, } one dram.
Sublimed Sulphur, half a dram.
Lard, one ounce.
Triturate together.

CHEVALLIER'S ANTI-HERPETIC OINTMENT. (HENRI)

R. Sub-deuto-sulphate of Mercury, two drams.
Chloruret of Lime, three drams.
Oil of Almonds, six drams.
Lard, two ounces.
Mix.

SULPHATE OF MERCURY. (BRUGNATELLI; HENRI; VAN MONS)

R. Mercury, three ounces.
Sulphuric Acid, four ounces and a half.

Heat in a retort placed in a sand-bath, till sulphureous acid begins to rise; then remove it from the fire, collect the saline mass which is formed, and preserve it in a glass bottle. (BRUGNATELLI)

VAN MONS directs: first, to mix together ten parts and a half of red oxide of mercury and five of sulphuric acid, adding water very gradually, and when the union is effected, to evaporate to dryness:—second, to boil together an equal part of each, of mercury and of acid to perfect dryness:—third, to heat together nine parts and a half of mercury and ten of acid, to evaporate to dryness, and to triturate the mass, still hot, with 4,75 of mercury.

These four processes give deuto-sulphate mixed with a certain quantity of proto-sulphate, with the exception perhaps of the second.

Proto-sulphate of mercury is obtained by decomposing the pure proto-nitrate, dissolved in water acidulated with nitric acid, by sulphate of soda or of potass, carefully washing the precipitate and drying it in a dark place. (HENRI)

Neither of these are used in medicine, but they are employed, the first especially, in several pharmaceutic operations.

AMMONIACO-MERCURIAL SUB-DEUTO-SULPHATE. (BRUGNATELLI; SWEDIAUR)

R. Mercury, two pounds.
Sulphuric Acid, three pounds.

Digest in a water-bath till dissolved, then dilute the liquor with water; separate the precipitate of sub-deuto-sulphate of mercury which forms, and pour gradually into the supernatant mercurial super-sulphate, solution of subcarbonate of ammonia or of pure ammonia till no more precipitate is formed; wash this perfectly, and dry it with slow heat.

This preparation is only used in making that which BRUGNATELLI improperly calls ammoniuret of mercury.

HYPOSULPHITE OF MERCURY AND OF POTASS, SULPHURETTED SULPHITE OF POTASS AND OF MERCURY. (VAN MONS)

R. Hyposulphite of Potass, two parts and a half.
Red Oxide of Mercury, one part.

Dissolve the salt in boiling water, and the powdered oxide, and stir till it is dissolved; filter the liquid while hot, and set it apart to cool. Wash the crystals which form with a little cold water, dissolve them in the least possible quantity of hot water, and crystallize again.

In the same manner are prepared the *Hyposulphite of mercury and of soda*, and the *H. of mercury and of ammonia*.

SUB-PROTO-NITRATE OF MERCURY.

Calx cinerea Mercurii, Mercurius solubilis Hahnemanni, Hydrargyrum oxydatum nigrum seu oxydulatum nigrum Oxydulum Hydrargyri Hahnemanni, Oxydum Hydrargyrum nigrum seu ammoniacale seu nitrico-ammoniacale seu salinum, Oxodes Hydrargyri nigrum, Oxynitras Hydrargyri et Ammonia, Præcipitatus solubilis niger, Protoxydum Hydrargyri nitrico-ammoniacale, Pulvis Hydrargyri cineris, Turpetium nigrum.—Black Precipitate, Black Turbith Mineral. (ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; GALL.; FORM. HÔP. FR.; HAMB.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; BRUGNATELLI; CADET DE GASSICOURT; SWEDIAUR; VAN MONS)

R. Nitric Acid, five ounces.

Pour it into a matrass plunged into a large vessel full of cold water in a cool place; add

Purified Mercury, half an ounce.

When the metal is dissolved, add one ounce and afterwards three ounces and a half more, adding cold water to the bath in proportion as it becomes heated. When crystals are formed which the acid refuses to dissolve, let the mass evaporate spontaneously, and pour off the liquid. Wash the crystals with a little distilled water, and dry them on a filter without heat. Then dissolve them in five pounds of distilled water, filter the solution, saturate it with solution of ammonia, shaking it; then set it apart; pour off the supernatant liquor; wash the precipitate with distilled water, dry it in the air without heat, and pulverize it. (LIPP.)

This is the process as HAHNEMANN himself has detailed it. The process is essentially the same in every pharmacopœia which gives it after him.

R. Mercury, three parts.
Nitric Acid (33 degrees), two parts.

Pour them into a phial placed in a sand-bath; heat moderately, and keep up ebullition till crystals begin to form; then remove it from the fire, agitate to determine a confused crystallization, pour the matter into a glass mortar, triturate, adding a small quantity of distilled water mixed with a little nitric acid. Set aside, decant, and repeat the same operation on the residuum till no more mercury remains undissolved. Mix the liquors, and add solution of ammonia drop by drop, constantly shaking. Finally, set aside, wash the powder several times, then let it drip on a filter, and dry it in a stove. (FORM. HÔP. FR.)

The operatory process is very nearly the same every where; but BRUNS. prescribes the solution of the sub-carbonate of ammonia.

HOLZE has proposed the following process:

R. Crystallized Proto-nitrate of Mercury, eight parts.
Nitric Acid (1.250), one part.
Boiling Distilled Water, eighty parts.

To the solution when cold, add

Solution of Ammonia (0.95), four parts,

diluted with

Distilled Water, eight parts.

Agitate, filter, and dry the precipitate. (FEE)

SOUBEIRAN has ascertained that by the process indicated in the greater number of pharmacopœias, a salt is obtained composed of sub-nitrate of mercury with a quantity of ammoniaco-mercurial proto-nitrate varying according to the process followed, and depending on the greater or less acidity of the proto-nitrate employed, making the product more or less deep in colour, and also more or less soluble in nitric acid, the single sub-nitrate being black, and the double being white. He concludes from thence that the proto-nitrate should be dissolved in the smallest quantity possible of nitric acid, pouring at many times upon the crystallized nitrate, water weakly acidulated with it, with less, in fact, than is known to be necessary to dissolve it entirely, prolonging the trituration in order to saturate the liquor perfectly before decanting it. These observations are of very great importance in a chemical point of view, but in no other respect do they interest us. It matters little to the practitioner whether the sub-nitrate of mercury contain ammoniaco-mercurial proto-nitrate or not, or whether it even contain ammoniaco-mercurial deuto-nitrate. The chemical properties of the latter are very similar to those of the double proto-nitrate,

and all these preparations act in the same manner on the system, namely, as stimulants, by inducing irritation, differing only from each other in an imperceptible degree, or in a degree not to be perceived in the excitement which they induce.

The sub-proto-nitrate of mercury has been very celebrated, particularly in Germany; it is administered in the dose of one grain with five or ten grains of any mucilaginous powder; half a grain or one grain of opium is usually added.

MERCURIAL TROCHISCS.

Trochisci Hydrargyri. (SWEDIAUR)

R. Sub-proto-nitrate of Mercury, one ounce.
Oil of Orange, twenty drops.
White Sugar, two ounces.
Mucilage of Gum Arabic, a sufficient quantity

to make trochiscs of ten grains each.

Dose, one, twice or thrice a day.

MERCURIAL BOLUS.

Bolus ex Hydrargyri. (SWEDIAUR)

R. Sub-proto-nitrate of Mercury, three grains.
Extract of Liquorice, a sufficient quantity.
Mix.

HAHNEMANN'S PILLS.

Pilula Hydrargyri seu ex Hydrargyro oxydulato nigro seu ex Mercurio solubili Hahnemanni. (WURCEB.; RATIER; SWEDIAUR; PH. SYPH.)

R. Sub-proto-nitrate of Mercury, fifteen grains.
Extract of Liquorice, half an ounce.
Simple Syrup, a sufficient quantity.
Make one hundred and fifteen pills. (WURCEB.)

SWEDIAUR; and PH. SYPH. prescribe one dram of sub-proto-nitrate, and a sufficient quantity of the extract to make a mass.

R. Sub-proto-nitrate of Mercury, twenty grains.
Gum Arabic, } of each.
Sugar, } half a dram.
Make thirty pills. (RATIER)

MERCURIAL DIURETIC PILLS.

Pilula ex Hydrargyro diuretica. (SWEDIAUR)

R. Sub-proto-nitrate of Mercury, half a dram.

Conserve of Squill,
from twenty to thirty grains.

Sulphuretted Oil, three drops.

Powdered Squill, a sufficient quantity
to make pills of four grains each.

They are recommended in mucous diseases of the lungs, dropsy, obstinate catarrh, and at the suspected commencement of phthisis.

Dose, three or four, every two days, at night.

MERCURIAL HONEY. (SWEDIAUR)

R. Sub-proto-nitrate of Mercury, one dram.
Despumated Honey, one or two ounces.
Triturate together.

SYRUP OF SUB-PROTO-NITRATE OF MERCURY.

Syrupus ex Oxydulo Hydrargyri Hahnemanni. (NIEMANN)

R. Sub-proto-nitrate of Mercury, twenty grains.
Gum Arabic, one dram.
Honey of Roses, a sufficient quantity.
Triturate, and add
Syrup of Marshmallows, three ounces.
Mix.—Dose, a spoonful, morning and evening.

PROTO-NITRATE OF MERCURY.

Calx Mercurii, Hydrargyrum nitricum, Mercurius nitrosus, Nitras Hydrargyri.

I. In the Solid State.

ANT.; DAN.; FERR.; LIPP.; WIRTEM.; HERBIP.; SWEDIAUR; VAN MONS.

R. Nitric Acid, (26 degrees), any quantity.

Pour it into a vessel surrounded with cold water or ice, and add to it at several times,

Mercury, a sufficient quantity,
till crystals appear in the liquor; set it aside for twenty-four hours, then collect, wash, and dry the crystals. (ANT.; DAN.; HISP.; FERR.; LIPP.)

R. Mercury, one ounce and a half.
Nitric Acid, one ounce.

Shake frequently the mixture for about four days, then leave it to spontaneous evaporation to obtain crystals; wash these and dry them in bibulous paper. (WIRTEM.)

HERBIP. directs an equal part each, of nitric acid and of mercury to be digested in a sand-bath; the liquor to be then evapo-

rated over a slow fire till there remains only a white dry mass.

R. Mercury, one part.
Nitric Acid, (28 degrees), two parts.

Dissolve the greater part of the metal in the acid with the aid of a moderate heat, then evaporate the solution, and set it apart to crystallize. (GALL.)

As by following this process a great deal of deuto-nitrate is obtained, HENRI has proposed the following :

R. Mercury, two hundred parts.
Nitric Acid (25 degrees), one hundred and eighty parts.

Dissolve till no more nitrous vapours appear, and add

Hot Distilled Water, one hundred parts.

Boil a little, decant, and crystallize. (HENRI)

BAVAR. gives a process analogous to this: place in a sand-bath a phial containing three parts of mercury, two of nitric acid, and two of distilled water, gradually augmenting the heat till no more aqueous vapours are disengaged, and then to let it crystallize.

VAN MONS requires mercury to be boiled in nitric acid diluted with thrice its weight of boiling water, taking care that undissolved metal always remains; the liquor to be decanted and crystallized.

II. In the Liquid State.

Liquor Hydrargyri nitrici, Nitras Hydrargyri liquidus, Liquor Nitratiss Hydrargyri seu Bellosti. (AMST.; ANT.; BRUNS.; DAN.; GALL.; FORM. HÔP. FR.; HANN.; OLDEN.; POL.; BORUS.; BRUGNATELLI.; NIEMANN.; PIERQUIN.; RATIER.; SPIELMANN.)

R. Mercury, one hundred and twenty parts.
Nitric Acid (33 degrees), one hundred and fifty parts.

Dissolve at a moderate heat, and add

Distilled Water, nine hundred parts.

Strain at the end of a few hours, and preserve. (GALL.; FORM. HÔP. FR.)

HANN. indicates two processes:—1. to put one ounce each, of mercury, of nitric acid, and of distilled water into a phial, to leave it plunged in cold water for eight or fifteen days till bubbles cease to be disengaged from it; the liquor thus obtained to be preserved, in a well-stopped flask, on a little metallic mercury, and to add a few drops of nitric acid, if any yellow powder happens to be formed.—2. to take one ounce of mercury and two ounces of nitric

acid or a sufficient quantity to dissolve the metal with the aid of a moderate heat.

ANT.; and BRUNS. direct two ounces of mercury to be dissolved in six of acid, and one ounce of the product to be dissolved in three pounds of distilled water.—SPIELMANN—the metal to be dissolved in an equal weight of nitric acid, and one dram of the salt thus obtained to be dissolved in two ounces of water.—OLDEN.—one ounce of metal to be dissolved in a sufficient quantity of nitric acid, and a sufficient quantity of water to be added to obtain four ounces of liquid.—POL.; and BORUS. to act in the same manner, to carry the total quantity to five ounces.—AMST. to dissolve, with heat, in the acid, as much mercury as it will take up, and to add to the liquor twelve parts of hot water.—LIPP.; NIEMANN.; and PIERQUIN to dissolve at a moderate heat one dram of mercury in two drams of acid, and to add twelve ounces of spring water.—DAN. to dissolve two drams of crystallized salt in one ounce of distilled water.

This preparation is scarcely used now, but it was formerly given in the dose of two or three drops in any sudorific drink.

It is occasionally employed in cases of chronic syphilitic ulceration. The salt itself has been lately recommended as an efficacious escharotic in the same affections.

MERCURIAL LIQUOR. (PIERQUIN)

R. Mercury, } of each, one ounce.
Nitric Acid. }

Dissolve and add

Plantain Water, two pounds.
Rose Water, } of each,
Gum Lac, } a sufficient quantity.

Mix.—For lotions and fomentations to old and atonic ulcers, pustules, aphthæ, and syphilitic herpetic eruptions.

MANNY'S MERCURIAL LOTION. (RATIER)

R. Mercury, two drams.
Nitric Acid, four ounces.

Dissolve, and add

Pure Water, ten pounds.

Mix.—For lotions in the treatment of itch.—Half an ounce to be used morning and evening.—It is a convenient and not expensive remedy, and does not stain the linen. It has also been found of service in prurigo formicans and in the morbus pedicularis. Two drams of camphor may be added to every two pounds, which is said to correct its irritating property.

GOHL'S GREY MERCURIAL LIQUOR.

Aqua grisea seu grysea Gohlii seu ad Synoviam. (BRUNS.; HASS.; PALAT.;

SARD.; WIRTEM.; CADET DE GASSICOURT; NIEMANN; PIDERIT; SPIELMANN; VAN MONS)

R. Mercury, } of each, one ounce.
Nitric Acid, }

Dissolve the metal in the acid, at a moderate heat, and add to the liquor the following decoction:

R. Root of Great Celandine, }
—— Round Birthwort, } of each,
Herb of Rue, } one
—— Scordium, } ounce.
Petals of Red Roses, }
Water, } five pounds.

Boil till reduced to four pounds. (BRUNS.; PALAT.; SARD.; WIRTEM.; NIEMANN.)

SPIELMANN directs a decoction of half an ounce of root of celandine, of one handful each, of round birthwort, of black nightshade, of rue, and of pale roses, in ten pounds of water, reduced to eight, to be added to a solution of one ounce of mercury in one ounce of nitric acid.

R. Mercury, } one ounce.
Nitric Acid, } two ounces.

Dissolve at a moderate heat, and to every half ounce of liquid add the expressed juice or decoction of

Root of Great Celandine, } of each,
—— Round Birth- } one pound.
wort, }

Mix. (HASS.; CADET DE GASSICOURT; PIDERIT)

R. Solution of Proto-nitrate of Mercury, }
Depurated Juice of Great Celandine, } one part.
twelve parts.

Mix. (VAN MONS)

This preparation used formerly to be frequently employed in lotions to syphilitic ulcers.

BELLET'S SYRUP.

Syrupus Belleti mercurialis. (AUGUSTIN; CADET DE GASSICOURT; HENRI; NIEMANN)

R. Proto-nitrate of Mercury, thirty parts.
Water, weakly acidulated, }
two hundred parts.

Dissolve, and add

Simple Syrup, three thousand parts,
previously concentrated; then add

Alcoholized Nitric Ether, ten parts.
Preserve. (HENRI)

CADET DE GASSICOURT directs after BOUILLON-LAGRANGE one dram and a half of nitrate of mercury to be dissolved in the least possible quantity of water, and the liquor to be mixed with one pound of simple

syrup and half a dram of nitric ether.—NIEMANN,—after VIREY, one dram of acetate of mercury, to be dissolved in a little distilled water, and two drams of nitric ether, and one pound of syrup of gum to be added.—AUGUSTIN—one dram and a half of solution of nitrate, half a dram of HOFFMANN'S liquor, and sixteen ounces of simple syrup, to be mixed together.

This syrup is very different to that which BELLOSTE prepared, and which was only a kind of sweetened nitric ether, without an atom of mercury. It is, like every other æthereal preparation of mercury, very inconstant and uncertain in its effects, on account of its tendency to become decomposed. It should not therefore be prescribed; it would be easy to substitute for it an extemporaneous mixture of a few drops of the solution, and three or four ounces of water sweetened with syrup of ether.

ZELLER'S PILLS.

Pilule ex Hydrargyro Nitrico. (AUGUSTIN; (SAINTE-MARIE)

R. Proto-nitrate of Mercury, ten grains.
Extract of Liquorice, forty grains.

Mix, and make sixty pills. (SAINTE-MARIE)

AUGUSTIN prescribes ten grains of crystallized nitrate and one hundred and ninety grains of crumb of bread, to make pills of two grains each.

They are anti-syphilitic.—Dose, one or two, in twenty-four hours, gradually carrying the dose to six daily.

MERCURIAL POTIÖN.

Potus mercurialis Belleti. (BRERA)

R. BELLET'S Syrup, } of each,
Distilled Water, } half an ounce.

Mix.—To be taken in the morning, in rachitis and scrofula.

ÖINTMENT OF NITRATE OF MERCURY.

Balsamum mercuriale, Unguentum ad Scabiem seu citrinum seu Hydrargyri citrinum seu Hydrargyri compositum seu Hydrargyri nitrati seu Nitratiss Hydrargyri seu mercuriale citrinum seu Supernitratiss Hydrargyri. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPE.; OLDEN.; POL.; BORUS.; SUEC.; BRUGNATELLI; CADET DE GASSICOURT; COXE; PIDERIT; SAUNDERS; SPIELMANN; SWEDIAUR; PH. SYPH.; VAN MONS)

Unguentum Hydrargyri Nitratus.
(LOND.)

- R. Purified Mercury, by weight, one ounce.
Nitric Acid, eleven fluid drams.
Prepared Lard, six ounces.
Olive Oil, four fluid ounces.

Dissolve the mercury in the acid, and mix the solution while hot, with the lard and oil melted together.

Unguentum Nitratus Hydrargyri fortius, vulgo U. Citrinum. (EDINB.)

- R. Purified Mercury, one part.
Nitrous Acid, two parts.
Olive Oil, nine parts.
Hog's Lard, three parts.

Dissolve the mercury in the acid, then beat the solution strongly with the lard and oil, previously melted together and nearly cold, in a glass mortar, to form an ointment.

EDINB. has also an *U. Nitratis Hydrarg. mitius*, which is prepared in the same manner, but with a triple proportion of oil and lard.

Unguentum Nitratis Hydrargyri, vulgo U. Citrinum. (DUBL.)

- R. Purified Mercury, one ounce.
Nitric Acid, eleven drams and a half.
Olive Oil, one pint.
Lard, four ounces.

Dissolve the mercury in the acid, then mix the solution with the lard and oil melted together, and form an ointment in the manner directed for the preparation of that of nitrous acid.

PH. SYPH.; and SAUNDERS prescribe one ounce each, of mercury and of nitric acid, four ounces of olive oil and eight of lard.—SWEDIAUR—the same as DUBL.—AMER.; and COXE—the same as EDINB.

- R. Mercury, sixty-four parts.
Nitric Acid (32 degrees), ninety-six parts.

To the cold solution, add

Melted Lard, one thousand parts, gradually and with constant trituration. (GALL.)

HISP. prescribes one part of mercury, one of acid, and ten of lard.—AMST.; AUSTR.; DAN.; FENN.; FULD.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; BORUS.; SUEC.; BRUGNATELLI; PIDERIT—one of the metal, two of acid, and twelve of lard.—ANT.; and SPIELMANN—three of the metal, four of acid, and thirty-two of lard.—VAN MONS—one of the metal, a sufficient quantity of acid to dissolve it, and fifteen parts of lard.—GENEV. and FERR.—two ounces of the metal, a sufficient quantity of acid to dissolve it, and two pounds of lard.

- R. Solution of Proto-nitrate of Mercury, one part.
Melted Lard, four parts.

Heat the solution, add it gradually to the lard, and triturate till cold. (POL.)

BATAV.; and BELG. prescribe three parts of the liquor and sixteen of lard.

The composition of this ointment is not well known; but it is ascertained that the decomposition which the salt and fatty substances undergo, varies according to the quantity of acid employed, its degree of concentration, and especially the age of the ointment: when fresh it is a compound of nitrous ointment, and of very subdivided subnitrate of mercury; but it appears that, in proportion as it becomes old, the nitrate totally disappears, and an acetate is produced.

It is stimulant, resolvent, detergent, siagagogue, anti-syphilitic, anti-psoric, and is very frequently employed in cutaneous diseases. In general it is reduced with an equal part of simple ointment.

DUBOIS' OINTMENT. (RATIER)

- R. Proto-nitrate of Mercury, two-drams.
Ointment of Cucumbers, half an ounce.
Mix.

It is applied to herpetic eruptions.

SATURNO-MERCURIAL OINTMENT. (AUGUSTIN; VAN MONS)

- R. Mercury, four parts.
Lead, one half part.

Dissolve each metal separately, in Nitric Acid (34 degrees), a sufficient quantity,

then mix with

Oil of Almonds, twenty-four parts.
Lard, forty-eight parts,
first the mercurial solution, and then the saturnine. Triturate well, and at the moment of cooling pour it into paper moulds.

MERCURIAL BOUGIES. (VAN MONS)

- R. Yellow Wax, six ounces.

Olive Oil, one ounce.
Melt together, and add,
Solution of Proto-nitrate of
Mercury, two drams.
Mix carefully, and make bougies.

OINTMENT AGAINST TINEA. (ELLIS)

R. Ointment of Nitrate of Mercury, half an ounce.
———— Tar, one ounce.
Mix.

DEUTO-NITRATE OF MERCURY. (BATAV.; BELG.; VAN MONS)

1. In the Solid State.

R. Red Oxide of Mercury, any quantity.
Nitric Acid, a sufficient quantity
to dissolve the oxide with the aid of a gentle
heat; then pour off the liquor, evaporate,
and crystallize. (VAN MONS)

2. In the Liquid State.

R. Red Oxide of Mercury, three parts.
Nitric Acid, diluted with an equal
quantity of water, six parts.
Dissolve with the aid of a gentle heat,
gradually adding the oxide to the acid, and
preserve the liquid. (BELG.)

LIPP. prescribes three parts of oxide and
four of acid, then the addition to the pro-
duct of twelve parts of pure water.—VAN
MONS—three parts of red oxide to be tritu-
rated with three of water, the mixture to be
slightly heated, and a sufficient quantity of
nitric acid to be added to effect solution,
and the liquor to be decanted at the end of
twenty-four hours.

The solution of the deuto-nitrate of the
latter pharmacopœias differs from the first
with respect to chemical composition, but
their medicinal effects are the same, or very
nearly so.

PROTO-NITRATE OF MERCURY AND OF AMMONIA. (PH. SYPH.; SWEDIAUR)

R. Diluted Nitric Acid, one pound.
Solution of Subcarbonate of
Ammonia, seven ounces.
Mix gradually, till effervescence ceases,
then add,

Mercury, eight ounces.

Dissolve in a sand-bath, and evaporate to
the point of crystallization.

WARD'S WHITE DROPS.

Guttulæ albæ Wardii, Solutio Nitratis Hy-
drargyri ammoniacalis. (PH. SYPH.;
NIEMANN; SWEDIAUR)

R. Proto-nitrate of Mercury and of
Ammonia, one ounce.
VOL. II.

Rose Water, three ounces.
Dissolve.

This preparation is [was] celebrated in
England.—Dose, two or three drops, in de-
coction of barley, sarsaparilla, or gruel.

PROTO-IODATE OF MERCURY. (NIEMANN; VAN MONS)

R. Proto-nitrate of Mercury, any quantity.
Iodate of Potass, a sufficient quantity.
Add the iodate to the nitrate by portions,
but quickly, and dry the precipitate.

HYDROCHLORATE OF MERCURY AND OF AMMONIA.

I. In the Solid State, but soluble in water and crystallizable.

Sal Alembroth, Sal Sapientiae, Sal Vitæ,
Sal Scientiæ. (ANT.; HASS.; GALL.;
VAN MONS)

R. Deuto-chloruret of } of each,
Mercury, } an
Hydrochlorate of } equal part.
Ammonia, }

Mix, and sublime. (GALL.)

SOUBEIRAN remarks that this process
gives a very variable mixture of deuto-
chloruret of mercury and of hydrochlorate
of ammonia. He recommends an equal
part of each of the two salts to be dissolved
in water, the liquor to be evaporated and
crystallized, bearing in mind that combina-
tion does not immediately ensue, and that
the crystals which form first, are a mixture
of the proper hydrochlorate of the two salts
and of a greater or less quantity of hydro-
chlorate of ammonia. According to him,
the preparation is a compound of one atom
of hydrochlorate of mercury, and of four
atoms of hydrochlorate of ammonia.

R. Precipitated Hydrochlorate of Mer-
cury and of Ammonia, any quantity.
Hydrochloric Acid, a sufficient quantity

to effect the solution with the aid of a mo-
derate heat; evaporate slowly, put it into a
cool place to crystallize, and wash the crys-
tals with cold water. (AMER.; BELG.;
VAN MONS)

This last preparation is denominated
Murias Hydrargyro-ammoniacale acerrimum,
and *Hydrargyrus muriatus ammoniacalis*
fortior.

II. In the Solid State, but insoluble in water.

Calx Hydrargyri alba, Hydrargyrum præ-
cipitatum album seu ammoniato-muriati-
cum seu muriaticum præcipitatum, Mer-
curius cosmeticus seu præcipitatus albus,
Murias Oxydi Hydrargyri ammoniacalis

seu *Ammoniacæ super-hydrargyricus*, *Submurias Hydrargyri ammonialis* seu *præcipitatus*.—White Precipitate. (LOND.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; DAN.; GALL.; FERR.; FULD.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; SUEC.; HERBIP.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS)

Hydrargyrum præcipitatum album.
(LOND.)

- R. Oxy muriate (Deuto-chloruret)
of Mercury, half a pound.
Muriate of Ammonia, four ounces.
Solution of Subcarbonate of
Potass, half a pint.
Distilled Water, four pints.

First dissolve the muriate, and then the oxy muriate, in the water, and add to the mixed solutions, that of the subcarbonate of potass. Wash the precipitated powder till it is tasteless, and then dry it.

Hydrargyri Submurias ammoniatum.
(DUBL.)

Add to the liquor poured off in the preparation of precipitated calomel, a sufficient quantity of water of caustic ammonia to precipitate the whole of the salt which it holds in solution. Wash the precipitate with cold distilled water, and dry it in bibulous paper.

AMER.; and COXE give the same formula as LOND.—VAN MONS—one part of deuto-chloruret and one and a half of hydrochlorate of ammonia, and a sufficient quantity of water to dissolve them, and also enough of solution of subcarbonate of potass to effect entire precipitation.—ANT.; BATAV.; FERR.; LUSIT.; SAX.; and SUEC. prescribe an equal part each, of deuto-chloruret and of hydrochlorate.—BATAV.—one of hydrochlorate and two of deuto-chloruret.

The other pharmacopœias which give this process prescribe eight ounces each, of the deuto-chloruret and of the hydrochlorate, to be dissolved in eight pounds of water, and precipitated with solution of subcarbonate of soda; these are (BELG.; HANN.; OLDEN.; POL.; BORUS.; SWEDIAUR)

- R. Deuto-chloruret of Mercury,
any quantity.

Hot Water, a sufficient quantity,
to dissolve; pour into the solution,

Solution of Ammonia,
a sufficient quantity,

or till no more precipitate is formed:
wash this with distilled water, and dry it.
(SWEDIAUR; VAN MONS)

- R. Mercury, } of each,
Nitric Acid, } eight ounces.
Distilled Water, } eight pounds.
Hydrochlorate of Ammonia,

two ounces.
Subcarbonate of Potass, one ounce.

First dissolve the metal in the acid, and add half of the water; then add the filtered solution of latter salts in the remainder of the water, till no more precipitate is effected. Collect this on a filter, wash, and dry it. (HASS.)

FULD. directs six ounces of mercury to be dissolved in twelve of nitric acid, seventy-two ounces of distilled water to be added, and then a solution of twelve ounces of hydrochlorate of ammonia in twenty-four of distilled water, and finally a solution of subcarbonate of potass to be added, till no more precipitate is formed.

- R. Nitrate of Mercury, eight ounces.
Distilled Water, a sufficient quantity.
Dissolve, and filter; add
Hydrochlorate of Ammonia
dissolved in Water, four ounces.

Instil gradually into the liquor,

Solution of Subcarbonate of Soda,
a sufficient quantity,
or till no more precipitate is formed; wash
and dry this. (DAN.)

These latter preparations must be considered precisely the same in effect as that required by DUBL.: the process is the same given by AMST., and COXE.

SOUBEIRAN considers the insoluble hydrochlorate of mercury and of ammonia to be a mixture of one atom of deuto-chloruret of mercury and of three atoms of mercuriate of ammonia, that is to say of a compound in which the oxide of mercury assumes the property of an acid. GUIBOUT had previously called a mixture of deuto-chloruret of mercury and of ammoniuret of deutoxide of mercury.

III. In the Liquid State.

Aqua mercurialis composita, *Liquor Muriatis hydrargyro-ammonialis*, *Mixtura seu Solutio mercurialis*, *Murias hydrargyro-ammoniacalis salutus*. (AUST.; BATAV.; GALL.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; SUEC.; AUGUSTIN; SWEDIAUR; PH. SYPH.; VAN MONS)

Dissolve, and filter. (AUSTR.)

R. Deuto-chloruret of Mercury, three grains.
Hydrochlorate of Ammonia, one or two grains.
Water, four ounces.
Mix.—It is recommended in gonorrhœa.

WEIKARD'S SOLUTION. (CADET DE GASSICOURT; PLANCHE)

- R. Deuto-chloruret of Mercury, } of each,
Hydrochlorate of } four grains.
Ammonia, }
Distilled Water, two ounces.
SYDENHAM'S Laudanum, thirty-six drops.
Oil of Cinnamon, one drop.

Mix.—Recommended in syphilis.—Dose thirty or forty drops, morning and evening in milk or gum water.

ANTI-SYPHILITIC LOHOCH. (VAN MONS)

- R. Hydrochlorate of Mercury and of Ammonia, half a dram.
Opium, four grains.
Sugar, four scruples.
Honey, four ounces.

Mix.—Dose, two drams, twice or thrice a day.

MERCURIAL LIQUOR.

Aqua mercurialis; Mercurius liquidus.
(SPIELMANN)

- R. Subcarbonate of Lead, one ounce.
Semi-vitreous Oxide of Lead, half an ounce.
Vinegar, six ounces.
Boil till both are dissolved: add
Alum, six drams.
Deuto-chloruret of Mercury, } of each,
Hydrochlorate of } one dram.
Ammonia, }
Spring Water, twelve ounces.
Rose Water, four ounces.

Mix, and preserve with straining.—It is recommended in an infinity of cutaneous affections. It is a bad preparation and should be rejected.

LARREY'S ALTERNATIVE AND ANTI-SYPHILITIC SYRUP. (BRUGNATELLI; CADET DE GASSICOURT)

- R. Sarsaparilla, two pounds.
Water, ten pounds.

Macerate for twelve hours; then boil for two and strain: boil the residue a second and third time with a similar quantity, and strain each time. Mix the three liquors, and set aside. On the other hand:

- R. Petals of Red Roses, half an ounce.
Senna, two ounces.
Anise Seeds, two drams.
Boiling Water, one pound and a half.
Depurated Juice of Borage, half a pound.

Infuse in a covered vessel placed in a water-bath, for twelve hours, then strain

with expression and set aside. During this time, evaporate the decoction of sarsaparilla till reduced to six pounds, and add to it besides the infusion,

- White Sugar, } of each, four pounds.
Honey, }

Clarify the syrup with white of egg, strain it while boiling, and evaporate it till very thick. Then

- R. Of this Syrup, two pounds and a half.
Hydrochlorate of Ammonia, } of each,
Deuto-chloruret of Mer- } ten
cury, } grains.
Extract of Opium, }
HOFFMANN'S Liquor, a sufficient quantity.

Dissolve the opium in a small quantity of water, and add the solution to the syrup, triturate the two salts with a sufficient quantity to dissolve them; add gradually, and with trituration, the whole of the syrup. Preserve it in well-stopped bottles.

It is alterative, anti-syphilitic, and anti-herpetic.—Dose, one ounce, in the morning, fasting, in a cupful of decoction of sarsaparilla.

MURIATICO-HYDRARGYRO-CUPRIC AMMONIACAL LIQUOR. (VAN MONS)

- R. Hydrochlorate of Mercury } of each,
and of Ammonia, } one dram
Copper } and a half.
and of Ammonia, }
Water, six ounces.

Dissolve each separately, mix the solutions, and filter.

WHITE PHAGEDENIC LIQUOR. (VAN MONS)

- R. Soluble Hydrochlorate of Mercury and of Ammonia, two scruples.
Lime Water, twenty-eight ounces.
Mix.

COMPOUND WHITE PHAGEDENIC LIQUOR. (VAN MONS)

- R. White Phagedenic Liquor, twenty-four parts.
Honey of Roses, one part.
Mix.

ZELLER'S OINTMENT.

Unguentum ad Scabiem Zelleri seu mercuriale album seu Hydrargyri album seu mundificans Zelleri seu Muriatis Hydrargyri ammoniacalis seu rosatum compositum. (LOND.; DUBL.; AMER.; AMST.; BATAV.; BRUNS.; HISP.; FERR.; FULD.; HANN.; HASS.; LUSIT.; PALAT.; POL.; CAST. BORUS.; SAX.; WIRTEM.; BRUGNATELLI; CADET DE

GASSICOURT; COXE; HUFELAND; PIDERIT; SWEDIAUR; PH. SYPH.; VAN MONS)

Unguentum Hydrargyri præcipitati albi. (LOND.)

R. White Precipitate of Mercury
(Precipitated Hydrochlorate
of Mercury and of Ammonia), one dram.
Prepared Lard,
one ounce and a half.

Melt the lard slowly, and add the first.

Unguentum Submuriatis Hydrargyri ammoniati. (DUBL.)

R. Prepared Lard,
one ounce and a half.
Ammoniated Submuriate of
of Mercury (Precipitated
Hydrochlorate of Mercury
and of Ammonia), one dram.
Melt the lard, and mix.

PH. SYPH.; and SWEDIAUR prescribe one part of the salt and from eight to sixteen parts of lard.—VAN MONS—one to four.—BATAV.; and HUFELAND—one to eight.—AMER.; AMST.; and COXE—one to twelve.

R. Precipitated Hydrochlorate of Mercury and of Ammonia, one part.
White Cerate, eight parts.
Triturate. (POL.)

R. Precipitated Hydrochlorate of Mercury and of Ammonia, one part.
Rose Ointment, eight parts.
Triturate. (HISP.; FERR.; FULD.; HASS.; SAX.; CADET DE GASSICOURT; PIDERIT)

BRUGNATELLI prescribes one part of precipitated hydrochlorate and twelve parts of ointment.—PALAT.; WIRTEM.; and SPIELMANN—one and twelve.

R. Precipitated Hydrochlorate of Mercury and of Ammonia, three parts.
Rose Ointment, twenty-four parts.
Oil of Lavender, one part.
Triturate. (LUSIT.)

BRUNS.; HANN.; and CAST. BORUS. prescribe sixteen parts of the first, one hundred and twenty-eight of lard, and one of oil of lemon.

It is stimulant, and is reputed cosmetic; it is applied in almost every affection of the skin. That which contains one-eighth of the precipitated hydrochlorate is used in the

quantity of from half a dram to one dram at each application.

ANTI-HERPETIC OINTMENT. (ELLIS; RATIER)

R. Precipitated Hydrochlorate of Mercury and of Antimony,
one dram and a half.
Camphor, one scruple.
Lard, two ounces.

Mix. (RATIER)

It is recommended in pustular tetters, after the inflammation has subsided.

R. Precipitated Hydrochlorate of Mercury and of Ammonia,
fifteen grains.
Nitrate of Potass, half a dram.
Sublimed Sulphur, one dram.
Lard, two ounces.

Triturate. (ELLIS)

OINTMENT AGAINST TÆNIA. (PIERQUIN)

R. Precipitated Hydrochlorate of Mercury and of Ammonia, half a dram.
Juniper, one dram.
Rose Ointment, one ounce.
Mix with a little oil of bay.

JADELOT'S OINTMENT. (RATIER)

R. Precipitated Hydrochlorate of Mercury and of Ammonia, two drams.
Camphor, eight grains.
Prepared Tutty, fifteen grains.
Fresh Butter, two drams.
Cocoa-nut Oil, half a dram.

Triturate.—It is recommended in scrofulous ophthalmia, to be applied to the margin of the eye-lids, every two or three days, at bed time.

JANIN'S OINTMENT. (FERR.; BORIES; CADET DE GASSICOURT)

R. Precipitated Hydrochlorate of Mercury and of Ammonia, one dram.
Tutty, of each,
Armenian Bole, } two drams.
Lard, half an ounce.

Mix.—For the same cases as the preceding.

ANTI-PSORIC OINTMENT. (PALAT.; WIRTEM.)

R. Turpentine, washed in Rose Water,
two ounces and a half.
Yolks of Eggs, three.
Mix, and add
Fresh Butter, six ounces.
Oil of Bay, one ounce.
Liquid Storax, half an ounce.

Precipitated Hydrochlorate of Mercury and of Ammonia, two drams.
Subcarbonate of Lead, one ounce.

Mix.—One dram for each application.—
It is a bad preparation.

MERCURIAL AMMONIACAL PILLS.

Pilule specifica. (AMER.; FENN.; GENEV.; LUSIT.; SUEC.; WURCEB.; BORIES; BRUGNATELLI; COXE; NIEMANN; SAUNDERS; SAINTE-MARIE; SWEDIAUR; VAN MONS)

R. Deuto-chloruret of Mercury, ten grains.
Arrow-root, one scruple.
Solution of Muriate of Ammonia, a sufficient quantity.

Make a mass. (AMER.; COXE)

AMER. erroneously gives the title of *Pilule Hydrargyri Oxymuriatis* to these pills. The formula is bad, and should be rejected.

R. Deuto-chloruret of Mercury, } of each,
Hydrochlorate of Ammonia, } fifteen grains.
Hot Distilled Water, a sufficient quantity
to dissolve the salts. Add to the solution

Powdered Root of Marshmallows, three drams.
Despumated Honey, two drams.

Make a mass, and divide into one hundred and twenty pills. (LUSIT.; WURCEB.)

Each contains one eighth of a grain of the deuto-chloruret.

R. Deuto-chloruret of Mercury, } of each,
Hydrochlorate of Ammonia, } one scruple.

Dissolve them in hot water, and add

White Sugar, one scruple.
Crumb of Bread, two drams.

Make one hundred and sixty pills. (FENN.; SUEC.)

They contain the same proportion of the deuto-chloruret as the preceding.

R. Distilled Water, two drams.
Hydrochlorate of Ammonia, a sufficient quantity

to saturate the water. Add

Deuto-chloruret of Mercury, twenty grains.
Crumb of Bread, a sufficient quantity.

Make eighty pills. (SARD.)

Each contains a fourth of a grain of the deuto-chloruret.

R. Hydrochlorate of Ammonia, one dram and a half.
Distilled Water, half an ounce.
Dissolve in the solution
Deuto-chloruret of Mercury, half a dram.

Add enough of crumb of bread to absorb the liquor, and make two hundred and eighty-eight pills. (GENEV.)

NIEMANN; SAUNDERS; SAINTE-MARIE; and SWEDIAUR give another formula which only differs from this in the proportion of the deuto-chloruret: from which are obtained pills, of which each contains half a grain of deuto-chloruret; they are called *Swediaur's Anti-syphilitic Pills*.

R. Deuto-chloruret of Mercury, twelve grains.
Hydrochlorate of Ammonia, half an ounce.

Syrup of Sarsaparilla, } of each,
Powdered Sarsaparilla, } a sufficient quantity

to make two hundred and forty pills.—
(BORIES)

Each contains one-twentieth of a grain of deuto-chloruret.

R. Soluble Hydrochlorate of Mercury and of Ammonia, one dram.
Powdered Biscuit, one ounce.
Simple Syrup, a sufficient quantity.

Make one hundred and twenty pills.
(VAN MONS)

Each contains half a grain of deuto-chloruret.

VAN MONS prescribes one scruple of sugar, two scruples of soluble hydrochlorate, two drams of crumb of bread, and a sufficient quantity of syrup.

The dose and mode of administration are the same as for HOFFMANN'S *Major Pills*, which ought to be preferred.

NOLTE'S MERCURIAL PILLS. (VAN MONS)

R. Soluble Hydrochlorate of Mercury and of Ammonia, twelve grains.
Opium, nine grains.
Extract of Liquorice, a sufficient quantity

to form a pilular mass.

ESCHAROTIC POWDER. (SAINTE-MARIE)

R. Precipitated Hydrochlorate of Mercury and of Ammonia, six grains.
Sub-carbonate of Magnesia, one scruple.
Powdered Savine, one dram.

Mix.—It is not very active.

OXALATE OF MERCURY.

Oxalas Hydrargyri. (BRUGNATELLI; NIE-MANN)

R. Aqueous Solution of Nitrate of Mercury, made without heat, and diluted with water, any quantity.
Concentrated Solution of Oxalic Acid, a sufficient quantity,
or till no more precipitate is formed; wash this with distilled water, and dry it in the shade.

The mode of administration is the same as that for the proto-chloruret. It is not much employed, but it has been tried in syphilitic cases.

PROTO-ACETATE OF MERCURY.

Acetas Hydrargyri, Hydrargyrum acetatus, Hydrargyrum acetatum seu aceticum, Mercurius aceticus, Sal Mercurii acetosus seu mercuriatus acetosus, Spuma Mercurii, Terra foliata Mercurii. (EDINB.; DUBL.; ANT.; BATAV.; BAVAR.; BRUNS.; GALL.; FERR.; HANN.; LUSIT.; PALAT.; BORUS.; WIRTEM.; BRUGNATELLI; COXE; SPIELMANN; SWEDIAUR; VAN MONS)

Acetas Hydrargyri. (EDINB.)

R. Purified Mercury, three ounces.
Dilute Nitrous Acid,
four ounces and a half,
or a little more than is sufficient to dissolve the metal.

Acetate of Potass, three ounces.
Boiling Water, eight pounds.

Mix the mercury with the acid, and towards the end of the effervescence, digest, if necessary, till the metal is completely dissolved. Then dissolve the acetate in water, and while hot add the other solution, and mix with agitation. Set aside the mixture to crystallize, wash the crystals, placed in a funnel, with cold distilled water, and finally dry them with the aid of a very gentle heat.

In the preparation of acetate of mercury, it is necessary that every vessel and implement made use of, should be of glass.

Hydrargyri Acetas. (DUBL.)

R. Purified Mercury, nine parts.

Dilute Nitrous Acid,

eleven parts.

Acetate of Potass, nine parts.

Boiling Distilled Water,
a sufficient quantity.

Add the acid to the metal, and when the effervescence has ceased, digest that the metal may be dissolved; then, having mixed the acetate with the water, immediately mix the solutions together. Then strain the mixture through a folded cloth, and put it in a cold place that crystals may form: having washed these with cold distilled water, dry them in bibulous paper with a very gentle heat. Throughout this process, glass vessels only should be used.

FERR.; and VAN MONS give the same formula as DUBL.—GALL. prescribes one part of proto-nitrate of mercury, six parts of distilled water, and a sufficient quantity of solution of acetate of potass to effect a complete precipitation.—COXE gives the same as EDINB.—BORUS.—five ounces of solution of acetate of potass, four pounds of boiling water to dilute it, and four ounces of solution of nitrate of mercury.—BRUGNATELLI—half a pound of mercury, half a pound of diluted nitric acid, and eight ounces of acetate of potass dissolved in two pounds and a half of hot distilled water.

R. Mercury, one pound.
Nitric Acid, diluted with water,
two pounds.

Dissolve the metal in the acid in a water-bath; add gradually to the liquor

Solution of Sub-carbonate of Potass,
a sufficient quantity

to precipitate all the oxide; wash this with pure water, and dry it in a moderate heat: after which dissolve it in

Distilled Vinegar, a sufficient quantity.

Filter this solution, evaporate it to a pellicle, and crystallize. (BATAV.)

LUSIT.; and SPIELMANN direct the oxide to be precipitated with solution of potass.

GAROT recommends the hydrate of protoxide of mercury, obtained by decomposing by caustic soda the proto-nitrate freed from deuto-nitrate, to be treated with acetic acid of the density of four degrees. He also gives another process, as being most likely always to give an identical product, and the proportions of which would be constantly the same. It is this:

R. Crystallized Proto-nitrate of Mercury,
seven parts.

Distilled Water, sixty parts.
Nitric Acid, one part.

Dissolve the salt in the water and acid; then add another solution made with

Acetate of Soda, five parts.
Distilled Water, sixty parts.

Mix.

There are two other processes:

R. Proto-chloruret of Mercury, } of each,
Solution of Potass, } two ounces.

Triturate together in a stone mortar, and add

Distilled Water, four ounces.

Boil for a quarter of an hour, constantly stirring; wash the residuum with distilled water till this is no longer made turbid by nitrate of silver: then, having dried it with a gentle heat, boil it with

Distilled Vinegar, thirty-six ounces, for three quarters of an hour. Filter the boiling liquor, and let it cool that it may crystallize. (HANN.; VAN MONS)

R. Red Oxide of Mercury, 9.75 parts.
Mercury, 9. parts.

Triturate together, adding from time to time a little distilled vinegar, to the complete extinction of the metal; then saturate with hot distilled vinegar, filter rapidly, and crystallize. (VAN MONS)

This process gives a mixture of proto-nitrate and of deuto-nitrate.

SOLUTION OF ACETATE OF MERCURY. (NIEMANN)

R. Proto-acetate of Mercury, ten grains.
Rose Water, five ounces.

Dissolve.—It is recommended in lotions in some diseases of the skin.

MERCURIAL SYRUP. (VAN MONS)

R. Proto-acetate of Mercury, one part.
Alcohol, as little as possible
to dissolve the salt with heat. Add to the solution
Simple Syrup, forty-eight parts.
Mix.

HYDRARGYRO-ACETOSE LINIMENT. (NIEMANN; VAN MONS)

R. Proto-nitrate of Mercury, one part.
Lard, six parts.
Olive Oil, two parts.

Triturate the salt with a little lard, and when the mixture is uniform, add the remainder, and then the oil. (VAN MONS)

NIEMANN prescribes from two scruples

to one dram of the salt, and one ounce of olive oil.

It is in rather general use in herpes.

KEYSER'S TROCHISCS. (PH. SYPH.; CADET DE GASSICOURT; COXE; VAN MONS)

R. Mercury, one ounce.
Nitric Acid, a sufficient quantity
to dissolve the metal. Add

Acetate of Potass, dissolved, a sufficient quantity
to effect precipitation. Triturate the precipitate with manna or honey and liquorice powder, and make trochiscs. (PH. SYPH.)

R. Acetate of Mercury, one part and a half.
Starch, } of each,
Gum Arabic, } one half part.
White Sugar, seven parts.
Simple Syrup, a sufficient quantity.

Make a mass. (VAN MONS)

R. Acetate of Mercury, } of each,
Manna, } one scruple.
Gum Arabic, }
Rose Water, a sufficient quantity.

Mix, and divide into twenty. (ELLIS)

R. Acetate of Mercury, two ounces.
Sugar, six ounces.
Gum Arabic, one ounce.
Powdered Marsh-mallows, } of each,
Starch, } half an ounce.

Mucilage of Gum Arabic, a sufficient quantity
to make comfits of one grain each. (CADET DE GASSICOURT)

Dose, two, morning and evening, augmenting the number progressively till twenty-five, or even more, are taken at each dose.

DEUTO-ACETATE OF MERCURY. (FERR.; VAN MONS)

R. Red Oxide of Mercury, two ounces.
Acetic Acid, seven ounces,
or a sufficient quantity to effect the solution of the oxide by the aid of a moderate heat; then evaporate slowly to dryness.

HYDRARGYRO-ACETIC LIQUOR. (VAN MONS)

R. Red Oxide of Mercury, ten parts.
Concentrated Acetic Acid, five parts and a half.

Pour the acid on the oxide in a bottle; agitate, and when the whole has become white, heat it in a water-bath to complete liquefaction. When it is decanted and be-

come cold, dissolve the white crystalline mass in

Rectified Alcohol, sixty parts.
Filter, and preserve.

TARTRATE OF MERCURY.

Hydrargyrum tartarisatum, Tartarus Mercurialis, Pulvis Constantini seu mercurialis argenteus. (NIEMANN; SWEDIAUR)

R. Black Oxide of Mercury, any quantity.
Tartaric Acid, a sufficient quantity.

Boil together to the perfect solution of the oxide; then evaporate, and crystallize.—Dose, one or two grains, twice a day.

TARTRATE OF MERCURY AND OF POTASS.
(BATAV.; GALL.; FERR.; CADET DE GASSICOURT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Saturated and boiling Solution of Mercury in Nitric Acid, any quantity.

Pour into it drop by drop an aqueous solution, also saturated and boiling, of neutral tartrate of potass till no more precipitate is formed: decant the liquor, wash it twice in succession with a little cold water; then dissolve it in a sufficient quantity of boiling water, and leave the solution in a cold place that it may crystallize. (BATAV.; GALL.; SWEDIAUR; VAN MONS)

R. Oxide of Mercury, precipitated from the Nitrate, and well washed, two ounces.
Super-tartrate of Potass, four ounces.

Triturate together for four hours; then boil it with

Distilled Water, forty-eight ounces.

Evaporate the liquor to a pellicle, and leave it to crystallize. (SPIELMANN)

R. Carbonate of Mercury, one ounce.
Super-tartrate of Potass, six ounces.

Triturate, and pour upon the powder

Boiling Water, four pounds.

Evaporate, and crystallize. (FERR.)

VAN MONS directs 10.25 parts of oxide of mercury, and 19.5 of super-tartrate of potass to be triturated together, a sufficient quantity of boiling water to be poured upon the mixture to effect a perfect solution: this to be heated for a quarter of an hour to the point of boiling, and then strained and crystallized.

He also proposes:

1. To triturate together 12.7 parts of deuto-chloruret of mercury, and 23. of neu-

tral tartrate of potass; then to continue the trituration with a little water in order to take up the hydro-chlorate of potass; to dissolve the residuum in boiling water, and finally to evaporate, filter, and crystallize.

2. To decompose a saturated solution of deuto-chloruret of mercury with the tartrate of potass and of soda.

3. To triturate together 10.25 parts of oxide of mercury, and 25. of neutral tartrate of potass, to take up the potass set at liberty, by washing; to dissolve the residuum, and crystallize the solution.

R. Mercury, twelve ounces and a half.
Red Oxide of Mercury, one ounce five drams and a half.

Super-tartrate of Potass, one ounce two drams and a half.
Water, a sufficient quantity

to make a paste of the whole. Triturate for a long while in a mortar placed in hot ashes, and then add boiling water to effect solution. Finally, filter the liquor, and evaporate to a pellicle. (CADET DE GASSICOURT)

VAN MONS also indicates this process, but with other proportions: viz. five parts and a half of mercury, six of oxide, and twenty-seven of super-tartrate of potass; again,—nine and a half of mercury, ten and a fourth of oxide, and nineteen and a half of super-tartrate. He also proposes the following processes:

1. To triturate together 22.2 parts of proto-chloruret, and 23. of tartrate of potass; to instil water, apply heat, and then to add boiling water to effect the solution, and finally to filter and crystallize.

2. To triturate to extinction 12.7 parts of deuto-chloruret of mercury, and 9.5 of mercury; and then to add 23. parts of super-tartrate of potass.

3. To make a sulphate with 9.5 of red oxide of mercury, and 10. of concentrated sulphuric acid; to triturate this salt with 9.5 of mercury, then to moisten the mass with 10. of water, and to mix it with 23. of neutral tartrate of potass: then to dissolve the whole in hot water, and to crystallize.

This double salt forms the base of PRES-AVIN's *Vegeto-mercurial Liquor*; this is prepared by dissolving mercury in nitric acid, precipitating by an alkali, taking up the precipitate by distilled vinegar, precipitating again, adding super-tartrate of potass, and finally dissolving it in hot water. (VAN MONS)

This salt is scarcely different from the preceding, and may be prescribed in the same circumstances. It was once very celebrated.

SYRUP OF TARTRATE OF MERCURY.
(BORIES; PIERQUIN)

- R. Stems of Bitter-sweet, two pounds and a half.
Water, a sufficient quantity.
Boil, and add to the decoction
Sugar, ten pounds.
Boil *à la plume*, and add to every two pounds,
PRESSAVIN's Liquor, twelve ounces.
Mix.—Dose, two ounces or more, daily.

OLEO-MARGARATE OF MERCURY.

Sapo Hydrargyri. (ROSS.; PH. SYPH.; AUGUSTIN; BORIES; NIEMANN; SWEDIAUR; VAN MONS)

- R. White Soap, one part.
Water, four parts.

Dissolve, and instil into the liquor
Weak Solution of Proto-nitrate of Mercury, a sufficient quantity, or, till no more precipitate is effected; apply heat till the thick parts (*magma*) reunite; then decant the liquid, and wash the precipitate with warm water; then mix it with eight parts of water at 30 degrees R., and add repeatedly solution of potass, to perfect solution: finally evaporate till the mass no longer adheres to the fingers.

It is applied as a resolvent to buboes. It

has been also recommended against rheumatism and chronic exanthemata.

CHAUSSEIER'S MERCURIAL SOAP. (CADET DE GASSICOURT)

- R. Mercurial Ointment (at equal parts), three ounces and a half.
Solution of Caustic Soda, three ounces.

Add this gradually, triturating in a glass mortar, till the mixture has acquired considerable tenacity; then put it into moulds, and leave it to dry.

It is for the same purposes as the preceding.

OINTMENT OF GREEN PRECIPITATE.
(VAN MONS)

- R. Deuto-chloruret of Mercury, two parts.
Hydrochlorate of Copper, one part.

Dissolve each separately in water, mix the solutions, add to it a solution of white soap, collect the precipitate on a filter, and wash it well.

It is recommended to be applied to syphilitic ulcers.—It is an oleo-margarate of mercury and of copper. The discoverer has doubtless considered it as fortunate that he has been able thus to combine the general method received from VAN SWIETEN with that, scarcely known, from CYRILLO.

HYDRASTES CANADENSIS, LINN.—YELLOW ROOT.

ELLIS.

A plant of the United States. (*Polyandria Polygynia*, LINN.; *Ranunculaceæ*, JUSS.; *Fig. Mill. Dict. et Icon.* p. 290. t. 285.)

The root is employed; it consists of fleshy tubercles, of a deep yellow colour within. It is extremely bitter.

HYDROGENIUM.—HYDROGEN; INFLAMMABLE GAS.

BRUNS.; NIEMANN; SWEDIAUR.

- R. Zinc, one part.
Put it into a bottle furnished with two tubes, one of which is bent, and points under a bell full of water or mercury; then pour upon the zinc

Sulphuric Acid, diluted with four times its weight of water,
one half part.

Receive the gas which is disengaged.

It is recommended to be inspired in phthisis pulmonalis.

PERCARBONATED HYDROGEN GAS.

Gas Hydrogenium carbonatum seu hydrocarbonatum. (BRUGNATELLI; NIEMANN; SWEDIAUR)

- R. Alcohol, any quantity.

Heat it in a retort; let the vapour which is exhaled pass through an earthen or stone

ware tube surrounded with burning charcoal, and receive in a pneumato-chemical apparatus the gas which is disengaged.

This gas is recommended to be inspired in

hæmoptysis, in ulcerose phthisis, and chronic catarrh, mixing it with atmospheric air; but if it be respired carelessly or for a long time, it may occasion fatal apoplexy.

HYOSCYAMUS.—HENBANE.

Two species of this genus of plants are used in medicine :

1. *Hyoscyamus albus*, LINN.—White Henbane.

ANT.; GALL.; FERR.; FORM. HÔP. FR.; GUIBOURT; MURRAY and GMELIN.

A plant of the middle of Europe. (*Pentandria Monogynia*, LINN.; *Solanææ*, JUSS.; *Fig. Zorn. Ic. pl. t. 218.*)

The herb (*Herba Hyoscyami albi*) is employed. It consists of rather branchy, very hairy stems, furnished with oval, petiolated leaves, the superior ones being entire, and the inferior sinuous or angular. It has a stupefying odour, and a vapid taste.

2. *Hyoscyamus niger*, LINN.—Henbane.

SYNONIMES : — *Jusquiame noire*, FR.; *Schwarzes Bilsankraut*, GERM.; *Buzirulbunj Urmanikun*, SIKRAN, ARAB.; *Blyn*, BOHEM.; *Korassanie*, CYNG.; *Khorasansie-ajuan*, DUK.; and *HIND.*; *Fandensosser*, SODBONNE, BULMEURT, DAN.; *Beleno*, SPAN.; *Bilsenkruid*, DUT.; *Giusquiamo*, ITAL.; *Adas*, JAV.; *Adas pedas*, MAL.; *Bielun*, SZALEY, LULEK, POL.; *Yosciamo*, PORT.; *Belena*, ROSS.; *Bolmært*, SUEC.; *Korassanie-omum*, TAM.

LOND.; DUBL.; EDINB.; AMER.: AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COKE.; GUIBOURT; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

This plant is spread all over Europe. (*Fig. Flore Medic. IV. 211.*)

The root, herb, and seeds are employed.

The root (*Radix Hyoscyami seu Jusquiami*) is thick, wrinkled, rather branched, brown without, white within, and of a sweet and disagreeable taste.

The herb consists of a hairy, thick, branching, cylindrical stem, and of large, alternate, amplexicaul, cottony, oval, lan-

ceolate, sinuous leaves, deeply notched on the border. It has a disagreeable and virous odour, and a vapid and repulsive taste, becoming bitter after desiccation.

The seeds are small, nearly reniform, wrinkled, rather compressed, and of an ash-grey colour.

The seeds contain, according to BRANDE, a peculiar alkaloid, *Hyoscyaminum*, *Hyoscyama*, which PESCHIER says he has also found in the leaves in combination with an equally peculiar acid, and which LINDBERGSON considers as ammoniaco-magnesian phosphate.

It is narcotic and poisonous, appearing to act principally on the nervous system.

Dose, from two to ten grains of the powder, twice or thrice a day.—REISINGER asserts that one drop of a solution of one grain of hyoscyamine in half a scruple of distilled water effects an extreme dilatation of the pupil without irritating the eye.

I. Preparations which contain Henbane in substance.

HYPNOTIC APPLICATION. (CADET DE GASSICOURT; PIERQUIN)

R. Powdered Leaves of	} of each,
Henbane,	
Flowers of	
Red Poppies,	
Opium,	ten grains.

Make a paste with a sufficient quantity of vinegar, to be applied to the forehead on folded linen, in cephalalgia.

ANODYNE CATAPLASM. (FORM. HÔP. FR.; CADET DE GASSICOURT; PIERQUIN; RATIER; SPIELMANN)

R. Powdered Leaves of Hen-	} of each,
bane,	
lock,	
To-	
bacco,	
den Nightshade,	} an equal part.
Linsed Meal,	
Decoction of Poppy Heads,	
a sufficient quantity.	

Make a paste. (CADET DE GASSICOURT; PIERQUIN)

RATIER gives the same formula, but replaces the meal with crumb of bread, and the decoction with water.

R. Leaves of Mallow, four handfuls.
 ——— Poppy, one handful.
 ——— Henbane, half a handful.
 Milk, a sufficient quantity.

Boil to a proper consistence, and add

Linseed Oil, two ounces.
 ——— Meal, one ounce.

Mix. (SPIELMANN)

FORM. HÔP. FR. prescribes an equal part each, of barley meal, of linseed meal, of powdered poppy leaves, and of powdered henbane leaves.

ANTI-HÆMOPTYSIC ELECTUARY. (NIEMANN)

R. Powdered White Poppy } of each,
 Seeds, } half
 ——— Henbane Seeds, } a dram.
 Syrup of Red Pop- } of each,
 pies, } one ounce and
 Conserve of Roses, } a half.

Mix.—Dose, the volume of a walnut, twice a day.

II. Preparations which contain a part of the constituent principle of Henbane.

A.

Extraction by mechanical means.

OIL OF HENBANE SEEDS. (BATAV.; BRUNS.; HISP.; FULD.; WIRTEM.; HERBIP.; VAN MONS)

R. Seeds of Henbane, any quantity.

Bruise them in a wooden mortar, enclose them in a linen bag, and express without heat. (BATAV.; HISP.; FULD.; HERBIP.)

VAN MONS recommends the seeds to be bruised first alone, then with an equal quantity of sweet almonds, and afterwards to be expressed.—BRUNS.; and WIRTEM.—to tonify them slightly before expression.

It is pectoral and anodyne.

PAREGORIC OIL. (FULD.)

R. Oil of Henbane Seeds, half an ounce.
 ——— White Petroleum, two drams.
 ——— Juniper, } of each, ten drops.
 ——— Amber, }

Mix.—It is anodyne and resolvent, and is used externally.

HÆMORRHOIDAL OINTMENT. (FULD.)

R. Oil of Henbane Seeds, two drams.

Unguentum Nutritum, six drams.
 Camphor, one scruple.
 Powdered Saffron, fifteen grains.

Mix.—It is anodyne, discutient, and cooling.

B.

Extraction by Water.

AQUEOUS EXTRACT OF HENBANE. (LOND.; EDINB.; DUBL.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SWEDIAUR; VAN MONS)

Extractum Hyoscyami, LOND.; *Succus spissatus Hyoscyami nigri*, EDINB.; *Succus spissatus Hyoscyami*, DUBL.

R. Fresh Leaves of Henbane, one pound.

Bruise them in a stone mortar, sprinkling them with a little cold water; then express the juice, without separating the sediment, and evaporate it to a proper consistence.

AMER.; AUSTR.; BRUNS.; DAN.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; HERBIP.; BRUGNATELLI; COXE; and SWEDIAUR all give the same formula.

R. Fresh Leaves of Henbane, any quantity.

Bruise them in a mortar and express the juice; evaporate this, without skimming till the addition of one fourth of powdered dry herb of henbane is sufficient to give it the consistence of extract. (BATAV.; BELG.; FERR.; FENN.; SUEC.)

R. Fresh Leaves of Henbane, any quantity.

Bruise with a little water, and strain through a sieve; set aside, decant, coagulate by heat, and strain again; evaporate to the consistence of pilular mass, remove it from the fire, incorporate the fecula, set apart, and finally evaporate to the required consistence. (ANT.; GALL.; FORM. HÔP. FR.; SAX.; WIRTEM.; VAN MONS)

R. Fresh Herb of Henbane, one pound.

Bruise it in a stone mortar, and express the juice; boil the residue for one hour, with three pounds of water; express, in the juice with the decoction, and evaporate. (HASS.; PIDERIT)

BAVAR. directs the juice to be expressed, the residue to be digested for half an hour

in one half of boiling water; the two liquors to be depurated by repose, and evaporated together.

Dose, from one grain to fifteen grains or more.

FECULA OF HENBANE.

Fecula Hyoscyami. (VAN MONS)

R. Fresh Leaves of Henbane, any quantity.

Bruise them in a stone mortar with a wooden pestle, and express; bruise the residuum with a little water, and express again. Mix the two liquors, set aside, decant, and put it over the fire to coagulate the fecula: immediately remove it, strain, wash the fecula, and having dried it with a gentle heat, pulverize it.

TINCTURE OF FECULA OF HENBANE.
(VAN MONS)

R. Fresh Fecula of Henbane, one part.
Alcohol, four parts.
Filter after several days' maceration.

SYRUP OF HENBANE. (HENRI)

R. Leaves of White Henbane, any quantity.

Bruise, express and filter the juice, and having evaporated it to the consistence of syrup, when cold, dilute it with distilled water, and evaporate it to dryness. Then dissolve thirty-two grains of this extract in a little distilled water, and add,

Simple Syrup, two pounds.

Mix by agitation.—It is recommended in chronic bronchitis. It is given alone, from a tea-spoonful to one ounce daily.

NARCOTIC BOLUSES. (BRERA)

R. Extract of Hemlock, two grains.
Mass of Pills of Dog-rose, four grains.
Liquorice Powder, a sufficient quantity

to make a bolus.—To be taken in the evening, at bed time.

SEDATIVE PILLS. (BRERA; ELLIS)

R. Extract of Henbane, eight grains.
Gum Arabic, six drams.
Rob of Elder, a sufficient quantity.

Make six pills.—Dose, one every three hours. (BRERA)

R. Opium, four grains.
Extract of Henbane, } of each,
Hemlock, } fifteen grains.

Mix, and divide into six pills. (ELLIS)

MEGLIN'S PILLS. (CADET DE GASSICOURT; HENRI; RATIER)

R. Extract of Henbane, } of each,
Valerian, } an equal part.
Oxide of Zinc, }

Make pills of three grains each. (CADET DE GASSICOURT; HENRI)

RATIER adds extract of fumitory.

They are recommended against tic dolo-reux.—Dose, from one to eight, gradually augmented.

ANTI-HYSTERIC PILLS. (PIERQUIN)

R. Extract of Hemlock, one dram.
Colocynth, a sufficient quantity.

Make pills of six grains each.—Dose, one morning and evening, augmenting it every five or six days, till four or five are taken at each dose.

SEDATIVE PILLS. (PIERQUIN)

R. Extract of Henbane, } of each,
Opium, } one ounce.
Belladonna, }
Hemlock, }

Make pills of one grain each.—They are prescribed in tic dolo-reux; to be augmented by one daily, till six are taken at each dose, which is to be continued till vertigo and general debility are produced.

ANODYNE AND RESOLVENT LINIMENT.
(RATIER)

R. Extract of Henbane, half a dram.
White Soap, two drams.
Linseed Oil, six ounces.

Mix.—It is employed by DUPUYTREN in glandular engorgements.

ANTI-SPASMODIC ELIXIR. (HAMB.; AUGUSTIN)

R. Cold and Strained Infusion of
Valerian, six ounces.
Extract of Henbane, half a dram.

Dissolve, and add,
Stimulant Liquor, two drams.

Mix. (HAMB.)

R. Extract of Henbane, ten grains.
Antimonial Wine, two drams.

Mix.—Dose, from five to ten drops; in whooping cough. (AUGUSTIN)

COLLYRIUM OF HENBANE.

Collyrium Hyoscyami. (AUGUSTIN)

R. Extract of Henbane, ten grains.
Distilled Water, half an ounce.

Mix.—It has been recommended by HIMLEY to dilate the pupil in cataract.

- R. Extract of Henbane, one scruple.
Water, three ounces.

Mix.—It is useful in slight ophthalmia with spasmodic constriction of the eye-lids.

SEDATIVE POTION.

Haustus sedans narcoticus. (BRERA)

- R. Extract of Henbane, five grains.
Lettuce Water, three ounces.
Simple Syrup, half an ounce.

Mix.—To be taken gradually in the course of the day.

SEDATIVE LAVEMENT.

Enema Hyoscyami seu anti-spasticum.
(AUGUSTIN)

- R. Herb of Henbane, }
 Rosemary, } of each,
Tops of Milfoil, } an equal part.
Orange Leaves, }
Root of Valerian, }

Mix.—Two drams of this mixture are to be boiled in one ounce or one ounce and a half of water.

NARCOTIC INJECTION. (FORM. HÔP. FR.; ELLIS)

- R. Leaves of Henbane, fifteen parts.
Water, a sufficient quantity
to obtain one hundred parts of strained decoction. Boil slightly, and add, if required, one half part of tincture of opium. (FORM. HÔP. FR.)

ELLIS prescribes half a dram of henbane, and one pound of boiling water to obtain a liquor which he recommends to be used externally, in cancer and indolent ulcers.

NARCOTIC EMULSION.

Emulsio sedativa. (AUGUSTIN; BRERA; NIEMANN; VAN MONS)

- R. Extract of Henbane, two drams.
Blanched Sweet Almonds, half an ounce.
Cherry Water, eight ounces.

Mix with the emulsion

- Calced Magnesia, two drams.
White Sugar, six drams.

Mix. (VAN MONS)

NIEMANN directs six ounces of emulsion to be made with twenty grains of henbane seeds, two drams of sweet almonds, and with pure water.

- R. Extract of Henbane, half a scruple.
Emulsion of Poppy Seeds, one pound.

Mix. (BRERA)

- R. Oil of Almonds, one ounce.
Water, six ounces.
Mucilage of Gum Arabic, a sufficient quantity

to make an emulsion. Add

- Amygdaline Syrup, one ounce.
Extract of Henbane, ten grains.

Mix. (AUGUSTIN)

Dose, one spoonful every two hours.

ANTI-ODONTALGIC GARGLE. (PIERQUIN)

- R. Leaves of Henbane, } of each,
 Plantain, } half a handful.
Flowers of Violet, } of each,
Petals of Red Roses, } one handful.
Poppy Heads, one ounce.
Leaves of Sage, six drams.
Water, a sufficient quantity.

Boil; and add, when strained,

- Vinegar, four ounces.

Mix.

NARCOTIC CATAPLASM. (GALL.)

- R. Leaves of Henbane, two ounces.
Poppy Heads, one ounce.
Water, a sufficient quantity.

Boil for an hour and a half; then add to the strained liquor

- Emollient Meals, four ounces.

Boil, constantly stirring, to the consistence of a cataplasm.

C.

Extraction by Alcohol.

ALCOHOLIC EXTRACT OF HENBANE. (RATIER)

- R. Dried Herb of Henbane, one part.
Alcohol (12 degrees), four parts.

Macerate for four days at a temperature of 20 degrees R., then filter, distil three-fourths from the liquor, and evaporate the remainder to the consistence of an extract.

ALCOHOLIC TINCTURE OF HENBANE.

Tinctura seu Essentia Hyoscyami seu Hyoscyami nigri. (LOND.; DUBL.; EDINB.; AMER.; HANN.; POL.; SAX.; SUEC.; ELLIS; HUFELAND; NIEMANN)

Tinctura Hyoscyami.

- R. Dried Leaves of Henbane, four ounces.

Proof Spirit, two pints.

Macerate for fourteen days, and filter. (LOND.)

R. Dried Leaves of Henbane, in
coarse powder, five ounces.
Proof Spirit, two pints.
Digest for seven days, and filter.
(DUBL.)

Tinctura Hyoscyami nigri. (EDINB.)

R. Dried Leaves of Henbane,
one ounce.
Proof Spirit, eight ounces.
Digest for seven days, and filter.

AMER.; and COXE give the same formula as DUBL.; and SUEC.; and NIEMANN the same as EDINB.

R. Leaves of Henbane, one ounce.
Rectified Spirit, four ounces.
Distilled Water, two ounces.

Digest with a gentle heat, express, and filter. (HAMB.; POL.)

R. Juice of Henbane, recently expressed, } of each,
Concentrated Alcohol, } an
equal part.

Digest without heat for several days, and filter. (SAX.)

R. Root of Henbane, two ounces.
Alcohol, one pound.

Macerate for three days, and filter. (HUFELAND)

It is narcotic.—Dose, from ten to thirty drops.

TINCTURE OF HENBANE SEEDS. (VAN MONS)

R. Seeds of Henbane, one part.
Proof Spirit, six parts.

Digest in a warm water-bath, strain with expression, and filter.

D.

Extraction by Æther.

ÆTHEREAL TINCTURE OF HENBANE.

Tinctura Hyoscyami ætherea. (HANN.)

R. Leaves of Henbane, two ounces.
Sulphuric Æther, eight ounces.

Digest for three days in a covered vessel, frequently shaking, and decant.—Dose, ten drops, several times a day.

E.

Extraction by Fat Substances.

OIL OF HENBANE.

Oleum Foliorum seu Herbæ Hyoscyami infusum seu coctum. (AMER.; BATAV.; BAVAR.; BELG.; GALL.; HAMB.; LIPP.;

BORUS.; SAX.; SUEC.; HERBIP.; SWEDIAUR.; VAN MONS)

R. Leaves of Henbane, one part.
Olive Oil, eight parts.

Digest for several hours, and express. (BORUS.; SAX.)

ANT.; BATAV.; BELG.; and LIPP. prescribe one part of herb and four parts of oil.—SUEC.—one of herb and three of oil.

R. Bruised Seeds of Henbane,
one part and a half.

Olive Oil, three parts.

Infuse, and express strongly. (VAN MONS)

R. Leaves of Henbane, one part.
Olive Oil, eight parts.

Boil slowly to the consumption of the humidity, and express. (HANN.)

HAMB. prescribes one of leaves and four of oil.—BAVAR.; DAN.; and SWEDIAUR—one of leaves and three of oil.

R. Leaves of Henbane, one part.
Olive Oil, two parts.

Digest for three days in a warm place, then boil to the consumption of the humidity, and express. (HERBIP.)

R. Bruised Leaves of Henbane, one part.
Olive Oil, two parts.

Digest on hot ashes for twenty-four hours, and express; repeat the operation a second time, then finally boil again, and express. (GALL.)

It is employed externally in frictions.

BUSCH has recommended, in hæmoptysis, three or four tea spoonfuls, thrice a day, of a mixture of two parts of oil of almonds and one part of an oil prepared by boiling two ounces of fresh herb of henbane in eight of olive oil.

GREEN OIL OF HENBANE.

Oleum Hyoscyami viride. (VAN MONS)

R. Oil of Henbane by infusion,
three parts.

Washed Fecula of Henbane,
one part and a half.

Dissolve. (VAN MONS)

It is rubbed on the epigastrium in colic, and on the anus in painful piles.

OINTMENT OF HENBANE.

Unguentum Hyoscyami. (SARD.; WIRTEM.; VAN MONS)

R. Fresh Leaves of Henbane, one pound.
Lard, ten ounces.

Boil to the consumption of the humidity, express, and add

Oil of Henbane Seeds, two ounces.
 Mix. (SARD.; WIRTEM.)
 R. Oil of Henbane Seeds, four parts.
 Green Fecula of Henbane, one part.
 Boil to the consumption of the humidity and add
 White Wax, one part.
 Melt slowly. (VAN MONS)
 It is to be rubbed on the abdomen and near the anus, in colic and piles.

HENBANE PLASTER.

Emplastrum Hyoscyami seu de Hyoscyamo.
 (BATAV.; BRUNS.; DAN.; HAMB.; OLDEN.; LUSIT.; POL.; BORUS.; SARD.; SUEC.; WIRTEM.; PIERQUIN.; SPIELMANN.; SWEDIAUR.; VAN MONS)

R. Oil of Henbane Seeds, half a pound.
 Juice of Henbane, one pound.

Boil to the consumption of the humidity; add

Yellow Wax, fourteen ounces.
 Venice Turpentine, two ounces.

After fusion, and when nearly cold

Powdered Leaves of Henbane
 two ounces.

Mix. (SPIELMANN.; SWEDIAUR.)

BRUNS.; SARD.; WIRTEM.; and PIERQUIN prescribe one pound of oil of henbane seeds, one pound of juice of henbane, fifteen ounces of wax, one ounce and a half of turpentine, and two ounces of powdered henbane.—VAN MONS—twelve parts of henbane seeds, six of green fecula of henbane, fifteen of yellow wax, one of turpentine, and one of resin.

R. Green Fecula of Henbane, twelve parts.
 Resin, four parts.
 Yellow Wax, two parts.
 Turpentine, one part.

Boil slowly to the consumption of the humidity. (VAN MONS)

R. Yellow Wax, } of each, one part.
 Resin, }
 Olive Oil, }

To the melted and half cold mixture, add

Powdered Leaves of Henbane,
 two parts.

Mix. (SWEDIAUR.)

DAN.; HANN.; POL.; BORUS.; and SUEC. prescribe one part of wax, one half part of resin, and one half part of olive oil.—POL.—also allows lard to be used, and adds one part of powdered henbane.—SAX.—one each of wax, of resin, and of oil, and two of the powder.

R. Simple Plaster, } of each.
 Yellow Wax, } one pound.

Olive Oil, six ounces.
 Add to the melted and half cold mass,
 Powdered Herb of Henbane,
 one pound

Mix. (BATAV.)

R. Yellow Wax, fourteen ounces.
 Resin, } of each,
 Olive Oil, } half a pound.

Add to the melted and half cold mixture
 Powdered Herb of Henbane,
 one pound.
 Extract of Henbane,
 one ounce.

Mix. (OLDEN.)

R. Strong Vinegar, sixteen ounces.
 Gum Ammoniac, eight ounces.

Strain the solution through a close linen cloth, and add

Inspissated Juice of Henbane,
 one pound.

Evaporate over a slow fire to the required consistence, stirring continually. (LUSIT.)

This plaster is applied as resolvent to schirrous and indolent tumours.

OINTMENT FOR BURNS.

Unguentum ad Ambusta. (SPIELMANN)

R. Leaves of Henbane, }
 ——— Stramonium, } of each,
 ——— Bitter-sweet, } one handful.
 ——— Elder, }
 Lard, thirty-two ounces.

Boil to the consumption of the humidity, and strain.

ANODYNE OINTMENT.

Unguentum anodynum. (WIRTEM.; SPIELMANN)

R. Fresh Leaves of Henbane, }
 ——— Chamomile, } of each,
 ——— Mallow, } one
 ——— Melilot, } ounce.
 ——— Lesser Mul-
 lein, }
 ——— Dill, }

— Root of Marshmallows, two ounces.

Fenugreek, } of each,
 Linseed Meal, } three drams.
 Olive Oil, } two pounds.

Boil over a slow fire to the consumption of the humidity, and melt in the strained mass,

Yellow Wax, four ounces.

Mix.—For frictions to painful parts.

BALSAMUM TRANQUILLUM.

Balsamum Tranquillum, Oleum Narcoticorum seu Bufonum seu Stramonii compositum. (HISP.; GALL.; SARD.; CADET DE GASSICOURT.; PIERQUIN.; SPIELMANN.; VAN MONS)

R. Fresh Leaves of Henbane }
 ————— Hounds- } of each,
 ————— tongue, } one pound.
 ————— Tobacco.

Wine, four pounds and a half.

Boil down to three pounds, and express strongly, add to the liquor

Olive Oil, three pounds.

Boil down to one half over a slow fire, and pour into bottles when cold. (CADET DE GASSICOURT; PIERQUIN)

R. Olive Oil, six pounds.
 ————— Leaves of Stramonium,
 ————— Garden Night-
 ————— shade, } of each,
 ————— Belladonna, } four
 ————— Tobacco, } ounces.
 ————— Henbane,
 ————— White Poppy,

Boil slowly to the consumption of the humidity, express, and pour the liquid upon

Dried Tops of Rosemary,
 ————— Sage,
 ————— Rue,
 ————— Wormwood,
 ————— Roman Worm-
 ————— wood, } of
 ————— Hyssop, } each,
 ————— Lavender, } one
 ————— Thyme, } ounce.
 ————— Marjoram,
 ————— Water-mint,
 ————— Elder,
 ————— St. John's-

wort,

Macerate in the same for several months, in a covered vessel, and strain. (GALL.; CADET DE GASSICOURT)

R. Fresh Leaves of Stramonium, }
 ————— Garden }
 ————— Nightshade, }
 ————— Nightshade (Phytolacca } of each,
 ————— decandra), } four
 ————— Belladonna, } ounces.
 ————— Mandrake,
 ————— Henbane,
 ————— Poppy, eight ounces.
 ————— Persicaria, one ounce.

Living Toads, five.

Olive Oil, sixty ounces.

Boil to the consumption of the humidity, and strain the liquid upon

Fresh Tops of Rosemary, }
 ————— Sage, } of
 ————— Rue, } each,
 ————— Hyssop, } one
 ————— Lavender, } ounce.
 ————— Thyme,
 ————— Marjoram,
 ————— Astmary (Bal-
 ————— samita odorata, DESF.)
 ————— St. John's-wort,

Fresh Tops of Wormwood, two ounces.

Fresh Leaves of Elder, one ounce.

Decant after fifteen days digestion. (HISP.; SPIELMANN)

R. Fresh Leaves of Wormwood, }
 ————— Lavender, } of
 ————— Marjoram, } each,
 ————— Water-mint, } one
 ————— Rue, } part.
 ————— Rosemary,
 ————— Sage,
 ————— Tansey,
 ————— Thyme,
 ————— Belladonna,
 ————— Henbane,
 ————— Mandrake,
 ————— Garden } of
 ————— Nightshade, } each,
 ————— Tobacco, } four
 ————— White Poppy, } parts.
 ————— Stramonium,
 ————— Elder,

Bruise, mix with the mass a little water, express the juice, set it aside, decant, coagulate at the fire, set apart the green fecula, wash it with cold water, and heat it slightly with

Beef Marrow, ninety-eight parts.

Dissipate the humidity completely, set aside, and decant. (VAN MONS)

R. Living Toads, two.
 ————— Leaves of Garden Nightshade, } of each,
 ————— White Henbane, } two
 ————— Mandrake, } hand-
 ————— Tobacco, } fuls.
 ————— Dwarf Elder,
 ————— Willow,

Olive Oil, six pounds.

Boil slowly to the consumption of the humidity, and infuse in the expressed oil,

Living Toads, two.
 ————— Tops of Wormwood, }
 ————— Mountain Calamint, }
 ————— Hyssop, }
 ————— Marjoram, } of each,
 ————— Rue, } one
 ————— Thyme, } handful
 ————— Flowers of Rosemary,
 ————— Sage,
 ————— Lavender,
 ————— St. John's-wort,
 ————— Elder,

After twenty-four hours' digestion in a water-bath, strain with expression. (SAUNDERS)

It is used externally in gout and rheumatism.

ACOUSTIC BALSAM. (CADET DE GASSICOURT)

R. Balsamum Tranquillum, two drams.
 Oil of Rue, by infusion, half an ounce.

Sulphuretted Balsam of Turpentine,	} of each, two drams.
Tincture of Assafoetida,	
— Ambergis,	
— Castor,	
Rectified Oil of Amber,	

Mix.—It is applied in accidental deafness where there is no inflammation nor acute pain.

SEDATIVE LINIMENT. (SAINTE-MARIE)

R. Balsamum Tranquillum,	} of each, three ounces.
Oil of Henbane,	
— Chamomile,	
— Camphor,	

Mix.—To be used as a liniment to pained parts.

UNGUENTUM POPELUM; OINTMENT OF POPLAR.

Unguentum Gemmarum Populi compositum.
(AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; FULD.; HANN.; PALAT.; SARD.; WIRTEM.; HERBIP.; SPIELMANN; VAN MONS.)

R. Buds of Poplar,	two parts.
Fresh Herb of Henbane,	one part.
Lard,	four parts.

Boil slowly to the consumption of the humidity, and express. (HANN.)

R. Fresh Buds of Poplar,	one pound and a half.
Lard,	three pounds.

Preserve till the season at which can be added to it,

Fresh Leaves of Henbane,	} of each, four ounces.
— Bella-donna,	

After sufficient maceration, boil to the consumption of the humidity, and express. (AMST.)

This formula presents very numerous but unimportant varieties, both with regard to the proportions and to the number and choice of the plants associated with the henbane: for instance—BATAV. prescribes two pounds of buds, four of lard, six of henbane, and six of belladonna.—HISP.—three pounds of buds, eight of lard, one and a half of henbane, one each, of houseleek, of violet, and of navel-wort, and two of garden nightshade.—SARD.—two pounds of buds, four of lard, three handfuls each of white henbane, of red poppies, of houseleek, and of small houseleek, and six ounces of garden nightshade.—BRUNS.—one pound and a half of buds, six pounds of lard, four ounces each, of white poppy, of henbane, of garden nightshade, of lettuce, of violet, and of houseleek.—FORM. HÔP. FR.; and GALL.—four parts of buds, twelve of lard,

one each of poppies, of belladonna, of henbane, and one of garden nightshade.—FULD.—one pound of buds, two pounds and a half of lard, four ounces each of henbane and of garden nightshade, and six ounces of leeks.—HERBIP.—two pounds of buds, five of lard, four ounces each of garden nightshade, of henbane, of poppies, of elder, and of violet, and three ounces of houseleek.—POL.—one pound and a half of buds, six pounds of lard, six ounces of henbane, three each, of poppies, of garden nightshade, of houseleek, of wall pepper, of lettuce, of burdock, of violet, of naval-wort, and of tops of mountain bramble.—VAN MONS.—two parts of buds, four of lard, and one part each, of the dried feculæ of burdock, houseleek, small houseleek, henbane, lettuce, mandrake, garden nightshade, mountain bramble, and rose-wort.

R. Dried Buds of Poplar,	one pound.
Lard,	four pounds.
Fresh Leaves of Henbane,	} of each, six ounces.
Belladonna,	

Boil to the consumption of the humidity, and strain with expression. (BELG.)

ANT. directs three ounces each of the tops of mountain bramble, of rose-wort, of small houseleek, of houseleek, of henbane, of belladonna, and of poppy, and six ounces of garden nightshade, to be bruised together, and put into a pan with four pounds of lard, and boiled slowly till the lard has acquired the colour of the plants; then eight ounces of dried poplar buds to be added, and left to infuse for some time; then to be expressed, and when cold melted again, and the heat to be continued to be applied till the humidity is entirely consumed, when it is to be finally strained through a linen cloth.

It will be seen that all the processes given for the preparation of this ointment may be reduced to two; one in employing fresh poplar buds and waiting the proper season to macerate in the lard charged with their resinous principle, the other plants prescribed, as they appear; the other, in macerating the dried buds in the lard at the same time with the fresh leaves. Both these processes have been lately modified. HENRI recommends the buds to be kept in melted lard till all their humidity is dissipated, the mixture to be poured into a pot and covered with a stratum of lard, and afterwards to be made into an ointment in the usual manner. BOULLAY requires an equal part each, of buds and of lard to be heated together to the perfect consumption of the humidity, the mixture to be strained and set aside; an ointment to be then prepared with lard and narcotic herbs: and finally the two ointments to be melted together. BRIAND has proposed to reduce the narcotic herbs to pulp, to put them into a pan and evaporate the greater part of their

moisture, then to add lard, and when it is well coloured, to add the dried buds; then, having kept the mixture in a fluid state for about half an hour, to strain it with expression and to keep it again in infusion for five or six hours that the heavier parts may subside, then to let it cool, to scrape it, and finally to liquify it again and pour it into a pot. GERMAIN recommends one hundred pounds of lard, twelve pounds each, of the feculæ of henbane, black poppy, garden nightshade, and belladonna, to be put into a pan which they may fill to three fourths; to stir incessantly to the consumption of the humidity; then to add seventeen pounds of dried buds of poplar, and to apply a very moderate heat for half an hour, constantly stirring; finally to strain it through linen, and to separate the deposit by the ordinary means.

BOULLAY'S process is the best of all. But we do not see the necessity for all this anxiety about the mode of preparing this ointment. It is trouble and care absolutely thrown away; for the ointment itself, already becoming disused, will soon be for-

gotten, as it should have been long ago. The plaster of henbane reduced with oil would replace it perfectly well, where a compound of resinous and narcotic bodies may be thought useful; or there might be substituted in its place a simple resinous ointment mixed with oil of henbane or of garden nightshade.

ANTI-HÆMORRHOIDAL LINIMENT.

Lotus seu Litus ad Hæmorrhoides. (BRERA; BORIES; CADET DE GASSICOURT; PIERQUIN; SPIELMANN)

R. Unguentum Populeum, one ounce.
Balsamum Tranquillum, half an ounce.
Oil of Eggs, two drams.

Mix. (SPIELMANN)

R. Unguentum Populeum, }
Rose Ointment, } of each, an
Marshmallows Ointment, } equal part.
Honey,

Mix. (BRERA; BORIES; CADET DE GASSICOURT; PIERQUIN)

HYPERICUM.

Three species of this genus of plants are employed in medicine.

1. *Hypericum Androsæmum*, LINN.—Common Tutsane or All-heal.

SYNONIMES:—Toute-saine, Androsème, FR.; Conradskraut, GERM.; Mannsblød, DUT.

GALL.

A plant of the middle of Europe. (*Polyadelphia Polyandria*, LINN.; *Hypericeæ*, JUSS.; Fig. Flore Lond. t. 206.)

The herb (*Herba Sicilianæ seu Totæ-Sanæ*) is employed. It consists of a woody and slightly angular stem, and of oval, sessile, glabrous leaves. It has a resinous odour, and taste.

It is diuretic and anthelmintic, according to DALE.

2. *Hypericum quadrangulare*, LINN.

GALL.; HISP.; HASS.; BERGIUS; SPIELMANN.

This plant is spread over all Europe. (Fig. CEd. Fl. Dan. t. 640.)

The tops (*Summitates Hyperici*) are employed. It consists of a square stem, and of oval, glabrous, simple leaves, presenting numerous semi-transparent glands, with a

row of black hairs; at its extremity is a close corymb of yellow flowers.

3. *Hypericum perforatum*, LINN.—St. John's-wort.

SYNONIMES:—Millepertuis perforé, FR.; Johanniskraut, GERM.; St. Jana bylina, Zwonacek cerweny, BOHEM.; Jordhumle, Perikum, DAN.; Hypericono, ITAL.; and SPAN.; Mannsblød, DUT.

AMST.; ANT.; BATAV.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; BORUS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A common plant of Europe. (Fig. Zorn. Ic. pl. t. 31.)

The flowering tops and seeds are employed.

The tops (*Summitates Hyperici perforati*) consist of a slightly triangular stem, and of oval, oblong leaves dotted with semi-transparent points, and of yellow flowers disposed in corymbs. They have a feeble but agreeable odour, and a balsamic, bitter, and slightly styptic taste.

The seeds are small, of a deep brown

colour, without odour, but of a bitter and resinous taste.

OIL OF THE SEEDS OF ST. JOHN'S-WORT.

Oleum Seminum Hyperici. (SARD.)

R. Seeds of St. John's-wort, any quantity.

Bruise them, enclose the paste in a bag, express the oil without heat, and filter.

EXTRACT OF ST. JOHN'S-WORT. (WIRTEM.)

R. Flowers of St. John's-wort,

one pound.

Alcohol, five or six pounds.

After sufficient extraction, express, digest, and strain; then boil the residuum in three pounds of water, and clarify the decoction with white of egg. Mix the two liquors together, distil off the alcohol, and evaporate to the consistence of an extract.

SYRUP OF ST. JOHN S-WORT.

Syrupus Hyperici. (WIRTEM.)

R. Fresh Flowers of St. John's-wort,

one pound and a half.

Boiling Water, one pound.

Macerate for twenty-four hours in a cold place; then express, and add to every nine ounces of the liquor,

White Sugar, sixteen ounces.

Mix.

COMPOUND VINOUS TINCTURE OF ST. JOHN'S-WORT.

Balsamum catholicum ex Vino seu innocentianum. (SARD.)

R. Benzoin, three ounces.

Storax, two ounces.

Dry Balsam of Peru, one ounce.

Aloes, } of each,

Myrrh, } half an ounce.

Olibanum, }

Root of Angelica, }

Dried Flowers of St. John's-wort,

four handfuls.

Red Wine, four pounds.

Digest the flowers and root in the wine for three days, on warm ashes; then, when cold, express, and digest the other substances in the tincture for fifteen days, frequently shaking the vessel, and strain.

TINCTURE OF ST. JOHN'S-WORT.

Tinctura seu Essentia Hyperici. (BRUNS.; FULD.; LIPP.; WIRTEM.)

R. Flowers of St. John's-wort,

four ounces.

Alcohol, one pound and a half.

Digest for three days, express strongly, and filter. (WIRTEM.)

BRUNS. directs three ounces of flowers to be digested for three days in sixteen ounces of alcohol; then two more ounces of flowers to be infused in the strained tincture, and finally filtered.—FULD.; and LIPP.—one part of flowers to be digested in four parts of alcohol for three days.

It is excitant, and was formerly extolled in melancholy and in verminal disorders.—Dose, from fifty to eighty drops.—It was also applied externally as a vulnerary.

COMPOUND TINCTURE OF ST. JOHN'S-WORT.

Tinctura alcoholica Hyperici vulneraria, Balsamum catholicum cum Spiritu Vini. (HISP.; SARD.)

R. Dried Flowers of St. John's- } of each,
wort, } half an
Root of Sweet-flag, } ounce.
Alcohol, three pounds.

Digest with a moderate heat in a covered vessel for several days, shaking from time to time; then add

Benzoin, three ounces.

Storax, } of each, two ounces.

Balsam of Tolu, }

Myrrh, half an ounce.

Strain after sufficient digestion. (HISP.)

R. The dry residue of the preparation of the Compound Vinous Tincture of St. John's-wort.

Root of Angelica, half an ounce.

Dried Flowers of St. John's-wort,

one ounce.

Alcohol, four pounds.

Digest for eight days, and strain. (SARD.)

It is stimulant, but is more used externally than internally.—Dose, from six drops to half a scruple.

STAHL'S SPLENETIC ESSENCE.

Essentia splenetica Stahl. (WIRTEM.)

R. Flowers of St. John's-wort,

one ounce.

Herb of Vervain,

— Rue,

— Wall-rue,

— Sage,

— Plantain,

of each,
half an ounce.

Alcohol, one pound.

Infuse, express, and filter.

It is excitant, and is recommended in hypochondria, and also as resolvent and diuretic.—Dose, from sixty to eighty drops.

OIL OF ST. JOHN'S-WORT BY INFUSION.

Oleum Hyperici seu Hyperici per infusum. (AMST.; BATAV.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HANN.;

HASS.; LIPP.; OLDEN.; BORUS.; SAX.;
SARD.; WIRTEM.; HERBIP.; PIDERIT;
SPIELMANN; VAN MONS)

- R. Fresh Flowers of St. John's-wort,
three parts.
Olive Oil, four parts.

Digest for eight days in a warm place,
and express. (BATAV.; BELG.)

BRUNS.; FERR.; FULD.; and LIPP. pre-
scribe one part of flowers and two parts of
oil.—BORUS.; and SAX.—one of flowers
and eight of oil.

- R. Flowers of St. John's-wort, one part.
Olive Oil, four parts.

Expose them to the rays of the sun for
one week, then boil slightly, and strain;
repeat the operation two other times; the
second with dried flowers, and for a month.
(GALL.)

OLDEN.; and SARD. indicate the same
process, but OLDEN. prescribes one of herb
and three of oil, and SARD.—one of herb
and two of oil each time.

- R. Flowers of St. John's-wort, one part.
Olive Oil, two parts.

Macerate for eight days, boil to the con-
sumption of the humidity, and express.
(BRUNS.; HERBIP.)

AMST.; and DAN. prescribe one of flow-
ers and three of oil.—HISP.—two of flowers
and three of oil.—HANN.—one of flowers
and eight of oil.

- R. Flowers of St. John's-wort,
three pounds.
White Wine, half a pound.
Olive Oil, two pounds.

Digest for several days in a warm place;
then express, and add

Venice Turpentine, one pound.

Mix. (PIDERIT)

- R. Flowers of St. John's- } of each,
wort, } three pounds.
Olive Oil, }
White Wine, } six ounces.

Boil to the consumption of the humidity,
express, and add

Venice Turpentine. one pound.

Saffron, two drams.

Strain at the end of twenty-four hours.
(SPIELMANN)

- R. Flowers of St. John's-wort,
three pounds.
Saffron, four scruples.
Olive Oil, two pounds.
Wine, half a pound.

Macerate for three days, boil to the con-
sumption of the humidity, strain, and add

Turpentine, one pound.

Mix. (SARD.; WIRTEM.)

- R. Flowers of St. John's-wort,
seventy-two parts.

Olive Oil,

one hundred and forty-four parts.

Turpentine, seventy-two parts.

After twelve hours' digestion in a warm
place, set aside to cool, express, and add

Powdered Saffron, one part.

Triturate together. (VAN MONS)

It is slightly excitant, and is considered
vulnerary, anodyne, and pectoral.

It is applied in sprains and ulcers.

OIL OF FLOWERS.

Oleum Florum Slotani. (WIRTEM.;
SPIELMANN)

- R. Olive Oil, six pounds.
Blind Puppies, (*Chiens nouvellement*
nés), two.
Frogs, twelve.
Earth Worms, four ounces.
Spring Water, three pounds.

Boil slowly to the consumption of the
humidity, strain, and add

Human Fat, }
Fat of the Marmot, } of each,
Oil of Almonds, } six ounces.
— Scorpions, }

Fresh Flowers of Primrose

— Coltsfoot,

— Betony,

— Dyer's

Broom,

— Chamo-

mile,

— Violet,

— Lily of the

Valley,

— Lavender,

— Elder,

— Dill,

— Sage,

— Wallflower,

— Rue,

— Wild

Thyme,

— Rosemary,

— Lesser Cen-

taury,

— Veronica,

— Origanum,

— Tansey,

— Privet,

— Agrimony,

— Penny-

royal,

— Melilot,

— Lesser

Mullein,

— Worm-

wood,

— Roses,

— St. John's-wort,
four pounds.

Digest in a water-bath, express, and filter. (WIRTEM.)

The leaves and calices should be removed, if not the oil will be green instead of red.—SPIELMANN'S formula differs very little from this.

It used formerly to be esteemed anodyne, nervine, vulnerary, anti-spasmodic, and anti-nephritic.—The two following preparations come near it:

1. *Oleum Catellorum*.—Oil of Puppies.

R. Blind Puppies, three.
Olive Oil, forty-eight ounces.

Boil to the consumption of the humidity, and strain. Add

Herb of Origanum,	} of each, two ounces.
— Wild Thyme,	
— Pennyroyal,	
— St. John's-wort,	
— Marjoram,	

Infuse for fifteen days, and express.

2. *Oleum Vulpinum*.—Oil of Fox. (WIRTEM.)

R. Fox, flayed and emptied, (*Rénard écorché et vidé*), one.
Spring Water, six pounds.
Olive Oil, four pounds.
Muriate of Soda, six ounces.
Dill, } of each, one pound.
Thyme, }

Boil till the flesh becomes detached from the bones, and the humidity is dissipated; then strain and preserve.

GREEN OINTMENT.

Unguentum viride potabile. (PALAT.; WIRTEM.; SPIELMANN)

R. Herb of Chervil, three ounces.
— Sage, }
— Ground Ivy, }
— Scabious, }
— Coltsfoot, }
— Horehound, }
— Betony, }
— Ox-tongue, }
— Fennel, } of each,
— Veronica, } one ounce
— Comfrey, } and
— Golden Rod, } a half.
— Ladies-mantle, }
— Rue, }
— Milfoil, }
— Agrimony, }
— Plantain, }
— Nettle, }
— Hyssop, }
— St. John'swort, }
Black Snails, } of each, fifteen.
Shrimps, }
Fresh Butter, ten pounds.

Boil to the consumption of the humidity, and express. (PALAT.)

R. Fresh Herb of Southern-wood, }
— Mugwort, }
— Easter }
Daisy, }
— Betony, } of each,
— Fennel, } one ounce.
— Hyssop, }
— Mint, }
— Liver- }
wort, }
— Milfoil, }
— Marjoram, }
— Plantain, }
— Veronica, }
— Vervain, }
— Sage, }
— Rue, }
Green Juniper Berries, three ounces.
Butter, forty-eight ounces.

Boil, and strain. (WIRTEM.; and SPIELMANN)

It was formerly prescribed externally to disperse ecchymosis, and to appease pain; internally, after falls and blows, in the dose of one or two spoonfuls, in broth.

COMPOUND OIL OF ST. JOHN'S-WORT.

Oleum seu Balsamum Apparitii. (ELLIS)

R. Olive Oil, six pounds.
Tops of Rosemary, } of each,
— St. John's-wort, } one pound
— Rue, } and a half.

Digest for three days in a warm place; then add

Earth Worms, half a pound.

Boil to the consumption of the humidity, and dissolve in the strained oil,

Powdered Sandarac,	} of each, two ounces.
— Myrrh,	
— Olibanum,	
— Mastic,	one ounce.

Strain, and preserve.

VISCERAL LIQUOR. (WIRTEM.)

R. Root of Comfrey, } of each,
— Liquorice, } five ounces.
— Coltsfoot, }
Fresh Herb of Bugle, }
— Chervil, }
— Ground Ivy, } of each,
— St. John's- } four
wort, } handfuls.
— Plantain, }
— Scabious, }
— Veronica, }

Anethum Stellatum,	six drams.
Mace,	one dram.
Mastic,	three drams.
Living Craw-fish,	thirty.
Calf's Lung, chopped,	one.
Water of Bugloss,	one pound and a half.

Water of Chervil,	} of each,
Ground Ivy,	
Veronica,	
	one pound and a half.

Distil four pounds in a water-bath.—Dose, one or two ounces.

HYPOCHÆRIS MACULATA, LINN.—HUNGARIAN HAWK-WEED.

SYNONIMES:—Porcelle tachetée, FR.; Geflecktes Ferkelkraut, Fleckenferkelkraut, GERM.

BRUNS.

A plant of the North of Europe. (*Syn-
genesia Polygamia Æqualis*, LINN.; *Synan-
therea*, CASS.; *Fig. Zorn. Ic. pl. t. 271.*)

The herb (*Herba Costæ* seu *Costi nos-
tratis*) is employed. It consists of a star
of oval, oblong, dentate, entire radical
leaves. It has a nauseous and bitterish
taste.

HYSSOPUS OFFICINALIS, LINN.—HYSSOP.

SYNONIMES:—Hysope, FR.; Ysop, Isop,
Hysop, GERM.; Yzop, BOHEM.; Isop,
DAN.; and SUEC.; Hisopo, SPAN.;
Yzoop, DUT.; Issopo, ITAL.; Yzopeck,
POL.; Hysopo, PORT.

DUBL.; EDINB.; AMST.; ANT.; AUSTR.;
BATAV.; BAVAR.; BRUNS.; DAN.;
HISP.; GALL.; FERR.; FENN.; FORM.
HÔP. FR.; FULD.; GENEV.; HANN.;
HASS.; LIPP.; OLDEN.; LUSIT.; POL.;
BORUS; ROSS.; SAX.; SUEC.; WIRTEM.;
HERBIP.; BERGIUS; GUIBOURT; MUR-
RAY and GMELIN; PIDERIT; SPIEL-
MANN; ZARDA.

A plant of the middle of Europe. (*Didy-
namia Gymnospermia*, LINN.; *Labiæ*, JUSS.;
Fig. Jacq. Fl. Austr. III. t. 254.)

The herb (*Herba Hyssopi*) consists of a
square stem furnished with lanceolated,
linear leaves, more or less glabrous.

It has an agreeable odour, with a bitter,
warm, and aromatic taste.

It is stimulant and expectorant.

CONSERVE OF HYSSOP. (ROSS.; VAN
MONS)

R. Fresh Leaves of Hyssop, bruised,
one part.
Powdered Sugar, two parts.

Mix, and keep in a cool place. (ROSS.)

VAN MONS gives the same proportions,
but requires the mixture to be slightly
heated.

INFUSION OF HYSSOP. (FORM. HÔP. FR.;
RATIER)

R. Leaves of Hyssop, two drams and a half.
Grated Liquorice, two drams.
Hot Water, a sufficient quantity
to obtain two pounds of infusion. (FORM.
HÔP. FR.)

R. Tops of Hyssop, two drams.
Boiling Water, two pounds.

Infuse, and strain. (RATIER)

It is slightly aromatic, and is given to
favour expectoration in catarrhs.

SYRUP OF HYSSOP.

Syrupus Hyssopi seu *de Hyssopo*. (HISP.;
GALL.; SARD.)

R. Dried Tops of Hyssop, one ounce.
Distilled Water of Hyssop, two pounds.

Digest in a water-bath for two hours, in
a covered vessel; strain, filter, and add to
the infusion,

White Sugar, double the quantity.

Melt in a water-bath, in a covered vessel.
(GALL.)

HENRI proposes to mix one part of the
infusion with three parts of simple syrup,
previously boiled till very thick.

R. Dried Tops of Hyssop, two ounces.
Water, two pounds.

Infuse, and add to the strained liquor,

White Sugar, four pounds.

Strain the syrup. (HISP.)

SARD. prescribes one part of sugar and two parts of infusion, or an equal part each, of simple syrup and of depurated juice of syrup.

Dose, from half an ounce to two ounces.

COMPOUND SYRUP OF HYSSOP. (BRUNS.; WIRTEM.)

- R. Maidenhair, half an ounce.
 Root of Elicampagne, one ounce.
 ——— Smallage, half an ounce.
 Water, two pounds and a half.

Boil, and pour sixteen ounces of the strained decoction upon

Dried Tops of Hyssop, two ounces.

Digest in a warm place, and express; clarify, and add

White Sugar, two pounds eight ounces.

SCILLITIC SYRUP OF HYSSOP.

Syrupus Hyssopi cum Scilla seu scilliticus. (AMST.)

- R. Dried Tops of Hyssop, two ounces.
 ——— Squill, one ounce.
 Ginger, bruised, half an ounce.
 Water of Hyssop, two pounds.

Digest for twenty-four hours, and express; set the liquor aside to depurate by repose; and finally add fifteen parts of white sugar to every eight of the strained infusion, and make a syrup with a gentle heat.

OIL OF HYSSOP.

Oleum Hyssopi æthereum, Ætheroleum Hyssopi. (FERR.; WIRTEM.)

- R. Herb of Hyssop, twenty-five parts.
 Water, seventy-five parts.

Muriate of Soda, three parts.
 Digest for three days, and distil.

DISTILLED WATER OF HYSSOP. (AUSTR.; BAVAR.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SARD.; SAX.; PIDERIT; SWEDIAUR; VAN MONS)

- R. Fresh and Flowering Herb of Hyssop, one part.
 Water, a sufficient quantity.
 Distil ten parts. (HANN.; OLDEN.; POL.; BORUS.; SWEDIAUR)

AUSTR. prescribes one part of herb and twelve parts of water; four parts to be distilled.—FERR.; and PALAT.—one of herb and four of water; two to be distilled.—SAX.—one of herb and fifteen of water; one third to be distilled.—BAVAR.—one of herb and sixteen of water; four to be distilled.—HASS.; and PIDERIT.—one of herb and eight of water; four to be distilled.—GALL.—five of herb and twenty of water; ten to be distilled.—SARD.—one of herb and four of water; one half to be distilled, and the distillation to be repeated with fresh herb.

- R. Herb of Hyssop, three parts.
 Put it into a bag, suspended in the centre of a cucurbit containing

Water, twelve parts,
 not allowing it to touch this. Distil six parts. (FULD.; LIPP.)

- R. Flowering Herb of Hyssop, ten parts.
 Proof Spirit, one half part.
 Water, one hundred and sixty parts.
 Distil forty parts, and separate the oil. (VAN MONS)

Dose, two or three ounces.

JACEA NIGRA, MÆ.—COMMON Knapweed.

SYNONIMES: — Jacée, FR.; Gemeine schwarze Flockenblume, GERM.; Knopurt, DAN.; Knoopkruid, DUT.; Knapper, SWED.

GALL.; GENEV.

A very common plant of Europe. (*Syn- genesia Polygamia Frustranea*, LINN.; *Synantherea*, CASS.; *Fig. Cæ. Fl. Dan.* t. 519)

The root and herb are employed.

The root (*Radix Jaceæ nigrae*) is woody and possesses a slightly astringent taste.

The herb consists of a branching, hard, nearly glabrous stem, and of leaves of which the lower are dentated, and the upper sessile, oval, lanceolated, and usually entire. It has a sweetish yet rather bitter taste.

It is astringent and tonic.

IGNATIA AMARA, Lf.

BRUNS.; HISP.; GALL.; FERR.; SAX.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SAUNDERS; SPIELMANN.

A shrub of the Philippine Islands. (*Pentandria Digynia*, LINN.; *Apocynæ*, JUSS.; *Fig.* Philos. Trans. XXI. t. 1.)

The seed (St. Ignatius' Bean; *Faba Sancti Ignatii* seu *Indica* seu *febrifuga*) is employed. It is of the volume of a filbert, and of a variable figure, rather wrinkled, fawn or bistre coloured externally, and hav-

ing the appearance of being sprinkled with a sort of silvery farina which is very adherent; internally it is of a greenish-brown colour, and of a horny consistence.

It contains, according to PELLETIER and CAVENTOU, strychnine and strychnic acid.

Its composition is the same as that of the *nux vomica*, but it contains much more strychnine and less brucine. Its properties are also the same.

Dose, from two to six grains of the powder.

ILEX.

Two species of this genus of plants are met with in pharmacopœias:

1. *Ilex Aquifolium*, LINN.—Holly.

SYNONIMES:—Houx Commun, FR.; Stechpalme, Hælsendorn, GERM.; Chrishorn, Charetorn, Stovtidse, Beenved, DAN.; Acebo, SPAN.; Steckpalmen, Hulst, DUT.; Agrifolio, ITAL.; Ostrozkow, Kzewina, POL.

BRUNS.; DAN.; GALL.; OLDEN.; ROSS.; SAX.; BERGIUS; GUIBOUT; MURRAY and GMELIN.

A little tree of the middle of Europe. (*Tetrandria Tetragynia*, LINN.; *Rhamnæ*, JUSS.; *Fig.* Zorn. Ic. pl. t. 372.)

The leaves (*Folia Aquifolii*) are employed. They are petiolated, oval, coriaceous, shining, of a fine green colour, most frequently waved, angulous, dentated, and spiny. They are without odour, and have a slightly astrigent, mucilaginous, and bitter taste.

It is tonic, and is particularly extolled in rheumatism. It has also been employed in intermittent fevers.—Dose, from half a dram to one dram of the powder. A decoction may also be prepared with from half a dram to one ounce of the leaves to two pounds of water; and a wine, with one part of leaves and forty parts of wine. The latter is reputed febrifuge: it is given in the dose of from four to five ounces.

2. *Ilex vomitoria*, AIT.

SYNONIMES: Apalachine, Thé de la mer du Sud, FR.

BRUNS.; GALL.; WIRTEM.; COXE; SPIELMANN.

A plant of Florida. (*Fig.* Herb. Kew. I, p. 170.)

The leaves are employed. They are elliptical, lanceolated, obtuse, serrated on the borders, glabrous, and shining.

It is very strongly emetic and diuretic.

ILlicium ANISATUM, LINN.—INDIAN ANISE.

SYNONIMES:—Badiane, Anis étoilé, FR.; Sternanis, GERM.; Badianehuttaie, ARAB.; Badian, BOHEM.; Pa-co-hu-huei-hiam, CHIN.; Stierneanys, DAN.; Anaspul, DUK.; Anis de la China, SPAN.; Sternanys, DUT.; Anice stellato, ITAL.; Skimmi, Somo, JAP.; Gwiazdkowy, POL.; Anis estellado, PORT.; Badyan, Anyz, ROSS.; Stjernanis, SWED.; Anasipu, TAM.
VOL. II.

AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A shrub of China and of Japan. (*Polyan-*

dria Polygynia, LINN.; *Magnioleæ*, JUSS.; *Fig. Flore Medic. I. 30.*)

The fruit (*Anisum stellatum* seu *Sinense*, *Semen Anisi Stellati*, *Capsulæ seminalis Anisi Sinensis*) is employed. It consists of several oval, compressed, bivalve capsules, disposed in the form of an orbicular star, each containing a small lenticular, smooth nut of a reddish-grey colour, composed of a thin fragile shell covering a white kernel. Its odour is penetrating, aromatic, very agreeable, and resembles that of anise as it does in taste.

It is excitant and stomachic.—Dose, one

scruple of the powder. An infusion may be made with one or two drams, to one pound of water.

ANTI-CATARRHAL DRINK. (PIERQUIN ;
SAINTÉ-MARIE)

R. Honey, four ounces.
Water, two pounds.

Boil for a quarter of an hour, remove from the fire, and add

Indian Anise, two drams.
Root of Elecampane, one scruple.

Strain when cold.—To be taken by glassfuls.

IMPERATORIA OSTRUTHIUM, LINN.—MASTER-WORT.

SYNONIMES:—Imperatoire, FR.; Meisterwurz, Kaiserwurz, Ostranz, GERM.; Wasedobr, BOHEM.; Mesterwurt, DAN.; Imperatoria, ITAL.; PORT.; and SPAN.; Mester-wortel, DUT.; Mæsterrot, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of the temperate climates of Europe. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Flore Medic. IV. 200.*)

The root (*Radix Imperatoria*, seu *Ostruthii* seu *Magistrantia* seu *Astrantia*) is employed. It is thick, branching, nearly tuberculous, ringed, and furnished with long fibres. It is of a greenish-brown without,

and white within; its odour is strong and aromatic; and its taste is acrid, bitter, and disagreeable.

It is excitant, carminative, diaphoretic, and sialagogue. HOFFMANN called it a *divinum remedium*, but there is nothing peculiar about it, and is now almost out of use.—Dose, from twenty to thirty grains of the powder.

MASTICATORY POWDER. (SPIELMANN)

R. Root of Master-wort, } of each,
Elder Bark, } half an ounce.
Bay Leaves, one dram and a half.
Seeds of Mustard, } of each,
Stavesacre, } one dram.
Cloves, four scruples.
Myrrh, six drams.

Mix.—Dose, one scruple, enclosed in a bag. It is chewed in aphonia and paralysis of the tongue.

INULA.

Two species of this genus of plants are used in medicine:

1. *Inula dysenterica*, LINN.—Middle-sized Flea-bane.

SYNONIMES:—Aunée anti-dysenterique, Herb de Saint-Roch, FR.; Dürrwurz, Ruhralant, GERM.; Rooloops alant, DUT.

BRUNS.; GALL.; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe, growing in moist places. (*Syngenesia Polygamia Su-*

perflua, LINN.; *Synanthereæ*, CASS.; *Fig. Zorn. Ic. pl. t. 272.*)

The herb (*Herba Conyza mediæ* seu *Arnica spuria* seu *Suedensis*) is employed. It consists of a rounded, tomentose stem, furnished with amplexicaul, oblong, heart-shaped leaves, serrated, undulated, wrinkled, and sub-tomentose. It is destitute of odour, and has an acrid, aromatic, and asstringent taste.

2. *Inula Helenium*, LINN.—Elecampane.

SYNONIMES:—Aunée commune, FR.;

Alant, Alantwurz, Alantwurz, Boustaland, Oland, Helenenkraut, GERM.; U-sululrasun, ARAB.; Women, BOHEM.; Alant Sanet Ellensron, DAN.; Enula campana, SPAN.; and PORT.; Alants-wortel, DUT.; Enula campana, Elenio, ITAL.; Bekhisanjubischami, PERS.; O-ma-nowy, POL.; Alant, SWED.

DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PIDE-RIT.; SPIELMANN.; ZARDA.

A plant of every part of Europe. (*Fig. Flore Medic. I. 48.*)

The root (*Radix Enulæ seu Enulæ campanæ seu Helenii*) is employed. It is fusiform, of the thickness of the thumb, fleshy, branching, furnished with a few fibrils, fawn-coloured or brown without, and white within. When fresh, it has, a strong and penetrating odour, but after desiccation, one resembling that of violets. Its taste is peculiar, bitterish, aromatic, and lively.

It contains a volatile oil, a kind of fecula called *Inuline*, a bitter substance, and a resinous matter.

It is tonic, diuretic, sudorific, expectorant, anthelmintic, and emmenagogue.—Dose, from half a dram to two drams of the powder.

I. Preparations which contain Elecampane in substance.

CONSERVE OF ELECAMPANE.

R. Fresh Root of Elecampane, one pound.
Water, eight pounds.

Macerate for several days; then soften the root by boiling; and clarify in the decoction,

White Sugar, half a pound.

Digest the root in the syrup for three days; then boil till this becomes very thick, and preserve the root in it, or dry it in a stove. (*SARD.*)

R. Root of Elecampane, any quantity.

Boil it in a sufficient quantity of water, till it may be bruised between the fingers; pound it in a marble mortar; strain the pulp through a silken sieve, and add to it twice its weight of syrup, boiled *à la plume*. (*SARD.*)

GALL. directs one part of pulp and four parts of sugar boiled to the consistence of

an electuary in decoction of the root.—VAN MONS—one of pulp and three of white sugar.

AROMATIC POWDER OF ELECAMPANE. (SAUNDERS)

R. Powdered Root of Elecampane, six parts.
Angelica, one part.

Mix.

ANTI-LYSSIC POWDER.

Pulvis anti-lyssicus Ormskirckianus. (NIEMANN)

R. Root of Elecampane, one dram.
Chalk, half an ounce.
Armenian Bole, three drams.
Alum, ten grains.
Oil of Anise, five drops.

Pulverize and mix.

This powder was formerly celebrated, but like all those of the same kind, its pretended virtues have all gradually become obsolete.

OINTMENT OF ELECAMPANE.

Unguentum Inulæ seu Helenii seu enulatum. (BRUNS.; HISP.; WIRTEM.; AUGUSTIN.; HUFELAND.; NIEMANN.; SPIELMANN.; SWEDIAUR)

R. Root of Elecampane, half a pound.
Spring Water, one hundred and twenty-eight ounces.

Boil, reduce to pulp, and add

Fresh Butter, four ounces.

Mix. (NIEMANN)

SWEDIAUR directs half a pound of root to be boiled in one pound of water, and three ounces of butter to be added to the pulp.—AUGUSTIN.; and HUFELAND—the root to be boiled in a sufficient quantity of water, and reduced to pulp, and then a sufficient quantity of lard to be added.

R. Root of Elecampane, one pound.
Lard, half a pound.
Olive Oil, four ounces.

Triturate in a stone mortar; then boil over a slow fire to the consumption of the humidity, express, strain, and add,

Yellow Wax, } of each, one ounce.
Turpentine, }

Mix. (BRUNS.; WIRTEM.; SPIELMANN)

R. Compound Oil of Elecampane, four pounds.
Yellow Wax, one pound.
Lard, four ounces.

Melt together, and add to the mass when half cold,

Liquid Storax, eleven drams.
 Powdered Mastic, six drams.
 Stir till cold. (HISP.)
 It has been recommended in itch.

OFFICINAL THERIAC. (LIPP.)

R. Elecampane, two ounces.
 Zedoary, }
 Myrrh, } of each,
 Round Birthwort, } one ounce and
 Angelica, } a half.
 Rob of Juniper, six ounces.
 Despumated Honey, } of each,
 Syrup of Poppies, } eight ounces.

Mix.—Dose, from one dram to half an ounce.

PASTILLES OF ELECAMPANE. (BORIES;
CADET DE GASSICOURT)

R. Powdered Elecampane, half an ounce.
 Florentine Iris, one dram.
 Mucilage of Tragacanth,
 a sufficient quantity.

They are recommended in scurvy, dropsy, and chlorosis, in the dose of half an ounce daily.

II. Preparations which contain only a part of the Constituent Principles of Elecampane.

A.

*Extraction by Distillation.*DISTILLED WATER OF ELECAMPANE.
(GALL.)

R. Fresh Root of Elecampane, one part.
 Water, five parts.
 Distil two parts.

ANTI-ASTHMATIC LIQUOR. (BRUNS.;
PALAT.; WIRTEM.; SPIELMANN)

R. Root of Elecampane, }
 Liquorice, } of each,
 Florentine Iris, } two ounces.
 Herb of Ground Ivy, } of each,
 Coltsfoot, } half
 Jerusalem Oak, } a pound.
 Hyssop, }
 Horehound, } of each,
 Pennyroyal, } three ounces.
 Sage, }
 Anis, } of each, one ounce.
 Fennel, }
 Alcohol, } one pound.
 Spring Water, } ten pounds.

Digest for three days, and distil six or seven pounds. (WIRTEM.)

R. Root of Elecampane, two ounces.
 Burnet Saxi- }
 frage, } of each,
 Florentine Iris, } one ounce
 and a half.
 Herb of Hyssop, }
 Horehound, } of each,
 Sage, } two ounces.
 Anis, }
 Fennel, } of each, one ounce.
 Juniper, }
 Bay Berries, }
 Lesser Cardamoms, } of each,
 Cinnamon, } half an ounce.
 Ginger, }
 Storax, } six drams.
 Alcohol, } twelve pounds.
 Water of Acacia, } eight pounds.

Digest for three days, distil twelve pounds, and add to the product,

Rose Julep, two pounds.
 Fennel Water, eight pounds.

Mix. (BRUNS.)

R. Root of Elecampane, }
 Liquorice, } of each,
 Florentine Iris, } one ounce.
 Angelica, }
 Burnet Saxifrage, } of each,
 six drams.
 Herb of Maidenhair, }
 Hyssop, } of each,
 Lung-wort, } half an ounce.
 Ground-Ivy, }
 Sage, }
 Coltsfoot, }
 Seeds of Anise, }
 Fennel, }
 Nettle, }
 Saffron, } two scruples.
 Alcohol, } one pound.
 Water of Scabious, }
 Horehound, } of each,
 Speedwell, } half a pound.

Digest for three days, and distil in a water-bath. (PALAT.)

R. Root of Elecampane, two ounces.
 Florentine Iris, one ounce and a half.
 Herb of Hyssop, }
 Rosemary, } of each,
 Sage, } one ounce.
 Anise, }
 Fennel, } of each, half an ounce.
 Cinnamon, }
 Lesser Cardamoms, } two drams.
 Saffron, } two scruples.
 Alcohol, } twelve ounces.
 Hyssop Water, } twenty-four ounces.

Digest for three days and distil. (SPIELMANN)

It is recommended in the mucous coughs of aged people.—Dose, one or two ounces in the morning.

B,

Extraction by Mechanical Means.

SYRUP OF ELECAMPANE.

Syrupus Enulæ. (WIRTEM.)

R. Depurated Juice of Elecampane
Root, one pound and a half.
White Sugar, two pounds and a half.
Dissolve, and strain.

C.

Extraction by Water.

PTISAN OF ELECAMPANE. (GALL.)

R. Root of Elecampane, one ounce.
Boiling Water, two pounds.
Infuse, strain, and add
Honey, or Syrup of Maidenhair,
one ounce.
Mix.

AQUEOUS EXTRACT OF ELECAMPANE.

(AMST.; ANT.; AUSTR.; BATAV.; BRUNS.;
HISP.; GALL.; GENEV.; LIPP.; PALAT.;
SUEC.; VAN MONS)

R. Fresh Root of Elecampane,
any quantity.

Having washed it well, rasp it; and express the juice; bruise the residue with a little water, and express: strain the liquor through flannel, and evaporate it with a gentle heat to the consistence of an extract. (GENEV.; VAN MONS)

R. Root of Elecampane, one pound.
Spring Water, six pounds.

Macerate for four days in a warm place, then boil a little: express and evaporate the decoction. (BRUNS.; LIPP.; PALAT.)

R. Root of Elecampane, one pound.
Cold Water, four pounds.

Macerate in a covered vessel, stirring occasionally, and strain; pour more water on the residue and macerate again; then express slightly, mix the two liquors, and having strained, evaporate them in a water-bath. (GALL.)

R. Root of Elecampane, one part.
Boiling Water, eight parts.

Digest for twenty-four hours, then boil for a quarter of an hour, and strain; boil the residue for a quarter of an hour in four parts of fresh water, and strain. Mix the two liquors, set aside, decant, and evaporate. (ANT.)

BATAV. prescribes twenty-four hours' digestion and two hours' ebullition with ten parts of water, and a second ebullition for

two hours with eight parts of water.—AMST.—boiling first with twenty, then with fifteen, parts of water, to be reduced each time to one half.—HISP.—twenty-four hours' digestion and two hours' boiling in ten parts of water, then a second boiling with a fresh quantity of water to be reduced to one half.

R. Root of Elecampane, any quantity.
Water, a sufficient quantity.

Exhaust the root by several successive boilings, mix the decoctions, and after decantation, evaporate. (AUSTR.; SUEC.)

Dose, from a scruple to half a dram.

PECTORAL MIXTURE. * (CADET DE GASSI-COURT)

R. Extract of Elecampane, two scruples.
Gum Arabic, two drams.
Hyssop Water, half an ounce.
Oxymel of Squill, } of each, one ounce
Syrup of Hyssop, } and a half.

Mix.—To be taken by spoonfuls.

ANTI-ASTHMATIC MIXTURE. (SPIELMANN)

R. Extract of Elecampane, } of each,
Cascarilla, } one scruple.
Tincture of Burnet Saxifrage, } of each,
Liquorice, } one dram.
WEDEL'S Pectoral Elixir, two drams.
Oxymel of Squill, half an ounce.
Water of Coltsfoot, }
Lungwort, } of each,
Hyssop, } one ounce.
Fennel, }
Speedwell, }
Syrup of Marshmallows, }

Mix.—To be taken by spoonfuls.

DECOCTION OF ELECAMPANE.

Decoctum Helenii. (NIEMANN)

R. Root of Elecampane, half an ounce.
Spring Water, a sufficient quantity
to obtain six ounces of strained decoction.
—Dose, one or two spoonfuls, every two hours.

COMPOUND DECOCTION OF ELECAMPANE.

Hydromel composé. (RATIER)

R. Root of Elecampane, one ounce.
Tops of Hyssop, } of each,
Leaves of Ground Ivy, } two drams.
Water, a sufficient quantity.

Boil, and add when strained,

Syrup of Honey, two ounces.

Mix.—It is slightly stimulant, and is recommended in chronic catarrh, asthma, and aneurisms of the heart.

CORDIAL DIURETIC POTION. (RATIER)

- R. Decoction of Elecampane, four ounces.
 Tincture of Digitalis, half an ounce.
 ——— Potass, eighteen drops.
 Compound Syrup of Smalage, one ounce.

Mix.—It is stimulant and diuretic.—To be taken by spoonfuls.

ALTERATIVE BROTH. (PIERQUIN)

- R. Neck of Mutton, four ounces.
 Root of Elecampane, half a dram.
 ——— Wild Patience, one dram.
 Bruised Crabs, two.
 Leaves of Cichory, } of each,
 ——— Fumitory, } one handful.
 ——— Cress, }

Boil in a sufficient quantity of water.

PECTORAL OXYMEL. (BRUNS.; CADET DE GASSICOURT; SPIELMANN)

- R. Root of Elecampane, one ounce.
 ——— Florentine Iris, half an ounce.
 Spring Water, thirty-six ounces.

Boil down to twenty-four ounces, strain, set aside, and add

- Honey, sixteen ounces.
 Gum Amoniac, one ounce.

Dissolved in

- Vinegar, eight ounces.

Boil to the consistence of honey.

It is recommended in humid cough and asthma.—Dose, a spoonful frequently.

D.

Extraction by Vinegar.

THERIACAL VINEGAR.

Acetum theriacale. (PALAT.)

- R. Root of Elecampane, }
 ——— Angelica, } of each,
 ——— Cyperus, } half
 ——— Zedoary, } an ounce.
 ——— Avens, }
 ——— Contrayerva, }
 ——— Master-wort, }
 ——— Valerian, }
 ——— Viper's Bugloss, }
 Fresh Orange Peel, }
 ——— Lemon Peel, }
 Cloves, }
 Cinnamon, } of each,
 Galanga, } two drams.
 Juniper Berries, }
 Bay Berries, }
 Tops of Sage, }
 ——— Rosemary, }
 ——— Rue, }
 Vinegar, seven pounds.

Macerate for a month in a warm place; then express strongly, and add

- Theriac, seven ounces.

Macerate for one month, and filter.

It is recommended internally, in the dose of from half a spoonful to one spoonful, as a sudorific and as a preservative against contagion; it is also applied externally as a corroborant.

E.

Extraction by Wine.

WINE OF ELECAMPANE.

Vinum Enulae seu enulatum. (FORM. HÔP. FR.; SARD.; NIEMANN; VAN MONS)

- R. Fresh Root of Elecampane, one part.
 White Wine, sixteen parts.

Digest without heat for several days; then express, and filter. (VAN MONS)

SARD. prescribes two ounces of dried root, four pounds of white wine, and twenty-four hours' maceration.

- R. Tincture of Elecampane, one part.
 Red Wine, ten or twenty parts.

Mix. (FORM. HÔP. FR.)

F.

Extraction by Alcohol.

TINCTURE OF ELECAMPANE.

Tinctura Helenii. (AMST.; AUSTR.; ANT.; BATAV.; PALAT.)

- R. Root of Elecampane, one part.
 Alcohol (15 degrees), six parts.

Digest for three days in a moderate heat, and filter. (BATAV.)

AUST. prescribes one part of root and six parts of alcohol (0.910).—AMST.—one of root and six of alcohol (0.917).—PALAT.—one of root and four of alcohol.

- R. Root of Elecampane, one ounce.
 Alcohol (20 degrees), four ounces.

Infuse in a water-bath for four days; strain, and pour upon the residue

Alcohol (20 degrees), two ounces.

Macerate again for two days, strain, mix, and filter.

It is stimulant, carminative, and diaphoretic, and is principally recommended in affections of the chest.—Dose, from six drops to one dram.

ALCOHOLIC EXTRACT OF ELECAMPANE.
(BAVAR.; FULD.; HANN.; OLDEN.;
POL.; SAX.; WIRTEM.; HERBIP.)

R. Root of Elecampane, } of each,
Alcohol, } two pounds.
Water, } nine pounds.

Digest for twelve hours, express, distil off the alcohol, and evaporate the residue to the proper consistence. (POL.)

BAVAR.; HANN.; OLDEN.; BORUS.; and SAX. prescribe two parts of root, three of alcohol, and nine of water.—FULD.—one of root, six of alcohol, and three of water.

R. Root of Elecampane, one pound.
Alcohol, five or six pounds.

After sufficient extraction, strain with strong expression; digest, and then boil the residue in three pounds of water. Having clarified the decoction with white of egg, mix the two liquors, distil off the alcohol, and evaporate to the proper consistence. (WIRTEM.)

R. Root of Elecampane, any quantity.
Alcohol, a sufficient quantity

to cover it by four fingers' breadth.
Digest in a warm place, and decant; boil the residue for an hour with water, strain, and evaporate the liquor to the consistence of honey, adding the tincture towards the end, stirring constantly that the mass may be homogeneous. (WIRTEM.)

The dose is the same as that of the aqueous extract.

ANTI-HYSTERIC ELIXIR. (CADET DE GASSICOURT; VAN MONS)

R. Root of Asarabaca, three parts.
—— Florentine Iris, five parts.
—— Elecampane, } of each,
—— Sweet Flag, } ten parts.
—— Liquorice, fifteen parts.
Seeds of Anise, five parts.
Proof Spirit, eighty parts.

Infuse without heat for several days; strain, and add

Camphor, one part.
Dissolve. (VAN MONS)

CADET DE GASSICOURT prescribes one dram and a half of liquorice, one dram of sweet flag, half a dram each of iris and of anise, eighteen grains of asarabaca, six grains of camphor, and half a pound of alcohol.

It is stimulant, and is principally recommended in humid asthma.—Dose, from ten drops to thirty.

ELZNER'S PECTORAL ELIXIR. (VAN MONS)

R. Root of Liquorice, } of each,
Saffron, } four parts.
Indian Anise, }

Root of Elecampane, } of each,
—— Florentine Iris, } eight parts.
Spirit of Hyssop, } ninety-six parts.

Infuse in a water-bath; then filter, and add

Balsam of Peru, one part.
Honey, sixteen parts.

Set aside, and filter.

COURCELLE'S AMERICAN ELIXIR. (CADET DE GASSICOURT; NIEMANN)

R. Alcohol (32 degrees), } of each,
Brandy (21 degrees), } twelve pounds.
Root of Elecampane, four pounds.
—— Sugar Cane, } of each,
—— Round Birth- } three pounds.
wort, }
—— Arundo Donax, } of each,
Leaves of Avocatoria, } two pounds.
Flowers of St. John's-wort, one pound.
—— Elder, half a pound.
Bark of Iron Wood, six ounces.
Leaves of Orange, } of each,
Croton Balsamiferum, } four ounces.
Juniper, three ounces.
Opium, two ounces and a half.
Justicia Assurgens, } of each,
Flowers of Orange, } two ounces.
—— Lime, }
Tops of Rosemary, } of each,
Root of Asarabaca, } one ounce.
—— Palma Altissima, }
Calebash, }

Macerate, and strain. Burn the residue and infuse the ashes in the liquor with

Flowers of Red Poppies, six ounces.
Madder, three ounces.

Filter.

CADET DE GASSICOURT has corrected this formula in the following manner:

R. Alcohol, one hundred and twenty pounds.
Root of Elecampane, sixteen pounds.
Flowers of St. John's-wort, eight pounds.
Leaves of Orange, six pounds.
Flowers of Elder, five pounds.
Leaves of Balm, four pounds.
Flowers of Lime, two pounds and a half.
Root of Provence Cane, } of each,
Juniper, } two pounds.
Flowers of Rosemary, }
Opium, }
Root of Asarabaca, one pound.
Water, a sufficient quantity,

that the liquor may mark twenty-four degrees. Colour with red poppies.

This is a highly esteemed stimulant in America; it is administered in many affections attributed either to nervous spasms or to debility, and is regarded as an almost universal panacea.—Dose a large spoonful,

either alone or in a proper drink ; it may be taken twice or thrice a day.

G.

Extraction by Oil.

COMPOUND OIL OF ELECAMPANE.

Oleum Enulæ compositum. (HISP.)

R. Root of Elecampane,	} of each, three ounces.
—— Valerian,	
—— Burdock,	
Tops of St John's-wort,	
—— Southernwood,	
—— Wormwood,	
—— Sweet Basil,	
—— Mountain Balm,	
—— Astmary,	

Tops of Marjoram,	} of each, three ounces.
—— Mint,	
—— Savine,	
Leaves of Sage,	
Flowers of Elder,	} of each, two ounces.
Flowering Tops of Chamomile,	
Tops of Melilot,	
—— French Lavender,	
Fresh Leaves of Bay,	} of each, six ounces.
—— Dwarf Elder,	
—— Rosemary,	
—— Rue,	
Seeds of Cumin,	} of each, one ounce.
—— Fenugreek,	
—— Nettle,	
Olive Oil,	twelve pounds.

Macerate for four days ; then boil to the consumption of the humidity, and set aside to depurate by repose.

IODIUM.—IODINE.

SYNONIMES :—*Iodinium* (DUBL.).—Iode, Fr.

DUBL. ; BAVAR. ; FERR. ; MAGENDIE ; NIEMANN ; VAN MONS.

R. Mother Liquors of the Soda obtained from Bladder-Wrack, any quantity.

Add concentrated sulphuric acid to it in excess, and boil the liquor slowly in a glass retort furnished with a receiver ; collect the brilliant slender crystals which form in the latter, wash them in a little cold water, and dry them slowly.

It is stimulant, powerfully irritant, and even poisonous in large doses. Under certain circumstances, it exercises upon the mammary and thyroid glands, the latter particularly, a very remarkable action, making them waste away more or less. It has been recommended in the treatment of scrofula and of goitre, of lymphatic swellings of every kind, gonorrhœa, chronic leucorrhœa, enlargement of the testicle, &c. It is used both internally and externally ; but the greatest circumspection is required in its administration, and it should be discontinued as soon as wasting of the flesh is seen to come on : this wasting is said to be, in general, the first symptom of its acting noxiously.—Dose, from the eighth of a grain to one grain, twice a day, in pills.

As we have done with regard to arsenic and hydrocyanic acid, we shall collect into one article all the preparations of this substance, which appear to have an identical

action ; we except only the mercurial iodurets which are placed in the article on mercury.

TINCTURE OF IODINE.

Tinctura seu Alcohol Iodii. (BAVAR. ; FERR. ; BORIES ; FEE ; MAGENDIE ; NIEMANN ; PIERQUIN ; RATIER ; VAN MONS)

R. Iodine, one part.
Alcohol (35 degrees), twelve parts.

Dissolve. (FERR ; BORIES ; FEE ; MAGENDIE ; RATIER)

PIERQUIN prescribes one part of iodine, and eight parts of alcohol.—BAVAR.—one of iodine, and nineteen of alcohol.—VAN MONS—one of iodine, and a little more than ten of alcohol.—NIEMANN—one of iodine, and seventeen of alcohol.

Twenty drops of the tincture made according to the first formula contains about one grain of iodine, and one drop weighs two thirds of a grain.—Dose, for adults, from four to ten drops thrice a day in a little sweetened water, augmenting it progressively to twenty drops or more.

SYRUP OF IODINE. (FERR.)

R. Tincture of Iodine, two drams.
Simple Syrup, four ounces.

Mix without heat.

ÆTHEREAL TINCTURE OF IODINE.

Tinctura Iodii ætherea. (FERR.)

- R. Iodine, six grains.
 Æther (66 degrees), one dram.
 Dissolve.

PILLS OF IODINE.

Pilule Iodatæ. (FERR.; BRERA)

- R. Iodine, one grain.
 Liquorice Powder, one scruple.
 Rob of Elder, a sufficient quantity.
 Make two pills.

OINTMENT OF IODINE.

Unguentum Iodatum. (FERR.; BRERA)

Unguentum Iodinii. (DUBL.)

- R. Iodine, one scruple.
 Prepared Lard, one ounce.
 Triturate, and form an ointment.

IODIC ACID.

Acidum Iodicum. (VAN MONS)

- R. Pulverized Iodate of Potass, any quantity.

Pour upon it a sufficient quantity of tartaric acid dissolved in the least possible quantity of water. Decant the liquid after a short time, and evaporate it slowly, allowing it to cool very frequently, till the acid ceases to be deposited.

IODATE OF POTASS. (FERR.; VAN MONS)

- R. Iodine, any quantity.
 Caustic Potass, a sufficient quantity.

Evaporate the solution to dryness; treat the residuum with alcohol, and preserve what is not dissolved. (FERR.)

- R. Soda from Bladder-Wrack, sifted
 and powdered, one part.
 Water, slightly alkaliized by potass, two parts.

Boil, and repeat the operation with the same quantity of water; mix the liquors, filter, evaporate, remove the crystals as they form, leave them to drip, and then pulverize them: having washed them in hot water, mix the waters in which they have been washed with that which dripped from them, evaporate the whole to dryness, and preserve the residuum. (VAN MONS)

The latter contains much chloruret of sodium and of potassium.

IODATE OF LIME.

The preparation of this salt is not indicated in any pharmacopœia.

RESOLVENT BOLUSES.

Boli resolventes despumentes. (BRERA)

- R. Iodate of Lime, from six to ten grains.
 Extract of Aconite, twenty-four grains.
 Make six boluses.—Dose, one, every four hours; in chronic bronchitis and tuberculous phthisis.

EMMENAGOGUE BOLUSES.

Boli emmenagogi resolventes. (BRERA)

- R. Iodate of Lime, ten grains.
 Extract of Savine, twelve grains.
 Make four boluses.—Dose, one every four hours; in amenorrhœa complicated with scrofula.

HYDRIODIC ACID.

Acidum hydriodicum. (FERR.; HENRI)

- R. Phosphuret of Iodine, any quantity.
 Put it into a retort, and having moistened it a little, apply a gentle heat; collect the acid which is disengaged, by the aid of a long bent tube, at the bottom of a narrow test-glass.

It may also be prepared by passing a current of hydrosulphuric acid gas through a solution of iodine in distilled water, till the liquor becomes colourless; and then carefully evaporating this under the receiver of a pneumatic machine, after having filtered it.

PROTO-HYDRIODATE OF POTASSIUM.

Hydriodas Potassæ. (DUBL.; FERR.; FEE; HENRI; MAGENDIE; NIEMANN)

- R. Iodine, one part.
 Sulphuret of Iron, in coarse powder, five parts.
 Sulphuric Acid, seven parts.
 Distilled Water, forty-eight parts.
 Solution of Carbonate of Potass, a sufficient quantity.
 Rectified Spirit, six parts.

Triturate the iodine with sixteen parts of water, and put the mixture into a glass vessel; then put the sulphuret into a matrass, and pour over it the acid previously diluted with thirty-two parts of water; then, having adapted a tube which dips to the bottom of the vessel containing the solution of iodine, to the neck of the matrass, pass the gas through the solution till the iodine disappears. Eva-

porate the filtered solution at a high temperature, and filter again: then add gradually the solution of carbonate of potass to saturate the acid, which may be known by the effervescence ceasing. Set the mixture apart in a warm place till a dry white salt remains; pour the spirit upon this, and dissolve with heat. Finally, pour off the liquid from the salt, evaporate to dryness, and preserve the residuum in a close vessel.

Turner's Process.

R. Solution of Potass (1.150),
any quantity.

Add to it gradually, and at a slow heat,
Iodine, a sufficient quantity
to neutralize the alkali. Evaporate to dryness, calcine strongly, dissolve the residuum in water, and crystallize. (FEE; HENRI; MAGENDIE)

NIEMANN recommends potass to be saturated with hydriodic acid, and then evaporated to a thick liquid.

Caillot and Baup's Process.

R. Iodine, ten parts.
Distilled Water fifty parts.

Put them into a matrass, and add by small portions

Pure Iron Filings, five parts,
shaking the vessel each time. Heat till the liquor becomes colourless; filter it, and wash the filter with boiling water till this comes off insipid. Mix the liquor with the waters, and heat the whole in a sand-bath; when nearly at the boiling point, pour a sufficient quantity of solution of sub-carbonate of potass to precipitate all the oxide of iron. Filter, wash the filter thoroughly, mix all the liquors, test the mixture with turnsol, and if there is excess of base, add a little hydriodic acid. Evaporate in a sand-bath, till a slight pellicle forms; then remove the fire, and leave the liquor to crystallize: wash the crystals with a little water, and dry them. (HENRI)

Taddei's Process.

R. Iodine, any quantity.
Alcohol (25 degrees B.),
a sufficient quantity

to effect solution: add to it, at several times, hydrosulphate of potass, till it becomes of a milky whiteness; then set apart the liquor, and when it has become clear, add a few more drops of hydrosulphate; finally, filter,

evaporate to dryness, and crystallize. (HENRI)

This preparation only takes the name of hydriodate of potass when it is dissolved; when crystallized it is *Ioduret of Potassium*.

R. Hydriodic Acid, one part.
Water, four parts.

Add sub-carbonate of potass gradually to perfect saturation; then filter, evaporate, and crystallize. (FERR.)

SOLUTION OF HYDRIODATE OF POTASS.

Potio resolvens ex Iodio. (FERR.; BRERA; BORIES; FEE; HENRI; MAGENDIE; PIERQUIN; RATIER)

R. Hydriodate of Potass, thirty-six grains.
Distilled Water, one ounce.

Dissolve.

One drop of this solution weighs more than one grain. It is more generally preferred to be used than the tincture of iodine. It is prescribed in the same manner, in the same doses, and in the same cases.

STIMULANT POTION.

R. Hydriodate of Potass, two grains.
Sulphate of Magnesia, half an ounce.
Tartarized Antimony, half a grain.
Water, six ounces.

Mix.—Dose, a tea-spoonful, three or four times a day, in scrofula.

PILLS OF HYDRIODATE OF POTASS.
(PIERQUIN)

R. Hydriodate of Potass, five drams.
Distilled Water, six drams.
Biscuit, a sufficient quantity
to make a mass to be divided into three hundred pills, each of which contains one grain of the salt.

They have seemed to be useful in goître, leucorrhœa, and white swellings.

OINTMENT OF HYDRIODATE OF POTASS.
(FERR.; BRERA; FEE; HENRI; BORIES; ELLIS; MAGENDIE; NIEMANN; PIERQUIN; RATIER)

R. Hydriodate of Potass, half a dram.
Lard, one ounce and a half.

Mix with trituration.

The volume of a filbert to be rubbed upon goitrous tumours night and morning.

LOZENGES OF HYDRIODATE OF POTASS.

Tablettes au Moka. (PIERQUIN)

R. Sugar, four ounces.
Perphyrized Mocha Coffee, half a dram.

Hydriodate of Potass, seventy-five grains.
Mucilage of Tragacanth made with strong Infusion of Coffee, a sufficient quantity.

Make three hundred lozenges.—In leucorrhœa, goitre, amenorrhœa, tabes mesenterica, and scrofula.

IODURETTED HYDRIODATE OF POTASS.
(FERR. ; MAGENDIE)

R. Hydriodate of Potass, twenty parts.
Pure Iodine, six parts.

Mix in a glass mortar, and triturate for a long time, till the mixture becomes homogeneous, and of a deep red colour.

SOLUTION OF IODURETTED HYDRIODATE OF POTASS.

Solution de Coindet. (FERR. ; BRERA ; CADET DE GASSICOURT ; MAGENDIE)

R. Hydriodate of Potass, thirty-six grains.
Distilled Water, one ounce.
Dissolve, and add
Iodine, ten grains.

Mix.—One drop weighs from one grain and a half to two grains.

OINTMENT OF IODURETTED HYDRIODATE OF POTASS. (FERR. ; BORIES ; CADET DE GASSICOURT ; MAGENDIE ; PIERQUIN)

R. Ioduretted Hydriodate of Potass, half a dram.
Iodine, from ten to fifteen grains.
Lard, one ounce and a half.

Triturate well together. (FERR. ; BORIES ; CADET DE GASSICOURT ; MAGENDIE)

R. Hydriodate of Potass, one dram.
Iodine, twenty grains.
Calf's Gastric Juice, a sufficient quantity

to effect solution, in the space of twenty-four hours. Add

Lard, two ounces.

Mix. (FERR. ; BRERA ; PIERQUIN)

All these preparations are more active

than those in which only the hydriodate enters.

RESOLVENT POTION.

Mistura resolvens. (BRERA)

R. Rest Harrow, one ounce.
Water, a sufficient quantity

to obtain six ounces of strained decoction. Add to it

Solution of Ioduretted Hydriodate of Potass, forty drops.
Simple Syrup, one ounce.

Mix.—To be taken gradually in scrofulous diseases.

HYDRIODATE OF IRON.

The mode of preparing this salt is not given in any pharmacopœia.

ANTI-STRUMOUS SOLUTION. (PIERQUIN)

R. Hydriodate of Iron, four drams forty grains.
Distilled Water, four ounces.

Dose, ten drops, morning and evening, augmenting it progressively.

ANTI-SCROFULOUS OINTMENT, (PIERQUIN)

R. Hydriodate of Iron, sixty grains.
Lard, one ounce.

Mix.—It is said to have been used with success in leucorrhœa, white swellings, and glandular engorgements.

EMMENAGOGUE PASTILLES. (PIERQUIN)

R. Hydriodate of Iron, thirty grains.
Powdered Saffron, half a dram.
Sugar, four ounces.
Mucilage of Tragacanth, } of each
Infusion of Cinna- } a sufficient
mon, } quantity.

Make two hundred and forty pastilles.

They are extolled in amenorrhœa and leucorrhœa.—Dose, fifteen or twenty, daily, to be augmented progressively.

IPECACUANHA.

SYNONIMES :—*Ipecacuanha*, Fr.; Brech-
wurzel, GERM.; *Hipecacuana*, Bejuguillo,
SPAN.; *Ipekakuany*, POL.; Kræktrot,
SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.;
ANT.; AUSTR.; BATAV.; BAVAR.;
BELG.; BRUNS.; DAN.; MIL. COPEN.;
HISP.; GALL.; FERR.; FORM. HÔP. FR.;
FENN.; FULD.; GENEV.; HAMB.;
HANN.; HASS.; LIPP.; OLDEN.; POL.;
LUSIT.; CAST. BORUS.; BORUS.; ROSS.;
SAX.; SUEC.; WIRTEM.; HERBIP.;
WURCEB.; BERGIUS.; BRUGNATELLI;
COXE.; GUIBOURT.; MURRAY and GME-
LIN.; PARIS.; PIDERIT.; SPIELMANN.;
ZARDA.

The true *ipecacuanha* is the root of the
Cephalis Ipecacuanha, RICH., a plant of the
forests of Brazil. (*Pentandria Monogynia*,
LINN.; *Rubiaceæ*, JUSS.; *Fig. Dict. des*
Sc. Med. t. xxvi. pl. 1.)

This root (*Radix Brasiliensis*) presents
three varieties in commerce :

1. *Grey, ringed, blackish-grey, or brown Ipe-*
cacuanha : the diameter of this is not greater
than that of a writing quill. It presents ir-
regular rings, and is of a blackish-grey ex-
ternally; its ligneous axis is less consider-
able than its cortical portion; its fracture
is whitish and resinous; and its taste
bitter and rather acrid.

2. *Reddish-grey, red-grey, or grey Ipe-*
cacuanha. This has irregular rings, and
presents a resinous fracture. It is reddish
externally and slightly reddish-white inter-
nally, and of a more bitter taste than the
preceding.

3. *White-grey, white, or larger ringed*
Ipecacuanha : the rings of this sort are nearly
regular, and scarcely project; it is thicker
than either of the two preceding; its colour
is whitish-grey; its fracture whitish and re-
sinous, and its taste bitter.

The two former enter, the first in the pro-
portion of two-thirds, and the second in
that of one-third, into the *ipecacuanha* of
commerce, in which the third is rarely
found.

Many other roots also bear the name of
ipecacuanha : in America, according to DE-
CANDOLLE, this name is given to all roots
having emetic properties. In the number
are distinguished that of the *Psychotria eme-*
tica, MUT., a shrub of Peru (*Fig. Flore*
Medic. IV. 201), called *Black, striped, ash-*
grey, or liquorice Ipecacuanha. It is with-
out irregular rings, or has but few, but pre-
sents longitudinal striæ. Its fracture is
resinous, but less compact than that of the

preceding, and is of a very decided black-
grey, becoming still stronger when mois-
tened: it is without taste. It is not much es-
teemed, and is rarely met with in commerce.

There is also a *White or amylaceous Ipe-*
cacuanha, the root of the *Richardsonia Bra-*
siliensis, GOM. (*p. m.*), which presents ir-
regular semicircular rings. This root is of the
diameter of a pigeon's quill. It is wrinkled,
twisted, of a white-grey colour, without
taste, but exhales a mouldy smell. Its frac-
ture is white, amylaceous, and not at all
resinous.

This is never met with among the *ipecacu-*
anha of commerce.

The true *ipecacuanha* contains, according
to PELLETIER, a fat matter, gum, starch,
and particularly an alkaloid called *Emetine*,
to which it owes its properties. This al-
kaloid is less abundant in the root of *Psycho-*
tria than in that of the *Cephalis*, and there
is very little of it in that of the *Richard-*
sonia.

It is irritant, occasioning vomiting and
purging; but it acts only as a stimulant and
tonic when given in small doses.—Dose, as
a stimulant, from one grain to four grains;
as an emetic, from twelve to thirty grains, in
four ounces of water.

I. Preparations which contain *Ipecacuanha* in substance.

ANTI-DIARRHŒIC POWDER. (SAINTE-
MARIE)

R. *Ipecacuanha*, two grains.
Sugar, four drams.
Oil of Nutmeg, two drops.

Make a powder, and divide into four
doses, to be taken in twenty-four hours, for
fifteen or twenty days.—It is recommended
by FRANK, in chronic gonorrhœa.

EMETO-CATHARTIC POWDER. (ELLIS;
RATIER)

R. *Ipecacuanha*, twenty-four grains.
Rhubarb, twelve grains.

Mix. (RATIER)

ELLIS prescribes a scruple of *ipecacuanha*
and the same of rhubarb, to be taken to-
gether in syrup.

NAUSEANT POWDER.

Pulvis motum peristalticum invertens. (AU-
GUSTIN; BRERA)

R. *Ipecacuanha*, one fourth of a grain.
Sugar, fifteen grains.

To be taken every two hours.—It is recommended by RICHTER, in strangulated hernia. (AUGUSTIN)

R. Ipecacuanha, } one grain.
Gum Arabic, } of each,
Sugar, } one scruple.

Mix.—To be taken every two hours, in chronic catarrhs. (BRERA)

• EMETIC POWDER. (AUGUSTIN)

R. Ipecacuanha, } of each, one scruple.
Sugar, }

Divide into four parcels.—Dose, one every quarter of an hour till vomiting takes place.

R. Ipecacuanha, } of each, fifteen grains.
Sugar, }
Red Sulphuret of Antimony, } five grains.

Mix, and divide into four parcels.

R. Ipecacuanha, } of each, fifteen grains.
Sugar, }
Tartarized Antimony, } one grain.

Divide into four doses.

FEBRIFUGE POWDER. (SAINTE-MARIE)

R. Ipecacuanha, one scruple.
Subcarbonate of Magnesia, one ounce.

Divide into twenty-four parcels.—Dose, one every three hours in a cup of tea. It is recommended by WICHMANN.

EXPECTORANT POWDER. (ELLIS; PARIS; RATIER)

R. Ipecacuanha, twenty-four grains.
Squill, twelve grains.

Make a powder. (RATIER)

R. Ipecacuanha, six grains.
Myrrh, twelve grains.
Nitrate of Potass, half a dram.

Make four parcels.—Dose, one every quarter of an hour. (ELLIS; PARIS)

CARMINATIVE POWDER. (PIERQUIN)

R. Ipecacuanha, one grain.
Amber, twelve grains.

Mix.

DIGESTIVE BOLUSES. (CADET DE GASSICOURT; SWEDIAUR)

R. Ipecacuanha, from two to four grains.
Aromatic Powder, ten grains.
Syrup of Cinnamon, a sufficient quantity.

Mix.

ALEXITERIC POWDER. (FULD.)

R. Ipecacuanha, } of each,
Extract of Opium, } six grains.
Sulphate of Potass, } three drams.
Mix.—Dose, ten grains.

ANTI-DYSENTERIC BOLUSES. (CADET DE GASSICOURT; SAINTE-MARIE)

R. Ipecacuanha, two grains.
Theriac, one scruple.
Prepared Chalk, a sufficient quantity.

Mix.

ASTRINGENT BOLUSES. (PIERQUIN)

R. Ipecacuanha, one grain and a half.
Alum, six grains.
Diascordium, a sufficient quantity.

Make a bolus.

R. Ipecacuanha, one grain and a half.
SYDENHAM'S Laudanum, one grain.
Conserve of Hips, a sufficient quantity.

Make a bolus.

EMETIC BOLUSES. (RATIER)

R. Ipecacuanha, from eighteen to twenty-four grains.
Honey, a sufficient quantity.

Make three boluses; one to be taken every half hour.

NAUSEANT PILLS.

Pilule Nauseam cientes. (BRERA)

R. Ipecacuanha, six grains.
Rob of Elder, a sufficient quantity
to make six pills.—Dose, one every two hours.

EXPECTORANT PILLS. (ELLIS)

R. Extract of Hemlock, } of each,
Powdered Ipecacuanha, } ten grains.

Make four pills.—Dose, two.

R. Ipecacuanha, fifteen grains.
Soap, a sufficient quantity.

Make fifteen pills.—Dose, one four times a day.

PECTORAL MARMALADE. (BORIES)

R. Ipecacuanha, twelve grains.
Sublimed Sulphur, one scruple.
Florentine Iris, one dram.
Manna, } of each,
Syrup of Marshmallows, } two ounces.

Mix.—It is recommended in whooping cough.—Dose, a tea spoonful twice or thrice a day.

ANTI-DYSENTERIC ELECTUARY. (CADET DE GASSICOURT)

R. Ipecacuanha, half a dram.

Root of Tormentil,	one dram.
Opium,	four grains.
Conserve of Roses,	of each,
Syrup of Barberries,	six drams.
Mix.	

EMETIC ELECTUARY.

Electuarium Ipecacuanhæ moschatum. (AUGUSTIN)

R. Rhubarb,	} of each, one dram.
Borated Tartar,	
Ipecacuanha,	half a dram.
Musk,	four grains.
Red Sulphuret of Antimony,	ten grains.
Honey,	a sufficient quantity.

Mix.—Dose, a tea spoonful, repeated till slight vomiting comes on.—BAERENS recommends it in hooping cough.

TROCHISCS OF IPECACUANHA. (ANT.; BAVAR.; BELG.; HISP.; GALL.; FERR.; HANN.; SUEC.; BRUGNATELLI; CADET DE GASSICOURT; NIEMANN; VAN MONS)

R. Ipecacuanha,	one part.
White Sugar,	forty parts.
Mucilage of Tragacanth, made with	
Rose Water,	a sufficient quantity.

Make trochiscs of twelve grains each. (GALL.; FERR.; CADET DE GASSICOURT)

BRUGNATELLI prescribes one part of ipecacuanha, twelve parts of sugar, and mucilage of gum arabic to make ninety-four trochiscs.—HISP.—one of ipecacuanha, forty-four of sugar, three of tragacanth, and a sufficient quantity of water to make a hundred and forty-four trochiscs.—NIEMANN; and BAVAR.—one of ipecacuanha, sixty of sugar, and a sufficient quantity of mucilage of tragacanth to make a mass, one dram of which is to be divided into ten trochiscs.—ANT.; and BELG.—one of ipecacuanha, seventy-two of sugar, and a sufficient quantity of mucilage of tragacanth to make trochiscs of ten grains each.—SUEC.—one of ipecacuanha, ninety-six of sugar, and a sufficient quantity of mucilage of tragacanth to make sixty trochiscs.

R. Ipecacuanha,	thirty-six grains.
Mucilage of Gum Arabic,	
	a sufficient quantity

to make one hundred and forty-four trochiscs. (HANN.)

R. Ipecacuanha,	one part.
Sugar,	thirty-two parts.

Incorporate the powdered root triturate with an equal quantity of sugar, in the rest of the sugar boiled *à la grande plume*. (VAN MONS)

OPIATE TROCHISCS OF IPECACUANHA. (HENRI)

R. Ipecacuanha,	sixteen parts.
White Sugar,	six hundred and forty parts.
Extract of Opium,	five parts.
Gum Tragacanth,	eight parts.
Orange-flower Water,	a sufficient quantity.

Dissolve the opium in a little of the water, add it to the mucilage, and make trochiscs.

COMPOUND TROCHISCS OF IPECACUANHA. (NIEMANN)

R. Ipecacuanha,	one scruple.
Muriate of Ammonia,	half an ounce.
Powdered Chocolate,	one ounce and a half.
Gum Arabic,	two drams.
Syrup of Maidenhair,	a sufficient quantity.

Make eighty trochiscs.

EMETIC POTION. (FORM. HÔP. FR.; DISP. CHA. PARIS; BORIES; RATIER; SAINTE-MARIE; SWEDIAUR)

R. Ipecacuanha,	twelve grains.
Water,	four ounces.

Mix. (RATIER)

FORM. HÔP. FR.; and BORIES prescribe twenty grains of ipecacuanha, and four ounces of water.

R. Ipecacuanha,	twenty-four grains.
Water,	nine ounces.
Syrup of Maidenhair,	one ounce.

Mix. (GALL.)

DISP. CHA. PARIS prescribes ten grains of ipecacuanha, four ounces of water, and one ounce of simple syrup.

R. Ipecacuanha,	twenty-four grains.
Oxymel of Squill,	one dram.
Water,	one ounce.

Mix. (SWEDIAUR)

R. Ipecacuanha,	ten grains.
Proto-hydrosulphate of Antimony,	one grain.

Hordeated Cinnamon Water,	
Water,	four drams.
	six ounces.

Mix. (SAINTE-MARIE)

R. Ipecacuanha,	one scruple.
Tartarized Antimony,	one grain.
Syrup of Honey,	half an ounce.
Water,	four ounces.

Mix. (RATIER)

R. Ipecacuanha,	twenty grains.
Water,	four ounces.

Tartarized Antimony, two grains.
 Mix. (FORM. HÔP. FR.)
 To be taken at two or more doses, or even at once.

ANTI-SPASMODIC POTION. (HAMB.)

R. Ipecacuanha, three grains.
 Distilled Water of Valerian, } of each,
 ————— Pepper- } three
 mint, } ounces.
 Mix.

ANTI-VOMITIVE POTION. (SAINTE-MARIE)

R. Ipecacuanha, from two to four grains.
 Subcarbonate of Soda, ten grains.
 Syrup of Poppies, one ounce.
 Mint Water, six ounces.

Mix. — Against spasmodic vomiting.—
 Dose, a spoonful, every hour, or every two or three hours.

EMETIC MIXTURE. (NIEMANN)

R. Ipecacuanha, twenty grains.
 Oil of Almonds, one ounce.
 Mix.

II. *Preparations which contain the Active Principle of Ipecacuanha, without the vehicle employed to extract it.*

EXTRACT OF IPECACUANHA. (VAN MONS)

R. Powdered Bark of the Root of Ipecacuanha, any quantity.

Put it into a REAL's press, moistening it with cold water; at the end of twenty-four hours, charge the apparatus with a column of from twenty-two to twenty-five feet of boiling water, and receive the liquid so long as it flows coloured; strain it through flannel, and evaporate it to the consistence of an extract.

III. *Preparations which contain the Active Principle of Ipecacuanha, with the vehicle employed to extract it.*

A.

Extraction by Water.

INFUSION OF IPECACUANHA. (AMST.; FULD.; HAMB.; LIPP.; NIEMANN; VAN MONS)

R. Ipecacuanha, two drams.
 Boiling Water, six ounces.
 Infuse for a quarter of an hour, and strain. (FULD.; LIPP.)

R. Ipecacuanha, one dram and a half.
 Orange Peel, two drams.
 Water, five ounces.

Infuse for half an hour in a covered vessel. Add

Supertartrate of Potass, two drams.

Cover the vessel, and when cold, strain without expression. Add to four ounces of this liquid

Oxymel of Squill, half an ounce.

Mix. (AMST.; HANN.; NIEMANN; VAN MONS)

Dose, one or two spoonfuls, every quarter of an hour.

TONIC INFUSION. (BORIES)

R. Ipecacuanha, twenty-five grains.
 Orange Peel, one dram.
 Water, a sufficient quantity
 to obtain six ounces of strained infusion.
 Add to this

Syrup of Orange-flowers, one ounce.

Mix.—Dose, a spoonful, half an hour before each meal.

DECOCTION OF IPECACUANHA.

Decoctum Ipecacuanha. (SPIELMANN)

R. Ipecacuanha, coarsely powdered, two drams.
 Spring Water, four ounces.

Boil down slowly to one half, and strain; boil the residue twice again in the same manner, and mix the three strained decoctions.

It is recommended in dysentery.—To be taken at thrice, for three successive days.

AQUEOUS SYRUP OF IPECACUANHA. (GALL.; LIPP.; VAN MONS)

R. Ipecacuanha, coarsely powdered, half a pound.
 Water, seven pounds.

Boil in a covered vessel down to six pounds; set aside, clarify, filter, and add,

White Sugar, twelve pounds

Boil to the consistence of syrup. (GALL.)

R. Ipecacuanha, bruised, one part.
 Cold Water, a sufficient quantity
 to moisten it. At the end of twelve hours, add

Boiling Water, twenty parts.

When cold, strain, and add,

White Sugar, twenty-four parts.

Clarify, and evaporate to the consistence of syrup. (VAN MONS)

FEE proposes the following.

R. Ipecacuanha, one pound.
 Water, six pounds.

Macerate for twelve hours; pour an equal quantity of water upon the residue; mix and filter the two infusions, and make a syrup with

Sugar, twenty-four pounds.

R. Ipecacuanha, one pound.
Water, six pounds.

Boil for eight or ten minutes; strain, and add,

White Sugar, two pounds and a half.

Boil to the consistence of syrup; then add

Orange-flower Water, one ounce.

Mix. (LIPP.)

EXPECTORANT SYRUP. (FERR.; BORIES;
PIERQUIN; VAN MONS)

R. Ipecacuanha, two drams.
Florentine Iris, } of each,
Lichen Pyxidatus, } one ounce.
Sugar, two pounds.

Make a syrup. (VAN MONS)

BORIES; and PIERQUIN prescribe one ounce of iris, and half an ounce each, of lichen and of seneka.—They also quote the following syrup under the name of COURTI'S *Pectoral Syrup*.

R. Seneka, } of each, two ounces.
Iceland Moss, }
Red Cinchona, half an ounce.
Florentine Iris, two drams.
Ipecacuanha, one dram.
Sugar, two pounds.

Make a syrup.

R. Ipecacuanha, two ounces.
Cinchona, four ounces.
Opium, two scruples.
Water, seven pounds.

Macerate for two or three days, and decant; repeat the operation with fresh water so long as this comes off coloured; filter the liquors, and add to it,

White Sugar, nine pounds.

Evaporate in a water-bath, to the consistence of syrup. (FERR.)

EMETIC POTION. (FULD.)

R. Supertartrate of Potass, half a dram.
Boiling Water, four ounces.

Add to the hot strained liquor,

Powdered Ipecacuanha, two drams.
Orange Peel, one dram.

Filter after a short time.—To be taken by spoonfuls.

FEBRIFUGE POTION. (CADET DE GASSICOURT; SAINTE-MARIE)

R. Ipecacuanha, one dram and a half
Orange Peel, two drams.
Supertartrate of Potass, half an ounce.

Water, four ounces.

Boil and add to the strained liquor,

Oxymel of Squill, half an ounce.

Mix. (CADET DE GASSICOURT)

SAINTE-MARIE prescribes from half a dram to one dram of ipecacuanha, two drams of orange-peel, two drams of supertartrate, twelve ounces of water, to be reduced to eight, and two ounces of oxymel of squill.

To be taken by spoonfuls.

EXPECTORANT AND INCISIVE POTION.
(GALL.; BORIES; CADET DE GASSICOURT)

R. Ipecacuanha, one dram.
Follicles of Senna, two drams.
Boiling Water, six ounces.

After twelve hours' infusion, strain and add

Oxymel of Squill, } of each, one ounce.
Syrup of Hyssop, }

Mix.—For whooping cough.—Dose, six tea-spoonfuls to children; in the course of the morning.

EXPECTORANT AND TONIC MIXTURE.
(PIERQUIN)

R. Ipecacuanha, ten grains.
Orange Peel, } of each,
Iceland Moss, } two drams.
Water, six ounces.

Boil for half an hour, and infuse in the decoction,

Herb of Hyssop, } of each,
Root of Seneka, } one handful.

Strain, and add

Syrup of Borage, one ounce.

Mix.—To be taken by spoonfuls, in catarrhal affections.

B.

Extraction by Wine.

WINE OF IPECACUANHA.

Vinum seu Tinctura Ipecacuanhæ seu Ipecacuanne, Infusum Ipecacuanhæ vinosum, Vinum Psychotria seu Psychotriæ emetica. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BELG.; FENN.; LIPP.; LUSIT.; SUEC.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS)

Vinum Ipecacuanhæ.

R. Bruised Ipecacuanha, two ounces.
Proof Spirit, twelve fluid ounces.

Distilled Water,
twenty fluid ounces.

Macerate for fourteen days, and
filter. (LOND.)

R. Ipecacuanha, bruised, one part.
Spanish White Wine, fifteen parts.

Macerate for seven days, and filter.
(EDINB.)

R. Ipecacuanha, bruised,
two ounces.

Spanish White Wine, two pints.

Digest for seven days, and filter.
(DUBL.)

AMST.; BATAV.; BELG.; LUSIT.; BRUG-
NATELLI; and SWEDIAUR give the same
formula as EDINB.—AMER.; and COKE—
the same as DUBL.—FENN.; and SUEC.
prescribe six ounces of ipecacuanha and half
a pound of Spanish white wine.—VAN MONS
—one part of ipecacuanha and thirty-two
parts of Madeira wine.—LIPP.—six drams
of ipecacuanha and six ounces of Spanish
wine.

R. Bruised Ipecacuanha, one ounce.
White Wine, eight ounces.
Alcohol (20 degrees), two ounces.

Digest for three days in a sand-bath;
strain, and pour upon the residue,

White Wine, four ounces.
Alcohol (20 degrees), one ounce.

Digest for two days; strain, mix, and
filter the two liquors. (ANT.)

COMPOUND WINE OF IPECACUANHA.

Vinum emeticum. (HERBIP.; NIEMANN)

R. Ipecacuanha, two ounces.
Orange Peel, half an ounce.
Wine, two pounds and a half.

Infuse without heat for several days, and
strain. (HERBIP.)

R. Ipecacuanha, four ounces.
White Sugar, two ounces.
Alcohol (32 degrees), two pounds.

Digest in a sand-bath for eight days, then
add,

White Wine, eight pounds.
Indian Anise, one ounce.
White Sugar, four ounces.

Filter at the end of six days. (NIEMANN)

One ounce is equivalent to eighteen
grains of ipecacuanha.

COMPOUND SYRUP OF IPECACUANHA. (PIERQUIN)

R. Ipecacuanha, five drams, one scruple.
White Wine, one pound.

Infuse for a quarter of an hour, and add

Boiling Water, four pounds.
Subcarbonate of Potass, four ounces.
Wild Thyme, six drams.
Orange Peel, five drams one scruple.

Infuse for four hours; strain and add,

Syrup of Marshmallows, two pounds.
Orange-flower Water, twelve ounces.

Mix.

DESSESSART'S SYRUP. (BRERA; CADET
DE GASSICOURT; PIERQUIN; PLANCHE)

R. Ipecacuanha, one ounce.
Senna, three ounces.
White Wine, twenty-four ounces.

Macerate for two hours; then decant,
filter, and set apart the liquor. Add to the
residue,

Sulphate of Magnesia, three ounces.
Tops of Wild Thyme, one ounce.
Flowers of Red Poppies, four ounces.
Boiling Water, six pounds.

Infuse for four hours; then decant, fil-
ter, and add

Orange-flower Water,
twenty-four ounces.
White Sugar, fifteen pounds.

And the vinous tincture. Mix without
heat.

C.

Extraction by Alcohol.

TINCTURE OF IPECACUANHA. (VAN MONS)

R. Ipecacuanha, one part.
Proof Spirit, }
Madeira Wine, } of each, four parts.

Infuse without heat for several days; ex-
press, and filter.

JEROMEL has proposed the following
process:

R. Grey Ipecacuanha, in coarse powder,
four ounces.
Alcohol (37 degrees B.), one pound.

Digest in a stove for twenty-four hours,
at a temperature of 34 degrees, shaking the
vessel from time to time; then let it cool
and filter. Pour upon the residue

Alcohol (22 degrees), one pound.

Digest again in the same manner for
twenty-four hours, and filter. Pour again
upon the residue

River Water, filtered and made hot,
one pound.

Infuse for twenty-four hours; set apart
to cool, and then filter. Mix the three li-
quors, agitate the mixture, and set aside.
An abundant precipitate is formed, which
is to be separated at the expiration of twelve
hours. Distil the filtered liquor, and draw
off two pounds of alcohol. Filter the liquor

S

remaining in the water-bath, and add to it one ounce of alcohol at 39 degrees B.

In every two ounces of this tincture there are contained fifty-six grains of solid matter.

SYRUP OF IPECACUANHA.

Syrupus Psychotriæ emeticæ seu Ipecacuanhæ. (ANT.; BELG.)

R. Bruised Ipecacuanha, one ounce.
Alcohol (20 degrees),
nine ounces and a half.

Macerate the root for eight days in one half of the alcohol; then decant, and pour the remainder of it upon the residue. Digest in a sand-bath, and then mix the filtered tinctures together. Make a syrup by adding five drams of the mixture to one pound of warm simple syrup. (ANT.)

BELG. gives the same formula, but requires alcohol at 10 degrees.—JEROMEL has proposed to prepare the syrup by mixing two ounces of his tincture of ipecacuanha with one pound two ounces of simple syrup, boiling it for a moment to give it the proper consistence.

ANISATED TINCTURE OF IPECACUANHA.
(BORIES; CADET DE GASSICOURT; RATIER)

R. Ipecacuanha, in powder, one ounce.
Spirit of Anise, four ounces.

Digest. (CADET DE GASSICOURT)

RATIER prescribes one ounce of ipecacuanha and two ounces of spirit.—BORIES four ounces each, of spirit and of sugar, twelve grains of tartarized antimony, one ounce of ipecacuanha, twelve pounds of white wine, and fifteen days' infusion.

It is given as an emetic, principally to children.—Dose, one or two ounces.

WILLIS' SUDORIFIC ELIXIR. (CADET DE GASSICOURT)

R. Ipecacuanha,	}	of each,
Balsam of Tolu,		half an ounce.
Benzoic Acid,	}	of each, two drams.
Purified Opium,		
Saffron,		
Oil of Anise,		one dram.
Camphor,		two scruples.
Alcohol,		two pounds.

It is stimulant and sudorific.—Dose, from one to two drams.—Half an ounce contains two grains of opium.

IRIS.

Many species of this genus are quoted in pharmacopœias; we shall notice the following:

1. *Iris Germanica*, LINN.—Blue Flower-de-Luce.

SYNONIMES:—Flambe vraie, FR.; Deutsches Schwerdtel, Blaue Schwertel, GERM.; Kosatec, BOHEM.; Sverdlilie, DAN.; Lirio de Alemania, SPAN.; Blaawe Iris, DUT.; Giglio celeste azurro, ITAL.; Mieczyk ziele, POL.; Lirio roxo dos montes, PORT.; Sværas lilja, SWED.

AMST.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; PALAT.; WIRTEM.; HERBIP.; BERGIUS; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of Europe. (*Triandria Digynia*, LINN.; *Irideæ*, JUSS.; Fig. Zorn. *Ic. pl. t.* 188.)

The root (*Radix Ireos nostratis seu Gladioli cærulei*) is employed. It is long, cylindrical, articulated, thick, fleshy, wrinkled, brownish and fibrillous without, and white within. It has a disagreeable odour when fresh, and is feebly violaceous when dried.

Its taste in the fresh state is acrid and biting, but is bitter and nauseous after desiccation.

2. *Iris Pseudacorus*, LINN.—Yellow Iris.

SYNONIMES:—Flambe bâtarde, Faux acore, Acore bâtarde, FR.; Ackermann, Wasserschwerdwurzel, Gelbe Teichlilie, Falsche Acornswurz, Unächte Kalmuschwertel, GERM.; Sværdlilie, DAN.; Acore bastardo, SPAN.; Geel lisch, DUT.; Irida gialla, ITAL.; Mieczyk zolty, POL.; Sværdslilja, SWED.

BRUNS.; GALL.; WIRTEM.; BERGIUS; GUIBOUT; MURRAY and GMELIN; SAUNDERS; SPIELMANN.

A plant of Europe. (Fig. Zorn. *Ic. pl. t.* 187.)

The root (*Radix Acori palustris seu vulgaris seu Pseudacori seu Acori adulterini seu Ireos palustris seu Gladioli lutei*) is employed. It is long, slightly compressed, of about the thickness of the thumb, wrinkled and of a blackish-brown colour without, and spongy and reddish within. It has a

strong and disagreeable odour, and an acrid and styptic taste.

Both of these sorts are purgative.

3. *Iris fœtidissima*, LINN.—Stinking Gladwyn.

SYNONIMES:—*Iris fetide*, Glaycul puant, FR.; Waldcœusekraut, Stinkende Schwardffilie, GERM.

GALL.

A plant of Europe. (*Triandria Monogynia*, LINN.; *Irideæ*, JUSS.; *Fig.* Blackw. Herb. t. 158.)

The root (*Radix Xyridis seu Spathulæ fœtidæ*) is employed. It has a disagreeable odour, and acrid taste.

4. *Iris versicolor*, LINN.—Blue Flag.

SYNONIME.—*Iris variée*, FR.

AMER.; COXE.

A plant of North America. (*Fig.* Hort. Elth. tab. 155. *fig.* 187.)

The root is employed. It is fibrous.

5. *Iris Florentina*, LINN.—Florentine Iris, Orris.

SYNONIMES:—*Iris de Florence*, FR.; Florentinische Schwertel, Violenwurzel, Veilchenwurz, GERM.; Ussul, Ussosunul, Asman junie, Irsa, ARAB.; Trialowy koren, BOHEM.; Firolod, DAN.; Lirio de Florencia, SPAN.; Irsa, HIND.; Florentynse Iris, DUT.; Iride di Firenze, ITAL.; Korzen fiolkowy, POL.; Lirio de Florencia, PORT.; Firolot, SWED.

EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FULD.; GENEV.; HAMB.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PIDERIT.; SAUNDERS.; SPIEL-MANN.; ZARDA.

A plant of the middle of Europe. (*Fig.* Flore Medic. IV. 204.)

The root (*Radix Ireos seu Iridis Florentinæ*) is employed. It is thick, knotty, compact, brownish without, and white within. It has an acrid, bitter, and persistent taste, and a very pleasant odour similar to that of violets. It is usually met with in the market freed from the epidermis, in cylindrical, but rather flattened tuberculous pieces, whitish, and nearly insipid.

It contains, according to VOGEL, gum,

a brown extract, starch, a fixed oil, a volatile oil, solid and crystallizable, and woody fibre.

In the fresh state it is purgative, and is also esteemed incisive and errhine. It is sometimes prescribed in catarrhs.—Dose, one scruple of the powder; and from one to two ounces of the juice, in wine.

DIAIREOS SPECIES. (ANT.; BRUNS.; PALAT.; WIRTEM.; NIEMANN.; SPIEL-MANN.; VAN MONS)

R. Root of Florentine Iris, }
Sugar Candy, } of each,
Compound Powder of } six drams.
Tragacanth,

Make a powder. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

NIEMANN prescribes half an ounce of iris, and two drams each, of the powder and of sugar.—ANT.—two parts of iris, one part of tragacanth, and one of sugar.—HANN.—half an ounce of iris, and two drams each, of tragacanth and of sugar.

CITRINE POWDER.

Pulvis Puerorum citrinus. (BRUNS.)

R. Florentine Iris, eight ounces.
Liquorice, six drams.
Anise, } of each, three drams.
Fennel, }
Saffron, two drams and a half.
White Sugar,
an equal weight of the whole.

Make a powder.

SPECIES FOR PERFUME.

Species pro Odoramento. (SPIELMANN)

R. Fresh Petals of Red Roses, half an ounce.
— Flowers of Lavender, two drams and a half.
Rhodium Wood, half an ounce.
Root of Florentine Iris, two ounces.
Fresh Herb of Sweet Basil, } of each,
Mountain Ca- } two
lamint, } drams.
— Marjoram,
— Peppermint, four scruples.
— Origanum, }
— Pennyroyal, } of each,
— Sage, } two
— Savory, } drams.
Orange Peel, }
Lemon Peel, }
Cinnamon, }
Tops of Rosemary, } of each,
Cloves, } three drams.
Mace, }
Bay Leaves, half an ounce.

Musk, triturated with a little Sugar,
two grains.
Ambergris, triturated with Sugar,
four grains.

Make a coarse powder.

CEPHALIC SPECIES.

Species cephalica pro Cucuphis. (BRUNS.;
PALAT.; WIRTEM.; SPIELMANN)

R. Root of Round Cyperus,	}	of each,
—— Florentine Iris,		half
		an ounce.
Flowers of Lavender,	}	of each,
—— Rosemary,		half
—— Roses,		six drams.
Rhodium Wood,	}	of each,
Yellow Sanders Wood,		half
Aloe Wood,		an ounce.
Cloves,	}	of each,
Cinnamon,		half
Nutmegs,		three drams.
Yellow Amber,		
Storax,		

Cut, bruise, and mix. (BRUNS.; WIRTEM.)

R. Root of Florentine Iris,	three ounces.	
Herb of Lavender,	of each,	
—— Marjoram,	one ounce.	
Rhodium Wood,	of each,	
Yellow Sanders Wood,	six drams.	
Cinnamon,	two drams.	
Orange Peel,	half an ounce.	
Lemon Peel,	one ounce and a half.	
Cloves,	two drams and a half.	
Benzoin,	}	of each, half an ounce.
Mastic,		
Storax,		
Amber,		

Cut, bruise, and mix. (SPIELMANN)

R. Root of Florentine Iris,	one ounce and a half.	
Herb of Betony,	}	of each,
—— Marjoram,		half an ounce.
—— Rosemary,		
—— Sage,		two drams.
Flowers of Red Roses,		one ounce.
—— Lavender,		of each,
—— Clove Pink,		two drams.
Storax,	}	of each, three drams.
Amber,		
Indian Spikenard,	}	of each,
Yellow Sanders Wood,		two drams.
Rhodium Wood,	}	of each,
Aloe Wood,		four scruples.
Cloves,		
Round Cyperus,		
Cinnamon,		
Benzoin,		

Make a powder. (PALAT.)

EXTRACT OF FLORENTINE IRIS. (SARD.;
WIRTEM.)

R. Root of Florentine Iris, one part.
Boiling Water, six parts.

Infuse for three days, then boil a little, strain with expression, and evaporate the liquor to the consistence of an extract. (WIRTEM.)

SARD. directs the infusion to be reduced to two thirds before straining and expression.

FECULA OF IRIS.

Fecula Ireos. (GALL.; GENEV.; PALAT.)

R. Fresh Root of Iris, any quantity.

Having rasped and enclosed it in a bag, submit it to the press; add a little water to the juice, set it aside, decant the clear liquor, dry the fecula with a moderate heat, and pulverize it.

CONFECTION OF IRIS. (VAN MONS)

R. Root of Florentine Iris, one part.
White Sugar, three parts.

Triturate together.

MARMALADE OF IRIS.

Conditum Iridis saccharatum. (VAN
MONS)

R. Root of Florentine Iris, one pound.
Sub-carbonate of Potass, one ounce.
Squill, one dram.
Water, four pounds.

Macerate over a slow fire till soft; then strain through a sieve: pour fresh water upon the residue, macerate, and strain again. Repeat this operation till the whole has passed through the sieve. Then add

Simple Syrup, } of each,
Sugar, } three pounds.

Triturate to the consistence of thick jelly, and add

Pomated Extract of Iron, two ounces.

Mix.

It is tonic, and expectorant, and is recommended in chronic pulmonary catarrhs.—
Dose, four, five, or six tea-spoonfuls daily.

TROCHISCS OF IRIS.

Rotulae Diaireos, Tabellae Iridis Florentinae compositae, Trochisci bechici albi. (ANT.;
BRUNS.; WIRTEM.)

R. Diaireos Species, one ounce.
White Sugar, dissolved in
Violet Water, and boiled
à la plume, twelve ounces.

Make trochiscs. (BRUNS.; PALAT.)

- R. Root of Florentine Iris,
 one ounce and a half.
 Starch, two ounces and a half.
 Sugar, twenty ounces.
 Mucilage of Tragacanth,
 a sufficient quantity.

Make trochiscs. (ANT.)

PASTE OF FLORENTINE IRIS. (VAN MONS)

- R. Root of Florentine Iris, one part.
 Cold Water, sixteen parts.

Infuse for some hours, and strain; heat the liquor with

Gum Arabic, twenty parts.
 and when dissolved, add

Sugar Candy, fifteen parts.
 Clarify with white of egg, strain, evaporate to the consistence of jelly, and pour upon an oiled slab.

VIOLET WATER.

Aqua violacea. (SPIELMANN)

- R. Root of Florentine Iris, four ounces.
 Alcohol, thirty-two ounces.

Infuse for fifteen days.

It is used principally as a cosmetic.

OIL OF IRIS.

Oleum Iridis seu Ireos seu irinum. (BRUNS.; HISP.; PALAT.; SARD.)

- R. Root and Flowers of Iris,
 one part and a half.
 Olive Oil, three parts.

Digest for two days in a water-bath; express when cold, and repeat the maceration on fresh herb. (SARD.)

- R. Root and Flowers of } of each,
 Iris, } an equal part.
 Olive Oil, }

Boil slowly to the consumption of the humidity. (PALAT.)

- R. Fresh Root of Iris,
 one pound and a half.
 Olive Oil, three pounds.

Digest for twenty-four hours; then boil to the consumption of the humidity, strain with expression, pour the liquid upon half an ounce of flowers of iris, digest for twelve hours, boil to the consumption of the humidity, and strain. (HISP.)

- R. Juice of the Root and } of each,
 Flowers of Iris, } an equal part.
 Olive Oil, }

Boil to the consumption of the humidity, and express. (BRUNS.)

PECTORAL OINTMENT.

Unguentum Pectorale. (SARD.; WIRTEM.; SPIELMANN)

- R. Oil of Almonds, four ounces.
 — Chamomile, } of each,
 — Violet, } three ounces.
 Butter, } six ounces.
 Chicken's Fat, } of each,
 Duck's Fat, } two ounces.
 White Wax, three ounces.

Add to the melted mass,

Powdered Florentine Iris, two drams.
 — Saffron, half a dram.

Mix.—This was formerly recommended to children with cough and difficult respiration.

AROMATIC VINEGAR. (FERR.)

- R. Vinegar, six pounds.
 Root of Iris, } of each,
 Petals of Roses, } one ounce.
 Flowers of Clove-pink, }
 Cinnamon, }
 Nutmegs, } of each,
 Cloves, } two drams.
 Orange Peel, }
 Storax, }
 Benzoin, }
 Marjoram, }
 Mint, } of each,
 Thyme, } one ounce.
 Dittany of Crete, }

Digest for one month, and distil with a moderate heat.

ISATIS TINCTORIA, LINN.—WOAD.

SYNONIMES:—Pastel, FR.; Færberward, GERM.; Wede, DAN.; Pastel, SPAN.; and PORT.; Verfwede, DUT.; Guado, ITAL.; Linilo, POL.; Vejde, SWED.

BRUNS.; GALL.; BERGIUS; GUIBOURT.

A plant of Europe. (*Tetradynamia Siliculosa*, LINN.; *Cruciferae*, JUSS.; *Fig. Zorn. Ic. pl. t. 191.*)

The herb (*Herba Glasti seu Isatis*) is em-

ployed. It consists of glabrous, rather glaucous leaves, the radical ones being lanceolated and narrowed into petioles at the base, and caulinary being sessile, semi-amplexicaul, and prolonged into two appendices at the base. It has an acrid and bitterish taste.

A colouring principle is derived from it, resembling indigo, *Indigotine* or *Isatine*.

JASMINUM OFFICINALE, LINN.—JESSAMINE.

SYNONIMES:—Jasmin, FR.; Gelsomino, SPAN.

BRUNS.; HISP.; GALL.; FERR.; ROSS.; SAX.; WIRTEM.; BERGIUS.; BRUGNATELLI.; MURRAY and GMELIN.; SPIELMANN.

A shrub of the middle of Europe and of the east. (*Diandria Monogynia*, LINN.; *Jasmineæ*, JUSS.; *Fig. Zorn. Ic. pl. t. 153.*)

The flowers (*Flores Jasmini*) are employ-

ed. They are white, pedunculated, disposed in corymbs. When fresh, they exhale an agreeable odour; their taste is bitterish.

SPIRIT OF JESSAMINE. (SPIELMANN)

R. Oil of Jessamine, sixteen ounces.
Alcohol, eighteen ounces.

Mix with agitation; then expose, to the cold, to congeal the oil, and pour off the spirit.

JATROPHA CURCAS, LINN.—ANGULAR-LEAVED PHYSIC-NUT.

SYNONIMES:—Pignon des Barbades, Medicinier, Grand Haricot du Pérou, FR.; Grosser Wunderbaum, Barbados-Nussbaum, GERM.; Mara harulu, CAN.; Rata endaru, CYNG.; Bagbarenda, HIND.; Dscharrak pagger, JAV.; Caat amunak, TAM.; Nepala, Adivze amida, TEL.

BRUNS.; GALL.; GUIBOURT.; MURRAY and GMELIN.

A shrub of Brazil, Surinam, and Jamaica. (*Monœcia Monadelphica*, LINN.; *Euphorbiaceæ*, JUSS.; *Fig. Desc. Flore Ant. II. 141.*)

The seeds (*Semines Ricini majoris, Nux cathartica Americana, Nux Barbadiensis*) are

employed. They are oval, somewhat compressed, of the size of a filbert, rugose, and blackish. Their taste is acrid and burning. They are purgative.

OIL OF PHYSIC NUT.

Oleum Ricini Americani. (VAN MONS)

R. Seeds of *Jatropha Curcas*, any quantity.

Bruise them in a mortar, enclose the paste in a linen bag, and express.

It is drastic and emetic; it is recommended in ileus, worms, and obstinate piles. —Dose, from half an ounce to two ounces.

JUGLANS.—WALNUT TREE.

Two species of this genus are employed in medicine:

1. *Juglans cinerea*.—Butter-nut or White Walnut Tree.

AMER.; COXE.

A tree of South America. (*Monœcia Polyandria*, LINN.; *Juglandææ*, CAND.; *Fig. Big. Med. Bot. II. 32.*)

The internal bark of the trunk and of the root is employed; it is purgative. The internal bark of the root is said, when fresh, to be vesicant.

2. *Juglans regia*, LINN.—Walnut Tree.

SYNONIMES:—Noyer commun, FR.; Nussbaum, Wallnus, GERM.; Akirut, Jowz,

Khusif, ARAB.; Walsky arech, BOHEM.; Ho-tao, CHIN.; Valnodd, DAN.; Aktrut, DUK.; and Hind., Nogal, SPAN.; Noce, ITAL.; Olass-dio, MAL.; Girdigan, Charminghz, Jouzirumic, PERS.; Orzeszina wloska, POL.; Nogueira, PORT.; Greziak orechi, RUSS.; Walnø Hrød, SWED.

ANT.; AUSTR.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

This tree is originally of Persia, but is cultivated in every part of Europe (*Mo-*

nœcia Polyandria, LINN.; *Terebinthaceæ*, JUSS.; *Fig. Flore Medic. V. 250.*)

The fruit (*Nux Juglans* seu *Juglandis*) is employed. It is an oval drupe. Its exterior envelope (*Cortex* seu *Putamen Nucis Juglandis*) is green, fleshy, and of a bitter and extremely acerb taste. The nut which it covers (*Nucleus Nucis Juglandis*) consists of a hard, shining, woody, yellow grooved shell, inodorous and insipid, containing a four-lobed oleaginous kernel, very irregular in its surface, covered with a yellowish, thin, astringent pellicle (*Epidermis Nuclei Juglandis*). It has an agreeable taste.

ANTI-HYDROPHOBIC ELECTUARY. (NIEMANN)

R. Ripe Walnuts, thirty.
Herb of Rue, one handful.
Honey, eight ounces.
Bruise together.

It is extolled by STRACHWITZ.—A spoonful to be taken morning and evening, before meals.

OIL OF WALNUTS.

Oleum Nucum juglandium seu *Juglandis regiae*. (ANT.; BRUNS.; HISP.; GALL.; HASS.; LIPP.; PALAT.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; VAN MONS)

R. Ripe Walnuts, any quantity.
Remove the shells and partitions; then having bruised them in a stone mortar with a wooden pestle, enclose the paste in a linen bag, and express it between two tin plates without heat.

When fresh, this oil may be substituted for that of almonds.

ROB OF WALNUTS.

Rob Nucum seu *Corticis Nucum Juglandis regiae*, *Rob Diacarii Galeni*, *Diacaryon*. (AUSTR.; BRUNS.; FULD.; PALAT.; WIRTEM.; HERBIP.; NIEMANN; SPIELMANN)

R. Green Walnuts, one pound.
After having bruised them in a mortar, boil them for several hours with two pounds of water; express, and evaporate the liquor to the proper consistence, with an equal weight of despumated honey. (BRUNS.; FULD.; WIRTEM.; SPIELMANN; SWEDIAUR)

R. Expressed Juice of Green Walnuts, two pounds.
Honey, one pound.
Boil to a proper consistence over a slow fire, skimming. (AUSTR.; PALAT.; HERBIP.; NIEMANN; SPIELMANN)

It is slightly astringent and resolvent, but is principally designed for external use.

EXTRACT OF GREEN WALNUTS.

Extractum Nucum seu *Juglandis* seu *Nucum juglandinum viridum*. (AMER.; AUSTR.; BAVAR.; BRUNS.; DAN.; FENN.; HANN.; HASS.; LIPP.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; COXE; NIEMANN; VAN MONS)

R. Walnuts gathered before maturity, any quantity.

Bruise them in a stone mortar, adding a little water; express the juice, and immediately evaporate it in a water-bath, constantly stirring towards the end with a spatula. (AUSTR.; BAVAR.; BRUNS.; DAN.; FENN.; HANN.; POL.; BORUS.; SAX.; SUEC.; NIEMANN; VAN MONS)

R. Walnut Rinds, one part.
Pure Water, six parts.

Boil together; and evaporate the strained liquor. (AMER.; HASS.; LIPP.; ROSS.; WIRTEM.; COXE; PIDERIT)

It is excitant, tonic, bitter, anti-scorbutic, and anthelmintic.—Dose, from half a scruple to half a dram.—It used formerly to be applied to cleanse foul ulcers, and also to syphilitic ulcers.

The extract of walnuts forms the basis of *Mittie's Anti-syphilitic Remedy*, of which ALYON and CADET DE GASSICOURT have given the two following different formulæ:

1. Alyon's formula.

R. Walnut Leaves, } of each, an
Leaves of Smallage, } equal weight.
Buck-bean, }

Bruise together, express the juice, and evaporate it to the consistence of pilular mass.—Dose, from twelve to thirty-six grains daily.

2. Cadet de Gassicourt's formula.

R. Walnut Leaves, one ounce and a half.
Jalap, one ounce.
Leaves of Angelica, } of each,
Blessed Thistle, } four
Soap-wort, } ounces.

Make an infusion; evaporate, and divide the extract into pills of six grains each.

CADET DE GASSICOURT also prescribes the following syrup:

R. Walnut Leaves, twenty pounds.
Infuse them for twenty-four hours in hot water; pour the infusion upon

Leaves of Angelica, sixteen pounds.

Infuse again for twenty-four hours; keep the infusion hot in a water-bath, and after a time pour it upon

Leaves of Soap-wort, sixteen pounds.
 Musk Roses, twelve ounces.
 Common Roses, six ounces.
 Senna, twenty-six ounces.

Leave to infuse for twenty-four hours;
 then dissolve in the infusion

Sugar, thirty-seven pounds.

Boil, clarify with white of egg, and reduce the liquor to the consistence of syrup.

MITTIE directed the patient to take eight pills in the morning, two hours before breakfast; and four, two hours before supper; he also directed him to take four spoonfuls of syrup in a glass of water in the morning, fasting, and two spoonfuls, two hours before supper.

ANTHELMINTIC DROPS.

Mixtura Nucum Juglandium. (HAMB.; AUGUSTIN)

R. Extract of Walnuts, two drams.
 Cinnamon Water, one ounce.

Mix.—Dose, sixty drops to a child of two years, and one hundred to an adult.

TINCTURE OF WALNUT RINDS.

Essentia Corticis Viridis Nucum Juglandis. (DAN.)

R. Walnut Rinds, fresh and sliced, six ounces.
 French Brandy, two pounds.
 Digest in a moderate heat for four days, and filter.

It is bitter, astringent, tonic, and stimulant.

DECOCTION OF WALNUT RINDS.

Decoctionum Drupæ Juglandis regię seu Corticis viridis Nucum juglandium. (GENEV.; AUGUSTIN; NIEMANN)

R. Green Walnut Rinds, one ounce.
 Boiling Water, ten ounces.

Boil for a quarter of an hour, and strain. (GENEV.; AUGUSTIN; NIEMANN)

R. Walnuts, gathered a short time before their maturity, and dried, three ounces.
 Spring Water, four ounces.

Boil down to one half, and add to the strained liquor

Honey, two ounces.

Dissolve. (GENEV.)

This drink is extolled in syphilitic disorders which do not yield to mercury, and in those which attack very irritable subjects. FRANK recommends it in lotions to herpetic ulcers.

COMPOUND INFUSION OF WALNUT RINDS.

Infusum Juglandis compositum. (VAN MONS)

R. Dried Walnut Rinds, } of each,
 Walnut Leaves, } four handfuls.
 Water, a sufficient quantity

to obtain eight pounds of strained infusion. Having let it stand for an hour, add

Muriate of Lime, one ounce and a half.

Mix.—It is astringent, and is employed externally in the form of baths, at the temperature of from 36 to 38 degrees R., in scrofula, tubercles of the lungs, and generally in all diseases of the lymphatic system.

POLLINI'S DECOCTION.

Decoctionum Drupæ Juglandis regię. (AUGUSTIN; FEE; NIEMANN; RATIER; SAUNDERS; SWEDIAUR; PH. SYPH.)

R. Walnut Rinds, ten ounces.
 Sarsaparilla, } of each,
 China Root, } half
 Sulphuret of Antimony, } an ounce.
 Pumice Stone, }
 Water, sixty-four ounces.

Boil down to one half in a lightly-covered vessel. (AUGUSTIN; FEE; NIEMANN)

R. Sarsaparilla, } of each, two ounces.
 China Root, }
 Dried Walnut Rinds, one pound.
 Sulphuret of Antimony, } of each,
 Pumice Stone, } two ounces.
 Water, twenty pounds.

Boil down to one half. (NIEMANN)

R. Sarsaparilla, three ounces.
 China Root, one ounce and a half.
 Sulphuret of Antimony, two ounces.
 Entire Walnuts, fifty.
 Water, ten pounds.

Boil. (SAUNDERS)

R. Walnut Rinds, eighteen ounces.
 Sarsaparilla, } of each,
 China Root, } half
 Sulphuret of Antimony, in } an ounce.
 a bag, }
 Pumice Stone, in a bag, }
 Water, eight pounds.

Macerate for twelve hours; then having boiled them in a covered vessel, take out the bags, and continue the boiling till only four pounds of liquid remain; then strain without filtering. (PH. SYPH.)

PH. SYPH. gives also under the name of *Decoctionum Syphiliticum roborans*, another formula,—four ounces of sulphuret of antimony, two each, of pumice stone, of sarsaparilla, and of china root, forty dried unripe walnuts, and twenty pounds of water, boiled till reduced to ten.

- R. Dried Walnut Rinds, one ounce.
 Sarsaparilla, } of each, two ounces.
 China Root, }
 Sulphuret of Antimony, four ounces.
 Pumice Stone, two ounces.
 Water, twenty pounds.

Boil down to one half. (FEE; SWEDIAUR)

- R. Grey Cinchona, } of each,
 Sarsaparilla, } three ounces.
 Pumice Stone, } of each,
 Sulphuret of Antimony, } one ounce
 and a half.

- Walnut Rinds, one pound.
 Water, eight pounds.

Boil down to two pounds, and add when strained

- Subcarbonate of Potass, two drams.

Dissolve. (RATIER)

It is stimulant and tonic, and is recommended in leprosy, tetters, and obstinate syphilitic disorders. — Dose, one pound morning and evening. It is also employed externally, in lotions.

JUNIPERUS.

Three species of this genus are employed in medicine:

1. *Juniperus communis*, LINN.—Juniper.

SYNONIMES:—Genevrièr commun, FR.; Wachholder, Kaddig, GERM.; Galowec, BOHEM.; Ene, Eneboer, DAN.; Enebro, SPAN.; Geneser, DUT.; Genepro, ITAL.; Jalowice, POL.; Zimbrow, PORT.; En, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.: ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

A tree or shrub of Europe. (*Diœcia Monadelphica*, LINN.; *Conifera*, JUSS.; *Fig. Florè Medic. IV. 181*)

The wood, tops, and fruit are employed.

The wood (*Lignum Juniperi*, *Lignum cedrinum*) is solid, heavy, and very resinous; the bark is grey or reddish-brown without, and yellowish or reddish-yellow within. It has an agreeable and balsamic odour, and an aromatic, resinous, and bitterish taste.

The tops, or extremities of the branches, are slender, with sessile leaves, usually grouped in threes, narrow, hard, and pointed.

The fruit consists of spherical berries, blackish, of a resinous odour, and balsamic, bitterish taste, accompanied with a degree of sweetness.

Every part of the plant, but particularly the fruit, is considered powerfully diuretic. To the wood also are attributed anti-syphilitic properties. — Dose, from two to six drams, of the powder.

lentic properties. — Dose, from two to six drams, of the powder.

CONSERVE OF JUNIPER. (SWEDIAUR)

- R. Fresh Juniper Berries, eight ounces.
 Bruise them, adding gradually

- White Sugar, four ounces.

Heat slightly, that the sugar may be dissolved.

DIURETIC POWDER. (AUGUSTIN; NIEMANN)

- R. Grated Juniper Berries, } of each,
 Root of Burnet Saxifrage, } five grains.
 Antimoniated Nitre, three grains.

Mix.—It is recommended by THEDEN. (NIEMANN)

- R. Grated Juniper, } of each,
 White Sugar, } an equal part.

Mix.—Dose, a tea spoonful, every two or three hours. (AUGUSTIN)

EXTRACT OF JUNIPER.

Extractum Juniperi. (ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; WIRTEM.; HERBIP.; PIDERIT; SWEDIAUR; VAN MONS)

- R. Herb of Juniper, one part.
 Warm Water, four parts.

Infuse for forty-eight hours, occasionally stirring; then strain, set aside, and decant the liquor: finally evaporate it slowly to the consistence of an extract. (BAVAR.; GALL.; FORM. HÔP. FR.)

- R. Juniper, any quantity.
 Bruise it, moistening it with a little warm T

water and adding a little yeast; then let it ferment at a temperature of from 15 to 20 degrees, R., and strain with slight expression: reduce the liquor to one half by boiling, clarify with white of egg, and evaporate to the proper consistence. (VAN MONS)

R. Residue of the distillation of Spirit of Juniper, any quantity.

Clarify it, and evaporate to the proper consistence. (SWEDIAUR)

R. Bruised Juniper, eight pounds.
Hot Water, twelve pounds.

Digest for three days; then distil in an alembic till essential oil ceases to pass over with the water; set the oil apart, and express the residue strongly: evaporate the liquor to the consistence of an extract, and mix the oil intimately with it. (SARD.)

R. Bruised Juniper, four pounds.
Boiling Water, twelve pounds.

Digest for two or three days, and then boil till the berries become soft. Express, set aside the liquor, and evaporate it in a water-bath to the consistence of an extract. (HISP.)

GENEV.; HANN.; LIPP.; OLDEN.; BORUS.; and SAX. direct the berries to be softened by boiling them in water, then to be gently expressed, and the liquor to be evaporated to the consistence of honey.—POL.—to macerate them in water for two days without boiling, and to evaporate the expressed juice to the same degree.—FENN. to repeat the maceration with the same water till the berries become nearly tasteless, and to evaporate the liquor, without previous boiling, to the consistence of honey.

R. Juniper, one pound.
Water, four pounds.

Infuse in a water-bath; and to the strained infusion, depurated by repose, add

White Sugar, one pound.

Evaporate slowly to the consistence of honey. (ANT.)

VAN MONS directs to be macerated, for twelve hours, twelve parts of bruised juniper in forty-eight parts of water; then to strain the liquor through a sieve without expression, and to macerate the residue in twenty-four parts of water: to strain again; to mix the two infusions; and to add one part of white sugar; finally to clarify with white of egg, and to evaporate to the consistence of honey.—FERR.—to macerate, for two days, six pounds of juniper in twenty-four of water, and to repeat the maceration with fresh water till this comes off insipid; the mixed infusions to be strained, and two pounds of sugar added; the solu-

tion to be clarified with white of egg, and evaporated to the proper consistence.

R. Juniper, any quantity.

Reduce to a pulp with a sufficient quantity of water, and then boil in a sufficient quantity of water for half an hour; strain through a sieve and express the residue: dissolve one part of sugar in four parts of liquor, and evaporate to the consistence of honey. (AUSTR.; BATAV.; BELG.; DAN.; HAMB.; PALAT.; HERBIP.)

BRUNS.; FULD.; HANN.; ROSS.; WIRTEM.; PIDERIT; and SWEDIAUR prescribe one pound of sugar to eight pounds of berries submitted to boiling.

This extract when only reduced to the consistence of honey, and particularly when sweetened, takes the name of Rob of Juniper; *Rob Juniperi seu Baccarum Juniperi, Succus Juniperi inspissatus.*

It is stimulant, aromatic, carminative, stomachic, diaphoretic, and diuretic.—Dose, from half an ounce to one ounce, in water.

DIURETIC ELECTUARY.

Electuarium de Junipero. (SARD.)

R. Rob of Juniper,	}	of each,
Despumated Honey,		one pound.
Juniper,	}	four ounces.
Cinnamon,		of each,
Dried Orange Peel,	}	half an ounce.

Melt the honey and rob together, with a gentle heat, and add the other substances in powder.

SOLUTION OF ROB OF JUNIPER.

Solutio Rob Juniperi. (NIEMANN)

R. Rob of Juniper, four ounces.
Juniper Water, two pounds.

Mix intimately, and add

Tincture of Juniper, two ounces.
Nitrous Æther, half an ounce.

Mix.—It is excitant, and is extolled by VAN SWIETEN in ascites.—Dose, one or two ounces, every three hours.

ELIXIR OF JUNIPER.

Elixir Juniperinum seu Malvaticum Juniperinum. (SPIELMANN)

R. Rob of Juniper, four ounces.
Madeira Wine, twelve ounces.

Mix well, and strain through linen.

It is excitant and is classed among the carminatives and diuretics.—Dose, one spoonful.

HYDRAGOGUE POTION.

Potio Juniperina. (AUGUSTIN; CADET DE GASSICOURT; NIEMANN; SAUNDERS)

R. Rob of Juniper, four ounces.
Juniper Water, two pounds.

Mix, and add

Spirit of Juniper, two ounces.

Mix. (AUGUSTIN; CADET DE GASSICOURT; NIEMANN)

R. Rob of Juniper, one ounce.
Juniper Water, six ounces.
Spirit of Juniper, half an ounce.
Spirit of Nitric Æther, half a dram.

Mix. (SAUNDERS)

Dose, two ounces, every two or three hours.

OIL OF JUNIPER.

Oleum Baccarum Juniperi æthereum, Ætheroleum Baccarum Juniperi. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FULD.; GENEV.; HANN.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; SWEDIAUR)

Oleum Juniperi. (LOND.)

Oleum Volatile Juniperi communis. (EDINB.)

Oleum Baccarum Juniperi. (DUBL.)

R. Juniper Berries, bruised, any quantity.

Water, a sufficient quantity to prevent empyreuma. Distil, and separate the oil.

The same is required by AMER.; AMST.; BATAV.; BELG.; FENN.; FERR.; GENEV.; LUSIT.; ROSS.; SUEC.; and COXE.

BAVAR. prescribes one part of juniper and three parts of water.—FERR.; FULD.; and SWEDIAUR—one of juniper and four of water.—DAN.; and VAN MONS—one of juniper and six of water.—HANN.; OLDEN.; POL.; BRUNS.; and SAX.—one of juniper and eight of water.—GALL.—five of juniper and seven of water.—ANT.; and BRUGNATELLI—three of juniper and ten of water.—HISP.—one of juniper and forty of water.

R. Juniper Berries, four parts.
Water, sixteen parts.
Muriate of Soda, one part.

Macerate for three days, and distil.—(BRUNS.; PALAT.; SARD.; WIRTEM.)

R. Juniper Berries, any quantity.

Juniper Water, a sufficient quantity.

Macerate for twelve hours, and distil. (HERBIP.)

Dose, from two to ten drops, in an appropriate vehicle.

OIL OF JUNIPER WOOD.

Oleum Ligni Juniperini æthereum, Ætheroleum Juniperini Ligni. (AUSTR.; HANN.; PALAT.)

R. Juniper Wood, one part.
Water, eight parts.

Distil. (HANN.)

PALAT. prescribes one part of wood and six parts of water.—AUSTR.—one of wood and twelve of water.

SARD. speaks of another oil obtained by distilling the rasped wood at the open fire, and separating from the fetid oil the clear liquor which passes over with it.

OLEO-SACCHARUM OF JUNIPER. (CAST. BORUS.)

R. Essential Oil of Juniper, twelve drops.
Sugar, half an ounce.

Triturate together

STIMULANT POWDER. (AUGUSTIN)

R. White Sugar, half an ounce.
Essential Oil of Juniper, three drops.

Mix intimately, and add

Muriate of Soda, one dram.

Divide into twelve parcels.—Dose, three daily; in white swellings of the joints.

JUNIPER WATER.

Aqua Juniperi. (AUSTR.; BAVAR.; GALL.; HASS.; LIPP.; CAST. BORUS.; SAX.; HERBIP.; BRUGNATELLI; NIEMANN; PIDERIT; SWEDIAUR; VAN MONS)

R. Juniper, two parts.
Water, fifteen parts.

Distil four parts. (GALL.)

BAVAR. and NIEMANN prescribe one of juniper and five of water; two to be distilled.—HERBIP.—one of juniper and twelve of water.—AUSTR.—one of juniper and ten of water; six to be distilled.—BRUGNATELLI—one of juniper and twelve of water; four to be distilled.—HASS.; LIPP.; and PIDERIT—the same, but six to be distilled.—SAX.—one of juniper and twenty-two of water; fifteen to be distilled.—SWEDIAUR—one part of juniper and a sufficient quantity of water; twenty parts to be distilled.

R. Fresh Leaves of Juniper, one part.
Proof Spirit, one half part.
Water, one hundred and sixty parts

Distil forty parts, draw off the oil, and preserve the water. (VAN MONS)

R. Oleo-saccharum of Juniper, one dram.

Distilled Water, eight ounces.

Dissolve. (CAST. BORUS.)

SPIRIT OF JUNIPER.

Spiritus Juniperi. (ANT.; BRUNS.; FULD.; HANN.; HASS.; LIPP.; PALAT.; WIRTEM.; HERBIP.; PIDERIT; VAN MONS)

R. Juniper, one part.
Alcohol, twelve parts.

Macerate for twenty-four hours, and distil. (FULD.; HASS.; PIDERIT)

ANT.; and LIPP. prescribe one part of juniper and twelve parts of alcohol.—BRUNS.; PALAT.; WIRTEM.; and HERBIP.—one of juniper and two of alcohol.

R. Juniper, one part.
Alcohol, four parts.
Water, a sufficient quantity.

Distil at the end of twenty-four hours. (HANN.)

R. Juniper, fifteen pounds.
Sugar, half a pound.
Yeast, two ounces.

Add a sufficient quantity of warm water to mix the whole well together, expose it to the temperature of from + 15 to + 18 degrees R.; after the fermentation, pour it into an alembic: add twice its quantity of water, and eight pounds of spirit from grain. Distil a spirit marking from 18 to 20 degrees.

COMPOUND SPIRIT OF JUNIPER.

Spiritus Juniperi compositus, Alcohol Juniperi compositum. (LOND.; EDINB.; DUBL.; AMER.; BATAV.; BELG.; SUEC.; COXE; SWEDIAUR; VAN MONS)

R. Juniper Berries, bruised, one pound.

Caraway Seeds, } of each,
bruised, } one ounce
Fennel Seeds, } and a half.
bruised, }

Proof Spirit, one gallon.

Water, a sufficient quantity

to prevent empyreuma. Macerate for twenty-four hours, and distil one gallon with a moderate heat. (LOND.)

DUBL. gives the same formula, but requires the maceration to be continued for two days.

EDINB. requires nine pounds of

alcohol and the same quantity of the other ingredients as LOND.; and nine pounds to be distilled after two days maceration.

AMER.; BATAV.; BELG.; SUEC.; COXE; and SWEDIAUR give the same formula.

R. Juniper Berries, six parts.
Fresh Leaves of Black Currant,

Seeds of Wild Carrot, } one part.
Black Currant, } of each,
two parts.

Spirits from Grain, thirty-six parts.
Water, one hundred and eight parts.

Distil twenty-four parts. (VAN MONS)

It is carminative and diuretic.—Dose, from two to four drams.

LINIMENT OF JUNIPER.

Linimentum Juniperinum. (AUGUSTIN)

R. Spirit of Juniper, two ounces.
Oil of Cloves, } of each,
Expressed oil of Nutmeg, } half a dram.

It is extolled by ROSENSTEIN in ecchymoses and cold tumours.—CHRESTIEN employs it in frictions to the spine and loins in women disposed to abortion.

DIURETIC LINIMENT.

Linimentum resolvens et diureticum.
(BRERA)

R. Spirit of Soap, three ounces.
Juniper, one ounce.

Mix.—To be rubbed upon the abdomen, morning and evening, in ascites with debility.

HYDRAGOGUE TINCTURE. (BORIES)

R. Root of Round Birthwort, } of each,
Long Birthwort, } one ounce
Ginger, } and a half.
Horse-radish, } of each,
Dried Squill, } half an ounce.
Tops of Wormwood, } of each,
Lesser Centaury, } one ounce.
Juniper, }
Subcarbonate of Potass, } half an ounce.
Spirit of Juniper, six pounds.

Macerate for eight days, and filter.

It is excitant and diuretic. BOERHAAVE recommended it in dropsy.—Dose, one ounce, four times a day.

INFUSION OF JUNIPER.

Infusum Baccarum Juniperi. (CAST. BORUS.; WURCEB.; AUGUSTIN; NIEMANN; RATIER)

R. Bruised Juniper Berries, one ounce and a half.
Boiling Water, one pound and a half.

Digest in a covered vessel for half an hour, and strain. (CAST. BORUS.; WURCEB.; AUGUSTIN)

RATIER prescribes two drams of juniper and two pounds of water.—NIEMANN—one ounce of juniper and a sufficient quantity of water to obtain three ounces of strained infusion.

It is aromatic and slightly diuretic; and is recommended in the treatment of dropsies.—Dose, a glassful.

COMPOUND INFUSION OF JUNIPER.
(SAUNDERS)

R. Bruised Juniper Berries, two ounces.
Juniper Water, two pounds.

Digest for three hours in a moderate temperature; then strain, and add

Rob of Juniper, two ounces.
Oxymel of Squill, one ounce and a half.

Mix.—It is diuretic.

DECOCTION OF JUNIPER LEAVES.

Decoctum Juniperi. (SAUNDERS; SWEDIAUR)

R. Tops of Juniper, three ounces.
Water, four pounds.

Boil down to two pounds, and add towards the end,

Juniper, one ounce.

Strain.—It is stimulant and diuretic; and is recommended in dropsy, asthma, and diseases of the skin.—Dose, from four to eight ounces, thrice a day.

DECOCTION OF JUNIPER WOOD.

Decoctum Ligni Juniperi. (NIEMANN)

R. Shavings of Juniper Wood, half an ounce.
Spring Water, sixteen ounces.

Boil down to one half, and strain.

This used formerly to be recommended in the treatment of syphilitic diseases: the eight ounces were taken in the course of the day.

DIURETIC PTISAN.

Potus diureticus. (SAINTE-MARIE; SWEDIAUR)

R. Bruised Juniper Berries, one handful.
Water, one pound and a half.

Boil for half an hour, and add when strained,

Syrup of Seneka, two ounces.
White Wine, eight ounces.

Mix. (SAINTE-MARIE)

R. Juniper, } of each,
Mustard Seeds, } one ounce.

Carrot Seeds, six drams.
Leaves of Wormwood, three drams.
Beer, four pounds.

Infuse without heat, and strain. (SWEDIAUR)

CARMINATIVE DECOCTION.

Decoctum Fructuum Juniperi carminativum, Potus carminativus. (HISP.)

R. Juniper, } of each,
Seeds of Anise, } one ounce.
Muriate of Ammonia, }
Chamomile Flowers, two ounces.
Water, four pounds.

Boil slightly, and add to the strained decoction

Proof Spirit, one pound.
Mix.

DIURETIC WINE.

Vinum Juniperinum. (AUGUSTIN)

R. Juniper, } of each,
Mustard Seeds, } half an ounce.
Horse-radish, }
Sub-carbonate of Potass, half a pound.
Rhenish Wine, eight pounds.

Digest for several days, and strain.

R. Root of Zedoary, } of each, two drams.
Squill, }
Juniper, } three drams.
Cinnamon, }
Sub-carbonate of Potass, one dram and a half.
Rhenish Wine, one pound and a half.

Digest, and strain.

OINTMENT OF JUNIPER. (VAN MONS)

R. Bruised Juniper Buds, one part.
Melted Lard, two parts.

Macerate for two hours, and strain with strong expression.

2. *Juniperus Virginiana*, LINN.—Virginian Juniper, Common Red Cedar Tree.

AMER.; COXE.

A tree of North America. (*Diaccia Monadelphica*, LINN.; *Conifera*, JUSS.; Pluk. *Alm.* 201. t. 197. f. 4.)

The leaves are very simple, lancet-shaped and pointed. When young they may be used for savine.

CERATE OF VIRGINIAN JUNIPER.

Ceratum Juniperi Virginianæ. (AMER.; COXE)

R. Leaves of Virginian Juniper, one part.
Cerate of Resin, six parts.

Mix.

3. *Juniperus Sabina*, LINN.—Savine.

SYNONIMES:—Sabine, FR.; Sadebaum, Sevenbaum, GERM.; Savina neb klasferska chwogka, BOHEM.; Sevenbom, DAN.; Sabina, ITAL.; SPAN.; and PORT.; Sevenboom, DUT.; Sawina, POL.; Sæfwenbom, SWED.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GME-LIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

A shrub of the middle of Europe and of the Levant. (*Diœcia Monadelphica*, LINN.; *Conifera*, JUSS.; *Fig. Zorn, Ic. pl. t. 525.*)

The leaves (*Folia seu Herba Sabinae*) are employed. They are oval, opposite, rather acute, and convex at the back. They have a resinous, aromatic, and very penetrating odour. Their taste is very bitter.

It is powerfully stimulant, irritant, and emmenagogue.

CONSERVE OF SAVINE.

Conserva Sabinae. (HANN.; POL. SAX.; SWEDIAUR)

R. Bruised Leaves of Savine, one part.
White Sugar, two parts.

Triturate together. (POL.; SAX.; SWE-DIAUR)

HANN. prescribes one part of leaves, and three parts of sugar.

EMMENAGOGUE POWDER. (ELLIS)

R. Savine, } of each, one dram.
Ginger, }
Sulphate of Potass, two drams.

Make six parcels.—Dose, one twice a day.

VERMIFUGE POWDER. (ELLIS)

R. Savine, twelve grains.
Senna, } of each, two scruples.
Indian Pink, }

Make six parcels.—One to be taken for three successive days, in the morning; and afterwards a dose of castor oil. The treatment to be re-commenced, if necessary.

EMMENAGOGUE BOLUSES. (BRERA; CA-DET DE GASSICOURT)

R. Powdered Savine, one dram.
Syrup of Mugwort, a sufficient quantity.

Extract of Birthwort,	} of each, half a dram.
———— Mugwort,	
Powdered Saffron,	
———— Castor,	

Make thirty-six boluses.—Dose, one three or four times a day, in amenorrhœa. (CADET DE GASSICOURT)

R. Extract of Savine, four grains.
Powdered Yew Leaves, six grains.
Despumated Honey, } of each,
Powdered Liquorice, } a sufficient
quantity.

Make six boluses.—Dose, one every three hours. (BRERA)

EMMENAGOGUE PILLS. (NIEMANN)

R. Extract of Savine, half an ounce.
Powdered Savine, two drams.
Oil of Savine, a sufficient quantity.

Make pills of two grains each.—Dose, eight morning and evening.

R. Powdered Savine, } of each,
Extract of Cham- } one dram and
mile, } a half.
Black Oxide of Iron, } of each,
Extract of Aloes, } fifteen grains.

Make pills of two grains each.—Dose, six.—In chlorosis.

ANTI-ARTHRITIC ELECTUARY. (HAMB.)

R. Powdered Savine, one dram.
———— Sweet Flag, two drams.
Honey, two ounces.
Mix.

AQUEOUS EXTRACT OF SAVINE.

Extractum Sabinae aquosum. (FERR.; PA-LAT.; SAX.; WIRTEM.)

R. Tops of Savine, one pound.
Water, six pounds.

Macerate for four days in a warm place; then boil for a short time, express, and evaporate to the consistence of an extract. (PALAT.; WIRTEM.)

FERR. directs one part of savine to be boiled in three parts of water, and the decoction to be clarified and evaporated.

R. Tops of Savine, one part.
Water, eight parts.

Infuse for twenty-four hours; then boil for a quarter of an hour, and strain with strong expression: boil the residue with four parts of water. Mix the two decoctions, and after twenty-four hours' repose, evapo-rate to the proper consistence. (SAX.)

ALCOHOLIC EXTRACT OF SAVINE.

Extractum Sabinae alcoholicum. (HERBIP.)

R. Tops of Savine,	}	of each,
Alcohol,		one pound.
Water,		eight pounds.

Digest for three days in a moderate temperature; then express, set aside and decant the liquor: distil off the alcohol, and evaporate the residue to the proper consistence.

DECOCTION OF SAVINE.

Decoctum Sabinae. (NIEMANN)

R. Herb of Savine,	one ounce.
Boiling Water,	one pound.

Boil down to six ounces; then strain, and add

Syrup,	one ounce.
--------	------------

Mix.—Dose, two spoonfuls every two hours.

INFUSION OF SAVINE.

Infusum Sabinae. (AUGUSTIN)

R. Herb of Savine,	one dram.
Boiling Water,	seven ounces.

Infuse, and strain.—Dose, a spoonful every two hours.—RAU extolled it in asthenic inflammations of the uterus and bladder.

ALKALINE TINCTURE OF SAVINE.

Tinctura Sabinae kalina seu arthritica. (AUGUSTIN; NIEMANN; SWEDIAUR; VAN MONS)

R. Fresh Leaves of Savine,	one ounce.
Alcoholic Solution of Potass,	six ounces.

Digest for four days. (AUGUSTIN; NIEMANN; VAN MONS)

This tincture may be prepared extemporaneously by dissolving two drams of essential oil of savine in six ounces of alkaline tincture.

SWEDIAUR prescribes one ounce of savine, and eight ounces of alkaline tincture.

TINCTURE OF SAVINE.

Tinctura seu Essentia Sabinae. (HAMB.)

R. Leaves of Savine,	one part.
Alcohol,	six parts.

Digest for several days, express, and filter.—Dose, thirty drops several times a day.

EMMENAGOGUE POTION. (ELLIS)

R. Tincture of Savine,	one ounce.
------------------------	------------

Tincture of Black Hellebore,
 ————— Castor, half an ounce.
 ————— two drams.
 Mix.—Dose, forty drops thrice a day.

CERATE OF SAVINE.

Ceratum seu Unguentum Sabinae. (LOND.; DUBL.; EDINB.; AMER.; BAVAR.; SUEC.; COXE; NIEMANN; SWEDIAUR; VAN MONS)

Ceratum Sabinæ. (LOND.)

R. Fresh Leaves of Savine, bruised,	one pound.
Yellow Wax (140 degrees),	half a pound.
Prepared Lard (97 degrees),	two pounds.

Melt the lard and wax together, and boil the leaves in the mixture; then strain through a linen cloth.

Unguentum Juniperi Sabinæ. (EDINB.)

R. Fresh Leaves of Savine,	two parts
Yellow Wax,	one part.
Lard,	four parts.

Melt the wax and lard together, boil the leaves in the mixture, and strain through a cloth.

Unguentum Sabinae. (DUBL.)

R. Fresh Leaves of Savine, freed from the stalks, and bruised,	half a pound.
Lard,	two pounds.
Yellow Wax,	half a pound.

Boil the leaves in the lard till they are crisp, and express; then melt the wax in the impregnated lard.

BAVAR.; SUEC.; NIEMANN; SWEDIAUR; and VAN MONS give the same formula as EDINB.

AMER.; and COXE direct six parts of cerate of resin to be mixed with one part of powdered savine.

It is employed as an epispastic.

OIL OF SAVINE.

Oleum Sabinae æthereum, Ætheroleum Sabinae. (EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BRUNS.; GALL.; FERR.; FULD.; HANN.; HASS.; OLDEN.; PALAT.; POL.;

BORUS.; ROSS.; SAX.; HERBIP.; SARD.; COXE.; PIDERIT.; SWEDIAUR.; VAN MONS)

Oleum Herbæ Juniperi Sabinæ.
(EDINB.)

Oleum Foliorum Sabinæ. (DUBL.)

R. Leaves of Savine, any quantity.
Pure Water,
a sufficient quantity
to prevent empyreuma. Distil.

AMST.; FERR.; ROSS.; and COXE give the same formula.

FULD.; and SARD. prescribe one part of savine and three parts of water.—SWEDIAUR.—one of savine and four of water.—VAN MONS.—one of savine and six of water.—HANN.; OLDEN.; POL.; BORUS.; and SAX.—one of savine and eight of water.—AUSTR.—one of savine and twelve of water.—ANT.—six of savine and twenty of water.—GALL.—five of savine and seven of water.

R. Herb of Savine, twenty-five parts.
Water, sixty-four parts.
Muriate of Soda, three parts.

Distil. (BRUNS.; HASS.; PALAT.; PIDERIT)

R. Herb of Savine, any quantity.
Distilled Water of Savine,
a sufficient quantity.

Distil. (HERBIP.)

Dose, from four to twenty drops, in common drink.

EMMENAGOGUE POTION. (CADET DE GASSICOURT.; RATIER)

R. Mugwort, five ounces.
Syrup of Orange-flowers, one ounce.
Orange-flower Water, half an ounce.
Oil of Rue, } of each, six drops.
— Savine, }

Mix.—To be taken by spoonfuls.

SAVINE WATER. (HERBIP.; PIDERIT)

R. Herb of Savine, one part.
Water, eight parts.

Macerate for twenty-four hours, and distil eight parts. (PIDERIT)

HERBIP. prescribes one part of savine and six parts of water.

SPIRIT OF SAVINE.

Spiritus Sabinæ. (HANN.; AUGUSTIN)

R. Herba Sabinæ, one part.
Alcohol, four parts.
Pure Water, a sufficient quantity.

Macerate for twenty-four hours, and distil.

STIMULANT LINIMENT. (AUGUSTIN)

R. Spirit of Savine, six ounces.
Oil of Savine, one dram.

Mix.—To be used in gout.

JUSTICIA.

Two species of this genus of plants are mentioned in pharmacopœias:

1. *Justicia Adhatoda*, LINN.—Malabar Nut.

SYNONIMES:—Adhatoda, Noyer de Ceylan, FR.; Adhatoda, CYNG.; Urus Vassica, Attaruscha, SANS.; Adatodey alley, TAM.; Adasara paku, TEL.

WIRTEM.; AINSLIE.; SPIELMANN.

A shrub of Ceylon. (*Decandria Monogynia*, LINN.; *Acanthaceæ*, JUSS.; *Fig. Pluk. Alm. IX. t. 173.*)

The leaves are employed; they are wide, oval, lanceolated, acuminate, green above, whitish beneath, and marked with large nervures. They have a bitter taste.

They are reputed emmenagogue and ecoprotic.

2. *Justicia pectoralis*, LINN.

SYNONIMES:—Carmantine pectorale, Herbe aux charpentiers, FR.

GALL.

A plant of the Antillas. (*Diandria Monogynia*, LINN.; *Acanthaceæ*, JUSS.; *Fig. Jacq. Austr. III. tab. 3.*)

The herb is employed; it consists of quadrangular, nodose, glabrous stems, and of opposite, linear, lanceolated, pointed, entire, glabrous leaves.

It is slightly astringent.

KADMIUM, ZINCUM SILESIACUM.

This is a very rare metal, and has not been very long discovered. Among its compounds, the following have been proposed to be employed in medicine.

OXIDE OF KADMIUM.

Oxydum Kadmi. (NIEMANN)

R. Kadmium, any quantity.
Diluted Sulphuric Acid, a sufficient quantity

to dissolve the metal. Let a current of hydrosulphuric acid gas pass through the cold and filtered liquor: collect and wash the precipitate, treat it with hot hydrochloric acid, and evaporate the solution nearly to dryness: add water to the residue, and mix with solution of ammonia. Wash and dry the precipitate.

The action of this oxide is not well known, nor indeed is that of the following

SULPHATE OF KADMIUM.

Sulphas Cadmi. (NIEMANN)

R. Oxide of Kadmium, any quantity.
Sulphuric Acid, a sufficient quantity

to dissolve it. Evaporate, and crystallize the liquor.

OPHTHALMIC LIQUOR. (NIEMANN)

R. Sulphate of Kadmium, from two to eight grains.
Distilled Water, one ounce.

Mix.—It is recommended externally, in chronic ophthalmia, by GRAAFE.

KALMIA LATIFOLIA, LINN.—MOUNTAIN LAUREL.

COXE.

A plant of Carolina and Virginia. (*Decandria Monogynia*, LINN.; *Rhodoraceae*, JUSS.; *Fig. Trew. Ehret*, t. 38, f. 1.)

The leaves are oval, oblong, firm, very glabrous, entire, and from two to three inches in length and one inch wide.

It is astringent [and poisonous, EDIT.].

DECOCTION OF MOUNTAIN LAUREL.

Decoctum Kalmie latifoliae. (NIEMANN)

R. Leaves of Mountain Laurel, one ounce.
Spring Water, eight ounces.

Boil down to one half, and strain.

It is recommended in chronic diarrhoea.—Dose, thirty drops, sixty times a day, till the patient becomes affected with vertigo.

KINO.

SYNONIMES:—*Gummi Kino*, *Gummi Gambiense*, *Gummi adstringens Fothergilli*.—Kino, Gomme Kino, Gomme de Gambie, FR.; Kinoharz, Kinogummi, Gambien-sergummi, GERM.; Tumble loan, TAM.; Dummulackwayn, Kadamurgarittum, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; DAN.; GALL.; FERR.; FENN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; COXE; GUI-VOL. II.

BOURT; MURRAY and GMELIN; PARIS; SPIELMANN; ZARDA.

Three principal kinds of kino are distinguished.

1. In irregular, dry, brittle masses of a deep-brown colour, opaque, inodorous, of a brilliant and nearly black fracture, and of an astringent and bitterish taste. It is supposed to be produced by evaporating to dryness, a decoction of the branches of the *Nauclea longiflora*, POIR. (*Pentandria Monogynia*, LINN.; *Rubiaceae*, JUSS.), a shrub of the Indies (*Fig. Rumph. Amb. V. tab. 34. f. 1.*)

U

2. In very small fragments of a brilliant black colour, opaque, (but translucent and a deep red when reduced to thin scales), inodorous, and of a strongly astringent taste. It is supposed to be obtained by the decoction of the wood of the *Coccoloba urifera* (*Octandria Digynia*, LINN.; *Polygoneæ*, JUSS.), a tree of Jamaica.

3. In porous masses, dull, and reddish in some spots, and shining, brilliant, and almost black in others, inodorous, and of an astringent taste. It is believed to flow from the trunk of the *Eucalyptus resinifera*, IM. (*Icosandria Monogynia*, LINN.; *Myrtææ*, JUSS.), a tree of New Holland (Fig. Whit. *Ilin.* 23).

This substance contains much tannin and extractive, that is to say, we do not thoroughly know the nature of it. It is incompatible with the salts of iron, of silver, and of lead, and with gelatine.

It is an energetic astringent.—Dose from ten grains to half a dram of the powder.

COMPOUND POWDER OF KINO.

Pulvis Kino compositus seu stypticus seu Kino gummosus. (LOND.; AUGUSTIN; CADET DE GASSICOURT; COXE)

R. Kino, fifteen drams.
Cinnamon, half an ounce.
Hard Opium, one dram.

Triturate separately, and then mix the powders. (LOND.)

The same formula is given by COXE; and CADET DE GASSICOURT.

Dose, one scruple, in cases of internal hæmorrhages.

R. Kino, half a dram.
Gum Arabic, one dram.

Make a powder, to be applied to external hæmorrhages. (AUGUSTIN)

CAMPHORATED PILLS OF KINO.

Pilule e Kino et Camphora. (AUGUSTIN)

R. Kino, two scruples.
Camphor, one scruple and a half.
Aromatic Powder, one scruple.
Oxide of Zinc, one half scruple.

Make twenty pills.—Dose, two, morning and evening; in diabetes.

ANTI-DIARRHŒIC PILLS. (ELLIS)

R. Kino, twenty grains.
Opium, two grains.
Mucilage of Gum Arabic, a sufficient quantity.

Make four pills.—Dose, one every hour.

ASTRINGENT PILLS.

Pilule stimulantés ad Gonorrhæam. (PH. SYPH.)

R. Kino, half an ounce.
Canadian Balsam, two ounces.
Root of Tormentil, a sufficient quantity.

Make pills of five grains each.

ASTRINGENT ELECTUARY. (SWEDIAUR)

R. Kino, six drams.
Alum, } of each, two drams.
Cinnamon, }
Syrup, a sufficient quantity.

Mix.—For diarrhœa, chronic dysentery, and hæmorrhages.—Dose, one dram, twice or thrice a day.

ASTRINGENT POTION. (SPIELMANN)

R. Provence Roses, two drams.
Boiling Water, four ounces.

Infuse for two hours; strain, and add

Kino, half a dram.
RABEL's Water, twenty drops.
Syrup of Pomegranate Bark, one ounce.

Mix.

ASTRINGENT INJECTION. (SWEDIAUR; PH. SYPH.)

R. Kino, from twenty to thirty grains.
Boiling Water, one pound.

Infuse for an hour, and strain. (PH. SYPH.)

R. Kino, one or two drams.
Alum, twenty or thirty grains.
Boiling Water, two pounds.

Digest for an hour, and strain. (SWEDIAUR)
For chronic urethritis.

TONIC MIXTURE.

Mixtura roborans. (SWEDIAUR)

R. Kino, two drams.
Nitrate of Potass, one dram.
Mucilage of Gum Arabic, two ounces.
Astringent Electuary, two drams.

Triturate, and add gradually,

Vinegar, one ounce and a half.
Cinnamon Water, six ounces.
Syrup of Cinnamon, one ounce.

Mix.—For uterine hæmorrhage.—Dose, one or two spoonfuls, every four or six hours.

SYRUP OF KINO. (PLANCHE)

R. Powdered Kino, six hundred and ninety grains.
Water, seven pounds.
White Sugar, two pounds.

Triturate the kino for a quarter of an hour, in a stone mortar, with four pounds of boiling water poured upon it nearly all at once. Let the liquor cool; then filter and set it aside: pour upon the residue the three other pounds of water, heated to 60 degrees. Mix the two liquors. On the other hand, boil the sugar to the consistence of an electuary. Add the kinated liquor to it, and boil to the consistence of syrup.

TINCTURE OF KINO.

LOND.; EDINB.; DUBL.; AMER.; ANT.; BELG.; FERR.; FULD.; LIPP.; LUSIT.; SAX.; BRUGNATELLI; NIEMANN; SWEDIAUR; VAN MONS.

Tinctura Kino.

R. Kino, three ounces.
Proof Spirit, two pints.

Macerate for fourteen days, and strain. (LOND.)

R. Kino, two ounces.
Dilute Alcohol,
one pound and a half.

Digest for seven days, and filter.
(EDINB.)

R. Kino, three ounces.
Proof Spirit,
one pint and a half.

Digest for seven days, and filter.
(DUBL.)

AMER.; and COXE give the same formula as LOND.—NIEMANN—the same as EDINB.—FERR.; and SWEDIAUR—two ounces of kino, one pound of alcohol (0.935) and eight ounces of cinnamon water.—BELG.—one part of kino and eight parts of alcohol (20 degrees)—BRUGNATELLI—one of kino and sixteen of alcohol (0.935).—SAX.—one of kino and five of rectified spirit.—VAN MONS—one of kino and six of alcohol.—LUSIT.—two ounces of kino and one pound and a half of spirit.—FULD.—two drams of kino, one ounce and a half of simple cinnamon water, and half an ounce of vinous cinnamon water.—LIPP.—two ounces of kino, six of distilled water, and six of spirit of nitric ether.

It is astringent.—Dose, half an ounce.

KEKEKUNEMALO.

BRUNS; WIRTEM.; MURRAY and GME-LIN; SPIELMANN.

This resin is from America, it is greenish, semi-transparent, fragile, presenting yel-

low grains in its fracture, and of an agreeable odour and bitter taste.

It is, according to GUIBOURT, a kind of gum anime. It is now disused.

KRAMERIA TRIANDRIA, RU. and PAV.—RATANY.

SYNONIMES;—*Rhatania*, DUBL. — Ratan-hie, FR.; Ruiz para los dientes, SPAN.

LOND.; DUBL.; BATAV.; BELG.; HISP.; GALL.; FERR.; HANN.; SAX.; COXE; PARIS.

A plant of South America. (*Tetrandria Monogynia*, LINN.; *Polygaleæ*, JUSS.; *Fig.* Fl. Peruv. t. I. p. 61. Ic. 93.)

The root is employed; it is woody, hard, and rounded, consisting of a thickish bark, rather irregular on the outside, of a blackish-red colour, and of a bitter and astringent taste; it covers a compact, whitish or rather yellowish wood, which is without taste or smell.

PESCHIER has discovered in it a peculiar acid, *Krameric Acid*.

It is incompatible with gelatine, the salts of iron, and the mineral acids.

It is a powerful astringent, and is much used in chronic mucous discharges, and in passive hæmorrhages.

EXTRACT OF RATANY.

Extractum Ratanhiæ. (HISP.; GALL.; FERR.; SAX.; NIEMANN; VAN MONS)

R. Bark of Ratany, one part.
Pure Water, eight parts.

Macerate for twenty-four hours; then boil for a quarter of an hour, and strain with strong expression; boil the residue with four parts of water. Mix the decoctions, set aside for twenty-four hours, and

evaporate to the proper consistence. (FERR.; SAX.; NIEMANN)

R. Ratany Bark, in powder, one pound.
Water, twelve pounds.

Infuse for two days in a water-bath, in a warm-place, frequently stirring, repeat the infusion with fresh water, till it no longer acquires any taste: filter the mixed liquors, and evaporate, first over the fire, and then in a water-bath, to dryness. (HISP.)

R. Ratany Bark, in powder, any quantity.

Add water to it, and leave it to macerate for twenty-four hours; then add a quantity of cold water, equal in weight to that of the whole mass, and boil for fifteen or twenty minutes; strain the boiling liquor through flannel; pour upon the residue twice its weight of boiling water, and boil again for a quarter of an hour. Strain, mix the two liquors, set aside for an instant, decant, and evaporate with constant ebullition. (VAN MONS)

R. Saturated Alcoholic Tincture of Ratany, any quantity.

Distil one-fourth from it in a water-bath; then evaporate the residue slowly to the consistence of a dry extract. (GALL.)

Dose, from twenty grains to half an ounce.

DECOCTION OF RATANY.

Decoctum Radicis Ratanhiæ. (FERR.; NIEMANN)

R. Ratany Root, half an ounce.
Spring Water, two pounds.

Boil down to half a pound. (NIEMANN)

FERR. gives the same formula, but requires the subsequent addition of two drams of vinegar, and four of sugar.

It is astringent.—Dose, from one to three ounces.

AQUEOUS TINCTURE OF RATANY. (FERR.)

R. Dry Extract of Ratany, five drams.
Pepperwort, four ounces and a half.
Water, nine ounces.

Macerate for three days, and strain.

ACETOSE TINCTURE OF RATANY. (FERR.)

R. Dry Extract of Ratany, one dram.
Sulphate of Soda, one ounce.
Water, six ounces.

Dissolve, and add
Vinegar, one dram.

Mix.

TINCTURE OF RATANY.

Tinctura Ratanhiæ. (FERR.; NIEMANN; VAN MONS)

R. Powdered Ratany Bark, three ounces.

Alcohol, two pounds.

Digest for eight days, in a moderate heat, and strain. (NIEMANN)

VAN MONS prescribes one part of ratany and six parts of proof spirit.

R. Extract of Ratany, four drams.
Alcohol (24 degrees), four ounces.
Vinegar, eight ounces.

Digest for three days, and filter. (FERR.)

It is astringent.—Dose, from thirty to eighty drops.

COMPOUND TINCTURE OF RATANY.

Tinctura Ratanhiæ composita. (FERR.; NIEMANN)

R. Powdered Ratany Root, three ounces.
Orange Peel, two ounces.
Snake Root, half an ounce.
Saffron, one dram.
Alcohol, two pounds.

Digest for twelve days, and strain.

It is astringent.—Dose, from sixty to eighty drops.

ASTRINGENT GARGLE. (FERR.)

R. Extract of Ratany, four drams.
Water, twenty ounces.

Dissolve and add

Vinegar, four ounces.

Mix.

ASTRINGENT OINTMENT. (FERR.)

R. Extract of Ratany, } of each,
Olibanum, } four ounces.
Galbanum, two ounces.
Ointment of Armenian Bole, six ounces.
Turpentine, two ounces and a half.
Olive Oil, two ounces.

Mix.—To be applied to ruptures.

ASTRINGENT PILLS. (FERR.)

R. Extract of Ratany, one dram.
Water, a sufficient quantity
to soften the extract.—Make ten or twelve pills.

ASTRINGENT ELECTUARY. (FERR.)

R. Powdered Ratany, one ounce.
Ginger, one dram.
Syrup of Orange Peel,
a sufficient quantity.

Mix.—Dose a tea spoonful; in fevers and in epileptic fits.

ASTRINGENT BOLUSES. (CADET DE GAS-SICOURT)

R. Powdered Extract of Ratany,
one scruple and a half.

Powdered Alum, ten grains.
Conserve of Hips, one dram.
Syrup, a sufficient quantity.

Make twenty boluses.—Dose, two every three or four hours.

ASTRINGENT POTION. (FERR. ; BRERA ; CADET DE GASSICOURT ; RATIER)

R. Extract of Ratany, one dram.
Rose Water, four ounces.
Syrup of Cinchona, one ounce.

Mix. (RATIER)

R. Extract of Ratany, half a dram.
Alum, ten grains.
Infusion of Roses, five ounces.
Syrup of Roses, } of each,
Catechu, } one ounce.

RABEL'S Water, fifteen drops.

Mix. (CADET DE GASSICOURT)

R. Compound Tincture of Ratany, } of each,
Powdered Ratany, } six drams.
Water, six ounces.

Mix. (FERR.)

R. Infusion of Roses, one ounce.
Diluted Sulphuric Acid, forty drops.
Extract of Ratany, two drams.
Syrup of Mulberries, six drams.

Mix. (BRERA)

To be taken by spoonfuls.—The last particularly has been recommended in uterine hæmorrhage, and the last but one in fevers and in epileptic fits.

LAC.—MILK.

SYNONIMES :—Lait, FR. ; Milch, GERM. ; Lebu, ARAB. ; Leche, SPAN. ; Dud, HIND. ; Milk, DUT. ; Latte, ITAL. ; Schir, PERS. ; Khschira, PAYAS, DUGHDA, SANS. ; Jœlk, SWED.

AMST. ; BAVAR. ; HISP. ; GALL. ; FORM. HÔP. FR. ; HAMB. ; HASS. ; SUEC. ; WIRTEM. ; AINSLIE ; GUIBOUT ; PIDERIT ; SPIELMANN.

An opaque, white, slightly oleaginous liquid, rather heavier than water. It is secreted by the mammary glands of the females of mammiferous animals, the secretion commencing a short time after parturition. It has a bland, agreeable, and more or less of a sweet taste, with a peculiar odour, which becomes dissipated in proportion as it becomes cold.

The kinds of milk employed are

1. Cow's Milk ; *Lac vaccinum seu bubulum*.

SYNONIMES :—Lait de Vache, FR. ; Kuhmilch, GERM. ; Ellakerry, CYNG. ; Ghayka dud, DUK. ; Gaicha dud, MAH. ; Susu, MAL. ; Gokschira, SANS. ; Paschuin paal, TAM. ; Aupatu, TEL.

2. Goat's Milk ; *Lac caprinum*.

SYNONIMES :—Lait de Chèvre, FR. ; Ziegenmilch, GERM. ; Jlukirie, CYNG. ; Chaylie ka dud, DUK. ; Leche de cabra, SPAN. ; Ajakschira, SANS. ; Aata paal, Veltata paal, TAM. ; May ka palu, TEL.

3. Asses Milk ; *Lac asininum*.

SYNONIMES :—Lait d'ânesse, FR. ; Eselsmilch, GERM. ; Cotalukirie, CYNG. ; Ga-

diko dud, DUK. ; Ghadwa tscha dud, MAH. ; Kharakhira, SANS. ; Kalady paal, TAM. ; Gadilay paalu, TEL.

The two first very nearly resemble each other, but the last contains much more cream than they do.

Milk, left undisturbed for some time, separates into three portions :

1. Cream ; *Cremor*.

SYNONIMES :—Crème, FR. ; Milchrahm, GERM. ; Grædda, SWED.

2. Caseum.

3. Serum ; *Lac ebutyratum*.

SYNONIMES :—Sérum, FR. ; Buttermilch, GERM. ; Dogh, ARAB. ; Schaatsch, DUK. ; Mutha, HIND. ; Takram, Dandahatam, SANS. ; Moru, TAM. ; Isalla, TEL.

By beating the cream, butter, *Butyrum*, is obtained.

SYNONIMES :—Beurre, FR. ; Butter, GERM. ; Manteca de vacca, SPAN. ; Boter, DUT. ; Burro, ITAL. ; Smør, SWED.

AMST. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; HISP. ; GALL. ; HASS. ; LIPP. ; PALAT. ; SAX. ; SUEC. ; WURCEB. ; GUIBOUT ; PIDERIT ; SPIELMANN.

This is a fat and solid substance, varying in colour from white to yellow, of an agreeable taste, and of a slightly aromatic odour. It contains, according to CHEVREUL, caseous matter, serum, elaine, stearine, a co-

louring principle, a little butyric acid, and an oil.

The coagulum which is found in the stomach of the calf, (GALL.; GUIBOUT) is mixed with the acetic, hydrochloric, and butyric acids which are secreted by the internal surface of this stomach, and which give to it a sour smell.

By evaporating the serum or whey, the sugar of milk is obtained, *Saccharum Lactis*.

SYNONIMES:—Sucre de lait, FR.; Milchsucker, GERM.; Melksukker, DAN.; Melkzuiker, DUT.; Zuccherò di latte, SPAN.

AMST.; BAVAR.; BRUNS.; DAN.; FENN.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; GUIBOUT.; PIDERIT.; SPIELMANN.; VAN MONS.

It crystallizes in white, semi-transparent parallelopipedons; it has a slightly saccharine, rapid, earthy taste.

HYDROGALE. (RATIER)

R. Cow's Milk, four ounces.
Barley Water, two pounds.
Mix.

PECTORAL POWDER. (NIEMANN)

R. Sugar of Milk, one ounce.
Gum Arabic, five drams.
Oleo-saccharum of Fennel, three drams.
Mix.

PECTORAL PTISAN.

Potus Sacchari Lactis. (AUGUSTIN)

R. Sugar of Milk, half an ounce.
Boiling Water, two pounds.
Dissolve.

LACCA.—LAC, GUM LAC.

SYNONIMES:—*Lacca Gummi*; Καρχαμον. —Laque, Gomme Laque, FR.; Gummi-lack, Lack, GERM.; Laak, ARAB.; Kam-balo, BA.; Lakada, CYNG.; Lacca, PORT.; and SPAN.; Lak'h, HIND.; Balo, JAV.; Ainbalu, MAL.; Leka, POL.; Lakscha, SANS.; Lacka, SWED.; Komburruk, TAM.; Commolekka, TEL.

AMST.; ANT.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; GUIBOUT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.

There are three sorts of lacs.

1. Stick Lac; *Lacca in ramulis* seu *ramis* (Laque en batons, FR.; Stangenlack, GERM.). This is of a more or less deep red colour, nearly transparent, unequal, rough, hard, friable, forming a crust of some inches in length, and about a line in thickness, over a slender stick, which it rarely surrounds entirely. It is pierced at its surface with a great many holes communicating with oval cellules placed against each other without regularity, and from whence a white cottony substance comes out.

2. Seed Lac; *Lacca in granis* (Laque en grains, FR.; Kærnerlack, GERM.). This consists of fragments of the preceding, detached from the branches.

3. Shell Lac, Lamp Lac; *Lacca in ta-*

bulis seu *in massis* (Laque en masses ou en pains, FR.; Schellack, GERM.). This is in plates or thin scales, obtained by melting either of the preceding and thus presenting the lac free from foreign substances.

Lac is produced by the puncture of the female of the *Coccus lacca*, KERR, a hemipterous insect of the Indies (*Fig. Journ. compl. des Sc. med. t. X. p. 20*), which it makes in the bark of a great number of resinous trees, among which we distinguish particularly the *Ficus religiosa*, LINN.; the *Croton castaneifolium*, LINN.; the *Rhamnus Jujuba*, LINN.; the *Mimosa cinerea*, WILLD.; and the *Mimosa corinda*, WILLD.—VIREY thinks, in contradiction to the opinion of LATREILLE, and justly, according to appearance, that the insect does not elaborate this substance, but that the production of it is occasioned by the punctures which it makes in the bark. He thinks that the resinous matter owes its fine colour to the insect, from the purple-red colour with which the female and the eggs of the cochineal are impregnated.

It is composed, according to FUNKE, of resin, of a peculiar matter, *Laccin*, intermediate between wax and resin, and of an animal colouring matter. JOHN says that he has also found in it a peculiar acid, *Laccic Acid*.

AQUEOUS TINCTURE OF LAC.

Tinctura aquosa Laccæ seu *Gummi Laccæ aquosa*. (BRUNS.; HISP.; PALAT.;

ROSS.; SARD.; SAX.; WIRTEM.; SPIELMANN)

- R. Seed Lac, in powder, one part.
 Alum, one half part.
 Distilled Water, eight parts.

Boil together till reduced to one half.
 When cold, add

Water of Red Roses, } of each,
 Sage, } two parts.

Filter. (ROSS.; SAX.)

- R. Powdered Lac, one ounce.
 Alum, half an ounce.
 Rose Water, two pounds.

Digest in a covered vessel, and filter when it has become of a fine red colour. (HISP.)

- R. Powdered Lac, one ounce.
 Alum, one dram and a half.
 Sage Water, three ounces.
 Rose Water, one ounce and a half.

Boil together, and filter. (SPIELMANN)

- R. Lac, one ounce.
 Alum, one dram.
 Sage Water, } of each,
 Rose Water, } two ounces.
 Scurvy-grass Water, }

Boil slowly, and strain. (SARD.)

BRUNS. prescribes one ounce of lac, two drams of alum, and two ounces each, of the three waters.—WIRTEM.—one ounce of lac, one dram and a half of alum, and one ounce and a half each, of the three waters.

- R. Lac, six drams.
 Alum, two drams.
 Sage Water, } of each,
 Selfheal Water, } one pound.
 Scurvy-grass Water, }

Digest, without boiling, till the liquor becomes of a fine red colour, and strain. It may be flavoured by adding two scruples of cloves. (PALAT.)

It is excitant, and is reputed anti-scorbutic.—It is employed in lotions and gargles.

ALCOHOLIC TINCTURE OF LAC. (AMST.; FULD.; VAN MONS)

- R. Stick Lac, in powder, one part.
 Alcohol, six parts.

Infuse for twelve hours in a water-bath, and filter. (VAN MONS)

AMST. prescribes one part of lac, and eight parts of alcohol (0.884), to be infused for eight days.—FULD.—one of lac and sixteen of spirit of scurvy-grass.

It is recommended in affections of the mouth and gums attributed to weakness, and is most frequently employed in the form of gargle: but it is likely that it only acts by virtue of the alcohol which it contains.

COMPOUND ALCOHOLIC TINCTURE OF LAC. (BAVAR.; FERR.; LUSIT.; ROSS.; SARD.; SUEC.; HERBIP.)

- R. Powdered Lac, one ounce.
 Myrrh, three ounces.
 Spirit of Scurvy-grass, one pound and a half.

Digest for some days, and strain. (ROSS.; SUEC.)

LUSIT. prescribes one ounce of lac, three drams of myrrh, two pounds of the spirit, and eight days' maceration.—FERR.—one ounce of lac, two drams of myrrh, and one pound and a half of the spirit.

- R. Lac, in powder, two ounces.
 Alum, in powder, one dram and a half.
 Spirit of Scurvy-grass, one pound.

Digest in a sand-bath, and strain. (SARD.)

HERBIP. directs one ounce of lac and two scruples of alum to be digested in one pound of spirit of scurvy-grass, till the liquor assumes a fine red colour.—LUSIT.—one ounce of lac, three drams of myrrh, two pounds of spirit, and eight days' digestion.—BAVAR.—five parts of lac, one part of alum, thirty-six of water, and five of spirit.

It is slightly stimulant, and is recommended in lotions and gargles, to asthenic ulcerations of the mouth. The tincture prepared according to the second formula is more active and astringent than the other.

ALKALINE TINCTURE OF LAC.

Tinctura Laccæ kalica. (BRUNS.; HAMB.; HASS.; PIDERIT; VAN MONS)

- R. Seed Lac, one ounce.
 Myrrh, half an ounce.

Pulverize, and pour upon them

Deliquesced Sub-carbonate of Potass, half an ounce.

Reduce the whole to a soft paste; dry it over a slow fire, and pour upon the pulverized mass

Spirit of Scurvy-grass, six ounces.

Filter, after three or four days' digestion.

BOERHAAVE recommended this tincture internally, in gout and rheumatism; and externally, in scorbutic affections of the mouth and gums. He gave it in Spanish or Madeira wine. The sub-carbonate of potass is the principal source of its activity.

ALUMINATED TINCTURE OF LAC. (ANT.; HISP.; SARD.; HERBIP.)

- R. Powdered Lac, two ounces.
 Alum, one dram and a half.
 Spirit of Scurvy-grass, one pound.

Digest in a sand-bath, and strain. (SARD.)

HERBIP. prescribes half an ounce of lac, two scruples of alum, and one pound of

spirit.—ANT.—one ounce of lac, two drams of alum, and one pound of spirit.

- R. Pure Lac, one ounce.
 Alum, dissolved in water, one dram and a half.
 Spirit of Scurvy-grass, eight ounces.

Digest till the liquor is of a fine red colour, and filter. (Hisp.)

This is for the same uses as the simple tincture; but it is more astringent.

LACTUCA.—LETTUCE.

Three species of this genus of plants are employed:

1. *Lactuca sativa*, LINN.—Garden Lettuce.

SYNONIMES:—Laitue cultivée, FR.; Lattich, Gartensalat, GERM.; Salat, Locyka, BOHEM.; Laktuk, DAN.; RUSS.; and SWED.; Lechaga, SPAN.; Gewoone Salade, Lataw, DUT.; Lattuca, ITAL.; Loczyga, Letuga, POL.

LOND.; EDINB.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; WIRTEM.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

This plant is cultivated in gardens, where it has produced numerous varieties. (*Syn-genesia Polygamia Æqualis*, LINN.; *Synantherea*, CASS.; Fig. Flore Medic. IV. 213.)

The herb and seeds are employed.

The herb (*Herba Lactucæ*) is composed of oblong, oval, undulated leaves, of a watery sweetish taste.

The seeds are long, flattened, pointed at both ends, and of a grey colour.

2. *Lactuca Scariola*, LINN.

SYNONIME:—Laitue sauvage, FR.

AMER.; AUSTR.; GALL.; FERR.; LIPP.; WIRTEM.; GUIBOURT.

A plant of Europe. (Fig. Zorn, *Ic. pl. t. 294.*)

The herb (*Herba Lactucæ sylvestris* seu *scariolæ*) is employed. It consists of alternate, sessile, amplexicaul, long leaves, sagittate at the base, and acute at the summit. They are usually pinnatifid, bordered with some spinescent teeth, and glabrous. It has a disagreeable odour, and a bitter taste.

3. *Lactuca virosa*, LINN.—Vinous Lettuce, Strong-scented Lettuce, Hemlock Lettuce.

SYNONIMES:—Laitue vireuse, FR.; Giffiger Lattich, Stinkender Lattich, GERM.; Locyka lesnjneb plana, BOHEM.; Stinkende Laktuk, DAN.; Stinkende Salade,

DUT.; Salata jadowita, Locyga dzika, POL.; Alsace brava, PORT.

EDINB.; ANT.; BAVAR.; BRUNS.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; POL.; ROSS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of Europe. (Fig. Zorn, *Ic. pl. t. 285.*)

The herb (*Herba Lactucæ virosæ* seu *Intybi angusti*) is employed. It is composed of alternate, sessile, embracing, long, notched leaves, obtuse at the top, and the lower ones sinuated and toothed. It has a nauseous odour, and an acrid and very bitter taste.

The garden lettuce is considered as a sedative, but it undoubtedly acts only as an emollient. [I think M. JOURDAN is mistaken in this opinion.—EDIT.]—Dose, of the juice, from half an ounce to two ounces.—As to the viros, it is exceedingly stimulant: it acts on the nervous system, and is also diuretic.

EXTRACT OF GARDEN LETTUCE.

Extractum Lactucæ. (LOND.)

Succus spissatus Lactucæ sativæ.
 (EDINB.)

R. Fresh Lettuce Leaves, one pound.

Bruise them in a stone mortar, sprinkling them with a little water; then express the juice, and evaporate it unstrained till it acquires the proper consistence.

EXTRACT OF VIROUS LETTUCE.

Extractum Lactucæ virosæ. (EDINB.; ANT.; AUSTR.; BELG.; FULD.; HANN.; HASS.; LIPP.; ROSS.; WIRTEM.; HER-

BIP. ; BRUGNATELLI ; SWEDIAUR ; VAN MONS)

Succus spissatus Lactucæ virosæ.
(EDINB.)

R. Leaves of Virous Lettuce,
any quantity.

Bruise them in a stone mortar, sprinkling them with a little water ; then express the juice, and evaporate it in a water-bath, constantly stirring towards the end with a spatula.

FULD. ; HAMB. ; LIPP. ; ROSS. ; HERBIP. ; and BRUGNATELLI give the same formula.

SWEDIAUR directs ten pounds of leaves to be bruised, adding six pounds of water to them ; the juice to be expressed, to be brought to a single boil, and then to be evaporated immediately.

R. Herb of Virous Lettuce, one pound.

Bruise it in a marble mortar, and express the juice ; boil the residue for an hour with three pounds of water, and express. Mix the liquors, and evaporate. (HASS.)

R. Fresh Herb of Virous Lettuce,
any quantity.

Express the juice, and keep it for twenty hours in a cool place that it may not ferment ; decant, boil for a few moments, and strain through flannel ; clarify with white of egg, and strain again : then evaporate the liquor slowly to one half, let it cool, clarify again with white of egg, and strain : finally, evaporate it to the required consistence. (WIRTEM.)

R. Fresh Leaves of Virous Lettuce,
any quantity.

Bruise them with a little water, and pass through a sieve ; set aside, decant, coagulate at the fire, and strain again : then evaporate to the consistence of pilular mass, remove it from the fire, incorporate the fecula set apart, and evaporate again to the proper consistence. (ANT. ; VAN MONS)

R. Fresh Leaves of Virous Lettuce,
any quantity.

Bruise them in a mortar, and express the juice ; evaporate this till the addition of one fourth of powdered dry herb of lettuce is required to give the consistence of an extract. (BELG.)

R. Fresh Leaves of Virous Lettuce,
any quantity.
Water, a sufficient quantity.

Exhaust the herb by several successive ebullitions, mix the liquor, and after decantation, evaporate them in a water-bath. (AUSTR.)

Dose, from four to twelve or more grains.

WATER OF GARDEN LETTUCE. (GALL. ; SARD.)

R. Fresh Herb of Lettuce, one part.
Water, two parts.

Distil two thirds. (SARD.)

GALL. prescribes five thousand parts of fresh lettuce, and twelve thousand five hundred of water ; ten thousand parts to be distilled, and poured upon a similar quantity of fresh lettuce, adding ten thousand parts of water ; ten thousand parts to be again distilled, and the operation to be repeated a third time.

Dose, from half an ounce to two ounces.

SEDATIVE PEDILUVIUM. (NIEMANN)

R. Lettuces, eight or ten.
Water, a sufficient quantity.

Infuse for a short time, and boil. It is to be used without straining.

TINCTURE OF VIROUS LETTUCE.

Tinctura seu Essentia Lactucæ virosæ.
(SAX. ; NIEMANN)

R. Juice of Virous Lettuce, } of each,
recently expressed, } an equal
Concentrated Alcohol, } part.

Macerate without heat for several days, and filter. (SAX.)

R. Herb of Virous Lettuce, one ounce.
Cinnamon, one dram.
Concentrated Alcohol, } of each,
Distilled Water, } four ounces.

Digest for several days, express, and filter. (NIEMANN)

It is stimulant and narcotic.—Dose, fifteen drops, to be gradually augmented.

LACTUCARIUM.

SYNONIMES :—Lettuce-opium.—Thridace, Lactucarium, Fr.

EDINB.; DUBL.; AMER., COXE; NIEMANN; SPIELMANN.

This substance is hard and brittle, of a deep-brown colour, of a slightly virous odour, and of a strong bitter taste.

It is the solidified milky juice of the *Lactuca sativa*, LINN.; and is obtained by cutting the stems horizontally.

According to CAVENTOU, lactucarium does not contain morphine, but malic acid and lime.

The property is attributed to it of producing sleep without occasioning stupor, of diminishing the frequency of the pulse, and of lessening animal heat.—Dose, from two to four grains or more, progressively.

TINCTURE OF LACTUCARIUM.

Tinctura Lactucarii. (COXE; NIEMANN)

R. Lactucarium, one ounce.
Proof Spirit, one pint.

Digest, and filter.—Dose, from ten to sixty drops.

SYRUP OF LACTUCARIUM. (FEE)

R. Fresh juice of the Stems of Lettuce, deprived of their leaves at the time of flowering, one pound.
Sugar, two pounds.
Dissolve without heat and filter.

RESOLVENT PILLS.

Pilulæ torpentes ac resolventes. (BRERA)

R. Lactucarium, twelve grains.
Rob of Elder, } of each, a
Extract of Liquo- } sufficient quantity.
rice, }

Make four pills.—Dose, one every three hours; in obstinate cough without expectation, and in acute dropsy.

SEDATIVE POTION.

Potio nutriens et sedans. (BRERA)

R. Iceland Moss, two ounces.
Spring Water, a sufficient quantity,
to obtain eight ounces of decoction. Add when strained,

Lactucarium, half a dram.
Mucilage of Gum Arabic,

Simple Syrup, half an ounce.
one ounce.

Mix.—To be taken gradually, in obstinate cough and tuberculous phthisis.

LAMIUM ALBUM, LINN.—WHITE DEAD NETTLE;
ARCHANGEL.

SYNONIMES :—Ortie blanche, Fr.; Weisse Bienensang, Weisse Todlennessel, GERM.; Hlucha, Mrtwa, Smzadlawa Kopriva, BOHEM.; Dædnelde, DAN.; Ortiga muerta, SPAN.; Pokvzywa martiva, POL.; Ortiga morta, PORT.; Blind nesta, SWED.

ANT.; BRUNS.; GALL.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

A very common plant throughout Europe.

(*Didynamia Gymnospermia*, LINN.; *Labiata*, JUSS.; *Fig. Zorn, Ic. pl. t. 80.*)

The herb and flowers are employed.

The herb (*Herba Lamii albi seu Galeopsisidis seu Urticæ mortuæ.*) consists of a nearly glabrous, rather hairy stem, and of cordiform, petiolated, acuminate leaves, bordered with acute teeth. It has a disagreeable odour and a styptic taste. It is an astringent stimulant.

The flowers, of which the corollæ only are used, are white, of an unpleasant smell and sweetish taste. They are emollient.

LASERPITIIUM.—LASSERWORT

Two species of this genus of plants are mentioned in pharmacopœias.

1. *Laserpitium latifolium*, LINN.—Broad-leaved Lasserwort.

SYNONIMES :—Laser à large feuilles, Gentiane blanche, FR. ; Hirschwurz, Weiser Enzian, GERM.

BRUNS. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; Fig. Jacq. Fl. Austr. t. 146.)

The root (*Radix Gentiane alba* seu *Cervariæ albæ*) is employed. It is very long, cylindrical, of a yellowish-brown colour without, and of a yellowish-white within. It has an aromatic and agreeable smell, and an aromatic, acrid, and bitterish taste.

It is tonic and stimulant.

2. *Laserpitium Siler*, LINN.—Mountain Lasserwort.

SYNONIMES :—Laser sermontain, FR. ; Roskümelartiges Laserkraut, GERM. ; Eselskomin, DUT. ; Laserpicio Silerino, PORT.

BRUNS. ; GALL. ; WIRTEM. ; HERBIP. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of the Alps. (Fig. Jacq. Fl. Austr. t. 145)

The seeds (*Semines Sileris montani*) are employed. They are long, flattish, grooved and furnished with narrow wings. They have an aromatic and pleasant odour, and an aromatic, acrid, and bitter taste.

It is tonic and stimulant.

LAVANDULA.—LAVENDER.

Two species of this genus of plants are employed in medicine.

1. *Lavandula Spica*, LINN.—Làvender.

SYNONIMES :—Lavande, FR. ; Lavandel, GERM. ; DUT. ; DAN. ; and SWED. ; Espliego, Alhugema, SPAN. ; Lavendola, ITAL. ; Lawanda, POL. ; Alfazema, PORT.

LOND. ; EDINB. ; DUBL. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FORM. HÔP. FR. ; FULD. ; GENÈV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; CAST. BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; WURCEB. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOUT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN.

A plant of the middle of Europe. (*Diodynamia Gymnospermia*, LINN. ; *Labiata*, JUSS. ; Fig. Zorn, Ic. pl. t. 53)

The flowering tops (*Herba et Flores* seu *summitates* seu *spicae florentes Lavandulae* seu *spicae*) are employed. They consist of straight ramified stalks, furnished with linear, downy, whitish leaves, and terminated by blue or violet-coloured verticillated flowers whose clusters form an interrupted spike, furnished with narrow linear bractææ. They have a penetrating and very agreeable odour, and an aromatic and bitter taste.

It is an energetic stimulant.—Dose, from a scruple to half a dram, of the powder.

AROMATIC SPECIES. (FORM. HÔP. FR. ; HANN. ; POL. ; BORUS. ; SARD. ; HÜFELAND ; SPIELMANN ; SWEDIAUR)

R. Herb of Marjoram,	} of each, two parts.
—— Peppermint,	
—— Rosemary,	
—— Wild Thyme,	
Flowers of Lavender	} of each, one part.
Cloves,	
Cubebs,	

Make a coarse powder. (HANN. ; POL. ; BORUS. ; SARD.)

R. Herb of Rosemary,	} of each, an equal part.
—— Rue,	
—— Sage,	
—— Hyssop,	
—— Lavender,	
—— Wormwood,	
—— Origanum,	
—— Thyme,	
Leaves of Bay,	
Petals of Red Roses,	
Flowers of Chamomile,	} of each, an equal part.
—— Melilot,	
—— Elder,	
Muriate of Ammonia,	

Make a coarse powder. (SPIELMANN)

For fomentation.

R. Flowers of Lavender,
 ————— Chamomile,
 Herb of Peppermint,
 ————— Balm,
 ————— Thyme,
 ————— Marjoram,
 Cut and mix. (HUFELAND)

} of each, an
 equal part.

R. Flowers and Tops of Lavender,
 ————— Balm,
 ————— Peppermint,
 ————— Sage.
 ————— Thyme,
 ————— Wormwood,
 ————— Water-mint,
 ————— Origanum
 ————— Rosemary
 Cut and mix. (FORM. HÔP. FR.)

} of each,
 an
 equal part.

CEPHALIC SPECIES.

Species cephalica pro Fomentatione seu Epithemate. (PALAT.; WIRTEM.; HERBIP.; VAN MONS)

R. Herb of Marjoram,
 ————— Mint,
 ————— Balm,
 ————— Wild Thyme,
 Flowers of Lavender,
 ————— Arnica,
 Cut, and mix. (HERBIP.)

} of each,
 an equal part.

R. Herb and Tops of Betony,
 ————— Hyssop,
 ————— Lavender,
 ————— Marjoram,
 ————— Origanum,
 ————— Rosemary,
 ————— Rue,
 ————— Sage,
 ————— Wild Thyme,
 Flowers of Chamomile,
 ————— Melilot,
 ————— Roses,
 ————— Elder,
 Leaves of Bay,
 Muriate of Ammonia, half an ounce.
 Cut, bruise, and mix. (PALAT.; WIRTEM.; VAN MONS)

} of each,
 one ounce.

INFUSION OF LAVENDER. (FORM. HÔP. FR.)

R. Dried Tops of Lavender,
 ————— Grated Liquorice,
 Boiling Water,
 to obtain two pounds of strained infusion.

} of each,
 two drams and
 a half.

COMPOUND AROMATIC TINCTURE.
(FORM. HÔP. FR.)

R. Aromatic Species, one part.
 Alcohol (22 degrees), nine parts.

Infuse the species for eight days in one half of the alcohol; then express, and infuse the residue in the other half of the spirit; then mix and filter the tinctures.

It is employed externally in lotions and fomentations, usually mixed with wine.—It is stimulant.

VINEGAR OF LAVENDER.

Acetum Lavandulae. (GALL.; LUSIT.; PALAT.; WIRTEM.; HERBIP.; PIDERIT)

R. Flowers of Lavender, one part.
 Vinegar, six parts.

Macerate in the sun for several days, and express. (PALAT.; WIRTEM.; PIDERIT)

GALL. prescribes one part of lavender, sixteen parts of vinegar, and fifteen days' maceration.—HERBIP.—one of lavender, nine of vinegar, and three weeks' maceration.

R. Dried Flowers of Lavender, one pound.
 Vinegar, six pounds.

Macerate for six days; boil for a short time, and add when strained,

Alcohol, half a pound.

Mix. (LUSIT.)

It is aromatic, and is sometimes employed externally.

COMPOUND VINEGAR OF LAVENDER.

Acide acétique aromatique, Esprit aromatique acéteux, Vinaigre des quatre voleurs. (FENN.; BRUGNATELLI; NIEMANN; VAN MONS)

R. Cinnamon, } of each, one dram.
 Cloves, }
 Nutmegs, }
 Peppermint, half an ounce.
 Lavender, } of each, one ounce.
 Rosemary, }
 Vinegar, eight ounces.
 Proof Spirit, twenty ounces.

Distil twenty-eight parts. (VAN MONS)

R. Fresh Herb of Wormwood,
 ————— Mentha
 ————— Crispa,
 ————— Rosemary,
 ————— Rue,
 Dried Flowers of Lavender,
 two ounces.

} of each,
 one ounce
 and a half.

Root of Sweet Flag,	} of each, two drams.
Cinnamon,	
Nutmegs,	
Cloves,	
Vinegar,	ten pounds.

Macerate for ten days; express, and filter. (BRUGNATELLI)

R. Tops of Wormwood,	} of each, four drams.
——— Milfoil,	
Herb of Peppermint,	
Flowers of Lavender,	
Cloves,	
Ginger,	
Boiling Vinegar,	seven pounds and a half.

Acetic Acid, half a pound.

Macerate for four days; express, clarify with cow's milk, strain, and add

Spirit of Rosemary, four ounces.

Mix. (FENN.)

R. Cinnamon,	} of each, half an ounce.
Mace,	
Cloves,	
Rhodium Wood,	
White Sanders Wood,	} of each, two drams.
Seeds of Fennel,	
Vinegar,	four pounds.

Having digested these for two days, in a moderate heat, add

Herb of Wormwood,	} of each, half an ounce.
——— Rosemary,	
——— Peppermint,	
Flowers of Lavender,	
Herb of Marjoram,	one dram.

Digest again for two days; then boil, and strain when cold. (NIEMANN)

LAVENDER WATER. (AUSTR.; GALL.; PALAT.; SARD.; WIRTEM.)

R. Flowering Herb of Lavender, one part.
Water, four parts.

Distil one half. (GALL.; PALAT.; SARD.)

HERBIP. prescribes one part of lavender, and six parts of water.—AUSTR.—one of lavender, and twelve of water.

ESSENTIAL OIL OF LAVENDER.

Oleum Lavandulae aethereum, Aetheroleum Lavandulae. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR)

Oleum Lavandulae. (LOND.)

Oleum volatile Lavandulae Spicae. (EDINB.)

Oleum Florum Lavandulae. (DUBL.)

R. Flowers of Lavender,
any quantity.

Water, a sufficient quantity.

Distil, and collect the supernatant oil.

The same is required by AMER.; AMST.; BATAV.; BELG.; FERR.; FULD.; GENEV.; LUSIT.; ROSS.; SUEC.; BRUGNATELLI; and COXE.

SWEDIAUR prescribes one part of flowering herb, and four parts of water.—HISP.; HANN.; POL.; BORUS.; and SAX.—one of herb, and eight of water.—DAN.—one of herb, and six of water.—ANT.—six of herb, and twenty of water.—GALL.—five of herb, and seven of water.—BAVAR.—one of herb, and nine of water.—AUSTR.—one of herb, and twelve of water.

R. Lavender, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Distil. (BRUNS.; HASS.; PALAT.; SARD.; WIRTEM.; PIDERIT)

R. Lavender, any quantity.
Lavender Water, a sufficient quantity.

Distil. (HERBIP.)

Dose, from two to eight drops.

BALSAM OF LAVENDER.

Linimentum aromaticum. (BRUNS.; SWEDIAUR)

R. Oil of Lavender, one part.
——— Nutmegs, five parts.

Mix. (BRUNS.)

R. Oil of Lavender, } of each,
——— Nutmegs, } an equal part.
Cocoa Nut Oil, }

Mix. (SWEDIAUR)

SPIRIT OF LAVENDER.

Spiritus Lavandulae. (LOND.; EDINB.; DUBL.; AMER.; ANT.; AUSTR.; BAVAR.; BELG.; BRUNS.; DAN.; FENN.; FERR.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Spiritus Lavandulae.

R. Fresh Lavender Flowers,
two pounds.

Rectified Spirit, one gallon.
Water, a sufficient quantity
to prevent empyreuma. Macerate
for twenty-four hours, and then distil
a gallon with a gentle heat. (LOND.)

DUBL. gives the same formula, but
requires only five pints to be distilled.

Spiritus Lavandulæ Spicæ. (EDINB.)

R. Fresh Lavender Flowers,
two pounds.
Alcohol, eight pounds.

Distil seven pounds by the heat of
a water-bath.

R. Lavender, one part.
Alcohol (20 degrees,) four parts.

Distil a spirit marking 28 degrees.
(ANT.; BRUNS.; HASS.; PALAT.; WIR-
TEM.; PIDERIT)

BELG. prescribes one part of lavender
and sixteen parts of alcohol.—SUEC.—one
and a half of lavender and eight of alcohol.
—LUSIT.—one and a half of lavender and
ten of alcohol.—SARD.—one of lavender
and four of alcohol, and the product to
be poured upon fresh herb.—DAN.—one of
lavender and six of alcohol; and the pro-
duct to be poured upon fresh herb.—HER-
BIP.—one and a half of lavender, and ten of
alcohol; and the product to be poured upon
fresh herb.—FERR.—one and a half of herb
and eight of alcohol; six to be distilled.

R. Lavender, three parts.
Proof Spirit, six parts.
Water, thirty-six parts.

Distil nine parts.

HANN.; OLDEN.; POL.; BORUS.; and
SAX. prescribe one part of lavender, four
parts of alcohol, and four of water.—BA-
VAR.—one of lavender, four of alcohol
(0.900), and eight of water.—AUST.—one
of lavender, four and a half of alcohol
(0.910), and six of water.—FENN.—one and
a half of lavender, four of alcohol, and
three of water.—AMER.; and COXE—two
pounds of herb, eight pints of alcohol, and
a sufficient quantity of water.

R. Lavender, twelve parts.
Alcohol, six parts.
Water, two parts.

Distil six parts; re-distil the product with
twelve parts of lavender and two of water;
and draw off six parts again. (SWEDIAUR)

R. Lavender, half a pound.
Lavender Water, one pound.
Alcohol, four pounds.

Infuse for twenty-four hours, and distil
three pounds. (ROSS.)

COMPOUND SPIRIT OF LAVENDER.

*Tinctura Lavandulæ composita seu aromati-
ca, Spiritus Lavandula compositus, Alco-
hol cum Lavandula spica compositus.*
(LOND.; DUBL.; EDINB.; AMER.;
ANT.; DAN.; GALL.; FENN.; GENEV.;
HANN.; PALAT.; SUEC.; COXE; CA-
DET DE GASSICOURT; FEE; NIEMANN;
SWEDIAUR; VAN MONS)

R. Spirit of Lavender, three pints.
Rosemary, one pint.

Cinnamon, bruised, } of each,
Nutmegs, bruised, } half
an ounce.

Red Sanders Wood, one ounce.

Macerate for fourteen days, and
strain. (LOND.)

DUBL. gives the same formula,
adding two drams of cloves, and re-
quires only ten days' maceration.

R. Spirit of Lavender, three pounds.
Rosemary, one pound.

Cinnamon, bruised, one ounce.

Nutmegs, bruised, two drams.

Red Sanders Wood, three drams.

Macerate for seven days, and
strain. (EDINB.)

R. Spirit of Lavender, thirty-two parts,
Rosemary, sixteen parts.

Cinnamon, } of each, two parts.
Nutmegs, }

Infuse without heat for three days, and add
Red Sanders Wood, one part.

Macerate again for several days; then ex-
press and filter. (VAN MONS)

HANN. prescribes two pounds of spirit of
lavender, eight ounces of spirit of rose-
mary, two drams of cinnamon, as much of
nutmeg, and half an ounce of red sanders.
—NIEMANN; and FEE—three pounds of
spirit of lavender, one pound of spirit of
rosemary, half an ounce each, of cinnamon
and of nutmegs, and one ounce of red san-
ders.—PALAT.; and SUEC.—three pounds
of spirit of lavender, one pound of spirit
of rosemary, half an ounce each, of cinna-
mon and of nutmegs, and three drams of
red sanders.—CADET DE GASSICOURT—
three pounds of spirit of lavender, one
pound of spirit of rosemary, one ounce and
a half each, of cinnamon and of red sanders,
and one ounce of nutmegs.—VAN MONS.—
twelve ounces of spirit of rosemary, thirty-
six ounces of spirit of lavender, half an
ounce each, of cinnamon and of nutmegs,
and half a dram of cochineal.

- R. Spirit of Lavender, three pounds.
 ——— Rosemary, one pound.
 Cinnamon, } of each,
 Nutmegs, } half an ounce.
 Cochineal, bruised, half a dram.

Digest for three days in a moderate heat, and strain. (FENN.)

- R. Spirit of Lavender, two pounds.
 ——— Rosemary, six ounces.
 Powdered Cinnamon, } of each,
 Nutmegs, } half an ounce.
 Red Sanders, three drams.

Digest for several days; filter, and pour upon the residue,

- Spirit of Lavender, one pound.
 ——— Rosemary, six ounces.

Digest for three days, and mix the filtered tincture with the former. (ANT.)

- R. Spirit of Lavender, three pounds.
 ——— Sage, one pound.
 Cinnamon, one ounce.
 Nutmegs, half an ounce.
 Red Sanders Wood, three drams.

Infuse for eight days, and strain. (SWEDIAUR)

- R. Spirit of Lavender, three pounds.
 ——— Rosemary, one pound.
 Cinnamon, } of each,
 Nutmegs, } half an ounce.
 Yellow Sanders, }
 Ambergris, half a scruple.
 Saffron, one dram.

Macerate for eight days, and strain. (GENEV.)

- R. Spirit of Lavender, } of each,
 ——— Rosemary, } twenty-four ounces.
 Cinnamon, one ounce.
 Nutmegs, half an ounce.
 Cloves, two drams.
 Water, one hundred and twenty ounces.

Distil forty-eight ounces, and infuse in this product

Red Sanders Wood, three drams, and filter; or infuse the whole, without distilling, for several days, in the two spirits, and filter. (VAN MONS)

- R. Spirit of Lavender, eighty-four parts.
 ——— Rosemary, forty-eight parts.
 ——— Lemon, twelve parts.
 Cinnamon, three parts.
 Cubebs, } of each, one part.
 Cloves, }

Infuse for several days without heat, and filter. (VAN MONS)

AMER.; and COXE give the same as (LOND.)

- R. Tops of Lavender, twelve parts.
 ——— Rosemary, six parts.
 Cloves, } of each, three parts.
 Cinnamon, }

- Cubebs, one part.
 Proof Spirit,
 one hundred and forty-four parts.
 Water,
 two hundred and eighty-eight parts.

Distil one hundred and eight parts. (VAN MONS)

DAN. prescribes two pounds of lavender, one pound of rosemary, three ounces of fresh lemon peel and twelve pounds of alcohol; to be distilled to dryness, and one ounce of cubebs, one ounce of cloves, and three ounces of cinnamon, to be infused in the product.

It is stimulant.—Dose, from ten drops to two drams, in sweetened water.

ODORIFEROUS LIQUOR.

Aqua odorifera. (WIRTEM.; NIEMANN)

- R. Herb of Rosemary, } of each,
 ——— Marjoram, } one ounce.
 Flowers of Lavender, }
 ——— Lily of the Valley, } of each,
 ——— Roses, } two ounces.
 Rhodium Wood, } of each,
 Fresh Lemon Peel, } one ounce and a half.
 Cinnamon, two ounces.
 Cloves, one ounce and a half.
 Alcohol, eight pounds.
 Water, four pounds.

Digest for three days, and distil three-fourths of it. (WIRTEM.)

- R. Lavender, } of each, two pounds.
 Balm, }
 Root of Sweet Flag, one pound.
 Cloves, } of each,
 Herb of Rosemary, } half a pound.
 ——— Marjoram, } of each,
 ——— Peppermint, } one ounce.
 Cinnamon, }
 Alcohol, two pounds.
 Water, thirty pounds.

Distil twenty ounces. (NIEMANN)

This aromatic liquor differs very little from the *Eau des Carmes* and *Eau de Cologne*. It is used for the same purposes.

2. *Lavandula Stoechas*, LINN.—French Lavender.

SYNONIMES:—*Stoechas* d'Arabie, FR.; *Stoechas*, GERM.; DAN.; DUT.; and SWED.; *Cantnosa*, SPAN.; *Stecade*, ITAL.; *Rosmarinho*, PORT.

AMST.; ANT.; BATAV.; BRUNS.; HISP.; FERR.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

A small shrub of the middle of Europe. (*Fig. Zorn, Ic. pl. t. 485.*)

The flowers (*Flores Stæchadis purpureæ Arabicæ*) are employed. They are of a deep purple colour, crowded into an oblong oval spike, which is surmounted by a bunch of coloured leaves. They have an agreeable and camphorous odour, particularly when rubbed. Their taste is aromatic and bitter.

SYRUP OF FRENCH LAVENDER.

Syrupus de Stæchade. (GALL.)

- R. French Lavender, one ounce.
Distilled Water of French Lavender, two pounds.

Digest in a water-bath for two hours, and add to the strained liquor

White Sugar, twice the quantity.

Dissolve in a water-bath, and strain through flannel.—Dose, from half a dram to two drams.

COMPOUND SYRUP OF FRENCH LAVENDER.

Syrupus de Stæchade seu de Stæchade compositus. (GALL.; WIRTEM.; VAN MONS)

- R. Cinnamon, one part.
Ginger, } of each, two parts.
Sweet Flag, }
Fresh Leaves of Fennel, }
Rue, } of each,
Rosemary, } eight
Sage, } parts.
Thyme, }
French Lavender, }
Water, five hundred parts.

Distil seventy parts, with which, and

White Sugar, one hundred and twenty-eight parts.

Make a syrup in a water-bath.

- R. Dried Spikes of French Lavender, three ounces.
Flowering Tops of Thyme, four ounces and a half.
Tops of Sage, } of each,
Rosemary, } six drams.
Seeds of Rue, } of each,
Fennel, } four drams and a half.
Cinnamon, } of each, two drams.
Ginger, }
Sweet Flag, }
Water, eight pounds.

Macerate for two days in a covered vessel, then distil half a pound of liquid, and add to it

White Sugar, one pound.

Strain the residue with expression, and add to the liquor

White Sugar, four pounds.

Make a syrup, and mix it with the former when half cold. (GALL.)

- R. French Lavender, two ounces.
Flowers of Rosemary, two drams.
Mountain Calamint, } of each,
Origanum, } three drams.
Thyme, }
Betony, } of each, two drams.
Sage, }
Seeds of Anise, } of each,
Peony, } two drams and a half.
Sweet Flag, } of each, one dram.
Cinnamon, }
Ginger, }
Boiling Water, one pound.

Macerate for two days in a warm place; then express, and add to nine ounces of the liquor

White Sugar, sixteen ounces.

Boil, and strain. (WIRTEM.)

Dose, from half a dram to two drams.

LAURUS.

Five species of this genus are used in medicine:

1. *Laurus Cinnamomum*, LINN.—Cinnamon.

SYNONIMES:—*Cinnamomum Canella*, Κίναμον.—Cannelle, Cannelle de Ceylon, FR.—Zimmt, Zimmet, GERM.; Darsini, ARAB.; Skorice, BOHEM.; Kerrundu, CYNG.; Caneel, DAN.; Kalmie darchinie, DUK.; Canella, ITAL.; PORT.; and SPAN.; Darchinie, HIND.; and PERS.; Kaneel, DUT.; Kaimanis, MAL.; Cyna-

mon prawdziwy, POL.; Darasita, SANS.; Kanel, SWED.; Karruwa puttay, TAM.; Sanalinga putta, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; GENÈV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BRIGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PARIS.; SPIELMANN.; ZARDA.

This tree is originally of Ceylon. (*Enneandria Monogynia*, LINN.; *Laurineæ*, JUSS.; *Fig. Flore Medic. II. 92.*)

The bark (*Cortex Cinnamomi veri seu acuti seu Cannella Zeylanica*) is employed. It is smooth, thin, flexible, papyraceous, rolled in long quills of about the diameter of the finger, and containing other smaller rolls; it is ligneous, and exhibits a splintery fracture. It is of a reddish-yellow or fawn colour; its odour very agreeable, and its taste aromatic, rather acrid, sweet, and mixed with a slight astringency.

It contains a very acrid volatile oil, much tannin, an azoted colouring matter, an acid, mucilage, and starch.

It is a powerful stimulant.—Dose, from ten grains to a scruple of the powder.

I. Preparations which contain Cinnamon in substance.

AROMATIC NUTRITIVE POWDER. (NIEMANN; SPIELMANN)

R. Vanilla,	one part.
Cloves,	two parts.
Cinnamon,	four parts.
Rice,	sixty-four parts.
White Sugar,	ninety-six parts.

Pulverize each separately, and mix. (VAN MONS)

NIEMANN; and SPIELMANN prescribe one of cloves, two of vanilla, twenty-four of cinnamon, one hundred and twenty-eight of rice, and one hundred and ninety-two of sugar.

This powder is taken as chocolate.

COMPOUND POWDER OF CINNAMON; AROMATIC POWDER.

Pulvis aromaticus seu Cinnamomi compositus, Pulvis seu Species Imperatoris, Pulvis Vitæ Imperatoris, Pulvis Diaromaton, Species aromatica seu diacinnamomi seu diambra seu latifiantes Rhazis, Tragea aromatica, Species aromatica rosata. (LOND.; EDINB.; DUBL.; AMER.; BATAV.; BAVAR.; BELG.; BRUNS.; FULD.; GENEV.; HANN.; LUSIT.; PALAT.; POL.; BORUS.; SARD.; WIRTEM.; BRUGNATELLI; COKE; CADET DE GASSICOURT; HUFELAND; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Pulvis Cinnamomi compositus. (LOND.)

R. Cinnamon,	two ounces.
Cardamom Seeds,	one ounce and a half.

Ginger, one ounce.
Long Pepper, half an ounce.
Reduce them to a very fine powder.

Pulvis aromaticus.

R. Cinnamon,	} of each, an equal part.
Cardamom Seeds,	
Ginger,	
Reduce them to a fine powder, to be kept in a well-stopped bottle. (EDINB.)	

R. Cinnamon,	two ounces.
Lesser Cardamom	} of each, one dram.
Seeds, husked,	
Ginger,	
Long Pepper,	
Reduce them to fine powder. (DUBL.)	

AMER.; BATAV.; BELG.; and COXE give the same formula as EDINB.—BRUGNATELLI; and VAN MONS—the same as LOND.—GENEV.; HANN.; and BORUS.—the same as DUBL.—BAVAR. prescribes two parts of cinnamon, and one each of cardamoms and of ginger.

R. Cinnamon,	}	one part.
Pichurim Beans,		of each,
Lesser Cardamoms,		two parts.
Make a powder. (LUSIT.)		

R. Cinnamon,	}	of each,
Lesser Cardamoms,		two ounces.
Ginger,	}	of each, half an ounce.
Cloves,		

Make a powder. (PoL.)

R. Cinnamon,	two ounces.
Cardamoms,	} of each, one ounce.
Ginger,	
Nutmegs,	

Make a powder. (CADET DE GASSICOURT; SWEDIAUR)—This formula bears the title of *Pulvis diaromaton*.

AMST. requires an equal part each of the four substances.

R.	Root of Elecampane,	}	of each, one ounce.
	—— Sweet Flag,		
	—— Ginger,		
	Anise,		
	Orange Peel,		
	Black Pepper,	}	of each, half an ounce.
	Cloves,		
	Cinnamon,		
Make a powder. (HUFELAND)			

R. Cinnamon,	ten drams.
Ginger,	} of each, half an ounce.
Cloves,	

Galanga, }
Mace, } of each, two drams.
Nutmegs, }

Make a powder. (BRUNS.; PALAT.; SARD.; WIRTEM.; SPIELMANN)—This powder bears the title of *Pulvis seu Species Imperatoris*.

PIDERIT prescribes one ounce of galanga, ten drams of cinnamon, half an ounce each, of cardamoms and of cloves, and two drams of mace.

This formula resembles the three following: they bear the name of *Species aromatica rosata*.

R. Liquorice, seven drams.
Galanga, one dram.
Aloe Wood, } of each,
Yellow Sanders Wood, } two drams.
Cinnamon, six drams.
Cloves, } of each,
Mace, } two drams and a half.
Nutmegs, } of each,
Lesser Cardamoms, } one dram.
Petals of Roses, two ounces.

Make a powder. (PALAT.)

R. Cinnamon, one ounce.
Cloves, }
Ginger, } of each, three drams.
Nutmegs, }
Mace, two drams.
Red Sanders Wood, one ounce.
White Sugar, thirty ounces.

Make a powder. (PALAT.; WIRTEM.)

These are called *completæ* when to every two ounces are added

Ambergris, eight grains.
Musk, three grains.

R. Petals of Red Roses, one ounce.
Liquorice, one ounce and a half.
Galanga, half an ounce.
Aloe Wood, } of each, one
Yellow Sanders Wood, } dram and a half.
Cinnamon, two drams and a half.
Cloves, }
Mace, } of each,
Gum Arabic, } four scruples.
Tragacanth, }
Nutmegs, } of each,
Lesser Cardamoms, } half a dram.
Spikenard, half a scruple.

Make a powder. (SPIELMANN)

WIRTEM. also prescribes one dram of cinnamon, half a dram of ginger, fifteen grains each, of cloves, of galanga, of grains of paradise, of mace, and of nutmeg; half a dram each, of white and of red sanders, and eight ounces of white sugar.—SPIELMANN—two drams of cinnamon, one dram of

ginger, half a dram each, of cloves, of lesser galanga, of grains of paradise, of mace, and of nutmeg; six drams of red sanders, and sixteen ounces of sugar.

To this formula belongs the title of *Pulvis seu Tragea aromatica*; the second of WIRTEM. bears that of *Tragea aromatica Gabelchoveri*.

R. Root of Elecampane, half an ounce.
Galanga, one ounce.
Ginger, }
Cloves, }
Lesser Cardamoms, } of each,
Long Pepper, } three drams.
Mace, }
Nutmegs, }
Aloe Wood, }
Saffron, one dram.
Cassia Lignea, half an ounce.
Cinnamon, two ounces.

Make a powder. (WIRTEM.)—This powder is called *Species diacinnamomi*.

R. Root of Doronica, } of each,
Galanga, } one dram.
Ginger, } and a half.
Spikenard, one dram.
Aloe Wood, }
Yellow Sanders Wood, } of each,
Long Pepper, } two drams.
Cloves, }
Nutmegs, } of each, three drams.
Mace, }
Lesser Cardamoms, } of each,
Great Cardamoms, } one dram.
Cinnamon, three drams.

Make a powder. (BRUNS.; WIRTEM.)

This formula gives the *Species diambra*; they are called *completæ*, when half a dram of musk and half a dram of ambergris are added.

The following formula gives the *Species aromatica caryophyllatæ*:

R. Root of Liquorice, two drams.
Zedoary, }
Galanga, }
Aloe Wood, }
Yellow Sanders Wood, } of each,
Cinnamon, } one dram.
Mace, }
Long Pepper, }
Cardamoms, }
Cubebs, two scruples.
Petals of Roses, half an ounce.
Cloves, one ounce.

Make a powder. (WIRTEM.)

These species are called *completæ*, when to every ounce are added

Ambergris, eight grains.
Musk, half a grain.

R. Root of White Behen,	} of each, half an ounce.
—— Red Behen,	
—— Zedoary,	
—— Doronica,	
Herb of Balm,	
Seeds of Sweet Basil,	
—— Peony,	
Saffron,	
Cloves,	
Lesser Cardamoms,	
Cinnamon,	
Lemon Peel,	
Mastic,	

Make a powder. (WIRTEM.)—These species are called *Rhazes' Species latifoliantes*; they are called *completæ*, when half an ounce of nutmegs and a scruple of musk are added.

R. Root of Zedoary,	} of each, two drams.
—— Elecampane,	
—— Sweet Flag,	
—— Burnet Saxi-	
frage,	
—— Galanga,	} of each, one dram.
—— Avens,	
—— Liquorice,	
Herb of Houndstongue,	} of each, one dram.
—— Speedwell,	
—— Blessed Thistle,	
Tops of Lesser Centaury,	
Leaves of Senna, one ounce and a half.	
Seeds of Anise,	} of each, one dram and a half.
—— Fennel,	
—— Caraway,	
—— Parsley,	
Juniper Berries,	three drams.
Cinnamon,	one dram.
Cubebs,	of each,
Lesser Cardamoms,	four scruples.
Mace,	of each,
Burnt Hartshorn,	two drams.
Pontic Rhubarb,	half an ounce.
Turkey Rhubarb,	one dram and a half.
Super-tartrate of Potass,	three drams.
Sugar Candy,	four ounces and a half.

Make a powder. (WIRTEM.)

This last formula, which is designated by the title of *Pulvis Vitæ Imperatoris*, should have been placed in the article rhubarb, were it not for the evident analogy which, by insensible degrees, the formulæ which precede it establish between it and those in which cinnamon is the principal ingredient.

In general the dose of the first powders is from five to fifteen grains. For the latter, which are much less stimulant, and indeed now almost generally disused, the dose may be carried as far as one or two drams.

AROMATIC LOZENGES.

Morsuli seu Trochisci aromatici seu stomachici seu Imperatoris seu aromatico-stomachici. (BRUNS.; FENN.; FULD.;

LIPP.; PALAT.; SAX.; SUEC.; WIRTEM.; NIEMANN; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. White Sugar,	sixteen ounces.
Water,	four ounces.

Boil, and add a mixture of

Chopped Almonds,	two ounces.
—— Lemon Peel,	half an ounce.
Powdered Cinnamon,	} of each, one dram.
—— Nutmegs,	
—— Ginger,	
—— Cardamoms,	
—— Galanga,	
—— Cloves,	

Make lozenges. (LIPP.; NIEMANN)

R. Sweet Almonds,	two ounces.
Bitter Almonds,	} of each, one ounce.
Oleosaccharum of Lemon,	
—— Orange,	
Powdered Cinnamon,	one dram.
—— Cloves,	} of each, half a dram.
—— Ginger,	
—— Nutmegs,	
Mucilage of Gum Arabic,	a sufficient quantity.

Make trochises of two drams each. (SWEDIAUR)

R. Chopped Sweet Almonds,	eight parts.
—— Lemon Peel,	two parts
Powdered Cinnamon,	} of each, one part.
—— Ginger,	
Sugar, boiled <i>à la plume</i> ,	thirty-two parts.

Mix. (SAX.)

NIEMANN prescribes one pound of sugar, boiled *à la plume* with half a pound of rose water, one ounce each, of almonds, of lemon peel, and of cinnamon, and two drams of ginger.—SUEC.—one pound of sugar, a sufficient quantity of water, one ounce each, of almonds and of orange peel, half an ounce of cinnamon, and one scruple of ginger.—FENN.—half a pound of sugar, one ounce and a half of almonds, six drams of orange peel, half an ounce of cinnamon, and one scruple of ginger.

R. Preserved Orange Peel,	} of each, one ounce.
—— Lemon Peel,	
—— Pulp,	
Sweet Almonds,	one ounce and a half.
Aromatic Powder,	one ounce.
Sugar, boiled <i>à la plume</i> ,	sixteen ounces.

Make lozenges. (SPIELMANN)

R. Cinnamon,	two drams.
Aromatic Powder,	six drams.
Preserved Orange Peel,	} of each, one ounce.
—— Lemon Peel,	
—— Pulp,	

Sweet Almonds, one ounce and a half.
Rose Sugar, boiled à la plume,
sixteen ounces.

Make lozenges. (BRUNS.; WIRTEM.)

PIDERIT prescribes sixteen ounces of sugar boiled in eight of water, one ounce and a half of lemon juice, two ounces of almonds, one ounce of fresh orange peel, one ounce of preserved lemon peel, and three drams of aromatic powder.

R. Pine Apple Kernels, } of each,
Pistachio Nuts, } half an ounce.
Sweet Almonds, }
Aromatic Powder, three drams.
Cinnamon, one dram.
Conserve of Roses, half an ounce.
Sugar, dissolved in Rose Water, eight ounces.

Make lozenges. (PALAT.)

R. White Sugar, sixteen ounces.
Water, eight ounces.

Boil à la grande plume, and add

Chopped Sweet Almonds, two ounces.
Preserved Lemon Peel, } of each,
mon Peel } half
Lesser Galanga, } an ounce.

Make lozenges. (FULD.)

R. Powdered Cinnamon, } of each,
Preserved Lemon Peel, } one part.
Orange Peel, }
Blanched Sweet Almonds, two parts.
Sugar, dissolved, twelve parts.

Make lozenges. (VAN MONS)

AROMATIC ELECTUARY.

Electuarium aromaticum seu cardiacum, Confectio aromatica seu cordialis. (LOND.; DUBL.; EDINB.; AMER.; LUSIT.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS)

Confectio aromatica. (LOND.)

R. Cinnamon, } of each,
Nutmegs, } two ounces.
Cloves, one ounce.
Cardamom Seeds, half an ounce.
Saffron, two ounces.
Prepared Shells, sixteen ounces.
Refined Sugar, two pounds.
Water, one pint.

Reduce the dry ingredients together to a fine powder; then add the water gradually, and incorporate thoroughly.

Electuarium aromaticum. (EDINB.; AMER.; COXE)

R. Aromatic Powder, one part.
Syrup of Orange, two parts:

Mix, and incorporate them well together.

Electuarium aromaticum. (DUBL.)

R. Cinnamon, } of each,
Nutmegs, } half an ounce.
Purified Sugar, } of each,
Saffron, } one ounce.
Lesser Cardamom } of each,
Seeds, } two drams.
Cloves, }
Prepared Chalk, two ounces.
Syrup of Orange, a sufficient quantity.

Pulverize the dry ingredients together, and then incorporate the syrup.

R. Aromatic Powder, two parts.
Conserve of Orange Peel, three parts.
Syrup of Orange Peel, a sufficient quantity.

Mix. (LUSIT.)

R. Conserve of Orange Peel, three ounces.
Nutmegs, one ounce and a half.
Ginger, six drams.
Cinnamon, half an ounce.
Syrup of Orange Peel, a sufficient quantity.

Mix. (SWEDIAUR)

R. Conserve of Orange Peel, three ounces.
Cinnamon, } of each, half an ounce.
Nutmegs, }
Saffron, } of each, two drams.
Ginger, }
Sugar, one ounce.
Syrup of Orange Peel, a sufficient quantity.

Mix. (BRUGNATELLI)

R. Cinnamon, } of each, four parts.
Nutmegs, }
Saffron, }
Cardamoms, one part.
Cloves, two parts.
Prepared Chalk, } of each,
White Sugar, } thirty-two parts.
Simple Syrup, a sufficient quantity.

Mix. (VAN MONS)

Dose, twenty grains.—It is employed generally as a vehicle for more active substances.

COMMON CONFECTION.

Confectio communis. (GENEV.)

- R. Lemnian Bole, } of each,
 Crab's Eyes, } three ounces.
 Cinnamon, } one ounce.
 Dittany of Crete, } of each,
 Yellow Sanders, } half an ounce.
 Myrrh, } two drams.
 Saffron, } half an ounce.
 Syrup of Lemon, } two ounces.
 Honey, } two ounces and a half.
- Mix.

AROMATIC PLASTER.

Emplastrum aromaticum. (DUBL.; COXE;
 NIEMANN)

- R. Frankincense, three ounces.
 Yellow Wax, half an ounce.
 Cinnamon in powder, six drams.
 Oil of Pimento, } of each,
 — Lemon, } two drams.

Melt the two first together, and strain; when half cold add the others, mixed together, and form a plaster.

II. *Preparations which contain the active Principles of Cinnamon extracted by Distillation.*

ESSENTIAL OIL OF CINNAMON.

Oleum Cinnamomi athereum, Ætheroleum Cinnamomi.

- 1 That which is met with in Commerce. (LOND.; OLDEN.)
- 2 Prepared according to various Formulæ. (ANT.; AUSTR.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FULD.; HANN.; HASS.; LIPP.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; PIDERIT; SWEDIAUR)

- R. Cinnamon, in coarse powder, any quantity.
 Water, a sufficient quantity.
 Distil, and separate the oil which sinks to the bottom of the receiver. (FENN.; LUSIT.; ROSS.; SUEC.)

HERBIP.; and SWEDIAUR prescribe one part of cinnamon and four parts of water.—DAN.; FULD.; and SAX.—one of cinnamon and six of water.—AUSTR.; HANN.; POL.; and BORUS.—one of cinnamon and eight of water.—BAVAR.—one of cinnamon and nine of water.—ANT.—three of cinnamon and ten of water.—HISP.—one of cinnamon and forty of water.

- R. Bruised Cinnamon, ten parts.
 Muriate of Soda, one part.
 Water, twenty parts.

Macerate for twelve parts, and distil. (GALL.)

BRUNS.; HASS.; PALAT.; WIRTEM.; and PIDERIT prescribe two pounds of cinnamon, four ounces of muriate, and sixteen pounds of water.—SARD.—one pound of cinnamon, three ounces of muriate, and eight pounds of water.

- R. Cinnamon, in coarse powder, two pounds.
 Alcohol, eight ounces.

Macerate for two days, and add

- Muriate of Soda, one pound.
 Water, twelve pounds.

Distil. (LIPP.)

It is stimulant and nervine.—Dose, one or two drops.

OLEOSACCHARUM OF CINNAMON.

Oleosaccharum Cinnamomi. (AUSTR.; BRUNS.; DAN.; GALL.; FENN.; HANN.; OLDEN.; PALAT.; BORUS.; SUEC.; WIRTEM.; SWEDIAUR)

- R. White Sugar, one ounce.
 Oil of Cinnamon, eight drops.

Triturate together. (DAN.)

HANN.; OLDEN.; and BORUS. prescribe one ounce of sugar and twenty-four drops of oil.—AUST.—one dram of sugar and three grains of oil.—SWEDIAUR—one dram of sugar and one or two drops of oil.—GALL.—two drams of sugar and two drops of oil.—WIRTEM.; BRUNS.; FENN.; and SUEC.—one ounce of sugar and sixteen drops of oil.

- R. White Sugar, one ounce.
 Red Sulphuret of Mercury, two drams.
 Oil of Cinnamon, fifteen drops.

Triturate together. (PALAT.)

BALSAM OF CINNAMON.

Balsamum Cinnamomi. (WIRTEM.; SPIELMANN)

- R. Oil of Cinnamon, one dram.
 Expressed Oil of Nutmeg, two drams and a half.

Mix, and colour with Armenian bole. (WIRTEM.)

SPIELMANN prescribes one part of oil of cinnamon and five parts of oil of nutmeg.

AROMATIC CONFECTION.

Confectio Cinnamomi regia. (SPIELMANN)

- R. Chocolate, four ounces.
 Preserved Areca Nuts, three.
 Triturate together, and add
 Alkermes Confection, one ounce.
 Oil of Cinnamon, half a dram.

Syrup of Cinnamon, six ounces.
Mix.—It is stomachic.—Dose, one dram.

CARDIAC MIXTURE. (SWEDIAUR)

R. Oil of Cinnamon, three drops.
White Sugar, half an ounce.

Add, while tritulating,
Yolks of Eggs, two.
White Wine, six ounces.

Mix.—Dose, two or three spoonfuls every six hours.

OLEO-BALSAMIC MIXTURE.

Mixtura oleoso-balsamica. (BRUNS. ;
HANN. ; PALAT. ; POL. ; BORUS. ; WIR-
TEM. ; NIEMANN ; SPIELMANN)

R. Oil of Cinnamon,	} of each, one scruple.
— Lavender,	
— Marjoram,	
— Cloves,	
— Lemon,	
— Mace,	} of each, half a scruple.
— Rue,	
— Amber,	
Ambergris,	
Balsam of Peru,	
Alcohol,	ten ounces.

Dissolve. (BRUNS. ; PALAT. ; WIRTEM.)

R. Oil of Lemon,	four scruples.
— Bergamotte,	twelve drops.
— Cinnamon,	} of each, eighteen drops.
— Mint,	
— Cardamoms,	} of each, twelve drops.
— Cloves,	
— Moldavian Mint,	
— Mace,	
— Rosemary,	
— Lavender,	} of each, fifteen drops.
— Rue,	
— Sage,	

Concentrated Alcohol, six ounces.

Colour with alkanet, and add when filtered,

Essence of Amber, half a dram.

Mix. (SPIELMANN)

Dose, to twenty drops.

SIMPLE CINNAMON WATER.

Aqua Cinnamomi seu Cinnamomi simplex seu sine Vino. (LOND. ; EDINB. ; DUBL. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; GALL. ; FERR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SARD. ; WIRTEM. ; BRUGNATELLI ; COXE ; PIDERIT ; SPIELMANN ; SWEDIAUR ; VAN MONS)

Aqua Cinnamomi. (LOND. ; and
DUBL.)

R. Cinnamon, bruised, one pound.
Water, one pint.

Macerate for twenty-four hours ; then add a sufficient quantity of water, that, after the distillation, empyreuma may be prevented. Distil one gallon.

BRUNS. ; FERR. ; and ROSS. prescribe one part of cinnamon and twelve parts of water ; to be digested for three days, and two-thirds to be distilled.—AMER.—the same as LOND.—BORUS.—one pound of cinnamon and a sufficient quantity of water ; nine pounds to be distilled.—AMST. ; BATAV. ; BELG. ; and SWEDIAUR—one pound of cinnamon and a sufficient quantity of water ; to be macerated for twenty-four hours, and ten pounds distilled.—ANT.—half a pound of cinnamon and eight pounds of water ; to be macerated for twenty-four hours, and four pounds distilled.—SARD.—two ounces of cinnamon and one pound of water ; to be macerated for three days, and then distilled till the liquor begins to whiten.—HANN. ; OLDEN. ; and POL.—one pound of cinnamon and a sufficient quantity of water, nine pounds to be distilled.—SPIELMANN—four ounces of cinnamon and forty-eight of water ; to be macerated for three days and thirty-six ounces distilled.—FULD.—nine ounces of cinnamon and six pounds of water ; to be digested for twenty-four hours, and three pounds distilled.—AUSTR. ; HASS. ; LIPP. ; and PIDERIT—one part of cinnamon and a sufficient quantity of water, to be macerated for twenty-four hours, and six parts distilled.—GENEV.—one pound of cinnamon and a sufficient quantity of water ; four pounds to be distilled.—BRUGNATELLI—one part of cinnamon and six parts of water ; two thirds to be distilled.—DAN.—one part of cinnamon and nine parts of water ; to be macerated for two days, and six to be distilled.—BAVAR.—one of cinnamon and sixteen of water ; to be macerated for twenty-four hours, and eight distilled.—SAX.—one of cinnamon and twelve of water ; to be macerated for twenty-four hours, and nine distilled.—LUSIT.—one of cinnamon and twelve of water ; to be macerated for twenty-four hours, and eight distilled.—GALL.—two of cinnamon and sixteen of water ; to be macerated for twelve hours and eight distilled.—VAN MONS—one of cinnamon and twenty-four of water ; eight to be distilled.

Aqua Lauri Cinnamomi. (EDINB.)

R. Cinnamon bark, bruised,
one pound.
Water, a sufficient quantity
to prevent empyreuma. Distil off
one gallon.

R. Cinnamon, eight ounces.
Cherry Water, twelve pounds.

Macerate for three days, and distil two-thirds. (WIRTEM.; SPIELMANN)

R. Cinnamon, four ounces.
Decoction of Barley,
forty-eight ounces.

Macerate for three days, and distil thirty-six ounces. (SPIELMANN)

This formula gives the Hordeated Cinnamon Water, *Aqua Cinnamomi hordeata*.

R. Cinnamon, eight ounces.
Water of the four Cordial Flowers,
twelve pounds.

Macerate for three days, and distil two thirds. (WIRTEM.; SPIELMANN)

The product of this bears the name of Cordial Cinnamon Water *Aqua Cinnamomi cordialis*.

R. Cinnamon, four ounces.
Water of Bugloss, three pounds.

Macerate for three days, and distil two-thirds. (WIRTEM.; SPIELMANN)

R. Cinnamon, one pound.
Root of Bugloss, } of each, one ounce.
Leaves of Borage, }
Balm, }
Water of Bugloss, six pounds.
Roses, } of each,
Violet, } two pounds.

Digest, and distil in a water-bath. (PALAT.)

This formula and the preceding gives the Buglossated Cinnamon Water, *Aqua Cinnamomi buglossata*.

Dose, from one to two ounces or more.

COLD CORDIAL WATER. (WIRTEM.)

R. Cinnamon, two ounces and a half.
Peels of Fresh Lemons, eight.
Yellow Sander's Wood, } of each,
Bugloss Water, } one pound.
Sorrel Water, }
Cherry Water, } of each,
Rose Water, } one pound and
Raspberry Water, } a half.

Distil two thirds.—Dose, one or two ounces.

COMFORTING WATER.

Aqua comfortiva. (PALAT.)

R. Buglossated Cinnamon Water,
one ounce and a half.

Violet Water, }
Rose Water, } of each, six ounces.
Borage Water, }
Cherry Water, }

Mix.—It is stimulant and cordial.—Dose, a spoonful.

WATER OF LOHOCH SANUM ET EXPERTUM. (BRUNS.; PALAT.; WIRTEM.)

R. Root of Florentine Iris, }
Liquorice, } of each,
Herb of Hyssop, } two
Maidenhair, } ounces.
Mountain Calamint, }
Seeds of Anise, } of each,
Fennel, } one ounce.
Cinnamon, two ounces.
Spring Water, ten pounds.

Digest for three days, and distil till only four pounds remain in the alembic.—Dose, from one ounce and a half to two ounces.

SPIRIT OF CINNAMON, VINOUS CINNAMON WATER.

Aqua Cinnamomi vinosa, seu spirituosa, Spiritus Cinnamomi. (LOND.; EDINB.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BRUNS.; DAN.; HISP.; FENN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; COXE.; PIDERIT.; SPIELMANN.; VAN MONS.)

Spiritus Cinnamomi. (LOND.)

R. Oil of Cinnamon, by weight,
five scruples.

Rectified Spirit,
four pints and a half.

Mix them together, add as much water as will be sufficient to prevent empyreuma, and distil slowly one gallon. (DUBL.)

Spiritus Lauri Cinnamomi. (EDINB.)

R. Cinnamon, bruised, one pound.
Proof Spirit, nine pounds.

Macerate for two days in a close vessel; then add a sufficient quantity of water to prevent empyreuma, and distil nine pounds.

R. Cinnamon, one part.
Alcohol, six parts.

Distil; and re-distil the product with one part of cinnamon. (HERBIP.)

R. Cinnamon, six ounces.
 Alcohol, six pounds.
 Macerate for twenty-four hours, and add
 Water, six pounds.
 Distil six pounds. (LIPP.)

HISP. directs eight ounces of cinnamon to be macerated for one day with three pounds of water and three ounces of alcohol; then one pound of water to be added, and three pounds of spirit distilled.—HANN.; OLDEN.; POL.; BORUS.; SAX.; and FEE—one part of cinnamon to be macerated in sixteen parts of water and two of alcohol; and nine to be distilled.—FENN.—one part of cinnamon to be macerated in two parts of alcohol and a sufficient quantity of water; six to be distilled.—VAN MONS—one of cinnamon to be macerated in four of proof spirit and twenty of water; eight to be distilled.—BAVAR.—one of cinnamon to be macerated for twenty-four hours in sixteen of water and two of alcohol (0.910); eight to be distilled.—AMST.; and BATAV.—one and a half of cinnamon to be macerated for three days in eight of proof spirit; one half to be distilled.—LUSIT.—one of cinnamon to be macerated in ten of water for ten days, then four of water to be added; ten to be distilled.

R. Cinnamon, one pound.
 Alcohol, } of each,
 Balm Water, } four pounds.

Macerate for two days, and distil. (SARD.)

R. Cinnamon, one pound.
 Proof Spirit, two pounds.
 Water, a sufficient quantity.

Distil six pounds, and add to the product,
 Sugar, one ounce and a half.

Dissolve. (SUEC.)

HASS.; and PIDERIT prescribe one pound of cinnamon, two pounds of proof spirit and two of water; six pounds to be distilled; two pounds of water and two ounces of sugar to be added to the product.

R. Simple Cinnamon } of each,
 Water, } an equal part.
 Alcohol, }

Mix. (DAN.)

R. Cinnamon, bruised, four ounces.
 White Wine, six pounds.

Digest for three days in a covered vessel, and distil one half. (SARD.)

HERBIP. prescribes one part of cinnamon, eight parts of wine, and two days' maceration.—SPIELMANN—one of cinnamon, three of wine, and a few days' maceration.—HISP.—one of cinnamon, six of wine, and three days' maceration; two parts to be distilled.—PALAT.; and WIRTEM.—nine ounces of cinnamon and twelve pounds of wine, one half to be distilled.

R. Cinnamon, six ounces.
 White Wine, one pound.
 River Water, five pounds.

Macerate for twenty-four hours, and distil one half. (ROSS.)

BRUNS. requires sixteen ounces of cinnamon, twelve pounds of white wine, and two pounds and a half of alcohol; to be macerated for two days, and ten pounds to be distilled.

R. Cinnamon, four ounces.
 Juice of Quince, become vinous
 by fermentation, three pounds.

Infuse for three days, and distil two-thirds. (WIRTEM.; SPIELMANN)

BRUNS. prescribes six ounces of cinnamon and three of juice.—PALAT.—two ounces of cinnamon and three pounds of juice.—The product bears the name of Cydoniated Cinnamon Water, *Aqua Cinnamonomi cydoniata*.

R. Simple Cinnamon Water, two parts.
 Tincture of Cinnamon, one part.

Mix. (LIPP.)

Among so many variations in the mode of making this preparation, no general dose can be assigned, as it varies with almost every formula. It might very well be rejected; for the tincture of cinnamon would answer all its purposes with equal advantage.

VINOUS POTION. (RATIER)

R. Spirituous Cinnamon Water,
 two drams.

Red Wine, four ounces.
 Syrup of Clove Pink, one ounce.

Mix.—It is excitant, stomachic, and cordial.

AROMATIC SPIRIT.

Aqua aromatica, Spiritus aromaticus seu alexiterius, Alcohol aromaticum. (BAVAR.; BELG.; DAN.; HANN.; OLDEN.; PALAT.; BORUS.; SAX.; PIDERIT; SPIELMANN; SWEDIAUR)

R. Cinnamon, } of each, one part.
 Cloves, }

Nutmegs, } of each, two parts.
 Coriander, }

Fresh Lemon Peel, four parts.

Dried Leaves of Balm, six parts.

Alcohol (0.900), seventy-two parts.

Water, thirty-six parts.

Macerate for three days, distil seventy-two parts and rectify the product, re-distilling sixty parts. (BAVAR.)

R. Cinnamon, } of each,
 Nutmegs, } half an ounce.
 Cloves, }
 Marjoram, }

Coriander,	one ounce.
Proof Spirit,	thirty-two ounces.
Water,	six ounces.

Macerate for two days, and distil twenty ounces. (BELG.)

R. Cinnamon,	two ounces.
Lemon Peel,	of each,
Fennel,	one ounce.
Mace,	half an ounce.
Peppermint,	three ounces.
Proof Spirit,	twelve pounds.

Distil eight pounds. (DAN.)

SPIELMANN employs angelica seeds instead of fennel.

R. Cinnamon,	two ounces.
Fresh Lemon Peel,	one ounce.
Cloves,	half an ounce.
Jamaica Pepper,	two drams.
Alcohol,	of each,
Spring Water,	eight pounds.

Digest for two days; then distil eight pounds of spirit, dilute it with four pounds of water, and add eight ounces of white sugar. (PIDERIT)

R. Cinnamon,	
Unripe Oranges,	of each,
Herb of Peppermint,	two parts.
Leaves of Rosemary,	
Flowers of Lavender,	
Lesser Galanga,	of each, one part.
Sweet Flag,	
Fennel,	
Cloves,	one half part.
Alcohol,	of each,
Water,	fifty-two parts.

Distil one half, after three days' maceration. (SAX.)

R. Herb of Sage,	eight ounces.
— Rosemary,	of each,
— Peppermint,	four ounces.
Flowers of Lavender,	
Fennel,	of each, two ounces.
Cinnamon,	
Alcohol,	four ounces.
Water,	a sufficient quantity.

Distil twelve pounds after twenty-four hours' maceration. (HANN.; OLDEN.; POL.; BORUS.; SWEDIAUR)

CATARRHAL LIQUOR. (VAN MONS)

R. Cinnamon,	
Leaves of Rosemary,	of each,
— Sage,	two ounces.
Flowers of Lavender,	
Cloves,	
Nutmegs,	
Proof Spirit,	six pounds.
Water,	twelve pounds.

Distil till the product begins to whiten.

VOL. II.

EAU-DE-VIE DES FEMMES.

Aqua Vitæ Mulierum Bæcleri. (SPIELMANN)

R. Root of Florentine Iris,	of each,
— Zedoary,	three drams.
— Sweet Flag,	
Cloves,	of each,
Cinnamon,	one ounce.
Grains of Paradise,	
Seeds of Anise,	sixteen ounces.
— Fennel,	of each,
— White Poppy,	half an ounce.
— Cumin,	
Brandy,	seventy-two ounces.

Digest for eight days, distil, and add to the product

White Sugar, four ounces.

Mix.—It is recommended in the greater part of the diseases of the uterus.—Dose, a few drams.

TEMPERATE CORDIAL LIQUOR. (WIRTEM.)

R. Cinnamon,	three ounces	two drams.
Lemon Peel,		one ounce.
Yellow Sanders,		half an ounce.
Galanga,		
Cloves,	of each,	
Mace,	one dram and a half.	
Nutmegs,		
Lesser Cardamoms,		three drams.
Flowers of Lavender,		six drams.
— Borage,		one ounce.
Petals of Roses,	two ounces and a half.	
White Wine,		
Juice of Rennet Apples,	of each,	
— Raspberries,	one pound.	
Borage Water,	of each,	
Bugloss Water,	one pound and a half.	
Rose Water,	two pounds and a half.	
Raspberry Water,	of each,	
Cherry Water,	two pounds.	

Digest for four days, and distil six pounds.—Dose, from half an ounce to one ounce.

CHARLES THE FIFTH'S CEPHALIC LIQUOR. (PALAT.; WIRTEM.)

R. Tops of Marjoram,	
— Balm,	
— Origanum,	of each,
Flowers of Rosemary,	two ounces.
— Lavender,	
Petals of Red Roses,	
Flowers of Sage,	
Cloves,	
Cardamoms,	of each,
Cinnamon,	half an ounce.
Grains of Paradise,	
Mace,	
Nutmegs,	
Proof Spirit,	six pounds.

Digest for six days, and distil. (PALAT.)

Z

WIRTEM. prescribes two ounces each, of the seven first ingredients, and also two ounces each, of flowers of the lily of the valley, of cloves, and of nutmegs; half an ounce each, of cardamoms, of cinnamon, of cubebs, of grains of paradise, and of mace; eight pounds of proof spirit; the addition of a pound and a half of water after six days' digestion; and the further addition, at pleasure, of a few grains of ambergris and of musk.

Dose, from two drams to an ounce.

MATTHIOLE'S ELIXIR.

Elixir Vitæ Matthioli. (PALAT.; WIRTEM.; CADET DE GASSICOURT; SPIELMANN)

- R. Root of Galanga, }
 ——— Ginger, } of each, one ounce.
 ——— Zedoary, }
 ——— Sweet Flag, }
 Leaves of Marjoram, }
 ——— Mint, }
 ——— Thyme, } of each,
 ——— Wild Thyme, } half an ounce.
 ——— Sage, }
 ——— Rosemary, }
 Petals of Pale Roses, }
 Seeds of Anise, } of each,
 ——— Fennel, } two drams.
 Cinnamon, } three ounces.
 Cloves, }
 Nutmegs, } of each, one ounce.
 Mace, }
 Cubebs, }
 Aloe Wood, } of each,
 Yellow Sanders Wood, } half an ounce.
 Lesser Cardamoms, }
 French Orange Peel, } three ounces.
 Alcohol, } twelve pounds.

Macerate for three days; then add

Spring Water, four pounds.

Distil slowly, in a water-bath, about thirteen pounds. Then to obtain the elixir *complete*, two scruples of ambergris and the same of musk. It may be coloured at pleasure with saffron, and sweetened with sugar. (PALAT.; WIRTEM.; SPIELMANN)

- R. Fresh Lemon Peel, one ounce and a half.
 Cinnamon, one ounce.
 Galanga, }
 Ginger, }
 Zedoary, } of each, half an ounce.
 Cloves, }
 Nutmegs, }
 Mace, }

- Sweet Flag, }
 Marjoram, }
 Mint, }
 Thyme, }
 Wild Thyme, } of each,
 Sage, } two drams.
 Rosemary, }
 Provence Roses, }
 Cubebs, }
 Aloe Wood, }
 Yellow Sanders Wood, }
 Lesser Cardamoms, }
 Anise, } of each, one dram.
 Fennel, }
 Alcohol (20 degrees), six pounds.

Distil. (CADET DE GASSICOURT)

Dose, from one dram to half an ounce.—
 It is also employed externally.

AROMATIC TINCTURE.

Tinctura regia seu aromatica. (AUGUSTIN)

- R. Flowers of Orange, }
 ——— Wallflower, }
 ——— Lavender, } of each,
 ——— Roses, } three
 Nutmegs, } drams.
 Cloves, }
 Cinnamon, }
 Alcohol, one pound.

Digest for three days, filter, and distil two-thirds; infuse in the product

- Musk, half a dram.
 Saffron, one scruple.
 Peruvian Balsam, twenty drops.
 Oil of Cloves, } of each,
 ——— Cinnamon, } eight drops.

Filter at the end of three days.

Dose, from twenty to thirty drops.

AQUA CARBUNCULI. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

- R. Flowers of Rosemary, }
 ——— Lily of the Valley, } of each,
 ——— Violet, } three
 ——— Borage, } ounces.
 ——— Lavender, } one ounce.
 Herb of Marjoram, }
 ——— Sage, }
 Nutmegs, }
 Mace, }
 Ginger, } of each,
 Cloves, } half an ounce.
 Cinnamon, }
 Galanga, }
 Juniper Berries, }
 Mistletoe, }
 Seeds of Peony, }
 Cardamoms, } one ounce.

Madeira Wine,	}	of each, three pounds.
Rose Water,		
Strawberry Water,		
Lavender Water,		

Digest for three days; then distil, and add a few gold-leaves to the product. (PALAT.)

R. Herb of Marjoram,	}	of each, one ounce and a half.
— Sage,		
Flowers of Lavender,		four ounces.
— Lily of the Valley,	}	of each, one ounce.
— Roses,		
— Rosemary,		
Cinnamon,		
Lesser Cardamoms,		
Nutmegs,	}	of each, half an ounce.
Mace,		
Cloves,	}	of each, three drams.
Ginger,		
Galanga,		
Juniper,		
Alcohol,		one pound and a half.
Water of Raspberries,	}	of each, three pounds.
— Lavender,		

Digest for eight days, and distil one half. (BRUNS.; WIRTEM.; SPIELMANN)

This liquor used formerly to be applied to the region of the heart and upon the pulse in syncope; it was also administered internally to prevent carbuncles and pestilential buboes from receding, as well as to hasten their developement. Our predecessors multiplied these spirituous and aromatic liquors to infinity, each of which passed for a certain specific against every disease.

Of all this farrago, there scarcely remain to us any except the *Eau des Carmes*, the *Eau de Cologne*, and the *Thieves Vinegar*, the curative and prophylactic virtues of which are so believed in by ignorant people, that it would be impossible to do away with them, as has been done with the greater number of preparations similar to the one which has called forth these remarks.

III. Preparations which contain the active principle of Cinnamon extracted by Infusion.

A.

Extraction of Beer.

AROMATIC BEER. (VAN MONS)

R. Cinnamon,	}	of each, one part.
Ginger,		
Cloves,		
Nutmegs,		
Strong Beer,		sixty-four parts.

Macerate for several days, and strain.

B.

Extraction by Water.

INFUSION OF CINNAMON. (RATIER)

R. Cinnamon,	half an ounce.
--------------	----------------

Boiling Water,	two pounds.
----------------	-------------

Infuse, and strain.

It is excitant, and is recommended in dyspepsia attributed to inertia of the stomach, and in what are called nervous colics. It should be taken hot.

AQUEOUS SYRUP OF CINNAMON. (AMST.; BATAV.; BELG.; GALL.; VAN MONS)

R. Powdered Cinnamon,	four ounces.
Water,	one pound and a half.

Digest for three days in a covered vessel, strain with expression, set aside, decant, and add

Simple Cinnamon Water,	one pound and a half.
White Sugar,	three pounds.

Boil to a syrup with a gentle heat. (BATAV.; BELG.)

R. Simple Cinnamon Water,	one part.
White Sugar,	two parts.

Dissolve with a moderate heat in a covered vessel. (GALL.)

AMST.; and VAN MONS prescribe eight parts of the water and sixteen of sugar. HENRI proposes to add one part of the water to three parts of simple syrup previously boiled *au boulet*.

Dose, from two drams to two ounces.

C.

Extraction by Wine.

HIPPOCRAS; HIPPOCRATIC WINE.

Vinum Cinnamomi seu Calidum seu Hippocraticum. (HERBIP.; AUGUSTIN; NIEMANN; SPIELMANN)

R. Cinnamon,	four ounces.
French Wine,	six pounds.
Sugar,	a sufficient quantity,

to sweeten. Infuse without heat for twenty-four hours, and strain. (HERBIP.)

R. Cinnamon,	three ounces and a half.	
Ginger,	half an ounce.	
Cloves,	}	of each, two drams.
Lesser Cardamoms,		
Nutmegs,		
Red Wine,		thirty pounds.

Digest and strain through flannel, and dissolve in the liquor.

White Sugar,	eight pounds.
--------------	---------------

Strain again. (SPIELMANN)

AUGUSTIN directs half a dram each of cinnamon, of cloves, of mace, and of cardamoms, to be macerated in three pounds of wine, and to make a syrup with ten ounces of sugar.

R. Sweet Almonds,	four ounces.
Cinnamon,	one ounce and a half.

White Sugar, two pounds and a half.
 Proof Spirit, one pound.
 Madeira Wine, two pounds.

Digest without heat for several days, and add to the strained liquor.

Musk, } of each,
 Ambergris, } one grain and a half.
 Strain. (NIEMANN)

VINOUS SYRUP OF CINNAMON. (BAVAR.; HASS.; LIPP.; PIDERIT; VAN MONS)

R. Cinnamon, bruised, two ounces.
 White Wine, twenty ounces.

Digest for twenty-four hours in a moderate heat; filter, and add to the liquor.

White Sugar, boiled *à la plume*,
 two pounds.

Make a syrup. (HASS.; PIDERIT)

LIPP. prescribes two ounces of cinnamon, one pound and a half of wine, and the same of sugar.

R. Cinnamon, in powder, } of each,
 Sugar, } one part.
 Acidulous Wine, twelve parts.

Macerate for twelve hours at a temperature of from thirty to forty degrees. Add to one part of the strained liquor

White Sugar, one part and a half.
 Dissolve, (BAVAR.)

R. Cinnamon, in powder, one part.
 Red Wine, twenty-four parts.

Digest in a moderate heat for one day; express, filter, and add

Sugar, forty-six parts.

Dissolve. (VAN MONS)

CARDIAC SYRUP. (WIRTEM.; CADET DE GASSICOURT)

R. Cinnamon, three drams.
 Cloves, one dram.
 Ginger, half a dram.
 Rose Water, two ounces and a half.
 Wine, eight ounces.

Digest, express, and dissolve in the liquor
 Sugar, sixteen ounces.

D.

Extraction by Alcohol.

TINCTURE OF CINNAMON.

Tinctura seu Essentia Cinnamomi. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; DAN.; GALL.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; SAX.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS)

Tinctura Cinnamomi.

R. Cinnamon, bruised, three ounces:
 Proof Spirit, two pints.
 Macerate for fourteen days, and strain. (LOND.)

DUBL. prescribes three ounces and a half of cinnamon, and seven days' maceration.

Tinctura Lauri Cinnamomi. (EDINB.)

R. Cinnamon, bruised, three ounces.
 Dilute Alcohol, two pounds and a half.

Digest for seven days and filter.

AMER.; GALL.; FORM. HÔP. FR.; HASS.; LIPP.; and COXE prescribe one part of cinnamon, and four parts of alcohol (22 degrees); to be digested for six days.—AMST.—one of cinnamon, and eight of alcohol (0.907)—AUSTR.—one of cinnamon and six of alcohol (0.850).—BAVAR.—one of cinnamon and six of alcohol (0.900).—BATAV.; and BELG.—one of cinnamon and eight of alcohol (20 degrees).—BRUGNATELLI; and SWEDIAUR the same as EDINB.—HANN.; OLDEN.; POL.; CAST. BORUS.; and BORUS.—five ounces of cinnamon and two pounds of alcohol.—BRUNS.; GALL.; and VAN MONS—one ounce of cinnamon and six ounces of proof spirit.—DAN.; and SAX.—one part of cinnamon and five parts of alcohol.

R. Powdered Cinnamon, one ounce.
 Alcohol (20 degrees), four ounces.

Digest for four days in a water bath, and strain; pour upon the residue

Alcohol (20 degrees), two ounces.

Digest again for two days, strain, mix, and filter. (ANT.)

It is excitant, slightly astringent, and considered nervine.—Dose from two drams to one ounce.

AMST. prescribes also a *Tinctura Cassie Lignee* prepared with one part of cassia, and eight parts of alcohol; to be digested for eight days.

ALCOHOLIC SYRUP OF CINNAMON.

BRUNS.; HANN.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SARD.; WIRTEM.; HERBIP.; SWEDIAUR; VAN MONS.

R. Tincture of Cinnamon, one ounce.
 Simple Syrup, one pound.
 Mix well. (SWEDIAUR)

- R. Bruised Cinnamon, two ounces.
 Vinous Cinnamon Water, fourteen ounces.

Infuse, and add to eleven ounces of the strained liquor

White Sugar, one pound and a half.
 Dissolve. (POL.)

ROSS. prescribes five ounces of cinnamon, two pounds of vinous cinnamon water, twenty-four hours' maceration, and three pounds four ounces of white sugar.

- R. Powdered Cinnamon, one part.
 Vinous Cinnamon Water, ten parts.
 White Sugar, fifteen parts.

Moisten the cinnamon with one third of the water, twelve hours after, put it into a matrass with the rest of it: infuse for twelve hours in a water-bath. When cold, express, filter, and make a syrup with the sugar. (VAN MONS)

- R. Cinnamon, in coarse powder, five ounces.
 Vinous Cinnamon Water, one pound.
 Rose Water, two ounces.

Digest for twenty-four hours in a moderate heat, filter, and add to the liquor

Sugar, boiled *à la plume*, three pounds four ounces.

Make a syrup. (BRUNS.; HANN.; PALAT.; SARD.; WIRTEM.)

OLDEN.; BORUS.; and HERBIP. direct two ounces of cinnamon, and twelve ounces of vinous cinnamon water to be digested for twelve hours, and a syrup to be made with one pound and a half of sugar, and eleven ounces of the strained liquor.

TONIC POTION.

Vin cordial. (GALL.; FORM. HÔP. FR.; RATIER)

- R. Tincture of Cinnamon, one part.
 Red Wine, ten or twenty parts.
 Mix. (FORM. HÔP. FR.)

- R. Tincture of Cinnamon, two drams.
 Red Wine, five ounces.
 Simple Syrup, one ounce.
 Mix. (RATIER)

- R. Tincture of Cinnamon, } of each,
 Balm, } five parts.
 Red Wine, one hundred parts.
 Simple Syrup, thirty parts.
 Mix. (GALL.)

CARDIAC MIXTURE. (SWEDIAUR)

- R. Tincture of Cinnamon, } of each,
 Syrup of Cinnamon, } one ounce.
 Aromatic Electuary, } two drams.
 Peppermint Water, } five ounces.

Mix.—Dose, two or three spoonfuls.

STRENGTHENING TINCTURE.

Tinctura roborans. (SWEDIAUR)

- R. Tincture of Cinnamon, two ounces.
 Kino, one ounce.
 Acetic Acid, two drams.

Mix.—Dose, one or two drams.

COMPOUND TINCTURE OF CINNAMON.

Tinctura seu Essentia aromatica seu Cinnamomi composita, Alcohol cum Aromatibus composita. (LOND.; EDINB.; ANT.; BAVAR.; DAN.; MIL. COPEN.; GALL.; FENN.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; CADET DE GASSICOURT; HUFELAND; NIEMANN; SWEDIAUR; VAN MONS)

Tinctura Cinnamomi composita.

- R. Cinnamon, bruised, six drams.
 Cardamom Seeds, bruised, three drams.

Long Pepper, in } of each,
 powder, } two drams.
 Ginger, sliced, }
 Proof Spirit, two pints.

Macerate for fourteen days, and filter. (LOND.)

- R. Cinnamon, bruised, } of each,
 Lesser Cardamom } one ounce.
 Seeds, bruised, }
 Long Pepper, powdered, two drams.

Proof Spirit, two pounds and a half.

Digest for seven days, and filter. (EDINB.)

- R. Aromatic Powder, two ounces.
 Alcohol, twelve ounces.
 Digest for three days, and filter. (FULD.)

- R. Cinnamon, two ounces.
 Lesser Cardamoms, } of each,
 Cloves, } half an ounce.
 Galanga, }
 Ginger, }
 Alcohol, two pounds.

Infuse, and filter. (DAN.; HANN.; OLDEN.; POL.; BORUS.; SAX.)

VAN MONS prescribes one part each, of cardamoms, of ginger, of cloves, and of galanga, four parts of cinnamon, and thirty-six of proof spirit.—GENEV.; WIRTEM.; HERBIP.; COXE; and NIEMANN give the same formula as LOND.—SWEDIAUR—half an ounce each, of cinnamon, of lesser cardamoms, of cloves, and of ginger, and two

pounds of alcohol; to be digested for six days.—**BAVAR.**—four parts of cinnamon, one each, of cardamoms, of cloves, and of galanga, and forty-eight of alcohol (0.900)—**FENN.**; and **SUEC.**—half an ounce each, of cinnamon, of canella alba, and of galanga, two drams of cardamoms, and one pound of dilute alcohol.—**MIL. COPEN.**—two ounces of cinnamon, half an ounce each, of cloves, of galanga, and of ginger, two ounces of valerian, and twenty-four ounces of alcohol.—**LUSIT.**—one ounce and a half of aromatic powder, half an ounce of angelica root, two drams of pepper, and two pounds and a half of proof spirit.

R. Nutmegs, }
Cloves, } of each, two ounces.
Cinnamon, }
Pomegranate Flowers, }
Alcohol (22 degrees), } two ounces and a half.
two pounds.

Macerate for fifteen days; then express, and pour upon the residue,

Alcohol (22 degrees), two pounds.

Macerate for fifteen days, and strain. Mix and filter the two tinctures. (**GALL.**; **CADET DE GASSICOURT**)

R. Cinnamon, one ounce.
Lesser Cardamoms, }
Cloves, } of each,
Ginger, } two drams.
Alcohol (20 degrees), } eight ounces.

Macerate for eight days, occasionally shaking the vessel; strain, and pour upon the residue

Alcohol (20 degrees), four ounces.

Digest for two days, and strain. Mix and filter the tinctures. (**ANT.**)

R. Cinnamon, }
Root of Sweet Flag, } of each,
Lesser Galanga, } one ounce.
Herb of Peppermint, } of each,
Fresh Lemon Peel, } one ounce and
Lesser Cardamoms, } a half.
Ginger, } of each,
Alcohol, } two drams.
thirty ounces.

Digest for four days, express, and filter. (**LIPP.**)

R. Root of Sweet Flag, }
Angelica, }
Ginger, } of each,
Elecampane, } one ounce.
Peppermint, }
Anise, }
Cloves, } of each, half an ounce.
Cinnamon, }
Black Pepper, } three drams.
Alcohol, } two pounds.

Infuse. Add to the strained tincture,

Oil of Anise, three drams.

Mix. (**HUFELAND**)

It is excitant, stomachic, and carminative. Dose, from fifty to sixty drops.

ALKERMIC ELIXIR.

Alkermès liquide. (**FERR.**; **CADET DE GASSICOURT**)

R. Cinnamon, }
Mace, }
Cloves, } of each, two drams.
Nutmegs, }
Alcohol, } four pounds.

Macerate for seven days. Then add four pounds of sugar dissolved in four pounds of water, and colour the whole with kermesine syrup, or with a mixture of one dram and a half of alum, and one dram of cochineal.

By augmenting the proportion of the sugar by one fourth, and first distilling the alcoholic infusion, a more agreeable liquor is obtained. (**CADET DE GASSICOURT**)

R. Cloves, }
Cinnamon, } of each, three drams.
Vanilla, } two drams.
Alcohol, } three pounds.

Infuse for twenty-four hours; then add a tincture prepared with

Cochineal, four drams.
Alum, one scruple.
Alcohol, six ounces.

Having digested for three days, add further

Cinnamon Water, two pounds.
Sugar, five pounds.

Filter. (**FERR.**)

It is excitant, stomachic, and carminative.—Dose, from half an ounce to one ounce.

AROMATIC STOMACHIC TINCTURE. (**WIRTEM.**)

R. Cinnamon, one ounce.
Cloves, }
Nutmegs, } of each, half an ounce.
Saffron, }
Root of Sweet Flag, six drams.
Mace, two drams.
Peel of Fresh Lemons, two.
Orange, one.
Alcohol, one pound and a half.

Digest, express, and filter. Dose, from fifty to eighty drops.

ANTI-APOPLECTIC ELIXIR. (**CADET DE GASSICOURT**)

R. Cinnamon, one ounce five drams.

Anise Seeds,	} of each, one ounce.	
Juniper,		
Mace,		
Liquorice,		
Galanga,		
Masterwort,	} six drams.	
Cloves,		
Red Sanders Wood,		
White Sanders Wood,		
Yellow Sanders Wood,		
Root of Contrayerva,	} of each,	
Powdered Viper, [<i>Poudre de Vipère</i>].		five drams.
Seeds of Angelica,	} seven pounds.	
Alcohol,		

Digest for one month, and filter.—It is excitant and aromatic.

E.

Extraction by Æther.

ÆTHEREOUS TINCTURE OF CINNAMON.

Elixir Vitrioli seu Vitrioli aromaticum seu Vitrioli Mynsichti seu Vitrioli Edinburgensium seu Vitrioli cum Tinctura aromatica seu Vitrioli dulce seu acido-aromaticum, Tinctura acida aromatica, Alcohol cum Aromatibus sulphuricatus, Tinctura aromatica sulphurica, Elixir aromaticum acidum, Æther sulphuricus cum Alcohole aromaticus, Acidum Vitrioli aromaticum seu cum Alcohole aromatico, Alcohol aromaticum sulphuricum Tinctura Acidi sulphurici. (EDINB.; AMER.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; MIL. COPEN.; HISP.; GALL.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; CADET DE GASSICOURT; COXE; HUFELAND; PIDERIT; SAUNDERS; SPIELMANN; SWEDIAUR; VAN MONS)

Acidum sulphuricum aromaticum.

(EDINB.)

R. Alcohol,	two pounds.
Sulphuric Acid,	six ounces.

Mix very gradually. Digest the mixture in a covered vessel for three days, at a very gentle heat; then add

Cinnamon, bruised, one ounce and a half.

Ginger, bruised, one ounce.

Digest again in a covered vessel for six days: and finally filter through paper placed in a glass funnel.

R. Compound Tincture of Cinnamon,	six parts.
Sulphuric Acid,	one part.

Mix gradually, and filter after a few days. (VAN MONS)

BAVAR.; MIL. COPEN.; DAN.; FENN.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; and SWEDIAUR prescribe twenty-four parts of tincture, and one part of acid.—SARD.—an equal part of each.—HUFELAND—thirty-eight parts of tincture, and one part of acid.

R. Compound Tincture of Cinnamon,	eight parts.
Acid Elixir,	one part.

Mix. (LIPP.)

R. Alcohol,	two pounds.
Sulphuric Acid,	six ounces.

Mix gradually, and digest in a water-bath for three days, at a moderate heat; then add

Cinnamon,	one ounce and a half.
Ginger,	one ounce.
Dried Peppermint,	half an ounce.

Digest again for three days, at the same temperature, and filter. (BRUNS.; GENEV.; WIRTEM.; SPIELMANN)

AMST. gives the same formula, but omits the mint.—SWEDIAUR prescribes one ounce and a half of cinnamon, one ounce of ginger, half an ounce of sage, twenty-four ounces of alcohol, six hours' infusion, and the addition of six ounces of sulphuric acid to the strained liquor.

R. Alcohol,	one pound and a half.
Sulphuric Acid,	three ounces.

Set aside for two days in a sand-bath, at + 35 degrees R.; add then

Leaves of Peppermint,	} of each,
——— Mentha Crispa,	
	half an ounce.

Cinnamon,	} of each, three drams.
Cloves,	
Ginger,	

Infuse for five days at the ordinary temperature; express and filter. (BRUGNATELLI; CADET DE GASSICOURT)

R. Concentrated Alcohol,	sixteen ounces.
Sulphuric Acid,	three ounces.

Mix these very gradually; then digest for three days in a moderate heat, and add

Powdered Cinnamon,	six drams.
——— Calumba,	} of each,
——— Ginger,	
——— Peppermint,	half an ounce.
	two drams.

Infuse for three days, and strain. (HERBIP.)

FULD. prescribes sixteen ounces of alcohol, two ounces of sulphuric acid, eight drams of cinnamon, six drams of calumba, and half an ounce of peppermint.—VAN MONS—six drams each, of cinnamon, of ginger, and of cloves, one ounce and a half of alcohol, three days' infusion without

heat, the instillation of twenty-four ounces of RABEL's water into the strained liquor, and three days' subsequent maceration.—AMER.; LUSIT.; and COXE—the same as EDINB.

- R. Fresh Lemon Peel, } of each,
Nutmegs, } two drams.
Cubebbs, }
Cinnamon, }
Ginger, } of each three drams.
Cloves, }
Leaves of Mentha Crispa, } of each,
Sage, } half
an ounce.
Root of Sweet Flag, } of each,
Lesser Galanga, } one ounce
and a half.
White Sugar, } two ounces.
Concentrated Sulphuric Acid, } three ounces.
Proof Spirit, } thirty-two ounces.

Mix the acid with two ounces of the spirit, and add the sugar; infuse the six first ingredients with the remainder of the spirit, and then add the others. After sufficient extraction, express, add the acid, and filter at the expiration of a few days. (VAN MONS)

- R. Herb of Peppermint, } of each,
Sage, } one ounce
and a half.
Root of Sweet Flag, } of each,
Lesser Galanga, } one ounce.
Cinnamon, }
Lesser Cardamoms, } three drams.
Orange Peel, } half an ounce.
Rectified Alcohol, } thirty ounces.

Digest for three days, filter, and add
Diluted Sulphuric Acid, half a pound.
Preserve. (HASS.; PIDERIT)

- R. Sulphuric Acid (66 degrees), } two ounces.
Alcohol (38 degrees), } nine ounces.

Digest for three days in a covered vessel, at a temperature of from forty-five to fifty degrees, with frequent agitation. Then add

- Root of Lesser Galanga, } six drams.
Sweet Flag, } half an ounce.
Cloves, } of each,
Cinnamon, } one dram and a half.
Ginger, }
Lemon Peel, }
Nutmegs, } of each, one dram.
Cubebbs, }
White Sugar, } one ounce.

Macerate for twelve days, and strain; then infuse the residue for three days, in the remainder of the alcohol: express, mix together the two liquors, and filter. (ANT.)

- R. Sulphuric Acid, } three ounces.
Proof Spirit, } twenty-four ounces.
Root of Lesser Galanga, } one ounce and a half.
Sweet Flag, } one ounce.
Herb of Mentha Crispa, } of each,
Sage, } half an ounce.
Cinnamon, }
Cloves, } of each, three drams.
Ginger, }
Nutmegs, }
Cubebbs, } of each, two drams.
Lemon Peel, }

Infuse for eight days, and dissolve in the strained liquor

- White Sugar, } four ounces.

Filter. (SPIELMANN)

- R. Alcohol, } three pounds.
Sulphuric Acid, } four ounces.
Root of Sweet Flag, } of each,
Lesser Galanga, } two ounces.
Ginger, } six drams.
Leaves of Mint, } of each,
Sage, } one ounce.
Cubebbs, } of each, half an ounce.
Nutmegs, }
Orange Peel, }
White Sugar, } three ounces.

Infuse for three days in a sand-bath, and filter. (PIERQUIN)

- R. Root of Sweet Flag, } one ounce.
Lesser Galanga, } one ounce and a half.
Ginger, } three drams.
Herb of Mentha Crispa, } of each,
Sage, } half an ounce.
Cloves, } of each,
Cinnamon, } three drams.
Cubebbs, } of each,
Nutmegs, } two drams.
Aloe Wood, } of each,
Lemon Peel, } one dram.
Alcohol, } one pound and a half.

Macerate for fifteen days, and add

- Sulphuric Acid, } two ounces.

Digest for six days, express, and filter. Pour upon the residue

- Alcohol, } six ounces.

Infuse for some time, mix this with the preceding, and finally add

- White Sugar, } four ounces.

Digest, decant, and preserve. (BRUNS.; PALAT.; WIRTEM.)

- R. Root of Sweet Flag, } of each,
Lesser Galanga, } half an ounce.
Flowers of Chamomile, } of each,
Leaves of Sage, } two drams.
Wormwood, }
Mentha Crispa, }

Cloves, }
 Cinnamon, } of each,
 Cubebs, } one dram and a half.
 Nutmegs, }
 Ginger, }
 Aloe Wood, } of each, half a dram.
 Lemon Peel, }
 White Sugar, one pound and a half.
 Alcohol (22 degrees), three ounces.
 At the end of six hours, add
 Sulphuric Acid (66 degrees),
 two ounces.

Then, after twenty hours,

Alcohol (12 degrees), nine ounces.

Digest for four days, strain, express, and filter. (GALL.; CADET DE GASSICOURT)

R. Acid Sulphuric Æther, four ounces.
 Alcohol (20 degrees), eight ounces.
 Cinnamon, } of each, half an ounce.
 Ginger, }

Digest for eight days in a stopped flask, and filter. (BATAV.; BELG.)

HISP. prescribes eight ounces each, of cinnamon and of orange peel, two ounces of gentian, one pound of alcoholized sulphuric æther, and four days' digestion.—CADET DE GASSICOURT—three drams of cinnamon, one dram and a half of lesser cardamoms, one dram each, of long pepper and of ginger, and four days' maceration in a mixture of one part of sulphuric æther and two parts of alcohol.—NIEMANN; and SWEDIAUR—six drams of cinnamon, one ounce of lesser cardamoms, two drams of long pepper, three drams of angelica seeds, two pounds and a half of alcoholized sulphuric æther, and eight days' digestion.—SPIELMANN; and VAN MONS—six drams of cinnamon, one ounce of lesser cardamoms, three drams of angelica seeds, two drams of long pepper, and ten or twelve days' maceration.—AMST.—three drams of cinnamon, half an ounce of cardamoms, one dram and a half of angelica root, one dram of long pepper, five ounces of sulphuric æther, ten ounces of alcohol (0.835), and seven days' maceration.—ANT.—three drams of cinnamon, half an ounce of cardamoms, one dram and a half of angelica root, one dram of long pepper, eight ounces of alcoholized sulphuric æther, and six days' maceration, to be succeeded by the maceration of the residue for three days, with four ounces of æther, the two liquors to be then mixed together.

R. Oil of Cinnamon, } of each,
 ——— Lemon, } half a dram.
 ——— Mint, } one dram.
 Alcoholized Sulphuric Æther,
 eight ounces.

Digest in a moderate heat. (HERBIP.)

It is excitant, and was formerly celebrated as stomachic and tonic, and used to be
 VOL. II.

prescribed in every disease arising from debility and also in passive hæmorrhage.—Dose, from thirty to fifty drops, in wine or any other appropriate liquor.

AROMATIC MIXTURE.

Mixtura piperita. (GENEV.)

R. Vitriolic Elixir, two drams.
 Syrup of Ginger, two ounces.
 Peppermint Water, eight ounces.

Mix.

ANTI-SCORBUTIC LIQUOR.

Essentia anti-scorbutica. (BRUNS.; WIRTEM.)

R. WINTER'S Bark, two ounces.
 Subcarbonate of Potass,
 one ounce and a half.

Leaves of Buckbean, }
 Root of Arum, }
 ——— Patience, } of each,
 Bark of Ash, } one ounce.
 ——— Sassafras, }
 Myrrh,
 Herb of Wall pepper, [*se-*
dum acre]. } of each,
 Tops of Lesser Centaury, } six drams.
 Seeds of Patience,
 Spirit of Scurvy-grass, three pounds.

Digest for six days, express, and add

Essence of Amber, three ounces.
 Spirit of Sulphuric Æther,
 one ounce and a half.

Digest and filter.

It is excitant, and is reputed anti-scorbutic, attenuant, resolvent, diuretic, and stomachic.—Dose, from thirty to fifty drops.

2. *Laurus Cassia* LINN.—Cassia Bark.

SYNONIMES:—Cannelle de la Chine, des Indes, du Malabar ou de Coromandel, Casse en bois, Casse odorante, Fr.; Zimmtcassie, Zimmtkassie, Zimmtsorte, Mutterzimmt, Kaszienrinde, Sinesicscher, Englischer, Indianischer Zimmt, GERM.; Selikey, ARAB.; Kayu-maris, BA.; Mukalla, CYNG.; Darchinie, Mota darchinie, DUK.; Tuj, HIND.; Canellina, Canella del Coromandel, ITAL.; Kayu-legi, MAL.; Havanga, MALAB.; Singroula, N.; Lawanga puttay, TAM.

EDINB.; DUBL.; AMST.; ANT.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; FULD.; HANN.; LIPP.; OLDEN.; CAST. BORUS.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; AINSIE.; BERGIUS.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN;—ZARDA.

A tree of the East Indies. (*Fig.* Burm. Zeyl. p. 63. t. 28)

The bark and leaves are employed.

The bark (*Cassia cinnamomea*, *Cinnamomum Indicum* seu *Sinense*, *Cassia lignea* *Xylocassia*, *Cannella Malabarica*) is in pieces of about an inch and a half long, from three lines to an inch broad, and from three to six lines thick, either quilled or simply bent longitudinally, smooth, and of a light reddish-brown colour. It is thicker, redder, and much less aromatic than the preceding; it has a somewhat vapid taste, and after being chewed for a long time, leaves in the mouth a mucilage which dissolves in the saliva. Its fracture is not splinty.

It is stimulant, tonic, nervine, and anti-spasmodic.

The leaves (*Malabathrum*) are five or six inches long, lanceolated, pointed at both ends, of a reddish or purple hue beneath, and exhibit three longitudinal nerves. They are slightly aromatic and stimulant, but are little used.

3. *Laurus Culilawan*, LING.—Culilawan.

SYNONIMES:—Culilawan, FR.; Nelkenzimmet, GERM.; Koelilaban, DUT.; Culilabao, PORT.

AMST.; ANT.; BATAV.; BELG.; GALL.; FERR.; WIRTEM.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

A tree of the East Indies and of the Molucca Isles. (*Fig. Rumph. Amb. II. t. 14*)

The bark (*Cortex Culilaban* seu *Culilawan* seu *Caryophylloides*) is in flat or slightly curved pieces, of a brown or reddish colour, covered with grey, glabrous, rugose patches of epidermis. It has an agreeable odour, and an acrid, hot, and aromatic taste.

It is stimulant and stomachic.

TINCTURE OF CULILAWAN.

Tincturaseu Essentia Culilaban. (WIRTEM.)

R. Culilawan Bark, four ounces.
Concentrated Alcohol, one pound and a half.

Digest for four days, and filter.

It is excitant, aromatic, stomachic, and carminative.

4. *Laurus nobilis*, LINN.—Bay Tree.

SYNONIMES:—Common Sweet Bay or Laurel Tree.—Laurier commun, FR.; Lorbeer-baum, GERM.; Bob, BOHEM.; Laurbertra, DAN.; Laurel, SPAN.; Laurierboom, DUT.; Lauro, ITAL.; Bobek dizewo, POL.; Lawreiro, POL.; Lagerbærstrød, SWED.

EDINE.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.;

HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PIDE-RIT.; SPIELMANN.; ZARDA.

A tree originally of the Levant, and naturally in Europe. (*Fig. Zorn, Ic. pl. t. 52*)

The leaves and berries are employed.

The leaves (*Folia Lauri*) are glabrous, hard, coriaceous, somewhat waved on the borders, four or five inches long, of a strong aromatic odour when rubbed, and of a warm aromatic taste.

The berries (*Baccæ Lauri*) are oval, of about the size of small cherries, of a deep blue colour when fresh, and of a blackish-brown and wrinkled when dry. Each contains, under a thin brittle membrane, two yellow or yellowish-brown seeds. They have an aromatic and agreeable odour, and a bitterish, acrid, and aromatic taste.

The leaves are employed rather as a condiment than medicament. They are not more stimulant than the berries.

STIMULANT POWDER.

Pulvis Lauri carbonicus. (AUGUSTIN)

R. Bay Berries,	}	of each,
Nutmegs,		six ounces.
Burnt Hartshorn,		
Liquorice,		thirty-six ounces.

Mix.—Dose, a tea-spoonful in rachitis.

OIL OF BAY BERRIES.

Oleum laurinum seu *Baccarum Lauri* seu *Lauri expressum.* (AUSTR.; BRUNS.; HISP.; GALL.; FULD.; HASS.; LUSIT.; PALAT.; SARD.; WIRTEM.; SWEDIAUR)

R. Bay Berries, any quantity.

Bruise slightly in a mortar; then beat them on a heated slab, and express. (HISP.)

R. Bay Berries, any quantity.

Bruise in a mortar, beat them on a heated slab, then boil in water, and take off, when cold, the oil which is congealed on the surface. (AUSTR.; BRUNS.; GALL.; HASS.; LUSIT.; PALAT.; SARD.; SUEC.; WIRTEM.)

R. Bay Berries, any quantity.

Bruise in a mortar, inclose the paste in a linen bag, expose it to the vapour of boiling water, and then submit it to the action of a heated press. (SWEDIAUR)

HASS.; OLDEN.; POL.; BORUS.; and VAN MONS direct the oil which is met with in commerce to be used.

It is stimulant, resolvent, carminative, nervine, and corroborant; it is recommended in diseases of the uterus, spasmodic affections, cases of worms in children, pains and weaknesses of the joints, and in pain of the ear.

CARMINATIVE LAVEMENT. (LIPP.)

R. Chamomile Flowers, half an ounce.
Water, one pound.
Boil, and add when strained
Oil of Bay Berries, half an ounce.
Mix.

ÆTHEREOUS LINIMENT.

Linimentum æthereum. (AUGUSTIN)

R. Sulphuric Æther, two drams.
Alcohol, two ounces.
Oil of Bay Berries, half an ounce.
Mix.—It is recommended in frictions, to herniæ, principally to those of the umbilicus in children.

OINTMENT OF BAY.

Unguentum laurinum, Oleum laurinum infusum. (GALL.; SARD.; VAN MONS)

R. Bay Leaves, one part.
Lard, two parts.
Bruise together, boil slowly to the consumption of the humidity, and add towards the end

Bruised Bay Berries, one part.

Digest for ten hours in a water-bath, and express. (GALL.)

SARD. directs three pounds of fresh bay leaves to be digested for three days in a water-bath, in six pounds of olive oil, to strain with strong expression, and to digest the oil again, for three days, with one pound and a half of leaves and the same of berries.

R. Fresh Bay Leaves, } of each,
Oil of Almonds, } one part.
Fresh Bay Berries, four parts.
Lard, eight parts.

Bruise the leaves and berries, and boil them gently in the lard and oil for twelve hours; then strain, set aside to cool, remove the deposit, and melt again; or, wash with hot water till this comes off clear. (VAN MONS)

JEROMEL proposes to pour one pound of melted lard upon three ounces of dried and pounded leaves, and to keep the vessel at the same degree of heat for twelve hours; then to strain, and to melt with the ointment one pound of expressed oil of bay-berries.

It is stimulant and corroborant.

NERVINE OINTMENT.

Unguentum nervinum seu laurinum seu aromaticum seu Rosmarini compositum. (AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Herb of Sage, }
Ground Pine, } of each, one ounce.
Marjoram, }
Rosemary, }
Mint, }
Lavender, }
St. John's-wort, }
and Flowers of Primrose, }
Flowers of Chamomile, }
Melilot, }
Tops of Dill, }
Root of Pyrethrum, }
Bay Berries, }
Juniper Berries, }
Mastic, half an ounce.
Benzoin, six drams.
Turpentine, four ounces.
Oil of Earthworms, }
Dill, } of each,
St. John's-wort, } three ounces.
Fox's Fat, }
Human Fat, } of each, one
Wild Cat's Fat, } ounce and a half.
Dog's Fat, }
Venice Turpentine, one ounce.
Boil, and strain. (PALAT.)

R. Herb of Sage, }
Ground Pine, }
Marjoram, }
Rosemary, }
Mint, }
Rue, }
Lavender, }
St. John's-wort, }
and Flowers of Primrose, }
Flowers of Chamomile, }
Melilot, }
St. John's-wort, }
Tops of Dill, }
Bay Berries, }
Juniper Berries, }
Olive Oil, one pound.

Macerate for several days; then boil slowly to the consumption of the humidity, strain, and add

Venice Turpentine, two ounces.
Yellow Wax, four ounces and a half.
Common Turpentine, one ounce.
Oil of Earth-worms, } of each,
Bay Berries, } two ounces.

Benzoin, dissolved in three ounces of Concentrated Alcohol, one ounce.

Make an ointment. (BRUNS.; WIRTEM.)

- R. Root of Pyrethrum, }
 Fresh Leaves of Dill, }
 Bay, }
 Marjoram, }
 Mentha } of
 Crispa, } each,
 Rosemary, } four
 Sage, } parts.
 Flowers of Chamomile, }
 Lavender, }
 Melilot, }
 St. John's- }
 wort, }
 Benzoin, } of each, six parts.
 Mastic, }
 Bay Berries, } of each, eight parts.
 Juniper Berries, }
 Lard, } of each, sixteen parts.
 Turpentine, }
 White Wax, } forty-three parts.
 Olive Oil, } ninety-six parts.

Bruise the vegetables with the oil and lard, and macerate them in a covered vessel, at a moderate heat, for four hours; then strain, wash with warm water, add the mastic and benzoin melted with the turpentine: strain again, add the melted wax, and stir till perfectly cold. (VAN MONS)

- R. Fresh Herb of Rue, }
 Flowering Tops of Sage, }
 Marjoram, }
 Rosemary, }
 Mint, }
 Dill, }
 Lesser } of each,
 Mullein, } two
 Lavender, } ounces.
 St. John's- }
 wort, }
 Chamomile, }
 Melilot, }
 Leaves of Bay, } of each,
 Juniper, } one ounce.
 Oil of Earth-worms, } three pounds.

Macerate for two days; then boil to the consumption of the humidity, and add to the strained liquid

Yellow Wax, nine ounces.

Add when the mass is half cold

- Oil of Wax, } of each,
 Turpentine, } two ounces.
 Ointment of Marshmallows, } four ounces.

Mix. (HISP.)

- R. Herb of Rosemary, } two ounces.
 Rue, }
 Marjoram, }
 Mentha crispa, }
 Flowers of Melilot, } of each,
 Lavender, } one ounce.
 Chamomile, }
 Bay Berries, }

- Lard, one pound.
 Mutton Suet, half a pound.

Boil slowly to the consumption of the humidity, and melt in the strained mass

- Yellow Wax, } of each, two ounces.
 Turpentine, }
 Oil of Bay Berries, } four ounces.

Make an ointment. (DAN.)

- R. Herb of Rosemary, }
 Sage, } of each,
 Lavender, } one handful.
 Rue, }
 Fresh Bay Berries, } three ounces.
 Lard, } one pound.
 Mutton Suet, } four ounces.

Keep the mixture in a water-bath for four days; then express, and add

- Oil of Bay Berries, eight ounces.

Mix. (BRUGNTELLI)—This is the *Unquendum aromaticum*.

- R. Fresh Herb of Rosemary, } one pound.
 Marjo-ram, } of each,
 Rue, } half a pound.
 Bay Berries, } of each,
 Root of Pyrethrum, } four ounces.
 Lard, } four pounds.
 Mutton Suet, } two pounds.

Boil to the consumption of the humidity; express, and add

- Yellow Wax, } half a pound.
 Oil of Rosemary, } of each,
 Juniper, } three ounces.

Mix. (HANN.; OLDEN.; BORUS.; SAX.; NIEMANN)

- R. Flowers of Rosemary, } of each,
 Leaves of Tacamahac, } four ounces.
 Lard, } two pounds.

Digest for fifteen days, and strain; add, when cold,

- Distilled Water of Lavender, } of each,
 Chamomile, } half
 Rue, } an ounce.

Mix. (HASS.)

- R. Leaves of Tacamahac, } eight ounces.
 Melted Lard, } two pounds.

Strain at the end of fifteen days, and add

- Oil of Amber, } of each,
 Chamomile, } half an ounce.
 Rue, } of each,
 Mint, } two drams

Mix. (PIDERIT)

- R. Fresh Bay Berries, } eight ounces.
 Leaves, } two ounces.

Bruise, and macerate in

- Melted Lard, sixteen ounces.

At the end of twenty-four hours, express, wash with warm water, melt again, and set aside. (SWEDIAUR)

R. Mutton Suet, eight parts.

Melt, remove from the fire, and add when half cold, stirring well,

Oil of Bay,	twelve parts.
— Turpentine,	one part.
— Amber,	one half part.
— Peppermint,	} of each,
— Rosemary,	
— Juniper,	
	one eighth part.

Mix. (SAX.)—This is also called *Unguentum aromaticum*.

R. Lard, four pounds.
Mutton Suet, two pounds.
Yellow Wax, half a pound.

Add to the mixture when melted and half cold,

Oil of Bay,	two ounces.
— Rosemary,	} of each,
— Juniper,	
	three ounces.

Mix, and preserve. (POL.)

R. Lard, one ounce.
Oil of Bay, half an ounce.
— Rosemary, } of each,
Solution of Am- } one dram and a
monia, } half.

Mix. (WURCEB.)

R. Stag's Suet, six ounces.

Add to it when melted,

Oil of Bay,	six ounces.
— Turpentine,	one ounce.
— Amber,	} of each,
Muriate of Ammonia,	
	half an ounce.

Mix. (LIPP.)

R. Lard, three ounces.
Stag's Suet, one ounce.

Melt gradually, and add

Oil of Bay,	one ounce.
— Amber,	half an ounce.
— Mentha Crispa,	two drams.

Triturate till cold. (FULD.)

R. Mutton Suet, eight ounces.
Oil of Bay, ten ounces.

Melt slowly, and add when nearly cold,

Spirit of Turpentine,	one ounce.
Oil of Amber,	half an ounce.

Mix. (ANT. ; BELG. ; SUEC. ; SPIELMANN ; VAN MONS)

R. Mutton Suet, eight ounces.

Melt, and add

Oil of Bay,	one pound.
Spirit of Turpentine,	} one ounce and a half.

Mix. (LUSIT.)

R. Mutton Suet, eight ounces.

Add to the melted mass,

Oil of Bay,	one pound.
— Turpentine,	one ounce.
— Petroleum,	half an ounce.

Triturate till cold. (BATAV. ; ROSS.)

AMST. prescribes eight ounces of suet, one pound of oil of bay, one ounce of spirit of turpentine, and half an ounce of oil of cajeput.

R. Simple Ointment,

	two pounds and a half.
Oil of Bay,	} of each, three ounces.
Yellow Wax,	

Add to the mixture when melted and nearly cold,

Essential Oil of Juniper,	} of each,
— Wild	
Thyme,	two drams.

Mix. (AUST.)

This ointment is employed externally as a corroborant, and as a resolvent.

CARMINATIVE OINTMENT. (FULD. ; PALAT.)

R. Bay Berries, half an ounce.
Fresh Leaves of Chamomile, two handfuls.

— Mint,	} of each,
Fresh Tops of Southernwood,	
— Ground Ivy,	
— Mentha Crispa,	} one hand- ful.
— Tansey,	

Seeds of Cumin,	} of each,
— Lovage,	
Oil of Chamomile,	one pound.

— Dill,	} of each, half an ounce.
— Rue,	

Macerate for fifteen days ; then boil slightly to the consumption of the humidity, and add, when strained,

Yellow Wax,	two ounces.
Storax,	six drams.

Oil of Anise,	} of each, one dram.
— Caraway,	
— Mint,	

Mix. (PALAT.)

R. Lard, six drams.
Yellow Wax, one dram.

Add to the melted mixture when nearly cold,

Oil of Bay,	one dram.
— Anise,	eighteen drops.
— Caraway,	twelve drops.

Mix. (FULD.)

It is emollient, anti-spasmodic, and carminative.

CAMPHORATED AROMATIC OINTMENT.

Unguentum laurino-campboratum. (BAVAR. ; CADET DE GASSICOURT)

R. Mutton Suet, sixteen parts.

Oil of Bay, twenty parts.
 Heat slowly, and add when melted,
 Camphor, dissolved in a little
 Oil of Juniper, two parts.
 Mix. (BAVAR.)

CADET DE GASSICOURT directs two ounces each, of lard, of beef suet, of oil of bay, and of wax, to be melted together, and when nearly cold, half an ounce of camphor, dissolved in one ounce of alcohol, to be added.

UNGUENTUM MARTIATUM. (WIRTEM.; SPIELMANN; VAN MONS)

R. Fresh Leaves of Bay, eight ounces.
 ————— Rue, five ounces.
 ————— Marjoram, four ounces.
 ————— Rosemary, three ounces.
 ————— Myrtle, } of
 ————— Dwarf Elder, } each,
 ————— Mint, } one
 ————— Sweet Basil, } ounce.
 White Wine, four ounces.
 Olive Oil, one pound.
 Fresh Butter, }
 Bear's Grease, } of each, six drams.
 Chicken's Fat, }
 Stag's Marrow, }

Infuse for several days; then boil slowly to the consumption of the humidity, and add to the strained mass,

Yellow Wax, eight ounces.
 Powdered Storax, five drams.
 ————— Mastic, half an ounce.
 ————— Olibanum, three drams.

Mix. (WIRTEM.; SPIELMANN)

R. Fresh Leaves of Wormwood, }
 ————— Southernwood, }
 ————— Sweet Basil, }
 ————— Calamint, }
 ————— Costmary, } of
 ————— Germander, } each,
 ————— Bay, } two
 ————— Marjoram, } parts.
 ————— Water-mint, }
 ————— Rosemary, }
 ————— Rue, }
 ————— Savine, }
 ————— Sage, }

Bruise them in a mortar, express the juice, coagulate it over the fire, remove the fecula, wash it, and separate the water from it as well as possible; then dissolve it in

Lard, sixty-four parts,
 heated with
 Seeds of Cumin, } of each,
 ————— Fenugreek, } one part.
 Nutmegs, six parts.
 Then add

Liquid Storax, } of each, four parts.
 White Wax, }

Strain, and stir till cold. (VAN MONS)

It is stimulant, and resolvent.

PLASTER OF BAY BERRIES. (BRUNS.; PALAT.; SARD.; WIRTEM.; SPIELMANN)

R. Bay Berries, four ounces.
 Round Cyperus, }
 WINTER's Bark, } of each, one ounce.
 Olibanum, }
 Mastic, }
 Myrrh, }

Make a powder, and incorporate it with a melted mixture of

Yellow Wax, eight ounces.
 Resin, }
 Oil of Bay, } of each,
 — Turpentine, } two ounces.

Mix.—It is stimulant and resolvent.

ELECTUARY OF BAY BERRIES. (PALAT.; SARD.; SPIELMANN)

R. Bay Berries, two ounces.
 Mastic, }
 Myrrh, } of each, one ounce.
 Olibanum, }
 Arabian Costus, } of each,
 Round Cyperus, } half an ounce.
 Oil of Bay, } of each,
 Chio Turpentine, } three ounces.
 Honey, one pound and a half.

Having melted the honey, turpentine, and oil together, add the other ingredients in powder. (SARD.)

R. Bay Berries, one ounce and a half.
 Root of Sweet Flag, }
 Herb of Rue, }
 ————— Mint, }
 ————— Origanum, }
 Seeds of Anise, }
 ————— Cumin, } of each,
 ————— Nigella, } two drams.
 ————— Lovage, }
 ————— Caraway, }
 ————— Carrot, }
 Black Pepper, }
 Long Pepper, }
 Castor, }
 Sagapenum, } half an ounce.
 Opoponax, } three drams.
 Honey, fourteen ounces.

Mix. (PALAT.; SPIELMANN)

AROMATIC ALCOHOL. (ANT.)

R. Bay Berries, one ounce.
 Myrrh, }
 Storax, } of each, six drams.

Cinnamon,	} of each,	two drams and a half.
Ginger,		
Nutmegs,		
Cloves,		
Water,	} of each, fifty-	four ounces.
Alcohol (20 degrees),		

Macerate for twenty-four hours, and distil till the product marks 30 degrees.

5. *Laurus Sassafras*, LINN.—Sassafras.

SYNONIMES :—Sassafras, FR. ; Fenchelholz, GERM. ; Sasafras, ARAB. ; Cay, Vangdi, COCHIN.

LOND. ; EDINB. ; DUBL. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FORM. HÔP. FR. ; FENN. ; FULD. ; GENÈV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; WURCEB. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY and GME-LIN ; PARIS ; SPIELMANN ; ZARDA.

A tree of South America. (*Fig. Flore Medic. VI. 312.*)

The wood (*Lignum Sassafras*) is light, of a spongy texture, brittle, of an iron-grey colour, inodorous, and of a very feeble taste.

The bark is rather thick, light, brittle, and rugose. It is of an iron-brown colour, of an agreeable and fennel-like odour, and of an aromatic and slightly acrid taste.

It is stimulant, diaphoretic and diuretic ; and is employed in gout, rheumatism, and syphilitic disorders.—Dose, from half a dram to one dram, of the powder.

EXTRACT OF SASSAFRAS.

Extractum Ligni Sassafras. (WIRTEM.)

R. Sassafras Wood, rasped,	one pound.
Alcohol,	six pounds.

After sufficient extraction, strain with expression ; digest, and boil the residue for a short time in three pounds of water : clarify the decoction with white of egg, distil off the alcohol, and evaporate to the required consistence.

Dose, from a scruple to a dram.

INFUSION OF SASSAFRAS. (FORM. HÔP. FR. ; NIEMANN)

R. Sassafras Wood, rasped,	half an ounce.
Boiling Water,	one pound.

Infuse for six hours, and strain. (NIEMANN)

FORM. HÔP. FR. prescribes one ounce of sassafras and a sufficient quantity of boiling

water to obtain half a pound of strained infusion.

COMPOUND LIME WATER.

Aqua Calcis composita ; Eau bénite, Eau bénite composée.

R. Sassafras Bark,	half an ounce.
Guaiac Wood, rasped,	half a pound.
Liquorice Root,	one ounce.
Coriander Seeds,	three drams.
Lime Water,	six pints.

Macerate without heat, for two days, and strain. (DUBL.)

BRUNS. ; and SPIELMANN give the same formula.

R. Sassafras Wood,	two ounces.
Nutmegs,	three drams.
Liquorice,	one ounce.
Lime Water,	four pounds.

Macerate for twenty-four hours, and strain. (LUSIT.)

VAN MONS prescribes two ounces of sassafras, three drams of nutmegs, one ounce of liquorice, thirty-six ounces of lime water, and three days' infusion.

R. Sassafras,	half an ounce.
Liquorice,	one ounce.
Lime Water,	six pounds.

Infuse for two days, and strain. (PIERQUIN)

R. Root of Marshmallows,	} of each,	one ounce.
—— Soapwort,		
—— Sarsaparilla,	} of each,	four drams.
Sassafras Wood,		
Lime Water,		four pounds.

Macerate for two days, and add,

Compound Syrup of Smallage,
two ounces.

Mix. (BORIES)

It is given in scrofulous diseases.—Dose, several cupfuls in the course of the day.

SYRUP OF SASSAFRAS.

Syrupus e Corticibus Ligni Sassafras.
(WIRTEM.)

R. Sassafras Bark,	three ounces.
White Wine,	twenty-two ounces.

Infuse, and add to the strained liquor,
Sugar, thirty-two ounces.
Make a syrup.

TINCTURE OF SASSAFRAS.

Tinctura seu Essentia Ligni Sassafras.
(BRUNS. ; WIRTEM. ; VAN MONS)

R. Sassafras Wood, rasped,	two ounces.
----------------------------	-------------

Proof Spirit, twelve ounces.

Digest for three days, and filter. (VAN MONS)

BRUNS.; and WIRTEM. prescribe one ounce and a half of sassafras, eight ounces of alcohol, and eight days' digestion.

This used formerly to be considered anti-catarrhal and alterative; it is now given as a stimulant sudorific, and diuretic, and is principally employed in chronic rheumatism.—Dose, one dram four times a day.

COMPOUND TINCTURE OF SASSAFRAS.

Essentia Ligni Sassafras composita. (WIRTEM.)

R. Sassafras Bark,	four ounces.
Leaves of Sage,	two ounces.
Nutmegs,	one ounce.
Cinnamon,	half an ounce.
Indian Anise,	three drams.
Alcohol,	two pounds.

Digest for three days; strain with expression, and filter.

This is recommended in catarrhal affections, gout, flatulence, certain varieties of itch.—Dose, from fifty to eighty drops.

OIL OF SASSAFRAS.

Oleum Sassafras Ligni æthereum, Æther-oleum Sassafras Ligni. (EDINB.; DUBL.; AMER.; AMST.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; PALAT.; SARD.; WIRTEM.; HERBIP.; COXE; PIDERIT; SWEDIAUR)

Oleum volatile Lauri Sassafras.
(EDINB.)

Oleum Corticis et Lignorum Sassafras.
(DUBL.)

R. Bruised Root of Sassafras,
any quantity.
Water, a sufficient quantity
to prevent empyreuma. (EDINB.)

DUBL. prescribes the bark and wood.

AMER.; AMST.; and COXE; give the same formula as DUBL.—FULD.; HERBIP.; and SWEDIAUR prescribe one part of wood and four parts of water.—PALAT.—one of wood and six of water.—HAMB.—one of wood and eight of water.

R. Sassafras Wood,	ten parts.
Water,	twenty parts.
Muriate of Soda,	one part.

After twelve hours, distil, and separate the oil which falls to the bottom of the receiver. (GALL.)

SARD. prescribes three parts of wood, twenty-four of water, and one of muriate.—HASS.; WIRTEM.; and PIDERIT—six of wood, thirty-six of water, and one of muriate.—FERR.—six of wood, thirty of water, and three of muriate.

R. Sassafras Wood,	two pounds.
Alcohol,	eight ounces.

Digest for two days, and add,

Water,	twelve pounds.
Muriate of Soda,	one pound.

Distil. (LIPP.)

Dose, from two to eight drops.

OLEOSACCHARUM OF SASSAFRAS. (NIE-MANN)

R. Oil of Sassafras,	ten drops.
Sugar,	three drams.

Triturate together.

SASSAFRAS WATER. (GALL.; SARD.)

R. Sassafras Wood,	one part.
Water,	eight parts.

Distil four parts after twelve hours' maceration. (GALL.)

SARD. prescribes one part of wood, six parts of water, and a little muriate of soda.

Dose, from one to four ounces.

COMPOUND SASSAFRAS WATER. (WIRTEM.)

R. Sassafras Wood,	half a pound.
Cinnamon,	three ounces.
Wine,	ten pounds.

Digest for several days, and distil one pound and a half.

LAWSONIA INERMIS, LINN.

SYNONIMES:—Henné, FR.; Rothes Mundholz, Rothes Ägyptisches Färbekraut, GERM.; Pochar, JAV.; Pontolesche, MALAB.

BAVAR.; GALL.; WIRTEM.; AINSLIE; MURRAY and GMELIN; SPIELMANN.

A shrub of the East Indies and of Africa.

{*Ocandria Monogynia*, LINN.; *Salicaria*, JUSS.; *Fig. Hort. Malab. IV. p. 117. t. 57.*)

The root (*Radix Alcanne vera* seu *orientalis* seu *Cypri Antiquorum* seu *Ligustri Egyptiaci*) is employed. It is of a deep red colour, and of an austere styptic taste. It is slightly astringent.

In the East Indies the root of the *Lavsonia spinosa*, LINN. (Urkan, ARAB.; Ma-

ritondi, CYNG.; Mayndie, DUK.; Daun lacca, MAL.; Mail anschi, MALAB.; Henna, PERS.; Sakachera, SANS.; Marudanie, TAM.; Gorunta chettu, TEL.) is also employed. It much resembles the preceding, and is said to be useful in leprosy, as well as in other diseases of the skin; it is prescribed externally and internally, in the dose of half a tea-spoonful.

LEDUM PALUSTRE, LINN.—MARSH LEDUM.

SYNONIMES:—*Romarin sauvage*, FR.; *Heydebienkraut*, *Sumpfporsch*, GERM.; *Vild Rosmarin*, DAN.; *Wilde Rosmoryn*, DUT.; *Ledo*, ITAL.; *Rozmaryn czeski*, POL.; *Getporss*, *Sqwattram*, SWED.

AUSTR.; BRUNS.; DAN.; GALL.; FENN.; FERR.; HAMB.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; BERGIUS.; MURRAY and GME-LIN.; SPIELMANN.

A plant of the north of Europe. (*Decan-dria Monogynia*, LINN.; *Rhodoraceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 54.*)

The tops (*Herba seu Folia Ledi palustris cum Floribus* seu *Rorismarini sylvestris* seu *Anthos sylvestris*) are employed. It consists of hairy, reddish branches, furnished with linear, sessile leaves, revolute at the edges,

green above, reddish and cottony beneath, and terminated by sessile umbels of white flowers. Their taste is bitter and styptic.

This plant is reputed narcotic, and has been extolled in spasmodic diseases.

INFUSION OF MARSH LEDUM. (PH. SYFH.; NIEMANN.; SWEDIAUR)

R. Leaves of Marsh Ledum,
from half an ounce to one ounce.
Boiling Water, one pound.

Infuse for one hour, and strain.

It is recommended in leprosy, itch, tinea, hooping cough, and syphilitic diseases.

Dose, from half a pound to one pound, twice a day.

LEONURUS CARDIACA, LINN.

SYNONIMES:—*Agripaume*, *Cardiaque*, FR.; *Herzgespann*, *Wolfstrapp*, GERM.; *Hartgespan*, DUT.; *Cardiaca*, ITAL.

BRUNS.; GALL.; OLDEN.; WIRTEM.; BERGIUS.; MURRAY and GME-LIN.; SPIELMANN.

This plant grows throughout Europe. (*Didynamia Gymnospermia*, LINN.; *Labiata*, JUSS.; *Fig. Zorn, Ic. pl. t. 114.*)

The herb (*Herba Cardiaca* seu *Pata lupina*) is employed. It consists of a branching stem with petiolated, hairy leaves, the lower ones of which are oval, nearly cordiform, and trilobate, and the caulinary, cuneiform, trilobate, and dentate. It has a strong and disagreeable odour, and a very bitter and repugnant taste.

It is scarcely employed now, but used formerly to be given to children in cardialgia.

LEPIDIUM.—PEPPERWORT.

Two species of this genus of plants are mentioned in pharmacopœias:

1. *Lepidium latifolium*, LINN.—Broad-leaved Pepperwort.

SYNONIMES:—*Passerage à grandes feuilles*, VOL. II.

Grande Passerage, FR.; Pfefferkraut, GERM.; Peperur, DAN.; Broedblabig peperkruid, DUT.; Lepidio, ITAL.; Pe-parœr, SWED.

HISP.; GALL.; SPIELMANN.

A common plant of Europe. (*Tetrady-2 B*

namia Siliquosa, LINN.; *Cruciferae*, JUSS.; *Fig. Ced. Fl. Dan.* t. 557.)

The herb (*Herba Lepidii*) is employed. It consists of a cylindrical, glabrous, branching stem, and of oval, lanceolated leaves of a pale glaucous green colour, denticulated but sometimes entire; the inferior are narrowed into petioles at the base, and the superior are sessile and narrower. It has an acrid, aromatic, and peppery taste.

2. *Lepidium Iberis*, LINN.

SYNONIMES:—*Passerage ibéride*, *Petite Passerage*, FR.

GENEV.

A common plant of Europe. (*Fig. Zorn, Ic. pl.* t. 593.)

The herb (*Herba Iberidis*) is employed. It consists of a branching stem, and of petiolate, lanceolate, radical leaves, dentated or incised at the summit, pinnatifid or lyre shaped, and of linear, entire caulinary

leaves. It has an acrid and strong taste similar to that of cress.

It is stimulant and anti-scorbutic.

CONSERVE OF PEPPERWORT.

Conserva Lepidii. (HISP.)

R. Fresh Leaves of Pepperwort, one part.

Bruise them finely in a stone mortar, and add gradually

White Sugar, two parts.

INFUSION OF PEPPERWORT.

Infusum Lepidii. (NIEMANN)

R. Herb of Pepperwort, from half an ounce to one ounce. Boiling Water, a sufficient quantity to obtain six ounces of infusion.

This infusion is used in Russia, according to ERDMANN, against intermittent fevers. —Dose, a spoonful every hour.

LICHEN.

This is the official name of the nine following plants:

1. *Physcia nivalis*, CAND.—Snow-white Lichen.

SYNONIME:—*Lichen blanc de neige*, FR.

WIRTEM.

A cryptogamous plant of the Alps and Pyrenees. (*Fig. Dill. Musc.* t. 21. f. 56.)

It consists (*Lichen terrestris cinereus*) of white or yellowish leaves, rather broad at the base, and narrow at the summit, branchy, almost scalloped, crisped, terminated by flat, circular lobules, often bordered with black points: the shields are very sparse, sessile, brown, surrounded with a crenulated margin, and situated on the leaves.

This plant was formerly considered as a specific against hydrophobia.

2. *Scyphophorus cocciferus*, CAND.—Cocciferous Lichen.

SYNONIMES:—*Lichen coccifère*, FR.; *Scharlachfarbige Bechernflechte*, GERM.

GALL.

It consists (*Herba Ignis seu Musci pyridati*) of small leaves of a greenish-white

above, but more white beneath, producing cylindrical, frequently branching stems, which become hollowed into the form of a cup, the borders of which present red tubercles.

3. *Peltigera canina*, CAND.—Ash-coloured Ground Lichen or Liverwort.

SYNONIMES:—*Lichen contre la rage*, FR.; *Hundsflechte*, GERM.; *Hondsmoos*, DUT.; *Musgo canino*, PORT.; *Hiendmossa*, SWED.

BRUNS.; GALL.; MURRAY and GMELIN; SPIELMANN.

A cryptogamous plant very common in woods, growing on the earth. (*Fig. Vaill. Bot.* t. 21, f. 16.)

It consists (*Herba Musci canini seu Lichen cinereus terrestris*) of large, broad, round, coriaceous leaves of an ash-grey or brown colour, having red nervures underneath, hairy, branching, and anastomosed, presenting round lobes, some of which, ascending, bear red, even oblong shields without a free margin, and arranged vertically.

The *Peltigera venosa, horizontalis, polydactyla et aphthosa*, indicated by GALL. resemble this very much.

This plant was formerly thought useful against hydrophobia.

4. *Scyphophorus pyxidatus*, CAND.—Funnel-shaped Lichen.

SYNONYME :—Lichen en entonnoir, FR.

GALL.; FORM. HÔP. FR.; FULD.; HAMB.; LIPP.; MURRAY and GMELIN; SPIELMANN.

A cryptogamous plant found on the earth and on old walls. (*Fig. Dill. Musc. t. 14, f. 2.*)

It consists (*Herba Musci pyxidati*) of leaves which are greenish above, but rather white beneath, lobed or crenulated, presenting cups hollowed regularly from below upwards, the margin of which exhibits dull-brown tubercles.

It was formerly recommended against cough.

5. *Usnea plicata*, CAND.—Hairy Oak Lichen or Tree-moss.

SYNONYMES :—Lichen entrelacé, Mousse de chêne, FR.; Häarflechte, Wickelflechte, GERM.; Skinslav, DAN.; Musco arboreo, SPAN.; Worrigh schurffmoos, DUT.; Musgo dos carvalhos, PORT.; Taf, SWED.

HISP.; GALL.; WIRTEM.; MURRAY and GMELIN.

A cryptogamous plant common on the branches of old trees. (*Fig. Dill. Musc. t. 11, f. 1.*)

It consists (*Muscus albus quernus seu arboreus*) of a stem sometimes more than a foot long, rather firm, and tuberculous, having interlaced, whitish branches; its shields are nearly terminal, white, numerous, and bordered with capillary hairs.

It has an agreeable odour, and a rather astringent taste.

That which grew on the skull of a hanged person used anciently to be sold for its weight of gold.

It is slightly astringent; it was formerly considered anodyne, and was recommended as a tonic, to arrest discharges of all kinds, and to appease cough.

6. *Physcia Islandica*, CAND.—Iceland Lichen, Moss, or Liver-wort.

SYNONYMES :—*Lichen*, LOND.; *Lichen Islandicus*, EDINB.; *Cetraria Islandica*, DUBL.—*Lichen d'Islande*, FR.; *Lungenmoos*, Isländisches Mooss, GERM.; *Gaternjk*, BOHEM.; *Islanskmoos*, Fielgræs, DAN.; *Liquen Islandico*, SPAN.; *Ysländisch moos*, DUT.; *Lichene Islandico*, ITAL.; *Meek Islandzki*, POL.; *Musgo Islandico*, PORT.; *Islands mosse*, SWED.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

A very common cryptogamous plant of Europe, particularly in the north. (*Æd. Fl. Dan. t. 153, f. 879.*)

It consists (*Herba Lichenis Islandici seu Musci Islandicus seu catharticus*) of membranous expansions of a chesnut-brown or olive colour, brownish-red at their base, paler underneath, straight, branching, lobed, with the segments erect, sublinear, multifold, channelled, dentato-ciliate, the fructiferous ones are larger, the shields are plane, sitting, appressed, of the same or of a deeper colour than the expansion, with a raised, entire, and ciliated border.

According to PROUST, this plant contains, in one hundred parts, sixty-four of mucus, thirty-three of amylaceous matter, and three of bitter principle.

It possesses, according as the bitter principle may or may not have been removed, either simply alimentary, or alimentary with tonic properties. It has been particularly extolled in pulmonary consumption.

7. *Imbricaria parietina*, CAND.—Common Yellow Lichen.

SYNONYMES :—*Lichen des murs*, FR.; *Wandflechte*, *Goldgelbe Schuppenflechte*, *Plankenflechte*, GERM.; *Guul skorpe-moos*, DAN.; *Maurig schurffmoos*, DUT.; *Væggmoos*, SWED.

AUSTR.; BAVAR.; FERR.; FORM. HÔP. FR.; SAX.

A cryptogamous plant of Europe. (*Fig. Dill. Musc. t. 24, f. 76.*)

It consists of a membranous, imbricated thallus, whitish beneath, greenish above, then passing successively to a golden yellow, and to an ash-grey: it has rounded, lobated, crisped, broad, expanded folioles: the shields are central, numerous, with an entire border, and a disk of a deeper colour.

It is bitter, and has been represented as a succedaneum to cinchona, in the treatment of intermittent fevers. The analysis which SCHRADER has given of it, does not, however, show that there is any analogy between it and cinchona.

8. *Physcia Prunastri*, CAND.—Slow Tree Lichen.

SYNONIME :—Lichen de Prunellier, FR. ;
WIRTEM. ; SPIELMANN.

A cryptogamous plant of Europe. (*Fig. Dill. Musc. tab, 21, f. 54, 55 A.*)

It consists of a soft, membranous expansion, of a pale ash colour, sometimes greenish, wrinkled, embossed, of a milky whiteness beneath, unequally bifurcated, very branching, with the segments erect, linear, slender, and even ; the shields are scattered, brown, and marginal.

It is slightly astringent.

9. *Lobaria pulmonaria*, CAND.—Lungwort.

SYNONIMES :—Lichen pulmonaire, Pulmonaire de chêne, FR. ; Langenflechte, GERM. ; Lungemot, DAN. ; Pulmonaria de arbol, SPAN. ; Longachtig lungwort, DUT. ; Polmonaria di quercia, ITAL. ; Hepatica dos avores, PORT. ; Lungmosse, SWED.

BRUNS. ; GALL. ; FORM. HÔP. FR. ; GENEV. ; WIRTEM. ; HERBIP. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A cryptogamous plant of Europe common in the north. (*Fig. Zorn, Ic. pl. t. 494.*)

It consists (*Herba Pulmoniae arboreae seu Muscus pulmonarius*) of an expansion of a brownish-green above, and marked with cavities separated by a sort of projecting network ; beneath, it is embossed, white and smooth in the convexities, and brown and hairy in the interstices.

PECTORAL SPECIES. (FULD.)

R. Iceland Moss, one part.
Root of Marshmallows, two parts.
Cut, and mix.

ANTI-LYSSIC POWDER.

Pulvis anti-lyssus. (BRUNS. ; WIRTEM. ; SPIELMANN)

R. Snow-white Lichen, two ounces.
Black Pepper, one ounce.
Make a powder.

ICELAND MOSS-CHOCOLATE.

Chocolata Lichenis Islandici, Pasta cacao-tina lichenifera. (SAX. ; AUGUSTIN)

R. Roasted Cocoa Nut, } of each,
Sugar, } four pounds.
Iceland Moss, two pounds.
Jalap, three ounces.

These proportions are indicated by TROMMSDORF. (AUGUSTIN)

R. Iceland Moss, washed with hot water, and dried, one part.
Jalap, one-eighth part.
White Sugar, two parts.
Mix exactly, and incorporate with
Hot Cocoa Paste, one part.
Divide into cakes. (SAX.)

EXTRACT OF ICELAND MOSS. (HERBIP.)

R. Iceland Moss, washed, any quantity.
Alcohol, a sufficient quantity
to cover it by four fingers' breadth. Digest it in a moderate heat, and decant ; boil the residue with water for an hour, and express : evaporate the liquor in a water-bath to the consistence of honey, gradually adding the tincture, and constantly stirring that a uniform mass may be obtained.

INFUSION OF ICELAND MOSS. (SAUNDERS)

R. Iceland Moss, three drams.
Boiling Water, a sufficient quantity
to obtain, after half an hour's infusion, and a few minutes boiling, twenty ounces of strained liquor. Add to this

Syrup of Raspberries,
one ounce and a half.

Mix.

SYRUP OF ICELAND MOSS. (HANN. ; VAN MONS)

R. Iceland Moss, six ounces.
Boiling Water, six pounds.
Boil down to four pounds ; add when strained

White Sugar, five pounds.

Boil to a syrup. (HANN.)

VAN MONS prescribes one pound of lichen, thirty-five of water, to be reduced to twenty, and ten of sugar.

SYRUP OF HAIRY TREE MOSS.

Syrupus de Musco querno. (WIRTEM.)

R. Hairy Tree Moss, three ounces.
Herb of Mexican Tea, } of each, one
Corinth Raisins, } ounce and a half.
Root of Comfrey, one ounce.
Liquorice, half an ounce.
Boiling Water, one pound a half.

Infuse for one night, and then boil a little.

To ten ounces of the strained and clarified liquor, add

White Sugar, sixteen ounces.

Make a syrup. When cold, add to it

Balsam of Tolu, one dram and a half.

White Sugar, one ounce and a half.

Mix.

MUCILAGINOUS DECOCTION OF ICELAND MOSS. (LOND.)

Decoctum Lichenis Islandici. (SAX.; SUEC.; DISP. CHA. PARIS; NIEMANN; RATIER; VAN MONS)

R. Iceland Moss, one part.
Cold Water, eight parts.

Macerate for half an hour; then decant, and reject the liquor. Boil the lichen then with

Water, thirty-two parts,
till this is reduced to sixteen; strain with expression. (SAX.)

R. Subcarbonate of Potass, one dram.
Water, thirty-two ounces.

Dissolve and pour upon

Washed Iceland Moss, two ounces.

After twelve hours' maceration, pour off the liquor, wash the lichen with pure water, and boil it in

Water, forty-eight ounces,
till reduced to sixteen; strain with expression. (SUEC.; NIEMANN; VAN MONS)

R. Iceland Moss, half an ounce.

After having washed it for five minutes in boiling water, boil it with

Water, three pounds,
till reduced to two. (RATIER)

R. Iceland Moss, half an ounce.

Wash it well in hot water, and add

Water, two pounds and a half.

Boil till reduced to one pound and a half; add when strained

Simple Syrup, two ounces.

Mix. (DISP. CHA. PARIS)

BITTER DECOCTION OF ICELAND MOSS.

Decoctum Lichenis Islandici. (LOND.; EDINB.; DUBL.; AMER.; FORM. HÔP. FR.; SUEC.; AUGUSTIN; COXE; ELLIS; NIEMANN; SAUNDERS)

R. Iceland Moss, one ounce.
Water, one pint and a half.

Boil down to one pint and strain. (LOND.)

EDINB. prescribes two pounds of water.

R. Iceland Moss, half an ounce.
Boiling Water, one pint.

Digest for two hours; then boil for a quarter of an hour, and strain while hot. (DUBL.)

WURCEB.; ELLIS; and NIEMANN prescribe one ounce of lichen and sixteen ounces of water; to be boiled down to

twelve ounces.—AMER.; and COXE give the same formula as LOND.—SUEC.—prescribes one ounce of lichen and twenty-four ounces of water; to be boiled down to sixteen.

R. Iceland Moss,

from half an ounce to one ounce.

Water, two pounds.

Boil down to one pound, strain, and add

Honey, one ounce.

Dissolve. (FORM. HÔP. FR.)

R. Iceland Moss, half an ounce.

Water, a sufficient quantity,

to obtain one pound and a half of strained liquor after a quarter of an hour's boiling: add

Syrup of Marshmallows, } of each,
———Poppy, } one ounce.

Mix. (SAUNDERS)

R. Iceland Moss, two drams.

Jalap, ten grains.

Cinnamon, eight grains.

Water, twelve ounces.

Boil down to six, and add

Simple Syrup, *ad libitum*.

Mix. (AUGUSTIN)

It is recommended in pulmonary consumption, leucorrhœa, and whooping cough.—Dose, four ounces, three or four times a day.

LACTEAL DECOCTION OF ICELAND MOSS.

Decoctum Lichenis Islandici lacteum. (AUGUSTIN)

R. Iceland Moss, one ounce.

Whey, twenty ounces.

Boil down to twelve ounces, and add

Sugar of Milk, six drams.

Mix.

SEDATIVE POTION.

Potio nutriens et sedans. (BRERA)

R. Washed Iceland Moss, two ounces.

Hartshorn Shavings, one ounce.

Water, a sufficient quantity,

to make a gelatinous drink; boil, and add

SYDENHAM'S Laudanum,
fifteen drops.

Mix.—To be taken gradually in the course of the day.—It is recommended in pulmonary consumption, where there is much pain.

PECTORAL DECOCTION.

Decoctum pectorale nutriens. (BRERA)

R. Iceland Moss, two ounces.

Buds of Spruce Fir, half an ounce.

Water, a sufficient quantity,
to obtain one pound of strained decoction.

STIMULANT PECTORAL DECOCTION.

*Decoctum pectorale nutriens et incitans seu
Lichenis Islandici compositum.* (HERBIP.;
AUGUSTIN; BRERA; SWEDIAUR)

R. Washed Iceland Moss, one ounce.
Cinchona, half an ounce.
Serpentaria, one dram.
Water, a sufficient quantity,
to obtain four ounces of decoction, strain,
and add

Almond Emulsion, four ounces.

Mix. (BRERA)

R. Iceland Moss, } of each,
Ground Ivy, } half an ounce.
Cinchona, two drams.
Water, a sufficient quantity,

to obtain two pounds of decoction. Add
Cow's Milk, ten ounces.

Mix. (BRERA)

R. Seneka Root, two drams.
Iceland Moss, half an ounce.
Bitter-sweet, one dram.
Water, a sufficient quantity,

to obtain four ounces of strained liquor.
Add

Cow's Milk, six ounces.

Mix. (BRERA)

R. Iceland Moss, half an ounce.
Seneka Root, one ounce.
Water, three pounds.
Boil down to two pounds. (SWEDIAUR)

R. Iceland Moss, six drams.
Scarlet Mushroom, } of each,
Root of Comfrey, } half an ounce.
—— Seneka, }
Conserve of Roses, two ounces.
Seeds of Henbane, three drams.
Alum, one dram and a half.
Water, five pounds.

Boil down to three pounds and a half,
and add when strained

Honey of Roses, one ounce and a half.

Mix. (HERBIP.)

R. Quassia Wood, two drams.
Iceland Moss, six drams.
Water, thirteen ounces.

Boil down to ten ounces, and add

Extract of Myrrh, one dram.
Tartarized Antimony, two grains.
Oxymel of Squill, one ounce.

Mix.—Dose, a spoonful every two hours.
(AUGUSTIN)

R. Iceland Moss, two ounces.

Dædalea suaveolens, }
(PERSOON)
Tops of Milfoil, } of each,
Flowers of Willow } half an ounce.
Herb,
Corinth Raisins,

Boil half an ounce of these species in
one pound of water. (AUGUSTIN)

ICELAND MOSS JELLY.

Gelatina seu Mucilago Lichenis Islandici.
(AMST.; ANT.; BATAV.; BELG.; GALL.;
FENN.; FULD.; HANN.; LIPP.; LUSIT.;
CAST. BORUS.; SAX.; SUEC.; HERBIP.;
AUGUSTIN; CADET DE GASSICOURT;
HUFELAND)

1. Without the bitter Principle.

R. Iceland Moss, washed with
alkaline liquor, one pound.
Spring Water, twelve pounds.

Boil down to one half, strain with ex-
pression, and set apart to gelatinize in a
cool place. (SUEC.)

HANN. prescribes two ounces of lichen
washed with alkaline liquor, thirty ounces
of water, reduced to six, and the addition
of four pounds of sugar.

R. Iceland Moss, one part.
Cold Water, eight parts.

Macerate for half an hour, reject the li-
quor, and pour upon the residue

Water, thirty-two parts.

Boil down to one half; then evaporate to
the consistence of honey, and set it aside in
a cool place. (SAX.)

GALL.; and CADET DE GASSICOURT di-
rect two ounces of lichen to be boiled slightly
in a sufficient quantity of water; then the
liquor to be rejected, and the residue to be
boiled twice in water; the two decoctions to
be mixed together, and one dram of dissolved
isinglass and four ounces of white sugar to
be added: the mixture to be then strained,
clarified, and evaporated to one pound.

2. With the bitter Principle.

R. Iceland Moss, one pound.
Water, twenty-pounds.

Boil down to six, express strongly, and
set aside in a cool place. (FENN.)

CAST. BORUS. prescribes one ounce of
lichen, and twelve ounces of water; to be
reduced to eight.—HUFELAND—a sufficient
quantity of lichen to bring an arbitrary
quantity of water to the consistence of mu-
cilage.—AUGUSTIN—one ounce of lichen,
and twelve ounces of water; to be reduced
to eight.

- R. Iceland Moss, three ounces.
Water, four pounds.

Boil down to one third, express, evaporate to one pound, and add
Syrup, three ounces.

Mix. (AUGUSTIN)

- R. Iceland Moss, one ounce and a half.
Water, two pounds.

Boil down to six ounces, and add
White Sugar, four ounces.

Boil to the consistence of thick syrup; and set aside to cool. (AMST.; ANT.; BATAV.; BELG.; LIPP.; LUSIT.)

This is REGNAULT'S process; he prescribed eight ounces of lichen and three pounds of water; to be boiled for a quarter of an hour, and the strained liquor to be evaporated to one pound and a half; six ounces of sugar to be added, and formed into a thick syrup.—IDT and CHEVALIER propose to boil one ounce of lichen, washed in five ounces of hot water, in a tin pan placed in a water-bath, for three hours; to strain the decoction with expression, to add one ounce of sugar to the liquor, to evaporate it slowly to four ounces, and then to pour it into a pot, and to remove from it, when gelatinized, the pellicle which forms on its surface.

3. With the bitter and an astringent Principle.

- R. Iceland Moss, } of each,
Lungwort, } six drams.
Scarlet Mushroom, }
Water, a sufficient quantity.

Boil down to one pound and a half; then, having evaporated it to the consistence of a jelly, add

Rose Sugar, one ounce and a half.

Mix. (HERBIP.)

- R. Iceland Moss, one ounce and a half.
Hairy Tree Moss, } of each,
Petals of Roses, } half an ounce.
Root of Seneka, }
Isinglass, one ounce.
Water, a sufficient quantity.

Boil for two hours, and add when strained
Oleosaccharum of Lemon, two ounces.

Mix. (AUGUSTIN)

FULD. prescribes three ounces of Iceland moss, one ounce of scarlet mushroom, four pounds of water to be reduced to one third, the evaporation of the strained decoction to the proper consistence, and the addition of three ounces of simple syrup.

This jelly is analeptic: according to the first formulæ it is simply so; according to the second, it is tonic; and according to the

third, it is astringent, and even hæmastatic. —Dose, from two to six spoonfuls.

ICELAND MOSS JELLY WITH CINCHONA.

- R. Iceland Moss, two ounces.
Isinglass, one dram.
Water, a sufficient quantity.

Boil the lichen slightly in water, and reject the liquor; boil it again twice successively, mix the two decoctions, and the isinglass in solution, add

Vinous Syrup of Cinchona, six ounces.

After a slight boiling, strain, evaporate to one half, and set aside to cool.

ICELAND MOSS-JELLY WITH OPIUM.

- R. Iceland Moss, } of each,
Hartshorn Shavings, } half an ounce.
Water, eight ounces.

Boil down to four ounces, and add,

Syrup of Opium, one ounce.

Mix.

ICELAND MOSS-PASTE.

Pasta lichenifera. (SAX.; HENRI)

- R. Iceland Moss, two pounds.
White Sugar, } of each, eight pounds.
Gum Senegal, }
Aqueous Extract of Opium, two drams.
Orange-flower Water, half a pound.

Boil the lichen in water for some time, and reject the liquor; boil again, and mix the strained decoction with the sugar and gum dissolved separately; evaporate them to the consistence of thick honey, add the extract dissolved in one part of orange-flower water; continue the evaporation, adding the remainder of the water, and boil till the mass no longer adheres to the hand. (HENRI)

- R. Iceland Moss Jelly, twelve parts.
Powdered Sugar, one part.
Gum Arabic, two parts.

Mix, while the jelly is hot. (SAX.)

PASTILLES OF ICELAND MOSS. (HENRI; FEE)

- R. Iceland Moss, one pound.
Having washed in cold water, boil it with
Water, six pounds,
for twenty minutes. Add to the strained decoction,
White Sugar, one pound.

Evaporate slowly to dryness; pulverize, mix the powder with an equal part of white sugar, and make with a sufficient quantity of water, a paste, to be divided into pastilles.

INCRASSANT ELECTUARY. (HERBIP.)

- R. Iceland Moss Jelly, two ounces.
 Gum Tragacanth, one ounce.
 Root of Comfrey, three drams.
 Conserve of Roses, six drams.
 Syrup of Poppies, a sufficient quantity.

Mix.

ANTI-CATARRHAL HYDROMEL. (CADET DE GASSICOURT)

- R. Iceland Moss, washed in boiling water, two ounces.
 Leaves of Hyssop, half an ounce.

Hydromel, two pounds.

Macerate for several days, strain and add,
 White Sugar, three ounces.
 Mix, and filter.

PECTORAL BROTH. (CADET DE GASSICOURT)

- R. Iceland Moss, half an ounce.
 White Orchard Snails, six.
 Sheep's Heart, one half.
 Calf's Lung, chopped, four ounces.
 Water, three pints.

Boil down to two pints.

LIGNUM AGALLOCHUM.—ALOE WOOD.

SYNONIMES:—*Xyloaloes*, *Lignum Agallochi veri*.—Bois d'Aloes, Bois de Calambac, Bois de Calembouc, FR.; Aloeholz, Paradiesholz, GERM.; Calumbuk, Aghaluchie, ARAB.; Aggar, Agar, BENG.; and HIND.; Sukkiang, CHIN.; Chin hiam, COCHIN.; Agfir, DUK.; Udhindi, PERS.; Pae de Aloes, PORT.; Aguru, SANS.

BRUNS.; HISP.; FERR.; WIRTEM.; AINS-
 LIE; GUIBOURT; MURRAY and GMELIN;
 SPIELMANN.

This wood is nodose, heavy, compact, unctuous, and nearly entirely composed of resin; it is of a uniform reddish-brown externally, but on being cut, shows a grey interior marked with black spots. It has a strongly balsamic odour and a bitter taste. When burnt, it diffuses a very agreeable perfume.

It is attributed to the *Excacaria Agallocha*, LINN.; a tree of Cochinchina. (*Diæcia*, *Triandria*, LINN.; *Euphorbiaceæ*, JUSS.; *Fig.* Rumph, *Amb.* II. t. 79, f. 80)

It is tonic and excitant.

CORDIAL SPECIES.

Species cordiales temperatæ. (WIRTEM.)

- R. Burnt Raw Silk, } of each,
 Ivory, burnt to whiteness, } half
 Red Coral, } an ounce.
 White Coral, }
 Prepared Pearls, one ounce and a half.
 Root of Doronica, } of each,
 ——— White Behen, } three drams.
 ——— Red Behen, }
 Aloe Wood, half an ounce.
 Cinnamon, } of each,
 Cloves, } three drams.
 Bone of Stag's Heart, }

- Hyacinth, }
 Emerald, } of each, one dram.
 Sapphire, }
 Saffron, } two scruples.

Make a powder, and add,

- Gold Leaves, } of each, *ad libitum*.
 Silver Leaves, }

Mix.—This absurd preparation used formerly to be given in the dose of from half a scruple to half a dram in malignant fevers, palpitation of the heart, and fainting occasioned by fear.

EXTRACT OF ALOE WOOD.

Extractum Ligni Aloes. (BRUNS.; PALAT.;
 WIRTEM.)

- R. Aloe Wood, any quantity.
 Concentrated Alcohol, a sufficient quantity

to exhaust the wood completely; evaporate the tincture in a water-bath, and preserve the residue.

This extract was formerly added to cephalic pills and powders.—Dose, one or two grains.

TINCTURE OF ALOE WOOD.

Tinctura seu *Essentia Ligni Aloes.*
 (BRUNS.; HISP.; WIRTEM.)

- R. Aloe Wood, rasped, one ounce and a half.
 Concentrated Alcohol, half a pound.
 Macerate, and filter. (BRUNS.; WIRTEM.)

HISP. prescribes one part of wood and twelve parts of alcohol.

It is excitant, and is reputed stomachic, uterine, cephalic, and analeptic.—Dose, from twenty to forty drops.

On the first of January, 1832, will be published,

A NEW PERIODICAL WORK, ENTITLED

THE

PICTORIAL HISTORY

Of the Bible :

CONSISTING OF THE DIVINE INSPIRATIONS OF THE GREATEST MASTERS,
ARRANGED IN A CHRONOLOGICAL SERIES, AND

ENGRAVED BY THE FIRST ARTISTS OF THE PRESENT DAY.

CONDITIONS OF PUBLICATION.

I. The PICTORIAL HISTORY will be complete in 20 Numbers, price one shilling each, Demy Quarto, containing two Plates in each Number.

II. A limited impression will be printed on India paper, price 2s. each.

III. The Work, which is now far advanced, will be continued regularly every month till the whole is completed.

It is unnecessary to apologize for the appearance of a New Pictorial Work at a period when a taste for the Fine Arts, keeps pace so fairly, in this Country, with a love of Science, and the diffusion of Knowledge.

ENGRAVING is to *Painting*, what *Printing* is to *Writing*,—that valuable Art by which the Original Works of Great Authors and Artists are multiplied to infinity, for the Instruction and Delight of Mankind.

Painting (the Parent of Engraving) was the first attempt to represent Thought, Hieroglyphics being a kind of Stenography, or short hand of Painting, in common use in Egypt—

“The Queen of Nations, and the boast of Times,
Mother of Science, and the Land of Gods!”

long before the invention of the Alphabet by Cadmus. The wonderful genius of the Greeks carried every Art to perfection. The refinement to which this, in particular, appears to have been carried at Athens by Apelles, Zeuxis, Protogenes, and Polygnotus, which latter adorned the celebrated *Athenian Portico* with the most striking events of the Trojan War, may be shewn by the fastidiousness of *Zeuxis* in condemning a Painting of his own, because it was not *truly natural*. The subject was a boy holding a bunch of grapes, which the birds flew to peck at : but the Painter, who was the Critic, remarked, that if the *boy* had been as faithfully represented as the *grapes*, the birds would have been afraid of him. He contended with Parrhasius which of them was the better Painter ; and when he went to see his Rival's production, he begged him to remove the curtain and display it. How great was his astonishment, when he understood the *curtain* was absolutely the work in question. You have the advantage of me, he said, I have only deceived *birds*, but you have deceived the *Master* of the Art. These Anecdotes serve to shew the necessity of Genius in a Painter. Like the Poet he must be a Lover and an Admirer of Nature—a Child of Hyperion—whose soul is so deeply impregnated by her sublimities and beauties, that his pencil may call them forth to life. Many indeed have been the disputes between the Poets and the Painters for the *precedence* of their Art Divine. Simonides, Horace, Du Fresnoy and others have elegantly defined Painting, or Picture, as *silent Poetry*—Poetry as *speaking Picture*—and this old family quarrel for precedence, has of late years been revived by the present Talented President of the Royal Academy (Sir Martin Archer Shee) in his “Rhymes on Art ;” where he maintains, that “the *Narrative* of an Action is not comparable to the *Action itself* before the eyes.” The enthusiast Barry considers Painting as “Poetry realised,” and Richardson and Darwin assert, that the “Essence of Poetry is Picture.” On this very interesting subject, the great Philosopher Plato remarks, “God Almighty, in the fabric of the Universe, first contemplated himself, and reflected on his own Excellencies, from which he drew and constituted those first forms, which are called Ideas, so that every species which was afterwards expressed, was produced from that first perfect Idea, forming that wonderful texture of all created beings.” Nature therefore always intends a consummate beauty, while the inequality of matter tends always towards disproportion. “For which reason (he adds) the skilful Painter, and the Sculptor, imitating the Divine Maker, form to themselves, as well as they are able, a model of the superior beauties, and reflecting on them, endeavour to correct, and amend the common

nature, and to represent it as it was first created, without fault, either in colour, or in lineament." This *ideal beauty* descends equally upon the marble, the cloth, or the paper, of the Sculptor, the Painter, and the Poet. What Michael Angelo is in Painting, so is Praxiteles in Sculpture, and Milton in Poetry.

From Greece the Spirit of this Divine Art fled to Italy, and was cultivated to perfection by Raffaele, Titian, Correggio, and Michael Angelo: Holland next caught the inspiration, and Holbein, Reubens, and Vandyck appeared; and at the present day the English and French vie with the most celebrated Masters of the Italian and Flemish Schools.

But if Painting is so Divine, what enthusiasm must the Painter feel at the discovery of an Art, which, as in a looking-glass, would multiply his Original, and send it round to an Admiring World!—ENGRAVING is, in fact, this *Mirror*,—and hence we find, that the most celebrated Painters of these latter Ages have *themselves* cultivated the Art of Engraving, as the surest method of placing their labours upon record among Mankind. ENGRAVING expresses this *ideal beauty* of *Painting*, in like manner, as *Painting* expresses this *ideal beauty* of *Nature*. Good Engravings are equally as imposing as Paintings; hence *Galleries* have been formed by the Public, and *Collections* by Private Virtuosi—proofs of the delight which this sister and rival Art is capable of affording. Indeed, herein consists the *great advantage* of Engraving over Painting, that not only may *Galleries* of the same Engravings be multiplied, but they may be even compressed into a *Book*,—a *Portfolio*,—or an *Album*, and become a *Pictorial Gallery* for the Instruction of the Study, or the Amusement of the Drawing Room. And, as in Painting, the Historical Painters are the Princes of the Art, so in Engraving, have *Historical Engravings* ever been prized as the most valuable productions of the Graphium. History then—the natural scene of Man—is the grand Study of the Painter and Engraver. But if this is true, what History affords the most copious source? It is the *Sacred History*—a Volume allowed, not only by the whole Christian World, but even by the Sceptics and Infidels themselves, to abound with the *finest subjects* for Painting, whether Historical, Spiritual, or Allegorical. The Scriptures have ever been the Study of the greatest Masters in the Art; and, hence has been illustrated by the very Genius of Painting and Engraving, the most curious, and the most wonderful, as well as the most Ancient Book in the whole World.

But although there have been many Illustrations of the Holy Scriptures, there has never yet appeared a PICTORIAL HISTORY OF THE BIBLE;—a *Book*, which, by means of a Series of *Chronological Engravings* (executed in the best style), shall answer all the purposes of the *Text* of the Original, and the *Notes* of Commentators. In this Enlightened Age, when the Bible is known by heart, it is conceived that it will create a *new Intellectual Pleasure* to know it by *Scenic Illustration*, which is a species of Acting, or Dramatic Exhibition of the Scriptures (a practice much in use by the Christian Fathers of the Early Ages). The eye in wandering over such a *Chronological Series of Engravings* (as is now proposed), acts strikingly upon the mind, and, by that process of thought called concatenation, links its ideas together into a complete History. Thus the Eye is delighted by the Imagery, and the Mind improved by the Operation; the subject itself is strengthened in the Memory, and the whole Bible perused even in a few minutes. Such a Pictorial History, or Paraphrase, or Summary, or Analysis, or Abridgment, or Index, (for it is all of these) it is conceived will supply the want of the *Text* itself, and render the *Notes* of some commentators useless. Hogarth was the first who conceived a similar design to represent the History and Fortunes of one Individual, but the same idea has never been, until now, applied to the Inspired Writings of the Old and New Testaments.

The Projectors of this extensive Undertaking, convinced of its great importance and general utility, have resolved to supply such a *Desideratum* in Literature and Art, by furnishing a *Chronological Series of Engravings from the Old and New Testaments*, (Historical, Cosmographical, Geographical, and Topographical,) under the Title of THE PICTORIAL HISTORY OF THE BIBLE, taken from the most Interesting and Instructive Subjects, of the English and other Schools, and executed by the first Artists of the day.

The Work will possess two other advantages, viz. it will not only, as described, be in itself THE PICTORIAL HISTORY OF THE BIBLE, but it will serve both as the *Illustrations* to any *Family Bible* whatever, whether of a Folio or Quarto Edition, which may not be ornamented with Engravings; and it will form the best vehicle for the *Father of a Family* to put into the hands of his *Children* to allure them to the knowledge of the important truths of Holy Writ.

The Proprietors, influenced by these important considerations, have also decided to offer this Work to the Public, on the *most Liberal and Economical Plan*, trusting alone for remuneration to an *extensive sale*; and they pledge themselves that the whole Series of Engravings shall be executed in the same masterly manner as those contained in the First Number, which is offered as a *Specimen*.

LIGNUM ASPALATHI SEU AQUILÆ.

SYNONIMES :—Bois d'Aigle, FR. ; Adlerholz, GERM.

BRUNS. ; WIRTEM. ; BERGIUS ; GUIBOURT.

This wood is of a dull-yellow colour, rather resinous, fibrous, sometimes spongy, difficult to be divided by the teeth. Its

taste is aromatic ; its odour is feeble and musky.

It is attributed to the *Aquilaria ovata*, CAV., a tree of the East Indies. (*Decandria Monogynia*, LINN. ; *Fig. Cav. Diss. Bot.* 7, p. 377, tab. 224)

It is stimulant, but is rarely employed.

LIGNUM BRASILIENSE.—BRAZIL WOOD.

SYNONIMES :—*Lignum Brasilianum rubrum* seu *Fernambuci*.—Bois de Bresil, Bois de Fernambouc, FR. ; Brasiliensholz, GERM. ; Ibirapitanga, BRAZ.

BRUNS. ; GALL. ; LIPP. ; GENEV.

This wood is heavy, hard, inodorous, of

a scarcely sensible taste, and of a red colour presenting different shades.

It belongs to the *Cæsalpinia echinata*, LMK., a tree of Brazil. (*Decandria Monogynia*, LINN. ; *Leguminosæ*, JUSS. ; *Fig. Piso, Bras.* p. 164, 1c.)

It is astringent, and used formerly to be prescribed in diarrhœa, but is now almost abandoned.

LIGNUM CAMPECHIANUM.—LOGWOOD, CAMPECHY WOOD.

SYNONIMES :—*Lignum Campechiense* seu *Campescanum* seu *Brasilianum rubrum* seu *Sappan* seu *cæruleum*.—Bois de Campeche, FR. ; Campescheholz, Blauholz, GERM. ; Campesketræt, DAN. ; Palo de Campeche, SPAN. ; Campechehout, DUT. ; Legno de Campeggio, ITAL. ; Niebieskie drzewo, POL. ; Pao de Campeche, PORT. ; Campchetraed, SWED.

LOND. ; DUBL. ; EDINB. ; AMST. ; ANT. ; BATAV. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; FULD. ; GENEV. ; HAMB. ; HANN. ; LIPP. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY and GMELIN ; PARIS.

This wood comes to us deprived of both its outer and inner bark ; it is hard, rather solid, heavy, and of a greyish-brown colour externally. When recently cut it has a deep red colour, with a yellowish tinge and longitudinal blackish veins, when sawed transversely, it presents small waved rings of a deep-red colour. Its taste is sweetish, astringent, with a bitterish after-taste. It tinges the saliva with bluish-red.

VOL. II.

It belongs to the *Hæmatoxylum Campechianum*, LINN. ; a tree of America. (*Decandria Monogynia*, LINN. ; *Leguminosæ*, JUSS. ; *Fig. Desc. Fl. Ant.* 1. 73)

CHEVREUL has found in it a peculiar colouring principle, *Hæmatinum*, *Hæmatoxylum*.

It is astringent and tonic.

DECOCTION OF LOGWOOD.

Decoctum Hæmatoxyli seu *Ligni Campechiani* seu *campechensis*. (HISP. ; FULD. ; HERBIP. ; ELLIS ; NIEMANN ; SARD. ; SWEDIAUR)

R. Logwood, one ounce.
Water, four pounds.

Digest, boil to one half, and strain. (HISP.)

ELLIS ; and NIEMANN prescribe from two to four drams of wood and one pound of water ; to be reduced to six ounces. — SAUNDERS—one ounce of wood and a sufficient quantity of water to obtain one pound of strained decoction. — FULD.—one ounce of wood and two pounds of water, to be reduced to one pound.

- R. Logwood, one ounce.
 Water, one pound.
 Boil down to one half, and add, if required,
 Tincture of Cinchona, one ounce.
 Opium, thirty drops.

Mix.

- R. Logwood, two ounces.
 Spring Water, four pounds.
 Boil down to two pounds and a half, and add,

Alcohol, four ounces.
 Cinnamon, four scruples.

Boil for a few moments, and strain.
 (HERBIP.)

It is astringent, and is recommended in diarrhœa, chronic dysentery, and hæmorrhages.—Dose, two ounces, every four hours or more frequently.

EXTRACT OF LOGWOOD. (LOND.; DUBL.;
 EDINB.; AMER.; FULD.; HANN.; POL.;
 BORUS.; SAX.; HERBIP.; COXE)

Extractum Hæmatoxyli. (LOND.)

Ext. Hæmatoxyli Campechiani.
 (EDINB.)

Ext. Scobis Hæmatoxyli. (DUBL.)

- R. Logwood, one part.
 Water, eight parts.

Boil down to one half, after twenty-four hours' maceration; then strain while hot, and evaporate to the proper consistence.

AMER.; and COXE give the same formula.

- R. Logwood, one part.
 Water, eight parts.

Macerate for twenty-four hours, then boil for a quarter of an hour, and express; boil the residue with four parts of water. Mix the decoctions, set aside for twenty-four hours, and evaporate. (SAX.)

- R. Logwood, one part.
 Boiling Water, eight parts.

Boil for a quarter of an hour, and express; boil the residue with four parts of

water, and express again. Mix, decant after sufficient repose, and evaporate. (BRUNS.)

HANN. directs one part of wood to be boiled in six parts of water twice or thrice in succession, and the mixed strained liquor to be evaporated.—POL.—it to be boiled at four different times in eight pounds of water, and for a quarter of an hour each time.

- R. Logwood, any quantity.
 Alcohol, a sufficient quantity

to cover it by four fingers' breadth; digest in a moderate heat, and then decant: boil the residue for one hour in water, express, and evaporate the liquor in a water-bath to the consistence of honey, gradually adding the tincture, constantly stirring, that a uniform mass may be obtained. (FULD.; HERBIP.)

Dose, from one dram to two drams, in any proper vehicle.

ASTRINGENT POTION. (ELLIS)

- R. Extract of Logwood, three drams.
 Water, seven ounces.
 Tincture of Catechu, two drams.

Mix.—Dose, two spoonfuls, every three or four hours; in dysentery and diarrhœa.

ASTRINGENT ELIXIR. (HAMB.; AUGUSTIN)

- R. Extract of Logwood, two drams.
 Distilled Water, four ounces.
 HALLER'S Acid Elixir, one dram.

Mix. (HAMB.)

- R. Extract of Logwood, half an ounce.
 Red Wine, two ounces.

Mix.—Dose, sixty drops, every two hours. (AUGUSTIN)

ASTRINGENT MIXTURE. (SWEDIAUR)

- R. Extract of Logwood, three drams.
 Spirituous Cinnamon Water, one ounce and a half.

Mix, and add,

Water, seven ounces.
 Tincture of Kino, two drams.

Mix.—Dose, two spoonfuls every four or five hours.

LIGNUM EBENUM.—EBONY WOOD.

WIRTEM.; GENEV.; SPIELMANN.

A dense, heavy, black wood, of a sensible taste.

It is from the *Diospyros Ebenum*, LINN.,

a large tree of the East Indies. (*Polygamia Diæcia*, LINN.; *Ebenaceæ*, JUSS.; *Fig. Rumph. Amb. p. 13. t. 6.*)

It used formerly to be considered as a sudorific equal to Guaiac wood.

LIGNUM NEPHRITICUM.—NEPHRITIC WOOD.

SYNONIMES :—Bois nephritique, FR. ; Griesholz, GERM. ; Palo nefritico, SPAN.

BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; BERGIUS ; GUIBOURT ; MURRAY and GMELIN ; SPIELMANN.

This wood is very heavy. It is formed of a whitish, compact alburnum, and of a reddish, very hard axis ; the bark is light, fi-

brous internally, fungus and cracked externally. Its taste somewhat resembles that of pepper.

It is supposed to belong either to the *Guilandina Moringa*, LINN., to the *Mimosa unguis cati*, LINN., or, according to VIREY, to a *Cissampelos*. It comes from New Spain.

Powerful diuretic virtues were formerly attributed to it. It is rarely used at present.

LIGNUM RHODII.—RHODIUM WOOD.

SYNONIMES :—Bois de Rhodes, Bois de rose, Bois de Chypre, FR. ; Rosenholz, Rhodisherholz, GERM.

AMST. ; BATAV. ; BRUNS. ; HISP. ; FERR. ; SAX. ; SUEC. ; WIRTEM. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; SPIELMANN.

This wood is hard, heavy, of a fawn-yellow colour, deeper in the centre than at the circumference, and formed of very close concentric layers. Its taste is bitterish, and its odour strongly rosaceous, particularly when rasped.

It is generally believed to come from the root and sometimes from the stump of the *Convolvulus scoparius*, LINN., a large tree of the Canary Isles. (*Pentandria Monogynia*, LINN. ; *Convolvulaceæ*, JUSS.)—Some authors ascribe it to the *Genista Canariensis*, LINN. ; but it most probably belongs to the former.

It is stimulant, but is not much used.

OIL OF RHODIUM WOOD. (BRUNS. ; GALL. ; FULD. ; PALAT. ; WIRTEM.)

R. Rhodium Wood, rasped,	ten parts.
Muriate of Soda,	one part.
Water,	twenty parts.

Macerate for twelve hours ; then distil, and collect the oil which sinks to the bottom of the receiver. (GALL.)

BRUNS. ; and WIRTEM. prescribe six parts of wood, one part of muriate, and thirty-six parts of water.—FULD.—one of wood and four of water ; to be digested for four days.—PALAT.—one of wood and six of water ; to be digested for five or six days.

This oil is rather used as a perfume than as a medicament.

DISTILLED WATER OF RHODIUM WOOD. (GALL.)

R. Rhodium Wood, rasped,	two parts.
Water,	sixteen parts.

Infuse for twelve hours, and distil eight parts.

LIGNUM SANTALI.—SANDERS WOOD.

There are three kinds of this wood :

1. *Lignum Santali rubri*.—Red Sanders.

SYNONIMES :—Santal rouge, FR. ; Rothes Santelholz, Kallaturholz, GERM. ; Sundal ahmer, Undum, ARAB. ; Czerweny santal, Cerwené santalowe drewo, BOHEM. ; Rackta chandana, BENG. ; Honnay, CAN. ; Ruct handun, CYNG. ; Lal chudend, DUK. ; Sandalo rubro, SPAN. ; Ruck ut chundun, Undum, HIND. ; Root

zandelhout, DUT. ; Santilaparni, Ranjana, Rakta Chandana, SANS. ; Segahu chandannum, TAM. ; Kuchandanum, TEL.

LOND. ; DUBL. ; EDINB. ; ANT. ; BATAV. ; BRUNS. ; HISP. ; GALL. ; SUEC. ; WIRTEM. ; AINSLIE ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

This wood is met with in pieces of various sizes, without bark, cut longitudinally, of a fine red colour, grooved with fibres, between

which traces of concrete resinous juice may be seen. It has a marked aromatic odour, and a rather resinous taste.

It is furnished by the *Pterocarpus Santalinus*, LINN., a tree of the East Indies. (*Diadelphia Hexandria*, LINN.; *Leguminosae*, JUSS.)

PELLETIER has discovered in it a peculiar colouring matter, *Santalina*.

2. *Lignum Santali albi*.—White Sanders.

SYNONIMES :—Santal blanc, FR.; Weisses Sandelholz, GERM.; Ayasru, AMB.; Sandel abiez, ARAB.; Chandana, BENG.; Taw-muh, CHIN.; Frigunda, CAN.; Cay-huyndhdan, COCHIN.; Sandun, CYNG.; Sundel, DUK.; Chunden, Chandana, Katchandan, HIND.; Sandalo bianco, ITAL.; Tsjendana, MAL.; Sundul sufleid, PERS.; Chandana, Malayaja, SWED.; Chandanum, Chandana-cuttay, TAM.; Chendanum, TEL.; Aikamenil, TIMOUR.

BATAV.; BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

This wood is in pieces cut lengthwise, heavy; its bark is of a blackish-grey colour and rough. Its colour is yellowish white; it has a feebly aromatic odour, and is almost destitute of taste.

3. *Lignum Santali Citrini*.—Yellow Sanders.

SYNONIMES :—Santal citrin, FR.; Gelbes Sandelholz, GERM.; Zlutý santal, Zlute santalowe drewo, BOHEM.; Sandalo citrino, ITAL.

ANT.; BATAV.; BRUNS.; HISP.; GALL.; WIRTEM.; GUIBOURT.; ZARDA.

This wood has a yellowish colour, and is of a finer grain than the preceding. It is lighter, more bitter, and is usually met with free from bark. It has a very marked aromatic odour, but is almost destitute of taste.

The two latter are derived from the *Santalum album*, LINN., a tree of the East Indies. (*Tetrandria Monogynia*, LINN.; *Onagracea*, JUSS.; *Fig. Rumph. Amb. II. p. 42. t. 11*)

HERMANN considers the white kind as the alburnum and the yellow as the pith of the wood. According to MERAT, the former is the young wood and the latter the matured. Yellow sanders is stimulant and diaphoretic; the red is slightly astringent.

DECOCTION OF YELLOW SANDERS WOOD.

Decoctum Ligni Santali citrini. (NIEMANN)

R. Yellow Sanders, rasped, half an ounce.

Spring Water, two pounds.
Boil down to one half.—Dose, a glassful.

OIL OF YELLOW SANDERS WOOD.

Oleum Santali citrini æthereum Ætheroleum, Santali citrini. (WIRTEM.)

R. Yellow Sanders, rasped, six pounds.
Water, thirty-six pounds.
Muriate of Soda, one pound.

Macerate for four or five days; distil, and separate the oil, which is at the bottom of the receiver.

SANTALINE OINTMENT.

Unguentum santalinum, Emplastrum santalinum seu incognitum. (BRUNS.; SAX.; SARD.; WIRTEM.)

R. Oil of Myrtle, three pounds.
White Wax, nine ounces.
Dried Petals of Red } of each,
Roses, } four ounces.
Red Sanders, }
Camphor, half an ounce.

Melt the two first together, add the roses in powder, then the sanders in powder, and lastly the camphor dissolved in a little oil.

R. White Sanders, } of each, six drams.
Yellow Sanders, }
Red Sanders, }
Fresh Petals of Red Poppies, one ounce and a half.
Roses, two ounces and a half.
Rose Oil, one pound.
White Wine, four ounces.

Macerate for two days, boil to the consumption of the humidity, express, and add,

Armenian Bole, } of each,
White Wax, } three ounces.
Camphor, two drams.

Mix. (WIRTEM.)

R. Powdered Red Sanders, six parts.
Saffron, one part.
Olibanum, } of each,
Mastic, } one
Myrrh, } twelfth part.
Alum, }

Melt together, and add to the mass when nearly cold,

Resin, } of each,
Yellow Wax, } sixteen parts.

Mix. (SAX.)

R. Wax, } of each, four ounces.
Resin, }

Melt together, and add,

Red Sanders, one ounce and a half.
Saffron, two drams.

Olibanum, }
 Mastic, } of each,
 Myrrh, } one dram and a half.
 Alum, }

Mix. (BRUNS.; WIRTEM.)

SPECIES OF THE THREE SANDERS WOODS.

Species diatron santalon. (PALAT.; WIRTEM.)

R. White Sanders, }
 Yellow Sanders, } of each, two drams
 Red Sanders, } two scruples.
 Petals of Roses, }
 three ounces three drams.

Petals of Violets, } half an ounce.
 Rhubarb, }
 Seeds of Purslain, } of each,
 Ivory, burnt to whiteness, } two
 Extract of Liquorice, } drams.
 Starch, }
 Gum Arabic, } of each,
 — Tragacanth, } four scruples.

Mix.—A scruple of camphor may be added.
 —The older practitioners believed that the sanders woods corrected irregular heat of the liver, on account of which property they prescribed these species in slow fevers, jaundice, and dropsy, in the dose of from half a dram to one dram.

LIGUSTICUM.

Two species of this genus of plants are employed.

1. *Ligusticum Levisticum*, LINN.—Lovage.

SYNONIMES:—Liveche, FR.; Liebstöckel, Bærmutterkraut, GERM.; Apich welliky, Apich, Libecek, BOHEM.; Lovstilk, Lobstik, DAN.; Ligustico, SPAN.; Laveskruid, DUT.; Livistico, ITAL.; Lakotnegozicle, POL.; Levistico, PORT.; Libbsticke, SWED.

AMST.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of the temperate climates of Europe. (*Pentandria Digynia*, LINN.; *Umbellifera*, JUSS.; *Fig. Zorn, Ic. pl. t. 233*)

The root and seeds are employed.

The root (*Radix Levistici* seu *Lybistici* seu *Ligustici* seu *Laserpitii Germanici*) is about a foot long, fleshy, branching, of a brown-yellow colour without, and rugose and whitish within. It has a strong and penetrating odour, and a taste which is at first sweet, but afterwards acrid and nauseous.

The seeds are oblong, five-sided, and yellowish.

EXTRACT OF LOVAGE.

Extractum Ligustici seu *Levistici*. (OLDEN.; POL.; BORUS.; SAX.)

R. Root of Lovage, } of each, two pounds.
 Alcohol, }

Water, nine pounds.

Digest for twelve hours; then express, distil off the alcohol, and evaporate the residue to the proper consistence. (POL.)

OLDEN.; BORUS.; and SAX.—prescribe two parts of root, three of alcohol, and nine of water.

TINCTURE OF LOVAGE.

Essentia Levistici. (WIRTEM.)

R. Root of Lovage, two ounces.
 Seeds of Lovage, one ounce.
 Alcohol, ten ounces.

Digest, express, and filter.

It is excitant, carminative, diuretic, and reputed alexipharmic.—It has been recommended in dropsies.

2. *Ligusticum Meum*, Ro.—Bawdmoney.

SYNONIMES:—Meum, FR.; Bærwurz, Hundspetersilic, Herzwurzel Mutterwurz, Bærenfenchel, GERM.; Koprnik, BOHEM.; Biverne dild, DAN.; Meo, ITAL.; and SPAN.; Beelwortel, DUT.; Olesnik, POL.; Bjvern dill, SWED.

ANT.; BELG.; BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of Europe. (*Pentandria Digynia*, LINN.; *Umbellifera*, JUSS.; *Fig. Zorn, Ic. pl. t. 499.*)

The root (*Radix Mei seu Mei athaman-tici seu Anethi seu Feniculi ursini*) is employed; it is from six to eight inches long, of the thickness of the finger, wrinkled, of a blackish-grey colour without, and white

within; it has a strong balsamic odour, and an aromatic and warm taste.

It is stimulant, diuretic, and expectorant, but is not much used.

LIGUSTRUM VULGARE, LINN.—PRIVET.

SYNONIMES. — Troene, FR.; Liguster, DAN.; DUT.; GERM.; and SWED.; Alhena Germanica, SPAN.; Ligustro, ITAL.; Zoba, POL.; Altena, PORT.

WIRTEM.; SPIELMANN.

A shrub of Europe. (*Diandria Digynia*, LINN.; *Jasmineæ*, JUSS.; *Fig. Zorn Ic. pl. t. 112.*)

The leaves and flowers are employed.

The leaves (*Folia Ligustri*) are lanceo-

lated, very entire, smooth, and of a bitter and astringent taste.

The flowers are disposed in small racemes; they are white, and of an agreeable odour.

It is slightly astringent.

DISTILLED PRIVET WATER. (PALAT.)

R. Privet Flowers, one part.
Water, three parts.
Distil one part and a half.

LILIUM CANDIDUM, LINN.—WHITE LILY.

SYNONIMES:—Lis blanc, FR.; Weisse Lillie, GERM.; Lilium bjle, BOHEM.; Heide Lillie, DAN.; Azucena blanca, SPAN.; Witte Lellie, DUT.; Liglio bianco, ITAL.; Silia, POL.; Lirio branco, PORT.; Hurt Lilja, SWED.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; LUSIT.; SAX.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

This plant is originally of the Levant, but is cultivated in our gardens. (*Hexandria Monogynia*, LINN.; *Liliaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 462.*)

The bulb, flowers, and anthers are employed.

The bulb is oval, yellowish, scaly without, and furnished with large fasciculated fibres. It is destitute of odour, but has a sub-acrid taste.

The flowers consist of five white channelled petals; they have an agreeable odour, which is lost by desiccation.

The anthers are oblong and yellow.

CONSERVE OF LILY.

Conserva Florum Liliorum alborum.
(GENEV.)

R. Petals of Lillies, one part.
White Sugar, three parts.
Triturate together.

PULP OF LILY.

Pulpa Lili. (GALL.)

R. Bulbs of Lillies, any quantity.

Having cleansed them, wrap them in paper, and place them in hot ashes till they are sufficiently soft to be bruised between the fingers; then take off the exterior and scorched parts, pound the rest in a stone mortar, and pass the pulp through a close sieve.

DISTILLED WATER OF LILY. (GALL.; GENEV.; PALAT.)

R. Fresh Flowers of Lily, one part.
Water, four parts.

Distil one half. (GALL.)

PALAT. prescribes one part of flowers and three parts of water; one part and a half to be distilled.—GENEV.—four pounds of flowers and a sufficient quantity of water to prevent empyreuma; six pounds to be distilled.

OIL OF LILY.

Oleum Liliorum alborum. (AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; PALAT.; SARD.; WIRTEM.)

R. Fresh Petals of Lily, one part and a half.
Olive Oil, three parts.
Digest for two days in a water-bath; ex-

press when cold, and repeat the maceration. (SARD.)

GALL. prescribes one part of petals, four parts of oil, and three successive macerations, the last to be for three months.

R. Petals of Lily, one part.
Olive Oil, four parts.

Digest for some hours in a water-bath, and express. (ANT.; BATAV.; BELG.)

R. Petals of Lily, one part.
Olive Oil, three parts.

Macerate for a few days in a warm place;

then boil to the consumption of humidity, and express. (AMST.; HISP.)

R. Petals of Lily, } of each, an equal part.
Olive Oil, }

Boil to the consumption of humidity, and express. (PALAT.)

R. Petals of Lily, two parts.
Olive Oil, three parts.

Digest for eight days, boil, and express. (BRUNS.; WIRTEM.)

It is emollient, and reputed anodyne.

LINARIA.—TOAD-FLAX.

Three species of this genus are used in medicine :

1. *Linaria Cymbalaria*, MILL.—Ivy-leaved Toad-flax.

SYNONIMES:—Cymbalaire, Linaire de montagne, FR.; Zymbelkraut, GERM.

WIRTEM.; SPIELMANN.

A common plant of Europe. (*Didynamia Angiospermia*, LINN.; *Scrofularia*, JUSS.; *Fig. Curt. Fl. Lond.* t. 55.)

The herb is employed: it consists of a slender, glabrous stem, and of alternate, petiolated round leaves, heart-shaped at the base, and divided into five or seven lobes. Its taste is bitterish.

It is esteemed vulnerary.

2. *Linaria spuria*, MILL.—Bastard Toad-flax.

SYNONIME:—Linaire bâtarde, FR.

GALL.; SPIELMANN.

This plant is common all over Europe. (*Fig. CEd. Fl. Dan.* t. 913.)

The flowering herb is employed which consists of a branching stem, of hairy, oval, entire leaves, and of yellow flowers, with the inferior lip of each violet coloured. The flowers are produced solitary in the axillæ of the leaves, on long filiform peduncles.

3. *Linaria vulgaris*, MÆ.—Yellow Toad-flax.

SYNONIMES:—Linaire commune, FR.; Gemeines gelbes flachskraut, Leinkraut, GERM.; Vild torskemand, DAN.; Linaria, ITAL.; and SPAN.; Het gemeene

Vlachskruid, DUT.; Lennek pauny maigi, POL.; Flmysblomster, SWED.

BAVAR.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; BORUS.; SAX.; WIRTEM.; BERGIUS.; BRUGNATELLI.; MURRAY and GMELIN.; SPIELMANN.

A common plant of Europe. (*Fig. Zorn, Ic. pl.* t. 442.)

The flowering herb (*Herba et Flores Linariæ seu Antirrhini*) is employed: it consists of a simple glabrous stem, of numerous lanceolated, sessile leaves of a glaucous green colour, and of yellow flowers clustered into a terminal spike. It has a rather disagreeable odour, which is removed by drying. Its taste is bitter and nauseous.

OINTMENT OF TOAD-FLAX.

Linimentum Anti-hæmorrhoidale. (BAVAR.; BRUNS.; HANN.; LIPP.; OLDEN.; BORUS.; SAX.; SARD.; WIRTEM.; PIERQUIN.; SPIELMANN.)

R. Fresh flowering Herb of Toad-flax, one pound.
Lard, two drams.

Boil slowly to the consumption of the humidity, and express. (BAVAR.; HANN.; OLDEN.; BORUS.; SAX.)

BRUNS.; and WIRTEM. prescribe an equal part of each.—LIPP.—one of herb and three of lard.—PIERQUIN fixes no proportions.

R. Herb of Toad-flax, twelve ounces.
Houseleek, two ounces.
Henbane, one ounce.
Butter, eighteen ounces.

Macerate for several days; then boil to the consumption of the humidity, and express. (SPIELMANN)

R. Flowering Herb of Toad-
flax,
Lard, } of each,
 } one pound.

Boil to the consumption of the humidity,
express, and add

Oil of Eggs, two drams.
Expressed Oil of Henbane Seeds,
 three drams.

Mix. (SARD.)

LINUM, LINN.—FLAX.

Two species of this genus of plants are employed in medicine :

1. *Linum usitatissimum*, LINN.—Lint; Flax.

SYNONIMES:—Lin ordinaire, FR.; Lein, GERM.; Buzruk, ARAB.; Len, BOHEM.; and POL.; Hær, DAN.; Ulsikebinge, DUK.; Lino, ITAL.; and SPAN.; Tisi Pahaha, HIND.; Vlasch, DUT.; Bidgierammi, MAL.; Tokhemkutan, PERS.; Linhaca, PORT.; Uma, Atasi, SANS.; Lin, SWED.; Alleverei, Serru sanulverei, TAM.; Alivitulu, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

It is probable that this plant is originally from the East; it is spread over the greater part of Europe. (*Pentandria Digynia*, LINN.; *Linée*, CAND.; *Fig*, ZORN, *Ic. pl. t.* 251.)

The seeds (Linseed; *Semines Lini*) are brown, shining, oval, pointed, flattish, very smooth, acute at the margin. Under the cuticle, it contains a white kernel of a disagreeable taste.

They contain a considerable quantity of oil in their kernels, and of mucilage in their cuticle. They are emollient.

2. *Linum catharticum*, LINN.—Purging Flax.

SYNONIMES:—Lin purgatif, FR.; Purgierflachs, GERM.; Liven vild hær, DAN.; Cantilagua, SPAN.; Purgeervlasch, DUT.; Lino purgativo, ITAL.; Linho purgante, PORT.; Villhær, SWED.

LOND.; DUBL.; GALL.; GENEV.; LIPP.; ROSS.; SAX.; WIRTEM.; BERGIUS; COXE; MURRAY and GMELIN; PARIS; SPIELMANN.

A plant of Europe. (*Fig*, ZORN, *Ic. pl. t.* 210.)

The herb (*Herba Lini purgativi seu cathartici*) is employed. It consists of a slender stem, furnished with dichotomous branches, and with oval, oblong, opposite, glabrous leaves. It is without smell, but its taste is bitter and nauseous.

It is seldom used.

EMOLLIENT MEALS. (GALL.; SPIELMANN)

R. Linseed Meal, } of each,
Fenugreek Meal, } an equal part.

Mix. (SPIELMANN)

GALL. prescribes an equal part each, of linseed, rye, and barley meals.

EMOLLIENT SPECIES FOR CATAPLASM.

(CAST. BORUS.; WURCEB.)

R. Wheat Bran, } of each,
Bruised Linseed, } an equal part.

Mix. (CAST. BORUS.)

WURCEB. prescribes one part of linseed, one of rye meal, and two parts of wheat bran.

MUCILAGE OF LINSEED. (GALL.; SARD.; SWEDIAUR)

R. Linseed, two drams.
Warm Water, one ounce and a half.

Digest in warm ashes for twenty-four hours, occasionally stirring, and then express. (GALL.)

SWEDIAUR prescribes an equal part of each.—SARD.—one part of seeds to be macerated for twelve hours in ten parts of water, then to boil, and express.

EMOLLIENT CATAPLASM. (GALL.; FORM. HÔP. FR.; LIPP.; DISP. CHA. PARIS; RATIER; SWEDIAUR)

R. Linseed Meal, any quantity.
Boiling Water, or Decoction of Marshmallows, a sufficient quantity to make a cataplasm.

SWEDIAUR also directs an arbitrary quantity of linseed to be boiled in water for half an hour, and a sufficient quantity of crumb of bread to be added to make a cataplasm.—RATIER prescribes an equal part each, of

linseed and of barley meals, and a sufficient quantity of water.—FORM. HÔP. FR.—an equal part each, of barley, of rye, and of linseed meals, to be boiled in a sufficient quantity of water.—GALL.; DISP. CHA. PARIS—four ounces of emollient meals and a sufficient quantity of emollient decoction, to be boiled together.—LIPP.—an equal part each, of linseed meal and of boiling cow's milk.

INFUSION OF LINSEED.

Infusum Lini seu Lini usitatissimi seu Seminum Lini. (LOND.; EDINB.; AMER.; FORM. HÔP. FR.; SUEC.; AUGUSTIN; COXE; ELLIS; DISP. CHA. PARIS; NIEMANN; RATIER; SAUNDERS; SWEDIAUR)

Infusum Lini compositum. (LOND.; and DUBL.)

Inf. Lini usitatissimi, (EDINB.)

R. Bruised Linseed, one ounce.
 ——— Liquorice, half an ounce.
 Boiling Water, two pints.

Macerate for three hours, near the fire, in a covered vessel, and strain. (LOND.; DUBL.)

EDINB. prescribes only two drams of liquorice.

AMER.; SUEC.; AUGUSTIN; COXE; ELLIS; NIEMANN; SAUNDERS; and SWEDIAUR give the same formula as LOND.—RATIER; and DISP. CHA. PARIS—two drams of linseed, two drams of liquorice, and two pounds of water.—FORM. HÔP. FR.—two drams and a half of linseed, two drams of liquorice, and two pounds of water.

R. Liquorice Powder, three drams.
 Linseed, one ounce and a half.
 Cold Water, three pounds.
 Infuse, frequently stirring, and strain. (VAN MONS)

It is prescribed as emollient in affections of the chest and of the urinary passages. Syrup may be added to render it more agreeable.

INFUSION OF PURGING FLAX.

Infusum Lini cathartici. (NIEMANN)

R. Herb of Purging Flax, two drams.
 Boiling Water, two ounces.
 Infuse, and strain It is slightly purgative.—For one dose.
 VOL. II.

EMOLLIENT FOMENTATION. (RATIER)

R. Linseed, two drams.
 Leaves of Mallow, two ounces.
 Water, two pounds.
 Boil, and strain.

EMOLLIENT BATH. (CADET DE GASSI-COURT)

R. Emollient Species, four pounds.
 Linseed, half a pound.
 Water, a sufficient quantity.
 Boil, and strain.

MUCILAGINOUS INJECTION. (FORM. HÔP. FR.)

R. Linseed, six drams.
 Boil in a sufficient quantity of water to obtain two pounds of strained decoction.

DEMULCENT POTION. (SAINTE-MARIE)

R. Bruised Linseed, one dram.
 Boiling Water, six ounces.
 Infuse, strain, and add when cold,
 Syrup of Poppies, four drams.
 Orange-flower Water, two drams.
 Mix.

DEMULCENT LOHOCH. (SAINTE-MARIE)

R. Bruised Linseed, half an ounce.
 Water, one pound.
 Boil to the consistence of syrup, and add
 Honey of Roses, two ounces.
 Mix.

MUCILAGINOUS OIL.

Oleum mucilaginum seu de Mucilaginis.
 (GALL.; SARD.; SPIELMANN; VAN MONS)

R. Seeds of Fenugreek, } of each,
 Linseed, } one part.
 Olive Oil, } eight parts.

Bruise the seeds, infuse for two days in the oil, in a sand-bath; then strain, set apart, and decant. (VAN MONS)

R. Seeds of Fenugreek, } of each,
 Linseed, } one pound.
 Root of Marshmallows, }
 Hot Water, ten pounds.

Digest for twenty-four hours, occasionally stirring; then express strongly, and add
 Olive Oil, two pounds.

Boil to the consumption of the humidity, and strain without expression.

R. Fresh Root of Marshmallows, two ounces and a half.
 ————— Squill, half an ounce.

Linseed, } of each,
 Seeds of Fenugreek, } two ounces.
 Water, a sufficient quantity.

Boil, and add

Infused Oil of Dill, } of each,
 ————— Chamomile, } sixteen
 ————— Lily, } ounces.

Boil to the consumption of the humidity, and strain. (SPIELMANN)

R. Root of Marshmallows, } of each,
 Bulb of White Lily, } one ounce.
 Seeds of Fenugreek, }
 Water, }
 Olive Oil, } of each, two pounds.
 Linseed Oil, }

Macerate the three first in the water to extract the mucilage; then mix this with the oils, and boil slowly to dissipate the humidity. Strain, and add

Flowers of White Lily, } of each,
 ————— Chamomile, } one pound.

Digest for twenty-four hours in a water-bath, and express. (SARD.)

LINSEED OIL.

Oleum Lini expressum. (LOND.; DUBL.; EDINB.; AMST.; AUSTR.; BAVAR.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; BRUGNATELLI; PIDERIT; SWEDIAUR)

Oleum Lini. (LOND.; DUBL.)

O. Lini usitatissimi. (EDINB.)

R. Linseed, any quantity.

Bruise, and express the oil without heat.

This is the general process. But GALL. requires the bruised seeds to be exposed to the vapour of boiling water for eight minutes before expression.—LIPP.; and BORUS.—the press to be heated with boiling water.—BRUNS.; PALAT.; and WIRTEM.—the seeds to be heated.

This oil is considered a specific against pleuritic pains, as well internally, in the dose of from half an ounce to one ounce in broth, as externally, applied hot on flannel to the part in pain.

EMOLLIENT LAVEMENT.

Clisma involvens seu oleosum. (LIPP.; AUGUSTIN.; RATIER.; SPIELMANN.; SWEDIAUR)

R. Emollient Species, one ounce.
 Water, one pound.

Boil, strain, and add

Linseed Oil, two ounces.

Mix. (LIPP.; RATIER.; SPIELMANN.)

R. Cow's Milk, } of each, four ounces.
 Linseed Oil, }

Mix. (SPIELMANN)

SWEDIAUR prescribes two ounces of oil, and eight of decoction of linseed.

R. Herb of Mallow, two drams.
 Water, a sufficient quantity

to obtain ten ounces of strained decoction. Add

Linseed Oil, one ounce.

Mix. (AUGUSTIN)

LIRIODENDRON TULIPIFERA.—VIRGINIAN TULIP TREE.

SYNONIMES:—Tulipier de Virginie, FR.; Tulpenbaum, GERM.; Tulpboom, DUT. Tulpeiro, PORT.; Kamestræd, SWED.

AMER.; COXE.

A tree of North America. (*Polyandria Polygynia*, LINN.; *Magnolie*, JUSS.; *Fig. Cat. Carol. I. t. 48.*)

The bark is employed, that especially of the root; it is bitter and aromatic.

LITHOSPERMUM OFFICINALE, LINN.—GROMWELL.

SYNONIMES :—Gremil, Herbe aux perles, FR.; Steinsaame, GERM.; Steenklinte, DAN.; Ackenstein saad, DUT.; Stenfræ, SWED.

BRUNS.; HISP.; GALL.; WIRTEM.; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe. (*Pentandria Monogynia*, LINN.; *Borraginea*, JUSS.; *Fig. Zorn, Ic. pl. t. 341.*)

The seeds (*Semines Lithospermi* seu *Milii solis*) are employed. They are small, very hard, and covered with a brilliant pearl-grey envelope. They are destitute of odour, and have a sweetish, oleaginous, and sub-astringent taste.

It used formerly to be considered lithon- triptic; it has also been recommended in dysentery, where it may sometimes act usefully as demulcent.

LOBELIA.

Two species of this genus are used :

1. *Lobelia inflata*, WILLD.—Indian Tobacco.

SYNONIME :—Lobélie renflée, FR.

AMER.; COXE.

A plant of North America. (*Pentandria Monogynia*, LINN.; *Lobeliaceæ*, JUSS.)

The leaves are oblong, sessile, and slightly serrated. They are at first insipid, but soon develope a pungent taste, and excite an abundant secretion of saliva.

From ten to twenty grains of the powder are given as emetic and sudorific.

TINCTURE OF INDIAN TOBACCO.

Tinctura Lobeliæ inflatæ. (AMER.; NIEMANN)

R. Dried Herb of Indian Tobacco,

two ounces.
one pint.

Dilute Alcohol,
Digest for ten days, and filter.

It is stimulant, and is extolled by THATCHER, in spasmodic asthma.—Dose, a tea- spoonful.

2. *Lobelia syphilitica*, LINN.—Blue Cardinal Flower.

SYNONIMES :—Cardinale bleue, FR.; Ge-

meine Lobelie, Blaue Cardinalsblume, GERM.; Pokkige Lobelia, DUT.

AMST.; BRUNS.; FULD.; HASS.; LIPP.; ROSS.; SAX.; WIRTEM.; HERBIP.; MURRAY and GMELIN; PH. SYPH.

A plant of North America. (*Fig. Zorn, Ic. pl. t. 368.*)

The root (*Radix Lobeliæ*) is employed. It is composed of slender, white fibres, of several fingers' length. It has a disagreeable and virous odour. When fresh its taste is very acrid, but less so after desiccation, be- coming then similar to that of tobacco, and very persistent.

It is stimulant, emetic, and purgative; and has also been classed among the anti- syphilitics.

DECOCTION OF BLUE CARDINAL FLOWER.

Decoctum Lobeliæ syphiliticæ. (PH. SYPH.; (NIEMANN)

R. Dried Root of Blue Cardinal Flower,
half an ounce.
Spring Water, twelve pounds.

Boil down to two-thirds, and strain.—
Dose, a glassful.

LONICERA PERICLYMENUM, LINN.—HONEY-SUCKLE.

SYNONIMES :—Chèvre-feuille des bois, FR.; Hahnenfüsslein, Deutsches Geißblatt, GERM.; Gewoone Kamperfolie, DUT.

GALL.; GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

This plant is found in almost every part

of Europe. (*Pentandria Monogynia*, LINN.; *Caprifoliaceæ*, JUSS.)

The stalks, leaves, and flowers are em- ployed.

The stalks (*Stipites Periclymeni* seu *Ca- prifolii*) are slender, round, long, flexible,

and covered with an ash-coloured epidermis. They have a sweet and slightly bitter taste.

The leaves are oval, pointed, glabrous, and of a sub-astringent taste.

The flowers are yellowish-white, sometimes red externally.

It is slightly astringent and tonic.

It is used in gargles.

SYRUP OF HONEY-SUCKLE. (GALL.)

R. Fresh Petals of Honeysuckle,
Boiling Water, four pounds.
 eight pounds.

Infuse for twelve hours in a covered vessel; express lightly; set aside the liquor for a few hours, decant, and add

White Sugar, twice the weight.

Make a syrup with the heat of a water-bath.

LUPINUS ALBUS, LINN.—WHITE LUPINE.

SYNONIMES:—Lupin blanc, FR.; Wolfsbohne, GERM.; Kzjmsky aneb sskrka-wicny brach, BOHEM.; Witte tygebone, DUT.

BRUNS.; HISP.; GALL.; FERR.; SAX.; WIRTEM.; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

It is very likely that this plant is originally

from the Levant, but it is naturalized in the middle parts of Europe. (*Diadelphia Hexandria*, LINN.; *Leguminosæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 321.*)

The seed (*Semen Lupini*) is employed. It is large, discoid, flattish, and white. Its kernel is yellow, bitter, farinaceous, and disagreeable. It is inodorous.

It formerly entered into cataplasms.

LYCOPERDON BOVISTA, LINN.—PUFF BALL.

SYNONIMES:—Vesse-loup des bouviers, FR.; Bovist, Wolfsrauch, Bovistkugelschamm, GERM.; Ulvefus, DAN.; Cuesco de lobo, SPAN.; Bovist, DUT.; Vesica, ITAL.; Pruchawka, POL.; Buiffa de lobe, PORT.; Kineringsfis, SWED.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; GALL.; FERR.; LIPP.; ROSS.; SAX.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A common mushroom (*Bovista seu Crepitis Lupi seu Fungus Chirurgorum*) of Europe. (*Fig. Schæff. Fung. Bav. t. 3. t. 294.*)

It is round; the flesh being at first white, then becoming greenish-yellow, and lastly brownish-grey.

It is employed to arrest hæmorrhage.

LYCOPodium CLAVATUM, LINN.—CLUB MOSS.

SYNONIMES:—Lycopode commun, FR.; Stenpulvermoos, Bärlapp, GERM.; Mechoeve seweno, BOHEM.; Joramos, Ulvefod, Lusegræs, DAN.; Licopodio, SPAN.; ITAL.; and PORT.; Geknodstaardmos, DUT.; Prosek Koltunowy, POL.; Gulnicht, SWED.

DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; GENEV.; HAMB.; HASS.; LIPP.; OLDEN.; BORUS.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A very common plant of Europe. (*Cryp-*

togamiæ, LINN.; *Musci*, JUSS.; *Fig. Zorn, Ic. pl. t. 54.*)

The herb and seeds are employed.

The herb (*Herba Musci clavati seu terrestris*) consists of a stem, much branched, and of sparse, bent leaves without nervures, and terminated by a bristle. It is inodorous and insipid.

The seeds, *Lycopodium*, (*Farina seu Pulvis seu Semines seu Sulphur Lycopodii seu Sulphur vegetabile*; Bärlappsaamen, Hexenmehl, Trutenmehl, Alpmehl, Pillenmehl, Wurmmehl, Klopfpulver, Blitzpulver, Moospulver, Strenpulver, Bärlappenzpulver, GERM.) form an exceedingly fine powder of a pale-yellow colour, light, rather unctuous

to the touch, and adhering to the fingers, and destitute of taste and smell; these minute seeds are round, flattish, and semi-transparent; they are inflammable.

This substance is used externally as a desiccative, and internally as a diuretic.—Dose, ten grains, morning and evening, in the stranguy of children.

DIURETIC MIXTURE.

Mixtura Lycopodii. (AUGUSTIN; NIEMANN)

R. Lycopodium, two drams.
Pure Water, two ounces.
Syrup of Marshmallows,
one ounce and a half.

Mix.—Dose, half a spoonful every two hours. (NIEMANN)

R. Sub-carbonate of Magnesia, two drams.
Concentrated Vinegar,
a sufficient quantity
to saturate it. Add

Mint Water, } of each,
Syrup of Opium, } two ounces.
Lycopodium, a sufficient quantity.

Mix.—Dose, two spoonfuls every two hours. (AUGUSTIN)

R. Lycopodium, half an ounce.
Syrup of Poppies, a sufficient quantity
to make a lohoch.—Dose, a spoonful every two hours.

LYSIMACHIA.—LOOSE STRIFE.

Two species of this genus of plants are employed in medicine:

1. *Lysimachia Nummularia*, LINN.—Money-wort.

SYNONIMES :—Nummulaire, FR.; Pfen-nigkraut, GERM.; Pengekrud, DAN.; Numularia, ITAL.; PORT.; and SPAN.; Penningkruid, DUT.; Pieniecznik, POL.; Penninge aert, SWED.

BRUNS.; GALL.; GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of Europe. (*Pentandria Monogynia*, LINN.; *Primulaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 20*)

The herb (*Herba Nummulariæ seu Centummorbiæ*) is employed. It consists of a simple, slightly quadrangular stem, furnished with opposite, oval leaves, raised upon very short petioles. It is without odour, and its taste is rather styptic.

It used formerly to be prescribed in diarrhoea and hæmorrhages.

2. *Lysimachia vulgaris*, LINN.—Common Loose strife.

SYNONIMES :—Lysimaque commune, Chasse-bosse, FR.; Gelbe Werderich, GERM.; Bastard dueurt, DAN.; Lis de tintureros, SPAN.; Gemeine weiderik, SWED.

GALL.; WIRTEM.; SPIELMANN; ZARDA.

A plant common throughout Europe. (*Fig. (Ed. Fl. Dan. t. 689)*)

The flowering herb (*Herba et Flores Lysimachiae luteæ*) is employed. It consists of a branching stem, of lanceolate, opposite, ternate or quaternate, sessile leaves, and of golden-yellow flowers, disposed in a terminal panicle. It is almost destitute of odour, but its taste is bitterish and styptic.

It is slightly astringent.

LYTHRUM SALICARIA, LINN.—WILLOW HERB.

SYNONIMES :—Salicaire, FR.; Brauner Weidrich, Ackerweiderich, Weiderich-blutkraut, GERM.; Wrbinda brunatna, BOHEM.; Parciike, DUT.; Salicaria, ITAL.

DUBL.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; FULD.; HAMB.; LIPP.; SAX.; HERBIP.; WIR-

TEM.; BERGIUS; COXE; MURRAY and GMELIN; SPIELMANN; ZARDA.

An European plant. (*Dodecandria Monogynia*, LINN.; *Salicariæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 113*)

The root and herb are employed.

The root (*Radix Salicariæ seu Lysima-*

II. *Manufactured according to various formulæ.*

LOND.; DUBL.; EDINB.; AMST.; ANT.;
AUSTR.; BATAV.; BAVAR.; BELG.
BRUNS.; DAN.; HISP.; GALL.; FENN.;
FERR.; FULD.; GENEV.; HANN.; HASS.;
LIPP.; OLDEN.; LUSIT.; PALAT.; POL.;
BORUS.; ROSS.; SAX.; SARD.; SUEC.;
WIRTEM.; HERBIP.; BRUGNATELLI;
COXE.; PIDERIT.; SPIELMANN.; SWE-
DIAUR.; VAN MONS.

Magnesiæ Subcarbonas. (LOND.)

R. Sulphate of Magnesia, one pound.
Subcarbonate of Potass, nine ounces.
Water, three gallons.

Dissolve the subcarbonate in three pints of water, and the sulphate in five pints, and filter the solutions; then add the remainder of the water to the solution of the sulphate and boil, adding while boiling, the solution of the subcarbonate, constantly stirring, and strain through linen. Wash the powder frequently with boiling water, and dry it upon bibulous paper, at the temperature of 200 degrees.

Carbonas Magnesiae. (EDINB.)

R. Sulphate of Magnesia, four parts.
Subcarbonate of Potass, three parts.
Boiling Water, a sufficient quantity.

Dissolve the salts separately in twice their weight of water, and strain so as to free the solution from impurities; then mix, and immediately add eight times their weight of boiling water: boil for a short time, stirring, and then having let it remain till the heat is diminished a little, strain through linen, on which the carbonate of magnesia will remain. Having washed it well with pure water, dry it slowly.

Magnesiæ Carbonas. (DUBL.)

R. Sulphate of Magnesia,
twenty-five parts.

Carbonate of Potass, fourteen parts.
Boiling Water, forty parts.

Dissolve the salts separately, each in two hundred parts of water, and mix together the defecated liquors; then boil the mixture for a short time, and strain it while hot through linen. Wash the powder frequently with boiling water, so as to remove the sulphate of potass, and finally dry the carbonate of magnesia.

R. Sulphate of Magnesia,	two pounds.
Subcarbonate of Potass,	one pound.
Boiling Water,	eight pounds.
Spring Water,	four pounds.

Dissolve, separately, the sulphate in the boiling, and the subcarbonate in the spring, water, filter the solutions, and add the latter to the former till it ceases to render it turbid : wash the precipitate with spring water, and when this is no longer rendered turbid by the addition of acetate of lead, dry the precipitated powder.

With some insignificant variations, this process is universally followed; but PALAT.; SARD.; and WIRTEM. also give the following:

R. Mother Water of Native Nitre,
[nitriers] any quantity.

Having evaporated it to dryness, increase the heat till the mass enters into fusion, and then becomes calcined; throw the pulverulent mass into boiling water, and wash and dry the precipitate which forms.

The subcarbonate of magnesia is purgative in a large dose, but it is seldom prescribed except as an antacid. It has also been extolled as a preventive to the formation of calculi by uric acid.—Dose, from three to six drams as laxative, and from six to twenty grains as absorbent.

PURGATIVE POWDER. (FERR.; AUGUSTIN; RATIER)

R. Subcarbonate of Magnesia, two scruples.
Supertartrate of Potass, one scruple.

Mix. (RATIER)

FERR.; and AUGUSTIN prescribe one ounce of magnesia, and one dram of rhubarb.—Dose, from a scruple to a dram.

DIGESTIVE POWDER. (VAN MONS)

R. Subcarbonate of Mag- } of each,
nesia, } sixteen parts.
Tartaric Acid, }
Cinnamon, } one part.

Mix.

PAREGORIC POWDER. (SAUNDERS)

- R. Subcarbonate of Magnesia, two drams.
 Nitrate of Potass, one dram.
 Cascarilla, one scruple.
 Mass of Dog-rose Pill, six grains.

Mix.

ABSORBENT POWDER.

Pulvis absorbens seu antacida seu galactopocus seu nutriticum seu infantum seu infantum antacidus seu pro infantibus seu puerorum citrinus seu epilepticus Marchionum seu ad tormina infantum antacidus. (GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; ROSS.; WIRTEM.; AUGUSTIN.; ELLIS.; NIEMANN.; PIDERIT.; PIERQUIN.; SAINTE-MARIE.; SPIELMANN.; SWEDIAUR.)

- R. Subcarbonate of Magnesia, } of each,
 White Sugar, } one ounce.

Triturate for a long time in a glass mortar. (GALL.)

- R. Subcarbonate of Magnesia, eight grains.
 Cinnamon, one grain.

Mix. (ROSS.)

HAMB. prescribes six drams of magnesia and two of orange peel.—HISP.—one dram and a half of magnesia and one dram of calumba; to be divided into six parcels.

- R. Subcarbonate of Magnesia, six parts.
 Oleosaccharum of Peppermint, one part.

Mix. (VAN MONS)

- R. Subcarbonate of Magnesia, one ounce.
 Cinnamon, one dram.
 Sugar, half an ounce.

Mix. (ROSS.)

FULD. prescribes one ounce and a half of subcarbonate, one dram of cinnamon, and one ounce of sugar.—LIPP.; and WIRTEM.—two ounces of subcarbonate, one dram of cinnamon, and half an ounce of liquorice.

- R. Subcarbonate of Magnesia, one ounce.
 Fennel, one dram.
 White Sugar, half an ounce.

Mix. (HASS.; PIDERIT)

- R. Subcarbonate of Magnesia, one ounce.
 Orange Peel, } of each, two drams.
 Fennel, }

Mix. (FERR.; FULD.; LIPP.)

HISP. prescribes half an ounce of subcarbonate, one dram each, of fennel and of cinnamon.—AUGUSTIN prescribes one ounce of subcarbonate, and half an ounce each, of fennel and of orange peel.

- R. Subcarbonate of Magnesia, three ounces.

- Nutmeg, half a dram.
 Cinnamon, one dram and a half.

Mix. (SWEDIAUR)

- R. Subcarbonate of Mag- } of each,
 nesia, } two ounces.
 Sugar of Milk, }
 Cascarilla, two drams.
 Oleosaccharum of Fennel, six drams.

Mix. (SAINTE-MARIE)

- R. Subcarbonate of Magnesia, six drams.
 Root of Florentine Iris, } of each,
 Anise, } two drams.
 White Sugar, }
 Saffron, one dram.

Mix. (HANN.; NIEMANN.; SPIELMANN)

- R. Subcarbonate of Magnesia, half an ounce.
 Cumin, three drams.
 Florentine Iris, two drams.
 Saffron, half a dram.

Mix. (PIERQUIN.; SPIELMANN)

- R. Subcarbonate of Magnesia, two ounces.
 Rhubarb, half an ounce.
 Miseltoe, } of each, six drams.
 Liquorice, }

Mix. (LIPP.)

OLDEN. prescribes one ounce of subcarbonate of magnesia, two drams of rhubarb, and half a dram of analeptic powder.

- R. Subcarbonate of Magnesia, half an ounce.
 Soda, } of each,
 Ginger, } one scruple.

Mix. (HISP.)

- R. Subcarbonate of Magnesia, one ounce.
 Gentian, } of each, two drams.
 Fennel, }

Mix. (AUGUSTIN)

The dose is generally a tea spoonful.—Some pharmacopœias, amongst which are GALL.; and HISP.; substitute for the subcarbonate the pure magnesia, which, indeed, has the advantage of not inconveniencing the patient by the disengagement of carbonic gas in the stomach.

DEOBSTRUENT POWDER.

Pulvis ecphracticus Sellii. (AUGUSTIN)

- R. Subcarbonate of Magnesia, }
 Supertartrate of Potass, } of each,
 Sublimed Sulphur, } one ounce.
 Rhubarb, }
 Chamomile Flowers, }
 Oleosaccharum of Fennel, }

Mix.—Dose, one tea spoonful several times a day, in obstructions and debility of the abdominal viscera.

STOMACHIC BOLUSES. (FORM. Hôp. Fr.)

- R. Subcarbonate of Magnesia, five parts.
Saffron, three parts.
Cinnamon, two parts.
Simple Syrup, a sufficient quantity.
Mix.

ABSORBENT TROCHISCS.

Tabellæ absorbentes seu Magnesiae, Trochisci antacidi seu Magnesiae seu Carbonatis Magnesiae. (EDINB.; AMER.; ANT.; GALL.; LUSIT.; BRUGNATELLI; COXE; PIERQUIN; SWEDIAUR; VAN MONS)

Trochisci Carbonatis Magnesiae. (EDINB.)

- R. Carbonate of Magnesia, six ounces.
Refined Sugar, three ounces.
Nutmeg, one scruple.

Pulverize, and form trochiscs with mucilage of tragacanth.

PIERQUIN gives the same formula.—ANT.; and GALL. prescribes one part of subcarbonate, four parts of sugar, and a sufficient quantity of mucilage of tragacanth prepared with orange-flower water.—BRUGNATELLI—three ounces of subcarbonate, one ounce and a half of sugar, one scruple of ginger, and mucilage of gum arabic.—SWEDIAUR—four ounces of magnesia, two of sugar, one scruple of ginger, and mucilage of gum arabic.—LUSIT.—six ounces of magnesia, three of sugar, half a dram of pichurim bean, and mucilage of tragacanth.—AMER.; and COXE—four ounces of subcarbonate, two of sugar, one scruple of ginger, and a sufficient quantity of simple syrup.

- R. Subcarbonate of Sugar, one part.
White Sugar, eight parts.

Boil seven-eighths of the sugar *à la plume*, and incorporate the rest triturated with the subcarbonate. (VAN MONS)

CHEVALLIER has proposed the following pastilles:

- R. Chocolate, two ounces.
Sugar, five ounces.
Pure Magnesia, one ounce.

Make them into a paste with mucilage of tragacanth flavoured at pleasure, and divide the mass into pastilles of eight, sixteen, and twenty-four grains.

ANTACID ELECTUARY.

Electuarium anodynum pro infantibus seu antacidum seu mithridaticum. (BRUNS.; LIPP.; CADET DE GASSICOURT; SPIELMANN; SWEDIAUR)

- R. Subcarbonate of Magnesia, one ounce.
Anise, half an ounce.
Compound Syrup of Cichory, three ounces.

Mix. (BRUNS.)

CADET DE GASSICOURT; and SPIELMANN prescribe six drams of subcarbonate, half an ounce of anise, one dram of saffron, and a sufficient quantity of the syrup.

- R. Subcarbonate of Magnesia, six drams.
Fennel, } of each, three drams:
Charcoal, }
Saffron, one dram.
Simple Syrup, eight ounces.
Petals of Red Poppies, a sufficient quantity

to colour the whole reddish-brown. (LIPP.)

- R. Subcarbonate of Magnesia, one ounce and a half.
Herb of Mentha Crispa, one ounce.
Saffron, one dram.
Rhubarb, two drams.
Honey, six ounces.
Charcoal, a sufficient quantity

to colour it blackish-green. (LIPP.)

- R. Conserve of Mentha Crispa, one ounce.
Aromatic Powder, one dram.
Subcarbonate of Magnesia, two drams.
Syrup of Mentha Crispa, a sufficient quantity.

Mix. (SWEDIAUR)

Dose, one or two drams, or more.

WATER OF MAGNESIA.

Aqua Magnesiae. (AMER.; COXE; HENRI; NIEMANN)

- R. Water, eight pounds.
Subcarbonate of Magnesia, three drams.

By means of a forcing pump, pass into the mixture about ten times its volume of carbonic acid gas. (AMER.; NIEMANN)

- R. Hydrated Carbonate of Magnesia, four parts.
Water, charged with four times its volume of carbonic acid gas, five hundred parts.

Mix. (HENRI)

FACTITIOUS PYRMONT WATER. (GALL.)

- R. Water, charged with five measures of carbonic acid gas, twenty ounces and a half.
Muriate of Soda, two grains.
Sulphate of Magnesia, eight grains.
Subcarbonate of Magnesia, twelve grains.
Iron, one grain.

Mix.

2 E

ANTACID POTION.

Mixtura antacida, Emulsio Amygdalarum composita. (HAMB.; HANN.)

- R. Magnesia, half a pound.
 Fennel Water, one ounce and a half.
 Aqueous Tincture of Rhu-
 barb, } of each,
 White Sugar, } two drams.

Mix. (HANN.)

- R. Spanish Soap, one scruple.
 Magnesia, one dram.
 Distilled Water, three ounces.
 White Sugar, two drams.

Mix. (HAMB.)

- R. Sweet Almonds, half an ounce.
 Seeds of Henbane, one dram.
 Bird Cherry Water, eight ounces.

Make an emulsion, and strain. Add

- White Sugar, six drams.
 Subcarbonate of Magnesia, one dram.

Mix. (HANN.)

PURGATIVE POTION. (AMER.; COXE;
 ELLIS; PIERQUIN; SAINTE-MARIE)

- R. Subcarbonate of Magnesia, two drams.
 Decoction of Chick-peas, four ounces.
 Syrup of Maidenhair, two ounces.

Mix. (PIERQUIN)

- R. Rhubarb, one dram.
 Water, twelve ounces.

Boil down to one half, and add when strained

- Subcarbonate of Magnesia, one dram.
 Spirit of Anise, three drops.

Mix. (SAINTE-MARIE)

- R. Magnesia, half a dram.
 Rhubarb, six grains.
 Sugar, one dram.
 Essence of Peppermint, six drops.
 Water, one ounce and a half.

Mix.—Dose, a tea spoonful every two hours; for infants. (ELLIS)

- R. Subcarbonate of Magnesia, half a dram.
 Tincture of Assafoetida, forty drops.
 ——— Opium, twenty drops.
 Sugar, one dram.
 Water, one ounce.

Mix.—Dose, twenty-five drops, to an infant of from two to four months old.—This is from DEWEES.

- R. Subcarbonate of Magnesia, one dram.
 Solution of Subcarbonate of
 Ammonia, one ounce.
 Cinnamon Water, three ounces.
 Water, five ounces and a half.

Mix. (AMER.; COXE)

LITHONTRIPTIC POTION. (ELLIS)

- R. Subcarbonate of Magnesia, one dram.
 Infusion of Gentian, six ounces.

Mix.—This is recommended by BRANDE in cases of predominancy of uric acid.—Dose, a glassful thrice a day.

PROTO-PHOSPHATE OF MAGNESIA.

Phosphas Magnesiae. (NIEMANN)

- R. Pure Magnesia, any quantity.
 Phosphuric Acid, a sufficient quantity to saturate; filter, and evaporate the solution to dryness.

This salt has been extolled at the commencement of rachitis.—Dose, from ten grains to half a dram. In the quantity of a dram it purges.

PROTO-SULPHATE OF MAGNESIUM.

Magnesia sulphurica seu vitriolata, Sal amarum seu Anglicanum seu Anglicum seu Anglicanum catharticum seu catharticum amarum seu Ebsharnense seu Epsomense seu Scydtschutzensen seu Sardinizensesen Sedlitzense, Sulphas Magnesiae, Terra amara sulphurica seu vitriolata, Vitriolicum Magnesiae.—Sulphate of Magnesia, Epsom Salt.

SYNONIMES:—Sulfate de protoxide de magnesium, Sulfate de magnesie, Sel d'Epsom, Sel de canal, Vitriol magnesien, Fr.; Bittersalz, Schwefelsaure Talkerde, GERM.; Engelsk laxesalt, DAN.; Sal-amargo, Sal de la biguera, SPAN.; Sale amaro, ITAL.; Sol gorzka, POL.; Sal cathartico amaro, PORT.; Bitter salt, Eller engelski salt, SWED.

I. *That which is met with in Commerce.*

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; COXE.; GUIBOURT.; PARIS.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS.

It is in four-sided prisms, terminated by four-sided pyramids, or by a dihedral apex, white, efflorescent, and capable of aqueous solution. Its taste is saline and very bitter.

All the pharmacopoeias direct it to be purified, by dissolving it in boiling water, evaporating the solution till a pellicle forms, and setting it aside to re-crystallize.

II. *Prepared pharmaceutically.*

GALL.

- R. Subcarbonate of Magnesia, any quantity.

Sulphuric Acid (15 degrees),
a sufficient quantity
to saturate the magnesia; boil slightly,
strain, evaporate till the liquor marks 25
degrees, and set it apart in a cool place to
crystallize.

It is purgative, in the dose of from three
drams to one ounce and a half dissolved in
a pound of water; or, in the quantity of
one or two ounces or more in lavements.

It is incompatible with the hydrochlo-
rates of ammonia and of lime, the subcar-
bonates of soda and of potass, the acetate of
lead, and the nitrate of silver.

COMPOUND SALINE POWDER.

*Pulvis salinus compositus seu Cheltenham-
ensis.* (EDINB.; COXE; NIEMANN)

R. Muriate of Soda, } of each,
Sulphate of Magnesia, } four
parts.
Potass, three parts.

Dry the salts with a gentle heat,
previously reduced to a fine powder;
then triturate them together, and
preserve in a well-stopped bottle.
(EDINB.; COXE)

NIEMANN prescribes one hundred and
twenty grains of muriate of soda, sixty-six
of sulphate of magnesia, ten of sulphate of
potass, and half a grain of sulphate of iron.

Dose, one tea spoonful, in a pound of
water.

DIGESTIVE POWDER. (BRUNS.)

R. Sulphate of Magnesia, two ounces.
Nitrate of Potass, } of each,
Supertartrate of Potass, } one ounce.
Powdered Root of Arum, }
Muriate of Ammonia, six drams.

Make a powder.—For the same purposes
as the preceding.—It resembles BIRKMAN'S
stomachic powder.

PURGATIVE POWDER. (ELLIS)

R. Sulphate of Magnesia, } of each,
Cinchona, } six drams.

Make four parcels.—Dose, one every two
hours.

CATHARTIC POTION.

*Potio cathartica seu laxans, Liquor salinus
laxativus, Haustus catharticus salinus,
Mixtura e Magnesia sulphurica, Aqua
mineralis cathartica.* (FERR.; WIR-
TEM.; HERBIP.; AUGUSTIN; BRERA;
ELLIS; SAUNDERS; SWEDIAUR)

R. Sulphate of Magnesia, } of each,
Manna, } one ounce.
Warm Water, four ounces.

Dissolve, and strain. (HERBIP.)

R. Sulphate of Magnesia, one ounce.
Boiling Water, one pound.

Add to the solution

Subcarbonate of Magnesia, ten grains.

Mix.

SWEDIAUR gives in two places this for-
mula under the names of *Potio cathartica*
and *Aqua mineralis cathartica*. It may be
used instead of Seidlitz water.

R. Sulphate of Magnesia, } of each,
Aqueous Tincture of } one ounce.
Rhubarb, }
Bird Cherry Water, two ounces.

Mix.—To be taken at twice, or at four
times, morning and evening. (AUGUSTIN)

R. Sulphate of Magnesia, one ounce.
Tartarized Antimony, one grain.
Lemon Juice, one ounce.
Water, three ounces.

Mix.—Dose, one spoonful every two
hours. (ELLIS)

R. Leaves of Senna, half an ounce.
Water, a sufficient quantity
to obtain eight ounces of infusion. Add
when strained

Sulphate of Magnesia, one ounce.

Mix.—To be taken at twice. (BRERA)

R. Senna, half an ounce.
Water, a sufficient quantity
to obtain four ounces of infusion. Add
when strained

Aqueous Tincture of Rhubarb,

six ounces.

Sulphate of Magnesia, one ounce.

Mix.—To be taken at twice or thrice.
(BRERA)

R. Senna, six drams.
Manna, } of each,
Sulphate of Magnesia, } one ounce.
Lesser Cardamoms, two drams.
Boiling Water, one pound.

Infuse for a quarter of an hour, and strain.
—Dose, a cupful every hour. (ELLIS)

R. Senna, three drams.
Sulphate of Magnesia, six drams.
Boiling Water, six ounces.

Infuse for twelve hours, and strain.
(FERR.)

SALINE POTION. (HAMB.)

R. Tartarized Antimony, one grain.
Sulphate of Magnesia, one ounce.
Spring Water, eight ounces.
Glycyrrhized Honey, one ounce.

Mix.—To be taken at four or five doses.
(HAMB.)

CADET DE GASSICOURT gives the following under the title of *Eau de Trevez*.

R. Sulphate of Magnesia, one ounce.
Tartarized Antimony, half a grain.
Water, two pounds.

Mix.—Dose, a glassful every hour, till it purges.

ANTACID SALINE MIXTURE. (NIEMANN)

R. Sulphate of Magnesia, two drams.
Subcarbonate of Magnesia, one scruple.

Syrup of Marshmallows, one ounce and a half
Chamomile Water, two ounces.

Mix.—Dose, a spoonful every two hours; for children.

OLEOSO-SALINE MIXTURE. (HAMB.)

R. Sulphate of Magnesia, one ounce.
Water, half a pound.

Dissolve, and add

Castor Oil, triturated with Yolk of Egg, one ounce.

Mix.

ARTIFICIAL SEIDLITZ WATER. (GALL.)

R. Water, containing thrice its volume of Carbonic Acid Gas, twenty ounces and a half.
Sulphate of Magnesia, two drams.
Hydrochlorate of Magnesia, eighteen grains.

Dissolve.—It may be made stronger and more like the natural water, by dissolving four drams of sulphate and thirty-six grains of hydrochlorate in the same quantity of acidulated water.

PURGATIVE LAVEMENT.

Enema purgans seu catharticum. (DUBL.; WURCEB.; AUGUSTIN; ELLIS)

R. Manna, } of each,
Compound Decoction } ten
of Chamomile, } ounces.
Olive Oil, one ounce.
Sulphate of Magnesia, half an ounce.

Dissolve, and mix. (DUBL.; COXE)

WURCEB. prescribes a solution of six drams of sulphate of magnesia, to be dissolved in ten ounces of warm water.—ELLIS—a solution of two ounces of sulphate in a pint of infusion of senna; or, two ounces of the salt in a pint of warm decoction of starch, to which an ounce of olive oil is to be added.

R. Chamomile Flowers, two ounces.
Linseed, one ounce.
Water, a sufficient quantity,
to obtain six ounces of decoction. Add when strained

Sulphate of Magnesia, half an ounce.
Linseed Oil, one ounce and a half.

Mix. (AUGUSTIN)

R. Gruel, eight ounces.
Sulphate of Magnesia, two drams.
Soap, half an ounce.

Mix. (AUGUSTIN)

R. Sulphate of Magnesia, six drams.
Carbonate of Magnesia, six grains.
Water, three ounces.
Tincture of Cardamoms, one dram.

Mix. (SWEDIAUR)

R. Sulphate of Magnesia, six drams.
Water, three ounces.
Tincture of Senna, six drams.

Mix. (SAUNDERS)

R. Sulphate of Magnesia, three ounces.
Potass, } of each,
Nitrate of Potass, } half an ounce.
Water, fifteen ounces.

Dissolve and filter. (WIRTEM.)

CARBONIC SOLUTION OF SULPHATE OF MAGNESIA.

Aqua amara seu Sulphatis Magnesiae carbonicae. (FERR.; FULD.; SUEC.; RATIER; VAN MONS)

R. Sulphate of Magnesia, seven parts.
Water, thirty-six parts.

Filter, and saturate the solution with carbonic acid gas. (SUEC.)

R. Sulphate of Magnesia, three parts.
Water, twenty-two parts.

Dissolve, and add

Solution of Carbonic Acid, twenty-two parts.

Mix. (FENN.; VAN MONS)

R. Water, three pounds and a half.
Sulphuric Acid diluted with six parts of Water, ten drams.
Subcarbonate of Magnesia, three drams.

Dissolve in a bottle filled only to two thirds. (RATIER)

R. Sulphate of Magnesia, one ounce.
Water, four ounces.

Dissolve and add.

Solution of Subcarbonate } of each,
of Ammonia } one
Diluted Sulphuric Acid, } scruple.

Mix.—For one dose. (FERR.)—It is said that this dose repeated thrice, will cure, without cinchona, intermittent fevers accompanied with disorder of the digestive organs.

APERIENT WHEY. (SAINTE-MARIE)

- R. Clarified Whey, one pound.
 Root and Herb of Dandelion, }
 Herb of Fumitory, } of each,
 Leaves of Water-cress, } one handful.
 Chervil, }

Boil for about six minutes, express strongly, and add

Sulphate of Magnesia, one or two drams.
 Honey, six drams.

Mix. It is recommended by VAN SWIETEN, in engorgements of the abdominal viscera. To be taken every morning fasting, in four doses at intervals of an hour. At the end of a month the waters of Spa should be commenced with.

PROTO-HYDROCHLORATE OF MAGNESIUM. (VAN MONS)

- R. Sulphate of Magnesia, ten parts.
 Hydrochlorate of Potass, seven parts.

Mix the pulverized salts in an earthenware pan, pour upon them one half of their weight of warm water, decant when cold, and evaporate, separating the crystals which

form, and pour the highly concentrated liquor upon a dry slab.

SILICATE OF MAGNESIA.

1. *Creta gallica*; French chalk.

SYNONIMES:—Craie de Briançon, FR.; Franzäische Kreide, GERM.

BRUNS.

In solid masses, formed of small pearly or satin-like scales.

2. *Talcum Venetæ, Stella Terræ*; Talc.

SYNONIMES:—Talc, FR.; Gemeiner Talk, Talch, Talch, GERM.

BRUNS.; WIRTEM.; GUIBOURT; SPIELMANN.

In flexuous, brilliant laminæ of a greenish-white colour, unctuous to the touch, very tender, flexible, non-elastic and easily separated. [*faciles à rayer.*]

ARCANUM COSMETICUM. (NIEMANN)

- R. Talc, one dram.
 Camphorated Oil, two drams.

Digest in a water bath, and mix.

MAGNOLIA GLAUCA, LINN.—SWEET-SCENTED MAGNOLIA.

SYNONIME:—Magnolier glauque, FR.; AMER.; COXE.

A small shrub of North America. (*Polyandria Polygynia*, LINN.; *Magnoliaceæ*, JUSS. Dill. *Eth.* tab. 168. fig. 205.)

The bark has an aromatic odour resembling that of sassafras.

It is tonic, stimulant, and diaphoretic—The tincture of it is employed in chronic rheumatism, it is also prescribed in intermittent fevers.

MALTUM.—MALT.

SYNONIMES:—*Maltum*.—Malt, Maltum, FR.; Gerstenmalz, GERM.; Malt, DAN.; and SWED.; Grano tosto, ITAL.; Stod ieczmienny, POL.

BAVAR.; DAN.; HANN.; HASS.; POL. CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; BERGIUS; MURRAY and GME-LIN; SPIELMANN.

This is prepared from barley, by moistening it, and keeping it in a warm place till it begins to germinate; it is then dried with a regulated heat, and the germs separated.

Malt contains more starch, sugar, and gum than barley, and is preferable to it as

an emollient. It forms the basis of beer. It has been ranked among the anti-scorbutics, but probably without much reason.

INFUSION OF MALT.

Infusum Maltis seu *Bynes*. (LIPP.; SUEC.; ELLIS; SPIELMANN)

- R. Malt, grossly powdered, one pound.
 Boiling Water, twelve pounds.

Infuse for several hours, and strain. (SUEC.; SPIELMANN)

LIPP. prescribes four ounces of malt, and one pound of boiling water; to be infused

for four hours.—ELLIS—one pound of essence of malt and three pounds of boiling water.

When evaporated to the consistence of honey, this infusion takes the name of Essence of Malt, *Essentia Malti*. It is taken to sea in long voyages, to make beer with.

DECOCTION OF MALT.

Decoctum seu Apozema Malti, Mustum Hordei. (FORM. HÔP. FR.; CAST. BORUS.; WURCEB.; AUGUSTIN.; NIEMANN.; SAUNDERS.; SWEDIAUR)

R. Malt, coarsely powdered, four ounces.
Water, four pounds.

Boil down to three pounds. (CAST. BORUS.; AUGUSTIN)

R. Malt, six ounces.
Spring Water, four pounds.

Boil for a quarter of an hour; add

Seeds of Fennel, half a dram.

Digest at a moderate heat in a covered vessel, and strain. (SAUNDERS)

R. Ground Malt, four ounces.
Boil it with

Water, a sufficient quantity
to obtain two pounds of strained decoction,
adding towards the end,

Liquorice Root, one dram.
Decant. (FORM. HÔP. FR.)

NIEMANN prescribes six ounces of malt and six pounds of water, to be reduced to four pounds. Four scruples of fennel, and three drams of liquorice may be added.

R. Malt, six ounces.
Water, five pounds.

Boil down to four pounds, and add when strained,

Syrup of Citric Acid, four ounces.

Mix. (SWEDIAUR)

R. Malt, two ounces.
Water, four pounds.

Boil, and add when strained, if required,

Oxymel, three or four ounces.

Raisins may also be boiled with the malt. (HERBIP.)

Dose, from two to four pounds, daily.

ANTI-SCORBUTIC GARGLE. (SAINTE-MARIE)

R. Malt, one ounce.
Water Cresses, one handful.
Water, four pounds.

Boil down to two pounds, and add

Alum, two drams.
Muriate of Ammonia, one dram.

Mix.

MALUS COMMUNIS, CAND.—APPLE TREE.

SYNONIMES:—Pommier, FR.; Apfelbaum, GERM.; Missenski gablko, BOHEM.; Æbletræ, DAN.; Manzano, SPAN.; Appelboom, DUT.; Malo, ITAL.; Jablon, POL.; Maceiro, PORT.; Appeltræd, SWED.

AMST.; BAVAR.; BRUNS.; HISP.; GALL.; FULD.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; BERGIUS.; BRUGNATELI.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

This tree, is spread over every part of Europe. (*Icosandria, Monogynia*, LINN.; *Rosaceæ*, JUSS.; *Fig.* Blackwood, *Herb. t.* 178)

It has from cultivation produced numerous varieties yielding very different fruits, but all are called apples, *Poma*. Of these fruits the Rennet, *Poma Renetia*, is employed.

JUICE OF APPLES.

Succus Pomorum. (HISP.; GALL.; FULD.; SARD.)

R. Rennet Apples, any quantity.

Rasp them, remove the pips, express the juice from between two layers of rye straw chopped and washed with warm water; set it aside, strain, and preserve it in bottles covered with a stratum of oil.

SYRUP OF APPLES.

Syrupus Pomorum redolentium. (HISP.; FERR.)

R. Juice of Apples, one pound.
White Sugar, two pounds.

Dissolve in a water-bath. (HISP.)

FERR. directs, either to dissolve three pounds of sugar in four of juice, or to boil quarters of apples, peeled and freed from pips, in simple syrup.

DECOCTION OF APPLES.

Potus ex Pomis seu Pomorium. (AUGUSTIN; NIEMANN)

R. Rennet Apples, ten ounces.
Spring Water, two pounds and a half.

Boil for an hour in a covered vessel, and strain.

This pleasant drink is rendered still more

agreeable by the addition of a little sugar. If the indication requires or permits it, a dram of nutmeg in powder, half a dram of cinnamon, or three ounces of Rhenish wine may be added. In the latter case it should be considered only as the vehicle of some stimulant; in the former, only as a means of soliciting the patient to drink water, for its very slight stimulant properties do not deserve consideration.

MALVA.—MALLOW.

Three species of this genus of plants are used in medicine:

1. *Malva Rotundifolia*, LINN. — Round-leaved Mallow.

SYNONIMES: — Mauve à feuilles rondes, Petite Mauve, FR.; Rundblättrige Pappel, GERM.; Zagec slez, BOHEM.; Store katterst, DAN.; Malva de boja redonda, SPAN.; Rondbladige maluwe, DUT.; Malva minore, ITAL.; Szlaz ziele, POL.; Malva ordinaria, PORT.; Kattost, SWED.

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; PIDERIT.; SPIELMANN.; ZARDA.

This plant is common in every part of Europe. (*Monadelphia Polyandria*, LINN.; *Malvaceae*, JUSS.; *Fig. Zorn, Ic. pl. t. 237*)

The root, herb, flowers, and seeds are employed.

The root (*Radix Malvæ Minoris seu vulgaris*) is thin, cylindrical, fibrous, and white. It has a sweet, almost saccharine, taste.

The herb consists of a round, hairy stem, and of orbicular, heart-shaped, five-lobed leaves, of a mucilaginous and bitterish taste.

The flowers are rather small, of a very pale purple and almost white.

The seeds are small, reniform, and black.

2. *Malva Sylvestris*, LINN. — Wild Mallow; Common Mallow.

SYNONIMES: — Mauve sauvage, FR.; Wilde pappel, GERM.; Lillekatost, DAN.; Wilde maluwe, DUT.; Malvone, ITAL.

LOND.; DUBL.; EDINB.; AUSTR.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; DAN.;

MIL, COPEN.; HISP.; GALL.; FORM. HÔP. FR.; GENEV.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; BORUS.; WURCEB.; COXE; MURRAY and GMELIN.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 480*)

The root, herb, flowers, and seeds are employed.

The root (*Radix malvæ sylvestris*) is whitish, and of a sweet and viscous taste.

The herb consists of a cylindrical, slightly pubescent, branching stem, and of round leaves, heart-shaped at the base, crenelated at the borders, and notched into five or seven lobes.

The flowers are large, rose coloured, streaked with red, sometimes white.

The seeds are yellowish and smooth.

These two plants are emollient.

EMOLLIENT SPECIES. (FULD.; LIPP.; SUEC.)

R. Leaves of Mallow, twelve parts.
Chamomile Flowers, six parts.
Linseed, three parts.

Cut and bruise.

PECTORAL FLOWERS. (GALL.)

R. Flowers of Mallow, } of each,
—Amaranth, } an
—Coltsfoot, } equal part.
—Red Poppies, }

Mix.

CONSERVE OF MALLOW. (ANT.; SARD.; HERBIP.)

R. Flowers of Mallow, one part.
White Sugar, two parts.

Beat to a paste. (HERBIP.)

R. Flowers of Mallow, beat up with
a little sugar, one part.

Syrup, three parts.
Boil for an instant. (SARD.)

R. Dried Leaves of Mallow, in powder, one part.
White Sugar, five parts.
Distilled Water, two parts.
Triturate together. (ANT.)

EMOLLIENT CATAPLASM. (FERR.)

R. Manna, four ounces.
Root of Mallow, two ounces.
Decoction of Marshmallows, a sufficient quantity,
to make a soft paste by boiling.

DISTILLED WATER OF MALLOW. (SARD.)

R. Fresh Herb of Mallow, one part.
Water, two parts.
Distil two thirds.

INFUSION OF MALLOW. (FORM. HÔP. FR.; PH. SYPH.; DISP. CHA. PARIS; SPIELMANN)

R. Fresh Leaves of Mallow, three handfuls.
Boiling Water, four pounds.
Infuse, and strain. (PH. SYPH.)
R. Fresh Leaves of Mallow, two drams.
Liquorice, one dram.
Water, a sufficient quantity
to obtain one pound of strained liquor.
(FORM. HÔP. FR.; DISP. CHA. PARIS)
R. Flowers of Mallow, two drams.
Boiling Water, two pounds.
Infuse and strain. (RATIER)
It is emollient.

COMPOUND DECOCTION OF MALLOW.

Decoctum Malvæ compositum. (LOND.; NIEMANN)

R. Dried Leaves of Mallow, one ounce.
—— Chamomile Flowers, half an ounce.
Water, one pint.
Boil for a quarter of an hour, and strain.

It is used in fomentations and lavements.

PECTORAL POTION. (GALL.)

R. Pectoral Flowers, two drams.
Boiling Water, two pounds.
Infuse for a quarter of an hour; strain, and add,
Syrup of Maidenhair, one ounce.
Mix.

EMOLLIENT LAVEMENT. (HISP.; RATIER; SPIELMANN; SWEDIAUR)

R. Leaves of Mallow, half an ounce.
Boiling Water, one pound.
Boil, and strain. (RATIER)

R. Leaves of Mallow, one ounce.
Chamomile Flowers, half an ounce.
Water, one pound and a half.
Boil down to two thirds. (SWEDIAUR)

R. Leaves of Mallow, one ounce.
Chamomile Flowers, } of each,
Fennel, } half an ounce.
Water, twelve ounces.

Boil and strain. (SPIELMANN)

R. Leaves of Mallow, } of each,
—— Beet, } two ounces.
Water, two pounds.

Boil, and add,
Olive Oil, three ounces.

Mix. (RATIER)

HISP. prescribes ten ounces of decoction of mallow and two ounces each, of honey and of oil.

DEMULCENT GARGLE. (RATIER)

R. Decoction of Mallow, four ounces.
Cow's Milk, one ounce.
Mix.

ANTI-HÆMORRHOIDAL FOMENTATION. (PIERQUIN)

R. Leaves of Mallow, } of each,
—— White Mullein, } three handfuls.
Spring Water, three pounds.
Boil, and strain.

OINTMENT OF MALLOW. (VAN MONS)

R. Fresh Herb of Mallow, one pound and a half.
—— Butter, two pounds.
Boil slowly to the consumption of the humidity, and express.

3. *Malva Alcea*, LINN.—Vervain Mallow.

SYNONIMES:—Mauve alcée, Mauve sauvage, FR.; Feltrisswurz, Schlitsblättrige Malva, Alceenmalve, Sigmundswurz, Sigmarskraut, GERM.

BRUNS.; GALL.; HAMB.; HASS.; OLDEN.; WIRTEM.; GUIBOUT; SPIELMANN.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 219.*)

The root and herb are employed.

The root (*Radix Alceæ*) is thick, white, rather ligneous; it is branched, and has many fibrillæ.

The herb consists of a cylindrical stem, furnished with clustered hairs, and of petioled leaves, rough to the touch; the lower ones are divided into five round lobes, and the upper into deeper lobes, for the greater part incised and nearly pinnatifid. It has a sweet and mucilaginous taste.

It is emollient.

TINCTURE OF ALCEA.

Tinctura Alceæ. (BRUGNATELLI)

- R. Dried Petals of Alcea, one dram.
Alcohol, two ounces.

Digest in a covered vessel at a moderate temperature, or exposed to the sun for two days, shaking from time to time, and filter.

This is used as a test for discovering the presence of acids which redden it, and of alkalis which render it green.

MANGANESIUM.—MANGANESE.

The following compounds of this metal are employed in medicine:

TETROXIDE OF MANGANESE.

Magnesia nigra seu *Vitrariorum*, *Magnesium oxydatum nativum* seu *nigrum* *Manganesium ochraceum nigrum* seu *oxydatum nativum*, *Manganum oxydatum nativum*, *Molybdænum Magnesii*, *Oxodes Mangani nativa*, *Oxydum Mangani nativum*, *Oxydum Magnesii nigrum* seu *nigrum nativum*, *Peroxydum Manganesii nigrum nativum*, *Superoxydum manganicum*; *Acidum manganesticum*, BELG.—Peroxide of Manganese, Black Oxide of Manganese.

SYNONIMES:—Magnésie noire, Manganèse oxidé, H., Oxide noir de manganèse, Peroxide de manganèse, B., Savon des verriers, Savon de verre, FR.; Braunstein, GERM.; Brunsteen, DAN.; Magnesia, ITAL.; PORT.; and SPAN.; Bruinsteen, DUT.; Mangana, POL.; Manganazi, ROSS.; Brunsten, SWED.

DUBL.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSIE; COXE; GUIBOUT.

It is in dull masses, the colour of which varies from an almost pure black to a brown or violet-brown.

The medicinal uses of this oxide are but few. It has been employed, mixed with an equal part of lard, in tinea, itch, and tetters. It would be sufficient merely to cover the

affected parts with the powder. Its principal use is for the preparation of chlorine.

EMMENAGOGUE BOLUSES.

- R. Peroxide of Manganese, fourteen grains.
Extract of Savine, } of each, ten grains.
Aloes, }

Make six boluses, to be taken in the course of the day.

HYDROCHLORATE OF MANGANESE.

Murias Manganesii. (NIEMANN)

- R. Manganese, any quantity.
Diluted Hydrochloric Acid, a sufficient quantity
to dissolve the metal. Evaporate the solution.

PILLS OF HYDROCHLORATE OF MANGANESE.

Pilule Manganesii muriatici. (AUGUSTIN; NIEMANN)

- R. Hydrochlorate of Manganese, two scruples.
Gum Arabic, } of each, one scruple.
Liquorice, }

Make a pilular mass.—Dose, from twenty to forty grains, in herpetic diseases.

The aqueous solution of this salt has also been extolled as a gargle in aphthæ. That of the acetate, in the proportion of a scruple to three ounces of water, has been likewise recommended, as well also as an ointment prepared with an equal part of the sulphate and of lard. (AUGUSTIN)

MANNA.

SYNONIMES:—*μελία*.—Manne, FR.; Manna, GERM.; BOHEM.; DAN.; ITAL.; POL.; PORT.; RUSS.; and SWED.; Mun, Terinjebin, ARAB.; Mana, SPAN.; Schirkischt, HIND.; and PERS.; Gambia, JAP.; Kapurimba, MAL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FIED.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SAUNDERS; SPIELMANN; ZARDA

Several kinds of manna are distinguished:

1. Flake Manna; *Manna canellata* seu *camulata* seu *longa*. This is in pieces hollow on one side, convex on the other, light, whitish or pale-yellow, sometimes reddish, and of an agreeable taste; it is called manna in tears, *Manna lacrymata*, when it is in smaller fragments.

2. Common Manna or Manna in sorts; *Manna in sortis* seu *communis*. This is in round cakes or in oblong pieces, of a yellowish colour, soft, unctuous to the touch, heavy, adhering to the hands, and of a rather nauseous taste. It is called *Geracy* or *Capacy Manna*, as it may have come from Giaraci in Upper Calabria, or from Capachi in Sicily. The grained and white pieces of the first are separated under the name of Choice Manna or Manna in grains, *Manna electa* seu *granulosa*.

3. Fat Manna; *Manna crassa, spissa, sordida* seu *pinguis*. This is very moist, yellowish, soft, glutinous, and loaded with foreign substances.

Manna is obtained from incisions made in the *Fraxinus excelsior*, LINN.; *F. rotundifolia*, LMK.; *F. parviflora*, LMK.; *F. subrefescens*, MÉR.; *F. ornus*, LINN.; *F. argentea*, LOIS. That which exudes spontaneously is rare, and in grains of the size of wheat; it bears the name of *Mastichina*, *Manna difronde*.

Manna contains, according to THENAUD, sugar, a peculiar principle called Mannite, and a nauseous matter.

There are also:

1. Briançon Manna; *Manna Brigantina* seu *laricina*, in small white viscous grains. It exudes, in the South, from the leaves of the *Larix Europæa*.

2. Alhagi Manna; *Manna Persica* (Trunschibil, Tarandyubin, PERS.), in small yellow, reddish, or brown grains. It exudes, in the East, from the leaves and branches of the *Hedysarum alhagi*.

MANNA TROCHISCS. (FEE; HENRI; VAN MONS)

R. Tragacanth, one part.
White Sugar, ninety-six parts.

Triturate together, and add

Manna, twenty-four parts.
Orange-flower Water, a sufficient quantity

to make a paste to be divided into trochiscs. (VAN MONS)

R. Root of Marshmallows, ninety-six parts.
Water, two thousand parts.

Boil for six minutes, and add

Sugar, three thousand parts.
Calabrian Manna, one hundred and ninety-two parts.

Strain, clarify with white of egg, and add

Aqueous Extract of Opium, six parts.
Oil of Bergamotte, ten parts.
Orange-flower Water, ninety-six parts.

Pour the mass into oiled paper moulds, and divide into trochiscs before it becomes cold. (FEE; HENRI)

LAXATIVE JELLY. (PIERQUIN)

R. Veal, } of each,
Water, } two pounds.

Boil for two hours, and add

Manna in Tears, three ounces.

Boil again for two hours, strain, and set apart to cool.—A table-spoonful to be taken every hour.

ELECTUARY OF MANNA.

Electuarium seu *Pulticula de Manna*. (FENN.; PALAT.; SUEC.; WIRTEM.; SWEDIAUR)

R. Manna, three parts.
Oil of Almonds, one part.

Triturate together, and add gradually

Syrup of Manna, one part.

Mix. (VAN MONS)

R. Manna, two ounces.
Fennel Water, one ounce.

Triturate together without heat; strain, and mix with

Sugar,	two ounces.
Florence Iris,	one dram.
Oil of Almonds,	one ounce.

Mix. (VAN MONS)

FENN.; and SUEC. prescribe two ounces each, of manna, of sugar, and of fennel water, one dram of iris, and one ounce of oil of almonds.—PALAT.; and WIRTEM.—two ounces each, of sugar, of manna, and of sloe-flower water, one dram of iris, and one ounce of oil.—WIRTEM.—two ounces each, of manna, of sugar, and of rose water, one dram of iris, and half an ounce of oil.

It is mildly laxative, and is principally recommended for infants.—Dose, from one to four drams.

PECTORAL ELECTUARY. (FEE)

R. Manna, one ounce.
Orange-flower Water,
a sufficient quantity
to soften the manna by trituration. Add
Gum Arabic, in powder, half an ounce.
Soft Extract of Liquorice, one dram.
Syrup of Maidenhair,
a sufficient quantity.

Mix.

ANTI-HÆMORRHOIDAL ELECTUARY. (CADET DE GASSICOURT)

R. Manna in Tears, two ounces.
Sulphate of Potass, } of each,
Nitrate of Potass, } two drams.
Sulphur, }
Honey, a sufficient quantity.

Mix.—Dose, two or three drams daily.

LOHOCH OF MANNA. (HENRI)

R. Oil of Almonds, } of each,
Syrup of Marshmallows, } four parts.
Manna in Tears, }
Orange-flower Water, one part.
Water, two parts.

Dissolve the manna, with a moderate heat, in the water; add the solution to the mixture of the oil and syrup, and mix intimately.

This preparation is proposed by CHEVALIER and IDT; it differs very little from TRONCHIN'S marmalade. There are other analogous preparations under the names of *Linctus mannatus* (HERBIP.; SAUNDERS; SWEDIAUR), *Linctus aperiens* (SAUNDERS), *Mixtura lavans* (PH. SYPH.), and *Potion purgative* (PIERQUIN; SAINTE-MARIE), of which the following are the formulæ:

R. Manna, three ounces.
Oil of Almonds, one ounce and a half.
Triturate, and mix. (SWEDIAUR)

SAUNDERS prescribes an equal part each, of manna and of oil, and a sufficient quantity of yolk of egg.—HERBIP.—one part each, of manna, of oil, and of syrup of violets.—SAUNDERS—one ounce each, of oil and of manna, and half an ounce of syrup of pale roses.

R. Gum Arabic, one ounce.
Oil of Almonds, two ounces.
Decoction of Barley, ten ounces.
Manna, one ounce.
Honey, half an ounce.

Mix. (PH. SYPH.)

R. Castor Oil, triturated } of each,
with Yolk of Egg, } one ounce.
Manna,
Sub-carbonate of Magnesia, two drams.
Hyssop Water, eight ounces.

Mix. (SAINTE-MARIE)

R. Flowers of Peach Tree, } of each,
Violets, } half
a handful.
Spring Water, a sufficient quantity.

Boil for a short time, and dissolve in the strained liquor

Manna, two ounces.
Then add

Oil of Almonds, one ounce.
Mix. (PIERQUIN)

PURGATIVE POTION.

Cassia cum Manna, Decoctum Mannæ purgans seu laxativum seu carminativum, Potio angelica seu laxativa seu anticolica seu Manna tamarindinata seu salina mannata, Aqua angelica Viennensis seu laxativa. (HISP.; FORM. HÔP. FR.; AUGUSTIN; BREERA; RATIER; SPIELMANN)

R. Manna, three ounces.
Water, six ounces.

Dissolve. (RATIER)

R. Manna, two ounces.
Water, one pound.
Cassia Pulp, one ounce.

Mix.—To be taken at twice. (BREERA)

R. Supertartrate of Potass, two drams.
Boiling Water, five ounces.
Manna, two ounces.

Dissolve, and strain. (SPIELMANN)

R. Cassia Pulp, half an ounce.
Tartrate of Potass, one dram and a half.
Manna, two ounces.
Spring Water, five ounces.

Dissolve, and strain. (SPIELMANN)

- R. Tamarind Pulp, three ounces.
Manna, one ounce.
Chamomile Water, nine ounces.

Dissolve, and strain. (AUGUSTIN)

- R. Manna, one ounce.
Sulphate of Soda, one ounce and a half.
Water, six ounces.

Dissolve, and strain. (AUGUSTIN)

- R. Supertartrate of Potass, two drams.
Boiling Water, eight ounces.

Dissolve, and add

- Manna, two ounces.
Tartrate of Potass, one dram and a half.

Juice of Lemon, one.

Infuse the whole with the peel of the lemon; clarify with white of egg, and strain. (PALAT.; and SPIELMANN)

- R. Manna, two ounces.
Senna, six drams.
Sulphate of Magnesia, two drams.
Water, fourteen ounces.

Boil till reduced to twelve ounces, and strain. (HISP.)

- R. Senna, two drams.
Sulphate of Soda, half an ounce.
Water, eight ounces.

Boil for a few minutes, and add,

- Manna, four ounces.

Strain. (FORM. HÔP. FR.)

- R. Water, eight ounces.
Agitate it with white of egg, and add
Senna, three drams.
Manna, three ounces.
Supertartrate of Potass, one dram.

Clarify the liquor by boiling it; then strain it several times through flannel, and add

- Cinnamon Water, one scruple.

To be taken at one dose. (HISP.)

- R. Manna, one ounce.
Senna, } of each,
Supertartrate of Potass, } half
Seeds of Anise, } an ounce.
Water, four pounds.

Infuse for half an hour; then boil a little, and strain

Dose, six ounces. (HISP.)

PURGATIVE EMULSION. (FERR.; HISP.; AUGUSTIN; RATIER)

- R. Almonds, four drams.
Manna in Tears, two ounces.

- Syrup of Peach Flowers, one ounce.
Infusion of Liquorice, four ounces.
Orange-flower Water, four drams.

Make an emulsion. (RATIER)

- R. Almond Emulsion, six ounces.
Manna, three ounces.
Vinous Cinnamon Water, one dram.

Mix. (HISP.)

FERR. prescribes six ounces of emulsion, two ounces of manna, and one dram of the water.

- R. Almond Emulsion, nine ounces.
Manna, one ounce and a half.

Dissolve. (AUGUSTIN)

PURGATIVE WHEY.

Serum Lactis mannatum. (AUGUSTIN; BRERA)

- R. Manna, one ounce.
Supertartrate of Potass, half an ounce.
Clarified Whey, six ounces.

Dissolve.—Dose, half a glassful every two hours. (AUGUSTIN)

- R. Pulp of Tamarinds, } of each,
Manna, } one ounce and
a half.

Whey, a sufficient quantity to obtain two pounds of strained liquor. (BRERA)

ANTI-HYDROPIC POTION. (BRUNS.)

- R. Juice of Iris, three ounces.
Manna, one ounce and a half.

Dissolve.

SYRUP OF MANNA. (BAVAR.; BRUNS.; FULD.; FIDERIT)

- R. Manna, ten ounces.
Spring Water, twelve ounces.

Strain, and add

- White Sugar, one pound.

Boil to the consistence of syrup. (BRUNS.)

FIDERIT prescribes two ounces of manna, twenty ounces of water, and two pounds of sugar.—BAVAR.—one part of manna, four parts of water, and one part and a half of sugar to be added to one part of the strained liquor.—FULD.—one ounce of manna, one ounce and a half of water, and six drams of sugar.

COMPOUND SYRUP OF MANNA.

Syrupus e Manna seu Mannæ cum Senna seu de Manna laxativus. (BRUNS.; DAN.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; WIRTEM.; SPIELMANN)

- R. Senna, four ounces.

Seeds of Fennel, half an ounce.
Boiling Water, two pounds.

Infuse for several hours in a covered vessel, strain with expression, and add to sevenenteen ounces of the strained liquor

White Sugar, two pounds.
Manna, one pound and a half

Boil to a syrup. (OLDEN.)

ROSS. prescribes four ounces of senna, two drams of fennel, two pounds of boiling water, six ounces of manna, and twenty-two ounces of sugar.—HANN.; LIPP.; POL.; and BORUS.—four ounces of senna, half an ounce of fennel, two pounds of water, and the addition of two pounds of sugar and half a pound of manna to eighteen ounces of the strained liquor.

R. Senna, four ounces.
Seeds of Fennel, } of each, one dram.
Ginger, }
Boiling Water, two pounds.

Digest sufficiently, express, and add to the depurated liquor

Calabrian Manna, six ounces.
White Sugar, twenty-two ounces.

Boil to a syrup. (DAN.)

R. Root of Polypodium, one ounce and a half.
Species for the Decoction of Flowers and of Fruits. [No quantity given. EDIT.]
Corinth Raisins, two ounces.
Boiling Water, two pounds and a half.

Infuse during one night, in the morning boil down to two-thirds, and infuse in the strained liquor

Senna, four ounces.
Seeds of Anise, }
Fennel, } of each, one dram.
Galanga, }
Mace, }
Ginger, }

Having digested for twenty-four hours in a warm place; express, clarify with white of egg, and add

Calabrian Manna, six ounces.
White Sugar, twenty-two ounces.

Make a syrup. (BRUNS.; PALAT.; WIRTEM.)

R. Root of Polypody, six drams.
Florentine Iris, half an ounce.
Raisins, three drams.
Honey, four ounces.
Spring Water, twenty-four ounces.

Boil down to twelve ounces; infuse in the strained decoction for twelve hours,

The Four Pectoral Flowers, one handful
Senna, one ounce and a half.
Cinnamon, half a dram.

Strain, and dissolve in the liquor

Calabrian Manna, three ounces.
Sugar, twelve ounces.

Boil to the consistence of a syrup. (SPIELMANN)

MARCHANTIA POLYMORPHA, LINN.

SYNONIMES:—Hépatique des fontaines, FR.; Vielgestaltige Steinmoos, GERM.

GALL.; WIRTEM.; SPIELMANN.

A common plant of Europe. (*Cryptogamia*, LINN.; *Hepaticæ*, JUSS.; *Fig. Dill. Hist. musc.* t. 76 and 77)

The herb (*Herba Hepaticæ fontanæ seu Lichenis stellati seu Lichenis petrei latifolii*) is employed. It is a membranous, plane frond, from two to four inches long, and lobed; the lobes are obtuse, and traversed by a median nervure.

It is slightly astringent.

MARRUBIUM VULGARE, LINN.—WHITE HOREHOUND.

SYNONIMES:—Marrube blanc, FR.; Marrienwurzel, Weisser Andorn, GERM.; Gablenjk, bilj, BOHEM.; Hvid marrube, DAN.; Marrubio blanco, SPAN.; Witte andoorn, DUT.; Marrobio bianco, ITAL.; Szanto biala, POL.; Marroyo branco, PORT.; Andorn, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BELG.; BAVAR.; BRUNS.;

DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A common plant of Europe. (*Didynamia*

Gymnospermia, LINN. ; *Labiata*, JUSS. ;
Fig. Flore Medic. IV. 226)

The herb (*Herba Marrubii albi* seu *Prassii* seu *Phrasi*) is employed. It consists of a tomentose, branching stem, and of oval, petiolated, crenelated leaves, wrinkled above, and tomentose and whitish beneath.

It has a strong odour, which is dissipated by desiccation ; its taste is bitter and rather acrid.

It is stimulant, emmenagogue, anti-spasmodic, febrifuge, and sometimes laxative.

CONSERVE OF HOREHOUND. (AMST.)

- R. Fresh Tops of Horehound, one part.
White Sugar, three parts.
Beat together.

JUICE OF HOREHOUND. (HERBIP.)

- R. Fresh Herb of Horehound,
 any quantity.

Bruise it in a stone mortar, moistening it with a little water, and express strongly ; set apart the juice, decant, and preserve it in bottles, adding to each pound of it one dram of alcohol.—Dose, one or two ounces.

AQUEOUS EXTRACT OF HOREHOUND. (AMST. ; ANT. ; AUSTR. ; DAN. ; FERR. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LU- SIT. ; POL. ; BORUS. ; SAX. ; PIDERIT ; VAN MONS)

- R. Fresh Leaves of Horehound,
 any quantity.

Bruise them with a little water, and strain through a sieve ; set aside the juice, decant, coagulate it at the fire, and strain again : then evaporate to the consistence of pilular mass, remove it from the fire, add the fecula set apart, and evaporate again to the required consistence. (VAN MONS)

- R. Herb of Horehound, one part.
Water, eight parts.

Macerate for twenty-four hours ; then boil for a quarter of an hour, and strain with strong expression ; boil the residue with four parts of water. Mix the two liquors, and after twenty-four hours' repose, evaporate to the consistence of an extract. (LIPP. ; SAX.)

- R. Herb of Horehound, one part.
Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and express ; boil the residue with four parts of water, and express again. Mix, decant after sufficient repose, and evaporate slowly to the consistence of an extract. (DAN. ; HANN. ; OLDEN. ; BORUS.)

AUSTR. ; and POL. direct the herb to be exhausted by several successive ebullitions,

the liquors to be mixed, decanted, and evaporated in a water-bath.

- R. Herb of Horehound, one pound.
Spring Water, ten pounds.

Boil for two hours, express, set aside, strain, and evaporate till the mass no longer adheres to the fingers. (AMST.)

HASS. ; PALAT. ; and PIDERIT direct one part of herb to be boiled in six parts of water, and the expressed liquor to be decanted and evaporated after twenty-four hours' repose.—ANT.—two pounds of herb to be boiled in ten of water for a quarter of an hour, and the decanted decoction to be evaporated.—FERR.—one of herb to be boiled in three of water, the decoction to be evaporated with white of egg, and decanted.

Dose, from a scruple to a dram.

ALCOHOLIC EXTRACT OF HOREHOUND. (FULD. ; HERBIP.)

- R. Herb of Horehound, } of each,
Alcohol. } one pound.
Water, } eight pounds.

Digest for three days at a moderate temperature, and express ; having set aside, and decanted the liquor, distil off the alcohol, and evaporate the rest to the proper consistence.

PECTORAL POTION.

Mixtura Marrubii. (AUGUSTIN)

- R. Extract of Horehound, } of each,
Dog-grass, } two drams.
Dandelion, } ten ounces.
Oxymel of Squill, } of each,
Syrup of Fennel, } two ounces.

Mix.—Dose, half a glassful every two hours.

SYRUP OF HOREHOUND.

Syrupus de Marrubio seu *Marrubii* seu *Prassii albi.* (HISP. ; GALL.)

- R. Dried Leaves of Horehound, two ounces.
Water, two pounds.

Infuse, and add to the strained liquor

Sugar, four pounds.

Make a syrup, and strain. (HISP.)

- R. Dried Leaves of Horehound, one ounce.

Distilled Water of Horehound,
 two pounds.

Digest for two hours in a water-bath, in a covered vessel ; strain, filter, and add

White Sugar, twice the weight.

Make a syrup. (GALL.)

HENRI proposes to make the syrup by mixing one part of the infusion with three parts of simple syrup previously boiled *au boulet*.—Dose, from half an ounce to two ounces.

DISTILLED WATER OF HOREHOUND. (PALAT.; SARD.)

R. Fresh Herb of Horehound, one part.
Water, four parts.
Distil two parts.

MASTICHE.—MASTIC : MASTICH.

SYNONIMES :—*Mastix, Resina Mastichis* seu *Lentisci*, *μαστιχη*.—Mastic, FR.; Mastix, DAN.; GERM.; and SWED.; Arah, Auluk, Bagdadie, AR.; Almaciga, SPAN.; Rumie mustikie, DUK.; and HIND.; Kinneh, PERS.; Mastyx, POL.; Almededa, PORT.; Rumie mastike, TAM.; Sakes, TU.

LOND.; EDINB.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

It is in tears and small irregular masses, dry, friable, yellowish, semi-transparent, of an agreeable odour when heated, softening in the mouth, of an acrid and aromatic taste, and of a bright and vitreous fracture.

This resin flows from incisions made in the bark of the *Pistacia Lentiscus*, LINN., a tree of the middle of Europe. (*Diæcia Pentandria*, LINN.; *Terebinthinacea*, JUSS.; Fig. Blackw. *Herb.* t. 195)

It is tonic, stimulant, and stomachic.

ANTI-LEUCORRHEIC POWDER. (PALAT.)

R. Root of Florentine Iris, half an ounce.
Burnt Hartshorn,
Cuttle Fish Bone,
Prepared White Coral, } of each,
Rock Crystal, } one dram.
Olibanum,
Mastic,

Make a powder.—Dose, from half a dram to two scruples or more.

MASTICATORY TROCHISCS. (AUGUSTIN)

R. Mastic, }
White Wax, } of each, one part.
Ginger, }

Make trochiscs of half a dram each.

ODONTALGIC MASS. (PLANCHE; NIEMANN)

R. Mastic, } of each, two drams.
Sandarac, }
Kino, } ten grains.
Opium, } two grains.
Oil of Rosemary, } eight drops.
Spirit of Scurvy-grass,

a sufficient quantity

to make a soft mass. (NIEMANN)

PLANCHE prescribes two drams each, of mastic and of sandarac, half a dram of dragon's blood, six grains of opium, eight drops of oil of rosemary, and a sufficient quantity of the spirit.

It is used to fill carious teeth.

DECOCTION OF MASTIC. (BRUNS.)

R. Mastic, one ounce.
Crust of Bread, half a pound.
Mace, one dram.
Water, four pounds.

Boil down to two pounds, clarify with hartshorn shavings, and add,

Conserve of Roses, four ounces.
Cinnamon Water, two ounces.

Mix.

SPIRIT OF MASTIC.

Aqua mastichina. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

R. Mastic, four ounces.
Nutmeg, one ounce.
Alcohol, one pound.

Digest for three days, and add,

Water, four pounds.

Distil three pounds. (BRUNS.; WIRTEM.; SPIELMANN)

PALAT. prescribes four ounces of mastic, two ounces of mace, one ounce of cloves, half an ounce of cinnamon, and four pounds of wine.

It is stomachic.—Dose, to one ounce.

COMPOUND SPIRIT OF MASTIC.

Spiritus Mastiches compositus seu matricalis. (AMST.; BRUNS.; HANN.; OLDEN.; PALAT.; BORUS.; SAX.; WIRTEM.; NIEMANN)

R. Mastic, }
Myrrh, } of each, one part.
Olibanum, }
Alcohol, } twenty-four parts.

Macerate for twenty-four hours, and distil two parts. (HANN.; OLDEN.; BORUS.; SAX.; NIEMANN)

BRUNS. prescribes an ounce and a half each, of the three substances, and four pounds of alcohol.

R. Mastic, }
Myrrh, } of each, two ounces.
Olibanum, }
Amber, }
Alcohol (0.907), } three pounds.

Digest for four days, and distil one pound and a half. (AMST.)

PALAT. and WIRTEM. prescribe half an ounce each, of the four substances, and ninety-six ounces of alcohol.

R. Mastic, } four ounces.
Cinnamon, } half an ounce.
Cloves, }
Nutmegs, } of each, three drams.
Ginger, }
Galanga, } of each,
Lesser Cardamoms, } two drams.
WINTER'S Bark, }
Alcohol, } ten pounds.

Digest for several days, and add,
Spring Water, } two pounds.

Distil nine pounds. (BRUNS.; WIRTEM.)

It is employed in lotions, in cases of debility and paralysis.

ELIXIR OF MASTIC.

Elixir Mastichinum. (BRUNS.; WIRTEM.)

R. Spirit of Mastic, } two pounds.
Spring Water, } one pound.
White Sugar, } ten ounces.

Mix.

SYRUP OF MASTIC. (BRUNS.; WIRTEM.)

R. Mastic, in coarse powder, two ounces.
Spirit of Mastic, } twelve ounces.

Infuse for three days in a close vessel; then strain, and add,

White Sugar, } sixteen ounces.

Make a syrup.

OIL OF MASTIC.

Oleum Mastiches seu mastichinum. (BRUNS.; SARD.; WIRTEM.; VAN MONS)

R. Mastic, } one part.
Olive Oil, } four parts.

Dissolve in a water-bath. (SARD.; VAN MONS)

BRUNS. and WIRTEM. prescribe one part each, of mastic and of oil; and also one of mastic and four of oil of roses.

It used formerly to be recommended to arrest vomiting and diarrhoea, &c.

ANTI-ODONTALGIC LIQUOR. (SWEDIAUR)

R. Mastic, } any quantity.
Spirit of Turpentine, }
a sufficient quantity.

Dissolve.

ANTI-ODONTALGIC PLASTER. (BRUNS.; PALAT.)

R. Mastic, }
Gum Caranna, } of each,
— Elemi, } one ounce.
— Tacamahaca, }
— Storax, } half an ounce.
Oil of Nutmeg, } eight scruples.
Turpentine, } half an ounce.

Mix. (PALAT.)

BRUNS. prescribes one ounce and a half of mastic, half an ounce each, of gum anime, of elemi, and of tacamahaca, three ounces of yellow wax, half an ounce of oil of mastic, and two drams of turpentine.

TINCTURE OF MASTIC.

Tinctura seu Essentia mastiches seu vulneraria simplex seu mastichina. (AMST.; BRUNS.; FULD.; SWEDIAUR)

R. Mastic, } one part.
Alcohol (0.884), } eight parts.

Digest for eight days, and strain. (AMST.; FULD.)

BRUNS. prescribes one ounce of mastic and four ounces of concentrated alcohol.—SWEDIAUR—one ounce of mastic and one pound of alcohol (0.935).

It is slightly stimulant; and used formerly to be employed principally in cases of caries and wounds of the bones, and also in diarrhoea.

COMPOUND TINCTURE OF MASTIC. (CADET DE GASSICOURT)

R. Mastic, } of each,
Olibanum, } two ounces and a half.
Myrrh, }
Alcohol, } sixty ounces.

Digest, and filter.

It is stimulant, and is applied to chaps and chilblains of the hands and feet.

MATRICARIA PARTHENIUM, LINN.—COMMON FEVERFEW.

SYNONIMES :—Matricaire, FR. ; Mutterkraut, GERM. ; Matecnjk, Kzimbaba, BOHEM. ; Matrum, DAN. ; Matricaria, ITAL. ; PORT. ; and SPAN. ; Marumaziele, POL. ; Matram, SWED.

AMST. ; ANT. ; BATAV. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; SAX. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUNATELLI ; GUIBOUT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

This plant is common in all parts of Europe. (*Syngenesia Polygamia Superflua*, LINN. ; *Synantherea*, JUSS. ; *Fig. Zorn*, *Id. pl. t. 166*)

The herb (*Herba Matricariæ seu Parthe-*

nii) is employed. It consists of a grooved stem, and of alternate, petiolated, winged leaves, with pinnatifid pinnules the divisions of which are incised and rather obtuse.

The tops present besides, terminal corymbs of radiated cupped corollas, yellow in the centre, and white in the circumference.

Its odour is strong, penetrating, and disagreeable, but is dissipated by drying. Its taste is bitter and aromatic.

It is stimulant, anti-hysteric, emmenagogue, and vermifuge.—Dose, from a scruple to a dram, of the powder ; from one to two ounces of the juice ; and from one to three ounces of the distilled water.—One or two drams is sufficient to make two pounds of infusion.

MELIA AZEDERACH, LINN.—BEAD TREE.

SYNONIMES :—Azéderach Commun, FR. ; Glatter Zidrach, GERM. ; Azadarac, ITAL. ; and SPAN. ; Velsch vygenboom, DUT.

AMER. ; FERR. ; COXE ; SPIELMANN.

A large and beautiful tree of the East Indies, whence it has been transported to the south of Europe and to America. (*Hexandria Monogynia*, LINN. ; *Meliaceæ*, JUSS. ; *Fig. Flore Medic. I. 50*)

The fruit, leaves and bark of the root are employed.

The leaves are broad, twice-winged, having oval leaflets, pointed, dentated, and glabrous.

The fruit consists of round, fleshy nuts ;

the external rind is rather thick ; the noyau is grooved, it presents five monospermal cells ; the kernel is oleaginous.

It is anthelmintic, and slightly narcotic. THATCHER extols an ointment, prepared with the pulp of the fruit and lard, in tinea. The fruit, said erroneously by PERSOON to be poisonous, is eaten with impunity by children, in America, against worms.

DECOCTION OF BEAD TREE ROOT.
(NIEMANN)

R. Bark of Bead Tree Root, four ounces.
Spring Water, one pound.
Boil, and strain.

It is stimulant, and anthelmintic.—Dose, one ounce every hour.

MELILOTUS.—MELILOT ; MELILOT TREFOIL.

Two species of this genus of plants are mentioned in pharmacopœias :

1. *Melilotus cæruleus*, LMK.—Blue Melilot.

SYNONIMES :—Mélilot bleu, Faux baume du Perou, Lotier odorant, Tréfle musqué, FR. ; Blauer steinklee, Siebengezeit, GERM. ; bleauwe melilot, DUT.

VOL. II.

GALL. ; WIRTEM. ; SPIELMANN.

A plant of Europe. (*Diadelphia Decandria*, LINN. ; *Leguminosæ*, JUSS. ; *Fig. Blackw. Herb. t. 284*)

The flowering herb (*Herba Ægyptiaca seu Meliloti cærulei seu Loti odoratæ seu urbanæ*) is employed. It consists of a branching stem, and of leaves furnished at
2 G

their base with large dentated stipulæ, and composed of three oval folioles finely serrated; the flowers, of a pale blue colour, firm clustered oval spike under, and are placed upon long axillary peduncles. It has a strong but not unpleasant smell, and a rather acrid taste.

2. *Melilotus officinalis*, LMK.—Common Melilot.

SYNONIMES :—Melilot officinal, FR.; Steinklee, Melilotenklie, GERM.; Komonice, BOHEM.; Amur, Steenklæver, Meloten, DAN.; Meliloto, ITAL.; PORT.; and SPAN.; Meloten, DUT.; and SWED.; Komonica Swoyska, POL.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; GENEV.; HAMB.; HANN.; HASS.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN.; PIDERIT.; ZARDA.

A common plant of Europe. (*Fig. Zorn, Ic. pl. t. 204*)

The flowering herb (*Herba Meliloti citrini*) is employed. It consists of a smooth and branching stem, furnished with oval, serrated leaves; the flowers are yellow, and disposed in long racemes, proceeding from the axillæ of the superior leaves. Its odour is strong and agreeable, and its taste bitter and rather acrid.

These two plants are emollient.

RESOLVENT APPLICATION. (BORIES)

R. Powdered Flowers of Melilot, one ounce.
Camphor, one dram.

Mix.—To be applied in a bag.

OIL OF MELILOT. (GALL.; SARD.; VAN MONS)

R. Flowers of Melilot, one part.
Olive Oil, six parts.

Infuse in a moderate heat for three days; then strain and filter. (VAN MONS)

SARD. directs one part of flowers to be digested in two parts of oil for three days in a water-bath; the oil to be expressed when cold, and the maceration repeated a second time.—GALL.—one of flowers to be macerated in the sun for three days in four of oil; the operation to be repeated thrice, and the last time to be continued for three months.

It is employed externally, in scrofula and recent chilblains.

PLASTER OF MELILOT.

Emplastrum Meliloti seu Trifolii Meliloti officinalis. (ANT.; BATAV.; BELG.; DAN.; MIL. COPEN.; HISP.; FULD.; HAMB.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; WIRTEM.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

R. Yellow Wax, }
Resin, } of each, one part.
Olive Oil, }

Melt together, and add when nearly cold,
Powdered Herb of Melilot, two parts.

Mix. (SAX.; SWEDIAUR)

FENN.; HANN.; POL.; BORUS.; and SUEC. prescribe two parts of wax, one part each, of resin and of oil, and two parts of powdered melilot.—OLDEN.—fourteen ounces of wax, six ounces each, of resin and of oil, one pound of powdered melilot, and one ounce of extract of melilot.—ROSS.—three parts of wax, six of resin, four of mutton suet, one of oil, and twelve of powdered melilot.—ANT.—eight of wax, four of resin, two and a half of turpentine, and four of powdered melilot.

R. Flowering Tops of Melilot, three parts.
Goat's Suet, four parts.

Boil to the consumption of the humidity; strain, and add

Resin, six parts.
Wax, three parts.

Melt together. (SPIELMANN)

DAN. prescribes four parts of flowering tops, five of goat's suet, eight of resin, and four of wax.—FULD.—three of tops, two of suet, three of resin, and one and a half of wax.—VAN MONS—eight of yellow wax, four of resin, one of turpentine, and four of tops; to be digested for twelve hours, and one part of liquid storax added.

R. Plaster of Lead, one pound.
Yellow Wax, }
Olive Oil, } of each, six ounces.

Add to the melted mixture when half cold
Powdered Melilot, eight ounces.

Mix. (BELG.)

BATAV. gives the same formula, but requires one pound of melilot.

R. Resin Plaster, one pound and a half.
Olive Oil, one pound.
Resin, six pounds.
Yellow Wax, three pounds.
Gum Ammoniac, four ounces.

Melt together, and when nearly cold add gradually

Powdered Flowering Herb of Melilot, two pounds.
 ————— Worm- one pound.
 wood,

Mix. (DAN.)

MIL. COPEN. prescribes twenty-six ounces of mutton suet, thirty-six of resin, eighteen of yellow wax, six of rape oil, two of gum ammoniac, twelve of powdered melilot, and six of powdered wormwood.—HASS.; and PIDERIT—one pound and a half of resin plaster, five ounces of powdered melilot, two ounces of powdered marshmallows root, one ounce of gum ammoniac, and two ounces of Venice turpentine.

R. Oil of Melilot, two pounds.
 Goat's Suet, one pound.
 Yellow Wax, eight ounces.
 Turpentine, six ounces.

Melt together, and add

Gum Ammo- } of each,
 niac, } one ounce and a half.
 Sagapenum, }
 Powdered Melilot, six ounces.

Mix. (SARD.)

LIPP. prescribes one pound and a half of resin plaster, six ounces of powdered melilot, and one ounce of gum ammoniac.

R. Flowering Herb of Melilot, four ounces.

Flowers of Wormwood, }
 ————— Chamomile, } of each,
 Bay Berries, } six drams.
 Seeds of Smallage, }
 Storax, half an ounce.
 Gum Ammoniac, one ounce.

Incorporate with a melted mixture of

Yellow Wax, eight ounces.
 Goat's Suet, four ounces.
 Turpentine, } of each,
 Resin, } one ounce and a half.
 Oil of Chamomile, two ounces.

Mix. (BRUNS.; PALAT.; WIRTEM.)

R. Simple Melilot Plaster, three pounds.
 Turpentine, six ounces.

Melt slowly, and add, in powder,

Flowers of Melilot, } of each,
 ————— Chamomile, } three ounces.
 Root of Iris, } of each,
 ————— Marshmallows, } one ounce.
 Saffron, half an ounce.
 Gum Ammoniac, } of each,
 Bdellium, } two ounces.
 Fenugreek, one ounce.

Mix. (HISP.)

CAMPFORATED MELILOT PLASTER. (MIL. COPEN.)

R. Melilot Plaster, two ounces.
 Camphor, in powder, one dram.

Mix carefully.

MELISSA.—BALM.

Three species of this genus are used in medicine:

1. *Melissa calamintha*, LINN.—Calamint;
 Mountain Balm.

SYNONIMES:—Calament de montagne, Bergmüze, GERM.; Plana mata, Poleg, Natkowe korenz, Marulka hornj, BOHEM.; Berg calaminth, DUT.

BRUNS.; HISP.; GALL.; FERR.; GENEV.; WIRTEM.; HERBIP.; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the south of Europe. (*Didynamia Gymnospermia*, LINN.; *Labiate*, JUSS.; *Fig. Zorn, Ic. pl. t. 111*)

The herb (*Herba Calaminthæ seu Calaminthæ montanæ seu Calmontanæ*) is employed. It consists of a pubescent stem, furnished with oval, pubescent leaves, somewhat heart shaped at the base, and obtusely dentate. Its odour is strong and aromatic,

diminishing very much by desiccation; its taste is acrid and aromatic.

It is stimulant and nervine.

2. *Melissa Nepeta*, LINN.—Field Balm.

SYNONIMES:—Calament des Champs, Petit calament, FR.; Poleyartige Kalamint, GERM.; Veldkalaminth, DUT.

GALL.; HERBIP.

This plant is spread over all Europe. (*Fig. Blackw. Herb. t. 167*)

The herb (*Herba Nepetæ seu Calaminthæ Pulegii odore seu Calaminthæ agrestis*) is employed. It presents a rather hairy stem, furnished with nearly round, glabrous leaves, having two or three unequal teeth on either margin. It has an aromatic taste and smell.

It is stimulant and nervine.

3. *Melissa officinalis*, LINN.—Common Balm.

SYNONIMES:—Melisse officinale, FR.; Melisse, Melissenkraut, Citronenkraut,

GERM.; Bucklitulfaristum, ARAB.; Melissa Medunka, BOHEM.; Hiertenfryd, DAN.; Mekka subza, DUK.; Torongil, SPAN.; Citronenkruid, DUT.; Badrunj-buych, PERS.; Melisa cytrynowa, POL.; Herva cidreira, PORT.; Citron-meliss, SWED.

EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of the south of Europe. (*Fig. Zorn, Ic. pl. t. 135*)

The herb (*Herba Melissa citrina seu citrate seu hortensis seu Cedronelle*) is employed. It consists of a tetragonal, branching, nearly glabrous stem, and of petiolated, oval leaves, slightly serrated at the base, and crenelated at the borders. It has an agreeable lemon-like odour, and an aromatic and rather acrid taste.

It ranks among the stimulants and antispasmodics.

AROMATIC SPECIES.

Species pro Thea. (HUFELAND)

- R. Herb of Balm, one ounce.
Fennel, three drams.
Liquorice, two drams.
Cut, and mix.

CONSERVE OF BALM. (ROSS.)

- R. White Sugar, two parts.
Fresh Leaves of Balm, one part.
Mix, and preserve in a cool place.

AROMATIC CATAPLASM. (FORM. HÔP. FR.; BRUGNATELLI)

- R. Chamomile Flowers, three ounces.
Herb of Balm, six ounces.
Juniper Berries, four ounces.
Brandy, a sufficient quantity.

Boil to the consistence of a soft paste. (BRUGNATELLI)

FORM. HÔP. FR. directs one hundred parts each, of barley and of rye meals to be boiled in one hundred of red wine, and one third of aromatic powder to be added towards the end.

EXTRACT OF BALM. (FERR.; SARD.)

- R. Herb of Balm, one part.
Water, three parts.

Boil, express, clarify with white of egg, and evaporate the liquor to the proper consistence. (FERR.)

- R. Fresh Herb of Balm, eight pounds.

Bruise it and digest it for three days in an alembic; then distil from it about two ounces of water in a water-bath; pour thrice its weight of water upon the residue, boil down to two-thirds, strain, clarify, and evaporate the decoction to the consistence of thick electuary, and add the distilled water to it when it is only warm. (SARD.)

INFUSION OF BALM. (FORM. HÔP. FR.; RATIER)

- R. Leaves of Balm, half an ounce.
Boiling Water, two pounds.
Infuse, and strain. (RATIER)

- R. Dried Herb of Balm, } of each, two
Liquorice, } drams and a half.
Boiling Water, a sufficient quantity
to obtain two pounds of strained liquor. (FORM. HÔP. FR.)

It is slightly stimulant; and is recommended in dyspepsia attributed to debility of the stomach.

CARMINATIVE POTION. (CADET DE GASSICOURT)

- R. Leaves of Balm, two drams.
Anise, }
Fennel, } of each, half a dram.
Coriander, }
Boiling Water, two pounds.

Infuse for a quarter of an hour; strain, and add

- White Sugar, a sufficient quantity.
Mix.

SYRUP OF BALM. (WIRTEM.)

- R. Dried Leaves of Balm, four ounces.
Distilled Water of Balm, one pound.
Digest for two days in a covered vessel; express, and add

- White Sugar, sixteen ounces.
Boil for a few moments.

COMPOUND ESSENCE OF BALM. (VAN MONS)

- R. Leaves of Balm, two parts.
Rue, } of each,
Pennyroyal, } one part
Fresh Lemon Peel, } and a half.
Cinnamon, } one part.
Spirit of Balm, twenty-four parts.

Infuse without heat for several days; then express and filter.

TINCTURE OF BALM. (ANT. ; VAN MONS)

- R. Herb of Balm, one part.
 Alcohol (20 degrees), eight parts.
 Infuse for three days, and filter.
 It is stimulant.

OIL OF BALM.

Oleum Melissa æthereum, Ætheroleum Melissa. (BRUNS. ; FERR. ; HANN. ; PALAT. ; ROSS. ; WIRTEM. ; HERBIP. ; PIDERIT ; SWEDIAUR)

- R. Fresh Herb of Balm, any quantity.
 Water, a sufficient quantity.
 Distil, and preserve the oil. (FERR. ; ROSS.)

SWEDIAUR prescribes one part of herb and four parts of water.—HANN.—one of herb and eight of water.

- R. Fresh Herb of Balm, twenty-five parts.
 Water, seventy-five parts.
 Muriate of Soda, three parts.
 Distil. (BRUNS. ; PALAT. ; WIRTEM. ; PIDERIT)

- R. Fresh Herb of Balm, any quantity.
 Distilled Water of Balm, a sufficient quantity.

Distil. (HERBIP.)

DISTILLED WATER OF BALM. (AMST. ; ANT. ; AUSTR. ; BAVAR. ; BRUNS. ; DAN. ; GALL. ; FERR. ; FULD. ; GENEV. ; HASS. ; LIPP. ; OLDEN. ; PALAT. ; POL. ; BORUS. ; SAX. ; SARD. ; WIRTEM. ; WURCEB. ; BRUGNATELLI ; PIDERIT ; VAN MONS)

- R. Fresh Herb of Balm, one part.
 Water, a sufficient quantity.
 Infuse for twenty-four hours, and distil ten parts. (AMST. ; OLDEN. ; POL. ; BORUS.)

DAN. ; and FEE prescribe one part of herb and a sufficient quantity of water ; twenty parts to be distilled.—HANN.—the same, but three parts to be distilled.—GENEV.—the same, but six to be distilled.—GALL. ; and PALAT.—one of herb and four of water ; two to be distilled.—ANT.—one of herb and five of water ; four to be distilled.—HERBIP. ; and BRUGNATELLI—one of herb and six of water ; four to be distilled.—HASS. ; and PIDERIT—one of herb and eight of water ; four to be distilled.—AUSTR.—one of herb and twelve of water ; four to be distilled.—SAX.—one of herb and fifteen of water ; ten to be distilled.—BAVAR.—one of herb and sixteen of water ; four to be distilled.

- R. Fresh Herb of Balm, one part.
 Water, four parts.

Distil one half, and re-distil with fresh herb. (SARD.)

- R. Fresh Herb of Balm, three parts.
 Suspend it in a bag in the centre of a cucurbit containing

Water, twelve parts,
 in such a manner that it may not touch the water, and distil six parts. (FULD. ; LIPP.)

- R. Flowering Herb of Balm, one part.
 White Wine, four parts.
 Distil one half. (BRUNS. ; WIRTEM.)

VAN MONS prescribes ten parts of flowering herb, one half part of proof spirit, and one hundred parts of water ; forty parts to be distilled.

ANTI-HYSTERIC LIQUOR. (WIRTEM.)

- R. Fresh Leaves of Balm, one pound.
 Bay Berries, } of each, one ounce.
 Seeds of Cumin, }
 Myrrh, half an ounce.
 Castor, two drams.
 Wine, twelve pounds.

Digest, and distil one half.

SPIRIT OF BALM. (ANT. ; BRUNS. ; GALL. ; FULD. ; LIPP. ; LUSIT. ; PALAT. ; ROSS. ; SARD. ; SAX. ; WIRTEM. ; HERBIP. ; PIDERIT ; VAN MONS)

- R. Herb of Balm, one part.
 Alcohol (20 degrees), two parts.
 Water, a sufficient quantity.
 Distil a spirit marking 28 degrees.

VAN MONS prescribes one part of herb, three parts of alcohol, and six of water ; two and a half to be distilled.

- R. Herb of Balm, one part.
 Water of Balm, two parts.
 Alcohol, eight parts.

Macerate for twenty-four hours, and distil six parts. (ROSS.)

GALL. prescribes twelve parts of herb, twelve of water of balm, and thirty-six of alcohol (22 degrees) ; thirty to be distilled.

- R. Herb of Balm, one part and a half.
 Proof Spirit, ten parts.

At the end of twelve hours distil one half. (LUSIT.)

BRUNS. ; PALAT. ; SAX. ; WIRTEM. ; and PIDERIT prescribe one part of herb and four parts of proof spirit.—FULD. ; and LIPP.—one of herb and six of proof spirit.

- R. Fresh Herb of Balm, one part.
 Alcohol, four parts.

Digest for twenty-four hours, and distil in a water-bath almost to dryness ; digest and distil the product again with fresh herb. (SARD.)

HERBIP. prescribes one pound and a half

of herb and ten pounds of alcohol; and the product to be re-distilled with three ounces of fresh herb.

EAU DES CARMES.

Aqua aromatica, Spiritus aromaticus, Aqua Melissa composita seu Carmelitana seu Carmelitarum seu Melissa spirituosa composita. (AMST.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; LUSIT.; PALAT.; SARD.; WIRTEM.; BRUGNATELLI; CADET DE GASSICOURT; HUFELAND; SPIELMANN; SWEDIAUR; VAN MONS.)

R. Fresh Herb of Balm, one pound and a half.
 — Lemon Peel, four ounces.
 Nutmegs, } of each, two ounces.
 Coriander, }
 Cinnamon, }
 Cloves, } of each, one ounce.
 Alcohol, six pounds.
 Water of Balm, three pounds.

Digest for three days, and distil six pounds. (BRUNS.; DAN.; SARD.; WIRTEM.)

R. Herb of Balm, twelve ounces.
 Lemon Peel, two ounces.
 Nutmegs, one ounce.
 Coriander, four ounces.
 Cloves, } of each, one ounce.
 Cinnamon, }
 Root of Angelica, half an ounce.
 Alcohol, sixty-four ounces.

Infuse for several days, and distil eighty-eight ounces. (FERR.; SPIELMANN)

HANN. prescribes eighteen ounces of balm, four of lemon peel, three each, of nutmegs and of angelica root, one ounce each, of cloves and of cinnamon, and three pounds each, of alcohol and of balm water; to be digested for three days, and seven pounds distilled. — BRUGNATELLI — two pounds of balm, four ounces of lemon peel, two of nutmegs, eight of coriander, half an ounce each, of cinnamon and of cloves, and one hundred and twenty ounces of alcohol; to be digested for three days in a water-bath, and one hundred and twenty ounces distilled. — FULD. — sixteen ounces of balm, four of lemon peel, two of cinnamon, one of nutmegs, two drams of cloves, and eight pounds of white wine; to be digested for twenty-four hours, and four pounds distilled. — LUSIT. — two pounds of balm, four ounces of lemon peel, two of cinnamon, one of nutmegs, two drams of cloves, and ten pounds of proof spirit; to be macerated for three days, and five pounds distilled. — AMST.; BATAV.; and BELG. — four ounces of balm, one ounce each, of lemon peel, of nutmegs, and of coriander, half an ounce

each, of cloves and of cinnamon, and five pounds of proof spirit, to be digested for three days, and three pounds distilled. — HISP. — eight ounces of balm, two each, of lemon peel and of orange peel, one ounce each, of nutmegs and of coriander, half an ounce of cinnamon, and four pounds of proof spirit; to be macerated for twenty-four hours, and one half distilled. — SWEDIAUR — one ounce of nutmegs, four ounces of coriander, one ounce and a half each, of cinnamon and of angelica root, one ounce of cloves, two ounces of lemon peel, two pounds of water of balm, and four of alcohol; to be digested for eight days, and distilled. — AUSTR. — two pounds of balm, the peels of ten lemons, three ounces each, of cinnamon and of angelica root, six ounces of coriander, eleven pounds of alcohol (0.910), and six pounds of spring water; ten pounds to be distilled.

R. Herb of Balm, }
 — Peppermint, } of each,
 — Sage, } three ounces.
 — Rosemary, }
 Flowers of Lavender, }
 Fennel, }
 Root of Angelica, two ounces.
 — Sweet Flag, four ounces.
 Alcohol, four pounds.
 Water, a sufficient quantity.

Macerate for twenty-four hours, and distil twelve pounds. (HUFELAND; NIEMANN)

R. Flowers of Red Roses, } of each,
 — Rosemary, } four ounces.
 — Lavender, } of each,
 — Water Lily, } three
 handfuls.
 Herb of Marjoram, } of each,
 — Balm, } one handful.
 Lesser Cardamoms, }
 Cloves, } of each,
 Mace, } one ounce.
 Cubebs, }
 Cinnamon, two ounces.
 Spirit of Bird Cherry, six pounds.

Digest for six days, and distil. (PALAT.)

R. Cinnamon, }
 Cloves, } of each, one part.
 Nutmeg, }
 Anise, }
 Coriander, }
 Hyssop, }
 Rosemary, } of each, two parts.
 Marjoram, }
 Sage, }
 Thyme, }
 Fresh Lemon Peel, } of each, four parts.
 Root of Angelica, }
 Fresh Balm, twelve parts.
 Proof Spirit, one hundred and ninety-two parts.
 Water, five hundred and seventy-six parts.

Distil one hundred and twenty-eight parts.
(VAN MONS)

- R. 1. Balm, twenty parts.
 Angelica, five parts.
 Alcohol (22 degrees), two hundred and fifty parts.
 2. Cinnamon, five parts.
 Cloves, } of each, three parts.
 Nutmegs, }
 Alcohol, one hundred and ten parts.
 3. Anise, two parts.
 Coriander, five parts.
 Dried Lemon Peel, one part.
 Alcohol, eighty parts.
 4. Rosemary, } of each, two parts.
 Marjoram, }
 Hyssop, }
 Thyme, }
 Sage, four parts.
 Alcohol, one hundred and twenty parts.

Distil separately in a water-bath; then mix together the four products, add one tenth of the total weight of water, and distil four-fifths in a water-bath. (FORM. HÔP. FR.)

- R. Cinnamon, }
 Cloves, } of each,
 Nutmegs, } one part.
 Anise, }
 Coriander, }
 Dried Lemon Peel, }
 Alcohol (22 degrees), one thousand parts.

Macerate each separately for three days in ninety-six parts of alcohol; then distil in a water-bath, and receive the products separately. On the other hand, infuse for two days, in the same quantity of alcohol, distilling each separately,

- Herb of Angelica, }
 Flowering Herb of Rose- }
 mary, } of each,
 Mar- } one part.
 joram, }
 Hyssop, }
 Thyme, }
 Sage, }

Macerate and distil in the same manner,
 Fresh Leaves of Balm, one part.
 Then take three vessels; pour into the first

- Spirit of Cinnamon, } of each,
 Coriander, } three parts and
 a half.
 Cloves, } of each,
 Nutmeg, } three parts.
 Anise, two parts.
 Lemon Peel, one fourth part.

Into the second

- Spirit of Angelica, ten parts.
 Rosemary, six parts.
 Marjoram, } of each,
 Thyme, } seven parts.
 Hyssop, eight parts.
 Sage, fifteen parts and a half.

Pour into the third the spirit prepared above. Then mix five parts of the contents of the first vessel with five of the second and five and a half of the third; pour this mixture into a cucurbit, add one-tenth of the whole of spring water, and one-eightieth of sugar in powder, and distil till only one-fifth remains in the cucurbit. (GALL.; CADET DE GASSICOURT)

This last formula offers one example, among many, of the anxious care of the French codex for the preservation of empirical formulæ, and that too with all their most insignificant minutiae. The *Eau des Carmes* will no doubt keep its place for a long while in domestic medicine, but it is time that such toys should be expunged from the pages of every pharmacopœia, and left entirely to the perfumer's receipt-book. If it is thought necessary to have a compound spirit of balm, which does not, however, appear evident, one of the formulæ placed at the beginning of this article would do very well.

DENTIFRICE LIQUOR. (PIERQUIN)

- R. Alcohol, twenty ounces.
 Spirit of Cinnamon, }
 Lavender, } of each,
 Lemon, } five drams.
 Compound Spirit of Balm, }

Mix.

AROMATIC VINEGAR.

Acetum odoratum seu bezoardicum seu odoriferum. (BRUNS.; FULD.; NIEMANN)

- R. Herb of Lovage, two ounces.
 Leaves of Balm, } of each, four ounces.
 Rue, }
 Green Walnuts, sliced, two pounds.
 Lemons, sliced, one pound.
 Vinegar, ten pounds.

Digest for twelve hours, and distil almost to dryness. (BRUNS.)

- R. Vinegar, two ounces.
 Spirit of Balm, half an ounce.
 Oil of Lemon, } of each, ten drops.
 Lavender, }
 Cloves, four drops.

Mix. (FULD.; NIEMANN)

MELITTIS MELISSOPHYLLUM, LINN.—BASTARD BALM.

SYNONIMES :—Melitte à feuilles de mélisse, FR. ; Wanzenmelisse, GERM. ; Vild hier-teus fryd, DAN. ; Melissebladig kruis, bloem, DUT.

GALL. ; WIRTEM. ; HERBIP. ; MURRAY and GMELIN ; SPIELMANN.

This plant is rather common in Europe. (*Didynamia Gymnospermia*, LINN. ; *La-biata*, JUSS. ; Fig. Jacq. *Fl. Austr.* t. 26)

The herb (*Herba Melissophylli* seu *Lamii montani* seu *Melissæ humilis*) is employed. It consists of a tetragonal, hairy stem, and of petiolated, oval, oblong, pointed, crenelated leaves. It has a disagreeable odour, in the fresh state, but develops an agreeable aroma when dried. Its taste is bitterish.

It is stimulant ; it was administered as an emmenagogue, and also to effect the expulsion of urinary gravel.

MELOE.

Two insects of this genus have been introduced into medicine :

1. *Meloë majalis*, OLIV.—Oil Beetle.

SYNONIMES :—Meloë de Mai, FR. ; May-kæfer, Maywurm, GERM. ; Olie bille, DAN. ; Carralejas, Aradijos, SPAN. ; Meywurm, DUT. ; Majbagyl, SWED.

FERR. ; FULD. ; HANN. ; SAX. ; WIRTEM. ; BRUGNATELLI ; GUIBOURT.

A coleopterous heteromorous insect common in Europe. (*Trachelidæ Cantharidæ*, LATR. ; Fig. Leach, *Trans. Soc. Lin.* XI. VI. 1. 2)

The colour of its body is a mixture of bronze and of cupreous-red.

2. *Meloë proscarabeus*, LINN.

BRUNS. ; DAN. ; HISP. ; FERR. ; FULD. ; HANN. ; ROSS. ; SAX. ; WIRTEM. ; GUIBOURT ; SPIELMANN.

An insect (*Vermis majalis* seu *Scarabeus unctuaris*) of a violet black colour, shagreened and shining.

ANTI-LYSSIC ELECTUARY. (FULD. ; WIRTEM. ; NIEMANN)

R. Meloës, without heads, twenty-four.
Theriac, two ounces.
Wood of Yew, two drams.
Root of Serpentaria, one dram.
Bedeguar of the Ash, twenty grains.
Honey, in which the insects have been suffocated, a sufficient quantity.
Make an electuary. (WIRTEM.)

R. Meloë, one.
Honey, in one pound of which eighty insects have been suffocated, a sufficient quantity to make a pulp.

Theriac, forty grains.
Oil of Meloës, by infusion, six drops.
Wood of Ebony, a little.
Rob of Elder, a sufficient quantity.

Mix. (NIEMANN)

For one dose.

This formed a celebrated remedy for hydrophobia up to the period of 1782, when FREDERIC II. purchased the secret of it ; from which time we believe it has not been employed. A specific can only live in the dark, and dies immediately when it is exposed to light.

Although meloës will not cure hydrophobia, yet they may be useful, by virtue of their acridity, for they are almost vesicant, as an external application ; in Sweden, they are said to be used with success in the cure of rheumatic affections.

FULD. prescribes a *Mel Vermium majalium* ; it is prepared by digesting forty or fifty insects in a pound of honey.

SPIELMANN, under the title of *Oleum Scarabeorum*, gives a preparation considered nervine and resolvent ; it is obtained by boiling, to the consumption of humidity, eight ounces of dusky beetles (*scarabées de fumier*) in sixteen ounces of oil of bay.

ANTI-LYSSIC POTION. (AUGUSTIN ; NIEMANN)

R. Meloës, eight.
Theriac, half an ounce.
Sub-carbonate of Ammonia, two drams.
Camphor, one dram.
Solution of Acetate of Ammonia, eight ounces.

Mix.—The authority of SELLE has not been sufficient to keep this remedy in credit.

MENTHA.—MINT.

Of the species of this genus of plants which are used in medicine, we shall quote the ten following:

1. *Mentha gentilis*, LINN.

SYNONIMES :—Menthe apparente, Baume des jardins, FR.

GALL.; BERGIUS; GUIBOURT; SPIELMANN.

A plant (*Didynamia Gymnospermia*, LINN.; *Labiata*, JUSS.; *Fig. Céd. Fl. Dan.* t. 736)

The herb (*Herba Menthae balsaminæ*) is employed. It consists of a rather hairy stem, very branched, and furnished with oval, dentated, petiolated leaves. It has an agreeable odour, and an acrid and bitter taste.

2. *Mentha aquatica*, LINN.—Water Mint.

SYNONIMES :—Menthe aquatique, FR.; Wassermünze, GERM.; Cywerwenamata, BOHEM.; Hestermynthe, DAN.; Yerba buena dos rios, SPAN.; Menta acquatica, ITAL.

AUSTR.; GALL.; FERR.; FORM. HÔP. FR.; WURCEB.; GUIBOURT; SPIELMANN; ZARDA.

A plant of Europe. (*Fig. Blackw. Herb.* t. 32)

The herb (*Herba Menthae aquaticæ seu rubræ seu Balsami palustris*) consists of glabrous stems, bearing glabrous, oval, dentated, petiolated leaves. It has a strong and not very agreeable odour, and a bitter and rather acrid taste.

3. *Mentha arvensis*, LINN.—Corn Mint.

SYNONIMES :—Menthe des Champs, Pouliot thym, FR.; Ackermünze, GERM.

GALL.

This is a common plant of Europe. (*Fig. Céd. Fl. Dan.* t. 512)

The herb (*Herba Menthae equinæ seu sylvestris*) is employed. It consists of a square, feeble, branching stem, and of opposite, oval, pointed, obtusely dentated, petiolated, greyish leaves. It has a disagreeable odour, and a bitter and sub-acrid taste.

4. *Mentha Crispa*, LINN.—Curled Mint.

SYNONIMES :—Menthe crêpue, FR.; Krause münze, Blumenkopfkrausemünze, GERM.; Kaderawa mata, Balssam zahradny, BOHEM.; Krusemyute, DAN.; Menta rizada, SPAN.; Kuizenmunt, DUT.; Menta crispa, ITAL.; Faka, JAP.; Fodor

mentha, M.; Mienta kedzierzawa, POL.; Hortela vulgar, PORT.; Miata, RUSS.; Krusmyuta, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; LUSIT.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A plant of Europe. (*Fig. Flore Medic.* V. 232)

The herb (*Herba Menthae crispæ*) is employed. It consists of tomentose stems, and of leaves wrinkled above, tomentose and white beneath, of an oval or rounded figure, and bordered with large unequal teeth. Its odour is strong, balsamic, peculiar, and not very agreeable; its taste is warm, aromatic, and bitterish.

5. *Mentha sativa*, LINN.—Spear Mint.

SYNONIMES :—Menthe cultivée, FR.; Zahme münze, Frauenmünze, GERM.; Nana, Hibbuk, ARAB.; Pudina, DUK.; and POL.; Menta Romana, ITAL.; Widatillum, TAM.

DUBL.; HISP.; AINSLIE; PARIS.

A plant of Europe. (*Fig. Céd. Fl. Dan.* t. 794)

The herb (*Herba Menthae verticillatæ seu crispæ verticillatæ*) is employed. It consists of a simple or slightly branching, glabrous stem, and of oval, acute, dentate, petiolated leaves. Its odour and taste, resemble those of the *M. Crispa*.

6. *Mentha rotundifolia*, LINN.—Round-leaved Mint.

SYNONIMES :—Menthe à feuilles rondes, Baume Sauvage, FR.; Mastranzo, SPAN.

HISP.; GALL.

A plant of Europe.

The herb (*Herba Menthae sylvestris folio rotundiore*) is employed. It consists of a tomentose stem, and of sessile, oval or rounded leaves, wrinkled above, tomentose and whitish beneath, and toothed at the borders.

Its odour is strong and agreeable.

7. *Mentha piperita*, LINN.—Peppermint.

SYNONIMES:—Menthe poivrée, FR.; Pfeffermünze, GERM.; Peprowa mata, BOHEM.; Pebermynte, DAN.; Verhabuena de sabor de pimienta, SPAN.; Pepermin-do, DUT.; Mienta pieprzna, POL.; Hortela apimentada, PORT.; Peparmynta, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; COXE.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

This plant appears to belong originally to England, but it is now spread over the whole of Europe. (*Fig. Flore Medic. V. 233*)

The herb (*Herba Menthae piperita* seu *piperitidis*) is employed. It consists of a branchy, rather hairy stem, and of petioled, oval, acute or lanceolated, dentated leaves, glabrous and of a deep-green colour above.

It has an aromatic, penetrating, agreeable odour. Its taste is balsamic, pungent, and camphorous, and leaves a sensation of coldness in the mouth.

8. *Mentha sylvestris*, LINN.

SYNONIMES:—Menthe sauvage, FR.

GALL.; BERGIUS.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

A common plant of Europe. (*Fig. Zorn, Ic. pl. t. 354*)

The herb (*Herba Menthae sylvestris* seu *longifoliae*) is employed. It is tomentose, and consists of a stem furnished with sessile, oblong, lanceolated, unequally toothed, whitish leaves. It has a strong and agreeable odour, and a bitter taste.

9. *Mentha viridis*, LINN.—Green Mint.

SYNONIMES:—Menthe verte, Baume verte, FR.

LOND.; AMER.; FERR.; GENEV.; SAX.; BERGIUS.; COXE.; GUIBOURT.

A plant of Europe. (*Fig. Blackw. Herb. t. 290*)

The flowering herb (*Herba et Flores Menthae tri*) is employed. It consists of a

glabrous stem, furnished with lanceolated, sessile, sparsely-toothed leaves, and terminated by a long spike of purple verticillated flowers.

Its odour is agreeable and aromatic.

10. *Mentha Pulegium*, LINN.—Pennyroyal Mint.

SYNONIMES:—Pouliot, FR.; Poley, DAN.; DUT.; and GERM.; Polec, BOHEM.; Poleo, SPAN.; Puleggio, ITAL.; Poly, POL.; Poejo, PORT.; Poleja, SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; LUSIT.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 490*)

The herb (*Herba Pulegii*) is employed. It consists of a nearly cylindrical, pubescent, and very branching stem, bearing oval, obtuse, scarcely dentated leaves. Its odour is strongly penetrating; its taste is warm and is succeeded by a sensation of coolness.

All these plants are more or less stimulant; but the *M. piperita*, is the most active.—Dose, from a scruple to half a dram, of the powder.

CEPHALIC SPECIES.

Species cephalica seu *nervina externa*. (LIPP.)

R. Herb of <i>Mentha Crispa</i> ,	} of each, three ounces.
—— Marjoram,	
—— Origanum,	
—— Wild Thyme,	} of each, one ounce and a half.
Flowers of Chamomile,	
—— Lavender,	
Bay Berries,	
Juniper Berries,	
Caraway Seeds,	

Mix.

CONSERVE OF MINT.

Conserva Foliorum Menthae. (AMST.; ANT.; BATAV.; BELG.; FULD.; GENEV.; PALAT.; LUSIT.; ROSS.; SAX.; WIRTEM.; HERBIP.; SWEDIAUR.; VAN MONS)

R. Fresh Leaves of Mint, one part.
White Sugar, two parts.

Beat together, and preserve in a cool place. (BATAV.; BELG.; FULD.; LUSIT.; PALAT.; ROSS.; WIRTEM.; HERBIP.; SWEDIAUR.)

AMST.; and GENEV. prescribe one part of mint and three parts of sugar.—SARD.—an equal part of each.

R. Fresh Herb of Mint, one part.
Sugar, two parts.

Bruse the herb, triturate the pulp with the sugar and heat in a water-bath till this is melted. (VAN MONS)

R. Powdered Leaves of Mint, one dram.
White Sugar, five drams.
Water, two drams.

Mix. (ANT.)

These formulæ are for the *M. crispa*, *piperita*, and *viridis*.

AROMATIC APPLICATION. (CADET DE GASSICOURT)

R. Flowers of Chamomile, } of each,
Mentha Crispa, } three ounces.
Rue, }
Sweet Flag, }
Camphor, six drams.

Make a gross powder, sprinkle it with hot brandy, and apply it to the head.

INFUSION OF MINT.

Infusum Menthe. (DUBL.; MIL. COPEN.; FORM. HÔP. FR.; RATIER)

Infusum Menthe simplex. (DUBL.)

R. Dried Leaves of Spearmint, two drams.

Boiling Water,

a sufficient quantity

to obtain six ounces of strained infusion.

R. Dried Herb of Pepper- } of each,
mint or of Watermint, } half
Liquorice, } an ounce.
Boiling Water, a sufficient quantity
to obtain two pounds of strained infusion.
(FORM. HÔP. FR.)

R. Leaves of Mint, half an ounce.
Boiling Water, two pounds.

Infuse. (RATIER)

MIL. COPEN. prescribes one dram of mint and twelve ounces of water.

It is useful in cases where the stomach requires to be slightly stimulated.

COMPOUND INFUSION OF MINT. (DUBL.; COXE; NIEMANN)

Infusum Menthe compositum.
(DUBL.)

R. Dried Leaves of Spearmint, two drams.

Boiling Water,

a sufficient quantity

to obtain six ounces of strained infusion. Digest for half an hour in a covered vessel. To the cold infusion add,

Refined Sugar, two drams.

Oil of Spearmint, three drops.

Compound Tincture of

Cardamoms, half an ounce.

Mix intimately.

NIEMANN; and COXE give the same formula.

SYRUP OF MINT.

Syrupus Menthe seu Menthe crispe.
(AMST.; ANT.; BATAV.; BRUNS.; HISP.; GALL.; HANN.; PALAT.; SARD.; WIRTEM.)

R. Fresh Leaves of Mentha Crispa, two ounces.
Water, two pounds.

Infuse, strain, and add

White Sugar, four pounds.

Make a syrup, and strain. (HISP.; SARD.)

HANN. directs eight ounces of mint to be infused in three pounds of water, and four pounds of sugar to be dissolved in twenty ounces of the strained infusion.

R. Dried Leaves of Mentha Crispa, one ounce.
Distilled Water of Mentha Crispa, two pounds.

Digest for two hours in a water-bath, in a covered vessel; strain, filter, and add,

White Sugar, twice the quantity.

Dissolve. (GALL.)

BRUNS.; and WIRTEM. prescribe four ounces of mint, one pound of distilled water; two days' digestion in a water-bath, and sixteen ounces of sugar.

R. Juice of Mentha Crispa, } of each,
depurated by repose, } three
White Sugar, } pounds.

Boil together. (PALAT.)

SARD. prescribes simple syrup instead of sugar.

R. Dried Herb of Mint, half a pound.
Water, five pounds.

Macerate for twelve hours; then distil four ounces, and dissolve in them eight ounces of sugar. To the expressed and strained residue add,

White Sugar, two pounds four ounces.

Boil to the required consistence, and add the other syrup when cold. (AMST.)

R. Dried Flowering Tops of Mint, four ounces.
Water, three pounds.

Infuse for twelve hours in a covered vessel; express slightly, set aside, filter, and add,

White Sugar, four pounds.

Clarify, and evaporate to the proper consistence. To the cold syrup add another prepared without heat, with

Water of *Mentha Crispa*, one pound.

White Sugar, twenty ounces.

Preserve. (ANT.)

BATAV. directs four ounces of dried herb to be infused in four pounds of water for twenty-four hours; three ounces of liquor to be distilled, and six ounces of sugar to be dissolved in them; the residue to be then expressed, two pounds and a half of sugar added to the liquor, and boiled to a syrup: the preceding to be finally mixed with it when cold.

Dose, from half an ounce to two ounces.

SYRUP OF PEPPERMINT.

Syrupus Menthe piperitidis seu de Mentha piperita. (BATAV.; BELG.; GALL.)

R. Dried Tops of Peppermint, one ounce.
Distilled Water of Peppermint, two pounds.

Digest for twenty-four hours in a water-bath, in a covered vessel; strain, filter, and add,

White Sugar, twice the quantity.

Dissolve in a water-bath, and strain through flannel when cold. (GALL.)

HENRI proposes to mix one part of infusion with three parts of simple syrup previously boiled *au boulet*.

R. Tops of Peppermint, four ounces.
Water, one pound and a half.

Digest for three days in a covered vessel; express, depurate by repose, strain, and add Peppermint Water,

one pound and a half.

White Sugar, three pounds.

Boil slowly to a syrup. (BELG.)

R. Dried Herb of Peppermint, four ounces.
Water, four pounds.

Digest for twenty-four hours in a covered vessel; then distil about three ounces of liquid, with which and

Sugar, two pounds and a half.

Make a syrup, and, when it is cold, mix it with the preceding. (BATAV.)

VINOUS SYRUP OF PEPPERMINT. (VAN MONS)

R. Vinous Peppermint Water, five parts.

White Sugar, nine parts.
Dissolve in a covered vessel with the heat of a water-bath.

TINCTURE OF PEPPERMINT.

Tinctura seu Essentia Menthe piperitæ. (AMER.; ANT.; BRUNS.; COXE)

R. Herb of Peppermint, one part.
Alcohol (20 degrees), eight parts.

Digest for three days, and filter. (ANT.)

R. Flowering Tops of Peppermint, four ounces.
Rectified Spirit of Wine, two pounds.

Digest, express, and pour the tincture upon Flowering Herb of Peppermint, two ounces.

Digest again, express, and filter. (BRUNS.)

R. Oil of Peppermint, two drams.
Alcohol, one pound.

Mix. (AMER.; COXE)

It is excitant.—Dose, one or two drams.

TINCTURE OF MENTHA CRISPA. (ANT.; VAN MONS)

R. Herb of *Mentha Crispa*, one part.
Alcohol (20 degrees), eight parts.

Digest for three days, and filter.

TINCTURE OF GREEN MINT.

Tinctura Menthe viridis. (AMER.; COXE)

R. Oil of Green mint, two drams.
Alcohol, one pound.

Mix.

OIL OF MINT BY INFUSION. (BRUNS.; OLDEN.; SAX.; SARD.; WIRTEM.)

R. Herb of Mint, one part.
Olive Oil, eight parts.

Digest, and express. (SAX.)

R. Herb of Mint, one part.
Olive Oil, three parts.

Digest for four days in a warm place, and express; repeat the operation once or twice with fresh herb and filter. (OLDEN.)

SARD. prescribes one part of herb and two parts of oil.

R. Herb of Mint, } of each, one part.
Olive Oil, }

Boil to the consumption of the humidity, and express. (BRUNS.; WIRTEM.)

It is stomachic, and carminative.

ESSENTIAL OIL OF MINT.

R. Fresh Herb of Mint, any quantity.

Spring Water,
a sufficient quantity
to prevent empyreuma. Distil, and
collect the supernatant oil. (LOND.;
DUBL.; EDINB.)

The same process is required by AMER.;
AMST.; BATAV.; BELG.; FENN.; FERR.;
GENEV.; LUSIT.; ROSS.; SUEC.; BRUG-
NATELLI; and COXE.

FULD.; and LIPP. prescribe one part of
herb and three parts of water.—ANT.—
three of herb and ten of water.—GALL.—
five of herb and seven of water.—DAN.—
one of herb and six of water.—HISP.;
HANN.; OLDEN.; POL.; BORUS.; and
SAX.—one of herb and eight of water.—
BAVAR.—one of herb and nine of water.—
AUSTR.—one of herb and twelve of water.
—SWEDIAUR.—one of herb and four of
water.

R. Herb of Mint, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Distil. (BRUNS.; PALAT.; SAX.;
WIRTEM.; PIDERIT)

R. Herb of Mint, any quantity.
Distilled Water of Mint,
a sufficient quantity.

Distil. (HERBIP.)

There are required to be prepared, an
Oleum Menthe Sativæ, by DUBL.; an *Ol.*
M. viridis, by LOND.; AMER.; GENEV.;
COXE; and SWEDIAUR; an *Ol. M. crispæ*,
by AMST.; ANT.; AUSTR.; BATAV.; BA-
VAR.; DAN.; FULD.; HANN.; OLDEN.;
LUSIT.; POL.; BORUS.; ROSS.; SAX.;
WIRTEM.; HERBIP.; PIDERIT; and SWE-
DIAUR; an *Ol. M. piperitæ*, by LOND.;
EDINB.; DUBL.; AMER.; AMST.; ANT.;
AUSTR.; BATAV.; BAVAR.; BELG.; DAN.;
GALL.; FENN.; FULD.; GENEV.; HANN.;
HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.;
POL.; BORUS.; SARD.; SAX.; WIRTEM.;
HERBIP.; BRUGNATELLI; COXE; PIDE-
RIT; and SWEDIAUR; and lastly, an *Ol.*
Pulegii, by LOND.; DUBL.; AMER.;
BRUNS.; HANN.; PALAT.; SARD.; WIR-
TEM.; HERBIP.; and COXE.

Dose, from two to six drops.

OLEOSACCHARUM OF PEPPERMINT. (AUSTR.;
DAN.; HANN.; OLDEN.; CAST. BORUS.;
BORUS.; WIRTEM.; PLANCHE)

R. White Sugar, one ounce.
Essential Oil of Peppermint,
eight drops.

Triturate together. (DAN.; PLANCHE)

HANN.; OLDEN.; CAST. BORUS.; and
BORUS. prescribe half an ounce of sugar,
and twelve drops of oil.—AUSTR.—one
dram of sugar and three grains of oil.—

WIRTEM.—one ounce of sugar and sixteen
drops of oil.

TROCHISCS OF PEPPERMINT.

*Rotulæ seu Morsuli seu Trochisci Menthe
Piperitæ.* (AUSTR.; BAVAR.; GALL.;
FULD.; HANN.; HASS.; LIPP.; OL-
DEN.; SAX.; WIRTEM.; NIEMANN)

R. White Sugar, } of each,
Mint Water, } two ounces.

Boil to the consistence of a soft elec-
tuary, and add to it an oleosaccharum pre-
pared with

White Sugar, four ounces.
Essential Oil of Mint, half a dram.

Pour the mixture by drops upon a mar-
ble slab, and when the drops have solidified,
expose them, placed on a sieve, to a mode-
rate heat for several hours. (GALL.)

LIPP.; and WIRTEM. prescribe fourteen
drams and a half of sugar, half an ounce
of mint water, and an oleosaccharum of one
dram and a half of sugar and ten drops of
oil.—SAX.—an arbitrary quantity of sugar,
about one fourth of mint water, and such a
quantity of oleosaccharum that a total mass
of two pounds may contain about two
drams of essential oil.—OLDEN.—half a
pound of sugar, a sufficient quantity of
mint water, and an oleosaccharum of half
a dram of sugar and thirty drops of oil.—
AUSTR.—one ounce of sugar, a sufficient
quantity of mint water, and twenty-four
grains of essential oil.

R. White Sugar, half a pound.
Distilled Water, one ounce.

Heat till a drop solidifies on being placed
upon a cold body; then add to the mass
when half cold,

Essential Oil of Mint, half a dram.

Make trochiscs. (HASS.)

HANN. prescribes four ounces of sugar,
a sufficient quantity of water, and thirty-six
drops of oil.—BAVAR.—an equal part each,
of sugar and of oleosaccharum, to be boiled
together, with a little mint water.

R. Sugar, boiled *à la plume*, six ounces.
Powdered Herb of Peppermint,
six drams.

Mix. Add to the mass when half cold,

Oil of Peppermint, thirty drops.

Make thirty trochiscs. (FULD.)

R. Sugar, one ounce.
Oil of Peppermint, six drops.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs. (FULD.)

FERR. prescribes half a dram of oil, six
ounces of sugar, and a sufficient quantity of
mucilage made with mint water.

R. Isinglass, one dram.
Water, three ounces.

Dissolve, and add,
Oleosaccharum of Peppermint,
one pound.

Make trochiscs. (NIEMANN)

DISTILLED WATER OF MENTHA CRISPA.
(AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; MIL. COPEN.; FENN.; FERR.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; HERBIP.; BRUGNATELLI; PIDERIT; SWEDIAUR; VAN MONS)

R. Fresh Herb of Mentha Crispa, one part.
Water, a sufficient quantity.

Digest for twenty-four hours, and distil ten parts. (AMST.; BATAV.; BELG.; SWEDIAUR)

DAN.; OLDEN.; POL.; and BORUS. prescribe one part of herb, and a sufficient quantity of water; ten parts to be distilled.—HANN.—eight of herb, and a sufficient quantity of water; twenty-four to be distilled.—FENN.; and SUEC.—four of herb, and a sufficient quantity of water; six to be distilled.—LUSIT.—six of herb, and a sufficient quantity of water; eight to be distilled.—FERR.; and PALAT.—one of herb, and four of water; two to be distilled.—ANT.—one of herb, and five of water; four to be distilled.—HERBIP.; and BRUGNATELLI—one of herb, and six of water; two to be distilled.—HASS.; ROSS.; and PIDERIT—one of herb, and eight of water; four to be distilled.—AUSTR.—one of herb, and twelve of water; four to be distilled.—SAX.—one of herb, and fifteen of water; one to be distilled.—BAVAR.—one of herb, and sixteen of water; four to be distilled.

R. Herb of Mentha Crispa, one part.
Water, four parts.

Distil one half, and re-distil with fresh herb.

R. Fresh Herb of Mentha Crispa, ten parts.
Water, one hundred and sixty parts.
Proof Spirit, one half part.

Distil forty parts. (VAN MONS)

BRUNS. prescribes one part of herb, and four parts of white wine; one part and a half to be distilled.

DISTILLED WATER OF PEPPERMINT.
(LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FULD.; GENEV.; HANN.;

HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; HERBIP.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

R. Herb of Peppermint, one pound and a half.
Water, a sufficient quantity

to prevent empyreuma Distil one gallon. (LOND.; DUBL.; AMER.)

EDINB. prescribes three pounds of the herb, and ten pounds to be distilled.

AMST.; BATAV.; BELG.; DAN.; and SWEDIAUR prescribe one part of fresh herb, and a sufficient quantity of water; to be macerated for twenty-four hours, and ten parts to be distilled. HANN.; OLDEN.; POL.; and BORUS.—one of herb, and a sufficient quantity of water; seven to be distilled.—GENEV.—one of herb, and a sufficient quantity of water; six to be distilled.—FENN.; and SUEC.—two of herb, and a sufficient quantity of water; three to be distilled.—LUSIT.—three of herb, and a sufficient quantity of water; four to be distilled.—GALL.—one of herb, and four of water; two to be distilled.—ANT.—one of herb, and five of water; four to be distilled.—AUSTR.; BRUNS.; HERBIP.; and PIDERIT—one of herb, and six of water; two to be distilled.—HASS.; and ROSS.—one of herb, and eight of water; four to be distilled.—SAX.—one of herb, and fifteen of water; ten to be distilled.—BAVAR.—one of herb, and sixteen of water; eight to be distilled.

R. Fresh Herb of Peppermint, three parts.

Enclose it in a bag suspended in the middle of a cucurbit containing

Water, twelve parts,
in such a manner that it do not touch the liquid, and distil six parts. (FULD.; LIPP.)

R. Herb of Peppermint, ten parts.
Proof Spirit, one half part.
Water, one hundred and sixty parts.

Distil forty parts. (VAN MONS)

R. Oleosaccharum of Peppermint, one dram.
Distilled Water, eight ounces.

Triturate to dissolve. (CAST. BORUS.)

R. Essential Oil of Mint, one part.
Sugar, } of each,
Alcohol (22 degrees), } three parts.
Water, twelve hundred parts.

Mix. (GALL.)

Dose, from one ounce to four ounces.

DISTILLED WATER OF SPEAR MINT.
(LOND.; AMER.; NIEMANN; SWEDIAUR)

Aqua Mentha viridis. (LOND.)

R. Herb of Spear Mint,
 one pound and a half.
Water, a sufficient quantity
to prevent empyreuma. Distil one
gallon.

AMER.; and NIEMANN prescribe three
pounds of herb, and a sufficient quantity of
water; ten pounds to be distilled.—SWE-
DIAUR—one part of herb, and a sufficient
quantity of water; ten parts to be distilled.

DISTILLED WATER OF PENNYROYAL MINT.

Aqua Pulegii. (LOND.; EDINB.; AMST.;
AUSTR.; FERR.; FULD.; GENEV.;
HANN.; HASS.; PALAT.; SARD.; SAX.;
HERBIP.; PIDERIT; SWEDIAUR; VAN
MONS)

R. Herb of Pennyroyal,
 one pound and a half.
Water, a sufficient quantity
to prevent empyreuma. Distil one
gallon. (LOND.)

EDINB. prescribes three pounds of
herb, and the distillation of ten
pounds of water.

HANN. prescribes two parts of herb, and
a sufficient quantity of water; fourteen
parts to be distilled.—AMST.; and SWE-
DIAUR—one of herb, and a sufficient quan-
tity of water; ten to be distilled.—PALAT.—
one of herb, and four of water; two to be
distilled.—HERBIP.—one of herb, and six
of water.—HASS.; and PIDERIT—one of
herb, and eight of water; four to be dis-
tilled.—AUSTR.—one of herb, and twelve of
water; four to be distilled.—SAX.—one of
herb, and fifteen of water; two-thirds to be
distilled.—GENEV.—one pound of herb,
and a sufficient quantity of water; six pounds
to be distilled.—FERR.—four pounds of
herb, and sixteen of water; eight to be
distilled.

R. Fresh Herb of Pennyroyal, one part.
Water, four parts.
Distil one half, and re-distil with fresh
herb.

R. Fresh Herb of Pennyroyal, three parts.
Suspend it, in a bag, in the middle of a
cucurbit containing

Water, twelve parts,
so that the bag may not touch the water.
Distil six parts. (FULD.)

R. Fresh Herb of Pennyroyal, ten parts.
Proof Spirit, one half part.
Water, one hundred and sixty parts.
Distil forty parts. (VAN MONS)

SPIRIT OF MINT.

*Spiritus Menthae, Aqua Menthae spirituosae
seu vinosa seu alcoholica.*

R. Oil of Mint, by weight,
 six scruples and a half.
Proof Spirit,
 four pints and a half.
Water, a sufficient quantity
to prevent empyreuma. Mix the spi-
rit with the oil; then add the water,
and distil slowly one gallon. (LOND.;
DUBL.)

R. Fresh Herb of Mint, one part.
Alcohol (20 degrees), four parts.
Distil a spirit marking 28 degrees. (ANT.)

BELG. prescribes one part of herb and six
parts of alcohol; a spirit marking 16 de-
grees to be distilled.—BRUNS.; PALAT.;
and WIRTEM.—one of herb and four of al-
cohol; two to be distilled.—AMST.—two of
herb and nine of alcohol; to be distilled.
—LUSIT.—one and a half of herb and ten
of alcohol; five to be distilled.—SWEDIAUR
—one of herb and fourteen of alcohol; fif-
teen to be distilled.—BRUNS.—one pound
ten ounces of herb and eight pounds of
alcohol.—FERR.—one part and a half of
herb and eight parts of alcohol; six to be
distilled.

R. Fresh Herb of Mint, one part.
Alcohol (0.910), four parts and a half.
Water, six parts.

Macerate for twenty-four hours, and distil
four parts. (ANT.)

COXE prescribes one part and a half of
herb, nine parts of proof spirit, and a suffi-
cient quantity of water; nine parts to be
distilled.—VAN MONS—one of herb, three
of alcohol, and six of water; two and a half
to be distilled.—HANN.; OLDEN.; POL.;
and BORUS.—one of herb, one and a half
of alcohol, and a sufficient quantity of
water; six to be distilled.—BAVAR.—one
of herb, two and a half of alcohol, and six-
teen of water; eight to be distilled.

R. Fresh Herb of Mint, one part.
Alcohol, four parts.

Digest for twenty-four hours, and then
distil to dryness. Macerate with fresh herb,
and distil a second time. (SARD.)

HERBIP. prescribes one pound and a half
of herb, ten pounds of alcohol, and the
product to be re-distilled from three ounces

of fresh herb.—SWEDIAUR—twelve parts of herb, six of alcohol, and two of water; six parts to be distilled, and the product to be re-distilled with two parts of water and twelve of fresh herb.

R. Fresh Herb of Mint, two pounds.
Alcohol, } of each, an equal part,
Water, }

and in sufficient quantity to obtain eight pounds of product. Add to this,

Water, }
White Sugar, } of each, two ounces.

Preserve. (HASS.; PIDERIT)

R. Flowering Herb of Mint, twelve parts.
Alcohol, thirty-six parts.
Distilled Water of Mint, twelve parts.

Infuse for four days, and distil thirty parts in a water-bath. (GALL.)

ROSS. prescribes one half part of herb, four parts of alcohol, and one part of mint water; to be infused for twenty-four hours, and three parts to be distilled.

R. Mint Water, one pound.
Alcohol, one ounce.

Mix. (BRUGNATELLI)

LOND.; COXE; and SWEDIAUR require a *Spiritus Menthe viridis*; AMST.; ANT.; AUSTR.; BATAV.; BRUNS.; GALL.; LIPP.; LUSIT.; ROSS.; and SWEDIAUR, a *Sp. M. crispæ*; LOND.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BRUNS.; GALL.; HANN.; HASS.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SARD.; HERBIP.; WIRTEM.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR; and VAN MONS, a *Sp. M. Piperitæ*; and LOND.; SARD.; COXE; and VAN MONS, a *Sp. M. Pulegii*.

STOMACHIC LIQUOR.

Aqua Stomachica seu Menthe composita.
(WIRTEM.; SPIELMANN)

R. Fresh Herb of Mentha Crispa, three ounces.

Galanga, }
Nutmegs, } of each, one ounce.
Cinnamon, }
Wine, eight pounds.

Distil one half. (WIRTEM.)

R. Fresh Herb of Mint, twenty-four ounces.

——— Wormwood, three ounces.
——— Sweet Basil, } of each,
——— Pennyroyal, } two
——— Rosemary, } of each,
Flowers of Lavender, } two drams.

Cinnamon, half an ounce.
Cloves, }
Coriander, } of each, one dram.
Alcohol, sixteen ounces.
Mint Water, sixty ounces.

Distil nearly to dryness. (SPIELMANN)

Dose, from half an ounce to one ounce.

SACCHARINE SPIRIT OF MINT. (SAUNDERS; SAINTE-MARIE; SWEDIAUR; VAN MONS)

R. Spirit of Peppermint, seventy-four ounces.

Simple Syrup, twelve ounces.
Tincture of Peppermint, three drams.

Filter the mixture at the end of several days. (VAN MONS)

R. Mint Water, six ounces.
Spirit of Peppermint, } of each,
Syrup of Peppermint, } one ounce.
Oil of Mint, six drops.
Sugar, half an ounce.

Mix. (SAINTE-MARIE)

R. Mint Water, eight ounces.
Spirit of Mint, one ounce and a half.
Simple Syrup, half an ounce.

Mix. (SAUNDERS)

SWEDIAUR prescribes eight ounces of spear mint water, one ounce of tincture of mint, and one ounce of syrup of mint.

To be taken by spoonfuls.

STOMACHIC POTION. (SAINTE-MARIE)

R. Distilled Water of Mentha Crispa, four ounces.
——— Pepper- }
mint, } of each,
Hordeated Cinnamon Water, } one
Syrup of Poppies, } ounce.

Mix.

ANTI-SPASMODIC POTION.

Aqua ad tormina infantum. (SAINTE-MARIE; SPIELMANN)

R. Mint Water, } of each,
Chamomile Water, } two ounces.
Syrup of Red Poppies, } of each,
——— Chamomile, } half an ounce.

Mix. (SPIELMANN)

R. Oil of Cinnamon, two drops.
——— Peppermint, three drops
Spirit of Peppermint, one ounce.
Syrup of Marshmallows, three ounces.
Peppermint Water, four ounces.

Mix. (SAINTE-MARIE)

To be taken by spoonfuls.

MERCURIALIS ANNUA, LINN.—ANNUAL MERCURY.

SYNONIMES:—Mercuriale annuelle, Jähriges Bingelkraut, GERM.; Mercurialis, neb pluna bazalicka, BOHEM.; Bingelurt, DAN.; Jaarlyks bingelkruid, DUT.; Mercorella, ITAL.; Mercuryasez, POL.; Bin-gelært, SWED.

AMST.; ANT.; BELG.; BRUNS.; HISP.; GALL.; GENEV.; HERBIP.; WIRTEM.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

This plant is common in Europe. (*Diœcia Enneandria*, LINN.; *Euphorbiaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 103, 104*)

The herb (*Herba Mercurialis*) is employed. It consists of a branching, glabrous stem, bearing oval, lanceolated leaves, dentated on the borders, and of a pale-green colour. It is destitute of odour, and has a mucilaginous and rather saline taste.

It contains, according to FENEULLE, a bitter, slightly purgative principle, chlorophylle, albumen, a fatty matter, a volatile oil, pectic acid, salts, &c.

PURGATIVE LAVEMENT. (RATIER)

R. Herb of Annual Mercury, half an ounce.
Water, one pound.
Boil, and strain.

DECOCTION OF ANNUAL MERCURY.
(SARD.)

R. Senna, six drams.
Annual Mercury, one handful and a half.
Water, a sufficient quantity
to obtain one pound and a half of strained
decoction. Boil for half an hour, and add
Pectoral Honey, two ounces.
Mix.

HONEY OF ANNUAL MERCURY.

Mel mercuriale seu mercurialis annuæ.

(ANT.; GALL.; PALAT.; SARD.; WIRTEM.; SPIELMANN.; VAN MONS.)

R. Juice of Annual Mercury, } of each, an
Honey, } equal part.

Boil, removing the scum, to the consistence of syrup, and strain through flannel. (GALL.; SARD.)

ANT.; PALAT.; and WIRTEM. prescribe two parts of honey and three of juice.—SPIELMANN—three of juice and two of honey.—VAN MONS—one of juice and one and a half of honey.

It used to be employed in lavements.—Dose, one or two ounces, in emollient decoction.

COMPOUND HONEY OF ANNUAL MERCURY.

Syrupus mercurialis seu prophylacticus seu longæ vitæ. (GALL.; WIRTEM.; SPIELMANN.)

R. Juice of Annual Mercury, four pounds.
—— Borage, } of each,
—— Bugloss, } one pound.
Honey, six pounds.

Boil, removing the scum; then

R. Gentian, two ounces.
Root of Iris, three ounces.
White Wine, one pound and a half.

Macerate for twenty-four hours; then express; mix this with the preceding liquor, boil slowly to the consistence of syrup, and strain.

SPIELMANN gives the same process, but prescribes eighty-six ounces of the juice of annual mercury, twenty-four of borage juice, twenty-four of bugloss juice, thirty-two of honey, four of gentian, eight of iris, and forty-eight of wine.—GALL.—four pounds of the juice of annual mercury, one pound each, of borage juice, and of bugloss juice, six pounds of honey, two ounces of gentian, four of iris, and two pounds of wine.

It is purgative, stomachic, and emmenagogue.

MESEMBRYANTHEMUM CRYSTALLINUM, LINN.—ICEPLANT,
DIAMOND FIG MARIGOLD.

SYNONIMES:—Glaciale, Cristalline, FR.; Eiskraut, Mittagsblume, Eispflanze, GERM.; Isplant, DAN.; Hierba de la plata, SPAN.; Yskruid, DUT.; Erba cristallina, ITAL.; Agclada, PORT.; Isært, SWED.

VOL. II.

ANT.; BATAV.; BAVAR.; FULD.; MURRAY and GMELIN.

A plant of the Archipelago. (*Icosandria Pentagynia*, LINN.; *Ficoides*, JUSS.; *Fig. Dill. Elth. 231, t. 180, f. 221*)

2 I

The herb is employed; it consists of a branching stem, and of small, undulated, oval, opposite or alternate leaves; the whole plant is covered with brilliant crystalline vesicles. It has a rapid and rather nauseous taste.

It is sedative and resolvent; and has been recommended in hooping cough and dysury. It has been extolled as a specific diuretic.—Dose, a tea spoonful every two hours, of the juice.

CONSERVE OF ICEPLANT.

Conserva Mesembryanthemi crystallini.
(FULD.)

R. Fresh Herb of Iceplant, one part.

White Sugar, two parts.

Mix.

LOHOCH OF ICEPLANT. (NIEMANN; AUGUSTIN)

R. Juice of Iceplant, } of each,
Syrup of Oak Lichen, } one ounce.

Mix.—Dose, half a spoonful, six times a day, in hooping cough.

SYRUP OF ICEPLANT. (BAVAR.)

R. Fresh Juice of Iceplant, depurated
by boiling, one part.
Sugar, one part and a half.

Make a syrup.

MESPILUS.

Two plants of this name are used in medicine:

1. *Mespilus Oxyacantha*, CAND. — Hawthorn.

SYNONIMES:—Aubepine, Epine blanche, FR.; Weissdorn, GERM.; Hagetorn, DAN.; Espina majuolo, SPAN.; Hangdorn, DUT.; Bianco spino, ITAL.; Böldak, POL.; Espinheiró alvar, PORT.; Hagtorn, SWED.

HISP.; GALL.; SPIELMANN.

A common shrub of Europe. (*Icosandria Digynia*, LINN.; *Rosaceæ*, JUSS.; *Fig. Austr. III. t. 292, f.*)

The flowers (*Flores Oxyacanthæ seu Spinæ albæ*) are employed. They are white, large, and disposed in bunches. They have an agreeable odour, which becomes almost entirely dissipated by drying. Their taste is feebly styptic.

Infused in warm water, they form a domestic remedy for cough. No loss would be endured by their being banished from the materia medica.

2. *Mespilus Germanica*, LINN. — Dutch Medlar.

SYNONIMES:—Neflier d'Allemagne, FR.; Mispelbaum, GERM.; Mispeltræ, DAN.; Mispero, SPAN.; Mispelboom, DUT.; Niesplik, POL.; Nespereira, PORT.; Mesp-telträd, SWED.

AMST.; HISP.; GALL.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A tree originally of the mountains of Europe. (*Icosandria Pentagynia*, LINN.; *Rosaceæ*, JUSS.; *Fig. Nouv. Duh. IV. 38*)

The fruit and seeds are employed.

The fruit *Medlar* (*Fructus Mespili*), is round, depressed, fleshy, umbilicated at the summit, comprising several cells, each of which contains a seed. Its taste is acerb and astringent, but becomes, by ripening, sweet, vinous, and agreeable.

The seed is oblong; osseous, and very hard.

The fruit is astringent, and the seed is reputed diuretic.

MOMORDICA.

Two species of this genus of plants are employed in medicine:

1. *Momordica balsamina*, LINN. — Balsapple.

SYNONIMES:—Balsamine, Pomme de mer-

veille, FR.; Balsamapfel, GERM.; Balsamæple, DAN.; Momordica, ITAL.; and SPAN.; Balsam appel, DUT.

BRUNS.; GALL.; FERR.; BRUGNATELLI; SPIELMANN.

A plant of the East Indies. (*Monæcia*

Monadelphia, LINN.; *Cucurbitaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 45*)

The fruit (*Fructus Momordicæ* seu *Pomum Hierosolymitanum*) is employed. It is a sort of berry, of about the size of a pigeon's egg, and of an orange or scarlet colour.

It used formerly to be reputed vulnerary.

2. *Momordica Elaterium*.—LINN.—*Elatarium*, Squirting Cucumber.

SYNONIMES:—*Elatarium*, Concombre sauvage, FR.; *Eselskürbis*, *Eselsgurke*, Springgurke, GERM.; *Strikawa anel plana tykvice*, BOHEM.; *Cohombrillo*, amargo, SPAN.; *Ezelskommers*, DUT.

LOND.; EDINB.; DUBL.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; SAX.; SUEC.; WIRTEM.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; SAUNDERS.; SPIELMANN.; ZARDA.

A plant of the middle of Europe. (*Fig. Zorn, Ic. pl. t. 144*)

The root and the fruit are employed.

The root (*Radix Cucumeris agrestis* seu *Asini* seu *Momordicæ Elaterii*) is about two or three inches in thickness, and one foot in length; it has a disagreeable odour, and a repugnant and burning taste.

The fruit is an oval, oblong berry, of a green colour; it is gathered when only half ripe.

This fruit contains, according to PARIS, a peculiar principle, *Elatin*, *Elatarin*, *Elatarium*, which is the source of its activity.

It acts as a very violent drastic purgative.

It used formerly to be much employed in chronic dropsies, but is scarcely prescribed at the present day.

JUICE OF ELATERIUM.

Succus Elaterii. (HISP.; SARD.)

R. Fruit of *Elatarium* not yet ripe, any quantity.

Having removed the seeds, bruise the remainder in a stone mortar, then express the juice, set it aside, decant, and preserve the limpid part.

EXTRACT OF ELATERIUM.

Elatarium, *Inspissamentum Cucumeris agrestis*, *Extractum Fructus Momordicæ Elaterii*. (LOND.; DUBL.; BRUNS.; HISP.; HANN.; SARD.; HERBIP.; WIRTEM.; SPIELMANN.; SWEDIAUR.; VAN MONS)

Extractum Elaterii.

Slice ripe fruit of *elaterium*, ex-

press the juice very gently, and pass it through a very fine hair-sieve into a glass vessel; then set it apart for some hours, till the thicker part shall have subsided. Reject the supernatant thinner portion, and dry the other with a gentle heat. (LOND.)

Slice ripe fruit of *elaterium*, and strain the juice very lightly, letting it pass through a very fine hair-sieve into a glass vessel; then set it aside for some hours till the thicker part shall have subsided. Reject the supernatant liquor, and dry the fecula, placed in a folded linen cloth, with a moderate heat. (DUBL.)

R. Ripe Fruit of *Elatarium*, any quantity.

Cut round the insertion of the footstalks with the point of a knife, and then take out the seeds; bruise the pericarps in a stone mortar with a wooden pestle, adding a little water: strain the pulp through a sieve, and evaporate it over a slow fire to the consistence of an extract. (GALL.; SARD.; WIRTEM.; HERBIP.; SPIELMANN.; SWEDIAUR.; VAN MONS)

BRUNS. directs the deposition of the fecula to be hastened by digesting the expressed juice in a water-bath.

This used generally to be prescribed in combination with other hydragogues, in dropsies.—Dose, three or four grains.

FECULA OF ELATERIUM.

Fæcula Cucumeris agrestis. (GENEV.; VAN MONS)

R. Green Fruit of *Elatarium*, any quantity.

Bruise it in a stone mortar, with a wooden pestle, and express the juice; bruise the residue with a little water, and express again. Mix the two liquors, set aside, decant, and coagulate the fecula over the fire; then remove the vessel from the fire, strain, wash well the fecula, and, having dried it with a gentle heat, reduce it to powder.

HYDRAGOGUE PILLS. (SAUNDERS)

R. Extract of *Elatarium*, ten grains.
Gentian, three drams.
Make pills of four grains each.

COMPOUND PILLS OF ELATERIUM. (SUEC.)

R. Gum Ammoniac, two ounces.
Aloes, } of each, two drams:
Gamboge, }

Extract of Elaterium, half a dram.
Tincture of Wormwood,
a sufficient quantity.
Make six hundred and fifteen pills.

OIL OF ELATERIUM.

Oleum Momordicæ. (BRUNS.; SARD.;
WIRTEM.; HERBIP.)

R. Ripe Fruit of Elaterium, one part.
Olive Oil, two parts.

Digest for two days in a water-bath, and express; then having let it cool, repeat the maceration. (SARD.)

BRUNS.; and WIRTEM. direct one part of the fruit and one part and a half of linseed oil to be digested in hot ashes; the mixture to be then boiled slowly for an hour to the consumption of the humidity; and the digestion and boiling to be repeated with a similar quantity of fresh fruit.—HERBIP.—one part of fruit and two parts of olive oil to be boiled slowly together to the consumption of the humidity.

It is recommended to be applied to chapped nipples, to chilblains, and to painful piles.

COMPOUND OIL OF ELATERIUM.

Oleum Conyzæ compositum. (HISP.)

R. Leaves of Erigeron viscosum, }
——— Spurge Laurel, } of each,
——— Elaterium, } four
——— Calamint, } ounces.
——— Rosemary, }

Leaves of Marjoram,	} of each, four ounces.
——— Thyme,	
——— Rue,	
——— Bay,	
——— Savine,	
——— Sage,	} of each, two ounces.
Root of Bryony,	
——— Arum,	
——— Elaterium,	} ten pounds.
Olive Oil,	

Macerate for four days; then boil slowly to the consumption of the humidity, and express.

HYDRAGOGUE MIXTURE.

Solutio Elaterii. (AUGUSTIN; COXE;
HUFELAND)

R. Extract of Elaterium, one grain.
Nitric Æther, two ounces.
Tincture of Squill, } of each,
Oxymel of Colchicum, } half an ounce.
Simple Syrup, one ounce.

Mix.—Dose, a spoonful, four or five times a day, in a little water. (AUGUSTIN; ELLIS)

R. Extract of Elaterium, one grain.
Parsley Water, six ounces.
Nitric Æther, two drams.
Oxymel of Squill, } of each,
Syrup of Buckthorn, } half an ounce.
Rob of Juniper, one ounce.
Compound Tincture of Cinnamon, one dram.

Mix.—Dose, two spoonfuls every two or three hours. (HUFELAND)

MONARDA PUNCTATA, LINN.—AMERICAN MOUNTAIN BALM.

SYNONIME:—Monarde ponctuée, FR.

AMER.; COXE.

A plant of North America. (*Diandria Monogynia*, LINN.; *Labiata*, JUSS.; *Fig. Pluk. Alm. tab. 24. fig. 1*)

The herb is employed; it consists of a whitish, downy stem, and of acute, narrow leaves covered with numerous brilliant hairs.

It is probable that it acts as an excitant.

OIL OF AMERICAN MOUNTAIN BALM.

Oleum Monardæ æthereum. (AMER.;
COXE)

R. Herb of American Mountain Balm, any quantity.
Water, a sufficient quantity
to cover the herb. Distil.

MORINGA OLEIFERA, LMK.—BEN; BEHN.

SYNONIME:—Ben ollifère, FR.

BRUNS.; GALL.; FULD.; LIPP.; WIRTEM.;
GUIBOURT; SPIELMANN.

A tree of Malabar. (*Hexandria Monogynia*, LINN.; *Leguminosæ*, JUSS.; *Fig. Flore Medic. III. 63*)

The root and seeds are employed.

The root (*Radix Moringæ*) is in pieces of various thicknesses, of a yellowish colour and resinous fracture, and of an aromatic taste and smell.

It is doubtful whether it should be attributed to this vegetable.

It is anthelmintic, but scarcely ever used.

The seed is called *Ben Nut* (*Nux Ben seu Been*; *Balsamus myrepsica*, *Glans unguentaria*). It is an ovoid nut, furnished with three membranous wings, and contains under a hard cartilaginous cuticle, a whitish oleaginous kernel.

OIL OF BEN.

Oleum Been seu balatinum. (GALL.; FULD.; LIPP.; BRUGNATELLI)

R. Ben Nuts, blanched, any quantity.

Bruise them in a marble mortar with a wooden pestle; enclose the paste in a woollen bag, express without heat, and strain the oil. (GALL.; FULD.; BRUGNATELLI)

LIPP. directs the bag containing the paste to be suspended over boiling water before submitting it to the press.

It is emollient, rubefacient, anti-spasmodic, and vermifuge, but is rarely used.

MORPHINA; MORPHEUM.—MORPHINE.

GALL.; FERR.; NIEMANN.

1. *Sertuerner's Method.*

R. Extract of Opium, five hundred parts.
Water, a sufficient quantity,
that the solution may mark eight degrees by the areometer. Then add to it

Solution of Ammonia (22 degrees),
a sufficient quantity,

or so much that it may remain in slight excess. Separate the gelatinous precipitate which forms, and then collect on a filter the crystals which collect at the bottom; dry them, and dissolve them in sulphuric acid (66 degrees) warm and previously diluted with eight parts of water: pour solution of ammonia into the liquor, collect the precipitate which forms, and wash it with alcohol (22 degrees); then dissolve it in boiling alcohol (36 degrees), strain the solution, and finally, set it aside to crystallize.

2. *Hottot's Method.*

R. Opium, any quantity.

Macerate it at several times, and then concentrate the liquors, mixed together, to two degrees of the areometer, add ammonia to the liquor in very slight excess, and separate the glutinous precipitate which forms; then filter, and again add solution of ammonia in excess: collect the precipitate upon a close cloth, wash it with alcohol (20 degrees), dry it, reduce it to powder, and finally treat it with alcohol (36 degrees) boiling, mixed with a small proportion of animal carbon. (HENRI)

3. *Robiquet's Method.*

R. Pure Opium, sliced, three hundred parts.
Water, one thousand parts.

Macerate for five days, strain, and add

Pure Magnesia, fifteen parts.

Boil for ten minutes, and filter; wash the sediment with cold water, dry it, and macerate it at a moderate heat in

Alcohol (22 degrees),
a sufficient quantity.

Wash again what remains with cold alcohol of the same strength, and let the residue be finally boiled for a few minutes in alcohol (32 degrees). Strain the liquor, collect the crystals which form on cooling, dissolve them in boiling alcohol (36 degrees), and set aside to re-crystallize. (GALL.; FERR.; NIEMANN)

4. *Winkler's Method.*

R. Opium, three parts.
Vinegar, one part.
Distilled Water, eight parts.

After several days' digestion, separate the clear part, and wash the residue at several times with water acidulated with vinegar; then, having mixed these liquors together, add solution of ammonia to them; collect the precipitate upon a filter, and digest it for several days in alcoholized æther. Separate from the supernatant liquor the morphine, now freed from the matters which had been precipitated with it, dissolve it in boiling alcohol (36 degrees), and set the solution aside, that crystals may form.

Morphine is rarely employed, on account of its being scarcely soluble in water.

NARCOTIC POTION.

Potio narcotica. (BRERA)

R. Morphine, half a grain.

Distilled Water, one ounce.
Syrup of Poppies, one dram.
Mix.—To be taken at bed-time.

SEDATIVE INJECTION.

Injectio leniens. (BRERA)

R. Oil of Almonds, } of each,
— Lily, } half an ounce.
Morphine, two grains.

Mix.—It is employed to appease pains of the ear, and in gonorrhœa, and also to relieve the tenesmus sometimes accompanying piles.

R. Oil of Chamomile, } of each,
— Poppy Seeds, } one ounce.
Yolk of Egg, one.
Morphine, two grains.

Mix.—It is employed in ear-ache-otalgia.

SULPHATE OF MORPHINE. (FERR.; HENRI; MAGENDIE)

R. Pure Morphine, four ounces.

Dissolve it in alcohol, or mix it with distilled water, and add carefully

Sulphuric Acid, diluted with its weight of Water, a sufficient quantity to saturate the alkali. Filter, evaporate slowly, and set apart to crystallize.

Dose, from the fourth of a grain to one grain, in potion or pills.

SYRUP OF SULPHATE OF MORPHINE. (FERR.; FEE; MAGENDIE)

R. Sulphate of Morphine, four grains.

Dissolve it in a little water slightly acidulated with sulphuric acid, and add

Warm Simple Syrup, one pound.

Mix.—To be taken in the dose of a teaspoonful.

SEDATIVE PILLS.

Pilule deprimentes. (BRERA)

R. Sulphate of Morphine, one grain.
Olive Oil, a sufficient quantity
to dissolve the salt. Add
Ipecacuanha, three grains.
St. Ignatius' Bean, two grains.
Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make six pills.—Dose, one, every two hours.

SEDATIVE BOLUSES. (BRERA)

R. Sulphate of Morphine, two grains.
Oil of Almonds, a sufficient quantity
to dissolve the salt. Add

Sulphate of Iron, four grains.
Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make eight boluses.—Dose, one every two hours.

R. Sulphate of Morphine, one grain.
Ipecacuanha, three grains.
Oil of Almonds, twelve drops.
Extract of Aconite, six grains.
Liquorice Powder, } of each,
Honey, } a sufficient quantity.

Make six boluses.—Dose, one every three hours.

ACETATE OF MORPHINE.

Acetas Morphei seu Morphineæ. (FERR.; HENRI; MAGENDIE; NIEMANN)

R. Morphine, any quantity.

Mix it with distilled water, and add

Acetic Acid (10 degrees),
a sufficient quantity

to saturate the alkali. Evaporate the solution to dryness.—Dose, from the fourth of a grain to one grain.

The preparations which will be found in the article on opium, in the paragraph of extraction by acetic acid, ought, strictly speaking, to be placed here, for it is the acetate of morphine, more or less impure, which forms their base.

SOLUTION OF ACETATE OF MORPHINE. (ELLIS)

R. Acetate of Morphine, sixteen grains.
Water, six drams,
Acetic Acid, two drams.

Mix.—Dose, from six to twenty-four drops.

SEDATIVE DROPS. (FERR.; BORIES; CADET DE GASSICOURT; ELLIS; HENRI; MAGENDIE; RATIER)

R. Acetate of Morphine, sixteen grains.
Acetic Acid, four drops.
Distilled Water, one ounce.
Alcohol, one dram.

Mix.—Dose, from six to twelve drops, in a sedative potion.

ANTI-CATARRHAL BOLUS. (CADET DE GASSICOURT; PIERQUIN)

R. Cocoa Nut Oil, one dram.
Gum Tragacanth, }
Saffron, } of each, half a dram.
Honey, }

Anisated Balsam of Sulphur, } of each,
Extract of Liquorice, } eighteen
grains.

White Henbane, six grains.
Acetate of Morphine, two grains.

Make thirty-six boluses.—Dose, one, every two or three hours.

SEDATIVE PILLS. (BRERA; CADET DE GASSICOURT)

R. Acetate of Morphine, one grain.
Conserve of Orange Flowers, sixteen grains.

Make eight pills.—Dose, one or two, every six hours, in neuralgia and cancer. The dose may be gradually carried to three grains daily, carefully observing its effects. (CADET DE GASSICOURT)

R. Acetate of Morphine, one grain.
Digitalis, eight grains.
Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make six pills.—Dose, one, every two hours. (BRERA)

R. Acetate of Morphine, one grain.
Red Sulphuret of Antimony, two grains.
Extract of Aconite, four grains.
Liquorice Powder, } of each, a
Honey, } sufficient quantity.

Make eight pills.—Dose, one, every three hours. (BRERA)

EMMENAGOGUE PILLS.

Pilula leviter emmenagogæ deprimentes.
(BRERA)

R. Acetate of Morphine, one grain.
Olive Oil, ten drops.

Dissolve, and add

Ipecacuanha, two grains.
Extract of Henbane, four grains.
Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make six pills.—Dose, one, every three hours; in amenorrhœa from uterine irritability.

SEDATIVE BOLUSES. (BRERA)

R. Acetate of Morphine, one grain.
Olive Oil, ten drops.

Dissolve, and add

Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make six boluses.—Dose, one, every two hours.

SEDATIVE POTION.

Haustus sedans narcoticus. (BRERA)

R. Infusion of Chamomile, five ounces.
Extract of Virous Lettuce, ten grains.
Sedative Drops, twenty drops.
Syrup of Marshmallows, half an ounce.

Mix.—To be taken gradually.

SYRUP OF ACETATE OF MORPHINE. (FERR.; BORIES; CADET DE GASSICOURT; FEE; HENRI; MAGENDIE)

R. Simple Syrup, one pound.
Acetate of Morphine, four grains.

Dissolve the salt in half a dram of water acidulated with a few drops of acetic acid; filter, and add the syrup warm.

To be taken by tea spoonfuls.

SEDATIVE MIXTURE.

R. Syrup of Acetate of Morphine, one ounce.

Infusion of Ground Ivy, four ounces.

Mix.

SEDATIVE EMULSION.

Emulsio sedativa simplex. (BRERA)

R. Almond Emulsion, eight ounces.
Syrup of Acetate of Morphine, one ounce.

Mix.—To be taken gradually.

CITRATE OF MORPHINE. (MAGENDIE; PARIS)

R. Pure Morphine, sixteen grains.
Crystallized Citric Acid, eight grains.

Dissolve both in

Distilled Water, one ounce.

Colour the solution with

Tincture of Cochineal, two drams.

Mix. (MAGENDIE)

PORTER prepared this salt by triturating four ounces of crude opium with two of crystallized citric acid, adding sixteen of water, and filtering the liquor after twenty-four hours' maceration. (PARIS)

This liquor has the same uses as the solution of the acetate of morphine.

MORUS NIGRA, LINN.—MULBERRY TREE.

SYNONIMES :—Mûrier noir, FR.; Maulbeerbaum, GERM.; Marusse, BOHEM.; Morbærtræ, DAN.; Moras, SPAN.; Mørbezieboom, DUT.; Gelso, ITAL.; Amoreira, PORT.; Mulbærstræd, SWED.

LOND.; AMST.; ANT.; AUSTR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A tree said to be originally of Persia, and cultivated in Europe. (*Montecia Tetrandria*, LINN.; *Urticææ*, JUSS.; *Fig. Nouv. Duh. IV. t. 22*)

The fruit, mulberries (*Baccæ Mori*), is a species of berry, produced by the agglomeration and union of the calices of the female catkins, become succulent and fleshy, and each containing one seed. They are oval, oblong, tolerably large, of a blackish purple colour, and of a sweet taste.

JUICE OF MULBERRIES.

Succus Mororum. (BRUNS.; HISP.; GALL.; PALAT.; SAX.; SUEC.; WIRTEM.)

R. Ripe Mulberries, any quantity.

Bruise them with the hands, and leave them in a cool place till a clear liquor appears on the surface of the mass; then express the juice, depurate it, and preserve it in long-necked bottles, under a stratum of oil.

ROB OF MULBERRIES.

Rob Mororum seu Succus Mororum inspissatus. (ANT.; BRUNS.; DAN.; LIPP.; LUSIT.; PALAT.; SARD.; SAX.; WIRTEM.; FEE.; NIEMANN.; SWEDIAUR.)

R. Ripe Mulberries, any quantity.

Bruise them in a stone mortar, and leave the mass in a cellar for two days, in a covered earthen vessel; then express strongly, set aside the juice in a cool place, strain through flannel, and evaporate it, constantly stirring, to the consistence of honey. (LUSIT.; NIEMANN.)

R. Depurated Juice of Mulberries, four parts.
White Sugar, one part.

Boil to the consistence of honey. (BRUNS.; DAN.; LIPP.; WIRTEM.; SWEDIAUR.)

SARD. prescribes five pounds of juice and one pound of despumated honey; or, two of juice and one of sugar.—PALAT.—one

of sugar and five of juice.—AUSTR.—one of sugar and nine of juice.—SAX.—one of sugar and sixteen of juice.—FEE.—one of juice and two of sugar.

It is principally employed externally, as a detergent to aphthæ.

HONEY OF MULBERRIES.

Mel Fructuum Mororum. (HISP.)

R. Juice of Mulberries, one pound.
Honey, two pounds.
Water, a sufficient quantity.

Boil to the consistence of thick syrup, and strain.—Dose, one or two ounces.

SYRUP OF MULBERRIES.

Syrupus Mororum seu de Moribus seu Mori nigre. (LOND.; AMST.; ANT.; BELG.; BRUNS.; DAN.; GALL.; GENEV.; HANN.; OLDEN.; PALAT.; POL.; BORUS.; WIRTEM.; HERBIP.; COXE.; SWEDIAUR.; VAN MONS.)

Syrupus Mori. (LOND.; COXE.)

R. Strained Juice of Mulberries, one pint.
Refined Sugar, two pounds.

Make a syrup.

BRUNS.; DAN.; PALAT.; WIRTEM.; and HERBIP. prescribe ten ounces of juice and sixteen of sugar.—ANT.—four pounds of juice and six of sugar.—HANN.; and BORUS.—twenty ounces of juice and three pounds of sugar.—SWEDIAUR.—sixteen ounces of juice and three pounds of sugar.—OLDEN.; and POL.—twenty ounces of juice and three pounds of sugar.—GENEV.—one pound and a half of juice and two pounds and a half of sugar.

R. Mulberries, before perfectly ripe, } of each
White Sugar, } an equal part.

Mix, boil slowly for a short time, and strain through a silk-sieve. (GALL.)

VAN MONS directs an equal part each, of mulberries and of sugar to be heated together without boiling, and the mass while hot to be placed on a sieve and slightly expressed; the liquor to be then strained through close flannel, and a sufficient quantity of sugar to be added to give it the consistence of syrup.

MAGNES proposes the following process:

R. Mulberries, before perfectly ripe, } of each
Blackberries, } an equal part.

Bruise them in a stone mortar, and put them into a silver pan with one-fiftieth of distilled water; boil for a few moments, strain through a hair-sieve, without expression, and set apart the juice for two or three days; then decant and pour it into bottles containing about one-tenth by weight, of the juice of very ripe mulberries. Having stopped the bottles, plunge them into a pan of water; heat for a quar-

ter of an hour, and withdraw and set aside the bottles. Then

R. Of this depurated and filtered juice,
one pound.

White Sugar, one pound ten ounces.

Dissolve slowly.

VAN MONS has also proposed to dissolve in a water-bath thirteen parts of sugar in eight of juice.

Dose, from one ounce to four ounces.

MOSCHUS.—MUSK.

SYNONIMES:—Musc, Fr.; Bisam, GERM.; Mischk, ARAB.; DUK.; and PERS.; Rutta urula, CYNG.; Desmer, DAN.; Almi-zele, SPAN.; Kustowrie, HIND.; Muscus, DUT.; Moschio, ITAL.; Dedes, JAV.; Jebat, MAL.; Pizmo, POL.; Misca, PORT.; Casturi, SANS.; TAM.; and TEL.; Desman, Mysk, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FULD.; FORM. HÔP. FR.; GENEV.; HAMB.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINS-LIE; BRUGNATELLI; COXE; GUIBOURT; PARIS; PIDERIT; SAUNDERS; SPIEL-MANN.

An almost solid, granular substance, of a blackish-brown colour, of a bitter and aromatic taste, and of a very strong peculiar odour. It is secreted by the inner surface of a pouch situated before the prepuce of the *Moschus moschiferus*, LINN., a ruminating mammiferous animal of Asia.

Musk contains, according to BLONDEAU and GUIBOURT, ammoniac, elaine, stearine, cholesterine, a volatile oil, gelatine, albumen, an intermediate acid, &c. It decomposes the nitrate of silver, sulphate of iron, deuto-chloruret of mercury, and the infusion of yellow cinchona.

It is a diffusible stimulant, and anti-spasmodic, and has been extolled in hysteria, epilepsy, hooping cough, hydrophobia, and tetanus. — Dose, from twenty to thirty grains.

ANTI-SPASMODIC POWDER.

Pulvis Moschi opiatu. (AUGUSTIN)

R. Musk, thirty-two grains.
Opium, four grains.
VOL. II.

Sugar, one dram.

Divide into twelve parcels.—Dose, one, every hour or oftener.

APHRODISIAC POWDER. (PIERQUIN)

R. Cinnamon, half an ounce.
Ginger, }
Nutmeg, } of each, one dram.
Long Pepper, }
Musk, one scruple.
White Sugar, four ounces.

Mix.

ANTI-SPASMODIC BOLUSES. (BORIES;
CADET DE GASSICOURT; ELLIS; SAUNDERS;
SAINTE-MARIE; SWEDIAUR)

R. Musk, from ten to fifteen grains.
Conserve of Orange Peel, half a dram.
Simple Syrup, a sufficient quantity.

Mix.—Eight grains of camphor may be added. (SWEDIAUR)

SAUNDERS prescribes half a scruple of musk, half a dram of conserve, and simple syrup.

R. Musk, ten grains.
Red Sulphuret of Mercury, five grains.
Conserve of Orange Flowers,
a sufficient quantity.

Make one or two boluses. (CADET DE GASSICOURT; SAINTE-MARIE)

R. Musk, }
Nitrate of Potass, } of each, six grains.
Camphor, four grains.
Conserve of Valerian,
a sufficient quantity.

Make four boluses. (BORIES)

POWDER OF MUSK. (SARD.)

R. Musk, four grains.
Sugar of Milk, eight grains.

Mix.—For one dose.

2 K

COMPOUND POWDER OF MUSK.

Pulvis Moschi compositus seu alexipharmicus sinicus seu anti-lyssicus Tunchinensis. (HASS.; ROSS.; WIRTEM.; HERBIP.; HENRI; NIEMANN; AUGUSTIN; CADET DE GASSICOURT; PIDERIT; SPIELMANN; VAN MONS)

R. Musk, sixteen grains.
Red Sulphuret of Mercury, twelve grains.

Mix. (WIRTEM.; NIEMANN)—The Chinese take this dose in a spoonful of spirit, as a remedy against hydrophobia.

SPIELMANN prescribes one part of musk and two parts of sulphuret.—PIDERIT has modified this formula in the following manner:

R. Musk, sixteen grains.
Root of Valerian, one scruple.
Camphor, six grains.

Make a powder. (HASS.; ROSS.; WIRTEM.; AUGUSTIN; CADET DE GASSICOURT; HENRI; NIEMANN; PIDERIT)

VAN MONS prescribes eight parts of musk, ten of valerian, and three of camphor.

R. Musk, one dram.
Oxide of Zinc, twenty-four grains.
Sugar, two drams.

Mix. (HERBIP.)

R. Musk, fifteen grains.
Camphor, five grains.

Mix.—For a bolus. (ELLIS)

R. Musk, } of each,
Subcarbonate of } half a scruple.
Ammonia, }

Mix.—For a bolus. (ELLIS)

PLASTER OF MUSK.

Emplastrum moschatum. (AUGUSTIN)

R. White Wax, } of each, half an ounce.
Spermaceti, }
Camphor, half a dram.
Castor, }
Musk, } of each, one scruple.
Oil of Cajeput, ten drops.

Melt together.—It is anti-odontalgic.

PILLS OF MUSK.

Pilule anti-hystericae seu emmenagogae; Pilules contre la pourriture d'hôpital. (AUGUSTIN; BRERA; NIEMANN; RATIER; SAINTE-MARIE)

R. Musk, two drams.
Oxide of Zinc, one dram.

Make thirty pills. (RATIER)—They are useful in all nervous diseases.

R. Musk, half a grain.

Assafoetida, two grains.

Make one pill. (RATIER)

AUGUSTIN prescribes two drams of musk and six of assafoetida, to be made into pills of four grains each; the dose of which is from twenty to thirty, twice a day.

R. Musk, one grain.
Assafoetida, } of each, two grains.
Gum Ammoniac, }
Powdered Valerian, eight grains.
Simple Syrup, a sufficient quantity.

Make two pills. (RATIER)

R. Musk, fifteen grains.
Nitrate of Potass, } of each,
Assafoetida, } two scruples.
Extract of Cinchona, } of each,
Catechu, } two drams.

Make pills of six grains each. (RATIER)

R. Musk, eight grains.
Subcarbonate of Ammonia, five grains.
Mucilage of Gum Arabic, a sufficient quantity.

Make a pilular mass. (NIEMANN)

From the recommendation of WHITE, LENTIN employed these pills, and with success, in arresting the progress of sphacelus. DUPUYTREN also employs the following, against hospital gangrene (*pourriture d'hôpital*):

R. Camphor, twenty-four grains.
Musk, eight grains.
Extract of Opium, two grains.
Simple Syrup, a sufficient quantity.

Make a pilular mass. (RATIER)

R. Musk, } of each, one scruple.
Ambergris, }
Opium, three grains.
Peruvian Balsam, a sufficient quantity.

Make twenty-four pills. (SAINTE-MARIE)

Dose, two, three times a day.

R. Musk, half a scruple.
Camphor, one scruple.
Gum Ammoniac, two scruples.
Opium, four grains.

Make pills of four grains each. (SAINTE-MARIE)

Dose, three or four daily, in hysteria and hypochondria.

R. Musk, twenty-four grains.
Extract of Opium, twelve grains.
Valerian, twenty-four grains.

Make sixteen pills.

R. Musk, twelve grains.
Rob of Juniper, a sufficient quantity.

Make nine pills.—Dose, one every hour. (BRERA)

R. Musk, eight grains.
Salt of Hartshorn, five grains.

Mucilage of Gum Arabic,
a sufficient quantity.

Make one pill. (AUGUSTIN)

- R. Aqueous Extract of Myrrh, one dram.
Aloes, half a dram.
Camphor, ten grains.
Musk, twenty grains.
Peruvian Balsam, a sufficient quantity.

Make pills. (PIERQUIN)

BARTHEZ prescribed them, as emmenagogue, in the dose of twenty-four grains daily, at thrice, with decoction of rue.

- R. Calumba, half a dram.
Subcarbonate of Potass, one scruple.
Aqueous Extract of Opium, two grains.
Extract of White Henbane, } of each,
Ambergris, } six grains.
Musk, } four grains.
Oil of Mint, one scruple.
Syrup of Orange Flowers, a sufficient quantity.

Make twenty-five pills. (BORIES)

Dose, three, every two hours.

APHRODISIAC LOZENGES.

Tablettes de Vie. (BORIES; PIERQUIN; SAINTE-MARIE)

- R. Alkermes Confection, two drams.
Ambergris, } of each, one dram.
Musk, }
Sugar, one pound.

Make lozenges of one scruple each.—Dose, one, three times a day.

These lozenges may prove dangerous, particularly when they contain cantharides; but we cannot see why SAINTE-MARIE should designate as dangerous *les plaisirs de l'amour*; but to apply to the use what should be said of the abuse, is a fault which medical writers too frequently commit in their works.

MUSK BALLS.

Globuli moschati. (BRUNS.; FULD.; LIPP.; PALAT.; WIRTEM.; SPIELMANN)

- R. White Sugar, six ounces.
Starch, two drams.
Florentine Iris, one dram.
Musk, two grains.
Mucilage of Tragacanth, a sufficient quantity.

Make balls of various sizes, from that of a pea to that of a nut.

These were formerly used in the form of a suppository, to provoke alvine dejections.

MUSK TROCHISCS.

Trochisci Gallie moschatae seu Aliptæ moschatae. (WIRTEM.)

- R. Ambergris, three drams.
Aloe Wood, five drams.
Musk, one dram.
White Sugar, one ounce.
Mucilage of Tragacanth, a sufficient quantity.

Make trochisks. They are added to other preparations which require to be flavoured.

- R. Storax, four ounces.
Opium, } of each, one ounce.
Benzoin, }
Aloe Wood, six drams.
White Sugar, one ounce and a half.
Musk, one scruple.
Ambergris, one dram.
Mucilage of Tragacanth, a sufficient quantity.

Make trochisks.—They are now scarcely employed except for fumigations. They used formerly to be prescribed in asthma.

MOGUL TROCHISCS. (CADET DE GASSI-COURT)

- R. Sugar, three ounces and a half.
Gum Arabic, one ounce.
Hard Extract of Opium, seven drams.
Cloves, } of each,
Mace, } two drams and a half.
Nutmegs, }
Musk, five grains.
Rose Water, a sufficient quantity.

Make trochisks of six grains each. They are excitant and stomachic.—Dose, two trochisks, at bed-time.

GOA STONE. (WIRTEM.)

- R. White Coral, } of each, one ounce.
Red Coral, }
Oriental Bezoar, }
— Pearls, }
Prepared Hyacinth, } of each,
Topaz, } half an ounce.
Sapphire, }
Rubies, }
Emeralds, }
Musk, } of each, two drams.
Ambergris, }
Gold Leaves, forty.
Hartshorn Jelly, a sufficient quantity

to make a paste, with which oval balls are to be formed, to be dried, and then polished.

This absurd preparation was imported from the Indies, and was for a considerable time kept secret by a couple of Jesuits.

It was formerly considered anti-spasmodic and anti-epileptic.—Dose, from half a scruple to half a dram.

MIXTURE OF MUSK.

Mixtura moschata seu Moschi, Julapium Moschi seu moschatum. (LOND.; AMER.; HISP.; HANN.; LUSIT.; AUGUSTIN.; BRERA; BRUGNATELLI; CADET DE GASSICOURT; COXE; ELLIS; NIEMANN; PIERQUIN; RATIER; SAUNDERS; SWEDIAUR)

Mixtura Moschi. (LOND.)

R. Musk, } of each,
 Refined Sugar, } one dram.
 Gum Arabic, }
 Rose Water, six fluid ounces.

Triturate the musk with the sugar, then add the gum, and finally the rose water very gradually.

AMER.; LUSIT.; BRUGNATELLI; COXE; ELLIS; and NIEMANN prescribe two scruples of musk, six ounces of rose water, and one dram each, of gum and of sugar.

R. Musk, twelve grains.
 Sugar, two drams.
 Water, two ounces.

Mix. (RATIER)

HANN. prescribes twelve grains of musk, one dram of sugar, and six ounces of rose water.

R. Musk, two scruples.
 Gum Arabic, dissolved in water, one dram.

Compound Spirit of Juniper, one ounce.
 Simple Syrup, three drams.
 Spearmint Water, five ounces.

Mix. (SWEDIAUR)

SAUNDERS prescribes two scruples of musk, one dram of mucilage, half an ounce of compound spirit of juniper, three drams of syrup, and five ounces of pennyroyal water.

R. Camphor, half a dram.
 Musk, twenty-five grains.
 White Sugar, } of each,
 Mucilage of Tragacanth, } one dram.
 Vinegar, half an ounce.
 Lettuce Water, six ounces.
 Syrup of Lemon, one ounce.

Mix. (PIERQUIN)

R. Musk, six grains.
 Sugar, one dram.

Triturate, and add

Rose Water, } of each, two ounces.
 Balm Water, }
 Syrup of Æther, one ounce.

Mix. (PIERQUIN)

R. Rose Water, six ounces.
 Hordeated Cinnamon Water, }
 Compound Water of Peony, } of each,
 Syrup of Clove-pink, } two ounces.
 Orange-flower Water, one ounce.
 Alkermes Confection, one dram.
 Saffron, one scruple.
 Musk, } of each,
 Ambergris, } two
 Subcarbonate of Ammonia, } grains.
 Essence of Cloves, one drop.

Mix. (CADET DE GASSICOURT; PIERQUIN)

R. Orange Flowers, } of each,
 Cloves, } one scruple.
 Saffron, }
 Water, one pound.

Infuse for an hour, and add to the strained liquor

Musk, } of each, two grains.
 Ambergris, }
 Subcarbonate of Ammonia, one grain.
 Alkermes Confection, two drams.
 Simple Syrup, one ounce and a half.

Mix. (HISP.)

R. Camphor, two drams.
 Musk, one scruple.
 Salt of Hartshorn, sixteen grains.
 Syrup of Orange Peel, one ounce.
 Lemon Water, three ounces.

Mix.—Dose, one spoonful, every hour. (BRERA)

R. Peppermint Water, } of each,
 Lemon Water, } two ounces.
 Musk, } eight grains.
 HOFFMANN'S LIQUOR, } of each,
 Succinated Spirit of Ammonia, } half a dram.
 SYDENHAM'S Laudanum, twenty drops.

Mix.—To be taken gradually. (BRERA)

R. Musk, twelve grains.
 Peppermint Water, three ounces.
 Syrup of Orange Peel, one ounce.

Mix.—Dose, one tea-spoonful every hour. (BRERA)

R. Balm Water, } of each,
 Cinnamon Water, } two ounces.
 Camphor, } one scruple.
 Musk, } of each,
 Salt of Hartshorn, } fifteen grains.
 Syrup of Narcotine, one ounce.

Mix.—To be taken gradually. (BRERA)

R. Musk, one scruple.
 White Sugar, two drams.
 Mucilage of Gum Arabic, half an ounce.

Cinnamon Water,	}	of each,
Peppermint Water,		two ounces.
Aromatic Liquor,		two drams.

Mix.—To be taken by spoonfuls. (AUGUSTIN)

R. Musk,	one scruple.
Sugar,	one dram.
Rose Water,	eight ounces.
Sulphuric Æther,	two drams.
Oil of Cajeput,	twenty drops.

Mix.—To be taken by spoonfuls. (AUGUSTIN)

R. Musk,	six grains.
Sugar,	one dram.
Succinated Spirit of Ammonia,	sixty drops.
SYDENHAM'S Laudanum,	twenty drops.
Elder-flower Water,	five ounces.
Amygdaline Syrup,	one ounce.

Mix.—A spoonful to be taken every hour. (AUGUSTIN)

ANTI-SPASMODIC LAVEMENT. (ELLIS ; PIERQUIN ; SWEDIAUR)

R. Musk,	twelve grains.
Sugar,	two scruples.
Gum Arabic,	one scruple and a half.
Solution of Ammonia,	thirty drops.
Linseed Water,	four ounces.

Mix. (ELLIS)

R. Infusion of Valerian,	four ounces.
Musk,	one dram.

Mix, and add, if required, forty or sixty drops of tincture of opium. (PIERQUIN ; SWEDIAUR)

VINOUS TINCTURE OF MUSK.

Tinctura Moschi vinosa. (NIEMANN)

R. Musk, in powder,	one ounce.
Spanish Wine,	six ounces.

Triturate the musk with one ounce of wine, till it becomes dissolved; then add the rest of it. Macerate for twenty-four hours in a well-stopped bottle, and filter.

It is anti-spasmodic.

TINCTURE OF MUSK.

Tinctura seu Essentia Moschi seu moschata. (DUBL. ; AMER. ; MIL. COPEN. ; GALL. ; HANN. ; COXE ; NIEMANN)

Tinctura Moschi. (DUBL.)

R. Musk, in powder,	two drams.
Rectified Spirit,	one pint.

Digest for seven days, and filter.

AMER. ; COXE ; and NIEMANN give the same formula.—MIL. COPEN. ; and GALL. prescribe one part of musk, and four parts of alcohol (22 degrees).—HANN.—two drams of musk, and eight ounces of rectified spirit.

Dose, from thirty to sixty drops, several times a day.

ÆTHEREAL TINCTURE OF MUSK. (GALL.)

R. Musk,	two drams.
Sulphuric Æther, (46 degrees)	one ounce.

Macerate for two days, and decant.

CONDOM'S BALSAM.

Balsamum Lectorense ; Baume de Leictour, de Vinceguère, or de Condom. (SPIELMANN)

R. Camphor,	}	of each, one dram.
Saffron,		
Musk,		
Amber,	}	of each, half a dram.
White Petroleum,		
Oil of Lavender,		
— Turpentine,	}	of each, one ounce.
— Juniper,		
— Cloves,		
— Mace,	}	of each,
— Nutmeg,		two drams.
— Benzoin,		half an ounce.

Infuse, and pour off the liquid as it may be wanted.

It is anodyne, nerveine, and balsamic.

CEPHALIC OLEOSACCHARUM.

Pulvis sternutatorius albus, Oleosaccharum cephalicum. (WIRTEM.)

R. White Sugar,	twelve ounces.
Ambergris,	of each,
Musk,	half a scruple.
Oil of Cinnamon,	of each,
— Cloves,	fifteen drops.
— Anise,	
— Rosemary,	of each,
— Lavender,	half a dram.
— Marjoram,	
— Rhodium Wood,	of each,
— Amber,	one scruple.

Dissolve the ambergris in the oils, and the musk in a small quantity of oil of roses; then mix both with the sugar.

R. Oil of Marjoram,	}	of each,
— Cloves,		three drops.
— Rhodium Wood,		
— Cinnamon,		two drops.

Oil of Lemon,	}	of each, two drops.
— Fennel,		
— Lavender,		
Musk,		three grains.
White Sugar,		three ounces and a half.

Mix.

It is employed as a sternutatory, and also in spasmodic affections and paralysis of the tongue.—Dose, from a scruple to half a dram.

MUSCARI AMBROSIACUM.—MUSK HYACINTH.

SYNONIMES:—Muscare odorant, Jacinth musquée, FR.; Muscatenhyacinth, GERM.; Muskeerende hyacinth, DUT.

AUSTR.; FERR.

This plant is originally of the Levant.

(*Hexandria Monogynia*, LINN.; *Asphodelea*, JUSS.; *Fig. Kniph. Bot. orig. cent. 10. t. 52*)

The root is employed; it is thick. Its taste is bitter, and it is considered emetic.

MYRICA GALE, LINN.—DUTCH MYRTLE.

SYNONIMES:—Galé odorant, Piment royal, Myrte bâlard, FR.; Myrtenheide, Brabantische Post, Gerbemyrtenstrauch, GERM.; PORS, DAN.; and SWED.; Gagel, DUT.

AMST.; BRUNS.; GALL.; WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GME-LIN; SPIELMANN.

A shrub of Europe. (*Diœcia Pentandria*,

LINN.; *Amentacea*, JUSS.; *Fig. Nouv. Duh. II. t. 57*)

The herb (*Herba Myrti Brabantica* seu *Gales* seu *Chamaeleagni*) is employed. It consists of reddish-brown branches, furnished with oblong leaves, broadened and indented at their upper part, narrowed at their base, and supported upon short foot-stalks. Its odour is strong, agreeable, and aromatic; and its taste aromatic, bitter, and sub-astringent.

MYRISTICA AROMATICA MUR.—NUTMEG TREE.

An original tree of the Moluccas. (*Diœcia Monadelphica*, LINN.; *Myristicea*, CAND.; *Fig. Flore Medic. VI. 242*)

The seed and its arillus are employed.

The seed bears the name of *Nutmeg*; *Nux moschata* seu *Myristica*.

SYNONIMES:—Noix muscade, FR.; Muskatennuss, GERM.; Jowzalteih, ARAB.; Muskatowa kulka, BOHEM.; Buwahpa, BAL.; Jayaphala, BE.; Jatipullum sadikka, CYNG.; Muskad, DAN.; Japhul, DUK.; Nuez de especia, HISP.; Jaiphul, HIND.; Nooten, moskaat, DUT.; Noce moscada, ITAL.; Wohpala, JAV.; Japhul, MAH.; Buahpala, MAL.; Noz moscada, PORT.; Jatiphala, SANS.; Muskot, SWED.; Jadicaï, TAM.; Jajikaia, TEL.; Gasori, TERN.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.;

BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PARIS.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

This is oval, oblong, sub-rotund, covered with a reddish pellicle, whitish and dotted with brown points at the summit; its substance is white and oleaginous. It is contained in a hard, thin, brown or blackish shell, irregularly grooved. Its odour is very agreeable, and its taste warm and aromatic.

The arillus, or envelope between the nut and shell, is called *Mace*, or *Flowers of Nutmeg*; *Macis* seu *Flores Macis*.

SYNONIMES:—Macis, Fleurs de muscade, FR.; Muskatenblüthen, GERM.; Talzuffur, ARAB.; Bungapala, BAL.; Wassawasie, CYNG.; Jawatrie, DUK.; and HIND.; Macias, SPAN.; Kambang pala, JAV.; Bunga bua pala, MAL.; Bezbaz, PERS.; Flor de nozimoscada, PORT.; Jatipatri, SANS.; Jadiputrie, TAM.; Japatri, TEL.

EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; AINSLIE.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SAUNDERS.; SPIELMANN.

It is a laciniated, reticulated, thin, cartilaginous, semi-transparent membrane of a saffron-yellow colour, and of a taste and odour resembling those of nutmeg.

It is a very energetic stimulant, but is rather culinary than medicinal. — Dose, from ten grains to a scruple, of the powder.

Mace is much less active than nutmeg.

ANTI-SCROFULOUS POWDER.

Pulvis anti-hectico-scrofulosus seu ex Baccis Lauri compositus seu Nucum moschatarum compositus. (NIEMANN)

R. Nutmegs, } of each,
Torrefied Bay Berries, } one ounce.
Calcined Hartshorn, }
Liquorice Root, three ounces.

Pulverize, and mix.—Dose, half a teaspoonful, twice a day, to children of one year; and one spoonful, to those of two to four years.

AROMATIC BOLUS. (SAUNDERS)

R. Nutmeg, one scruple.
Diascordium, two scruples.
Syrup of Poppies, a sufficient quantity.

Mix.

SPIRIT OF NUTMEG.

Spiritus Nucis Moschatæ seu Myristicæ aromatica. (LOND.; DUBL.; EDINB.; LIPP.; COXE)

Spiritus Myristicæ, (LOND.)

Sp. Myristicæ moschatæ. (EDINB.)

Sp. Nucis Moschatæ. (DUBL.)

R. Nutmegs, bruised, two ounces.
Proof Spirit, one gallon.
Water, a sufficient quantity
to prevent empyreuma. Distil one gallon by a gentle heat. (LOND.; DUBL.)

EDINB. prescribes nine pounds of weaker alcohol, and nine pounds of spirit to be distilled.

LIPP. prescribes two ounces of nutmegs, four pounds each, of alcohol and of water; to be macerated for twenty-four hours, and four pounds distilled.

COMPOUND SPIRIT OF NUTMEG. (VAN MONS)

R. Nutmegs, two parts,
Lemon Peel, }
Orange Peel, } of each, one part.
Balm, }
Mentha Crispa, }
Proof Spirit, sixty-four parts.
Water, one hundred and ninety-two parts.

Distil forty-eight parts.

SPIRITUOUS AROMATIC LIQUOR. (FULD.)

R. Nutmegs, }
Lesser Cardamoms, } of each,
Orange Peel, } one ounce.
Ginger, }
Subcarbonate of Potass, }
Proof Spirit, six pounds.
Water, two pounds.

Digest for twenty-four hours, and distil four pounds in a water-bath.

It is stimulant, anti-spasmodic, carminative, and stomachic.—Dose, from forty to sixty drops.

SYRUP OF NUTMEG.

Syrupus Nucistæ. (WIRTEM.)

R. Nutmegs, in powder, two ounces.
Burgundy Wine, one pound.

Digest for three days in a covered vessel; then express, and dissolve in the liquor,

White Sugar, sixteen ounces.

Mix.—It is carminative and stomachic.

TINCTURE OF MACE.

Tinctura seu Essentia Macidis seu Macis. (BRUNS.; HANN.; PALAT.; SAX.)

R. Mace, one part.
Rectified Spirit, ten parts.

Digest for eight days in a moderate heat; then express and filter. (SAX.)

BRUNS. prescribes three parts of mace and sixteen of spirit.—HANN.—five ounces of mace and two pounds of spirit.

R. Mace, four ounces.
Solution of Subcarbonate of Potass, two ounces.

Digest to complete desiccation; then pour upon the residue,

Rectified Spirit, two pounds.

Digest for a sufficient time, and distil in

a sand-bath; cohobate several times, and never distil to dryness. Finally pour the product upon,

Mace, any quantity.

After some time, filter. (PALAT.)

It is excitant, stomachic, uterine, and anti-colical.—Dose, thirty drops or more.

EXPRESSED OIL OF NUTMEG.

Oleum Nucistæ seu Nucis moschatae, Oleum æthereo-pinguae Nucleorum Myristicæ moschatae, Oleum seu Balsamum Nucistæ expressum.

1. That which is obtained in commerce.

LOND.; DUBL.; DAN.; OLDEN.; WIRTEM.

2. Prepared expressly.

AUSTR.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HASS.; OLDEN.; PALAT.; BORUS.; SAX.; WIRTEM.; BRUGNATELLI; PIDERIT; SWEDIAUR; VAN MONS.

R. Nutmegs, rasped, any quantity.

Reduce to a soft paste by trituration in a heated marble mortar, enclose this paste in a hair-bag, press it between two heated tin plates, melt the oil and strain it through linen. (HISP.; FERR.; VAN MONS)

R. Nutmegs, any quantity.

Bruise them in a warm stone mortar, and expose them in a hair bag, to the vapour of boiling water; then submit it to the press between two heated tin plates, and melt the oil over the fire. (BAVAR.; BRUNS.; DAN.; FULD.; HASS.; OLDEN.; PALAT.; BORUS.; WIRTEM.; BRUGNATELLI; PIDERIT; SWEDIAUR)

R. Nutmegs, any quantity.

Bruise them in a mortar; then, having worked the mass upon a heated slab, boil it in water till the oil becomes separated, and remove it from the surface when it has become concrete by cold. (GALL.)

It is stimulant and stomachic.—Dose, one or two grains, to children; one scruple and a half, to adults.

ESSENTIAL OIL OF MACE.

Oleum Macidis æthereum, Ætheroleum Macidis. (BAVAR.; BRUNS.; DAN.; FULD.; HANN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; VAN MONS)

R. Mace, any quantity.

Water, a sufficient quantity.
Infuse, and distil. (ROSS.)

SAX.; and HERBIP. prescribe one part of mace and four parts of water.—DAN.; and FULD.—one of mace and six of water.—HANN.; PALAT.; and BORUS.—one of mace and eight of water.—BAVAR.—one of mace and nine of water.—VAN MONS—one of mace and ten of water.—PALAT.; and WIRTEM.—two pounds of mace, sixteen pounds of water, and four ounces of muriate of soda.

Dose, from four to twelve drops.

OLEOSACCHARUM OF MACE. (BRUNS.; PALAT.; and WIRTEM.)

R. Essential Oil of Mace, sixteen drops.
White Sugar, one ounce.
Triturate together.

BALSAM OF NUTMEG.

Balsamum Nucistæ. (BRUNS.; SAX.; WIRTEM.)

R. Expressed Oil of Nutmeg, two drams.
Essential Oil of Nutmeg, two scruples.
Triturate together.

It is stomachic, cephalic, and nervine.

ANTI-DIARRHOIC MIXTURE. (PIERQUIN)

R. Balsam of Nutmeg, one dram.
Theriac, two drams.
Oil of Cumin, six drops.

Mix.—To be applied to the umbilical region.

STIMULANT LINIMENT. (NIEMANN)

R. Essential Oil of Nutmeg, } of each,
Cloves, } half a dram.
Spirit of Juniper, } two ounces.
Mix.

AROMATIC SPARADRAP.

Sparadrap ad Corpus conditum involvendum. (SPIELMANN)

R. White Wax, one hundred and twenty-eight ounces.
Melt it, and when nearly cold, add to it
Expressed Oil of Nutmeg,

eight ounces.
Essential Oil of Lemon, } of each,
Neroli, } four
Lavender, } ounces.
Thyme, }
Ambergris, fifteen grains.
Musk, eight grains.
Civet, four grains.

Mix, and spread the mass upon strips of linen.

MYROBALANUS.

BRUNS.; HISP.; GALL.; WIRTEM.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

This name is given to four kinds of fruit.

1. *Myrobalani bellirica*.

SYNONIMES:—Myrobalans bellirics, FR.; Bellirische Myrobalanen, GERM.; Beleyluj, ARAB.; Bullu, CYNG.; Bulla, DUK.; Mirobalanos belliricos, SPAN.; Beheyra, HIND.; Beleyleh, PERS.; Bahira, SANS.; Tanikai, TAM.

These are oval, nearly globular drupes, of the size of an olive, of a deep greyish-yellow colour, indistinctly quincilateral, enclosing an osseous, thick, pentagonal, irregular stone, broad and obtuse at the base, and acuminate at the summit.

They are derived from the *Myrobalanus bellirica*, GÆRTN. (*Decandria Monogynia*, LINN.; *Myrobalanæ*, JUSS.), a tree of the East Indies. (Fig. Roxb. *Corom.* t. 198)

2. *Myrobalani chebula*.

SYNONIMES:—Myrobalans chebules, FR.; Grosse Schwarzbraune Myrobalanen, GERM.; Helelije Kabuli, ARAB.; Aralu, CYNG.; Huldah, DUK.; Mirobalanos quebulo, SPAN.; Har, Hara, HIND.; Helileh Kelan, PERS.; Haritaka, SANS.; Kadukai, TAM.; Curakai, TEL.

These are oval, long drupes, narrowed at each extremity, of a blackish-brown colour, marked externally by six angles, produced by five sides alternating with five grooves; it contains, under a hard fragile pulp of a resinous appearance, an osseous pentagonal shell, enclosing an oval, long, acuminate seed.

When less ripe, dried before their maturity, and probably punctured by some insect, they constitute the *Myrobalani indicæ* seu *nigræ*. (*Myrobalans indiques ou noir*, FR.; *Indianische Myrobalanen*, GERM.; *Myrobalanos indicos*, SPAN.) These are drupes of the size of a filbert, of a deep black colour, hard, compact, and scarcely presenting the vestige of a stone, in the

place of which there is generally only a small cavity. Their taste is bitter and astringent.

These fruits proceed from the *Myrobalanus chebula*, GÆRTN., a tree of the East Indies. (Fig. LMK., *Enc. Meth.* t. 849, f. 4)

3. *Myrobalani citrina* seu *flavæ*.

SYNONIMES:—Myrobalans citrins, FR.; Gelbe Myrobalanen, GERM.; Mirobalanos citrinos, SPAN.

They are long, oval, sub-pyriform drupes, of a pale yellow colour, with variable angles, and wrinkled between the angles. They very much resemble the preceding internally, and are very probably only a variety of them.

They come from the *Myrobalanus citrina*, GÆRTN., a tree of the East Indies. (Fig. LMK. *Enc. Meth.* t. 849)

4. *Myrobalani emblica*.

SYNONIMES:—Myrobalans emblics, FR.; Aschfarbene myrobalanen, GERM.; Hacin-san, Cay-boung-ngat, COCHIN.; Awasadanelli, Aunla, CYNG.; Mirobalanos emblicos, SPAN.; Anola, Amle, Aunwerd, Aongra, HIND.; Amleh, PERS.; Amalaka, SANS.; Nelli kai, TAM.; Wusoherikaia, TEL.

These are capsules formed of berries, with three lobes, rounded, of the size of a large nut-gall, of a blackish-grey colour, with six valves raised exteriorly at the sides, and containing internally, a fleshy pulp with whitish and angular seeds. They have a sour and astringent taste.

They are derived from the *Phyllanthus emblica* (*Monæcia Triandria*, LINN.; *Euphorbiacæ*, JUSS.), a tree of Malabar. (Fig. Rheed. *Malab.* I. 31)

All these fruits, disused at the present day, were formerly given as purgatives, in the dose of from half a dram to half an ounce, of the powder.

MYRTUS COMMUNIS, LINN.—MYRTLE.

SYNONIMES:—Myrte commun, FR.; and GERM.; Arajan, SPAN.; Myrtus, DUT.; Myrto, ITAL.; Marta, PORT.

VOL. II.

BRUNS.; HISP.; GALL.; FERR.; LUSIT.; WIRTEM.; BERGIUS.; BRUGNATELLI.; MURRAY and GMELIN.; SPIELMANN.

2 L

A shrub of the south of Europe. (*Icosandria Monogynia*, LINN.; *Myrtææ*, JUSS.; *Fig. Zorn, Ic. pl. t. 169*)

The leaves and fruit are employed.

The leaves (*Folia Myrti seu Baticæ sylvestris*) are oval or lanceolated, smooth, of a fine green colour, sprinkled with semi-transparent glands. They develop, particularly when rubbed, an agreeable odour; their taste is bitter, aromatic, and rather styptic.

The fruits are globular berries, of a blackish-blue colour, of an aromatic odour, and astringent and bitter taste.

It is astringent, but is seldom employed.

OIL OF MYRTLE.

Oleum Myrtinum. (HISP.; SARD.)

- R. Myrtle Berries, bruised, two pounds.
 ——— Leaves, chopped, two handfuls.
 Olive Oil, four pounds.
 Vinegar, one pound.

Boil to the consumption of the humidity, and strain. (SARD.)

- R. Leaves and Flowers of Myrtle,
 Olive Oil, one part.
 three parts.

Digest for three or four days; then boil to the consumption of the humidity, and strain with expression.

SYRUP OF MYRTLE.

Syrupus de Myrto. (GALL.; SARD.)

- R. Myrtle Flowers, one ounce.
 Distilled Water of Myrtle, two pounds.

Digest in a water-bath for two hours, in a covered vessel; then add to the strained and filtered liquor,

White Sugar, twice the quantity.

Dissolve in a water-bath. (GALL.)

SARD. directs an equal part each, of depurated juice and of simple syrup, to be boiled together à la grande plume, boiling slowly, and stirring constantly with a wooden spatula.

NAPHTHA.

BRUNS.; DAN.; GALL.; FERR.: SAX.;
 WIRTEM.; GUIBOUT.

A fluid bitumen, colourless or of a pale yellow, diaphanous, slightly unctuous to the touch; it diffuses a strong but not dis-

agreeable odour; it is much lighter than water, and is exceedingly combustible, burning with a bluish flame, and producing a very thick smoke.

It is stimulant.

NARCISSUS PSEUDO-NARCISSUS, LINN.—COMMON DAFFODIL NARCISSUS; WILD NARCISSUS.

SYNONIMES:—Narcisse sauvage, Narcisse des prés, Faux Narcisse, FR.; Gemeine Narcisse, GERM.; Gaal Narcisse, DAN.; Narcisso silvestre, SPAN.; Geele Narcis, DUT.; Narcisso giallo, ITAL.; Pæskilja, SWED.

FERR.; GALL.

A common plant of Europe. (*Hexandria Monogynia*, LINN.; *Narcissææ*, JUSS.; *Fig. Red. Lil. Vol. III. 158*)

The bulb and flowers are employed.

The bulb is viscid, and has a feebly acrid taste.

The flowers are yellow.

The bulb is purgative and emetic; dose, two drams.—The flowers are anti-spasmodic; dose, twenty-four grains.

EXTRACT OF WILD NARCISSUS.

Extractum Narcissi pratensis. (GALL.;
 VAN MONS)

- R. Fresh Flowers of Wild Narcissus,
 Water, one pound.
 two pounds.

Infuse without heat for twelve hours, and strain without expression; pour upon the residue the same quantity of hot water, and strain again. Clarify the mixed infusions with white of egg, and evaporate to the consistence of extract. (VAN MONS)

VAN MONS also directs one part of dried flowers to be digested in eight parts of warm water, and the clarified liquor to be evaporated.—GALL.—one pound of flowers

to be macerated in four pounds of water for twenty-four hours, and the operation to be repeated with the residue and fresh water; the mixed infusions to be then evaporated.

It is stimulant, nauseant, expectorant, and anti-spasmodic, and has been recommended in whooping cough, and spasmodic affections.—Dose, from four to twenty grains, according to age.

VINEGAR OF WILD NARCISSUS.

Acetum Pseudo-narcissi. (VAN MONS)

- R. Fresh Flowers of Wild Narcissus,
one part.

Vinegar, eight parts.

Infuse in a moderate temperature for several hours; then express, and filter.

It is used in preparing the following.

OXYMEL OF WILD NARCISSUS.

Oxymel Narcissi pratensis. (VAN MONS)

- R. Vinegar of Wild Narcissus, one part.
Honey, four parts.

Dissolve without heat.

It is nauseant, expectorant, and sedative; it is recommended in whooping cough and spasmodic asthma.—Dose, one or two tea spoonfuls, three or four times a day.

NARCOTINA.—NARCOTINE.

HENRI.

- R. Residue of Opium exhausted by
water, any quantity.

Dry, pulverize, and digest, without heat, with hydrochloric acid (4 degrees), or with pyro-acetic acid (4 or 5 degrees); express and filter the liquor, and add to it an excess of ammonia: wash the precipitate with cold water, and digest it in boiling alcohol (36 degrees); then having set apart the solution to crystallize, purify the crystals by means of animal carbon.

It is stupefactive, debilitating, and poisonous, according to ORFILA; powerfully excitant in large doses, according to MAGENDIE; and narcotic, according to BRERA.

NARCOTIC INJECTION. (CADET DE GASCOURT)

- R. Decoction of Virous Lettuce,
one pound.

Extract of Belladonna,

one dram and a half.

Narcotine, six grains.

Mix.—It is recommended in painful affections of the urethra and vagina.

BOLUSES OF NARCOTINE.

Boli e Narcotina. (BRERA)

- R. Narcotine, two grains.
Oil of Almonds, six drops.
Crumb of Bread, } of each,
Honey, } a sufficient quantity.

Make eight boluses.—Dose, one, every three hours.

SEDATIVE MIXTURE.

Mixtura incitans et sedans. (BRERA)

- R. Pure Narcotine, one dram.
Lemon Water, three ounces.

Mix.—Dose, one spoonful.

NEPETA CATARIA, LINN.—CAT MINT.

SYNONIMES:—Cataire, Herb aux chats, FR.; Katzenmünze, GERM.; Poleg, Natkowe korenje, Marulka polnj, BOHEM.; Katteurt, DAN.; Gatera, SPAN.; Kattenkruid, DUT.; Gattara, ITAL.; Mielka kotcza, POL.; Nevada dos gados, PORT.; Kattmynta, SWED.

ANT.; BATAV.; BRUNS.; GALL.; FERR.; GENEV.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GME-LIN; SPIELMANN; ZARDA.

A plant common throughout Europe. (*Didynamia Gymnospermia*, LINN.; *Labiata*, JUSS.; *Fig. Bull. Herb. t. 287*)

The herb (*Herba Nepetæ seu Catarisæ seu Menthæ Catarisæ*) is employed. It consists of a square, pubescent, whitish stem, and of cordiform, petiolated, serrated leaves, whitish underneath. It has an aromatic odour, and an acrid and bitter taste.

It is excitant, stomachic, incisive, carminative, emmenagogue, and nervine.

NEPHRETICUS LAPIS.—NEPHRITIC STONE.

SYNONIMES:—Pierre nephretique, Jade nephrite ou oriental, FR.; Griesslein, GERM.; Piedracananor, Piedra nefritica, SPAN.

HISP.; WIRTEM.; SPIELMANN.

A stone of a green colour and greasy aspect.

It used formerly to be worn as an amulet as a preservative from epilepsy.

NERIUM OLEANDER, LINN.—COMMON ROSEBAY.

SYNONIMES:—Laurier Rose, FR.; Rosen-sorbeer, GERM.

GALL.; FERR.

This shrub is said to be originally from the East, but it is naturalized in the South of Europe. (*Pentandria Monogynia*, LINN.; *Apocynæ*, JUSS.; *Fig. Blackw. Herb. t. 331*)

The leaves (*Folia Nerii seu Rhododaphnis seu Rosaginis*) are employed. They are long, lanceolated, pointed, rigid, coriaceous, and of a deep green colour. They have a bitterish taste.

They are reputed narcotic. The infusion is used in tetters; and the powder, incorporated with lard, applied in itch.

NICOTIANA.—TOBACCO.

Two species of this genus of plants are used in medicine:

1. *Nicotiana Tabacum*, LINN.—Tobacco.

SYNONIMES:—Tabac ordinaire, FR.; Toback, Tobak, DAN.; GERM.; and SWED.; Buijterbhang, ARAB.; Tambroco, BAL.; and JAP.; Petmne, BOHEM.; Youly, CARAMB.; Sang-yeu, CHIN.; Dunkola, CYNG.; Tumbaku, DUK.; and HIND.; Tabaco, HISP.; and PORT.; Tabak, DUT.; and POL.; Tabaco, ITAL.; Tambracu, MAL.; Quanyrell, MEX.; Dhumrapatra, SANS.; Poghei elley, TAM.; Tamer, TART.; Poghako, TEL.

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

This plant is originally of South America, but is cultivated in Europe. (*Pentandria Monogynia*, LINN.; *Solanæ*, JUSS.; *Fig. Flore Medic. VI. 336*)

The leaves (*Herba Tabaci seu Nicotianæ*) are employed. They are very large, sessile,

slightly decurrent at the base, oval, lanceolated, pointed, very entire, and nearly glabrous; they are green when fresh, but become brown after desiccation. Their odour is strong and disagreeable, and their taste, acrid, nauseous, and stupefying.

They contain, according to VAUQUELIN, a peculiar crystalline principle, *Nicotianine*, *Nicotine*, *Nicotianum*, which HERMBSTAEDT has succeeded in isolating.

2. *Nicotiana rustica*, LINN.

SYNONIME:—Tabac Rustique, FR.

GALL.

An original plant of America. (*Fig. Blackw. Herb. t. 437*)

The leaves (*Folia Nicotianæ rusticæ seu minoris*) are employed. They are oval, entire, obtuse, and sub-cordiform. They also contain nicotianine.

Tobacco is violently irritant, and poisonous in a large dose, occasioning vomiting, sanguinolent alvine dejections, vertigo and somnolency, acceleration of the pulse, diaphoresis, and diuresis. Besides its being in ordinary use as an errhine and masticatory, it is administered in chronic catarrhs and dropsy; employed in the form of lavement, it is useful to destroy ascarides, and also to induce a salutary irritation in the

rectum in cases of asphyxia and apoplexy: it is also useful externally, in tinea and itch.

Dose, from half a grain to five grains, of the powder.

ERRHINE POWDER. (HASS.; PIDERIT)

R. Tobacco, } of each,
Leaves of Marjoram, } one ounce.
Flowers of Lavender, }

Make a fine powder.

CEPHALIC POWDER. (BORIES)

R. Tobacco, half an ounce.
Marjoram, } of each,
Lily of the Valley, } half a dram.
Syrian Herb Mastic, }
Horse Chesnut, one dram.

Pulverize, and mix.

PILLS OF TOBACCO. (SUEC.; AUGUSTIN; VAN MONS)

R. Extract of Tobacco, one dram.
——— Liquorice, one ounce.
Vinegar of Squill, a sufficient quantity.
Make one hundred and eighty pills. (SWED.)

VAN MONS prescribes an equal part each, of tobacco and of liquorice-powder, with vinegar of squill.

R. Extract of Tobacco, one part.
——— Liquorice, six parts.
Liquorice Powder, two parts.

Make a pilular mass. (VAN MONS)

R. Virginian Tobacco, } of each,
Conserve of Roses, } one dram.
Mucilage of Gum Arabic, a sufficient quantity.

Make sixty pills.—Dose, from one to three, two or three times a day.

These pills are reputed diuretic, and are said to be serviceable in dropsy.

EXTRACT OF TOBACCO.

Extractum Tabaci seu Nicotianæ. (FENN.; FULD.; SAX.; SUEC.; HERBIP.; VAN MONS)

R. Fresh Leaves of Tobacco, any quantity.

Bruise them in a stone mortar, express the juice, and immediately evaporate it in a water-bath to the required consistence. (HERBIP.)

R. Fresh Leaves of Tobacco, any quantity.

Bruise them in a mortar, and express the juice; evaporate it, without removing the scum, till the addition of one fourth of powdered dry leaves of tobacco is sufficient to give it the consistence of an extract.

R. Leaves of Tobacco, } of each,
Cold Water, } one part.

Macerate for twelve hours, and add

Boiling Water, four parts.

Strain when cold, clarify with white of egg, and set the liquor upon the fire; when it begins to thicken, strain it through a sieve, and finally evaporate it to the proper consistence, constantly stirring. (SAX.; VAN MONS)

R. Leaves of Tobacco, one part.
Water, two parts.

Boil a little over a slow fire, and strain with expression; boil the residue with fresh water, and strain again. Mix the decoctions, and evaporate in a water-bath. (SUEC.)

FULD. directs one pound of tobacco to be boiled in eight pounds of water, the decoction to be expressed, and the residue to be macerated in cold water, and again expressed: the mixed liquors to be then evaporated.

Dose, from three to six grains.

MITIGATED EXTRACT OF TOBACCO. (BRUNS.; HANN.; LIPP.; WIRTEM.; VAN MONS)

R. Leaves of Tobacco, any quantity.

Deprive them completely of all taste by repeated infusions in cold water; then pour upon them twice their weight of boiling water: express when cold, clarify the liquor with white of egg, and evaporate. (VAN MONS)

BRUNS.; LIPP.; and WIRTEM. direct one pound of leaves, deprived of their acidity, to be infused and then boiled in ten pounds of water till reduced to five.

It is stimulant, and is recommended in asthma.—Dose, one grain, three or four times a day.

DECOCTION OF TOBACCO.

Decoction Tabaci seu Nicotianæ. (SWEDIAUR)

R. Dried Leaves of Tobacco, from two to four drams.
Boiling Water, one pound.

Boil for a few minutes; then digest for an hour, and strain.

It is employed in lavements, in ileus, obstinate constipation, and in cases of ascariides. It is also recommended by JUSTAMOND, in the form of lotion, in tinea; but it should be used with great caution, or it may occasion nausea and emesis.

ANTI-PSORIC LIQUOR. (FERR.)

- R. Dried Leaves of Tobacco, two pounds.
Water, fifteen pounds.

Boil for an hour, and add

Subcarbonate of Soda, four ounces.

Strain at the end of twenty-four hours.—

Dose, four ounces for each friction.

INFUSION OF TOBACCO.

Infusum Tabaci seu Nicotianæ Tabaci.
(LOND.; DUBL.; AMER.; COXE; NIEMANN)

- R. Leaves of Tobacco, one dram.
Boiling Water, one pint.

Macerate for an hour in a lightly covered vessel, and strain.

- R. Leaves of Tobacco, two drams.
Boiling Water, four ounces.

Infuse in a water-bath for four hours; express, and add

Alcohol, one ounce.

Mix. (NIEMANN)

It is stimulant, and is extolled in asthma and hydrothorax.—Dose, from twenty to forty drops.

FOWLER recommended an infusion of one ounce of Virginian tobacco in one pound of boiling water; to be digested in a water-bath for an hour, and two ounces of alcohol added to every fourteen ounces of the expressed liquor. He prescribed from sixty to one hundred drops of this liquor to adults, in asthma and dysury.—MARRYAT recommends another infusion, for tinea, prepared with two ounces of tobacco and sixteen of spring water; adding one dram of solution of potass to the strained liquor. The hair is to be removed and the ulcers fomented with this lotion, the head, at the same time, being anointed morning and evening with a mixture of one ounce of resin cerate and two drams of proto-chloruret of mercury.

SYRUP OF TOBACCO.

Syrupus de Nicotiana. (BRUNS.; FULD.; LIPP.; WIRTEM.; BORIES; NIEMANN; PIERQUIN; SPIELMANN)

- R. Juice of Tobacco Leaves, two pounds.
Hydromel, one pound and a half.
Oxymel, four ounces.

Digest for two or three days in a water-bath, in a warm place; pour off the clear liquor, set aside for twenty-four hours, and after the depuration, add,

White Sugar, three pounds.

Make a syrup. (WIRTEM.)

BRUNS. gives the same process, but pre-

scribes a decoction made with six ounces of tobacco and six pounds of water, to be boiled down to two.

NIEMANN; and SPIELMANN prescribe forty-eight ounces of tobacco juice, four of oxymel, twelve of hydromel, and forty-eight of sugar.—BRERA; and PIERQUIN—three ounces of juice, four of oxymel, one of hydromel, and two of sugar.

- R. Dried Leaves of Tobacco, half an ounce.
Water, six ounces.

Boil for an hour, and reject the decoction. Then add to the residue,

Root of Liquorice, one ounce.
Boiling Water, twelve ounces.

Boil for some time, till reduced to about three-fourths; express, and add to the liquor,

Despumated Honey, eight ounces.

Boil to the consistence of syrup. (FULD.; LIPP.)

IRRITANT LAVEMENT. (FORM. HÔP. FR.; ELLIS; NIEMANN; RATIER)

- R. Leaves of Tobacco, one dram.
Boiling Water, a sufficient quantity to obtain eight ounces of decoction. (ELLIS; NIEMANN)

- R. Leaves of Tobacco, one ounce.
Water, two pounds.

Boil, and add when strained,

Tartarized Antimony, twelve grains.

Dissolve. (FORM. HÔP. FR.; RATIER)

STIMULANT FOMENTATION.

Lotio Nicotianæ. (AUGUSTIN; RATIER)

- R. Leaves of Tobacco, two ounces.
Water, one pound.

Boil and strain.—MARRYAT; and NIEMANN extol this liquor in tinea.

DIURETIC DROPS.

Guttulæ diureticiæ. (HAMB.)

- R. Leaves of Tobacco, one dram.
Boiling Water, a sufficient quantity to obtain one ounce of infusion; add to the strained liquor

Vinous Cinnamon Water, half an ounce.

Mix.

EMETIC CATAPLASM. (ELLIS)

- R. Leaves of Tobacco, one ounce.
Water, a sufficient quantity

to make a cataplasm, to be applied to the region of the stomach, when an emetic cannot be administered by the mouth.

WINE OF TOBACCO.

Vinum Tabaci seu *Nicotianæ* seu *Nicotianæ Tabaci*. (EDINB.; AMER.; LUSIT.; AUGUSTIN; COXE; NIEMANN; SWEDIAUR; VAN MONS)

R. Leaves of Tobacco, one part.
Spanish White Wine, twelve parts.

Macerate for several days, and filter.

VAN MONS prescribes also two drams of tobacco and thirty ounces of wine.

It is powerfully stimulant and is recommended in dropsy.—Dose, from five to thirty drops.

TINCTURE OF TOBACCO. (AUGUSTIN; HUFELAND)

R. Leaves of Tobacco, two ounces.
Alcohol, one pound.
Infuse for three days, and strain. (HUFELAND)

AUGUSTIN prescribes one ounce of tobacco, and one pound of proof spirit.

It is diuretic.—Dose, from fifty to ninety drops, thrice a day.

DIURETIC MIXTURE.

Mixtura Nicotianæ diuretica. (AUGUSTIN)

R. Oxymel of Colchicum, }
Squill, } of each,
Tincture of Tobacco, } an equal part.
Nitric Æther,

Mix.—Dose, a tea-spoonful, four times a day, in dropsy.

VINEGAR OF TOBACCO.

Acetum Nicotianæ. (HERBIP.)

R. Dried Leaves of Tobacco, one ounce.
Cloves, six.
Vinegar, one pound.

Digest for four days, in a moderate heat, and strain.

OXYMEL OF TOBACCO.

Oxymel Tabaci. (HERBIP.)

R. Vinegar of Tobacco, one part.
Honey, two parts.

Boil slowly to the consistence of syrup.

OIL OF TOBACCO. (GALL.; WIRTEM.)

R. Fresh Leaves of Tobacco, one pound.
Olive Oil, sixteen ounces.

Boil to the consumption of the humidity, and express. (WIRTEM.)

GALL. directs one pound of tobacco to be

digested in two pounds of olive oil, in warm ashes for twenty-four hours, and the oil to be expressed; the maceration to be repeated with fresh leaves; then to boil slightly, and express.

It is recommended in itch, chilblains, paralysis, and indolent tumours.

TOBACCO LINIMENT.

Linimentum Tabaci. (AMER.; COXE)

R. Leaves of Tobacco, one ounce.
Lard, one pound.

Boil till the leaves become crisp, and strain.

TOBACCO OINTMENT.

Onguent anti-scrofuleux. (BRUNS.; SARD.; WIRTEM.; HERBIP.; AUGUSTIN; PIERQUIN; SPIELMANN; VAN MONS)

R. Leaves of Tobacco, three pounds.
Juice of Tobacco, nine ounces.
Lard, one pound and a half.

Macerate for twelve hours; then boil slowly to the consumption of the humidity; strain and add,

Resin, three ounces.
Yellow Wax, one ounce and a half.
Round Birthwort, in powder, three ounces.

Mix. (BRUNS.; SARD.; WIRTEM.; HERBIP.; PIERQUIN; SPIELMANN)

AUGUSTIN prescribes three ounces of the juice of tobacco, three of yellow wax, one ounce and a half of resin, one of turpentine, and a sufficient quantity of oil of myrrh.

R. Empyreumatic Oil of Tobacco, one part.
Resin, two parts.
Green Fecula of Tobacco, four parts.
Lard, twelve parts.

Dissolve the fecula in the lard over a slow fire, then add the resin, and at the moment of its becoming cold, incorporate the oil. (VAN MONS)

R. Green Fecula of Tobacco, one part.
Lard, five parts.

Dissolve. (VAN MONS)

It is recommended in scrofula, itch, tinea, and atonic ulcers.

DETERGENT OINTMENT.

Unguentum mundificans Poterii. (SPIELMANN)

R. Leaves of Tobacco, six handfuls.
Butter, twenty-four ounces.
Turpentine, four ounces.

Macerate for eight days; then add to the expressed liquid.

Gum Ammoniac, } of each, two ounces.
 Galbanum, }
 Sulphuret of Antimony, } of each,
 Red Sulphuret of Mer- } three drams.
 cury,
 Myrrh, } of each, two drams.
 Olibanum, }

Mix.—It is recommended in the cure of fistulous ulcers.

TOBACCO CERATE.

Ceratum Tabacinum. (NIEMANN)

R. Juice of Tobacco, } of each,
 Yellow Wax, } three ounces.
 Resin, one ounce and a half.
 Oil of Myrrh, a sufficient quantity.

Mix.—It is recommended by CONSRUCH in herpetic affections where there is neither inflammation nor pain.

TOBACCO PLASTER.

Emplastrum de Nicotiana. (SPIELMANN)

R. Leaves of Tobacco, four ounces.

Olive Oil, twelve ounces.
 Red Wine, five ounces.
 Boil to the consumption of the humidity,
 express, and add,
 Yellow Wax, } of each, twelve ounces.
 Resin, }

When liquified, add,

Powdered Tacamahac, }
 ——— Elemi, } of each,
 ——— Myrrh, } one ounce.
 ——— Aloes, }

Essential Oil of Tobacco, two ounces.

Mix.—TRILLER says that this plaster, applied to the abdomen, acts as a purgative.

ANTI-PSORIC OINTMENT. (FERR.)

R. Sulphur, } of each, two ounces.
 Tobacco, }
 White Hellebore, one ounce.
 Syrian Herb Mastic, } of each,
 Rose-bay, } half an ounce.
 Muriate of Soda, five drams.
 Rose Ointment, one pound.

Mix.

NIGELLA.

Three species of this genus of plants are met with in pharmacopœias :

1. *Nigella arvensis*, LINN.

SYNONIME :—*Nigelle des champs*, FR.

GENEV.

A common plant of Europe. (*Polyandria Pentaginia*, LINN.; *Ranunculacæ*, JUSS.; *Fig. Blackw. Herb. t. 126*)

The seed is employed; it is black, acrid, and aromatic.

2. *Nigella Damascena*, LINN.

SYNONIMES :—*Nigelle de Damas*, Barbiche, Barbe de Capucin, FR.

FERR.; BRUGNATELLI.

A plant of the South of Europe. (*Fig. Blackw. Herb. p. 558*)

The seed is employed; it is small, angular, black, and has a strong and aromatic odour.

3. *Nigella sativa*, LINN.—Small Garden Fennel Flower, Devil in a bush.

SYNONIMES :—*Nigelle des jardins*, Faux Cumin, FR.; *Zahmer shwarzkümmel*, GERM.; *Czerny neb zahradj kmjn*, BOHEM.; *Hangernes kornrose*, DAN.; *Neguilla sativa*, SPAN.; *Veldrigelle*, DUT.; *Nigella Romana*, ITAL.; *Nigellaordinaria* PORT.; *Swart kumin*, SWED.

BRUNS.; HISP.; GALL.; FERR.; LIPP.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant originally of the Levant, and common in the south of Europe. (*Fig. Zorn, Ic. pl. t. 119*)

The seed (*Semen Nigellæ seu Melanthii seu Cumini nigri*) is employed. It is small, oval or long, nearly triangular, obtuse at one end, rather compressed, wrinkled obliquely, black without, and greenish-white within. It has a strong and disagreeable smell, and an acrid and aromatic taste.

It is slightly stimulant, but is rather culinary than medicinal.

NITROGENIUM.—NITROGEN, AZOTE.

SYNONIMES :—*Gaz azoticum*.—Azote, Alcaligène, Nitrogène, Fr.

FERR. ; BRUGNATELLI ; SWEDIAUR.

R. Fresh Fibrine of Blood, washed,
one ounce.

Put it in a matrass, and pour upon it
Diluted Nitric Acid, two ounces.

Adapt a bent tube, apply heat, and receive the gas in a hydro-pneumatic apparatus. (BRUGNATELLI ; SWEDIAUR)

FERR. directs sulphuret of iron to be put under a bell glass, and, after some time, the gas to be received.

It has been recommended by some to be inspired in chronic diseases of the lungs.

OXYGENATED NITROGEN.

Gaz azoticum oxygenatum; *Gaz oxidé d'azote*. (BRUGNATELLI ; SWEDIAUR)

R. Nitrate of Ammonia, in powder,
six ounces.

Put it into a matrass to which is adapted a tube having a bladder at its extremity. Apply the flame of a spirit lamp to the bottom of the matrass, and receive the gas till white or red vapours appear.

This gas produces, when inspired, a kind of temporary delirium, with great propensity to muscular motion, from which latter property it has been recommended to be employed in paralysis.

NOSTOCHIUM COMMUNE, VAUCH.

SYNONIMES :—*Nostoc*, *Nostoch commun*, *Crachat de lune*, *Salive de Coucou*, Fr.

BRUNS.

An alga which appears on the ground in wet weather. It does not appear to have any root. (*Fig. Céd. Fl. Dan. t. 885. f. 1*)

It is composed of a frond, at first globular, then irregular, folded, undulated, lobed, and vesiculous. It is of an olive-green colour, but blackens as it becomes dry.

It is emollient.

NUX VOMICA.

SYNONIMES :—*Noix vomique*, Fr. ; Brechnuss, Krähenauge, GERM. ; Khanekulkeb, ARAB. ; Ma-tsin, CHIN. ; Kodak-addurualta, CYNG. ; Bræknøede, DAN. ; Kuchla, DUK. ; and HIND. ; Nuez vomica, SPAN. ; Breekroeten, DUT. ; Noce vomica, ITAL. ; Noz vomica, PORT. ; Culaka, Kutaka, Veschamustirijum, SANS. ; Røefkaka, SWED. ; Yettil cottay, TAM. ; Muscadi, Mustighenza, TEL.

DUBL. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FULD. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; WURCEB. ; AINSLIE ; BERGIUS ; BRUGNATELLI ; COKE ; GUIBOUT ; MURRAY and GMELIN ; PARIS ; SPIELMANN.

This is a flattish, button-shaped seed, of about half an inch in diameter and one or two lines thick, umbilicated in the centre, and of a horny consistence. Its colour is

grey, and it is silky on the surface ; it is inodorous, but has an acrid, bitter, and rather nauseous taste.

It comes from the *Strychnos Nux vomica*, LINN. ; a tree of the East Indies. (*Pentandria Monogynia* LINN. ; *Apocynæa*, JUSS. ; *Fig. Flore Medic. V. 249*)

According to PELLETIER and CAVENTOU, it contains strychnine and igasuric acid. Brucine has also been found in it.

It is one of the most powerful stimulants and poisons, exciting spasmodic and tetanic contraction. It is administered in palsy, in the dose of from four to fifteen grains.

POWDER OF NUX VOMICA.

Pulvis Nucis vomicæ. (AUGUSTIN)

R. Nux Vomica, three grains.
Gum Arabic, } of each,
White Sugar, } half a scruple.
2 M

Mix.—HUFELAND extols this powder in dysentery.—The dose to be given thrice a day.

STIMULANT PILLS.

Pilule nervosum et musculare systema vellantes. (BRERA; ELLIS)

R. Nux Vomica, half a dram.
Conserve of Roses, a sufficient quantity.

Make ten pills.—Dose, one, twice or thrice a day. (ELLIS)

R. Nux Vomica, from one to six grains.
White Sugar, half a scruple.
Rob of Elder, a sufficient quantity.

Make six pills.—Dose, one, every two hours. (BRERA)

They are prescribed in paralysis, epilepsy, chorea, hypochondria, hysteria, &c.

AQUEOUS EXTRACT OF NUX VOMICA.

Extractum Nucis vomice aquosum. (ANT.; BATAV.; BAVAR.; HASS.; SAX.; WIRTEM.; SWEDIAUR)

R. Nux Vomica, rasped, one pound.
Boiling Water, eight pounds.

Digest for twenty-four hours; then boil for a quarter of an hour, and strain: boil the residue for a quarter of an hour in four pounds of water, and strain again. Mix the two liquors, set aside, decant, and evaporate. (ANT.)

BAVAR.; HASS.; SAX.; WIRTEM.; NIEMANN; and SWEDIAUR give nearly the same process, requiring the nux vomica to be repeatedly boiled till the water ceases to acquire any bitterness; the decoctions to be then mixed together and evaporated.

Dose, from two to twenty grains, twice or thrice a day, augmenting it very gradually, in paralysis.

ALCOHOLIC EXTRACT OF NUX VOMICA.

Extractum Nucis vomice alcoholicum (DUBL.; BELG.; GALL.; FERR.; HANN.; SAX.; CADET DE GASSICOURT; MAGENDIE; PIERQUIN; VAN MONS)

Extractum Nucis Vomice. (DUBL.)

R. Nux Vomica, rasped, eight ounces.
Proof Spirit, two pints.

Digest for three days in a close vessel; then strain with strong expression; add to the residuum one pint and a half of proof spirit; digest for three days, and express again.

Mix the tinctures, distil till reduced to one fourth, and evaporate the residue, constantly stirring, to the proper consistence, with a medium heat obtained from steam. The process should be conducted so as to prevent as much as possible the contact of the air.

The softer extract should be sprinkled with rectified spirit.

GALL.; and FERR. direct one part of nux vomica to be digested in four parts of alcohol (22 degrees) for twenty-four hours in a covered vessel, and the expressed residue to be digested in three parts of alcohol (12 degrees); the expressed tinctures to be then distilled to three fourths and the remainder to be evaporated in a water-bath to the proper consistence.

R. Nux Vomica, rasped, any quantity.
Alcohol (22 degrees), a sufficient quantity.

Digest, renewing the alcohol from time to time till it ceases to acquire either colour or taste. Mix and filter the tinctures, reduce them one fifth by distillation, and evaporate the residue in a water-bath. (BELG.; GALL.; HANN.; SAX.; PIERQUIN; VAN MONS)

R. Nux Vomica, rasped, any quantity.

Exhaust it by alcohol (40 degrees) at the lowest possible temperature; then slowly evaporate the tincture to the proper consistence. (CADET DE GASSICOURT; MAGENDIE)

It is recommended in every species of paralysis.—Dose, one or two grains, in pills, increasing it, by one grain, every day till the desired effect is produced. The dose has in this way been carried to twenty-four or thirty grains daily before tetanic symptoms have supervened; but four or six grains are sufficient in the greater number of cases. This applies to the extract prepared with alcohol at 40 degrees, for that prepared with a weaker spirit, is not so strong, containing a great proportion of gummy matter.

HARD EXTRACT OF NUX VOMICA. (MAGENDIE)

R. Alcoholic Tincture of Nux Vomica, any quantity.
Filter it, and evaporate it in dishes.

The tincture employed should be as concentrated as possible, and prepared with alcohol at 36 degrees.

Dose, from one grain to ten grains progressively.

ANTHELMINTIC PILLS. (NIEMANN)

- R. Gum Ammoniac, }
 Soap, } of each, one scruple.
 Resin of Jalap, }
 Alcoholic Extract of Nux Vomica,
 half a scruple.

Essence of Amber,
 a sufficient quantity.

Make pills of one grain each.—Dose,
 twelve in the course of a week.

EXCITANT PILLS. (BORIES)

- R. Alcoholic Extract of Nux }
 Vomica, } of each,
 Conserve of Lime Tree } half a dram.
 Flowers,

Make pills of two grains each.—Dose,
 one every morning, fasting; encreasing one
 every week.

NIEMANN has successfully employed, in
 spasmodic cardialgia, [a very rare disease,
 EDIT.] pills of one grain, made with one
 part of extract and five parts of assafetida.

SOLUTION OF EXTRACT OF NUX VOMICA.

Emulsio cum Nuce vomica. (AUGUSTIN;
 NIEMANN)

- R. Extract of Nux Vomica, ten grains.
 Mucilage of Gum Arabic, one ounce.
 Spring Water, six ounces.

Mix.—It is extolled by HUFELAND, in
 dysentery.—Dose, two spoonfuls every two
 hours.—One ounce of syrup may be added.

ANTI-GONORRHOÏC MIXTURE.

*Decoctum Cucumerum cum Extracto Nucis
 vomica.* (AUGUSTIN)

- R. Extract of Nux Vomica, half a dram.
 Decoction of Cucumber, six ounces.
 Linseed Oil, triturated with Yolk of
 Egg, one ounce.
 Honey, half an ounce.

Mix.—Dose, two spoonfuls every two
 hours.—It is recommended by HANDEL, as
 an excellent remedy in every stage of go-
 norrhœa.

TINCTURE OF NUX VOMICA.

*Tinctura Strychnos seu Nucis vomica seu
 Nucum vomicarum.* (GALL.; FERR.;
 LIPP.; SAX.; BORIES.; ELLIS.; HUFEL-
 LAND.; MAGENDIE.; NIEMANN.; PIER-
 QUIN.; SWEDIAUR.; VAN MONS.)

- R. Nux Vomica, rasped, one part.
 Diluted Alcohol, eight parts.

Infuse without heat for three days; then
 having digested in a water-bath for six
 hours, express, and filter. (NIEMANN;
 VAN MONS)

SWEDIAUR prescribes one ounce and a
 half of nux vomica and two pounds of al-
 cohol (0.935).—SAX.—one part of nux vo-
 mica and five parts of proof spirit.—NIE-
 MANN, after WIEL—one of nux vomica
 and four of rectified spirit.—GALL.; and
 LIPP.—one of nux vomica and four of al-
 cohol (22 degrees).—HUFELAND—one of
 nux vomica and six of alcohol.

- R. Hard Extract of Nux Vomica,
 three grains.
 Alcohol (36 degrees), one ounce.

Dissolve. (FERR.; BORIES.; ELLIS;
 HENRI.; MAGENDIE.; PIERQUIN.)

This tincture was originally known under
 the title of *Gouttes uterines de la reine
 d'Espagne*.—That which VAN MONS gives
 is prescribed in the dose of from forty to
 eighty drops, and that of MAGENDIE also
 by drops. The indications are the same as
 for the alcoholic extract. It is also used
 externally, in liniments, in cases of par-
 alysis and partial atrophies.

BITTER DROPS. (RATIER)

- R. Spirit of Wormwood, two pounds.
 Nux Vomica, one pound.
 Solution of Subcarbonate of Potass,
 half an ounce.
 Soot, one dram.

Infuse for fifteen days in a water-bath;
 express, and filter.—Dose, from one drop
 to eight drops or more, in a glassful of
 bitter infusion; in flatulent colic.

NYPHÆA.—WATER LILY.

Two species of this genus of plants are
 used in medicine:

1. *Nymphæa alba*, LINN.—White Water
 Lily.

SYNONIMES:—Nénuphar blanc, FR.; Weisse
 Seerose, Seebume, Wasserlilie, Wasser-

rose, Haarwurz, GERM.; Seebloomster,
 DAN.; Nenufar blanco, SPAN.; Witte
 plompen, DUT.; Nenufaro bianco, ITAL.;
 Wodna lilia, POL.; Nymphæa branca,
 PORT.; Sjøblad, SWED.

BRUNS.; HISP.; GALL.; FERR.; GENEV.
 WIRTEM.; BERGIUS.; BRUGNATELLI,

GUIBOUT ; MURRAY and GMELIN ; SPIELMANN.

This plant is common throughout Europe. (*Polyandria Polygynia*, LINN.; *Nymphæa*, JUSS.; *Fig. Flore Medic. V. 247*)

The root and flowers are employed.

The root (*Radix Nymphææ albæ* seu *Nenupharis* seu *Nenufaris*) is very long, cylindrical, thick, nodose, fleshy, and of a spongy texture. It is brown without, and yellowish-white within; its taste is bitterish and rather styptic.

The flowers have an agreeable odour which is removed by desiccation. They contain much insipid mucilage.

The root contains a great deal of starch, and is very nutritious. The flowers are classed among the sedatives.

2. *Nymphæa lutea*, LINN.—Yellow Water Lily.

SYNONIME :—*Nénuphar jaune*, FR.

GALL.

A plant of Europe. (*Fig. Blackw. Herb. t. 491*)

The root resembles the preceding. The flowers are yellow.

COLD PEARL SPECIES.

Species diamargaritæ frigidaæ. (WIRTEM.)

R. Oriental Pearls,	six drams.	
White Coral,		} of each, two drams.
Red Coral,		
Flowers of Borage,		
White Water Lily,		
Red Roses,		
Kermes Grains,		
Aloe Wood,		
White Sanders,		
Yellow Sanders,		

Make a powder.—It used formerly to be recommended in phthisis and inconvalescence after malignant fevers.—Dose, from one to two scruples.

COLD SPECIES OF PRECIOUS STONES.

Species de Gemmis frigidaæ. (WIRTEM.)

R. Prepared Ivory,	one ounce.	
Oriental Pearls,	six drams.	
White Coral,		} of each, half an ounce.
Red Coral,		
Hyacinth,		} of each, three drams.
Emerald,		
Sapphire,		
Garnet,		
White Sanders,		
Yellow Sanders,		
Red Sanders,		

Flowers of Borage,	} of each, two drams.
Bugloss,	
Water Lily,	
Roses,	three drams.
Gold Leaves,	twenty.

Mix, and make a powder.—It used to be formerly administered in phthisis, to diminish nocturnal sweats.—Dose, from a scruple to half a dram.

CONSERVE OF WATER LILY. (SARD.)

R. Petals of Water Lily, bruised and reduced to a pulp with a little sugar,	one part.
Sugar,	three parts.
Apply heat, and mix.	

DISTILLED WATER OF WATER LILY. (GALL.; PALAT.; SARD.)

R. Flowers of Water Lily,	one part.
Water,	four parts.
Infuse for six hours, and distil one half.	

INFUSION OF WATER LILY.

Infusum Florum Nymphææ. (SARD.)

R. Fresh Flowers of Water Lily,	two pounds.
Boiling Water,	eight pounds.
Digest for eight hours, and express slightly; repeat the operation twice more with fresh flowers.	

SYRUP OF WATER LILY.

Syrupus Nymphææ seu de Floribus Nymphææ albæ. (HISP.; GALL.)

R. Fresh Petals of White Water Lily,	one pound.
Boiling Water,	four pounds.
Infuse for a quarter of an hour, and strain with slight expression; add to the liquor	
White Sugar,	six pounds.

Make a syrup. (HISP.)

GALL. directs four pounds of petals to be infused for twelve hours in eight of water; the liquor to be slightly expressed, set apart for a few hours, and then decanted; double the quantity of white sugar to be then added, and a syrup with the heat of a water-bath.

ANTI-SPASMODIC POTION. (PIERQUIN)

R. Almond Emulsion,	four ounces.
Syrup of Water Lily,	half an ounce.
Amber,	two drams.

Mix.

OCYUM.—BASIL.

Two species of this genus of plants are spoken of in pharmacopœias :

1. *Ocymum basilicum*, LINN.—Sweet Basil.

SYNONIMES :—*Basilic commun*, Grand Basilic, FR. ; *Basilienkraut*, *Königskraut*, *Herrnkraut*, GERM. ; *Basilic*, DAN. ; *Subzekebinge*, DUK. ; *Albahaca*, SPAN. ; *Kali tuls*, HIND. ; *Balsemkruid*, DUT. ; *Basilico*, ITAL. ; *Debanschab*, PERS. ; *Bazylico*, POL. ; *Alsavaca*, PORT. ; *Manjirika*, SANS. ; *Basilika*, SWED. ; *Tirnut patchie verie*, TAM. ; *Vepudipatsa vittilu*, TEL.

AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; HISP. ; GALL. ; FULD. ; HANN. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; WIRTEM. ; HERBIP. ; BERGIUS ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN.

A plant originally from the East. (*Didymia Gymnospermia*, LINN. ; *Labiata*, JUSS. ; *Fig. Zorn, Ic. pl. t. 26*)

The herb, flowers, and seeds are employed.

The herb (*Herba Basilici majoris seu citrati seu Ocymi vulgarioris seu medii seu magni*) consists of a slightly hairy stem, and of petiolated, oval, lanceolated leaves, somewhat ciliated on the borders, and

sparsely notched. It has a very agreeable odour and an aromatic taste.

The flowers are white, and are disposed in whirls.

The seeds are small, oblong, and black. They have an aromatic odour, but no taste.

It is stimulant.—Dose, two drams and a half, infused in two pounds of boiling water.

2. *Ocymum minimum*, LINN.

SYNONIME ;—*Basilic à petites feuilles*, FR. BAVAR.

A plant of the East Indies, cultivated in Europe. (*Fig. Schk. Bot. Hand. t. 166*)

The herb (*Herba Basilici minimi*) is employed. It consists of a branching stem, and of numerous acute, obtuse, or round leaves, rather thick, green or reddish, of a strong and agreeable odour, and of a very aromatic taste.

ESSENTIAL OIL OF SWEET BASIL.

R. Flowering Herb of Sweet Basil,

	five parts.
Water,	seven parts.

Distil, and separate the oil.

The distilled water is now frequently employed, in the dose of one or two ounces.

CESYPUS.—YELK OF WOOL.

SYNONIMES :—*Suint*, FR. ; *Hisopo humedo*, SPAN.

An oleagino-extractive substance of a fetid odour, with which the wool of sheep is impregnated.

It contains, according to VAUQUELIN, a fat animal matter and carbonate of lime,

with a little acetate and hydro-chlorate of potass.

It enters into several plasters.—HISP. directs it to be purified by boiling it in water in sufficient quantity to obtain a thick liquid, which is to be strained through a silk-sieve, and then evaporated to the consistence of soft extract.

OLEUM CAJEPUTI.—CAJEPUT OIL.

SYNONIMES :—*Oleum Cajuputi* seu *Cajaputi* seu *Cajeput*.—Huile de Cajeput, FR. ; *Cajaputöl*, GERM. ; *Cajaput olie*, DAN. ; *kyaputi ka tail*, DUK. ; *Cajaputa*, MAH. ; *Kayu putich*, MAL. ; *Kayaputi tayilam*, TAM.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; AINSLIE

BERGIUS ; COXE ; GUIBOUT ; MURRAY
and GMELIN ; PIDERIT ; SPIELMANN.

A transparent fluid oil, of a green colour,
lighter than water, and of a strong, aromatic,
and rather agreeable odour.

It is obtained by distilling the leaves of
the *Melaleuca Cajaputi*, MAT., a tree of the
East Indies. (*Polyadelphina Polyandria*,
LINN. ; *Myrtaceæ*, JUSS. ; *Fig. Rumph.*
Amb. II. p. 72. t. 6. and tab. 17. f. 1)

It has hitherto been attributed, and erroneously
according to AINSLIE, to the *Melaleuca leucodendron*,
LINN. It appears, however, that several species of
this genus, if not all, yield a similar oil.

It is excitant, nervine and diuretic.—Dose,
from two to twelve drops, on sugar.

OLEOSACCHARUM OF CAJEPUT. (SWEDIAUR)

R. White Sugar, one dram.
Cajeput Oil, two drops.
Triturate together.

STIMULANT DROPS.

Liquor Olei animalis et Cajeputi. (AUGUSTIN)

R. DIPPEL'S Animal Oil, one dram.
Cajeput Oil, half a dram.
Mix.—Dose, five, ten, or fifteen drops.

STIMULANT EMULSION.

Emulsio Cajeputi. (SWEDIAUR)

R. Thick Mucilage of Gum Arabic, three drams.
Cajeput Oil, one dram.
Water, four ounces.
Mix.

OLEUM OLIVARUM.—OLIVE OIL.

SYNONIMES:—Huile d'olive, FR. ; Baumöl,
Oliveneöl, GERM. ; Bomolie, DAN. ;
Aceyte comun, SPAN. ; Olyfoly, DUT. ;
Azeite, PORT. ; Bomolja, SWED.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ;
ANT. ; AUSTR. ; BATAV. ; BAVAR. ;
BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ;
FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ;
GENEV. ; HAMB. ; HANN. ; HASS. ;
LIPP. ; OLDEN. ; LUSTT. ; POL. ; BORUS. ;
ROSS. ; SAX. ; SUEC. ; WURCEB. ; BER-
GIUS ; COXE ; MURRAY and GMELIN ;
PARIS ; SAUNDERS ; SPIELMANN.

A fluid, fixed oil, nearly transparent, of
a yellowish or greenish colour, inodorous,
and of a bland taste. It is the lightest of
all fixed oils, and is the most esteemed for
alimentary uses. Its green colour is owing
to the presence of chlorophylle, its yellow,
to a vegetable principle, the presence of
which is manifested after exposure to the
air ; its odour is derived from another pecu-
liar principle.

It is obtained by expressing, boiling, or
fermenting the fruit of the olive tree, *Olea
Europæa*, LINN., a tree of the south of Eu-
rope. (*Diandria Monogynia*, LINN. ; *Jas-
mineæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 319*) It
appears to be originally from Asia Minor.

OILY POTION. (RATIER)

R. Gum potion, four ounces.
Olive Oil, half an ounce.
Mix.—For one dose.

OILY EMULSION.

Emulsio oleosa. (WURCEB.)

R. Olive Oil, one ounce.
Yolk of Egg, one.
Water, seven ounces.
Mix.

OILY LAVEMENT.

Clyisma oleosum. (BRERA ; SWEDIAUR)

R. Emollient Decoction, nine ounces.
Olive Oil, three ounces.
Honey, one ounce.

Mix.—Half an ounce of sulphate or of
muriate of soda may be added. (SWEDIAUR)

R. Decoction of Barley, } of each,
Olive Oil, } four ounces.
Mix. (BRERA)

BRICK OIL.

*Oleum Laterum seu de Lateribus seu Philo-
sophorum seu Olivarum empyreumaticum.*
(BRUNS. ; PALAT. ; SARD. ; WIRTEM.)

R. Bruised Brick, any quantity.

Impregnate it with olive oil, and distil from a glass retort, with the gradually raised heat of a sand-bath; separate the oil from the liquor which passes over with it,

and rectify it by re-distilling it with ashes.

It is stimulant, and was formerly reputed resolvent and antiseptic. It used also to be applied as a detergent to ulcers.

OLEUM PALMÆ.—PALM OIL.

SYNONIMES :—*Palmae Sebaceum*.—Huile de Palme, FR.

EDINB.; AMST.; BRUNS.; GALL.; SUEC.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A concrete oil, having the consistence of butter, of a golden-yellow colour, an agreeable violaceous odour, and very bland taste.

It is derived from the *Cocos butyracea*, LÄNN.; a tree of South America and of Africa (Pis. *Ind. res nat. et med.* p. 215), or, according to JACQUIN, of the *Elais Guineensis* (Fig. Jacq. *Stirp. Amer.* 360. t. 257)

It is not distinguished by any peculiar property from the other fixed oils, and it is now nearly disused.

OLIBANUM.

SYNONIMES :—Frankincense; *Thus*, *Thus verum*, *Gummi Olibani*; κέδρος.—Oliban, ENCENS, FR.; Weihrauch, GERM.; Luban, ARAB.; and MAL.; Kadidlo, BOHEM.; Virok, DAN.; Ayul cundur, DUK.; Inciense, SPAN.; Kunder zuchir, HIND.; Wierook, DUT.; Incenso, ITAL.; and PORT.; Kadzidlo biale, POL.; Wirach, SWED.; Labuniya, SYR.; Paringhi Sambrani, TAM.

LOND.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; LUST.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; BRUGNATELLI; COXE; GUBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SAUNDERS; SPIELMANN; ZARDA.

A gum-resin in oblong, irregular tears of the size of a pea and larger, isolated, agglomerated, or geminated, bearing in the latter cases the names of *Male Olibanum* and *Female Olibanum*, from their fancied resemblance to nipples and to testicles.

Olibanum is dry, fragile, semi-transparent, of a pale-yellow colour with an outer stratum of farinous powder. It softens in the mouth, and adheres to the teeth. Its odour is balsamic, particularly when burnt, and its taste is rather warm and bitter. The finest pieces are called *Olibanum electum*, and the impure, *Olibanum in sortis*.

We are not certain what plant furnishes this substance. It does not come from the *Juniperus lycia*, *J. phænicea*, or *J. thurifera*,

as many authors have supposed. COLBROOK says that it is the *Libanus thurifera*, a tree of the mountains of India. (*Decandria Monogynia*, LINN., *Terebenthaceae*, JUSS.). Others think it is a species of *Amyris*.

It is stimulant, and is esteemed diuretic and vulnerary.—Dose, half a dram.

It is more frequently employed in fumigations than administered internally.

COMPOUND POWDER OF OLIBANUM. (NIEMANN)

R. Olibanum, } of each, ten grains.
Orange Peel, }
Oak Bark, } five grains.

Pulverize and mix.—HEBERDEN extolled this in leucorrhœa.—Dose, one powder, every three hours, in concert with decoction of oak bark as drink and also in enemata.

The oak bark in this case is evidently the principal agent.

FUMIGATORY SPECIES.

Species Fumales pro Suffimento. (FULD.)

R. Olibanum, } of each, four ounces.
Juniper, }
Amber, } of each, one ounce.
Cascarilla, }
Flowers of Tansey, } two ounces.

Mix.—Recommended in asthenic rheumatism.

BALSAMIC PILLS. (AUGUSTIN; SAUNDERS)

R. Extract of Cascarilla, } two drams.

Extract of Liquorice, } of each,
Olibanum, } five drams.
Saffron, } four scruples.
Peruvian Balsam, } one scruple.

Make pills of two grains each. (AUGUSTIN)

R. Olibanum, } of each,
Gum Arabic, } two drams.
Extract of Liquorice, }
Peruvian Balsam, a sufficient quantity.

Make four pills. (SAUNDERS)

Dose, from four to eight, twice a day; in asthenic hæmoptysis.—They are recommended by REIL.

STIMULANT DRAUGHT.

Haustus ex Olibano. (SAUNDERS; SWEDIAUR)

R. Olibanum, } of each,
White Sugar, } half a dram.
Water, } ten drams.
Tincture of Orange Peel, } two drams.

Mix. (SAUNDERS)

R. Olibanum, } half a dram.
Honey, } a sufficient quantity.
Simple Alexiteric Water, } one ounce and a half.

Spirituous Water of }
Jamaica Pepper, } of each,
Syrup of Tolu, } two drams.

Mix. (SWEDIAUR)

To be taken twice a day in leucorrhœa.

ONISCUS ASELLUS, LINN.—WOOD-LOUSE, SLATER, MILLEPEDE.

SYNONIMES:—Cloporte, FR.; Kellerwurm, Kellersel, GERM.; Shuk ketrold, DAN.; Galminha Eucarrucha, SPAN.; Pissebedden, DUT.; Centogambo, Porcelletto, ITAL.; Stonog, POL.; Centopea, PORT.; Græsugga, SWED.

AMST.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAK.; WIRTEM.; AINS-LIE.; GUIBOURT.; SPIELMANN.

A crustaceous insect very common every where. (*Isopodes Cloportides*, LAT.; *Fig. Paaz. Faun. Germ.* fasc. IX. 21) It is smooth, and ash coloured, with yellowish and black spots.

Wood-lice are reputed deobstruent and diuretic. They were formerly much esteemed in dropsy and asthma.—Dose, of the expressed juice, half an ounce. FULD. requires those in *muris non sub trabibus collecti*.

CONSERVE OF WOOD-LICE.

Conserva Millepedum. (FULD.; LIPP.; WIRTEM.; NIEMANN)

R. Living Wood-lice, one pound.
White Sugar, two pounds and a half.
Triturate together. (LIPP.; WIRTEM.; NIEMANN)

FULD. prescribes one part of wood-lice and two parts of sugar.

Dose, one or two drams.

BROTH OF WOOD-LICE.

Jus Millepedum. (AUGUSTIN)

R. Wood-lice, one dram.
Aromatic Liquor, half an ounce.
Broth, nine ounces.

It is diuretic.

INFUSION OF WOOD-LICE.

Infusum Millepedum. (SARD.)

R. Living Wood-lice, bruised, six drams.
Water, three ounces.
Alcohol, one ounce.

Macerate for an hour, and express strongly.

SYRUP OF WOOD-LICE. (CADET DE GAS-SICOURT)

R. Sugar, twelve ounces.
Root of Asparagus, }
Liquorice, } of each,
Raisins, } two drams.
Leaves of Pellitory of the Wall, }
Manna, }

Make a syrup. Then

R. Juice of Borage, } of each,
Bugloss, } two ounces.
Bruised Wood-lice, } one ounce and a half.

Mix, filter, and add to four ounces of the liquor,

Sugar, half a pound.
Dissolve in a water-bath, and mix the syrup with the preceding.

It is recommended in hooping cough and in catarrhal cough.

DIURETIC WINE. (CADET DE GASSICOURT; SAINTE-MARIE; SPIELMANN)

R. Living Wood-lice, bruised, one ounce.
Acidulous White Wine, six ounces.

Triturate gradually together, and express.
(CADET DE GASSICOURT; SAINTE-MARIE)

SPIELMANN prescribes one ounce of wood-lice, one of sugar, and four ounces of wine.

DIURETIC LIQUOR. (PIERQUIN)

R. Wood-lice, from thirty to forty.
Alkekenge Berries,
from twenty to thirty.
Juniper Berries, six.

Bruise then in a stone mortar, and add gradually,

White Wine, a sufficient quantity.

Express.

TINCTURE OF WOOD-LICE. (BRUNS.)

R. Dried Wood-lice, half an ounce.
Rectified Spirit, six ounces.
Digest, express, and filter.

ONOPORDON ACANTHIUM, LINN.—COTTON THISTLE.

SYNONIMES:—Chardon aux Anes, FR.; Breite Wegdistel, GERM.; Witte Wegdistel, DUT.

BRUNS.; GALL.; HASS.; ROSS.; WIRTEM.; MURRAY and GMELIN.; SPIELMANN.; FIDERIT.

A common plant of Europe. (*Syngenesia Polygamia Equalis*, LINN.; *Synantherea*, CASS. *Fig. Zorn, Ic. pl. t. 359*)

The herb (*Herba Cardui tomentosi* seu *Acanthii vulgaris* seu *Spinæ albæ*) is employed. It consists of sessile, oval, long, dentated spinous leaves. It is destitute of smell, but has a bitter taste.

Its juice is esteemed useful in cancer of the face. The lint which is applied should be dipped in it.

EXTRACT OF COTTON THISTLE.

Extractum Cardui tomentosi. (FULD.; LIPP.)

R. Herb of Cotton Thistle, any quantity.

Bruise in a wooden or stone mortar, enclose the pulp in a linen bag, express the juice, and evaporate it, when depurated, in a water-bath to the consistence of an extract.

Dose, from ten to twenty grains.

ONOSMA ECHIOIDES, LINN.—YELLOW ALKANET.

SYNONIMES:—Orcanette jaune, FR.; Natertkopfartige Lotwurzel, GERM.; Wołowcy gazyk Cerweny, BOHEM.

GALL.; BRUGNATELLI.; ZARDA.

A plant of the south of Europe. (*Pen-*

tandria Monogynia, LINN.; *Borragineæ*, JUSS.; *Fig. Jacq. Austr. t. 295*)

The root is employed; the bark of which contains a yellow colouring principle.

It is slightly astringent.

OPHIOGLOSSUM VULGATUM, LINN.—ADDERSTONGUE.

SYNONIMES:—Ophioglosse, Langue de Serpent, FR.; Schlangenzunge, GERM.; Slangetunge, DAN.; Lingua de Sierpe, SPAN.; Natertong, DUT.; Lingua di serpente, ITAL.; Lingua de serpente, PORT.; Læke tunga, SWED.

VOL. II.

GALL.; WIRTEM.; SPIELMANN.

This plant is found all over Europe. (*Cryptogamia*, LINN.; *Filices*, JUSS.; *Fig. Ed. Fl. Dan. t. 147*)

The herb (*Herba Ophioglossi* seu *Echio-glossi* seu *Lingua serpentaria*) is employed. It consists of amplexicaul, oval, obtuse, very entire, glabrous fronds, without nervures.

It is without odour, and has a mucilaginous and sweetish taste.

It was esteemed vulnerary.

OPHIORRHIZA MUNGOS, LINN.—INDIAN SNAKE-ROOT.

SYNONIMES :—Mungos, FR. ; Indische Schlangenzur, GERM. ; Mendi, Wal eka-Weriya, CYNG. ; Slangerod, DAN. ; Raiz de Serpiente, SPAN. ; Ostindische schlangenzwortel, DUT. ; Radice di serpe, ITAL. ; Kajo ular, JAV. ; Colubrina, PORT. ; Ormot, SWED.

BRUNS. ; HISP. ; WIRTEM. ; AINSLIE ; MURRAY and GMELIN ; SPIELMANN.

A plant of the Indies. (*Pentandria Mo-*

nogynia, LINN. ; *Gentianeæ*, JUSS. ; *Fig. Plenk. Pl. med. t. 90*)

The root (*Radix Mungos* seu *Serpentum*) is employed. It is simple, contorted, and channelled ; the bark is wrinkled and porous, inodorous, and very bitter.

It used formerly to be esteemed a specific against the bites of venomous animals, and consequently as alexipharmic.

OPIUM.

SYNONIMES :—*Opium thebaicum*, *Meconium*.—Opium, FR. ; and POL. ; Mohnsaft, GERM. ; Ufiun, ARAB. ; and MAL. ; Hapium, BALL. ; Abim, CYNG. ; Afim, DUK. ; Opio, SPAN. ; ITAL. ; and PORT. ; Ufim, HIND. ; Turk Heulzop, DUT. ; Apium, JAV. ; Caruppa, MAL. ; Schirik haskasch, PERS. ; Apaynum, SANS. ; Apini, TAM. ; and TEL. ; Madjun, TURK.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; MIL. COPEN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; CAST. BORUS. ; ROSS. ; SAX. ; SARD. ; SUEC. ; WIRTEM. ; HERBIP. ; WURCEB. ; AINSLIE ; BERGIUS ; COXE ; MURRAY and GMELIN ; PARIS ; PIDERIT ; SAUNDERS ; SPIELMANN.

Opium is the concrete juice that has flowed from incisions made in the capsule of the poppy before it is ripe. There are two kinds of it known in commerce, the Turkish and the Indian. They are easily distinguished from each other ; the former is in cakes of a flat form, homogeneous, dry, compact, and heavy ; opaque, of a deep brown colour, and, when broken, shewing a bright fracture ; it has a nauseous, bitter, and pungent taste, and a very unpleasant odour peculiar to itself. The latter is much less consistent, it is always ductile, and sometimes is met with as soft as pitch ; its co-

lour is darker, its taste more nauseous but less bitter, and its odour is slightly empyreumatic. This is not so much esteemed as the former, and, of course, it is considerably cheaper.

Although many able chemists have made the composition of opium the subject of their researches for some years past, it appears that this is not yet perfectly known. It is very probable that there are several new principles in it which remain to be discovered. The most important of them that we are already acquainted with, are *Narcotine*, and an alkali discovered by SERTURNER, *Morphine*, which is found combined with an acid, equally peculiar to opium, called *Meconic Acid*.

The *modus operandi* of opium has long been a source of dispute ; by some it is represented to be a powerful stimulant, by others, a direct sedative that never stimulates at all. The fact is, that in a certain dose, variable according to idiosyncrasy, it acts as an excitant ; in a stronger dose, as a sedative ; and, in a still stronger, as a poison. But whether it acts by producing a simple revulsion, or by a direct action on the nervous system, remains yet a problem. The Persians, Turks, Syrians, Chinese, and Malays, who take great quantities of it at once, are affected in the same manner as we find the drinkers of ardent spirits are.

Opium, being in many instances adulterated with foreign substances, several pharmacopœias, LOND. ; DUBL. ; GALL. ; AMST. ;

and SWEDIAUR direct it to be purified.—LOND.; AMST.; and GALL. direct it to be softened with boiling water, the mass to be strained through linen, and then evaporated to the required consistence.—DUBL.; and SWEDIAUR direct it to be dissolved in proof spirit, the solution to be filtered, the spirit to be distilled off, and the residue evaporated to a proper consistence. The Crude Opium, *Opium crudum*, having undergone this operation, is then called Purified Opium, *Opium depuratum seu colatum*.

I. Preparations which contain only a part of the constituent principles of Opium, with or without the menstruum employed to extract it.

A.

Extraction by Distillation.

WATER OF OPIUM.

Aqua Opii. (SAX.; NIEMANN; VAN MONS)

R. Crude Opium, one part.
Water, two parts.
Distil one half of the liquid. (SAX.; NIEMANN)

VAN MONS directs six parts to be distilled from a mixture of one part of opium and twelve parts of water.—VAN MONS also directs opium to be dissolved in six parts of water, and distilled to dryness.

It is not yet known what the chemical composition of this distilled water is, but it is conjectured to contain what is called the virous principle of opium, and is said by some authors to possess powerful narcotic virtues; but it is probably not of much value.

Dose, from two to six drams, with simple syrup.

B.

Extraction by Acetic Acid.

VINEGAR OF OPIUM.

Acetum Opii. (DUBL.)

R. Turkish Opium, four ounces.
Distilled Vinegar, one pound.

Triturate the opium to a pulp with part of the vinegar; then add the remainder, and macerate in a stopped bottle for seven days, frequently shaking. Pour off the clear liquor and filter it.

ACETOSE TINCTURE OF OPIUM.

Acetum Opii. (VAN MONS)

R. Opium, sliced, one part.
Rectified Spirit, six parts.

Macerate for twenty-four hours, in a water-bath, then add,

White Wine Vinegar, six parts.

Put it in a cool place and let it macerate for several days; then express and filter.

BLACK DROP.

Essence noire Anglaise. (AMER.; COXE; FEE; NIEMANN; PARIS; VAN MONS)

R. Opium, half a pound.
White Wine Vinegar, three pounds.
Nutmegs, bruised, one ounce and a half.
Saffron, half an ounce.

Boil till it has acquired a proper consistence, and add,

White Sugar, four ounces.
Yeast of Beer, one ounce.

Having digested it for seven weeks, expose it to the air till it becomes of the consistence of syrup; then strain and put it into bottles with the addition of a little sugar. (AMER.; COXE; FEE; NIEMANN)

VAN MONS directs six parts of opium to be mixed with three of white sugar, one part and a half of nutmegs, one of saffron, and forty-eight parts of verjuice; one part of yeast of beer to be then added, and the whole put into a shallow open vessel and left exposed to the action of a dry air till the supernatant liquor shall have acquired the consistence of a syrup: it is then to be decanted and preserved.

This preparation probably owes its activity to the acetate of morphine which is formed during the operation.

We find a preparation somewhat similar to this in the formulary of the republic of Genoa (1791) under the name of *Ethiops Vegetal*. It is directed there to take any quantity of opium, *crucibulo vitreato immittatur: sub agitatione adspargatur pluries vinum album; refrigefactum et pulveratum comburatur, simulque densius huncetatur vino; ad dimidium pondus reductum, addatur acetum acerrimum, dein siccatur massa*. In this preparation morphine would be separated from the opium, and would unite with the acetic acid.

ACETOSE EXTRACT OF OPIUM. (PIERQUIN)

R. Opium, sliced, one ounce.
Distilled Vinegar, one pound.

Digest for forty-eight hours, in a sand-bath, shaking it from time to time; then decant, filter, and evaporate to a phular consistence.

LALOUETTE'S LAUDANUM. (PIERQUIN)

- R. Acetose Extract of Opium, two drams.
 Spanish Wine, ten ounces.
 Proof Spirit, two ounces.
 One dram of the solution, contains four grains of the extract.

SYLVIVS' MIXTURE. (PIERQUIN)

- R. Opium, two grains.
 Distilled Vinegar, half an ounce.
 Plantain Water, six ounces.
 Syrup of Poppies, one ounce.
 Mix.—It is recommended in hæmoptysis with spasm.—Dose, one spoonful.

ANTI-ODONTALGIC GARGLE. (NIEMANN)

- R. Root of Sneezewort, two drams.
 Muriate of Ammonia, one dram.
 Opium, two grains.
 Distilled Water of Sage, } of each,
 Vinegar, } two ounces.

Digest for an hour in a water-bath, and strain.—It is proposed by MAYR.

SEDATIVE APPLICATION. (BORIES)

- R. Leaves of Henbane, } of each,
 Petals of Red Poppies, } one ounce.
 Opium, dissolved in a sufficient quantity of Vinegar, six grains.

Bruise together to form a paste.

It is recommended to be applied to the forehead in cases of violent pains of the head.

C.

Extraction by Wine.

VINOUS TINCTURE OF OPIUM.

Tinctura Opii vinosa seu simplex seu thebaica genuina. (AMST.; DAN.; HANN.; HASS.; LIPP.; SUEC.; PIDERIT; SWEDIAUR; VAN MONS)

- R. Opium, one part.
 Spanish Wine, twelve parts.

Digest for three days, and filter. (DAN.)

VAN MONS prescribes Madeira wine and one month's infusion.—HASS.; PIDERIT; and SWEDIAUR—one of opium, eight of Spanish wine, and one or two days' digestion at a moderate heat.—SUEC.—one of opium, six of white wine, and eight days' maceration without heat.

- R. Opium, one part.
 Malaga Wine, eight parts.
 Alcohol (0.907), two parts.

Digest for two days, and strain. (AMST.; BELG.; HANN.)

LIPP. prescribes two ounces of opium,

four of proof spirit, and one pound of white wine.

OPIACEOUS WINE. (FORM. HÔP. FR.)

- R. Tincture of Opium, ten grains.
 Red Wine, one ounce.
 Mix.

SEDATIVE FOMENTATION. (PIERQUIN)

- R. Opium, one ounce.
 Red or White Wine, one pound and a half.

Boil down to one half.—It is a useful application in gout, rheumatism, and colic. It is to be applied by means of flannel.

VINOUS EXTRACT OF OPIUM.

Extractum Opii vinosum, Laudanum opiatum. (GALL.; HASS.; PIDERIT; VAN MONS)

- R. Opium, any quantity.
 White Wine, a sufficient quantity.

Digest in a water-bath, express strongly, set aside, decant, and evaporate in a water-bath. (GALL.)

HASS.; and PIDERIT fix the quantity of wine to twelve times the weight of the opium.

- R. Opium, finely sliced, one part.
 Madeira Wine, six parts.

Macerate without heat for several days, and express; repeat the maceration with an equal quantity of wine: then mix the strained liquors, and evaporate slowly, constantly stirring. (VAN MONS)

This contains more narcotine and resin than the aqueous extract, and it also contains all the elements of the wine except the alcohol, which renders it still worse; for the quantity of these matters varying according to the kind of wine, and being always unknown, it is impossible to ascertain the exact quantity of opium which enters into a given dose of the extract.

VINOUS SYRUP OF OPIUM. (POL.; BORUS.; SWEDIAUR; VAN MONS)

- R. Opium, one part.
 Madeira Wine, twelve parts.

Infuse for two days, in a moderate heat; then express strongly, and add

White Sugar, twenty parts.

Dissolve in a very gentle heat, and mix the syrup with

Simple Syrup, one hundred and sixty parts.

Preserve. (VAN MONS)

R. Extract of Opium, twenty-five grains.
Malaga Wine, one ounce.
Dissolve, and add
Syrup of Liquorice,
twenty-four ounces.

Preserve. (BORUS.)

POL. prescribes syrup of marshmallows instead of that of liquorice.—SWEDIAUR—simple syrup.

AROMATIC VINOUS TINCTURE OF OPIUM.

Vinum Opii. (LOND.; EDINB.; AMER.;
BRUGNATELLI; COXE)

R. Extract of Opium, one ounce.
Cinnamon, bruised, } of each,
Cloves, bruised, } one dram.
Proof Spirit, six fluid ounces.
Water, ten fluid ounces.

Macerate for eight days, and filter.
(LOND.; COXE)

R. Opium, one ounce.
Cinnamon, bruised, } of each,
Cloves, bruised, } one dram.
Weaker Alcohol, six ounces.
Water, ten ounces.

Macerate for seven days, and filter.
(EDINB.)

AMER. prescribes two ounces of opium.—
BRUNS.—two ounces of opium, one dram
each, of cinnamon and of cloves, and one
pound of wine.

R. Extract of Opium, in powder,
two ounces.
Cinnamon Water, ten ounces.
Alcohol, two ounces.
White Wine, four ounces.

Digest for six days, occasionally shaking
the vessel, and filter. (BRUGNATELLI)

CROCATED VINOUS TINCTURE OF OPIUM.

Laudanum liquidum seu crocatum seu compositum. (FULD.; ROSS.; SAX.)

R. Opium, in powder, one part.
Saffron, one half part.
Spanish Wine, nine parts.

Macerate for three days, in a moderate
heat, and strain.—One dram contains about
ten grains of opium.

ESSENCE OF POPPY.

Vinum Opii crocatum, Elixir Papaveris.
(FENN.; SUEC.; VAN MONS)

R. Opium, } of each, five drams.
Saffron, }
Extract of Liqueurice,
one ounce and a half.
Madeira Wine, thirty ounces.

Digest without heat; decant, and filter.
(VAN MONS)

FENN. prescribes five scruples of opium,
the same of saffron, half an ounce of extract
of liquorice, and ten ounces of Spanish
wine—SUEC.—five scruples each, of extract
of liquorice, of saffron, and of opium, and
ten ounces of wine.

SYDENHAM'S LAUDANUM.

*Tinctura laudanum seu anodyna, Laudanum
liquidum Sydenhami seu compositum, Vi-
num crocatum opiatum seu Opii compositum,
Tinctura Opii crocata seu vinosa Opii
Sydenhami, Vinum Opii aromaticum seu
paregoricum.* (AMST.; BATAV.; DAN.;
HISP.; GALL.; FERR.; HANN.; LIPP.;
POL.; SARD.; CADET DE GASSICOURT;
SWEDIAUR; VAN MONS)

R. Opium, one ounce.
Saffron, half an ounce.
Cinnamon, one dram.
Malaga Wine, nine ounces.

Digest for three days, and strain. (LIPP.)
One dram contains about ten grains of
opium.

R. Opium, two ounces.
Saffron, one ounce.
Cinnamon, two drams.
Alcohol, three ounces.
Spanish Wine, one pound.

Digest for several days, express, and
filter. (FERR.)

R. Opium, sliced, two ounces.
Saffron, one ounce.
Cinnamon, } of each, one dram.
Cloves, }
Malaga Wine, one pound.

Digest in a moderate heat for fifteen days,
occasionally shaking the bottle; then ex-
press strongly, and filter. (AMST.; BA-
TAV.; BAVAR.; BRUNS.; HISP.; GALL.;
PALAT.; SAX.; WIRTEM.; CADET DE
GASSICOURT; SPIELMANN; SWEDIAUR)

DAN. prescribes but half an ounce of
saffron.—HANN.; POL.; and BORUS. pre-
scribe six drams of it; in other respects
their formula is the same.

SARD. prescribes half an ounce of opium,
two drams of saffron, half a dram each, of
cinnamon and of cloves, and four ounces
and a half of Madeira wine.

This last formula contains a greater pro-
portion of opium than the others, twenty
drops or fifteen grains of which contain one
grain of opium.

R. Cinnamon, } of each, one part.
Cloves, }
Saffron, eight parts.
Madeira Wine,
one hundred and ninety-two parts.

Infuse for twenty-four hours in a warm water-bath; then infuse, without heat, in the expressed liquor

Opium, sixteen parts.

Decant at the end of ten or fifteen days, without having shaken the vessel during the infusion, and filter. (VAN MONS)

R. Saffron, half an ounce.
Cinnamon, } of each, one dram.
Cloves, }
Spanish Wine, eight ounces.

Digest for three days, with occasional agitation; then dissolve in the strained liquor

Depurated Opium, one ounce.

And filter after twenty-four hours' digestion. Pour upon the residue

Spanish Wine, four ounces.

Digest, express strongly, and filter; then mix this with the preceding. (ANT.; BELG.)

Every one knows the celebrity which this preparation has obtained. It is curious to observe what the inventor thought of it. These are the words of SYDENHAM: *Hanc nostram preparationem laudani officinarum solido anteferebam virtutibus quidem non obcesco, sed formam saltem commodiorem et majorem dosis certitudinem eidem antepono.* All the inventors of new preparations have not had the modesty of this English physician [*modeste bonne foi du medecin Anglais*].

ANODYNE CATAPLASM. (RATIER)

R. Linseed Meal, } of each, one pound.
Barley Meal, }
Decoction of Poppy Heads, a sufficient quantity.

Boil to the required consistence, and add to the cataplasm,

SYDENHAM'S Laudanum, one or two drams.

Mix.—It is recommended in phlegmasiæ accompanied by acute pain.

ANTI-CANCEROUS CATAPLASM.

Cataplasma ad Cancrum. (FULD.; VAN MONS)

R. Pulp of Carrots, twenty-four parts.
Green Fecula of Hemlock, four parts.
SYDENHAM'S Laudanum, one part.
Beat together in a mortar. (VAN MONS)

R. Rob of Carrot, one pound.
Powdered Hemlock, two ounces.
SYDENHAM'S Laudanum, three drams.

Mix well. (FULD.)

It is recommended to correct the fetor and allay the pain of ulcerated cancer.

SEDATIVE LAVEMENT. (PH. SYPH.)

R. SYDENHAM'S Laudanum, from forty to sixty drops.
Olive or Linseed Oil, four ounces.
Mix by agitation.

OPIATE LINIMENT. (DISP. CHA. PARIS; RATIER)

R. SYDENHAM'S Laudanum, one part.
Olive Oil, eight parts.
Mix.

DISP. CHA. PARIS prescribes one part of laudanum and sixteen parts of oil of almonds.

OPIATE CERATE. (RATIER)

R. SYDENHAM'S Laudanum, one ounce.
GALEN'S Cerate, one pound.
Mix.

OPIATE OINTMENT. (SAINTE-MARIE)

R. SYDENHAM'S Laudanum, from two to four drams.
GALEN'S Cerate, two ounces.
Wheat Flour, a sufficient quantity
to thicken the mixture.

This ointment is applied on fine linen to burns.—A very concentrated aqueous infusion of opium would be better to be employed than the laudanum.

NARCOTIC LINIMENT. (RATIER)

R. SYDENHAM'S Laudanum, one part.
Balsamum Tranquillum, thirty-two parts.
Mix well.

R. SYDENHAM'S Laudanum, one part.
Balsamum Tranquillum, four parts.
FIORAVENTI'S Balsam, eight parts.
Mix.

ANTI-HEMORRHOIDAL LINIMENT. (PIERQUIN; SAINTE-MARIE)

R. SYDENHAM'S Laudanum, four drams.
Unguentum Populeum, two ounces.
Yolks of Eggs, two.
Triturate together. (SAINTE-MARIE)

R. Unguentum Populeum, half an ounce.
Opium, dissolved in White Wine, fifteen grains.
Prepared Oyster Shells, half a dram.
Yolk of Egg, one.
Oil of Lily, a sufficient quantity.

Mix well. (PIERQUIN)

It is recommended to be applied to painful piles.

ANTI-OPHTHALMIC OINTMENT. (PIERQUIN; SAINTE-MARIE)

- R. SYDENHAM's Laudanum, one dram.
 White of Eggs, two.
 Rose Water, } of each,
 Plantain Water, } a sufficient quantity.
 Shake well together.

CAMPHORATED OPIATE LINIMENT. (RATIER)

- R. SYDENHAM's Laudanum, half an ounce.
 Camphor, two drams.
 Oil of Almonds, two ounces.
 Mix.

ANODYNE LINIMENT. (FULD.; CADET DE GASSICOURT; PIERQUIN)

- R. SYDENHAM's Laudanum, } of each,
 Camphor, } one part.
 Ointment of Marshmallows, sixteen parts.
 Triturate together.

ANTI-SPASMODIC LINIMENT. (HAMB.)

- R. SYDENHAM's Laudanum, } of each,
 Tincture of Cantharides, } two drams.
 Camphor, one dram.
 Oil of Henbane, one ounce.
 Mix.

SAPONACEOUS OPIATE LINIMENT. (FULD.; RATIER)

- R. SYDENHAM's Laudanum, one or two drams.
 White Soap, half an ounce.
 Olive Oil, four ounces.
 Mix. (RATIER)
 R. Balsam of Soap, two ounces.
 SYDENHAM's Laudanum, one dram.
 Mix. (FULD.)

It is recommended as being at the same time sedative and resolvent.

SEDATIVE PECTORAL PILLS. (PIERQUIN)

- R. SYDENHAM's Laudanum, twenty drops.
 Gum Ammoniac, one scruple.
 Make four pills.—Dose, two or more.

NARCOTIC PILLS. (PIERQUIN)

- R. Sugar Candy, one ounce.
 Cinnamon, two drams.
 Pepper, } of each,
 SYDENHAM's Laudanum, } one dram.
 Coriander, two scruples.
 Saffron, half a dram.
 Musk, half a scruple.
 Syrup of Poppies, a sufficient quantity.

Mix.—Dose, from half a scruple to half a dram.—They are reputed sedative and sudorific.

SEDATIVE POTION. (BORIES; PIERQUIN; RATIER)

- R. SYDENHAM's Laudanum, twelve drops.
 Water, one ounce.
 Syrup, two drams.
 Mix.—For one dose. (RATIER)
 R. Gum Julep, four ounces.
 SYDENHAM's Laudanum, half a dram.
 Mix. (RATIER)
 R. Syrup of Lemon, one ounce.
 SYDENHAM's Laudanum, fifteen or twenty drops.
 Aqueous Tincture of Ipecacuanha, one or two spoonfuls.
 Orange-flower Water, two ounces.
 Mix. (BORIES; PIERQUIN)

SEDATIVE LAVEMENT. (PH. SYPH.)

- R. Starch, one dram and a half.
 Add gradually, with constant trituration,
 Boiling Water, half a pound.
 Boil slightly, and add to the strained liquor
 SYDENHAM's Laudanum, one dram.
 Mix.

SEDATIVE INJECTION. (RATIER)

- R. SYDENHAM's Laudanum, one dram.
 Decoction of Linseed, one pound.
 Mix.

ANODYNE MIXTURE. (GENEV.; BORIES; PIERQUIN; SAINTE-MARIE)

- R. SYDENHAM's Laudanum, two scruples.
 Cinnamon Water, six ounces.
 Mix. (PIERQUIN; SAINTE-MARIE)
 R. SYDENHAM's Laudanum, one dram.
 Hordeated Cinnamon Water, six ounces.
 Syrup of Poppies, one ounce.
 Mix. (BORIES)

BARTHEZ prescribed it in spasmodic cardialgia.—Dose, one spoonful before meals.

- R. SYDENHAM's Laudanum, eighteen drops.
 Aperient Powder, three grains.
 Spring Water, one ounce.
 Syrup of Lemon, one dram.
 Mix. (GENEV.)

OPIATE MIXTURE. (SAUNDERS)

- R. SYDENHAM's Laudanum, half a dram.

Peppermint Water, six ounces.
Simple Syrup, one ounce.

Mix.—To be taken by spoonfuls.

PAREGORIC POTION. (SAUNDERS)

R. SYDENHAM's Laudanum, twelve drops.
Water of Red Poppies, one ounce.
Syrup of Poppies, one ounce and a half.

Mix.—To be taken at two or three times.

ASTRINGENT TONIC POTION.

Haustus tonicus adstringens. (BRERA)

R. Calumba, two drams.
Boiling Water, a sufficient quantity
to obtain eight ounces of infusion. Add to
the strained liquor

SYDENHAM's Laudanum, half a dram.
Sulphuric Æther, twenty drops.

Mix.—To be taken gradually; in dyspepsia.

R. Cascarilla, } of each, two drams.
Calumba, }
Boiling Water, a sufficient quantity
to obtain seven ounces of strained infusion.
Add to this

SYDENHAM's Laudanum, } of each,
Sulphuric Æther, } twenty drops.

Mix.—To be taken gradually; in chronic diarrhoea.

ANTI-HÆMORRHAGIC POTION. (SAINTE-MARIE)

R. SYDENHAM's Laudanum, fifteen drops.
Tincture of Cinnamon, four drams.
Syrup of Comfrey, two ounces.
Water, four ounces.

Mix.—It is recommended in uterine hæmorrhage arising from debility.—Dose, a spoonful every quarter of an hour, every hour, or every two hours, according to the case.

HYPAROTIC POTION. (PIERQUIN)

R. SYDENHAM's Laudanum, five drops.
Syrup of Water Lily, } of each,
Lime-flower Water, } two ounces.
Orange-flower Water, }

Mix.—To be taken at twice.

SEDATIVE POTION.

Aqua aromatica cum Opio. (AUGUSTIN;
PIERQUIN)

R. SYDENHAM's Laudanum, twelve drops.
Syrup of Poppies, one ounce.

Lettuce Water, one ounce and a half.
Orange-flower Water, half an ounce.

Mix.—To be taken at twice. (PIERQUIN)

R. Cinnamon Water, one ounce.
Balm Water, two ounces.
Tincture of Castor, twenty drops.
SYDENHAM's Laudanum, ten drops.
Syrup of Opium, half an ounce.

Mix.—To be taken by spoonfuls. (AUGUSTIN)

CARMINATIVE POTION. (SAINTE-MARIE)

R. SYDENHAM's Laudanum, ten drops.
Cinnamon Water, one ounce.
Almond Emulsion, six ounces.

Mix.—It is recommended by DE HAEN in flatulent colic.—Dose, a spoonful every quarter of an hour.

SEDATIVE LOHOCH. (RATIER)

R. SYDENHAM's Laudanum, from fifteen to thirty drops.
Orange-flower Water, two drams.
Simple Lohoch, five ounces.

Mix.—To be taken in divided doses, or in one dose.

COUGH LINCTUS. (SAUNDERS)

R. SYDENHAM's Laudanum, half a dram.
Oil of Almonds, one ounce.
Yolk of Egg, two drams.

Triturate together, in a glass mortar, and add

Syrup of Marshmallows, three ounces.

Mix.—To be taken by spoonfuls.

PECTORAL POTION.

Mixtura sedativo-narcotica pectoralis. (BRERA)

R. Almond Emulsion, six ounces.
SYDENHAM's Laudanum, fifteen drops.
Syrup of Poppies, } of each,
Marshmallows, } one ounce.

Mix.—To be taken gradually, in whooping cough.

NARCOTIC EMULSION. (PIERQUIN)

R. Four Major Cold Seeds, one ounce.
Bruise them, and add gradually
Spring Water, six ounces.

Add to the emulsion

Syrup of Poppies, from four to six drams.
SYDENHAM's Laudanum, from ten to twelve drops.

Mix.—To be taken in the evening.

ANTI-PERIODIC POTION. (AUGUSTIN ;
PIERQUIN)

- R. SYDENHAM's Laudanum, } of each,
Sulphuric Æther, } twelve
Orange-flower Water, } drops.
Lime-flower Water, } half an ounce.

Mix.—We are told that this potion will prevent the coming on of a paroxysm in intermittent fever. (PIERQUIN)

- R. SYDENHAM's Laudanum, fifteen drops.
Tincture of Ipecacuanha, forty drops.
Solution of Acetate of Ammonia, half an ounce.
Rose Water, one ounce.
White Sugar, two drams.

Mix.—Dose, a tea spoonful.—WHYTT extolled it in periodical pains of the head, occasioned by disordered digestion, or the suppression of some hæmorrhage.

ANTI-LETHARGIC MIXTURE. (CADET DE
GASSICOURT ; PIERQUIN)

- R. SYDENHAM's Laudanum, half dram.
Sulphuric Æther, six drams.
Peppermint Water, six ounces.

Mix.—A tea spoonful to be taken every quarter of an hour.

Under the name of *Mixtura opiato-ætherea*, AUGUSTIN gives the following preparation ; it is recommended by VAN GESSNER in strangulated hernia :

- R. SYDENHAM's Laudanum, twenty drops.
HOFFMANN's Liquor, one dram.
Mint Water, four ounces.

Mix.—Dose, a spoonful every five or six minutes till the spasms go off.

SEDATIVE MIXTURE. (VAN MONS)

- R. SYDENHAM's Laudanum, half a dram.
Sulphuric Æther, two drams.
Peppermint Water, six ounces.
Compound Bryony Water, two ounces.
Syrup of Poppies, one ounce.

Mix.—Dose, from half a spoonful to one spoonful, every hour.

SEDATIVE POTION. (RATIER)

- R. SYDENHAM's Laudanum, } of each,
Sulphuric Æther, } fifteen
Orange-flower Water, } drops.
Infusion of Lime Flowers, five ounces.
Syrup of Maidenhair, two ounces.

Mix.—To be taken by spoonfuls.

ANTI-DYSENTERIC POTION.

Mixtura opiato-anti-dysenterica. (AUGUSTIN)

- R. Peppermint Water. three ounces.

- Sulphuric Æther, sixteen drops.
SYDENHAM's Laudanum, thirty drops.
Oil of Chamomile, six drops.
White Sugar, two drams.
Gum Arabic, one dram.
Extract of Cascarilla, one scruple.
Catechu, six grains.
Powdered Cinnamon, half a dram.

Mix.—Dose, one spoonful, every hour.—It is extolled by BUCKING.

D.

Extraction by Alcohol.

ALCOHOLIC TINCTURE OF OPIUM.

Tinctura Opii seu sedativa seu Opii simplex seu thebaica seu Opii spirituosâ, Laudanum liquidum, Alcohol cum Opio. (LOND. ; EDINB. ; DUBL. ; AMER. ; ANT. ; BRUNS. ; DAN. ; MIL. COPEN. ; GALL. ; FERR. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; LUSIT. ; CAST. BORUS. ; BORUS. ; COKE ; SWEDIAUR ; VAN MONS)

Tinctura Opii.

- R. Hard Opium, in powder, two ounces and a half.
Proof Spirit, two pints.
Macerate for fourteen days, and strain. (LOND.)
R. Purified Hard Opium, in powder, ten drams.
Proof Spirit, one pint.
Digest for seven days, and strain. (DUBL.)

Tinctura Opii sive thebaica, vulgo Laudanum liquidum. (EDINB.)

- R. Opium, two ounces.
Weaker Alcohol, two pounds.
Macerate for seven days, and filter.
R. Aqueous Extract of Opium, one part.
Alcohol (22 degrees), twelve parts.
Dissolve in a well-stopped bottle, and strain. (GALL.)

FORM. HÔP. FR. prescribes one of opium, and eleven of alcohol.—DAN. ; and MIL. COPEN.—one of opium, and forty-eight of alcohol.—BRUNS. ; and VAN MONS—one of opium, and eight of alcohol.—AMER. ; BRUGNATELLI ; COKE ; and SWEDIAUR—the same as EDINB.

- R. Purified Opium, one dram.
Alcohol, one pound.
Water, two pounds.
Dissolve, and filter. (LUSIT.)
2 O

CAST. BORUS. prescribes half an ounce of opium, and three ounces each, of alcohol and of water.—**FERR.**—ten drams of hard extract of opium, ten ounces of alcohol, and six of water.

R. Purified Opium, one ounce.
Alcohol (15 degrees), eight ounces.

Triturate in a stone mortar, and digest for twenty-four hours, occasionally shaking the vessel; then decant the clear liquor, and pour four ounces of alcohol (15 degrees) upon the residue; after two days' digestion, strain. Then mix and filter the two liquors. (ANT.)

ALCOHOLIC EXTRACT OF OPIUM. (ANT.; BATAV.; BELG.; FERR.; FORM. HÔP. FR.; LUSIT.)

R. Opium, any quantity.
Alcohol (25 degrees), a sufficient quantity

to dissolve it. Filter the liquor, and precipitate it by pouring into it a sufficient quantity of distilled water; then filter again, and evaporate over a slow fire to the consistence of extract. (ANT.)

R. Opium, one pound.
Alcohol, a sufficient quantity

to dissolve it. Digest it over a slow fire, in a close vessel, frequently stirring; then filter, and distil the tincture till it assumes the consistence of extract. (LUSIT.)

R. Opium, two ounces.
Alcohol, four ounces.

Digest for two days in a moderate heat; then express and filter: triturate the residue with four ounces of water, and express. Filter the liquor, mix, and evaporate to the consistence of extract. (BATAV.)

R. Opium, sliced, one part.
Water, two parts.

Infuse for twelve hours in a covered vessel, occasionally stirring, and express strongly; infuse the residue in one part of water, and then in one part of alcohol (12 degrees): set aside, strain, and evaporate the two mixed liquors to the consistence of an extract. (FERR.; FORM. HÔP. FR.)

This extract is seldom prescribed now; it contains more narcotine and more resin than that prepared with cold water.

CHRESTIEN'S ANTI-SPASMODIC TINCTURE. (BORIES; PIERQUIN)

R. Opium, forty-eight grains.
Alcohol (20 degrees), four ounces.

Macerate for three or four days.

It is recommended in amenorrhœa, dysmenorrhœa, spasmodic strangury, hysteria, nervous colic, rheumatism, cardialgia, and

intermittent fevers. As an embrocation, in the quantity of an ounce at each application, twice or thrice a day.

It is sometimes used unstrained and sometimes strained.

ROUSSEAU'S LAUDANUM.

Laudanum opiatum Rousseau, Solutio Opii alcoholisata; Vin opiace préparé par la fermentation. (ANT.; GALL.; CADET DE GASSICOURT; VAN MONS)

R. Honey, twelve ounces.
Warm Water, three pounds.

Dissolve the honey, and put the liquor in a warm place; when it begins to ferment, add to it

Opium, four ounces,
previously dissolved in

Water, twelve ounces.

Leave the mixture to ferment for one month in a temperature of 24 degrees R.; then, having strained and filtered the liquor, evaporate it till only ten ounces remain: then strain again, add four ounces and a half of alcohol (20 degrees), and preserve. (GALL.; CADET DE GASSICOURT)

R. Extract of Opium, one part.
Spirit from Hydromel, eight parts.

Mix, and filter after several days' repose. (VAN MONS)

ROUSSEAU'S laudanum is thicker than SYDENHAM'S. Twenty drops or twenty-two grains contain three grains of opium. The same remark may be applied to this preparation as will be found applied to the extracts of opium prepared after the methods of DEYEUX and of LANGELOT, excepting, however, the formula by VAN MONS, which is merely a simple tincture of opium. But his formula is even objectionable on account of the indeterminate and variable strength of the spirit which he prescribes. It deserves, therefore, to be rejected, as well also as the preparation which it is intended to supersede.

SEDATIVE CATAPLASM. (SWEDIAUR)

R. Emollient Cataplasm, a sufficient quantity.

Tincture of Opium, one or two ounces.

Mix well.

SEDATIVE INJECTION. (SWEDIAUR)

R. Tincture of Opium, fifty drops.
Distilled Water, four ounces.

Mix.—It is recommended in certain cases of chronic gonorrhœa.

SEDATIVE SOLUTION. (SWEDIAUR)

- R. Aqueous Extract of Opium, one dram.
 Distilled Water, four drams.
 Alcohol, half a dram.

Mix with trituration.—SWEDIAUR erroneously compares this preparation to the English black drop, which contains, as has been seen, the acetate of morphine.

ANODYNE LAVEMENT. (ELLIS; SAUNDERS; SWEDIAUR)

- R. Tincture of Opium, forty drops.
 Infusion of Linseed, six ounces.

Mix.—Two ounces of olive oil may be added. (SAUNDERS)

ELLIS prescribes one dram of tincture, and six ounces of infusion of linseed.

- R. Starch, one dram and a half.
 Boiling Water, six ounces.
 Tincture of Opium, one dram.

Mix. (SWEDIAUR)

OPIATE GARGLE. (FORM. HÔP. FR.; AUGUSTIN)

- R. Tincture of Opium, one part.
 Decoction of Marshmallows, two hundred parts.

Mix. (FORM. HÔP. FR.)

AUGUSTIN prescribes one dram of tincture, and ten ounces of water.

VOGEL has extolled the efficacy of this gargle in laryngitis.

ANTI-SPASMODIC LAVEMENT. (SWEDIAUR)

- R. Tincture of Opium, five or ten drops.
 Infusion of Valerian, three ounces.

Mix.—For children.

ALCOHOLIC SYRUP OF OPIUM. (SÆC.)

- R. Tincture of Opium, sixteen drops.
 Simple Syrup, one ounce.

Mix.

OPIATE GARGLE. (FORM. HÔP. FR.)

- R. Tincture of Opium, one part.
 Syrup of Honey, twenty parts.

Mix.

ANODYNE COLLYRIUM. (GALL.)

- R. ROUSSEAU'S Laudanum, six drops.
 Gum Arabic, half an ounce.
 Rose Water, two ounces.

Mix.

OPIATE LINIMENT. (FORM. HÔP. FR.)

- R. Tincture of Opium, four parts.
 Olive Oil, thirty parts.

Mix.

EXCITANT LINIMENT. (ELLIS)

- R. Rectified Oil of Amber, two drams.
 Olive Oil, half an ounce.
 Tincture of Opium, two drams.
 Proof Spirit, two ounces.

Mix.—To be rubbed between the shoulders in whooping cough.

SEDATIVE OINTMENT.

Unguentum emolliens et sedans. (BRERA.)

- R. Spermaceti Ointment, two ounces.
 Tincture of Opium, one dram.

Mix.—To be applied morning and evening, to hæmorrhoidal tumours.

SEDATIVE FOMENTATION. (NIEMANN)

- R. Tincture of Opium, one part.
 Rectified Oil of Amber, two parts.

Mix.—It has been found useful in tic douloureux.

SEDATIVE POTION. (BRERA; ELLIS)

- R. Tincture of Opium, from fifteen to twenty-five drops.
 Syrup of Poppies, two drams.
 Spirit of Cinnamon, one dram.
 Water, one ounce and a half.

Mix. (ELLIS)

- R. HOFFMANN'S Liquor, three drams.
 Tincture of Opium, eighty drops.
 Cinnamon Water, six ounces

Mix.—Dose, a spoonful every hour. (ELLIS)

- R. Sulphuric Æther, two drams.
 Sugar, of each,
 Gum Arabic, one dram and a half.
 Tincture of Opium, sixty drops.
 Cinnamon Water, two ounces.

Mix.—Dose, a tea-spoonful. (ELLIS)

- R. Infusion of Roses, six ounces.
 Tincture of Opium, twenty drops.
 Syrup of Orange Peel, one ounce.

Mix.—To be taken at four times, one dose every three hours; in passive hæmoptysis with cough. (BRERA)

ABSORBENT POTION. (ELLIS)

- R. Prepared Chalk, three drams,
 Sugar, of each,
 Gum Arabic, one dram and a half.
 Tincture of Opium, sixty drops.
 ———— Kino, two drams.
 Mint Water, six ounces.

Mix.—Dose, a spoonful every two hours, in diarrhoea.

ASTRINGENT POTION. (BRERA)

- R. Simarouba Bark, half an ounce.
 Water, a sufficient quantity

to obtain eight ounces of decoction. Add when strained

Almond Emulsion, two ounces.
Tincture of Opium, twenty drops.

Mix.—To be taken at small doses, in atonic diarrhœa.

R. Simarouba Bark, half an ounce.
Water, a sufficient quantity,
to obtain eight ounces of decoction. Add to the strained liquor

Tincture of Opium, one scruple.
Alcohol, one ounce.
Cinnamon Water, half an ounce.

Mix.—To be taken gradually, in cases where it is supposed that atony is accompanied by torpidity.

R. Hemp Seed, one ounce.
Water, a sufficient quantity,
to obtain a pound of strained liquor. Add

Extract of Leopards-bane, one dram.
Catechu, one scruple.

Syrup of Orange Peel, } of each
Marshmallows, } one ounce.

Mix.—To be taken gradually, in hæmaturia and in other involuntary discharges.

R. Simarouba Bark, half an ounce.
Quassia Wood, two drams.
Water, a sufficient quantity,

to obtain eight ounces of strained liquor, after half an hour's boiling. Add

Extract of Gentian, one dram.
Tincture of Opium, half a dram.

Mix.—Dose, four spoonfuls, every two hours; in atonic diarrhœa and dysentery.

AROMATIC ALCOHOLIC TINCTURE OF OPIUM.

Tinctura Opii simplex seu sedativa seu hypnotica seu thebaica, Alcohol Opii dilutum. (BATAV.; FENN.; FULD.; GENEV.; POL.; BORUS.; SAX.; SUEC.; HERBIP.; SWEDIAUR)

R. Opium, sliced, one part.
Cinnamon Water, } of each,
Alcohol, } three parts.
Strain when dissolved. (POL.; BORUS.; SAX.)

Each dram of this tincture contains about ten grains of opium.

FENN.; and GENEV. prescribe one part of opium, and five parts each of cinnamon water and of alcohol.—BAVAR.—one of opium, two of cinnamon water, and four of alcohol.—SUEC.—one of opium, four of cinnamon water, and six of alcohol.

SWEDIAUR gives two formulæ;—one dram of opium, ten ounces of alcohol, and five of cinnamon water;—two ounces of opium, ten of cinnamon water, and six of alcohol. The former contains one grain of opium in half

an ounce of tincture; and the latter, about four grains in half a dram.

R. Purified Opium, half a dram.
Concentrated Alcohol, ten ounces.
Chamomile Water, twenty ounces.

Digest, and filter. (HERBIP.)

R. Opium, one part.
Chamomile Water, six parts.
Vinous Cinnamon Water, two parts.

Digest for three days, and filter.

OPIATE MIXTURE. (SAUNDERS)

R. Crude Opium, one grain.
Cinnamon Water, one ounce.
Vinous Cinnamon Water, two ounces.

Mix.

ANODYNE ESSENCE.

Essentia anodyna officinalis. (PALAT.; WIRTEM.; SPIELMANN.; VAN MONS)

R. Aqueous Extract of Opium, one ounce.
Vinous Cinnamon Water, nine ounces.

Digest, and filter.—Dose, five, ten, fifteen, or twenty drops.—One scruple contains two grains of opium.

ECCARD'S TINCTURE.

Tinctura Opii Eccardi. (HAMB.; HANN.; OLDEN.; CADET DE GASSICOURT.; NIEMANN.; VAN MONS)

R. Opium, two ounces.
Cloves, one dram.
Cinnamon Water, eight ounces.
Proof Spirit, four ounces.

Digest six days in a warm place; then express and filter.

One dram contains ten grains of opium.

AROMATIC SEDATIVE MIXTURE. (SWEDIAUR)

R. Peppermint Water, five ounces.
Spirit of Nutmeg, one ounce.
Aromatic Electuary, two drams.
Aromatic Tincture of Opium, from one to two drams.
White Sugar, three drams.

Mix.—In cardialgia.—Dose, three spoonfuls every half hour or hour.

SEDATIVE DRAUGHT. (SWEDIAUR)

R. Aromatic Tincture of Opium, twenty drops.
Water of Mentha Crispa, one ounce.
Orange-flower Water, two drams.

Mix.—To be taken at bed time.

ANTI-CATHARTIC MIXTURE. (VAN MONS)

R. Gum Arabic, three drams.
Syrup of Poppies, one ounce and a half.

Chamomile Water, six ounces.
Cinnamon Water, one ounce.
Aromatic Tincture of Opium,
one scruple.

Mix.—Dose, a spoonful every quarter of an hour.

COMPOUND ALCOHOLIC TINCTURE OF OPIUM.

Tinctura vinosa Opii anti-colica. (HISP.)

R. Opium, two ounces.
Root of Zedoary, one ounce.
—— Carline Thistle, } of each, half
—— Galanga, } an ounce.
—— Sweet Flag, }
Dried Orange Peel, }
—— Lemon Peel, } of each,
Seeds of Anise, } two drams.
—— Caraway, }
Flowers of Chamomile, }
Mace, one dram.
Cloves, }
Bay Berries, } of each, half a dram.
Juniper, }
Alcohol, three pounds.

Macerate for eight days, and filter.—Dose, two scruples.

LUDWIG'S ANODYNE ESSENCE.

Essentia anodyna Ludovici. (WIRTEM.)

R. Aqueous Extract of Opium, one ounce.
Apoplectic Liquor, eight ounces.
Mix.—Dose, from seven to fifteen drops.
—one dram contains six grains and a half of opium.

CROCATED ALCOHOLIC TINCTURE OF OPIUM.

Tinctura alcoholica Opii. (HISP.)

R. Opium, two ounces.
Saffron, half an ounce.
Cinnamon, two drams.
Alcohol, } of each, twelve ounces.
Water, }

Digest for eight days, and filter.

This very much resembles SYDENHAM'S laudanum.—One scruple contains two grains of opium.

CHRESTIEN'S ANTI-SPASMODIC TINCTURE. (BORIES)

R. Crude Opium, one dram.
Camphor, four drams.
Proof Spirit, one pound.

Macerate for three or four days, and filter.

This is extolled as an external application in rheumatic affections, intermittent fevers complicated with rheumatism, retention of urine, amenorrhœa, &c.—In the latter case, it is to be rubbed upon the fore and inner part of the thighs.

CAMPHORATED SEDATIVE LINIMENT. (SWEDIAUR)

R. Aromatic Tincture of Opium, one or two drams.
Camphor, one dram.
Liniment of Soap, one ounce and a half.

Mix.—It is recommended in frictions and fomentations, in colic.

ANODYNE LINIMENT. (AUGUSTIN; SAUNDERS)

R. Liniment of Soap, two ounces.
Tincture of Opium, half an ounce.

Mix. (SAUNDERS)

AUGUSTIN prescribes an equal part of each.

It is useful in the same cases as the preceding. LIND says that he has several times cured intermittent fevers by the application of this liniment.

CAMPHORATED OPIATE LINIMENT. (FORM. HÔP. FR.)

R. Tincture of Opium, two parts.
Camphor, one part.
Olive Oil, fifteen parts.

Mix.

DUMAS' ANTI-SPASMODIC TINCTURE. (PIERQUIN)

R. Opium, forty grains.
Camphor, } of each, half a dram.
Amber, }
Alcohol, six ounces.

Mix.—To be rubbed upon the spinal column, the thighs, and legs.

AMMONIACAL AND CAMPHORATED ALCOHOLIC TINCTURE OF OPIUM.

Tinctura Opii ammoniacalis seu volatilis anodyna, Spiritus ammoniacatus. (LUSIT.; AUGUSTIN; VAN MONS)

R. Camphor, saturated with alcohol, one part.
Opium, in powder, two parts.
Spirit of Ammonia, twenty-four parts.

Macerate for several days; and filter. (VAN MONS)

LUSIT. prescribes one dram of opium, two scruples of camphor, four ounces of aromatic spirit of ammonia, and one pound of alcohol.

R. Proof Spirit, one ounce.
Spirit of Ammonia, half an ounce.
Opium, two scruples.
Camphor, one scruple.

Digest for three days, and strain. (AUGUSTIN)

It is recommended in hysteria and con-

vulsive disorders.—Dose, from half a dram to two drams, in syrup.—VICAT recommended it in cephalalgia, odontalgia, and low spirits; being rubbed upon the hands and then held before the nose.

WARNER'S LAUDANUM.

Laudanum liquidum Warneri, Tinctura Opii ammoniacata. (DAN.; VAN MONS)

R. Opium, } of each, six drams
 Soap, }
 Nutmeg, } one dram.
 Camphor, } two drams.
 Saffron, } four grains.
 Spirit of Ammonia, } nine ounces.

Digest for ten days, occasionally shaking the vessel, and decant. (DAN.)

One dram contains five grains and a half of opium.

R. Opium, one ounce and a half.
 Nutmegs, two drams.
 Saffron, two scruples.
 Spirit of Ammonia, eighteen ounces.

Macerate, and add,

White Soap, one ounce and a half.
 Camphor, half an ounce.

Filter the tincture as it may be wanted. (VAN MONS)

TINCTURE OF SOAP AND OF OPIUM.

Linimentum saponis compositum seu camphoratum, Balsamum seu Linimentum anodynum, Tinctura seu Linimentum Saponis et Opii, Tinctura alcohólica Opii saponacea, Balsamum thebaicum. (EDINB.; AMER.; HISP.; GENEV.; HERBIP.; AUGUSTIN; COKE; NIEMANN; PIERQUIN; RATIER; SPIELMANN; VAN MONS)

Tinctura Saponis et Opii, vulgo Linimentum anodynum. (EDINB.)

R. Hard Soap, sliced, four ounces.
 Opium, one ounce.
 Camphor, two ounces.
 Oil of Rosemary, half an ounce.
 Alcohol, two pounds.

Digest the soap and opium in the alcohol for three days; then strain the tincture, and add to it the camphor and oil, frequently shaking.

COKE; and NIEMANN give the same formula.—AMER. prescribes twelve ounces of soap, two drams of oil of rosemary, and eight pounds of alcohol.

R. Opium, one dram.
 Spanish Soap, five ounces.

Alcohol, twenty-four ounces.

Digest for three days, and dissolve in the strained liquor,

Camphor, two ounces.
 Oil of Rosemary, half an ounce.

Preserve. (SPIELMANN)

HERBIP. prescribes two drams of opium, one ounce of soap, three drams of camphor, half a dram of the oil, and nine ounces of alcohol.—GENEV.—one ounce of opium, five ounces of soap, two ounces of camphor, half an ounce of oil of rosemary, and one pound and a half of alcohol.—AUGUSTIN—one dram of opium, one ounce of soap, half a dram of camphor, one dram of oil of rosemary, and a sufficient quantity of alcohol to dissolve the whole.

R. Opium, half an ounce.
 Camphor, six drams.
 White Soap, half an ounce.
 Saffron, one dram.
 Alcohol, one pound and a half.

Digest for eight days, and strain. (HISP.; and PIERQUIN)

R. Saffron, one part.
 Alcohol, forty-two parts.

Digest in a warm water-bath, and add,

Opium, sliced, two parts.

Digest again, and add,

White Soap, scraped, four parts.

When the solution is completed, express and filter. (VAN MONS).

R. Opium, one ounce.
 Spirit of Rosemary, twenty-four ounces.

Macerate for a few days, express, and filter; dissolve in the strained liquor,

Camphor, three drams.
 Marrow Soap, six ounces.

Set aside, and decant. (VAN MONS)

R. Opium, one dram.
 Hard White Soap, half an ounce.
 Alcohol, four ounces.

Digest for three days, strain, and add,

Camphor, two drams.

Shake well together. (RATIER)

Dose, thirty, forty, or fifty drops, in wine.—It is also employed externally, in rheumatism.—This preparation differs only from the liniment of soap, by the presence of opium.

SOMNIFEROUS BALSAM. (WIRTEM.)

R. Unguentum Populeum, } of each,
 Expressed Oil of Nutmeg, } one ounce.
 Opium, dissolved in alcohol, one dram.
 Oil of Rhodium, sixteen drops.

Mix.

CAMPHORATED ALCOHOLIC TINCTURE OF OPIUM AND OF BENZOIC ACID.

Tinctura Opii benzoicæ seu Camphoræ composita seu Opii camphorata seu sedativa camphorata, Elixir paregorica. (LOND.; DUBL.; EDINB.; AMER.; BATAV.; DAN.; HANN.; LIPP.; BORUS.; SAX.; SUEC.; BORIS.; BRUGNATELLI.; CADET DE GASSICOURT.; COXE.; FEE.; PIERQUIN.; SWEDIAUR.; VAN MONS.)

Tinctura Camphoræ composita. (LOND.)

R. Camphor, two scruples.
Hard Opium, powdered, } of each,
Benzoic Acid, } one dram.
Proof Spirit, two pints.
Macerate for fourteen days, and filter.

Tinctura Opii camphorata, vulgo Elixir paregoricum Anglorum. (EDINB.)

R. Camphor, two scruples.
Hard Purified Opium, } of each,
in powder, } one
Benzoic Acid, } dram.
Proof Spirit, two pounds and a half.
Digest for seven days, and filter.

Tinctura Opii camphorata, sive Elixir Paregoricum. (DUBL.)

R. Hard Purified Opium, } of each,
in powder, } one
Benzoic Acid, } dram.
Camphor, two scruples.
Oil of Anise, one dram.
Proof Spirit, two pints.
Digest for ten days, and strain.

COXE.; FEE.; and PIERQUIN give the same formula as LOND.—HANN.; LIPP.; BORUS.; BORIS.; BRUGNATELLI.; NIEMANN.; PIERQUIN.; and SWEDIAUR give the same as DUBL.—DAN.; SUEC.; and VAN MONS prescribe half a dram of oil of anise.—CADET DE GASSICOURT prescribes one dram of opium, one dram of benzoic acid, twenty-six grains of camphor, two scruples of the oil, and twenty-three ounces of alcohol (24 degrés).

R. Opium, }
Benzoic Acid, } of each,
Camphor, } one half part,
Oil of Anise, }

Alcohol, ninety-six parts.
Digest, and strain. (SAX.)

R. Opium, }
Benzoic Acid, } of each, one dram.
Oil of Anise, }
Extract of Liquorice, half an ounce.
Despumated Honey, two ounces.
Camphor, two scruples.
Alcohol (0.930), two pounds.

Digest for ten days, and strain. (AMER.; COXE.; NIEMANN.)

In mentioning the additions which embellish this formula, COXE asks, and with reason: *Cui bono?*

It is extolled in asthma.—Dose, from twenty to one hundred drops to adults, and from five to twenty to children.

ALCOHOLIC TINCTURE OF OPIUM AND OF BENZOATE OF AMMONIA.

Tinctura Opii ammoniata, olim Elixir paregoricum. (EDINB.)

R. Opium, two drams.
Saffron, } of each,
Benzoic Acid, } three drams.
Oil of Anise, half a dram.
Spirit of Ammonia, sixteen ounces.

Digest for seven days, and filter.

CADET DE GASSICOURT.; COXE.; FEE.; HENRI.; and PIERQUIN give the same formula.

R. Saffron, six drams.
Spirit of Ammonia, thirty-two ounces.

Infuse for two days without heat, and add, Opium, sliced, half an ounce.

Infuse again; and, at the end of several days, add,

Benzoic Acid, six drams.
Oil of Anise, one dram.

Express and filter. (VAN MONS)

R. Saffron, five drams.
Aromatic Spirit of Ammonia, four ounces.

Digest for two days; then strain and add to the tincture,

Benzoic Acid, five scruples.
Opium, two drams.

Digest for two days; strain and filter. Pour upon the residue,

Compound Spirit of Ammonia, two ounces.

Digest for twenty-four hours, strain, and mix with the tincture the matter remaining on the filter: strain again at the expiration

of twelve hours. Mix the two tinctures, and add

Oil of Anise, thirty-two drops.

Preserve. (ANT. ; BELG.)

R. Hard Opium, two drams.
Benzoic Acid, half an ounce.
Spirit of Ammonia, one pound.

Digest for five hours, and filter. (CADET DE GASSICOURT ; PIERQUIN)

It is principally recommended in asthma and whooping cough.—Dose, from thirty to forty drops, in a proper vehicle.

LETTSON'S SUDORIFIC ELIXIR.

Elixir sudoriferum, Tinctura Opii benzoica composita. (AUGUSTIN)

R. Ipecacuanha, } of each,
Balsam of Tolu, } half an ounce.
Benzoic Acid, }
Opium, } of each, two drams.
Saffron, }
Camphor, } four scruples.
Oil of Anise, } one dram.
Alcohol, } two pounds.

Digest, and strain.—Dose, from five to thirty drops, augmenting progressively; in whooping cough.

ANTI-PODAGRIC EPITHEM. (BORIES)

R. Hungary Water, eight ounces.
Spirit of Ammonia, half an ounce.
Camphor, two drams.
Opium, half a dram.
Venice Soap, one ounce and a half.

Mix.—It is to be applied hot by means of fine linen, and to be renewed every hour.

ANTIMONIATED SEDATIVE TINCTURE.

Tinctura stibiata. (AUGUSTIN)

R. Solution of Succinate of } of each,
Ammonia, } two drams.
Paregoric Elixir, }
SYDENHAM'S Laudanum, } of each,
Antimonial Wine, } one dram.

Mix.—Dose, from twenty to forty drops every two hours, in whooping cough. It is extolled by JAHN.

LEMORT'S ANTI-ASTHMATIC LIQUOR. (VAN MONS)

R. Benzoic Acid, } of each, one dram.
Opium, }
Subcarbonate of Potass, } one ounce.
Honey, } of each, four ounces.
Liquorice, }
Oil of Anise, } half a dram.
Camphor, } two scruples.
Proof Spirit, } thirty-four ounces.

Digest in the spirit, first the saffron, then the opium, and then the liquorice; then add

the subcarbonate, and express at the end of several days: dissolve successively in the tincture, the acid, oil, camphor, and honey. Set apart, pass through a sieve, and then let the tincture stand till it becomes perfectly transparent.

SEDATIVE ÆTHERIAL LINIMENT. (VAN MONS)

R. Aromatic Tincture of Opium, six drams.
Sulphuric Æther, two drams.

Mix.—It is recommended in ischury, and in spasmodic and rheumatic affections.

SEDATIVE MIXTURE. (SWEDIAUR)

R. Water, six ounces.
Aromatic Tincture of } of each,
Opium, } one dram.
Sulphuric Æther, }
Simple Syrup, } one ounce.

Mix.—Dose, a spoonful every hour or two hours.

ANODYNE ACID MIXTURE. (FULD. ; HAMB.)

R. Tincture of Opium, two scruples.
Sulphuric Acid, one scruple.
Distilled Water, six ounces.
Syrup of Violet, one ounce.

Mix. (FULD.)

R. Tincture of Orange, half an ounce.
Sulphuric Acid, one dram.
Tincture of Opium, half a dram.

Mix. (HANN.)

VERMIFUGE TINCTURE.

Tinctura fetida pyro-oleosa. (AUGUSTIN)

R. Assafetida, one dram.
Opium, } of each,
Subcarbonate of } two drams.
Ammonia, }
Tincture of Castor, } two ounces.

Mix.—ROSENSTEIN recommends it in cases of tænia.—Dose, from thirty to forty drops, in wine or brandy.

E.

Extraction by Water.

AQUEOUS INFUSION OF OPIUM. (RATIER ; PLANCHE)

R. Opium, one ounce.
Distilled Water, eight ounces.
Alcohol (36 degrees), half an ounce.

Infuse the opium in the water for two or three days, at the ordinary temperature, and in a dark place; then filter, add the alcohol, and preserve in well-stopped phials.

CHAUSSIER substitutes this infusion for

SYDENHAM'S Laudanum.—Dose, from sixteen to twenty-four or thirty drops; as an external application, from fifty to eighty drops in two ounces of infusion of mallow-flowers.

AQUEOUS EXTRACT OF OPIUM.

Extractum Opii seu Opii gummosum seu aquosum seu thebaicum, Laudanum opiatum seu simplex, Opium colatum seu depuratum. (LOND.; DUBL.; AUSTR.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; COXE; SAUNDERS.; SPIELMANN; SWEDIAUR; VAN MONS)

Extractum Opii. (LOND.)

R. Opium, sliced, sixteen ounces.
Water, one gallon.

Pour a little of the water upon the opium, and macerate for twelve hours that it may become soft; then add the remainder of the water, triturating together till the whole is well mixed, and set apart the mixture that the impurities may subside. Finally, strain and evaporate the liquor to the proper consistence.

Extractum Opii aquosum. (DUBL.; COXE)

R. Opium, two ounces.
Boiling Water, one pint.

Triturate the opium with the water for ten minutes, and after a short time pour off the liquor; rub the residue with an equal quantity of boiling water for the same space of time, and pour off the liquor again; then repeat this operation a third time. Mix together the decanted liquors, and expose the mixture for two days to the air in a broad open vessel. Finally, strain it through linen, and evaporate slowly.

R. Opium, in powder, any quantity.

Triturate it with a little cold water till this acquires a brown colour; then set aside, and decant: repeat the operation till the water ceases to become coloured. Mix the liquors together, and evaporate in a water-

bath to the consistence of an extract. (FERR.; FULD.; LIPP.)

R. Opium, any quantity.

Knead it under a stream of cold water till this ceases to dissolve any more of it; then strain the liquor, clarify it with white of egg, evaporate it over a brisk fire, and when the liquor begins to thicken, strain and evaporate to the required consistence. (BELG.)

R. Opium, sliced, one part.
Cold Water, four parts.

Digest in a stove for two days, occasionally shaking the vessel; then strain, and set aside the liquor for two days; filter to separate the pellicle, of a virous odour, which is formed, and evaporate slowly to one half: place it again for two days in a stove, then filter again to remove the second pellicle, and finally evaporate to the consistence of an extract. (GALL.)

R. Opium, one part.
Cold Water, three parts.

Macerate for three days, without agitation, and decant; pour two parts of water upon the residue, and decant again after a short time: having repeated this operation a third time, strain the mixed liquors; clarify with white of egg, and finally evaporate, boiling slowly, and decanting in proportion as precipitate is formed. (VAN MONS)

R. Opium, any quantity.
Cold Water, a sufficient quantity.

Macerate for twelve hours, then digest in a moderate heat for twelve hours more, and strain; treat the residue in the same manner, till the water ceases to acquire any colour: then mix the liquors, and evaporate them to the consistence of an electuary. Dissolve the mass in cold water, set aside the liquor, strain, and finally evaporate to the consistence of an extract. (SUEC.)

R. Opium, four ounces.
Water, acidulated with half a pound
of Lemon Juice, four pounds.

When dissolved, filter, and evaporate to the consistence of an extract. (BRUNS.; PALAT.; WIRTEM.)

R. Opium, two ounces.
Boiling Water, one pound.

Dissolve; strain the hot liquor, and add Cold Water, one pound.

Expose the mixture to the open air for two days; then filter, and evaporate, first over a slow fire and then in a water-bath, to the consistence of an extract. (BRUGNATELLI)

R. Opium, in coarse powder, one part.
Water, four parts.

Digest for two days at a moderate heat
2 P

and with frequent agitation; having then decanted and strained the liquor, evaporate it in a water-bath. (AUSTR.; DAN.; HISP.; HANN.; OLDEN.; LUSIT.; PALAT.; BORUS.; SAX.; SAUNDERS; SPIELMANN; SWEDIAUR)

FORM. HÔP. FR. directs the liquor to be set apart for twenty-four hours; then to be filtered, and boiled down to one half; to be then set aside again, strained, and finally evaporated in a water-bath.—BAVAR.—the maceration to be continued for twenty-four hours, and the operation to be repeated with the residue and one fourth of water; the mixed liquors to be then evaporated.

The process generally followed at the present day is similar to that required by GALL. It is this: cut or rasp the opium according to its consistence, and macerate it repeatedly in cold water till dissolved; then strain the mixed liquors, evaporate the whole to one half, and set aside to cool: then filter to remove the resin, which separates partly in the form of precipitate and partly in that of a pellicle, and evaporate the clear liquor: then let it cool, and filter again. Finally, when the extract has arrived at the consistence of thick honey, mix it with cold distilled water; then filter the liquor, and evaporate it to the consistence of pilular mass.

When thus prepared, the extract contains meconate of morphine, a small quantity of narcotine, colouring matter, a little resin, gum, and fecula.—Dose, from half a grain to three grains.

ESSENTIAL SALT OR HARD EXTRACT OF OPIUM. (GALL.)

R. Opium, sliced, two pounds.
Cold Water, twelve pounds.

Macerate for twenty-four hours, occasionally stirring, and strain; macerate the residue with eight parts of water, and strain again. Mix the liquors, and evaporate them in a water-bath to the consistence of syrup. Pour this into dishes about two lines in depth, evaporate to dryness in an oven or water-bath, and detach the scales.

AQUEOUS EXTRACT OF OPIUM WITHOUT NARCOTINE. (FERR.; FEE)

R. Aqueous Extract of Opium, any quantity.

Having mixed it with a little water, put it into a bottle with purified æther, and shake the mixture frequently; after a short time, decant, and add fresh æther: continue to do this till the decanted æther ceases to leave a residuum on being evaporated; then evaporate the extract to the consistence of pilular mass.

This is the process of ALPHONSE LEROY as given by ROBQUET. DUBLANC has declared it inefficient, and proposes the following:

R. Extract of Opium, prepared without heat, three hundred parts.
Distilled Water, one hundred and fifty parts.

Dissolve; pour the solution into a retort, and add to it

Pure Æther, two thousand parts.

Distil slowly about two hundred parts of æther; separate the apparatus, and pour off the supernatant æther from the retort: wash the extract while hot with distilled æther; then dissolve it in distilled water, and evaporate it again to the proper consistence.

Thus prepared, the extract of opium does not contain an atom of narcotine.

DIEST'S EXTRACT OF OPIUM. (GALL.)

R. Opium, one pound.
Water, four pounds.

Boil slightly for half an hour, and strain; then having boiled the residue twice more with an equal quantity of water, strain the three mixed decoctions: put the mixture into a tin cucurbit placed in a sand-bath heated by a lamp in such a manner that it may be kept slowly boiling, without ceasing, for six months, adding occasionally a little distilled water, and decanting to remove the sediment. At the end of the six months, let the liquor cool, strain it through flannel, and evaporate it in a water-bath to the consistence of pilular mass.

This process is justly abandoned, for it presents no real advantage over the more speedy ones.

EXTRACT OF OPIUM BY FERMENTATION. (GALL.; CADET DE GASSICOURT)

R. Opium, sliced, any quantity.
Water, } of each,
Yeast of Beer, } a sufficient quantity.

Ferment at a proper temperature, and when the liquor becomes clear, dilute it with water; then having filtered it, boil it till all virous odour becomes dissipated, and finally evaporate it to the consistence of thick extract.

This process is due to DEYEUX. It wants exactness; for the alterations which the opium undergoes must be more or less considerable as the fermentation may have been more or less perfect. No one has hitherto ascertained the kind of alteration which it must of necessity undergo.

LANGELOT'S CYDONIATED EXTRACT OF OPIUM.

Extractum Opii cydoniatum Langelotti.
(SPIELMANN)

R. Opium, twelve ounces.

Dissolve it in a mixture of

Juice of Quinces, } of each,
Spring Water, } sixty ounces.

Digest for several days in a warm place, and strain. Add to the liquor

White Sugar, four ounces.

Leave the mixture to ferment for several months, occasionally stirring it with a wooden spatula; on the fermentation ceasing, strain the liquor through linen, and evaporate it to the consistence of an extract.

The same remark may be made with regard to this as the preceding extract. It very probably contains acetate of morphine.

CROCATED EXTRACT OF OPIUM.

Laudanum opiatum. (BRUNS.; HISP.; PALAT.; SARD.; WIRTEM.)

R. Aqueous Extract of Opium, two ounces.

Extract of Saffron, } of each,
Balsam of Mecca, } half a dram.

Mix. (PALAT.; WIRTEM.)

BRUNS. requires powdered, instead of extract of, saffron.

R. Aqueous Extract of Opium,

two ounces.
Extract of Saffron, one ounce.
Prepared Mother of Pearl, } of each,
Hyacinth, } one dram
Coral, } and a half.
Terra Sigillata, one dram.
Bezoar, four scruples.
Hartshorn Shavings, two scruples.

Pulverize, reduce to a paste with a little water, and form it into trochiscs. (HISP.)

R. Extract of Opium, } of each,
Saffron, } one ounce.
Powdered Amber, } six drams.
Castor, three drams.

Mix the extracts with a little alcohol, and then incorporate the powders. (SARD.)

ANODYNE EXTRACT. (WIRTEM.)

R. Aqueous Extract of Opium, three drams.

Extract of Blessed Thistle, five drams.
Sweet Flag, } of each,
Angelica, } one ounce.
Oil of Cloves, ten drops.

Mix.—Each dram contains seven grains and a half of opium.

ANTI-HYSTERIC LAUDANUM. (WIRTEM.)

R. Extract of Opium, half an ounce.
Castor, one dram.
Saffron, one scruple.
Essential Oil of Mace, } of each,
Mint, } six drops.
Rue, }

Make a pillular mass.—It is recommended in hysterical colic, to appease the spasms.
—Dose, one or two grains.

GLYCYRRHIZED EXTRACT OF OPIUM.

Laudanum diureticum seu urinarium, Extractum Opii glycyrrhizatum. (HISP.; WIRTEM.)

R. Aqueous Extract of Opium, two drams.

Extract of Liquorice, one ounce and a half.
Trochiscs of Alkekenge, } of each,
Venice Turpentine, } half an ounce.
Camphor, one dram.
Saffron, four scruples.
Gum Tragacanth, } of each,
Mastic, } half a dram.

Mix the extracts with a little water, and having reduced them to the consistence of honey, incorporate the turpentine; then add the other ingredients in powder, and form the mass into trochiscs.

This preparation is extolled in spasmodic affections of the urinary apparatus.—Thirteen grains contain one grain of opium.

CYDONIATED EXTRACT OF OPIUM.

Laudanum opiatum cydoniatum, Extractum Opii cydoniatum. (PALAT.; WIRTEM.; BORIES)

R. Opium, sliced, half a pound.
Juice of Quinces, recently expressed, six pounds.

Digest in a moderate heat; then decant, filter, and evaporate the liquor: add to it

Essential Oil of Cinnamon, } of each,
Cloves, } half
Mace, } a scruple.

Mix. (PALAT.; WIRTEM.)

R. Opium, in powder, four ounces.
Juice of Quinces, recently expressed, four pounds.

Digest for three weeks, and add

Cloves, } of each, one ounce.
Nutmegs, }

Heat slightly, and then leave to digest for eight days. Add then

Saffron, one ounce.
White Sugar, four ounces.

At the end of twenty-four hours, filter, and evaporate in a water-bath, till the li-

quor is reduced to one third, and preserve it. (BORIES)

This preparation, now disused, differs from the aqueous extract of opium by the impurities of the quince juice and by the aromatics which are added to it. It was formerly classed among the astringent antispasmodics, and was thought useful in vomiting, diarrhoea, hiccup, and generally as an anodyne.—Dose, one grain to an adult, and from the twentieth to the tenth of a grain to children. This however, applies only to the product of the second formula, for that of the first contains one grain of opium in ten drops.

SEDATIVE PILLS.

Pilulæ sedativæ seu hypnoticæ. (PH. SYPH.; RATIER; SWEDIAUR; VAN MONS)

SWEDIAUR's formula is very simple; it consists of dividing the aqueous extract of opium into pills of one grain each.—One or more for a dose, according to the indication.

R. Extract of Opium, three grains.
Camphor, six grains.
Simple Syrup, a sufficient quantity.

Make six pills. (PH. SYPH.; RATIER; VAN MONS)

R. Extract of Opium, one grain.
Camphor, six grains.
Nitrate of Potass, four grains.
Simple Syrup, a sufficient quantity.

Make one pill. (RATIER)

OPIATE PECTORAL PILLS. (FULD.)

R. Extract of Liquorice, one ounce and a half.
Opium, half an ounce.
Syrup of Poppies, a sufficient quantity.
Mix.—Dose, from two to six grains.

OPIATE BALSAMIC PILLS.

Pilulæ balsamicæ cum Opio. (AUGUSTIN)

R. Extract of Opium, half a dram.
Balsam of Peru, one dram and a half.
Myrrh, three drams.

Make pills of one grain each.—Dose, two or three every hour, alternating with a dose of tincture of opium. They are recommended by MARCUS.

ANTI-DIARRHOIC BOLUS. (SAINTE-MARIE)

R. Aqueous Extract of Opium, one grain.
Pimento, } of each, four grains.
Cassia Lignea, }
Saffron, }
Conserve of Dog-rose, a sufficient quantity.

Mix.

LOZENGES OF OPIUM. (VAN MONS)

R. Extract of Opium, one part.
White Sugar, sixty-four parts.

Incorporate the extract with the sugar, melted at a very gentle heat, and form the mass into lozenges of six grains each.

PECTORAL PASTILLES. (FERR.)

R. Mucilage of Iceland Moss, } of each,
Gum Arabic, } one pound.
Sugar, two pounds.
Opium, one dram and a half.

Make pastilles of twenty-four grains each.—Dose, ten, daily.

ANODYNE POWDER. (HAMB.; WIRTEM.; CADET DE GASSICOURT; PIERQUIN)

R. Extract of Opium, one dram.
Prepared Red Coral, two drams.

Mix. (WIRTEM.)

R. Aqueous Extract of Opium, two grains.
Compound Powder of Tragacanth, two ounces.

Mix. (PIERQUIN)

R. Extract of Opium, } of each,
Red Sanders Wood, } one scruple.
Prepared Red Coral, four scruples.

Mix. (WIRTEM.)

ANODYNE CATAPLASM.

Cataplasma cum Opio. (AUGUSTIN)

R. Linseed Meal, four ounces.
Water, a sufficient quantity
to make a cataplasm. Add,
Opium, half a dram.

OPIATE LINIMENT. (RATIER)

R. Aqueous Infusion of Opium, four drams.
Olive Oil, two ounces.
Shake well together.

PH. SYPH. prescribes a *Sedative Lavement* which resembles this liniment; it is prepared with two or three grains of aqueous extract of opium, and four ounces of olive or linseed oil.

SWEDIAUR also gives a *Sedative Liniment*, composed of one dram of extract of opium, triturated first with a little hot water, and then from three to eight drams of simple cerate.

They are employed to appease local pain.

OPIATE COLLYRIUM. (RATIER)

R. Aqueous Extract of Opium, half a dram.
Water, one pound.

Mix.

RATIER also prescribes two grains of extract and four ounces of water. — The strength of the collyrium might be varied according to the indication.

SEDATIVE SOLUTION. (RATIER; SWEDIAUR)

R. Extract of Opium, from two to four drams.
Hot Water, one pound.
Triturate together. — For fomentations to pruriginous affections of the skin.

SEDATIVE GARGLE. (SWEDIAUR)

R. Extract of Opium, one or two drams.
Boiling Water, one pound.
Mix, and add,
Alcohol, one ounce.
It is useful in painful ulcers of the throat and tongue, in rheumatic odontalgia and otalgia, and in nervous head-ach.

OPIATE LAVEMENT. (SPIELMANN; SWEDIAUR)

R. Bruised Linseed, one ounce.
Boiling Water, six ounces.
Infuse for an hour; then add to the strained liquor,
Extract of Opium, two grains.
Mix.

SEDATIVE INJECTION. (CADET DE GAS-SICOURT)

R. Extract of Opium, sixteen grains.
Infusion of Linseed, one pound.
Mix.

SYRUP OF OPIUM.

Syrupus Opii seu de Opio seu opiatius seu thebaicus seu sedativus seu hypnoticus.
(DUBL.; ANT.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; COXE; VAN MONS)

Syrupus Opii. (DUBL.; COXE)

R. Aqueous Extract of Opium, eighteen grains.
Hot Water, eight ounces.
Macerate, and when dissolved, form a syrup with
White Sugar, twenty-seven ounces.
R. Extract of Opium, prepared with cold water, three drams sixty grains.
Water, two ounces.
Dissolve, and add,
Simple Syrup, nine pounds nine ounces and a half

Mix, boil for a few moments, and strain through flannel. (GALL.)

FORM. HÔP. FR. prescribes one part of extract, four parts of water, and six hundred of syrup. — ANT. — one ounce of purified opium, a little water, and fifteen pounds of boiling syrup. — HISP. — one scruple of extract, three ounces of water, and one pound of syrup, to be boiled to the proper consistence. — FERR. — three drams of extract, four pounds of sugar, and two pounds and a half of water.

R. Extract of Opium, made without heat, one part.
Simple Syrup, four hundred and eighty parts.

Incorporate gradually, triturating it in a marble mortar with a large pestle. (VAN MONS)

R. Opium, cut in small pieces, one ounce and a half.
Water, eighteen ounces.

Macerate without heat till perfectly dissolved, then strain, and to eight parts of the strained liquor, add

White sugar, fifteen parts.
Dissolve in a water-bath. (VAN MONS)

Dose, from two to four drams, twice or oftener in the day.

OPIATE INJECTION.

Injectio Lactis opiata seu anodyna. (ELLIS)

R. New Milk, six ounces.
Syrup of Opium, one ounce.
Mix.

NARCOTIC POTION. (DISP. CHA. PARIS)

R. Syrup of Opium, half an ounce.
Lime-flower Water, three ounces.
Orange-flower Water, one ounce.
Mix.

EMULSIVE SYRUP OF OPIUM. (VAN MONS)

R. Gum Arabic, one part
Opium, two parts.
Triturate together and make an emulsion with
Water, forty-six parts.
Set aside, and dissolve with a moderate heat in the decanted liquor,
White Sugar, two hundred and thirty-six parts.

SEDATIVE LAVEMENT. (ELLIS)

R. Opium, two grains.
Mucilage of Gum Arabic, half an ounce.
Warm Milk, six ounces.
Mix.

OPIATE POTION. (AUGUSTIN)

- R. Opium, one grain.
Water, one ounce.

Dissolve.—To be taken at one dose; for an adult.

- R. Water, half an ounce.
Aromatic Liquor, } of each, two drams.
Syrup of Opium, }
Tincture of Opium, ten drops.

Mix.—Dose, a tea-spoonful.

ANODYNE OINTMENT. (PIERQUIN)

- R. Lard, three ounces.
Syrup of Opium, one ounce.
Essence of Roses, four drops.

Mix.—It is useful in inflammations of the membrane of the mouth.

PECTORAL SYRUP. (CADET DE GASSICOURT)

- R. Senna, } of each,
Liquorice, } nine ounces three drams.
Cinchona, six ounces three drams.
Anise, five ounces.
Elder Flowers, three ounces one dram.
Tops of Lesser Centaury, one ounce and a half.
Ipecacuanha, five drams.
Water, a sufficient quantity.

Boil for a long time; and add to the strained liquor,

- White Sugar, eight pounds.
Aqueous Extract of Opium, four drams and a half.

Mix.—Dose, from half a tea-spoonful to a table spoonful, in infusion of hyssop.

MALOET'S PECTORAL SYRUP. (BORIES; CADET DE GASSICOURT; HENRI; PIERQUIN)

- R. Jujubes, } of each, one ounce.
Dates, }
Corinth Raisins, }
Water, a sufficient quantity.

Boil, and add, towards the end,

- Canadian Maidenhair, four drams.
Liquorice, two drams.

Strain with slight expression, and add,

- White Sugar, two pounds.

Having boiled to a syrup, add,

- Extract of Opium, mixed with a little water, six grains.

Boil again, and strain.—Dose, one ounce.

SEDATIVE INJECTION. (RATIER)

- R. Poppy Heads, two.
Stems of Solanum Nigrum, one dram.
Water, one pound.

Boil, and add when strained,

- Extract of Opium, from ten to twenty grains.

Mix.

ANTI-DYSENTERIC MIXTURE. (VAN MONS)

- R. Opium, two or three grains.
Syrup of Poppies, one ounce.
Chamomile Water, six ounces.

Mix.—Dose, from half a spoonful to one spoonful, every quarter of an hour.

NARCOTIC COLLYRIUM. (CADET DE GASSICOURT)

- R. Infusion of Henbane Leaves, half a pound.
Extract of Belladonna, eight grains.
Opium, four grains.

Mix.

FEBRIFUGE POTION. (BORIES)

- R. Mint Water, } of each,
Orange-flower Water, } two ounces.
Valerian Water, }
Tincture of Castor, } of each,
Amber, } twenty-five drops.
Assafoetida, }
Theriac, one dram.
Aqueous Extract of Opium, one grain and a half.

- Solution of Ammonia, twelve drops.
Syrup of Orange Peel, one ounce.

Mix.—To be taken at two equal doses, on the day of the paroxysm, one six hours before and the other an hour after its invasion. It is said to have subdued intermittents which had resisted cinchona.

SUCCINATED SYRUP OF OPIUM.

Syrupus Opii succinatus seu *Karabe*. (ANT.; CADET DE GASSICOURT; HENRI)

- R. Syrup of Opium, one pound.
Succinic Acid, dissolved in water, one scruple.

Mix. (ANT.)

CADET DE GASSICOURT prescribes two scruples of acid.

- R. Syrup of Opium, one pound.
Spirit of Amber, two scruples.

Mix. (HENRI)

Dose, from half an ounce to one ounce, in a potion.

ANTI-HÆMORRHOIDAL OINTMENT. (PIERQUIN)

- R. Extract of Opium, dissolved in a little water, half a dram.
Calcined Oyster Shells, one dram.
Unguentum Populeum, four ounces.

Incorporate the whole with yolk of egg.

ANODYNE LINIMENT. (RATIER)

- R. Aqueous Extract of Opium, three drams.
Ointment of Marshmallows, one ounce.
Balsamum Tranquillum, } of each,
Oil of Almonds, } two ounces.

Mix.—This liniment is employed by BOYER.

SEDATIVE MIXTURE. (SWEDIAUR)

- R. Orange-flower Water, six ounces.
Extract of Opium, three grains.
Sulphuric Æther, one dram.
White Sugar, half an ounce.

Mix.—Dose, a spoonful, every hour or two hours.

ANODYNE COLLYRIUM. (FULD.; ELLIS; VAN MONS)

- R. Mucilage of Quince, one ounce.
Aqueous Extract of Opium, one grain.
Camphor, triturated with a drop of Oil, two grains.
Rose Water, two ounces.

Mix the extract and camphor, and make an emulsion with a part of the mucilage; then add alternatively the water and the mucilage by portions. (FULD.; VAN MONS)

ELLIS prescribes one grain of opium, five grains of camphor, and one ounce of mucilage of gum arabic.

COMPOUND LOHOCH. (RATIER)

- R. Simple Lohoch, four ounces.
Extract of Opium, one grain.
Cinchona, four grains.
Camphor, six grains.

Mix.—It is employed in bronchitis accompanied with violent cough and inordinate secretion of mucus.

II. Preparations which contain Opium in substance, either crude or purified.

A.

Preparations in which the action of the Opium is not modified, at least sensibly, by other Substances.

OPIATE POWDER.

Pulvis opiatius seu thebaicus seu Cretæ compositus cum Opio seu Cornu Cervi usti cum Opio. (LOND.; DUBL.; EDINB.; MIL. COPEN.; GENEV.; HANN.; Bo-

RUS.; SUEC.; WURCEB.; BRUGNATELLI; COXE; ELLIS; VAN MONS)

Pulvis Cornu usti cum Opio. (LOND.)

- R. Hard Opium, in powder one dram.
Prepared Burnt Hartshorn, one ounce.
Cochineal, one dram.
Mix.

Pulvis opiatius. (EDINB.)

- R. Opium, one part.
Prepared Carbonate of Lime, nine parts.
Mix intimately.

Pulvis Cretæ compositus cum Opio. (LOND.; DUBL.)

- R. Compound Powder of Chalk, six ounces and a half.
Hard Opium, in powder, four scruples.
Mix.

COXE; and VAN MONS also give the latter formula.

- R. Opium, one part.
White Sugar, twenty parts.

Mix. (MIL. COPEN.; WURCEB.)

GENEV. prescribes one part of opium, and twenty-three parts of sugar.

- R. Opium, one part.
Compound Powder of Tragacanth, nine parts.

Mix. (HANN.; BORUS.)

- R. Opium, one part.
White Sugar, two parts.
Compound Powder of Tragacanth, twelve parts.

Mix. (SUEC.)

- R. Opium, one part.
Calcined Hartshorn, nine parts.

Mix. (BRUGNATELLI)

- R. Prepared Chalk, one dram and a half.
Cinnamon, one dram.
Long Pepper, eight grains.
Opium, six grains.

Make twelve parcels. (ELLIS)

OPIATE PILLS.

Pilulæ opiatæ seu ex Opio. (EDINB.; AMER.; MIL. COPEN.; FERR.; HANN.; CAST. BORUS.; WURCEB.; AUGUSTIN;

BRUGNATELLI; COXE; ELLIS; SAUNDERS; VAN MONS)

Pilulæ Opiatæ, olim *Pilulæ thebaicæ*.
(EDINB.; BRUGNATELLI)

R. Opium, one part.
Extract of Liquorice, seven parts.
Pimento, in powder, two parts.

Soften the two first with dilute alcohol, and then incorporate the whole.

R. Opium, ten grains.
Extract of Liquorice, one dram.

Make forty pills. (WURCEB.)

MIL. COPEN. prescribes ten grains of opium, and one scruple of extract of liquorice.—CAST. BORUS.; and AUGUSTIN—ten grains of opium, and four scruples of the extract.—HANN.—one part of opium, and four parts of the extract.—SARD.—one of opium, and three of the extract.—FERR.—one of opium, and six of the extract.

R. Opium, four parts.
Water, nine parts.

Dissolve, strain, and add

Crumb of Bread, twelve parts.

Set apart for some time; express, and add

Starch, three parts.
Sugar, one part.

Make pills. (VAN MONS)

R. Opium, one dram.
Soap, twelve grains.
Water, a sufficient quantity
to make sixty pills. (AMER.; COXE; ELLIS)

ANTI-EPILEPTIC CONFECTION.

Conditum anti-epilepticum. (VAN MONS)

R. Opium, half a dram.
Valerian, one ounce and a half.
Syrup of Poppies, a sufficient quantity.

Mix.—Dose, a tea-spoonful, twice or thrice a day.

OPIATE OINTMENT. (BATAV.; CADET DE GASSICOURT; ELLIS; PIERQUIN; RATIER; VAN MONS)

R. Opium, in powder, one or two drams.
Simple Cerate, four ounces.

Mix. (RATIER)

VAN MONS prescribes one part of opium, and eighteen parts of cerate.

R. Olive Oil, two ounces and a half.
Yellow Wax, two ounces.

Melt together, and add
Opium, two grains.

Mix. (BATAV.)

R. Opium, twenty grains.
Yolk of Egg, one.

Mix, and add
GALEN'S Cerate, two ounces.

Mix by trituration. (CADET DE GASSICOURT; PIERQUIN)

R. Opium, five grains.
Rhubarb, ten grains.
Simple Cerate, one ounce and a half.

Triturate together. (ELLIS)

This ointment is used to apply to wounds, tumours, excoriations, and pustules which occasion much pain.

NARCOTIC OINTMENT.

Unguentum sedativo-narcoticum. (BRERA)

R. Opium, one dram.
Calf's Gastric Juice, a sufficient quantity.

Digest for twenty-four hours in a moderate heat, and then add a sufficient quantity of lard to make a soft ointment.

Half a dram or one dram of this is rubbed upon the epigastric region, in gastrodynia, cardialgia, spasmodic colic, hysteria, and hypochondria.

ANODYNE OINTMENT. (GENEV.; PIERQUIN)

R. Olive Oil, three ounces one dram and a half.
Wax, six drams and a half.
Powdered Opium, one ounce.

Boil for a quarter of an hour, and strain with expression; boil the residue in four ounces of water till reduced to two, and incorporate the decoction with the cerate. (PIERQUIN)

GENEV. prescribes one dram of opium, ten drams of oil, and half an ounce of wax.

It is recommended in asthenic nervous affections.

SOOTHING OINTMENT. (SAINTE-MARIE)

R. Oil of Roses, }
Lard, } of each, two ounces.
Yellow Wax, }

Melt over a slow fire, and add

Purified Opium, one dram.

Mix.—TRONCHIN applied this to painful enlargements of the breast, renewing it every twelve hours.

SEDATIVE SUPPOSITORY. (ELLIS)

R. Opium, two grains.
White Soap, four grains.

Mix.—To be introduced into the rectum when lavements are inadmissible.

OPIATE PLASTER. (LOND. ; EDINB. ; FULD. ; LIPP.)

Emplastrum Opii. (LOND. ; EDINB.)

R. Opium, in powder, half an ounce.
Resin of Spruce Fir, three ounces.
Lead Plaster, one pound.

Melt the two last, and mix the whole.

R. Melilot Plaster, sixteen parts.

Melt slowly, and when it has become nearly cold, add

Opium, in powder, one part.

Mix. (LIPP.)

FULD. prescribes one part of opium, and twenty-four of plaster.

LINIMENT OF OPIUM WITH CALF'S GALL.

Unguentum opiatum. (AUGUSTIN; NIEMANN)

R. Opium, in powder, two drams.
Calf's Gall, two ounces.

Digest for two days, and add

Lard, two ounces.
Oil of Bergamotte, ten drops.

Mix.—It is recommended by BRERA, as an excellent anodyne.

B.

Preparations in which the action of Opium is more or less modified, and in some instances completely masked by that of stimulant substances.

OPIATE PECTORAL TROCHISCS. (EDINB. ; AMER. ; HERBIP. ; COXE)

Trochisci Glycyrrhizæ cum Opi.
(EDINB. ; AMER. ; COXE)

R. Opium, two drams
Tincture of Tolu, half an ounce.
Simple Syrup, eight ounces.
Extract of Liquorice, } of each,
softened with hot water, } five
Gum Arabic, in powder, } ounces.

Triturate the opium with the tinc-
VOL. II.

ture; then add gradually the syrup and the extract, and sprinkle in the powdered gum arabic: finally, dry the mass and form it into trochiscs of ten grains each.

R. Opium, } of each, one dram.
Balsam of Peru, }
White Sugar, two drams.
Mass of Black Pectoral Trochiscs, seven ounces.

Triturate the opium with the balsam and sugar; then add the mass softened with hot water, and make trochiscs of from ten to fifteen grains. (HERBIP.)

ANTI-EMETIC PILLS. (PIERQUIN)

R. Purified Opium, four grains.
Calumba, four scruples.
Oil of Peppermint, ten drops.
Syrup of Clove Pink, a sufficient quantity.

Make thirty pills.—Dose, two, thrice a day.

SEDATIVE POWDER.

Pulvis Nitri thebaicus seu sedativus. (BATAV. ; FULD. ; LIPP. ; SUEC. ; SPIELMANN)

R. Opium, one part.
Nitrate of Potass, fifteen parts.
Sugar, forty-five parts.

Mix. (FULD.)

BATAV. ; and SUEC. prescribe one part of opium, eleven parts of nitre, and forty-eight of sugar.—SPIELMANN—one of opium, ten of nitre, and forty-eight of sugar.—LIPP.—one of opium, twelve of nitre, and forty-eight of sugar.

ANODYNE POWDER. (HAMB.)

R. Opium, four grains.
Antimoniated Magnesia, two drams.

Divide into eight parcels. It is recommended in rheumatism.

ANTI-DYSENTERIC POWDER. (VAN MONS)

R. Opium, } of each,
Tartarized Antimony, } one part.
Gum Arabic, } of each, eight parts.
White Sugar, }

Mix.—For the same purpose as the preceding.

ASTRINGENT POWDER.

Pulvis adstringens. (BRERA)

R. Opium, } of each, one grain.
Ipecacuanha, }
Cascarilla, ten grains.

Mix.—To be taken three or four times a day, in obstinate diarrhœa from atony.

STORAX PILLS.

Pilulæ e Storace. (DUBL.; BRUNS.; GENEV.; PALAT.; SARD.; SUEC.; WIRTEM.; HERBIP.; SPIELMANN.; SWEDIAUR.; VAN MONS)

- R. Purified Storax, three drams.
Soft Purified Opium, } of each,
Saffron, } one dram.

Mix intimately together. (DUBL.)

- R. Storax, half an ounce.
Extract of Liquorice, three drams.
Olibanum, }
Benzoin, } of each,
Mastic, } two drams.
Extract of Opium, }
Saffron, } one dram.
Essence of Amber, a sufficient quantity.

Make a pilular mass. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN.)

SARD. prescribes half an ounce of storax, two drams of extract of liquorice, one dram each, of olibanum, of benzoin, of myrrh, and of opium, two scruples of saffron, and a sufficient quantity of syrup of poppies.—VAN MONS—one part each, of benzoin, of myrrh, of olibanum, of opium, and of saffron, one and a half of extract of liquorice, two parts of storax, and a sufficient quantity of simple syrup.

- R. Storax, two ounces.
Extract of Liquorice, } of each,
Saffron, } six drams.
Opium, } five drams.
Balsam of Peru, a sufficient quantity.

Make a pilular mass. (HERBIP.)

- R. Storax, } of each,
Saffron, } one ounce.
Extract of Liquorice, }
Opium, } five drams.
Tincture of Myrrh, a sufficient quantity.

Mix. (SWEDIAUR.)

- R. Storax, two drams.
Extract of Liquorice, three drams.
Opium, one dram.
Spanish Wine, a sufficient quantity.

Mix. (SUEC.)

- R. Storax, } of each, half an ounce.
Olibanum, }
Myrrh, }
Opium, } of each,
Saffron, } one dram.
Extract of Liquorice, }
Syrup of Poppies, a sufficient quantity.

Mix. (GENEV.)

It is stimulant and expectorant.—Dose, from four to twelve grains.

COMPOUND POWDER OF IPECACUANHA.

Pulvis alexiterius seu anodymus seu diaphoreticus seu diapnoicus seu sudorificus seu sudorificus Doveri, Pulvis Doveri, Pulvis Ipecacuanhæ opiatius seu thebaicus seu Ipecacuanhæ et Opii seu compositus seu Ipecacuanhæ cum Opio. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; GENEV.; HAMB.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BRERA.; BRUGNATELLI.; CADET DE GASSICOURT.; COXE.; ELLIS.; PIERQUIN.; SPIELMANN.; SAUNDERS.; SWEDIAUR.; PH. SYPH.; VAN MONS)

Pulvis Ipecacuanhæ compositus.
(LOND.; DUBL.)

Pulvis Ipecacuanhæ et Opii. (EDINB.)

- R. Ipecacuanha, } of each,
Pure Hard Opium, } one part.
Sulphate of Potass, eight parts.

Mix.

LOND. requires the ingredients to be separately powdered: EDINB.; and DUBL. direct them to be pulverized together.

AMER.; BAVAR.; FENN.; HAMB.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; SUEC.; WIRTEM.; BRUGNATELLI.; and COXE give the same formula.—AMST.; BELG.; LIPP.; ROSS.; SAX.; and HERBIP. prescribe one part each, of opium and of ipecacuanha, and nine of sulphate of potass.—BATAV.—one each, of opium, and of ipecacuanha, and ten of sulphate.

- R. Purified Opium, one part.
Ipecacuanha, two parts.
White Sugar, seven parts.

Pulverize and mix. (FERR.; FORM. HÔP. FR.)

AUSTR.; DAN.; and WIRTEM. prescribe one part each, of opium and of ipecacuanha, and eight parts of sugar.—MIL. COPEN.—one each, of opium and of ipecacuanha, and twenty of sugar.—PIERQUIN—one of opium, two of ipecacuanha, and eighteen of sugar.

- R. Nitrate of Potass, } of each,
Sulphate of Potass, } four parts.

Pulverize them, and fuse them in a crucible; pour the fluid mass into an iron mortar, and when nearly cold, add

Powdered Opium, } of each,
Ipecacuanha, } one part.

Triturate to a very fine powder. (GENEV.; PH. SYPH.; CADET DE GASSICOURT; SWEDIAUR; SAUNDERS)

HAMB.; and ELLIS prescribe eight parts each, of the two salts, and one part each, of the two other substances.—VAN MONS—four each, of the nitrate, of the sulphate, and of ipecacuanha, and one of opium.—HISP.—four of sulphate, six of nitrate, and one each, of ipecacuanha and of opium.

R. Sulphate of Potass, } of each,
Nitrate of Potass, } four parts.

Add to the melted and cool mass,

Powdered Extract of Opium, } of each,
Ipecacuanha, } one
Liquorice, } part.

Pulverize. (GALL.; SPIELMANN)

VAN MONS prescribes four parts of sulphate, three of nitrate, and one part each, of the three powders.

This last is the original formula of DOVER.

The compound powder of ipecacuanha, most frequently nauseous and emetic, becomes sudorific when the patient is warmly covered, and takes warm drinks. The dose, which, without speaking of individual or circumstantial conditions, varies according to the relative proportion of opium, is in general from ten grains to half a dram.

DIAPHORETIC AND SEDATIVE POWDER. (BRERA)

R. DOVER's Powder, } of each,
JAMES' Powder, } four grains.

Mix.—To be taken every four hours, in obstinate pains attending rheumatism, syphilis, or gout.

SUDORIFIC BOLUSES. (BRERA; SWEDIAUR)

R. DOVER's Powder, }
from twenty to twenty-five grains.
Syrup of Ginger, a sufficient quantity.

Make a bolus.—To be taken in the morning. (SWEDIAUR)

BRERA prescribes sixteen grains of the powder, and a sufficient quantity of elder rob to make four boluses; or, twelve grains of the powder, and six of extract of aconite, to make six boluses; one to be taken every two hours.

ANODYNE POWDER. (HAMB.)

R. Magnesia, } one dram.
Oxide of Zinc, } sixteen grains.
Ipecacuanha, }
Opium, } of each, two drams.

Mix.—This nearly resembles DOVER's powder.

ANTI-OTODYNIC MIXTURE. (PIERQUIN)

R. Opium, three or four grains.
Saffron, half a scruple.
Myrrh, half a dram.
Juice of Mallow, half an ounce.
Oil of Almonds, two ounces.

Mix.

PIERQUIN also gives another formula:—two grains each, of camphor and of opium, and one ounce and a half of oil.

Both of these are to be injected warm into the ear.

ANTI-ODONTALGIC MIXTURE. (SAUNDERS)

R. Opium, half a grain.
Oil of Cloves, ten drops.
HOFFMANN's Liquor, one dram and a half.

Mix.

ANODYNE OINTMENT. (FULD.; HERBIP.)

R. Opium, } of each, one scruple.
Saffron, }
Yolk of Egg, one.
Unguentum Populeum } of each,
Nutritum, } one ounce.

Mix. (FULD.)

HERBIP. prescribes three ounces each, of saturnine cerate and of unguentum populeum, one dram each, of opium and of saffron, and three yolks of eggs.

It is principally recommended as an application to painful piles.

SAPONACEOUS PILLS OF OPIUM. (LOND.; AMER.; HERBIP.; COXE)

Pilulæ Saponis cum Opio. (LOND.)

R. Hard Opium, in powder, half an ounce.
— Soap, two ounces.

Make a pilular mass.

R. Purified Opium, } of each,
Root of Liquorice, } half an
White Hellebore, } ounce.
Black Hellebore, }

STARKEY's Soap, a sufficient quantity.

Make a pilular mass. (HERBIP.)

ANTI-SPASMODIC CATAPLASM. (CADET DE GASSICOURT; PIERQUIN)

R. Cataplasm of Linseed Meal, made with decoction of Poppy Heads, four ounces.

Sprinkle it with

Powdered Opium, twenty grains.
Camphor, half a dram.

ANTI-SPASMODIC PLASTER. (AUGUSTIN ; ELLIS ; SAUNDERS)

- R. Cumin Plaster, } of each, one ounce.
 Yellow Wax, }
 Opium, } of each, two drams.
 Camphor, }

Mix. (SAUNDERS)

- R. Opium, } of each, half a dram.
 Camphor, }
 Litharge Plaster, } two drams.

Mix. (ELLIS)

- R. Opium, } of each, one dram.
 Camphor, }
 White Soap, }
 Tincture of Opium, a sufficient quantity
 to make a soft mass. (ELLIS)

- R. Opium, } of each,
 Extract of Henbane, } two drams.
 Hemlock, } half an ounce.

Mix. (AUGUSTIN)

- R. Opium, } of each,
 Sub-carbonate of } half a dram.
 Ammonia, }
 Camphor, } two drams.
 Petroleum, }
 Crocated Plaster of Galbanum, } one ounce.

Mix. (AUGUSTIN)

ANTI-ODONTALGIC BALSAM. (PALAT. ; WIRTEM. ; AUGUSTIN ; PIERQUIN ; SAINTE-MARIE ; SPIELMANN ; VAN MONS)

- R. Opium, } of each, two scruples.
 Camphor, }

Dissolve them in a little alcohol, and add

- Oil of Cloves, } one dram.
 Guaiac, } two drams.
 Nutmeg, } six drams.

Mix. (PALAT. ; WIRTEM. ; CADET DE GASSICOURT ; PIERQUIN ; SPIELMANN ; VAN MONS)

- R. Opium, } of each, one scruple.
 Camphor, }
 Cajeput Oil, } one dram.
 Oil of Nutmeg, } three drams.

Mix. (SAINTE-MARIE)

A small quantity of this balsam is to be put into the hollow of the carious tooth.

To this preparation may be resembled that which BRUNS. ; and SPIELMANN give under the title of *Sotirella parva*, of which the following is the formula :

- R. Opium, } of each,
 Species Diambrae, } one dram.
 Camphor, }
 Saffron, } of each, twelve grains.
 Oil of Cloves, } thirty drops.
 Nutmeg, } two drams.

Mix.—It is to be used in the same manner.

- R. Opium, } half a dram.
 Extract of Belladonna, } of each,
 Camphor, } six grains.
 Cajeput Oil, } of each,
 Tincture of Cantharides, } eight drops.
 Oil of Henbane, } one dram,

Mix.—To be used in the same manner. (AUGUSTIN)

- R. Powdered Mastic, } of each,
 Sandarac, } two drams.
 Dragon's Blood, } ten grains.
 Opium, } two grains.
 Oil of Rosemary, } eight drops.
 Spirit of Scurvy-Grass, } a sufficient quantity.

Mix.—A little to be applied to the painful gum. (AUGUSTIN)

ANTI-SPASMODIC PILLS. (FULD.)

- R. Opium, } of each, fifteen grains.
 Castor, }
 Camphor, } half a dram.
 Extract of Valerian, } of each,
 Fetid Goose- } two drams.
 foot, }
 Tincture of Milfoil, } a sufficient quantity.

Mix.—Dose, from ten to twenty grains.

ANTI-SPASMODIC BOLUS. (PIERQUIN)

- R. Opium, } half a grain,
 Camphor, } three grains.
 Nitrate of Potass, } ten grains.
 Conserve of Roses, } a sufficient quantity.

Mix.—For one dose.

CHAUSSIER'S PASTILLES. (CADET DE GASSICOURT)

- R. Opium, } six grains.
 Camphor, } twenty-one grains.
 Sugar, } three drams.
 Mucilage, } a sufficient quantity,

to make fifty pastilles.—They are recommended in croup and lazyngal phthisis.—Dose, five or six, daily.

CHAUSSIER'S TROCHISCS. (CADET DE GASSICOURT ; SWEDIAUR)

- R. Opium, } one scruple.
 Camphor, } one dram.
 Sugar, } nine drams.
 Gum Tragacanth, } half a dram.
 Water, } a sufficient quantity.

Make one hundred and fifty trochiscs.—Dose, four or six, daily.

ANTI-SPASMODIC POWDER. (VAN MONS)

- R. Opium, } half a scruple.
 Camphor, } two scruples.

Subcarbonate of Ammonia, four scruples
 Starch, fifteen grains.
 Divide into eighteen parcels.—Dose, one
 parcel every hour, in mint water.

ANTI-CANCEROUS BALSAM. (BORIES)

R. Opium, twenty-four grains.
 Acetate of Lead, four ounces.
 Camphor, one dram.
 Spirit of Turpentine, twelve ounces.
 Digest for two days, and strain.

RENAL OINTMENT. (PALAT.)

R. Opium, one scruple.
 Camphor, half a scruple.
 Oil of Scorpions, } of each
 Lemon Juice, } half an ounce.
 Unguentum Populeum, one ounce and a half.
 Mix.—To be applied to the loins in the
 passing of renal calculi.

ANODYNE PLASTER. (BRUNS.; SPIEL-
 MANN)

R. Soap Plaster, } of each, four ounces.
 Yellow Wax, }
 Melt together and add to the mass when
 nearly cold,
 Deliquescent Subcarbonate of
 Potass, five drams.
 Opium, in powder, half an ounce.
 Camphor, three drams.
 Oil of Henbane Seeds, five drams.
 Mix.—It is recommended in diseases of
 the joints.

ORIENTAL PILLS. (CADET DE GASSI-
 COURT)

R. Purified Opium, }
 Saffron, } of each,
 Cinnamon, } an equal part.
 Nutmeg, }
 Lesser Cardamoms, }
 Syrup of Orange Flowers, a sufficient quantity.
 Make pills of three grains each.—Dose,
 two or three in the evening.

TONIC AND HYPNOTIC PILLS. (PIERQUIN)

R. Opium, }
 Olibanum, } of each,
 Storax, } two scruples,
 Myrrh, }
 Extract of Liquorice, }
 Starch, two drams.
 Gum Tragacanth, } of each,
 — Arabic, } half an ounce.
 Mix.—Dose, from half a scruple to half
 a dram.

COMPOUND PILLS OF HOUNDSTONGUE.

Pilule de Cynoglossa seu Opii composite,
Catapotia ad Catarrhos; Pilules adou-
cissantes de Mésué. (ANT.; BELG.;
 BRUNS.; HISP.; GALL.; FORM. HÔP. FR.;
 HANN.; PALAT.; SARD.; WIRTEM.;
 CADET DE GASSICOURT; NIEMANN;
 PIERQUIN; SPIELMANN; VAN MONS)

1. *Without Castor.*

R. Dried Bark of the Root of Hounds-
 tongue, one ounce.
 Seeds of White Henbane, } of each,
 Opium, } two drams.
 Olibanum, six drams.
 Saffron, one dram.
 Peruvian Balsam, a sufficient quantity.

Mix the opium, pulverized separately,
 with the other substances, pulverized to-
 gether, and gradually add the balsam to
 form a pilular mass. (SARD.)

R. Root of Houndstongue, } of each,
 Seeds of Henbane, } half
 Opium, } an ounce.
 Myrrh, six drams.
 Olibanum, five drams.
 Storax, }
 Cloves, } of each, two drams.
 Cinnamon, }
 Saffron, one dram and a half.
 Syrup of Violets, a sufficient quantity

to make a pilular mass, one dram of which
 contains eight grains each, of opium and of
 henbane. (SPIELMANN)

BRUNS.; PALAT.; WIRTEM.; and NIE-
 MANN give the same formula, but NIEMANN
 prescribes but two drams each, of myrrh,
 and of olibanum.—BRUNS.; PALAT.; and
 WIRTEM. prescribe an equal part each, of
 syrup of violets and of juice of hounds-
 tongue.

R. Root of Houndstongue, }
 Seeds of Henbane, } of each,
 Opium, } one ounce.
 Myrrh, } of each,
 Olibanum, } one ounce and a half.
 Storax, }
 Cloves, } of each, half an ounce.
 Cinnamon, }
 Simple Syrup, a sufficient quantity.
 Make a pilular mass. (HANN.)

2. *With Castor.*

R. Aqueous Extract of Opium, three drams.
 Bark of the Root of Houndstongue, two ounces.
 Saffron, five drams.
 Castor, six drams.
 Simple Syrup, nine drams.

Make a mass, one scruple of which contains two grains of opium. (Hisp.)

This is the only pharmacopœia which omits the henbane.

- R. Root of Houndstongue, } of each,
 Seeds of Henbane, } four
 Vinous Extract of Opium, } parts.
 Castor, } of each,
 Saffron, } one part and a half.
 Olibanum, } five parts.
 Myrrh, } six parts.
 Simple Syrup, a sufficient quantity.

Make a pilular mass. (VAN MONS)

- R. Myrrh, one ounce and a half.
 Olibanum, ten drams.
 Root of Houndstongue, } of each,
 Seeds of Henbane, } one ounce.
 Opium, }
 Castor, } of each, three drams.
 Saffron, }
 Simple Syrup, a sufficient quantity.

Make a pilular mass. (BELG.)

- R. Root of Houndstongue, } of each,
 Seeds of White Henbane, } half
 Purified Opium, } an ounce.
 Myrrh, } six drams.
 Olibanum, } five drams.
 Saffron, } of each, two drams.
 Castor, }
 Syrup of Violets, } of each, a
 —Houndstongue, } sufficient
 quantity.

Make a pilular mass. (BRUNS.; PALAT.; WIRTEM.)

GALL.; FORM. HÔP. FR.; CADET DE GASSICOURT; and PIERQUIN give the same formula, but substitute for the opium the vinous extract, and the two syrups by that of opium; they also diminish the quantity of saffron and of castor by half a dram, and assign a weight of four grains to the pills.

- R. Myrrh, six drams.
 Olibanum, five drams.
 Bark of the Root of Houndstongue, half an ounce.
 Saffron, } of each,
 Castor, } one dram and a half.
 Seeds of Henbane, half an ounce.

Pulverize each separately; then mix and sift the whole. Then

- R. Purified Opium, half an ounce.
 Distilled Water, two drams.

Soften, in a marble mortar, with a slight heat, and add

Syrup of Opium, one ounce.

Finally incorporate the powder, adding more syrup of opium, so as to form a pilular mass. (ANT.)

These celebrated pills are in very great use as sedative, in catarrhs, hæmoptysis, and strangury.—Dose, from three to five grains.

ANTI-COLICAL PLASTER.

Emplastrum ad Tormina Ventris. (SWEDIAUR)

- R. Lead Plaster, two ounces.
 Galbanum, one ounce.
 Oil of Caraway, two drams.

Melt, and add

Opium, one dram.

Mix.—It is recommended in flatulent colic, diarrhœa, and dysentery.

ANODYNE PLASTER. (SPIELMANN)

- R. Emplastrum Diapompholigos, one ounce.
 Soot, four scruples.
 Opium, one dram.
 Subcarbonate of Ammonia, two scruples.

Knead the whole well with a sufficient quantity of the oil of solanum nigrum. DECKER recommended it in cancer.

ODONTALGIC PLASTER. (BRUNS.; SARD.; WIRTEM.)

- R. Tacamahac, } of each, one ounce.
 Elemi, }
 Mastic, }
 Opium, two drams.

Mix, by trituration, with

Venice Turpentine, two drams.

It is recommended to be applied in tooth-ach, head-ach, and catarrhal discharges. It is spread, with a warm spatula, upon silk, and thus applied to the cheeks or temples.

We may compare with this plaster, that which BRUNS. gives under the title of *Emplastrum ad Tempora*, of which the following is the formula:

- R. Gum Anime, }
 — Caranna, } of each, two drams.
 — Elemi, }
 Mastic, }
 Storax, }
 Amber, }
 Turpentine, three drams.
 Opium, } of each, one dram.
 Balsam of Peru, }
 Oil of Cloves, } of each, three drops.
 — Amber, }

Mix.

CADET DE GASSICOURT also gives the following formula:

- R. Hard Opium, one ounce.
 Mastic, } of each,
 Sandarac, } two drams seven grains.
 Dragon's Blood, half a dram.

Pulverize each separately, mix them together, and add eight drams of oil of rose-

mary; then triturate in a mortar, adding gradually a sufficient quantity of spirit of scurvy grass to obtain a soft mass.

It will be proper to place among these various preparations, those which are found described in BATAV.; BELG.; HANN.; OLDEN.; BORUS.; SARD.; SPIELMANN; and VAN MONS, under the names of *Emplastrum opiatum*, and of *Emplastrum cephalicum*, of which the following are the formula:

R. Elemi, }
Mastic, } of each, four ounces.
Olibanum, }
Benzoin, two ounces.
Turpentine, six ounces.

Melt together in a water-bath, and add

Balsam of Peru, half an ounce.
Opium, one ounce.

Mix. (BATAV.; BELG.; HANN.; OLDEN.; BORUS.; SAX.)

VAN MONS prescribes one part of Peruvian balsam, two parts of opium, four of benzoin, and eight each, of elemi, of mastic, of olibanum, and of turpentine.

R. Tacamahac, }
Storax, } of each, eight parts.
Mastic, }
Caranna, }
Peruvian Balsam, four parts.
Opium, }
Oil of Amber, } of each, one part.

First mix together the tacamahac, mastic, and caranna, in a heated mortar; then incorporate the storax and balsam, and finally add the opium and oil. (VAN MONS)

VAN MONS also prescribes one part of opium, two parts each, of Peruvian balsam, of benzoin, and of turpentine, and four each, of elemi, of olibanum, and of tacamahac.—SPIELMANN—one ounce each, of tacamahac, of caranna, of storax, and of mastic, two drams of turpentine, four scruples of opium, half an ounce of Peruvian balsam, and one dram of oil of amber.

HYPNOTIC OINTMENT. (WIRTEM.)

R. Alabastrine Ointment, } of each,
Rose Ointment, } half an ounce.
Poplar Ointment, two ounces.
Oil of Henbane, half an ounce.
Oil of Walnuts, one dram and a half.
Opium, dissolved in Rose Water, one dram.
Saffron, half a dram.
White Wax, half an ounce.

Make an ointment.—It used formerly to be applied to the temples and wrists spread upon linen.

BASILIC PLASTER. (WIRTEM.)

R. Yellow Wax, one pound and a half.
Resin of the Pine, half a pound.

Melt slowly together, and strain; when the mass is nearly cold, add, in powder,

Tacamahac, }
Elemi, } of each, one ounce.
Olibanum, }
Dragon's Blood, }

When these are mixed, add further,

Gum Ammoniac, }
Galbanum, }
Opoponax, } of each, one ounce.
Bdellium, }
Assafœtida, }
Gum Hedera, }
Extract of Opium, } of each,
Saffron, } two drams.

The whole dissolved in vinegar. Express, evaporate to the consistence of honey, and add, constantly stirring,

Venice Turpentine, one pound.
Gum Tragacanth, ten ounces.
Castor, half an ounce.
Subcarbonate of Ammonia, one ounce and a half.

Mix.—This used formerly to be principally applied in œdema.

TROCHISCS OF AMBER. (WIRTEM.)

R. Opium, } of each,
Saffron, } one dram fifteen grains.
Myrrh, }
Calcined Hartshorn, }
Red Coral, }
Gum Arabic, }
Tragacanth, } of each,
Mastic, } two drams.
Lac, }
Juice of Acacia, }
Hypocist, }
Flowers of Pomegranate, }
Seeds of Black Poppy, } six drams.
Amber, }
Mucilage of Fleaseed, }

a sufficient quantity to make a mass, to be divided into trochiscs.

These trochiscs, now nearly forgotten, resemble diascordium in their effects. They were extolled by MESUE in hæmorrhages, particularly in hæmoptysis and menorrhagia, and also in diarrhœa and excessive hæmorrhoidal discharge.—Each dram contains three grains of opium.

ANTI-DYSENTERIC PILLS. (LUSIT.)

R. Opium, } of
Extract of Catechu, } each,
Bistort, } two
Angelica, } drams.

Extract of Contrayerva,	} of each, two drams.	
— Sweet Flag,		
— Tormentil,		
— Zedoary,		
— Milfoil,		
— Scordium,		
— Pomegranate Bark,		
— Red Roses,		
— Pomegranate		
Flowers,		
— Saffron,		
— Myrrh,		
— Rhubarb,		
Oil of Nutmeg,	three drams.	
— Mace,	one dram.	
Balsam of Peru,	three drams and a half.	

Oil of Juniper, two drams.

Make a pilular mass.—Dose, from four to eight grains, three or four times a day.

MITHRIDATE.

Electuarium mithridaticum, Antidotus mithridatica Domocratis. (PALAT.; SARD.; WIRTEM.; SPIELMANN)

R. Myrrh,	} of each, ten ounces.	
— Saffron,		
— Agaric,		
— Ginger,		
— Cinnamon,		
— Spikenard,		
— Olibanum,		
Seeds of Thlaspi,		
— Carrot,	seven ounces.	
— Fennel,	} of each, five ounces.	
— Anise,		
— Parsley,		
— Seseli,	} of each, two ounces and a half.	
— St. John's-wort,		
Herb of Lemon-grass,	eight ounces.	
— Pennyroyal,	of each,	
— Scordium,	seven ounces.	
Root of Arabian Costus,	of each,	
— Gentian,	eight ounces.	
— Sweet Flag,	} of each, three ounces.	
— Asarum,		
— Valerian,		
— Ligusticum Meum,	two ounces and a half.	
Storax,	} of each, eight ounces.	
Long Pepper,		
White Pepper,	seven ounces.	
Castor,	two ounces and a half.	
Juice of Acacia,	of each,	
— Hypocist,	eight ounces.	
Leaves of Laurus Cassia,	ounces.	
Fruit of Amyris Opobalsamum,	seven ounces.	
Gum Arabic,	} of each, five ounces.	
Opium,		
Lesser Cardamoms,		
Petals of Red Roses,		
Cypheos Trochiscs,		seven ounces.

Pulverize, mix, and add,
 Galbanum, } of each, eight ounces.
 Opoponax, }
 Bdelium, five ounces.
 Sagapenum, three ounces.

mixed perfectly with
 Spanish Wine, forty-eight ounces.

Incorporate the mixture with
 Honey, eight hundred and sixty ounces.

previously triturated with
 Opobalsamum, } of each, eight ounces.
 Turpentine, }

Mix.—(SPIELMANN)

This formula is from GALEN.

R. Root of Round Birth-wort,	} of each, four ounces and a half.	
— Valerian,		
— Sweet Flag,	five ounces three drams.	
— Gentian,	seven ounces and a half.	
— Ligusticum Meum,	three ounces six drams.	
— Ginger,	fifteen ounces.	
Herb of Dittany of Crete,	seven ounces and a half.	
— Pennyroyal,	of each, ten ounces and a half.	
— Scordium,	ounces and a half.	
Leaves of Laurus Cassia,	twelve ounces.	
Flowers of St. John's-wort,	three ounces and a half.	
— French Lavender,	twelve ounces.	
— Red,	of each, seven ounces and a half.	
— Roses,	a half.	
— Celtic Nard,	fifteen ounces.	
— Spikenard,	thirteen ounces.	
— Lemon-grass,	fifteen ounces.	
Seeds of Thlaspi,	twelve ounces.	
— Seseli,	ten ounces and a half.	
— Carrot,	of each, seven ounces and a half.	
— Parsley,	of each, seven ounces and a half.	
— Fennel,	of each, four ounces and a half.	
— Anise,	one ounce.	
Juniper,	one ounce.	
Long Pepper,	twelve ounces.	
White Pepper,	of each, ten ounces and a half.	
Fruit of Amyris Opobalsamum,	half.	
Lesser Cardamoms,	seven ounces and a half.	
Saffron,	fifteen ounces.	
Cinnamon,	fifteen ounces and a half.	
Arabian Costus,	twelve ounces.	
Cassia Lignea,	ten ounces and a half.	
Trochiscs of Agaric,	fifteen ounces.	
Castor,	twelve ounces.	
Scincus Marinus,	three ounces and a half.	
Myrrh,	sixteen ounces.	
Olibanum,	fifteen ounces.	
Bdelium,	ten ounces and a half.	

Gum Arabic, seven ounces and a half.

Pulverize, mix, and sift. Then dissolve in eight pounds of wine,

Galbanum,	} of each, twelve ounces.
Opoponax,	
Sagapenum,	four ounces and a half.
Juice of Hypocist,	twelve ounces.
Acacia,	four ounces.
Opium,	seven ounces and a half.

Mix the solution with

Despumated Honey,
one hundred and six pounds.

Incorporate the powder gradually. Then pour into the mixture,

Storax, twelve ounces,
dissolved in

Turpentine, fourteen ounces.

Add, finally,

Opobalsamum, twelve ounces

Stir again for several hours, and leave it to ferment in a large vessel. (WIRTEM.)

In the following formula, the ingredients of the electuary are classed in the same manner as a dose in making theriac.

1st class.—

Anise,	two drams and a half.
Ginger,	ten drams.
Mace,	} of each, half an ounce.
Cloves,	
Gentian,	five drams.

The total weight is three ounces one dram and a half.

2nd class.—

Scordium,	seven drams.
Celtic Nard,	} of each, five drams.
Seeds of Parsley,	
Long Pepper,	one ounce.

The total weight is three ounces one dram.

3rd class.—

Pennyroyal,	seven drams.
Seeds of Fennel,	five drams.
Valerian,	two drams and a half.

The total weight is one ounce six drams and a half.

4th class.—

Cinnamon,	ten drams.
Zedoary,	} of each, one ounce.
Seseli,	
Red Roses.	five drams.

The total weight is three ounces seven drams.

5th class.—

Dittany of Crete,	five drams.
Cassia Lignea,	} of each, seven drams.
Cubebs,	
White Pepper,	
Seeds of Thlaspi,	
	ten drams.

The total weight is four ounces and a half.

VOL. II.

6th class.—

Seeds of Carrot,	seven drams.
Lesser Cardamoms,	five drams.
Sweet Flag,	} of each, two
Round Birthwort,	
Angelica,	
St. John's-wort,	
	drams and a half.

The total weight is two ounces six drams and a half.

Each of these mixtures is to be reduced to powder, then each of the following substances are to be separately reduced to powder :

Myrrh,	} of each, ten drams.
Olibanum,	
Agaric,	
Saffron,	
Castor,	one ounce.
Bdellium,	seven drams.
Scincus Marinus, chopped,	two drams and a half.

Then take

Spikenard,	} of each, ten drams.
Lemon-grass,	
French Lavender,	one ounce.

Chop them small, mix them with mucilage of tragacanth, and dry and pulverize them.

Pulverize also,

Cyphos Trochiscs, seven drams ;
or their ingredients taken in proper proportions. Then dissolve

Opium,	} of each, five drams.
Gum Arabic,	
Juice of Hypocist,	one ounce.
Acacia,	two drams and a half,

in a sufficient quantity of wine of Crete. Then add,

Despumated Honey,

thrice the weight of the whole ;

first triturating it with

Turpentine, one ounce.

Incorporate gradually the powders, and finally, add

Storax, one ounce,
dissolved in oil of nutmeg and wine, and leave it to ferment. (PALAT.)

R. Castor,	} of each,
Saffron,	
Oil of Nutmeg,	
Amber,	
Chio Turpentine,	
Galbanum,	one ounce and a half.
Myrrh,	} of each,
Gum Arabic,	
Sweet Flag,	
Arabian Costus,	
Round Birthwort,	
Valerian,	one ounce.
Zedoary,	

2 R

Leaves of Dittany of Crete,	} of each, ten drams.
White Horehound,	
Rue,	
Wild Thyme,	
Pennyroyal,	
Flowers of Sweet Rush,	
Chamomile,	
Pennyroyal,	
French Lavender,	
Tansy,	
Seeds of Small Sandwort (<i>Sium Ammi</i>),	} six drams.
Carrot,	
Lesser Cardamoms,	} six drams.
Cubeb,	
Opium,	
Honey, thrice the weight of the whole.	
Madeira Wine, a sufficient quantity.	

Dissolve the opium in wine, add the solution to the honey heated, and then the oil of nutmeg mixed with the turpentine; then incorporate the powders, adding lastly, that of the saffron mixed with wine, and leave the mass to ferment. (SARD.)

The invention of this preparation is ascribed to MITHRIDATES, king of Pontus, whose recipe POMPEY brought to Rome, where DAMOCRATIS sung in praise of it. From its origin it has been considered alexipharmic, stomachic, comforting, anodyne, in a word, a panacea. Every modern pharmacopœia omits it, and it is sought for in the older ones only as a memento: but certainly it is not to reason but rather to caprice that we are to ascribe this reform. Mithridate has experienced the same fate as theriac: it has been prepared after a different formula from that of the original inventor, longer, more complicated, and differing also in every country. Such is the history of all panaceas; they vary according to times, places, and circumstances, but people's faith in them continues the same, till fashion, (to the caprices of which even faith is subject,) gives it another direction and object.

Each ounce of the mithridate of WIRTEM. contains three grains of opium, and four of saffron.

JULEP OF MITHRIDATE.

Julapium mithridaticum. (SAUNDERS)

R. Mithridate,	two drams.
Common Julep,	five ounces.
Mix.	

THERIAC, OPIATE ELECTUARY.

Theriaca, Electuarium theriacale magnum seu Opii compositum seu aromatico-opiatum seu opiatum seu thebaicum, Confectio Opii. (LOND.; EDINB.; AMST.; ANT.; BATAV.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HANN.; OLDEN.; LUSIT.;

PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; BRUGNATELLI.; COXE.; SPIELMANN.; SWEDIAUR.; VAN MONS)

The invention of the electuary to which NICANDER gave the name of *Theriaca* is generally attributed to ANDROMACH. PLINY says that the formula of a similar composition was inscribed in verse, three centuries before the Christian æra, upon the temple of Æsculapius, and that ANDROMACH merely imitated, by order of NERO, the composition of the mithridate which had then been for a long time celebrated. The following is ANDROMACH's formula as GALEN has transmitted it to us:

R. Root of Florentine Iris,	of each,
Liquorice,	twelve ounces.
Arabian Costus,	} of each,
Pontic Rhubarb,	
Cinquefoil,	six ounces.
Ligusticum Meum,	} of each,
Rhubarb,	
Gentian,	four ounces.
Birthwort,	two ounces.
Herb of Scordium,	twelve ounces.
Lemon-grass,	} of each,
Horehound,	
Dittany of Crete,	six ounces.
Calamint,	} of each,
Pennyroyal,	
Ground Pine,	four ounces.
Germander,	
Leaves of Laurus Cassia,	twelve ounces.
Flowers of Red Roses,	twelve ounces.
French Lavender,	six ounces.
St. John's-wort,	four ounces.
Lesser Centaury,	two ounces.
Saffron,	six ounces.
Fruit of Amyris Opobalsamum,	four ounces.
Cinnamon,	twelve ounces.
Cassia Lignea,	} of each, six ounces.
Spikenard,	
Celtic Nard,	four ounces.
Long Pepper,	twenty-four ounces.
Black Pepper,	} of each, six ounces.
Ginger,	
Cardamoms,	four ounces.
Seeds of Rape,	} of each,
Agaric,	
Seeds of Macedonian Parsley,	twelve ounces.
	six ounces.

Anise,	} of each,
Fennel,	
Cress,	
Seseli,	
Thlaspi,	
Amomum,	
Sandwort,	} two ounces.
Carrot,	
Opium,	twenty-four ounces.

Opobalsamum,	twelve ounces.
Myrrh,	} of each, six ounces.
Olibanum,	
Turpentine,	
Storax,	} of each, four ounces.
Gum Arabic,	
Sagapenum,	
Asphaltum,	} of each, two ounces.
Opoponax,	
Galbanum,	
Juice of Acacia,	} of each,
— Hypocist,	
Castor,	four ounces.
Lemnian Bole,	two ounces.
Calcined Vitriol,	of each,
Trochiscs of Squill,	four ounces.
	forty-eight ounces.
— Viper,	} of each,
— Sweet	
Flag,	twenty-four ounces.

Triturate the balsams, resins, and gums, in Wine, a sufficient quantity, to form a thin paste, and incorporate the whole with

Honey, nine hundred and sixty ounces. (SPIELMANN)

Another longer formula directs the ingredients to be divided into six classes, thus :

1st class.—

Root of Illyrian Iris,	} of each,
Tops of Cretan Scordium,	
Seeds of Rape,	
— Anise,	} of each,
— Seseli,	
Cretan Pennyroyal,	
Zedoary,	half an ounce.
	six drams.

The total weight is half a pound six drams.

2nd class.—

Root of Gentian,	} of each,
Herb of Germander,	
Seeds of Cretan Fennel,	
— Amomum,	} of each,
Root of Cinquefoil,	
Tops of Horehound,	
Calamint,	half an ounce.
	six drams.

The total weight is four ounces two drams.

3rd class.—

Root of Angelica,	} of each,
— Valerian,	
Celtic Nard,	
Herb of Ground Pine,	} half an ounce.
Seeds of Thlaspi,	
Long Birthwort,	
Dittany of Crete,	two drams.
	six drams.

The total weight is three ounces and a half.

4th class.—

Red Roses,	one ounce and a half.
Flowers of St. John's-wort,	half an ounce.

Flowers of Lesser Cen- taury,	} of each, three ounces.
Seeds of Cretan Carrot,	
— Macedonian Parsley,	
	six drams.
Trochiscs of Viper,	three ounces.
— Squill,	six ounces.

The total weight is ten ounces two drams.

5th class.—

Seeds of Sandwort,	} of each,
Sweet Flag,	
Cinnamon,	
	one ounce and a half.
Cassia Lignea,	} of each, six drams.
Rhubarb,	
Long Pepper,	
	three ounces.

The total weight is seven ounces.

6th class.—

Ginger,	} of each, six drams.
Black Pepper,	
White Pepper,	
Leaves of Laurus Cassia,	} of each,
Cubebs,	
Cardamoms,	
	half an ounce.

The total weight is three ounces six drams.

Each of these mixtures is to be separately reduced to powder. Then each of the following substances is to be separately powdered.

Myrrh,	} of each, six drams.
Male Olibanum,	
Oriental Saffron,	
Trochiscs of Agaric,	one ounce and a half.
Castor,	} of each, two drams.
Asphaltum,	
Lemnian Bole,	
Calcined Chalcitis,	half an ounce.

Then cut small, mix with mucilage of tragacanth, and dry

Lemon-grass,	} of each, six drams.
Spikenard,	
French Lavender,	

Pulverize separately ; then mix the ingredients with

Trochiscs of Sweet Flag, three ounces.

Dissolve in Wine of Crete,

Gum Arabic,	} of each, half an ounce.
Sagapenum,	
Opoponax,	
Galbanum,	of each, two drams,

and strain the thick liquor through a silk sieve. Then mix together, separately,

Chio Turpentine,	six drams.
Storax,	four drams.

Then mix together the six classes of powder, the powdered trochiscs, and the other substances powdered separately, with the exception of the Lemnian bole, the saffron, and the chalcitis. Then put into a large boiler,

Warm Purified Honey, six ounces,
and, by the aid of a large wooden pestle,
spread it exactly on the sides of the vessel.
Then pour into the boiler, the turpentine
and storax melted with a little honey and
wine of Crete, then the solution of the
gums, also mixed with a little wine, and
warmed. Then add

Juice of Acacia, half an ounce.
——— Liquorice, one ounce and a half.
——— Hypocist, half an ounce.
Opium, purified with vinegar,
three ounces,

each dissolved separately in wine of Crete.

Triturate well together, add several
pounds of warm despumated honey, and in-
corporate the powder gradually, passing it
through a close sieve, adding honey from
time to time, and constantly stirring, that
a uniform mass may be obtained; then
add the Lemnian bole and the chalcitis,
mixed with wine and a little oil of nutmeg,
and then the saffron also mixed with wine.
After having well stirred the mixture, put
it into the boiler, which should be of such
capacity that the mass may not fill it to more
than two-thirds or three-fourths, in order
to allow space for its fermenting. First
stir it every day for one week, then once
a month for a whole year; after which it
may be taken out of the vessel for use.
(PALAT.)

The same formula is found in HISP., but
that pharmacopœia does not class the in-
gredients nor does it give the mode of pre-
paration.

R. Root of Fraxinella, } of each,
——— Squill, } two pounds.
——— Florentine Iris, one pound.
——— Cinquefoil, six ounces.
——— Sweet Flag, } of each,
——— Valerian, } five ounces.
——— Rhubarb, } of each, six ounces.
——— Ginger, }
——— Angelica, } of each,
——— Gentian, } four ounces.
——— Long Birthwort, two ounces.
——— Asarum, sixteen scruples.
Herb of Cretan Scordium, one pound.
——— Dittany of Crete, six ounces.
——— Germander, four ounces.
——— White Hore- } of each,
——— hound, } six ounces.
——— Calamint, }
——— Cretan Penny- } of each,
——— royal, } four ounces.
——— Ground Pine, }
——— Lesser Centaury, two ounces.
——— Marjoram, } of each,
——— Marum, } sixteen scruples.
Leaves of Laurus Cassia, five ounces.
Flowers of Red Roses, one pound.
——— French Lavender, six ounces.
——— St. John's-wort, four ounces.

Seeds of Rape, one pound.
——— Cardamoms, eight ounces.
——— Macedonian Parsley, six ounces.
——— Anise, }
——— Cretan Fennel, } of each,
——— Sandwort, } four
——— Seseli, } ounces.
——— Thlaspi, }
——— Cretan Carrot, two ounces.
Lemon-grass, seven ounces.
Spikenard, eight ounces.
Celtic Nard, four ounces.
Long Pepper, two pounds.
White Pepper, } of each, six ounces.
Black Pepper, }
Lesser Cardamoms, } of each,
Cubebes, } four ounces.
Cinnamon, eight ounces.
Arabian Costus, seven ounces.
Aloe Wood, one ounce.
Trochiscs of Agaric, one pound.
Myrrh, eight ounces.
Olibanum, six ounces.
Mastic, eight scruples.
Gum Arabic, four ounces.
Asphaltum, two ounces.
Lemnian Bole, four ounces.
Castor, } of each,
Italian Vipers, } two ounces.

Pulverize all these substances together,
excepting the lemon-grass and spikenard,
which are to be cut small and beat up with
mucilage of gum arabic, the mass dried
and reduced to powder, which is then to be
mixed with the preceding.

Set apart also the powdered roses, and
mix them with wine, adding,

Chalcitis, or Vitriol calcined to
whiteness, four ounces.
Safflower, powdered, eight ounces.

Dissolve also in a sufficient quantity of
Madeira wine,

Opium, two pounds.
Extract of Liquorice, one pound.
Juice of Hypocist, } of each,
——— Acacia, } four ounces.
Sagapenum, }

Having strained the solutions, mix them
with

Despumated Honey,
one hundred and sixty pounds.

Add the vinous liquor prepared with the
roses, safflower, and chalcitis, and incorpo-
rate the powder by means of a sieve. Fi-
nally, add,

Storax, four ounces.

Dissolved in

Turpentine, six ounces.
Opobalsamum, or, Oil of Nutmeg,
thirteen ounces.

Stir for one hour; then leave the mass to ferment in a large vessel, stirring it two or three times every month. (BRUNS.; WIRTEM.)

Besides these two classifications of the ingredients of the theriac, one founded on the equality of the weights, and the other on original resemblance; we also find in GALL. another which rests on analogy of nature and of properties in the ingredients: it embraces thirteen divisions, viz. :—

1st class.—*Acrid Substances.*

Pulp of Squill,	
three ounces four drams sixty grains.	
Root of Asarum,	eighty-eight grains.
White Agaric,	of each,
Seeds of Rape,	one ounce and a half.
Thlaspi,	half an ounce.

2nd class.—*Bitter Substances.*

Myrrh,	one ounce.
Tops of Lesser Centaury,	two drams.
Root of Gentian,	half an ounce.
Rhubarb,	six drams.
Herb of Scordium,	one ounce and a half.
Germander,	of each,
Ground Pine,	half
Tops of St. John's-wort,	an ounce.

3rd class.—*Astringent Substances.*

Petals of Red Roses,	one ounce and a half.
Root of Cinquefoil,	six drams.
Juice of Hypocist,	of each,
Acacia,	half an ounce.
Calcined Chalcitis,	

4th class.—*Exotic Aromatics.*

Cinnamon,	two ounces and a half.
Cassia Lignea,	one ounce.
Ginger,	six drams.
Long Pepper,	three ounces.
Black Pepper,	six drams.
Amomum,	one ounce.
Cardamoms,	half an ounce.
Leaves of Laurus Cassia,	six drams.
Lemon-grass,	fourteen drams.
Spikenard,	one ounce.
Celtic Nard,	half an ounce.
Arabian Costus,	seven drams.
Acorus Virus,	five drams.
Aloe Wood,	eighty-eight grains.

5th class.—*Indigenous Aromatics.*

Saffron,	one ounce.
Dried Lemon Peel,	
Calamint,	of each,
Dittany of Crete,	six drams.
French Lavender,	
White Horehound,	
Teucrium Montanum,	half an ounce.
Maram,	of each,
Marjoram,	eighty-eight grains.
Florentine Iris,	one ounce and a half.

6th class.—*Aromatics from Umbelliferous plants.*

Seeds of Macedonian Parsley,		} of each, half an ounce.
Sandwort,		
Fennel,		
Anise,		
Seseli,		
Root of Spignel (<i>Ligusticum Meum</i>)		
Seeds of Cretan Carrot,	two drams.	

7th class.—*Resins and Balsams.*

Xylobalsamum,	one dram.
Carpobalsamum,	half an ounce.
Opobalsamum,	fifteen drams.
Olibanum,	
Chio Turpentine,	of each, six drams.
Mastic,	twenty-four grains.
Asphaltum,	two drams.
Storax,	half an ounce.

8th class.—*Fetid Substances.*

Root of Valerian,	five drams.
Long Birthwort,	of each,
Galbanum,	two drams.
Opoponax,	
Sagapenum,	half an ounce.
Castor,	two drams.

9th class.—*Virous Substance.*

Opium,	three ounces.
--------	---------------

10th class.—*Inert Earth.*

Lemnian Bole,	four drams.
---------------	-------------

11th class.—*Gums, Feculae, &c.*

Gum Senegal,	four drams.
Crumb of Wheat Bread,	five drams fifty grains.
Tare Meal,	three drams, fifteen grains.
Viper's Flesh,	two drams twenty grains.

12th class.—*Sweet Substances.*

Extract of Liquorice,	one ounce and a half.
Honey,	ten pounds and a half.

13th class.—

Spanish Wine,	two pounds and a half.
---------------	------------------------

Divide the wine into three portions; mix the honey with one, the opium with another, and the gums and juices with the last: strain the liquors separately, then, having mixed them together, add the chalcitis, the balsams, and lastly, very gradually, the powders. Leave the mass to ferment for one year in a covered vessel.

The opium forms about one eighty-eighth part of the whole mass, which is nearly equivalent to rather less than, one grain in a dram.

- R. Long Birthwort, }
 Asphaltum, }
 Castor, } of each,
 Galbanum, } one part.
 Opoponax,
 Lesser Centaury,
 Seeds of Wild Carrot, }
 Amomum,
 Carpobalsamum,
 Lemon Peel,
 Gum Arabic,
 Leaves of Germander,
 ——— Ground Pine,
 ——— Laurus Cassia,
 ——— Mountain Penny-royal,
 Flowers of St. John's-wort,
 Celtic Nard,
 Sweet Flag,
 Root of Gentian,
 ——— Spiguel, (*Ligusticum*
Meum) } of each,
 ——— Valerian, } two
 Sagapenum, } parts.
 Seeds of Sandwort,
 ——— Anise,
 ——— Lesser Cardamoms,
 ——— Fennel,
 ——— Seseli,
 ——— Thlaspi,
 Storax,
 Juice of Acacia,
 ——— Hypocist,
 Sulphate of Iron, calcined to
 whiteness, }
 Lemnian Bole,
 Calamint,
 Cassia Lignea,
 Arabian Costus,
 Dittany of Crete,
 Leaves of Horehound,
 Ginger,
 Myrrh,
 Spikenard,
 Olibanum, } of each,
 Black Pepper, } three
 Root of Cinquefoil, } parts.
 ——— Pontic Rhubarb,
 Saffron,
 Seeds of Macedonian Parsley,
 Lemon-grass,
 French Lavender,
 Turpentine,
 Agaric,
 Balm of Mecca,
 Cinnamon,
 Leaves of Scordium,
 Petals of Red Roses,
 Florentine Iris,
 Squill, } of each,
 Seeds of Wild Rocket, } six parts.
 Extract of Liquorice,
 Seeds of Tare, } nine parts.
 Species of Trochiscs of }
 Sweet Flag, } of each,
 Opium, } twelve parts.
 Long Pepper, }

White Honey,
 six hundred and seventy-two parts.
 Malaga Wine, a sufficient quantity.

Dissolve the opium, the juices of acacia and of hypocist, and the extract of liquorice in a sufficient quantity of wine; emulsionize the galbanum, opoponax, and sagapenum, at first alone, then with the gum reduced to mucilage with the wine, then with the opobalsamum and turpentine: reduce to powder and sift the other substances. Lastly, incorporate the whole, adding a sufficient quantity of wine to give it the consistence of an electuary. (VANMONS)

- R. Pulp of Squill, one pound and a half.
 Chio Turpentine,
 Oil of Nutmeg, } of each,
 Extract of Liquorice, } nine ounces.
 Opium,
 Dried Viper,
 Cinnamon, } of each, eight ounces.
 Myrrh, }
 Pomegranate Flowers,
 Benzoin, } of each,
 Saffron, } six ounces.
 Gum Arabic,
 Olibanum,
 Angelica,
 White Dittany,
 Arabian Costus,
 Contrayerva,
 Serpentaria,
 Florentine Iris, } of each,
 Gentian, } five ounces.
 Valerian,
 Calamint,
 Ground Pine,
 Pennyroyal,
 Scordium,
 Bay Berries,
 Lesser Cardamoms,
 Seeds of Lemon, } of each,
 ——— Macedonian } four ounces.
 Parsley,
 ——— Seseli, }
 Sulphate of Iron, calcined to white-
 ness, ten drams.
 Spikenard, } of each, two ounces.
 Castor, }
 Honey, thrice the weight of the whole.
 Madeira Wine, a sufficient quantity.

Pulverize all the ingredients which are susceptible of being powdered; dissolve the opium in the wine, and strain the solution; then melt the honey over a slow fire, add gradually the solution of opium, the oil of nutmeg, and the turpentine melted together, and finally incorporate the powder. (SARD.)

- R. Dried Root of Squill, fourteen parts.
 Root of Florentine Iris, six parts.
 ——— Ginger, twenty-two parts.
 ——— Gentian, eleven parts.
 ——— Sweet Flag, twenty-four parts.
 ——— Valerian, } of each,
 ——— Pontic Rhubarb, } three parts.

Cinnamon,	fourteen parts.
Dittany of Crete,	eighteen parts.
Scordium,	eight parts.
Petals of Red Roses,	twelve parts.
Saffron,	four parts.
Seeds of Anise,	fourteen parts.
Gum Senegal,	twenty parts.
Olibanum,	three parts.
Turpentine,	six parts.
Galbanum,	four parts.
Balsam of Peru,	} of each, six parts.
Purified Opium,	
Extract of Liquorice,	
White Agaric,	fourteen parts.
Sulphate of Iron,	three parts.
Castor,	one part.
Honey,	six hundred and fifty-six parts.
Spanish Wine,	one hundred and fifty-eight parts.

Pulverize the solid substances, and mix the powders; divide the wine into three portions; dissolve the honey with a moderate heat in the first, mix the balsam and the turpentine in the second, and the opium in the third; mix together the two latter liquors, incorporate the mixture with the strained honey and wine, and finally incorporate the powders. (FORM. HÔP. FR.)

R. Squill,	four ounces.
Florentine Iris,	half an ounce.
Ginger,	} of each, six drams.
Dittany of Crete,	
Spikenard,	
French Lavender,	
Saffron,	
Myrrh,	
Gentian,	
Sweet Flag,	
Spiguel (<i>Ligusticum Meum</i>),	
Valerian,	
Celtic Nard,	} three ounces.
Amomum,	
Long Pepper,	
Scordium,	four ounces.
Cinnamon,	one ounce.
Lemon-grass,	one dram and a half.
Fennel,	one ounce and a half.
Opium,	eleven drams.
Castor,	two drams.
Storax,	half an ounce.
Balm of Mecca,	two ounces.
Sagapenum,	half an ounce.
Galbanum,	two drams.
Spanish Wine,	a sufficient quantity.
Honey,	four pounds.

Dissolve the opium in wine, and the honey in one pound of wine; pulverize the other ingredients, with the exception of the balsam, sagapenum, and galbanum; then, having melted these in a water-bath, add to them the mellated and opiate wines; and finally incorporate the powders. (ANT.)

R. Root of Ginger,	} of each, six drams.
— Florentine Iris,	
— Spikenard,	
— Celtic Nard,	
— Sweet Flag,	
— Gentian,	
— Valerian,	one ounce.
— Squill,	three ounces.
Leaves of Dittany of Crete,	six drams.
— Calamint,	} of each, three ounces.
— Horehound,	
— Scordium,	
Seeds of Anise,	} of each, one ounce.
— Fennel,	
— Lesser Cardamoms,	six drams.
Long Pepper,	} of each, two ounces.
Extract of Liquorice,	
Cinnamon,	
Asphaltum,	
Opium,	
Storax,	} of each, three drams.
Sagapenum,	
Galbanum,	
Castor,	one dram and a half.

Pulverize all these substances. Then melt slowly in a tinned vessel

Chio Turpentine, six ounces.

Add gradually warm despumated honey in the proportion of one ounce to every two ounces and a half of the powder; mix this gradually, and then add a mixture, prepared two days previously, of

Gum Arabic,	} of each, one ounce and a half.
Sulphate of Iron, dissolved in water,	
Red Roses,	
Nut Galls,	
Spanish Wine,	
	six ounces.

Stir the electuary for about six days. (BRUGNATELLI)

R. Root of Valerian,	} of each, one pound.
— Contrayerva,	
— Gentian,	
Leaves of Scordium,	
Chamomile Flowers,	
Cinnamon,	
Pepper,	
Juniper,	
Orange Peel,	
Seeds of Anise,	
Myrrh,	} four pounds.
Cinchona,	
Saffron,	
Sulphate of Iron, calcined to whiteness,	} of each, one pound.
Opium,	
Honey of Elder,	six pounds.
Honey,	forty-eight pounds.

Despumate the honey with a sufficient quantity of wine, add the honey of elder, then the saffron and the calcined sulphate,

then the opium dissolved in wine, and finally incorporate the powder. (HISP.)

Each dram contains two grains of opium.

R. Root of Angelica,	six ounces.
——— Serpentaria,	four ounces.
——— Valerian,	} of each, two ounces.
——— Squill,	
——— Zedoary,	
——— Cinnamon,	
Lesser Cardamoms,	} of each, one ounce.
Myrrh,	
Saffron,	
Cloves,	
Sulphate of Iron,	
Opium,	} six pounds.
Despumated Honey,	

To the honey slightly warmed add the opium dissolved in a sufficient quantity of Malaga wine; then incorporate the other pulverized ingredients. (HANN.; OLDEN.; POL.; BORUS.; FEE; NIEMANN)

R. Serpentaria,	} of each, six ounces.
Contrayerva,	
Angelica,	} of each, four ounces.
Valerian,	
Sweet Flag,	
Gentian,	
Zedoary,	} of each, two ounces.
Cardamoms,	
Myrrh,	
Mace,	} of each, two ounces.
Opium, dissolved in Spanish Wine,	
	nineteen drams.
Rob of Elder,	eighty ounces.
——— Juniper,	forty ounces.
Simple Syrup,	two pounds.

Mix. (ROSS.)

R. Root of Angelica,	six ounces.
——— Serpentaria,	four ounces.
——— Squill,	} of each, two ounces.
——— Valerian,	
Cinnamon,	} of each, one ounce.
Lesser Cardamoms,	
Myrrh,	
Saffron,	
Cloves,	
Sulphate of Iron,	
Opium,	
Despumated Honey,	six pounds.

Heat the honey a little, add the opium dissolved in Spanish wine, and lastly the other substances, in powder. (DAN.)

Each ounce contains about five grains of opium.

R. Root of Angelica,	eight ounces.
——— Gentian,	six ounces.
——— Valerian,	three ounces.
——— Zedoary,	} of each, two ounces.
Lesser Cardamoms,	

Saffron,	} of each, one ounce.
Extract of Liquorice,	
Myrrh,	
Opium,	
Honey,	seventy-five ounces.

Dissolve the extract, myrrh, and opium in Spanish wine, heat this liquor with the honey, and add the other ingredients in powder, constantly stirring. (FERR.; SUEC.; SWEDIAUR)

R. Root of Angelica,	six parts.
——— Serpentaria,	four parts.
——— Squill,	} of each, one part.
——— Zedoary,	
Cinnamon,	
Lesser Cardamoms,	
Myrrh,	} of each, one part.
Saffron,	
Cloves,	

Incorporate these with

Honey, twelve parts.

At the time of prescription, add to each ounce Tincture of Opium, half a dram.

Mix. (SAX.)

R. Hard Opium, in powder,	one part.
Wine,	four parts.

Macerate for two days in a covered vessel; then add

Despumated Honey, seventy-two parts.

Mix, and add, constantly stirring, a powder composed of

Myrrh,	} of each, one part.
Saffron,	
Proto-sulphate of Iron,	} of each, two parts.
Cloves,	
Lesser Cardamoms,	
Root of Valerian,	
——— Ginger,	} of each, six parts.
——— Angelica,	
——— Gentian,	

Mix. (BAVAR.)

R. Aromatic Powder, composed of one part of Pichurim Bean, one of Lesser Cardamoms, and one half part of Cinnamon, six ounces. Opium, dissolved in wine, three drams. Powdered Serpentaria, three ounces. Despumated Honey, a sufficient quantity.

Mix. (LUSIT.)

R. Nutmegs,	} of each, two ounces.
Lesser Cardamoms,	
Ginger,	
Serpentaria,	three ounces.
Opium, dissolved in Spanish Wine,	three drams.

Despumated Honey, thrice the weight of the powders.

Mix. (WIRTEM.)

Confectio Opii. (LOND.; DUBL.; NIEMANN)

- R. Opium, in powder, six drams.
 Long Pepper, one dram.
 Ginger, two ounces.
 Caraway Seeds, three ounces.
 Simple Syrup, one pint.

Triturate the opium with the syrup made hot, then add the other ingredients reduced to powder, and mix.

Electuarium opiatum, olim Elect. thebaicum. (EDINB.; COXE)

- R. Aromatic Powder, six ounces.
 Root of Serpentaria, in powder, three ounces.
 Opium, mixed with a sufficient quantity of Spanish White Wine, half an ounce.
 Syrup of Ginger, one pound.

Mix, and make an electuary.

BATAV.; and BELG. give the same formula, but require despumated honey instead of syrup.

- R. Aromatic Powder, composed of two parts of Cinnamon, and one part each, of Lesser Cardamoms, of Ginger, and of Long Pepper, six ounces.
 Serpentaria, three ounces.
 Opium, dissolved in Spanish Wine, three drams.
 Despumated Honey, thrice the weight of the powders.

Mix. (GENEV.)

- R. Cinnamon, } of each,
 Nutmeg, } one ounce and
 Lesser Cardamoms, } a half.
 Ginger, }
 Serpentaria, three ounces.
 Opium, dissolved in Spanish Wine, three drams.
 Despumated Honey, twenty-eight ounces.

Incorporate. (AMST.)

For a long time Venice engrossed the supplying of all Europe with theriac; it was prepared there every year with great ceremony: but at the present day the apothecaries of every country are required to make it according to the process indicated in its proper authorized pharmacopœias, excepting however those of Spain, in which country every branch of medical science is in nearly the same state as it was in the middle ages:

VOL. II.

it is there allowed to be prepared only at Madrid, before the assembled royal college of medicine.

This electuary occupies a distinguished place among the other disgusting substances and absurd preparations which crowd the Spanish pharmacopœia, in which Galenism and Arabicism seem to have sought their last refuge. After that of Spain, the French pharmacopœia is the only one where a reforming hand is withheld from clearing away the rubbish of oriental medicine; for those of the Palatinate and of Wirtemburgh are no longer of authority, though they have not yet been replaced. In every other country theriac has more or less undergone happy modifications; it has even disappeared totally from some dispensatories, which shows that it has ceased to be employed in the practice of the physicians of the countries to which they belong.

Nothing can be more contradictory than the opinions which have been entertained with regard to theriac. According to some, it should be respected on account of its antiquity, and should interest us from its having been held in esteem for so great a length of time; according to others, it is a monstrous farrago which is a disgrace to medicine. To the former we say, that long empire does not authorize usurpation; that folly with grey hairs, is no more to be preferred to reason than when possessing all the grace and freshness of youth; and lastly, that they themselves, to be consistent, should have kept to ANDROMACH'S original formula, which was itself sufficiently voluminous and did not require to be still more enlarged. To the latter we say, that there is, undoubtedly, truth in their opinion, but there is also exaggeration, for the theriac scarcely exceeds in complication many other official preparations which have not yet lost all credit; and, with regard to absurdity, a greater or less degree of it does not alter its ridiculousness.

In the year 1826 twelve of the twenty-one hospitals of Paris consumed sixty-two pounds of theriac, though the other nine establishments used none; and it has been thought proper to be introduced into the formulary of the military hospitals of France. From this it may reasonably be imagined that it will not immediately be erased from the French pharmacopœia; but it might be, it ought to be simplified, after the example of the surrounding countries. In doing this, two methods might be adopted; to combine opium with stimulants alone, or with stimulants and astringents at the same time. The latter method, although proposed with reference to the composition of the original formula, reduces the theriac to diascordium and its

modern derivatives: but if the other deserves to be preferred, what would be the use of being burthened with a preparation of which we already have the equivalent in SYDENHAM'S laudanum, the employment of which is daily becoming more and more limited in proportion as the modern discoveries in the chemical composition of opium come to be adopted in medical practice.

It has been said that the ancient theriac was of greater efficacy than the modern; but the French pharmacopœia, in saying that it has ascertained by analysis that there is no difference between them, we may properly appreciate the effect of the fermentation which has been so much extolled as a means of perfecting the preparation: this effect can scarcely be more than the production of alcohol, or at most of acetic acid.

Dose, from a scruple to two or three drams, alone or in wine.

THERIACAL EXTRACT.

Extractum theriacale. (SAUNDERS)

R. Theriac, any quantity.
Alcohol, a sufficient quantity

to cover it by three fingers' breadth. Digest for three days, then decant the alcohol and express the rest; mix the two liquors, and evaporate them in a water-bath to the consistence of an extract.

According to GUILBERT, the alcohol takes up from the theriac, volatile oil, resin, turpentine, balsam, and a green oil. This analysis is evidently very incomplete.

SIMPLE THERIACAL WATER.

Aqua theriacalis simplex. (FERR.; BRUGNATELLI)

R. Theriac, one ounce.
Pure Water, four ounces.

Distil three pounds at a moderate heat. (BRUGNATELLI)

FERR. prescribes one ounce of theriac and four pounds of juice of sour grapes.

This water is reputed alexipharmic, stomachic, carminative, and vermifuge.—It is principally administered in the vermical disorders of children, by spoonfuls, alone or with other compounds.

ROMAN THERIACAL WATER.

Aqua theriacalis Romana seu Scordii composita seu theriacalis frigida. (PALAT.; SARD.)

R. Theriac, half a pound.

Juice of Galega,	}	of each, six pounds.
—— Sorrel,		
—— Scordium,		
—— Lemon,		

Digest for twenty-four hours, and distil. (PALAT.)

SARD. prescribes four pounds of scordium water, two of sorrel juice, two of galega juice, one and a half of lemon juice, and four ounces and a half of theriac.

It is for the same uses as the preceding.

COMPOUND THERIACAL WATER. (WIRTEM.)

R. Theriac,	half a pound.	
Root of Angelica,	}	of each, one ounce.
—— Valerian,		
—— Carline Thistle,		
—— Masterwort,		
—— Zedoary,	}	of each, one ounce and a half.
Herb of Scordium,		
—— Rue,		
Fresh Lemon Peel,	two ounces.	
Juniper,	one ounce.	
Water,	ten pounds.	

Distil six pounds.

This does not differ from the compound spirit of angelica except by the absence of alcohol.

THERIACAL SPIRIT. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

R. Theriac, four ounces.
White Wine, sixty ounces.

Digest, and distil thirty ounces. (BRUNS.; WIRTEM.; SPIELMANN)

R. Concentrated Alcohol, ten ounces.
Theriac, five ounces.
Myrrh, two ounces and a half.
Saffron, half an ounce.

Digest for eight days; then distil in a water-bath, and add, if required, two drams of camphor. (PALAT.)

There is very little difference between this and the compound spirit of angelica.

THERIACAL ESSENCE OR TINCTURE.

Essentia seu Tinctura theriacalis. (BRUNS.; SARD.; WIRTEM.; VAN MONS)

R. Theriac, one part.
Proof Spirit, three parts.

Infuse for several days, and filter. (VAN MONS)

SARD. prescribes one part of theriac and six parts of proof spirit.

R. Herb of Scordium,	}	of each, three drams.
—— Dittany of Crete,		
Root of Angelica,		
—— Sweet Flag,		
—— Serpentaria,		
Spikenard,		

Cinnamon,	} of each, one	
Lesser Cardamoms,		dram and a half.
Cloves,		
Opium,	} of each, one dram.	
Saffron,		
Proof Spirit,		one pound.

Infuse for several days, without heat, and filter. (BRUNS.; WIRTEM.; VAN MONS)

It is stimulant and stomachic.—Dose, from forty to sixty drops.—One ounce of this tincture contains about five grains of opium.

STOMACHIC PLASTER.

Emplastrum stomachicum. (SWEDIAUR)

R. Theriac, one or two ounces.
Oil of Nutmeg, two drams.

Mix, and spread it upon leather having a margin of adhesive plaster.

It is recommended in cardialgia and spasmodic affections of the abdomen.

Scutum stomachicum. (PIERQUIN; SWEDIAUR)

R. Theriac, two ounces.
Nutmeg, in powder, two drams.

Mix.—It is to be applied as the preceding. (SWEDIAUR)

R. Theriac, } of each,
Powdered Maize, } a sufficient quantity.
Opium, three grains.
Camphor, two grains.

Mix. (PIERQUIN)—FOUQUET recommended it in colic, chronic vomiting, and spasmodic affections of the stomach.

ANTI-HÆMORRHOIDAL PLASTER. (PIERQUIN; SAINTE-MARIE)

R. Opium, ten grains.
Theriac, two drams.

Mix. (PIERQUIN)

SAINTE-MARIE prescribes six drams of theriac and six grains of opium.—This plaster is to be applied to the umbilicus.—VALSALVA is said to have derived the happiest effects from it in arresting excessive hæmorrhoidal discharge.

ANTI-SPASMODIC AND TONIC APPLICATION. (PIERQUIN)

R. Theriac, two drams.
Opium, } of each, six grains.
Camphor, }

Mix.—To be applied to the epigastrium.

THERIACAL CATAPLASM.

Cataplasma Fermenti Panis theriacale seu ad Singultum. (HISP.; SAINTE-MARIE)

R. Yeast, three ounces.
Theriac, half an ounce.
Powdered Juniper, } of each,
Rose Petals, } two drams.
Mint, }
Powdered Cloves, } of each,
Nutmeg, } one dram.
Mace, }
Rose Vinegar, a sufficient quantity
to make a cataplasm. (HISP.)

R. Theriac, two ounces.
Anise, in powder, two drams.
Oil of Cloves, eight drops.
Spirit of Camphor, a sufficient quantity
to mix the whole. (SAINTE-MARIE)

The former of these cataplasms has been recommended, applied to the region of the stomach, against hiccup. The second is recommended by ZIMMERMANN against the clavus hystericus.

DYSENTERIC ELECTUARY. (SPIELMANN)

R. Theriac, } of each, two ounces.
Diascordium, }
Gum Arabic, half an ounce.
Armenian Bole, three ounces.

Mix.

By adding two ounces of LOCATELLI's balsam, the *Balsamic Dysenteric Electuary* is obtained.

The first has been recommended in ordinary dysentery, and the latter in ulcerous dysentery.

ASTRINGENT ELECTUARY. (BORIES; CADET DE GASSICOURT; PIERQUIN)

R. Theriac, } of each, one ounce.
Diascordium, }
Distilled Water of Angelica, } of each,
Valerian, } one
Mint, } dram.
Gum Tragacanth, half a dram.
Syrup of Sulphate of Quinine,
a sufficient quantity.

Mix.—It is astringent, and is extolled by BALLY in dysentery and chronic diarrhoea.—Dose, one dram, twice or thrice a day. (CADET DE GASSICOURT; PIERQUIN)

R. LOCATELLI's Balsam, } of each,
Conserve of Roses, } one ounce.
Theriac, }
Cold Seeds, }

Mix.—Dose, a tea-spoonful every four hours, in a glassful of infusion of milfoil. (BORIES)

THERIACAL MIXTURE. (RATIER)

- R. Theriac, one dram and a half.
 Gum Arabic, one dram.
 Water, six ounces.
 Simple Syrup, one ounce.

Mix.—To be taken by spoonfuls.

CORDIAL POTION. (BORIES; RATIER)

- R. Theriac, one dram.
 Infusion of Balm, seven ounces.
 Tincture of Cinnamon, two drams.
 Syrup of Clove Pink, one ounce.

Mix. (RATIER)

- R. Water of Mugwort, } of each,
 ——— Motherwort, } two
 ——— Orange Flowers, } ounces.
 Hordeated Cinnamon Water, half an ounce.
 Theriac, } of each,
 Confection of Hyacinth, } one scruple.
 Alkermes Confection, }
 Trochiscs of Myrrh, twenty-five grains.
 Mace, fifteen grains.
 Compound Tincture of Aloes, thirty drops.

Mix. (BORIES.)—To be taken by spoonfuls.

TONIC POTION. (PIERQUIN)

- R. Theriac, one dram.
 Orange-flower Water, } of each,
 Water or Wine, } one ounce.

Mix.—For one dose.

ANTI-SPASMODIC POTION. (PIERQUIN)

- R. Theriac, one dram.
 Syrup of Poppies, six drams.
 HOFFMANN'S LIQUOR, } of each,
 Tincture of Amber, } twenty drops.
 Orange-flower Water, } of each,
 Water of Motherwort, } two ounces.
 ——— Rue, }

Mix.—To be taken by spoonfuls.

TRYPHERA MAGNA. (SPIELMANN)

- R. Opium, two ounces.
 Cinnamon, }
 Cloves, }
 Galanga, }
 Spikenard, }
 Ginger, }
 Arabian Costus, }
 Storax, }
 Sweet Flag, }
 Round Cyperus, }
 Florentine Iris, }
 German Iris, }
 Sulphurwort, }
 Mandrake, }
 Lavender, }
 Roses, }

of each,
one ounce.

- Pepper,
 Anise,
 Smallage,
 Fennel,
 Macedonian Parsley,
 Carrot,
 Sweet Basil,

of each,
one ounce.

Pulverize all these substances, and incorporate them with

Honey, a sufficient quantity.

This anodyne electuary was prescribed by GALEN; it differs very little from theriac.

ORVIETANUM. (WIRTEM.)

- R. Root of Long Birthwort, }
 ——— Round Birthwort, }
 ——— Angelica, }
 ——— Bistort, }
 ——— Carline Thistle, }
 ——— Contrayerva, }
 ——— Fraxinella, }
 ——— Gentian, }
 ——— Masterwort, }
 ——— Cinquefoil, }
 ——— Serpentaria, }
 ——— Tormentil, }
 ——— Valerian, }
 ——— Zedoary, }
 Leaves of Blessed Thistle, }
 ——— Pennyroyal, }
 ——— Rue, }
 ——— Savine, }
 ——— Scordium, }
 ——— Scabious, }
 Flowers of St. John's-wort, }
 Orange Peel, }
 Lemon Peel, }
 Cinnamon, }
 Dried Viper, } two ounces.

of each,
one
ounce.

Incorporate these, powdered, with

- Rob of Juniper, } of each,
 Despumated Honey, } three pounds.

Having well mixed, add

- Theriac, }
 Mithridate, } of each, two ounces.
 Oil of Rue, }
 ——— Amber, }
 ——— Cloves, }
 ——— Juniper, } of each,
one scruple.

Mix intimately.

This electuary is said to have been invented by JEROME FERRANTE of Orvieto, whose name it bears. Although it is better adapted than the two preceding to please the tastes and fancies of empirics, yet it is now totally forgotten; it differs from the preceding ones in its scarcely containing any opium. It may be said to be to the theriac, what the Japanese confection is to diascordium; the principal is

masked, and almost entirely nullified by the accessories.

Dose, from two drams to half an ounce.

SPECIES OF DIASCORDIUM.

Pulvis seu Species Diascordii. (SARD.)

- R. Dried Leaves of Scordium, one ounce and a half.
 Petals of Red Roses, one ounce.
 Armenian Bole, two ounces.
 Cassia Lignea, } of each,
 Leaves of Dittany of Crete, } half
 Galbanum, } an ounce.
 Storax, }
 Root of Bistort, } of each, five drams.
 ——— Gentian, }
 Gum Arabic, }
 Opium, two drams.
 Pulverize, and pass through a sieve.

DIASCORDIUM.

Diascordium, Electuarium diascordium seu e Scordio seu Scordii seu Scordii opiatum seu Fracastoris seu Mimosæ Catechu compositum seu compositum Catechu, Confectio Japonica seu Diascordii, Conditum catechucicum. (DUBL.; EDINB.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENÈV.; LIPP.; LUSIT.; PALAT.; SARD.; SUEC.; WIRTEM.; AUGUSTIN.; BRUGNATELLI.; CADET DE GASSICOURT.; COXE.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

1. *Without Catechu.*

- R. Root of Bistort, } of each,
 ——— Tormentil, } half
 ——— Gentian, } an ounce.
 Herb of Dittany of Crete, }
 ——— Scordium, one ounce.
 Seeds of Sorrel, one dram and a half.
 Cinnamon, } of each,
 Cassia Lignea, } half an ounce.
 Long Pepper, } of each, two drams.
 Ginger, }
 Opium, one dram and a half.
 Gum Arabic, } of each, half an ounce.
 Galbanum, }
 Storax, }
 Armenian Bole, one ounce and a half.
 Lemnian Bole, half an ounce.
 Conserve of Red Roses, one pound.
 Despumated Honey, thirty ounces.

Dissolve the opium, gum arabic, and galbanum in three ounces of white wine, mix the solution with the honey, add the hard substances in powder, and finally incorporate the conserve. (BRUNS.; PALAT.; SARD.; WIRTEM.; SPIELMANN.)

- R. Leaves of Scordium, one ounce and a half.

- Red Roses, }
 Root of Bistort, }
 ——— Gentian, } of each,
 ——— Tormentil, } half an ounce.
 Seeds of Barberries, }
 Cassia Lignea, }
 Cinnamon, }
 Dittany of Crete, }
 Storax, }
 Galbanum, }
 Gum Arabic, }
 Ginger, } of each,
 Long Pepper, } two drams.
 Vinous Extract of Opium, }
 Prepared Oriental Bole, two ounces.
 Spanish Wine, about half a pound.
 Honey of Roses, two pounds.

Dissolve the galbanum in a small quantity of the wine, and the honey in the rest of it; then add the powders gradually. (GALL.; CADET DE GASSICOURT.)

- R. Root of Bistort, } of each,
 ——— Gentian, } two parts.
 ——— Tormentil, }
 ——— Ginger, one part.
 Cinnamon, four parts.
 Herb of Dittany of Crete, two parts.
 Dried Leaves of Scordium, six parts.
 Petals of Red Roses, three parts.
 Long Pepper, one part.
 Seeds of Barberries, }
 Gum Arabic, } of each,
 Galbanum, } two parts.
 Balsam of Peru, }
 Opium, one part.
 Armenian Bole, eight parts.
 Spanish Wine, thirty parts.
 Honey, one hundred and twenty-five parts.

Make an electuary. (FORM. HÔP. FR.)

- R. Ginger, }
 Gentian, } of each, one dram.
 Tormentil, }
 Dittany of Crete, }
 Mastic, }
 Armenian Bole, }
 Opium, two drams.
 Cinnamon, three drams.
 Leaves of Scordium, } of each,
 Petals of Red Roses, } six drams.
 Honey of Roses, twenty-four ounces.

Mix the powders with the honey. (HISP.)

- R. Ginger, }
 Opium, } of each, one part.
 Long Pepper, }
 Gentian, } of each, two parts.
 Storax, }
 Cinnamon, }
 Dittany of Crete, }
 Gum Arabic, } of each, four parts.
 Red Roses, }
 Scordium, }
 Tormentil, }

Armenian Bole, eight parts.
Honey, one hundred and twenty parts.

Triturate the opium with Spanish wine, and emulsionize the liquor with the storax and gum; then add a little honey, then the powders mixed together, and finally the rest of the honey: heat slightly and stir till cold. (VAN MONS)

R. Gentian, seven drams and a half.
Cinnamon, seven drams.
Opium, three drams.
White Wine, six ounces.

Digest for some days, strain with slight expression, and add to the liquor

Powdered Herb of Scordium, three ounces.

Despumated Honey, twenty ounces.

Mix. (LIPP.)

2. With Catechu.

Electuarium Catechu compositum.

R. Extract of Catechu, four ounces.
Kino, three ounces.
Cinnamon, } of each,
Nutmegs, } one ounce.
Opium, triturated with a sufficient quantity of Spanish White Wine, one dram and a half.

Syrup of Red Roses, boiled to the consistence of Honey, two pounds and a quarter.

Reduce the solid ingredients to powder; then mix them with the others, and form an electuary. (EDINB.)

R. Catechu, four ounces.
Cinnamon, two ounces.
Kino, three ounces.

Reduce these to powder, and add Hard Purified Opium, mixed with Spanish White Wine, one dram and a half.

Syrup of Ginger, boiled to the consistence of Honey, two pounds and a quarter.

Mix. (DUBL.)

ANT.; BATAV.; BELG.; LUSIT.; COXE; CADET DE GASSICOURT; HENRI; and SWEDIAUR give the same formula as EDINB.

R. Cinnamon, } of each, half an ounce.
Long Pepper, }
Catechu, four ounces.
Herb of Scordium, } of each,
Root of Tormentil, } two ounces.
Gentian, }

Pulverize, mix, and add

Opium, diffused in wine, five drams.
Honey, two pounds and a half.
Spanish Wine, a sufficient quantity

to make an electuary. (BRUGNATELLI)

R. Catechu, three ounces.
Root of Tormentil, } of each,
Nutmegs, } two ounces.
Olibanum, }
Opium, dissolved in Spanish Wine, one dram and a half.

Syrup of Orange Peel, a sufficient quantity.

Mix. (GENEV.)

R. Leaves of Scordium, } of each,
Catechu, } two ounces.
Root of Tormentil, }
Nutmegs, } of each,
Cinnamon, } one ounce.
Gum Arabic, }
Opium, dissolved in Spanish Wine, one dram and a half.
Honey of Roses, twenty-eight ounces.

Mix the opium with the honey, and incorporate the other substances reduced to fine powder. (AMST.)

R. Herb of Scordium, three ounces.
Catechu, } of each,
Pichurim Bean, } two ounces.
Root of Tormentil, }
Opium, one dram and a half.
Honey, twenty-eight ounces.

Triturate the opium with Spanish wine, add the honey rather warm, and incorporate the mixture with the other ingredients in powder. (SUEC.; FEE)

R. Catechu, three ounces.
Root of Tormentil, } of each,
Nutmegs, } two ounces.
Olibanum, }
Opium, half a dram.
Syrup of Roses, a sufficient quantity.

Make an electuary. (AUGUSTIN)

R. Leaves of Scordium, } of each,
Catechu, } three ounces.
Cinnamon, } of each,
Root of Tormentil, } two ounces.
Opium, two drams.
Honey, twenty-eight ounces.
Spanish Wine, a sufficient quantity.

Mix. (FERR.)

This preparation is reputed sudorific, alexipharmic, anodyne, and styptic. It is principally prescribed in diarrhoea and dysentery, in the dose of one, two, or three drams. Its activity depends more upon the astringent substances which it contains than upon the opium.

ASTRINGENT ELECTUARY. (SAUNDERS)

R. Diascordium, } of each, six drams.
Dragon's Blood, }

Alum,	} of each, two drams.
Catechu,	
Cinnamon,	
Simple Syrup,	
a sufficient quantity	

to make an electuary.

ASTRINGENT OPIATE WINE. (FORM.
HÔP. FR.)

R. Diascordium, one part.
Red Wine, twenty-five parts.
Digest.

ASTRINGENT LAVEMENT. (BRERA)

R. Infusion of Chamomile, four ounces.
Yolks of Eggs, two.
Diascordium, one dram.

Mix.—To be exhibited in passive alvine discharges, emesis, gastrodynia, and spasmodic colic.

ANTI-EMETIC PILLS. (PIERQUIN)

R. Powdered Mint, two drams.
Wormwood, two scruples.
Pomegranate Flowers, one scruple.
Oil of Cinnamon, two drops.
Diascordium, one dram.
Syrup of Quince, a sufficient quantity

to make a pilular mass.—Dose, ten grains, thrice a day.

ASTRINGENT MIXTURE.

Julapium sistens, Mixtura sistens seu Opii adstringens Sanguinis seu ad Vomitum Sanguinis. (HISP.; SAUNDERS)

R. Compound Electuary of Catechu (FORM. OF EDINB.), three drams.
Tincture of Opium, forty drops.
White Sugar, half an ounce.
Alexiterious Water, four ounces.
Spirituos Cinnamon Water, two ounces.

Mix. (SARD.)

The two following formulæ should be placed here, although they do not contain the preparation in question :

R. Syrup of Poppies, one ounce.
Water of Red Poppies, seven ounces.
Catechu, one dram and a half.
SYDENHAM'S Laudanum, twenty drops.

Mix. (SAUNDERS)

R. Distilled Water, two ounces and a half.
Vinegar, half an ounce.
Opium, two grains.

Mix perfectly, and add

Powdered Red Coral,	half a dram.
———— Catechu,	half a scruple.
Simple Syrup,	one ounce.

Mix. (HISP.)

THERIACA CELESTIS HOFFMANNI.
(BRUNS.; SPIELMANN)

R. Extract of the Root of Ange-	} of each, three drams.
lica,	
Birthwort,	
trayerva,	
campane,	
tian,	
mentil,	
rian,	
low-wort,	
um Vulgare,	
ary,	} of each, one ounce.
Leaves of Blessed Thistle,	
Lesser Centaury,	} of each, one dram.
Scordium,	
Extract of Opium,	} one ounce.
Resin of Cascarilla,	
Labdanum,	} three drams.
Storax,	
Myrrh,	} half an ounce.
Galbanum,	
Mastic,	} of each, three drams.
Opoponax,	
Guaiac Resin,	} of each, one dram.
Saffron,	
Castor,	} half a dram.
Powdered Viper,	
Volatile Salt of Hartshorn,	} of each, one dram
Amber,	
Camphor,	} of each, one scruple.
Ambergris,	
Balsam of Peru,	} three drams.
Essential Oil of Cloves,	
Lemon,	} of each, twelve drops.
Juniper,	
Amber,	} of each, two scruples.
Cardamoms,	
Cinnamon,	} of each, two scruples.
Cubebs,	
Mace,	} of each, two scruples.
Nutmegs,	

Pulverize the hard substances, and triturate together the resins and balsams in a mortar heated in a water-bath; then add the extracts, the volatile salts in powder, and finally the essential oil. (BRUNS.)

R. Myrrh, half an ounce.

Galbanum,	} of each, two scruples.	
Mastic,		
Opoponax,		
Storax,		three drams.
Extract of Blessed Thistle,	} of each, half an ounce.	
Lesser Centaury,		
Scordium,		
Tormentil,		
Contrayerva,	} six drams.	
Angelica,		
Opium,		
Saffron,		
Swallow-wort,	} of each, three drams.	
Valerian,		
Gentian,		
Round Birthwort,		
Resin of Aloe Wood,	} of each, one dram.	
Guaiac,		
Labdanum,		
Deutoxide of Antimony,	} of each, half an ounce.	
Cinnabar,		
Prepared Amber,		
Powdered Viper,		
Essential Salt of Amber,	} of each, one dram and a half.	
Harts-horn,		
Oil of Cinnamon,	} of each, twelve drops.	
Zedoary,		
Cardamoms,		
Cubebs,		
Mace,		
Cloves,		
Juniper,		
Lemon,		
Expressed Oil of Nutmeg,	} of each, three drams.	
Balsam of Peru,		
Camphor,	} of each, one scruple.	
Ambergris,		

Mix. (SPIELMANN)

This differs principally from the theriac by the absence of sugar and by the presence of salts susceptible of re-action upon those of the opium. Each dram contains about four grains of opium. It has been extolled as anodyne and antiseptic, in the dose of from two to twenty-four grains, in pills.

PHILONIUM ROMANUM.

Electuarium Hyoscyami opiatum. (HISP.; PALAT.; WIRTEM.; SPIELMANN)

R. White Pepper,	five drams.
Opium,	two drams and a half.
Cassia Lignea,	one dram and a half.
Seeds of White Henbane,	five drams.
Smallage,	one dram.
Macedonian Parsley,	} of each, forty-five grains.
Fennel,	
Cretan Carrot,	
Saffron,	one scruple.
Spikenard,	} of each, fifteen grains.
Pyrethrum,	
Zedoary,	

Cinnamon,	one dram and a half.
Euphorbium,	} of each, one dram.
Myrrh,	
Castor,	
Honey,	
	eight ounces four drams and a half.

Mix. (PALAT.; WIRTEM.)

R. Root of Pyrethrum,	} of each, one ounce.
Zedoary,	
Doronica,	
Pomegranate Flowers,	} of each, twenty ounces.
White Pepper,	
Seeds of White Henbane,	
Opium,	ten ounces.
Euphorbium,	} of each, one ounce.
Castor,	
Pearls,	
Saffron,	
Honey,	five ounces.
	thirty ounces.

Mix. (SPIELMANN)

R. Seeds of Henbane,	} of each, six drams.
Anise,	
Fennel,	
Myrrh,	} five drams.
Castor,	
Opium,	
Cinnamon,	
Saffron,	
Chamomile Flowers,	
Despumated Honey,	
	two pounds.

Mix. (HISP.)

Each dram contains more than one grain and a half of opium and four grains of henbane. It is sedative, anodyne, and somniferous; it is principally recommended in flatulent colic, violent cough, and convulsions. — Dose, from half a dram, to one dram. It is also prescribed in lavements.

REQUIES NICOLAI.

Electuarium Reques Nicolai. (PALAT.; WIRTEM.; SPIELMANN; VAN MONS)

R. Petals of Red Roses,	} of each, two drams.
Violets,	
Opium,	} of each, one dram and a half.
Seeds of Henbane,	
White Poppy,	
Endive,	
Lettuce,	
Purslane,	
Flseseed,	} one dram.
Bark of Mandrake Root,	
Nutmegs,	
Cinnamon,	
Ginger,	
Sugar Candy,	
White Sanders,	
Red Sanders,	
Ivory burnt to whiteness,	} five grains.
Gum Tragacanth,	
Sugar,	
	ten ounces.

Mix with rose water. (PALAT.)

R. Petals of Red Roses,	}	of each,	six drams.
— Violets,			
Seeds of White Henbane,	}	of each,	half an ounce.
— Poppy,			
— Lettuce,			
— Fleaseed,	}	of each,	three drams.
Bark of Mandrake Root,			
Nutmegs,			
Cinnamon,			
Ginger,			
Purified Opium,	}	of each,	two drams.
White Sanders,			
Red Sanders,			
Gum Tragacanth,			
Ivory, burnt to whiteness,			
White Sugar,			twenty ounces.

Dissolve the opium in three ounces of white wine, add the powders, and incorporate the sugar, dissolved in ten ounces of rose water. (WIRTEM.)

R.	Root of Mandrake,	}	of each, one ounce and a half.
	Petals of Roses,		
	———— Violets,		
	Seeds of Henbane,		
	———— Poppies,		
	———— Endive,		
	———— Lettuce,		
	———— Purslane,		
	———— Fleaseed,	}	of each, six drams.
	Opium,		
	Nutmegs,		
	Cinnamon,		
	Ginger,		
	White Sanders,		
	Red Sanders,	}	
	Ivory, burnt to whiteness,		
	Gum Tragacanth,		

Sugar Candy,	one ounce.
Rose Julep,	twenty-four ounces.
Mix. (SPIELMANN)	
R. Gum Arabic,	} of each, two drams.
Opium,	
Ginger,	} of each, three drams.
Nutmegs,	
Cinnamon,	six drams.
Red Roses,	fourteen drams.
Seeds of White Poppy,	eighteen drams.
Honey,	fourteen ounces.

Mix the powdered gum and seeds with the opium; then triturate the whole with the honey, adding the other powders mixed together. (VAN MONS)

R. Honey,	forty-eight parts.
Seeds of Anise,	of each,
Subcarbonate of Lime,	eight parts.
Opium,	one part.
Cinnamon,	of each,
Nutmegs,	twenty-five parts.

Triturate the opium with the subcarbonate till it is reduced to powder, then add the other ingredients in powder, and finally incorporate the honey. (VAN MONS)

It is rather remarkable that no other virtue has ever been attributed to this electuary than that of being sedative and soporific. It is nearly disused at the present day, though it formerly enjoyed a great celebrity. It used particularly to be given to children as an emetic; and it used also to be applied to the temples.—Dose, from half a dram to two drams and a half.—Each ounce, in the formula of WIRTEM., contains five grains of opium.

OPOPONAX.

SYNONIMES :—*Gummi Panacis*. — Opopanax, FR.; Heilwurzgummi, Opopanaxharz; Panaxgummi, Opopanax gummi, GERM.; Panaxgummi, DAN.; Opopannaco, PORT.; and SPAN.

LOND.; DUBL.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; GENEV.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; SPELMANN; ZARDA.

This gum resin is in irregular pieces, seldom in tears, of a reddish-yellow colour externally, and of a dull white varied with red and yellow internally; it is opaque, dry, and friable. It has a rather strong, aromatic, but not very agreeable odour, and an acid and warm taste.

VOL. II.

It is the inspissated milky juice which is obtained from incisions made in the root of the *Pastinaca Opoponax*, LINN.; a plant of the south of Europe and of the East Indies. (Fig. Gouan, *Illustr.* t. 13 and 14). But it is only yielded by the plant which grows in the east, from whence it comes to us by Syria.

It contains, according to PELLETIER, resin, gum, and a little starch.

It is excitant.—Dose, from a scruple to half a dram.

DETERGENT FOMENTATION. (BRERA)

R. Root of Round Birthwort,	}	of each,	half an ounce.
— Long Birthwort,			
— Florentine Iris,			
Opopanax,	}	of each, two drams.	
Sagapenum,			
Guaiac,			four scruples.

2 T

Cloves, two drams.
Camphor, three drams.
Proof Spirit, ten ounces.

Strain after twenty-four hours' infusion.

It is recommended to be applied to sordid syphilitic ulcers.

ANTHELMINTIC EMULSION. (BORIES)

R. Opoponax, } of each, one dram.
Venetian Soap, }
Yolk of Egg, two drams.
Syrup of Mugwort, one ounce and a half.
Distilled Fennel Water, three ounces.

Mix.—Dose, a tea-spoonful every hour.

ORIGANUM.

Four species of this genus of plants are employed.

1. *Origanum Dictamnus*, LINN. — Dittany of Crete, Cretan Dittany, Dictamnus of Candia.

SYNONIMES :—Dictame, Dictame de Crète, FR. ; Cretischer Diptam, GERM. ; Krets-ky Dyptam, BOHEM.

ANT. ; BRUNS. ; HISP. ; FERR. ; FORM. HÔP. FR. ; HERBIP. ; BERGIUS ; GUIBOURT ; SPIELMANN ; ZARDA.

A small shrub of Crete. (*Didynamia Gymnospermia*, LINN. ; *Labiata*, JUSS. ; *Fig. Zorn, Ic. pl. t. 542*)

The herb (*Herba seu Folia Dictamni Cre-tici*) is employed. It consists of a branching, reddish, hairy stem, furnished with leaves, the lower ones of which are oval, rounded, petiolated, whitish, and cottony ; and the upper, rounded, sessile, glabrous, frequently reddish, and covered with numerous glandular points. It has a strong and balsamic odour, and an aromatic and bitter taste.

It is stimulant, and used to be esteemed emmenagogue.

OIL OF DITTANY OF CRETE.

Oleum Dictamni athereum, Atheroleum Dictamni. (SWEDIAUR)

R. Leaves of Dittany, one part.
Water, four parts.
Distil.

SYRUP OF DITTANY.

Syrupus de Ditamno. (GALL.)

R. Dried Tops of Dittany of Crete, one ounce.
Distilled Water of Dittany of Crete, two pounds.

Digest in a water-bath for two hours in a covered vessel ; then strain and filter. Add,

White Sugar, twice the weight,
and make a syrup in a water-bath.

2. *Origanum Marjorana*, LINN. ; *μυρρακον*. —Marjoram, Sweet Marjoram.

SYNONIMES :—Marjolaine, FR. ; Mayran, Majoran, GERM. ; Mirzunjach, ARAB. ; Maryanka, BOHEM. ; Merian, Mairan, DAN. ; Murwa, DUK. ; Meiorana, SPAN. ; Mariolein, DUT. ; Maggioreana, ITAL. ; Majeran, POL. ; Mangerona, PORT. ; Mejram, SWED. ; Mairu, TAM.

DUBL. ; EDINB. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; AINSLIE ; BERGIUS ; BRUGNATELLI ; COKE ; GUIBOURT ; MURRAY and GME-LIN ; PIDERIT ; SPIELMANN.

A plant common in Europe. (*Fig. Zorn, Ic. pl. t. 81*)

The tops (*Herba et Flores Majoranae seu Amoraci seu Sampsuchi*) are employed. They are composed of a quadrangular stem, furnished with oval, cottony, whitish leaves, and with white flowers disposed in short, rounded spikes, which, standing in threes or fours, form a kind of head. It has an agreeable odour, and an aromatic, acrid, and bitterish taste.

It is stimulant and errhine.

CONSERVE OF MARJORAM. (SARD. ; WIRTEM.)

R. Herb of Marjoram, one part.
Sugar, two parts.
Beat together. (WIRTEM.)
SARD. prescribes an equal part of each.

DISTILLED WATER OF MARJORAM. (AMST. ; BATAV. ; PALAT. ; SAX. ; SARD. ; WUR-CEB.)

R. Herb of Marjoram, one part.
Water, four parts.

Distil one half, and re-distil with fresh herb. (PALAT.; SARD.)

AMST.; and BATAV. prescribe one part of herb and a sufficient quantity of water; ten parts to be distilled.—WURCEB.—one of herb and six of water.—SAX.—one of herb and fifteen of water; two-thirds to be distilled.

Dose, from two to four ounces.

ESSENTIAL OIL OF MARJORAM. (EDINB.; BAVAR.; BRUNS.; DAN.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; SARD.; WIRTEM.; HERBIP.; SWEDIAUR)

R. Flowering Herb of Marjoram, any quantity.
Water, a sufficient quantity.
Distil, and separate the oil. (EDINB.)

FULD.; and LIPP. prescribe one part of herb and three parts of water.—SWEDIAUR—one of oil and four of water.—DAN.—one of herb and six of water.—HANN.; OLDEN.; POL.; BORUS.; and SAX.—one of herb and eight of water.—BAVAR.—one of herb and nine of water.

R. Herb of Marjoram, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Macerate for three days, and distil. (BRUNS.; PALAT.; SARD.; WIRTEM.)

R. Herb of Marjoram, any quantity.
Distilled Water of Marjoram, a sufficient quantity.

Distil. (HERBIP.)

BALSAM OF MARJORAM. (WIRTEM.; SPIELMANN)

R. Essential Oil of Marjoram, one dram.
Nutmeg, two drams and a half.

Mix. (WIRTEM.)

SPIELMANN prescribes an equal part of each.

BUTTER OF MARJORAM.

Butyrum seu Unguentum Majoranæ. (BRUNS.; DAN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; WIRTEM.; PIDERIT; SPIELMANN; VAN MONS)

R. Fresh Herb of Marjoram, } of each, an
Butter, } equal part.

Beat together, boil slowly to the consumption of the humidity, and express. (HASS.; SAX.; PIDERIT; SPIELMANN)

HANN.; OLDEN.; and BORUS. prescribe one part of herb and two parts of lard.

R. Fresh Herb of Marjoram, } of each,
Butter, } one pound.

Boil to the consumption of the humidity, strain, and add

Essential Oil of Marjoram, thirty drops.

Mix. (BRUNS.; DAN.; PALAT.; WIRTEM.)

R. Fresh Butter, three parts.
Herb of Marjoram, one part.

Digest for twelve hours in warm ashes; then express, and wash the product in warm water; then melt, decant, and stir till cold. (VAN MONS)

R. Fresh Butter, one ounce.
Distilled Water of Marjoram, two drams.

Triturate together; then melt over a slow fire, pour off the butter, and triturate it with

Essential Oil of Marjoram, eight drops.

Mix. (FULD.)

R. Fresh Butter, one dram.
Essential Oil of Marjoram, two drops.

Triturate together. (LIPP.)

POL. prescribes half an ounce of lard and ten drops of oil.

It is emollient, resolvent, and carminative.—To be used as a liniment.

OIL OF MARJORAM BY INFUSION.

Oleum Majoranæ infusum. (BRUNS.; WIRTEM.; HERBIP.)

R. Fresh Herb of Marjoram, } of each,
Olive Oil, } one pound.

Macerate for several days; then boil over a very slow fire to the consumption of the humidity, and express. (BRUNS.; WIRTEM.)

HERBIP. prescribes one part of herb and two parts of oil.

3. *Origanum vulgare*, LINN.—Origanum, Wild Marjoram, Common Marjoram.

SYNONIMES:—Origan commun, FR.; Wohlgemuth, Gemeine Doste, Dostenkraut, GERM.; Dobra mysl, BOHEM.; Vild marvian, Tost, DAN.; Oregano, SPAN.; Wild majolyn, DUT.; Origano, ITAL.; Lebiotka, POL.; Ouregao, PORT.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.;

BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 57*)

The tops (*Herba et Flores Origani vulgaris seu Cunila bulbosa*) are employed. They consist of a slightly quadrangular pubescent stem, branching only at the superior part, furnished with oval, petiolated leaves, rather hairy underneath, and with purpurine flowers, disposed at the summit in short spikes, which, by their approximation, form a paniculated corymb. It has an aromatic and agreeable odour, and an acrid, warm, and distinct taste.

INFUSION OF WILD MARJORAM. (FORM. HÔP. FR.)

R. Dried Tops of Marjoram, } of each,
Grated Liquorice, } two drams
Boiling Water, a sufficient quantity
to obtain two pounds of strained infusion.

ESSENTIAL OIL OF COMMON MARJORAM.

Oleum Origani æthereum, Ætheroleum Origani. (LOND.; DUBL.; AMER.; ANT.; AUSTR.; BRUNS.; DAN.; HANN.; OLDEN.; PALAT.; BORUS.; SARD.; SAX.; HERBIP.; WIRTEM.; COXE; SWEDIAUR)

Oleum Origani. (LOND.; DUBL.)

R. Flowering Tops of Marjoram, any quantity.
Water, a sufficient quantity
to prevent empyreuma.

AMER.; ROSS.; and COXE give the same formula.

SWEDIAUR prescribes one part of herb and four parts of water.—DAN.—one of herb and six of water.—HANN.; OLDEN.;

BORUS.; and SAX.—one of herb and eight of water.—AUSTR.—one of herb and twelve of water.—ANT.—six of herb and twenty of water.

R. Tops of Marjoram, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Distil. (BRUNS.; PALAT.; SARD.; WIRTEM.)

R. Tops of Marjoram, one part.
Distilled Water of Marjoram, twelve parts.

Distil four parts. (AUSTR.)

PALAT. prescribes one part of herb and four parts of water; two to be distilled.

4. *Origanum Creticum*, LINN.—Cretan Marjoram, Marjoram of Candia.

SYNONIMES:—Origan de Crète, FR.; Kelscher, Spanischer Dosten, Spamscher Hopfer, Candischer Majoran, GERM.; Ouregao de Creta, PORT.; Spanskhumle, SWED.

AMST.; ANT.; BRUNS.; GALL.; FENN.; FORM. HÔP. FR.; HANN.; HASS.; OLDEN.; BORUS.; SAX.; SUEC.; WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

A plant of Crete and of the south of Europe. (*Fig. Dod. Pempt. t. 284*)

The tops (*Spica seu Herba Origani Cretici*) are employed. They consist of reddish, hairy, branching stems, furnished with round, sessile, glabrous leaves, and purpurine flowers disposed in short sparse spikes. It has a strong, aromatic, and agreeable odour, and an aromatic and warm taste.

This plant may be used indiscriminately with the preceding. They are both stimulant, nervine, and carminative: an infusion of them is applied externally.

OROBANCHE VIRGINIANA.—VIRGINIAN BROOM-RAPE, BEECH DROPS, OR CANCER ROOT.

SYNONIME:—Orobanché de Virginie, FR. COXE.

A parasitical plant of North America. (*Didymamia Angiospermia*, LINN.; *Orobanchæ*, JUSS.; *Fig. Moris. p. 12, t. 16, f. 9*)

It is employed internally in dysentery, and externally in chronic ulcers. It is astringent, and possesses at the same time an extremely nauseous bitter taste.

ORYZA SATIVA, LINN.—RICE.

SYNONIMES:—Riz, FR.; Reiss, GERM.; ARUZ, ARAB.; Prinsch, ARM.; Men-ho, CHIN.; Lua, COCHIN.; Ziis, DAN.; Chawl, DUK.; and HIND.; Arroze, SPAN.; Ryst, DUT.; Riso, ITAL.; Motsj, GOME, KO, JAP.; Riskesa, MAL.; Tandul, MAH.; Bras, MALAB.; Barinje, PERS.; Ryz, POL.; Arroz, PORT.; Vrihi, SANS.; Ris, SWED.; Dugu, TART.; Arisi TAM.; Buim, TEL.

AMST.; ANT.; BATAV.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HASS.; LIPP.; ROSS.; WIRTEM.; AINSLIE; BERGIUS; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

This plant is originally of Ethiopia, but is now cultivated in all warm climates. (*Hexandria Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Flore Medic. VI. 299*)

The seed (*Semen seu Grana Oryzæ*) is employed. It is oblong, obtuse, grooved, white, hard, dry, friable, inodorous, and of a farinaceous taste. It comes to us deprived of its pellicle.

It contains, according to BRACONNOT, a considerable proportion of starch, a little animalized matter, a very little uncrystallizable sugar, gum, oil, and various salts.

It forms a very nutritious aliment; and when combined with much water is rather emollient and demulcent; it is erroneously supposed astringent.

DECOCTION OF RICE.

Decoctum seu Ptisana Oryzæ. (GALL.; FORM. HÔP. FR.; DISP. CHA. PARIS; NIEMANN; RATIER; SAINTE-MARIE)

R. Washed Rice, half an ounce.
Water, two pounds eight ounces.
Boil down to two pounds; strain, and add Syrup of Marshmallows, one ounce.
Mix. (GALL.)

NIEMANN directs two ounces of rice to be boiled in two pounds of water till reduced to sixteen ounces.—RATIER—half an ounce of rice to be boiled in three pounds of water till reduced to two pounds.

R. Washed Rice, two ounces.
Water, three pounds.

Boil down to two pounds, and add towards the end

Fresh Butter, one ounce.
Sugar, or Salt, a sufficient quantity to make it agreeable. Mix. (SAINTE-MARIE)

R. Washed Rice, one ounce.
Boil it till it has burst, in

Water, a sufficient quantity to obtain one pound and a half of strained liquor; add towards the end, and leave to infuse for a quarter of an hour,

Bitter Orange Peel, two drams.
Strain. (FORM. HÔP. FR.)

R. Rice, one ounce.
Water, two pounds and a half.

Boil down to two pounds, adding towards the end

Liquorice, four drams.
Strain when cold. (DISP. CHA. PARIS)

It is emollient, but is improperly considered astringent, and is prescribed in diarrhoea and dysentery.—Milk, decoction of catechu, RABEL's liquor, syrup of quince, or any acidulous syrup, may be added.

PECTORAL DECOCTION. (SPIELMANN)

R. Rice, one ounce.
Spring Water, one hundred and twenty-eight ounces.

Boil till the rice bursts; then infuse in the decoction

Root of Liquorice, } of each, half
—— Marshmallows, } an ounce.
Canadian Maidenhair, } of each,
Flowers of Coltsfoot, } two drams.
—— Red Poppy, one dram.

Pour off at the end of a quarter of an hour.

ACIDULATED DECOCTION OF RICE.

Potus Oryzæ citratus. (AUGUSTIN)

R. Rice, in powder, one ounce.
Water, four pounds.

Boil for a short time, and add to the strained liquor

Barley Sugar, half an ounce.
Lemon Juice, one ounce.

Mix.

OVUM GALLINACEUM, LINN.—PULLET'S EGG.

SYNONIMES :—Œuf de poule, FR.; Hühnereyer, GERM.; Bayzah, ARAB.; Biju, CYNG.; Anda, DUK.; Huevo, SPAN.; Hoenderey, DUT.; Ovo di gallina, ITAL.; Tukim, PERS.; Kurze iaia, POL.; Anda, SANS.; Hœnroegg, SWED.; Koruy mutay, TAM.; Guddu, TEL.

LOND.; AMST.; ANT.; AUSTR.; BAVAR.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FULD.; FORM. HÔP. FR.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; AINSLIE.; BRUGNATELLI.; COKE.; GUIBOURT.; PARIS.; PIDERIT.; SPIELMANN.

OIL OF EGG.

Oleum e Vitellis Ovorum seu Ovorum expressum. (ANT.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SARD.; SAX.; SUEC.; HERBIP.; WIRTEM.; BRUGNATELLI.; PIDERIT.; VAN MONS)

R. Fresh Yolks of Eggs, any quantity.

Put them in a silver pan, boil till reduced to one half the quantity, in a water-bath, when, on being pressed between the fingers, the oil will be easily separated; then enclose the mass in a close linen bag, express between two heated plates, and filter, at the heat of a water-bath, through bibulous paper.

CHAUSSIER proposed to mix the hard yolks with alcohol, to heat the mixture in a covered vessel, to filter, then to distil, and lastly, to separate the oil with water. (HENRI.; VAN MONS)—But the oil obtained in this manner always retains, according to HENRI, the odour of alcohol; and, according to VAN MONS, it is more subject to become rancid.

EMOLLIENT LAVEMENT.

Clyisma sedans et emolliens. (MIL. COPEN.; CAST. BORUS.; BRERA.; PIERQUIN)

R. Well-washed Bran, half a handful.
Pure Water, two pounds.

Boil down to one half, and add when strained,

Fresh Yolks of Eggs, three or four.

Mix. (PIERQUIN)

R. Yolks of Eggs, two.
Rose Oil, } of each,
Chamomile Oil, } three ounces.

Mix. (BRERA)

R. Yolk of Egg, half an ounce.
Olive Oil, two drams.

Mix. (MIL. COPEN.; CAST. BORUS.)

ANTI-STRUMOUS OINTMENT. (SAINTE-MARIE)

R. Lard, one ounce.
Fresh Egg, without shell, one.

Shake the egg well with the lard melted over a slow fire.

ALPHONSE LEROY applied it to goitrous tumours, during three or four months. It is said to have sometimes removed them.

YELLOW PLASTER.

Emplastrum flavum. (OLDEN.)

R. Hard Yolks of Eggs, six.
Yellow Wax, } of each, four ounces.
Lard, }

Mix.

ANALEPTIC MIXTURE. (SAINTE-MARIE)

R. Cream, six ounces.
Fresh Yolks of Eggs, two.
Sugar, from four drams to one ounce,
Hordeated Cinnamon Water, six drams.

Mix well.—To be taken by spoonfuls, in the debility occasioned by excessive venery.

ANTI-APHTHOUS MIXTURE.. (SAINTE-MARIE)

R. Cream, } of each, one ounce.
Rose Water, }
Fresh Yolk of Egg, one.
Syrup of Poppies, four drams.

Mix.—To be taken by spoonfuls, in twenty-four hours.

ALBUMINOUS ALCOHOL. (VAN MONS)

R. Fresh Whites of Eggs, four.
Beat them in a porcelain vessel, adding gradually,
Alcohol, two ounces.

OXALIS ACETOSELLA, LINN.—WOOD SORREL.

SYNONIMES:—Alleluia, Surelle, Pain de Coucou, FR.; Sauerklee, Gukkusklee, Buchsauerklee, GERM.; Alleluga, Detel kysclý zagecý, BOHEM.; Giogeurt, Giogemad, suurklover, Skovsyre, DAN.; Acederilla, SPAN.; Zuurklavor, DUT.; Alleluja, ITAL.; Szczawik, POL.; Trewo azedo, PORT.; Harszya, SWED.

LOND.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; GENEV.; HASS.; LUSIT.; BORUS.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; MURRAY and GMELIN; PARIS; SPIELMANN; ZARDA.

This plant is met with in almost every part of Europe. (*Hexandria Pentagynia*, LINN.; *Oxalideæ*, CAND.; *Fig. Zorn, Ic. pl. t. 9*)

The herb (*Herba Acetosellæ seu Lujulæ seu Alleluja seu Trifolii acetosi, seu Oxytriphylly*) is employed. It is composed entirely of radical leaves, having long footstalks, presenting three heart-shaped leaflets, the apex of the heart joining the footstalk; it is of a pale green colour. It presents floriferous scapes, at the middle part of which there are two small opposite bractæ. It is inodorous; its taste is acid and rather agreeable, but is lost by desiccation.

The *Yellow Wood Sorrel*, *Oxalis corniculata*, LINN., is also used, (*Chua miba chia*, CHIN.; and *COCHIN.*; *Umbuti*, DUK.; *Amrul*, HIND.; *Chukrika*, *Ambachta*, *Amlika*, *SANS.*; *Puliaray*, *TAM.*; *Pullie chinta*, *Pulla chinta*, *TEL.*)—HANN.; AINSLIE.—It much resembles the preceding. [Not at all.—EDIT.]

Both contain super-protoxalate of potassium.

It is acidulous and refreshing.

CONSERVE OF WOOD SORREL. (FULD.; LUSIT.; PALAT.; ROSS.; SARD.; SUEC.; WIRTEM.; HERBIP.; PIDERIT; SWEDIAUR)

R. Fresh Leaves of Wood Sorrel, Sugar, } of each, an equal part.

Triturate together. (SARD.; SUEC.)

FULD.; PALAT.; ROSS.; WIRTEM.; HERBIP.; PIDERIT; and SWEDIAUR prescribe one part of herb and two parts of sugar.—AMST.—one of herb and three of sugar.

R. Powdered Leaves of Wood Sorrel, Distilled Water of Wood Sorrel, } three ounces, eight ounces.

Macerate for six hours; then add, constantly stirring,

White Sugar, one pound and a half.
Mix. (LUSIT.)

DISTILLED WATER OF WOOD SORREL.

Aqua e Succo Acetosellæ. (PALAT.; SARD.)

R. Bruised Herb of Wood Sorrel, one part.
Water, two parts.

Distil two thirds. (SARD.)

PALAT. prescribes one part of herb and three parts of water; one half to be distilled.

It may also be obtained by distilling the depurated juice of the herb in a water-bath to dryness.

JUICE OF WOOD SORREL. (PALAT.; SARD.; HERBIP.; WIRTEM.)

R. Fresh Herb of Wood Sorrel, any quantity.

Bruise it in a stone mortar with a wooden pestle; express the juice, set it aside, decant, and preserve it in a cool place.

HERBIP. directs one dram of alcohol to be added to each pound of the juice.

SYRUP OF WOOD SORREL.

Syrupus Acetosellæ. (SARD.; WIRTEM.)

R. Depurated Juice of Wood Sorrel, ten ounces.
White Sugar, sixteen ounces.

Boil to a syrup. (WIRTEM.)

SARD. directs a sufficient quantity of depurated juice to be added to simple syrup, and the mixture to be evaporated slightly in a water-bath.

EXTRACT OF WOOD SORREL.

Extractum Acetosellæ. (SARD.; HERBIP.; PIDERIT)

Bruise it in a stone mortar, express the juice and evaporate it to the consistence of an extract, at first over a slow fire, and then in a water-bath. (SARD.; HERBIP.)

R. Fresh Herb of Wood Sorrel, one pound.

Bruise it in a mortar, express the juice, and pour upon the residue,

Water, three pounds.

Boil for one hour, express, and evaporate to the consistence of an extract; then add the juice, set aside, and evaporate again. (PIDERIT)

It is acidulous and bitter.—Dose, from a scruple to half a dram.—This extract owes its activity only, to the super-oxalate of potass which it contains.

OXYGENIUM.—OXYGEN.

Aer purus seu dephlogisticatus seu vitalis.
(FERR.; HANN.; BRUGNATELLI; NIE-
MANN; SWEDIAUR; VAN MONS)

R. Black Oxide of Manganese, any quantity.

Bring it gradually to a red heat in an earthen retort communicating with a pneumato-chemical apparatus. Having let the first portions of the disengaged gas escape, receive the remainder in glass bottles; cork the bottles under water, cover the mouths with wax, and preserve them in a cool place. (FERR.; HANN.; SWEDIAUR; NIE-
MANN; VAN MONS)

R. Peroxide of Manganese, three parts.
Concentrated Sulphuric Acid, two parts.

Mix without heat in a glass mortar, calcine the mixture to redness, collect the gas which is disengaged, and preserve it after having shaken it with lime water. (VAN MONS)

R. Nitrate of Potass, in powder, any quantity.

Heat it in a matrass till the bottom redens, and receive the disengaged gas, letting the first portions pass off. (BRUGNATELLI)

VAN MONS says that nitre mixed with sea salt, yields its oxygen more readily than when it is alone.

R. Peroxide of Manganese, one part.
Chlorate of Potass, two parts.

Triturate together, apply the heat of a spirit-lamp, and collect the gas which is liberated. (VAN MONS)

OXYGENATED WATER.

Aqua oxygenata. (BRUGNATELLI)

R. Water, any quantity.

Saturate it perfectly with oxygen gas, by the aid of a proper machine.

It is slightly stimulant, and has been recommended in defective appetite, amenorrhœa, asthenic dropsies, spasms of the stomach, and hysteria.—Dose, one or two bottles in the space of twenty-four hours.

In general the proportion of oxygen con-

tained in the water thus prepared is as one volume to two volumes of the latter.

Thus prepared the oxygenated water differs very much from that which THENARD obtained by a process as long as it was difficult and minute, and to which he gives the name of *Deutoxide of Hydrogen*. This, when it is in as concentrated a state as possible, yields four hundred and seventy-five times its volume of gas, under the pressure of 0,76, and at the temperature of fourteen degrees. It is rather surprising that no one has thought of applying it to practice either internally or externally, varying its degree of concentration by the addition of a greater or less quantity of water. It is also surprising that our experimenters should have neglected this, which would have afforded them so fine an opportunity for feeding the public curiosity for several months with the recital of their daring trials. It would be well to remember the important observation made by THENARD, that the animal organs and tissues effect the decomposition of the oxygenated water in the same manner as the greater part of the metals and metallic oxides do, in not yielding any part of their principles, absorbing not the smallest quantity of oxygen, and consequently not undergoing the least apparent alteration, unless the peroxide be highly concentrated. Can it be the same with regard to the living tissues? The fact would deserve to be examined were it only to ascertain whether THENARD is right or wrong in supposing that, the animal tissues having, like platina, gold, and silver, the property of disengaging oxygen from oxygenated water, may be referred to a similar action; for if it is so, it is then not unreasonable to think that the secretions are also effected by a similar action: and consequently it may be conceived how an organ, without absorbing any thing, without ceding any thing, may constantly act upon a fluid and transform it into new products. These reasonable deductions from a positive fact are worthy of an attentive examination, as they are of a kind to completely overturn all our present theories in physiology and therapeutics.

PÆONIA OFFICINALIS, LINN.—PEONY.

SYNONIMES: — Pivoine, FR.; Pæonie, Pflingstrose, Gichtrose, GERM.; Peonia, SPAN.; Piwoniec, POL.

AMST.; ANT.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; HANN.; POL.; BORUS.; ROSS.; SAX.; HERBIP.; WIRTEM.;

BERGIUS; BRUGNATELLI; GUIBOURT;
MURRAY and GMELIN; SPIELMANN.

A plant of the South of Europe. (*Pentandria Digynia*, LINN.; *Papaveraceæ*, JUSS.; *Fig. Flore Medic. VI. 274*)

The root, flowers, and seeds are employed.

The root (*Radix Pæoniæ*) is large, long, tuberoso, and fleshy.

In the fresh state, its smell is disagreeable and virous, but is very feeble after desiccation. Its taste is bitter, sweetish, rather acrid, and feebly astringent: its acidity becomes dissipated by drying.

The flowers (*Rosæ benedictæ seu regiæ*) are red. When fresh, their odour is disagreeable; their taste is mucilaginous and bitterish.

The seeds are inodorous and nearly insipid.

CONSERVE OF PEONY. (PALAT.; SARD.; WIRTEM.)

R. Petals of Peony, one part.
Sugar, two parts.

Triturate together. (PALAT.; WIRTEM.)

SARD. directs one part of petals to be beaten with a little sugar, and the pulp to be boiled a little with three parts of simple syrup.

FECULA OF PEONY.

Fecula Peoniæ. (WIRTEM.)

R. Fresh Root of Peony, any quantity.

Grate it, enclose the pulp in a linen bag, and submit it to the press. Add a little water to the juice, set it apart, decant the clear liquor, dry the remainder in the open air, and reduce it to powder.

DISTILLED WATER OF PEONY. (GALL.; PALAT.; SARD.)

R. Flowers of Peony, one part.
Water, four parts.

Distil one half. (GALL.; SARD.)

PALAT. prescribes one part of flowers, and three parts of water; one and a half to be distilled.

EXTRACT OF PEONY. (FERR.; WIRTEM.)

R. Root of Peony, sliced, one pound.
Boiling Water, six pounds.

Infuse for three days, then boil a little; strain with expression, and evaporate the decoction to the consistence of an extract. (WIRTEM.)

FERR. directs one pound of fresh root to be boiled in three pounds of water; the decoction to be clarified with white of egg, and evaporated.

VOL. II.

INFUSION OF PEONY FLOWERS. (HISP.)

R. Fresh Petals of Peony, six pounds.
Boiling Water, twelve pounds.

Infuse for eight hours; strain with expression, and decant the liquor after a short time.

SYRUP OF PEONY. (HISP.; PALAT.; WIRTEM.)

R. Root of Peony, bruised, two ounces.
Water, six pounds.

Boil down to four pounds, and add to the strained liquor

White Sugar, two pounds.

Clarify, and make a syrup. (HISP.)

R. Fresh Flowers of Peony, three pounds and a half.
Boiling Water, four pounds.

Infuse for twenty-four hours; express, and dissolve in eighteen ounces of the liquor

White Sugar, thirty-two ounces.

Preserve. (PALAT.; WIRTEM.)

Dose, from two to three drams, in a draught.

ANTI-SPASMODIC BROTH. (BORIES)

R. Lean of Veal, six ounces.
Root of Valerian, } of each,
Male Peony, } one dram.
Leaves of Cichory, } of each,
Lettuce, } half a handful.
Water, a sufficient quantity.

Boil, and add to the cold strained liquor

Powdered Orange Leaves, half a dram.

Sulphuric Æther, ten drops.

To be taken at once, in the morning, fasting.

ANTI-EPILEPTIC BROTH. (BORIES)

R. Neck of Mutton, four ounces.
Root of Cichory, } of each,
Strawberries, } half an ounce.
Leaves of Cichory, } of each,
Borage, } half a handful.
Lettuce, }
Agrimony, }
Balm, one pinch.
Seeds of White Poppy, half a dram.
Water, a sufficient quantity.

Boil, and add to the strained liquor

Root of Peony, in powder, two scruples.

To be taken in the morning, fasting.

ANTI-APOPLECTIC SPECIES.

Species contra Apoplexiam. (PALAT.; WIRTEM.)

R. Prepared Amber, half an ounce.
2 U

Prepared Hartshorn,	} of each,	
Aloe Wood,		half an ounce.
Root of Peony,	} one dram.	
Lemon Peel,		
Sugar,		three ounces.
Oil of Cinnamon,		sixteen drops

Make a powder.

They are called *completæ* if there is super-added

Ambergris,	half a scruple.
Musk,	five grains.

This powder was formerly in repute against head-ach, vertigo, apoplexy, and epilepsy.—Dose, from half a dram to two scruples.

ANTI-CATARRHAL POWDER.

Pulvis contra Tussim Infantum. (PALAT.)

R. Root of Peony,	half an ounce.
—— Florentine Iris,	six drams.
—— Serpentaria,	two drams.
Saffron,	one dram.
Sugar Candy,	one ounce.

Make a powder.

ANTI-EPILEPTIC POWDER.

Pulvis epilepticus seu Pæoniæ compositus seu de Gutteta seu ad Guttetam seu cephalicus. (BRUNS.; HISP.; PALAT.; WIRTEM.; CADET DE GASSICOURT; FEE; HENRI; PIERQUIN; SPIELMANN)

R. Root of Peony,	} of each, one part.	
Sub-carbonate of Ammonia,		
Prepared Hartshorn,	} of each,	
Root of Fraxinella,		two parts.
Miseltoe,	} four parts.	
Prepared Amber,		

Pulverize. (CADET DE GASSICOURT; FEE; HENRI)

This is the *Poudre de la princesse de Caughan*.—Dose, eighteen grains, to a child of one year old; thirty-six, to one of two years; forty-eight, to one of three years; and one dram, to one of four and above. It is prescribed in convulsive disorders.

R. Root of Peony,	} of each,	
—— Fraxinella,		half an ounce.
Miseltoe,	} two drams.	
Seeds of Orache,		
—— Peony,	} of each,	
Elk's Hoof,		half an ounce.
Prepared Human Skull,	} one ounce.	
—— Red Coral,		
—— Hyacinth,	} half an ounce.	
Gold Leaves,		fifty.

Triturate together. (WIRTEM.)

Dose, from half a scruple to half a dram.

R. Root of Peony, one ounce and a half.		
—— Contrayerva,	} of each,	
Miseltoe,		six drams.
Rasped Ivory,	} of each,	
—— Hartshorn,		
—— Elk's Hoof,		one dram
Prepared Human Skull,		and a half.
—— Red Coral,		
—— Hyacinth,	} Occidental Bezoar,	
Occidental Bezoar,		

Make a fine powder. (HISP.)

R. Miseltoe,	} of each,	
Root of Fraxinella,		one ounce.
—— Peony,	} of each,	
Seeds of Peony,		
—— Orache,		half an ounce.
Prepared Red Coral,		one ounce.
Elk's Hoof,	} one ounce.	

Make a fine powder. (SPIELMANN)

R. Miseltoe,	} of each,	
Root of Valerian,		half an ounce.
Magnesia,	} of each,	
Seeds of Peony,		two drams.
—— Knotgrass,		
Prepared Red Coral,		

Make a powder. (PIERQUIN)

These four formulæ give the powder which is more particularly designated *Pulvis ad Guttetam*.

R.	Root of Peony,	}	three ounces.
	Miseltoe,		
	Rasped Ivory,		
	—— Elk's Hoof,		
	—— Hartshorn,		
	Burnt Ivory,	}	of each,
	Red Coral,		
	White Coral,		
	Prepared Pearls,		
	Gold Leaves,		
			six drams.
			twenty.

Make a powder. (WIRTEM.; SPIELMANN)

BRUNS. prescribes three ounces of peony, six drams each, of miseltoe, of hartshorn, of magnesia, of calcined ivory, of each of the two corals, and of mother of pearl, and ten gold leaves.

This formula gives the *Pulvis epilepticus Marchionis*, *Pulvis Marchionis*.—Dose, from ten grains to half a dram.

R. Prepared Amber,	} of each,	
—— Mother of Pearl,		half an ounce.
—— Emerald,	} of each,	
Aloe Wood,		two scruples.
Root of Peony,	} of each,	
Sea Unicorn,		one dram and a half.
Miseltoe,	} two ounces.	
Sugar Candy,		

Mix. (WIRTEM.)

This is the *Pulvis cephalicus albus*.—Dose, from half a dram to two scruples.

R. Root of Peony,	half an ounce.
Miseltoe,	} of each, one dram.
Rasped Ivory,	
— Elk's Hoof,	
— Calcined Ivory,	
— Hartshorn,	
Red Coral,	}
White Coral,	
Prepared Pearls,	

Mix. (PALAT.)

This is the *Pulvis epilepticus albus*.

R. Root of Peony,	of each,
Seeds of Peony,	half an ounce.
Horn of Sea Unicorn,	of each,
Oriental Pearls,	two drams.
The five precious Stones,	of each, forty-five grains.

Castor,	} of each, two drams.
Miseltoe,	
Elk's Hoof,	
Red Coral,	
Prepared Human Skull,	one dram and a half.
Powdered Viper,	one dram.

Mix. (WIRTEM.)

This is the *Pulvis epilepticus Cellarii*.—
Dose, from one scruple to half a dram.

R. Root of Peony,	ten drams.
— Fraxinella,	of each,
Miseltoe,	five drams.
Aloe Wood,	of each,
Seeds of Peony,	three drams.
Red Coral,	of each,
Mother of Pearl,	half an ounce.
Elk's Hoof,	three drams.
White Amber,	of each,
Horn of Sea Unicorn,	two drams.
Charcoal of Lime Wood,	one ounce and a half.

Make a powder. (PALAT. ; WIRTEM.)

This is the *Pulvis epilepticus niger*.—
Dose, from five grains to half a scruple,
to children ; half a dram, to adults.

R. Root of Peony,	of each,
— Gentian,	two ounces.
Herb of Betony,	} of each, one ounce.
— Germander,	
— Cinquefoil,	
Flowers of Peony,	

Flowers of Lily of the Valley,	} of each, one ounce.
— Violets,	
— Lime Tree,	
Seeds of Coriander,	of each,
— Peony,	half an ounce.
Miseltoe,	} two ounces.
Rasped Elk's Hoof,	
— Asses Hoof,	five drams.
— Hartshorn,	of each,
Hartshorn prepared without fire,	one ounce.
Human Skull,	six drams.
Red Coral,	} of each,
White Coral,	
Oriental Pearls,	half an ounce.
Gold Leaves,	twenty-four.

Make a powder. (WIRTEM.)

This is the *Pulvis epilepticus Holsaticus*.
—Dose, from half a scruple to half a dram.

ANTI-EPILEPTIC ELECTUARY. (BORIES)

R. Saffron,	half an ounce.
Senna,	} of each, two drams.
Rhubarb,	
Seeds of Peony,	} of each, one dram.
Muriate of Ammonia,	
Jalap,	
Scammony,	
Black Hellebore,	
Succinic Acid,	} two scruples.
Cinnamon,	
Castor,	one scruple.
Conserve of Peony,	a sufficient quantity.

Mix.—Dose, one dram and a half, in the morning, fasting. It is very evident that the peony is of very secondary importance in such a purgative farrago.

ANTHELMINTIC ELECTUARY. (BORIES ; PIERQUIN)

R. Sulphate of Iron,	of each,
Corsican Moss,	two drams.
Root of Valerian,	of each,
— Peony,	half an ounce.
Orange Leaves,	five drams.
Aloes,	one dram.
Syrup of Peony,	a sufficient quantity.

Mix.—Dose, one dram, fasting, every morning, in epilepsy from worms. What was said of the preceding formula applies equally to this.

PANAX QUINQUEFOLIUM, LINN.—GINSENG.

SYNONIMES :—Ginseng, FR. ; Fünfblättrige Kraftwurzel, GINZENG, GERM. ; Jinchén, CHIN. ; Garent-oguen, IROQ. ;

Nindsin, Dsindsom, JAV. ; Orkhoda, MAND. ; Ginsao, PORT. ; BRUNS. ; GALL. ; FERR. ; WIRTEM. ;

AINSLIE; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN.

A plant of Chinese Tartary and of Canada. (*Polyandria Diœcia*, LINN.; *Araliaceæ*, JUSS.; *Fig. Flore Medic. IV. 184*)

The root (*Radix Ginseng*) is employed. It is fusiform, of the thickness of the finger, and two or three inches in length, reddish without and yellowish within. Its crown is surmounted by a twisted knotty tissue upon which the vestiges of stalks are apparent. Its substance is semi-transparent, compact, and horny. It is destitute of odour, but has a saccharine, bitterish, and feebly aromatic taste.

It contains much starch and gum.

The Asiatics regard it as a powerful aphrodisiac; but it has no greater right to this title than the many other substances on

which it has been bestowed by what is called popular observation.

PASTILLES OF GINSENG. (BORIES; CADET DE GASSICOURT; PIERQUIN)

R. Sugar,	ten pounds.
Vanilla,	ten ounces.
Ginseng,	five ounces.
Tincture of Cantharides,	five drams.
Oil of Cinnamon,	one drop.
Essence of Ambergris,	ten drops.
Mucilage,	a sufficient quantity.

Make pastilles of from twenty-four to thirty grains.—Dose, four or five daily. These pastilles may occasion serious consequences; at the same time they will be found of no service in reviving, even for a moment, faculties which have ceased to respond to impotent desires.

PANICUM MILIACEUM.—MILLET.

SYNONIMES:—Millet, FR.; Hirse, DAN.; and GERM.; Hajo, SPAN.; Geerst, DUT.; Miglio, ITAL.; Proso, POL.; Milho, PORT.

HISP.; GALL.; MURRAY and GMELIN.

This plant is cultivated in several countries of Europe, but it is not known from what

part it is originally. (*Triandria Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 349*)

The seed (*Semen Milii*) is employed. It is small and round; under a thin, crustaceous, yellow, shining, bivalve pellicle, it contains a white and farinous substance.

PANIS.—BREAD.

BREAD DRINK.

Ptisana ex Pane. (NIEMANN)

R. Wheat Bread, eight ounces.
Water, thirty-two ounces.

Boil for an hour, then strain, and add to the liquid a quantity of water equal to that which has evaporated.

Drink prepared with the toasted crust of rye bread is more agreeable and rather astringent.

It is to be taken by glassfuls, in irritations of the alimentary passages.

PANADO.

Panada Anglorum. (AUGUSTIN)

R. Toasted Rye Bread, six ounces.
Cinnamon, half a dram.
Fresh Lemon Peel, two drams.
White Wine, two pounds.

Boil, and strain. The yolk of an egg or some aromatics may be added.

SYDENHAM'S WHITE DECOCTION.

Decoctum album seu album Sydenhami seu Cornu Cervi compositum seu Cornu Cervi Elaphi compositum seu Cornu Cervi cum Mica Panis. (ANT.; BELG.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; SARD.; SAX.; DISP. CHA. PARIS; CADET DE GASSICOURT; NIEMANN; PIERQUIN; RATIER; SPIELMANN; VAN MONS)

R. Shavings of Hartshorn, two ounces.
Crumb of Bread, four ounces.
Spring Water, six pounds.

Boil down to one third, and strain without expression. (SARD.)

R. Hartshorn, } of each,
Crumb of Bread, } two drams.
Water, } three pounds.

Boil down to two pounds, and add

White Sugar, any quantity.

Dissolve. (NIEMANN)

R. Hartshorn, two ounces.
Water, three pounds.
Boil, and add towards the end

Crumb of Bread, two ounces.

Having boiled the whole to two pounds, strain, and add

White Sugar, one ounce and a half.

Dissolve. (ANT. ; BELG.)

R. Hartshorn Shavings, half an ounce.
Crumb of Bread, one ounce.
Water, three pounds.

Boil down to two thirds, and add to the strained liquor

Syrup of Quince, two ounces.

Mix. (RATIER)

R. Hartshorn Shavings, } of each,
Crumb of Bread, } one half part.
Water, } thirty-six parts.

Boil down to two thirds, and add to the strained liquor,

Gum Arabic, one fourth part.

Dissolve. (SAX.)

HANN. prescribes half an ounce each, of hartshorn and of crumb of bread, and three pounds of water; to be boiled down to two thirds, and two drams of gum arabic added to the strained decoction.

R. Hartshorn Shavings, } of each,
Crumb of Bread, } half an ounce.
Water, } thirty-six ounces.

Boil down to two thirds; strain, and add

Gum Arabic, one dram and a half.

Sugar, two drams.

Dissolve. (SPIELMANN)

BRUNS. directs half an ounce each, of hartshorn and of crumb of bread, and one dram and a half of gum arabic to be boiled in three pounds of water till this is reduced to two, and two drams of sugar to be dissolved in the strained liquor.

R. Crumb of Bread, one part.

Boil it for half an hour with

Water, a sufficient quantity,
to obtain sixty-four ounces of strained liquor, add towards the end,

Gum Arabic, one half part.

Burnt Hartshorn, one fourth part.

Sugar, one half part.

Strain through flannel. When cold, decant, and add

Cinnamon Water, one half part.

Mix. (VAN MONS)

GALL. ; and DISP. CHA. PARIS direct two drams of calcined hartshorn to be boiled with six drams of crumb of bread and one ounce of sugar, in two pounds of water for eight minutes; the boiling liquor to be strained through a close sieve, with gentle

expression; and half an ounce of orange-flower water, or two drams of cinnamon water to be added.—FORM. HÔP. FR.—six drams of crumb of bread and two drams of hartshorn, or of any bone calcined to whiteness, to be boiled for eight minutes in a sufficient quantity of water to obtain two pounds of decoction; this to be strained while boiling, with slight expression, and a little lemon water and simple syrup added.—HISP.—half an ounce of calcined hartshorn, two ounces of crumb of bread, and six pounds of water, to be boiled slightly together; the decoction to be strained, and four ounces of simple syrup added, with a little aromatic water.—RATIER—two ounces of crumb of bread, two drams of calcined hartshorn, two pounds of water, one ounce of syrup, and one dram of tincture of cinnamon.—CADET DE GASSICOURT; and PIERQUIN—two ounces of crumb of bread, two drams of calcined hartshorn, three ounces of sugar, two pounds of boiling water, and half an ounce of cinnamon water.

It is emollient and demulcent, and is employed with advantage in chronic irritations of the alimentary canal, particularly in obstinate diarrhœa. A glassful to be taken every hour. From ten to twenty drops of SYDENHAM's laudanum may be frequently added with advantage.

BREAD JELLY. (HENRI ; NIEMANN ;
VAN MONS)

R. Stale Crumb of Bread, four ounces.
Distilled Water, two pounds.

Boil strongly, pass through a sieve, and add

White Sugar, one pound.

Wine, four ounces.

Cinnamon Water, six drams.

Mix. (HENRI)

R. Biscuit, four parts.
Water, eighty parts.

Boil for half an hour, pass through a close flannel, boil till twenty-four parts only remain, and add

Cinnamon Water, one part.

Lemon Juice, two parts.

White Sugar, four parts.

Rhenish Wine, six parts.

Mix. (NIEMANN ; VAN MONS)

EMOLLIENT CATAPLASM.

Cataplasma de Mica Panis seu emolliens.
(GALL. ; FERR. ; FULD. ; BRERA ; BRUGNATELLI ; NIEMANN ; SPIELMANN)

R. Stale Crumb of Bread, grated
four ounces.

Cows Milk or decoction of

Marshmallows, one pound and a half.

Boil to the proper consistence. (GALL.)

BRUGNATELLI directs ten ounces of crumb of bread and fifteen of milk to be mixed together without boiling.—BRERA—four pounds of milk and one pound of bread to be boiled together.

- R. Crumb of Bread, sixteen ounces.
 Melilot Flowers, in powder, } of each,
 Elder Flowers, in powder, } two
 Milk, forty-eight ounces. drams.

Boil. (SPIELMANN)

- R. Crumb of Bread, eight ounces.
 Fresh Butter, two ounces.
 Cow's Milk, eight ounces.

Boil together. (FULD.; NIEMANN)

The bread should be boiled with a little water, before boiling it with the milk, to remove from it the acetic acid which it contains.

- R. Crumb of Bread, fifteen ounces.
 Milk, twenty ounces.

Boiling Water, a sufficient quantity.
 Soften the bread in the water, strain it through a hair-sieve, and boil it with the milk. (FERR.)

SPIRIT OF BREAD. (WIRTEM.)

- R. Wheat Biscuit, any quantity.
 Distil slowly from a glass retort placed in a sand-bath, gradually raising the heat. Separate the spirit from the supernatant oil, and rectify it.

SPIRITUOUS WATER OF BREAD. (WIRTEM.; SPIELMANN)

- R. Crust of New Bread, four ounces.
 Cinnamon, } of each, half an ounce.
 Cloves, }
 Nutmegs, }
 White Wine, ninety-six ounces.

Distil one half.

It is an agreeable stomachic.—Dose, half an ounce.

PAPAVER.—POPPY.

Two species of this genus are employed in medicine:

1. *Papaver Rhæas*, LINN.—Red Poppy,
 Corn Poppy.

SYNONIMES: — Coquelicot, FR.; Klatschroze, GERM.; Plany neb wlej mak, BOHEM.; Klapperose, DAN.; Amapola, SPAN.; Klapperroos, DUT.; Rosolaccio, ITAL.; Paposileira, PORT.; Kornros, SWED.

LOND.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; MIL. COPEN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A very common plant of Europe. (*Polyandria Monogynia*, LINN.; *Papaveraceæ*, JUSS.; *Fig. Flore Medic. III. 134*)

The flowers (*Flores Papaveris erraticiseu Rhæadis*) are employed. The petals are red, oval and entire. They have a nauseous smell, but become deprived of it by desiccation. Their taste is mucilaginous and bitterish.

They contain, according to RIFFARD, a yellow fatty matter, a red colouring matter, gum, and vegetable fibre.

They are considered sedative and slightly narcotic, but are scarcely more than emollient. They are of service in slight pulmonary catarrhs, but more from the water in which they are infused than from any virtue they may themselves possess.

CONSERVE OF RED POPPY. (PALAT.; WIRTEM.; CADET DE GASSICOURT)

- R. Fresh Flowers of Red Poppy, one part.
 White Sugar, two parts.
 Mix. (PALAT.; WIRTEM.)

- R. Fresh Flowers of Red Poppy, one part.
 Sugar, four parts.
 Water, a sufficient quantity.

Mix. (CADET DE GASSICOURT)

DISTILLED WATER OF RED POPPY.

Aqua Papaveris erratici. (GALL.; PALAT.; SARD.)

- R. Flowers of Red Poppy, one part.
 Water, four parts.
 Macerate, and distil two parts. (GALL.; SARD.)

PALAT. prescribes one part of flowers and three parts of water; one and a half to be distilled.

EXTRACT OF RED POPPY. (BRUNS.; FERR.; SARD.; PIDERIT.; VAN MONS)

- R. Heads of Red Poppy, before the fall of the petals, one pound.

Spring Water, eight pounds.

Boil for an hour; express the juice, and evaporate it to the consistence of an extract. (PIDERIT)

BRUNS. prescribes three days' digestion in three pounds of hot water, and then a slight boiling.

R. Fresh Flowers of Red Poppy, any quantity.

Bruise them in a mortar, and macerate them for twelve hours in twice their weight of hot water; then strain through a hair-sieve, slightly expressing with a wooden spatula: clarify the liquor with white of egg, and evaporate.

R. Fresh Petals of Red Poppy, bruised, eight pounds.

Macerate for three days in an alembic; then distil off about two ounces of water: pour upon the residue thrice its weight of water, and, after six hours' digestion, boil down to one third. Then clarify, and strain the liquor, evaporate it to the consistence of an electuary, and when this is merely warm, add the distilled water. (SARD.)

It is recommended in troublesome cough, diarrhœa, pulmonary consumption, and in all cases where a slight sedative is indicated.—Dose, from ten grains to a scruple, to an adult.

INFUSION OF RED POPPY. (SARD.; RATIER)

R. Flowers of Red Poppy, two drams.
Boiling Water, two pounds.
Infuse.

It is emollient, and is considered slightly narcotic, under which title it is employed in nervous pains.—Dose, a glassful.

SYRUP OF RED POPPY.

Syrupus Papaveris erratici seu rubri seu Rhæados seu Rhæadis. (LOND.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; GENEV.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; WIRTEM.; COXE; SWEDIAUR; VAN MONS)

Syrupus Rhæados. (LOND.)

R. Fresh Petals of Red Poppy, one pound.

Boiling Water, one pint two fluid ounces.

Refined Sugar, one pound and a half.

To the water, heated in a water-bath, add gradually the petals, stir-

ring occasionally; then remove the vessel, and macerate for twelve hours: then having expressed the liquor, set it apart that the fæces may subside; then add the sugar and form a syrup.

Syrupus Papaveris Rhæados. (DUBL.; COXE)

R. Fresh Petals of Red Poppy, one pound.

Boiling Water, twenty fluid ounces.
Sugar, twenty-nine ounces.

Add the petals gradually to the boiling water; then remove the vessel from the fire, and macerate at a low heat for twelve hours: express the liquor and set it apart that the fæces may subside, and finally, add the sugar and form a syrup.

GALL. directs four pounds of petals and eight of boiling water, to be infused for twelve hours; the liquor to be slightly expressed, and set aside for a few hours, and a syrup to be formed in a water-bath, with the decanted liquor and twice its weight of white sugar.—AMST. directs one pound of flowers to be macerated for twelve hours in four pounds of boiling water, the liquor to be boiled down to one half, one pound of sugar to be dissolved in it, and the solution boiled to the proper consistence.—BATAV.; BELG.; HANN.; OLDEN.; POL.; and BORUS.—one pound of flowers, two pounds of boiling water, twelve hours' infusion, and three pounds of sugar.—DAN.—one pound and a half of flowers, three pounds of boiling water, twelve hours' infusion, the reduction of the strained liquor to one half by boiling, and the addition of three pounds of sugar to form a syrup.—HISP.—one pound of flowers, two pounds of water, eight hours' infusion, and four pounds of sugar.—PALAT.—one pound and a half of flowers, three pounds of boiling water, twenty-four hours' infusion, and three pounds of sugar.—BRUNS.; and WIRTEM.—two pounds of flowers, three of water, twenty-four hours' infusion, and five pounds four ounces of sugar.—GENEV.—three ounces of flowers, three pounds of boiling water, a decoction followed by a maceration of twelve hours, and six pounds of sugar.—LIPP.—one pound of dried petals, five pounds of boiling water, twelve hours' digestion, the reduction of the strained liquor to one half by boiling, and the addition of three pounds of sugar.

ANT.; SWEDIAUR; and VAN MONS recommend the petals to be bruised in a mortar.—ANT.; and VAN MONS prescribe one part of flowers and one and a half of boiling water.—SWEDIAUR requires one part of flowers and three parts of water:—the latter requires twenty-four hours' infusion, but the others only twelve.—VAN MONS prescribes two parts of sugar to three of strained liquor.—ANT.—eight ounces of sugar to be dissolved in the strained liquor; and SWEDIAUR—two parts of sugar to one of strained liquor.

PECTORAL INFUSION. (SAINTE-MARIE)

- R. Flowers of Red Poppy, two ounces.
Dilute Sulphuric Acid, fifteen drops.
White Sugar, two ounces.
Decoction of Barley, one pound.

Infuse.—To be taken in the course of twenty-four hours, two or three spoonfuls at a time, particularly on the coming on of the cough.

PASTILLES OF RED POPPY. (FERR.)

- R. Infusion of Red Poppy, six pounds.
Sugar, four pounds.

Dissolve, clarify, boil to a proper consistence, pour the mass upon a marble slab, and cut it into proper forms.

VINEGAR OF RED POPPY.

Acetum Papaveris erratici. (WIRTEM.)

- R. Flowers of Red Poppy, half a pound.
Vinegar, three pounds.

Macerate in the sun for several days, and express.

AQUEOUS TINCTURE OF RED POPPY.

Tinctura Papaveris Rhæados. (PALAT.)

- R. Conserve of Red Poppy, one ounce.
Flowers of Red Poppy, sprinkled
with a sufficient quantity of sulphuric acid, three handfuls.
Distilled Water of the Juice of Red Poppy, thirty ounces.

Digest for twenty-four hours; express strongly and add,

- Syrup of Red Poppy, slightly acidulated, eight ounces.

Mix.—It is slightly excitant, and used formerly to be much prescribed in malignant fevers, in the dose of a spoonful, in a mixture, or common drink.

2. *Papaver somniferum*, LINN.—Poppy, White Poppy.

SYNONIMES:—Pavot, Pavot blanc, FR.; Mohn, GERM.; Khuschhasch, ARAB.;

and DUK.; Bilymak, BOHEM.; Ying-suh, CHIN.; Gaysagussa, CAN.; Abin atta, CYNG.; Valmue, DAN.; Adormidera, SPAN.; Post, HIND.; and SANS.; Tamme heul, DUT.; Papavero, ITAL.; Kes, JAP.; Kuknar, PERS.; Mak bialy, POL.; Wallmo, SWED.; Casa casa, TAM.; Cassa cassa, TEL.

LOND.; EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

An original plant of the East, but naturalized in the south of Europe. (*Polyandria Polygynia*, LINN.; *Papaveraceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 371*)

The fruit and seeds are employed.

The fruit (*Poppy Heads*; *Capita seu Capitula seu Capsula Papaveris*) are globular or ovoid capsules, with a flat crown of ten radii or more, under which there are as many pores as there are radii. Its odour, which is virous in the fresh state, becomes dissipated by desiccation. It taste is bitter and acrid.

These capsules are sedative and narcotic. They furnish opium in the East, and might also yield it in our countries; for they contain all the elements of Eastern opium, but in proportions which seem to vary considerably, and upon which the chemists who have latterly devoted themselves to this important enquiry have not yet agreed.

The seeds (*Semen Papaveris albæ seu nigri seu cœrulei*) are very small, reniform, rather wrinkled, of a greyish white or blackish. The oil (*Huile d'aillette*, FR.) which is expressed from them, is bland, and good for the table.

AQUEOUS TINCTURE OF POPPY FLOWERS. (BRUNS.; SAX.; WIRTEM.)

- R. Flowers of White Poppy, half an ounce.
Sulphuric Acid, one dram.
Water of Poppy Flowers, four ounces.

Digest for twenty-four hours; express, and filter. It is sedative.—Dose, from half an ounce to one ounce. (BRUNS.; WIRTEM.)

SAX. prescribes one ounce and a half of flowers, half an ounce of acid, and twelve ounces of boiling water; to be macerated for twelve hours.

POPPY OIL.

Oleum Papaveris albi seu somniferi.
(AUSTR.; BAVAR.; BRUNS.; HISP.;
GALL.; HANN.; PALAT.; POL.; BO-
RUS.; SARD.; SAX.; WIRTEM.; HERBIP.)

R. Poppy Seeds, any quantity.

Bruse them in a marble mortar with a wooden pestle; enclose them in a linen bag, and express the oil without heat.

BORUS. directs the press to be heated with boiling water.—BRUNS.; PALAT.; and WIRTEM.—the seeds to be slightly torrifried before expressing them.

This oil is reputed anodyne and sudorific.

PECTORAL POWDER. (PIERQUIN)

R. Seeds of White Poppy, five drams.
Gum Arabic, } of each, one
— Tragacanth, } dram and a half.
Starch, }
Seeds of Purslain, }
— Marshmallows, } of each,
— Cotton, } two drams
— Wild Mallow, } and a half.
Four Cold Seeds,
Quince Rind, }
Prepared Tutty, } of each, one
Amber, } dram and a half.
Liquorice Powder, }
Sugar, }

Mix.—Dose, two drams, in syrup of poppy or of jujubes.

INCRASSANT ELECTUARY. (BORIES)

R. Conserve of Roses, one ounce.
Syrup of Violets, two ounces.
Seeds of White Poppy, three drams.
Oil of Nutmeg, six grains.

Mix.—For obstinate cough.—Dose, one tea spoonful every two hours.

SEDATIVE EMULSION. (PIERQUIN)

R. Four Cold Seeds, one ounce.
Poppy Seeds, two ounces.
Triturate, adding gradually,
Water of Lily, } of each,
— Rose, } three ounces.
Strain, and add
Syrup of Lemon, } of each,
— Water Lily, } one ounce.

Mix.

ANODYNE SPECIES. (FULD.; PALAT.; HERBIP.; SWEDIAUR)

R. Poppy Heads, one pound.
Root of Marshmallows, } of each,
Linsced, } half a pound.

Cut and bruise. (FULD.)

For lavement.

VOL. II.

R. Poppy Heads, six ounces.
Chamomile Flowers, } of each,
Leaves of Lesser } three ounces.
Mullein, }

Cut and mix. (FULD.)

For fomentation.

R. Poppy Heads, two drams.
Leaves of Peppermint, } of each,
Chamomile Flowers, } half a handful.

Cut and mix. (SWEDIAUR)

R. Leaves of Henbane, } of each,
Flowers of Elder, } one ounce.
Saffron, } two drams.
Poppy Heads, } four ounces.

Cut and mix. (HERBIP.)

R. Poppy Heads, bruised with } of each,
the seeds, } two
Seeds of Dill, } ounces.
Herb of Henbane, } of each,
— Houndstongue, } two
— Solanum nigrum, } handfuls.
Leaves of Chamomile, }

Cut and bruise. (PALAT.)

For fomentation.

SPECIES AGAINST PLEURISY.

Species pro Fotu laterali. (PALAT.)

R. Root of Marshmallows, } of each,
Poppy Heads, bruised with } two
the seeds, } ounces.
Linseed, } of each,
Seeds of Fenugreek, } one ounce.
— Cumin, }
Bay Berries, } of each,
Herb of Wild Mallow, } three handfuls.
— Motherwort, }
Chamomile Flowers, } two handfuls.

Mix.

EXTRACT OF POPPY.

Extractum Papaveris albi. (LOND.; EDINB.;
BRUNS.; HANN.; COXE; VAN MONS)

Extractum Papaveris. (LOND.)

Ext. Papaveris somniferi. (EDINB.)

R. Poppy Heads, freed from
seeds, and bruised, one pound.
Boiling Water, one gallon.

Macerate for twenty-four hours, and boil down to one half; then strain the hot liquor, and evaporate it to the proper consistence.

It is sedative and narcotic.—Dose, one or two grains, once or twice a day.

2 X

DECOCTION OF POPPY.

Decoctum Papaveris. (LOND.; DUBL.; NIEMANN)

R. Poppy Heads, bruised, four ounces.
Water, four pints.

Boil for a quarter of an hour, and strain.

It is employed in fomentations, particularly to painful abscesses and excoriations.—With regard to this, NIEMANN make the a remark which every practitioner should always bear in mind, that narcotic substances when applied to surfaces denuded of epidermis, very frequently exasperate the pain which they are applied to relieve.

Under the titles of *Anodyne Lavement*, *Poppy Lavement*, and *Sedative Lavement*, RATIER directs decoctions to be made with half an ounce of poppy heads and one pound of water; with three drams of poppy heads and one pound of water; and with one poppy head and two pounds of emollient decoction. He also prescribes a *Sedative Fomentation*, prepared by boiling two poppy heads in two pounds each, of water and of emollient decoction. Finally, we find in SWEDIAUR a *Sedative Lotion*, which is a decoction of half a pound of poppy heads in two pounds of water boiled down to one pound; to which, the author says, twenty grains of deuto-chloruret of mercury, or from one to three drams of sub-acetate of lead is sometimes added.

ANODYNE FOMENTATION. (PALAT.; SAUNDERS)

R. Anodyne Species (form. of PALAT.)
[Quantity not given, EDIT.]

Water, six pounds.
Boil down to four pounds, and strain. (PALAT.)

R. Poppy Heads, two ounces.
Spring Water, two pounds.
Boil down to one pound; strain, and add
Vinegar, one ounce.
Mix. (RATIER)

ANTI-PLEURITIC FOMENTATION. (PALAT.)

R. Species against Pleurisy (form of PALAT.) [Quantity not given, EDIT.]
Cow's Milk or Water, eight pounds.
Boil down to one half, and strain.

It is emollient, resolvent, and anodyne.—It is recommended in pleurisy when the inflammation is not acute.

SEDATIVE GARGLE. (RATIER)

R. Poppy Heads, two.
Linseed, one dram.
Boiling Water, six ounces.

Infuse, strain, and add
Syrup of Honey, two drams.
Mix.

EMOLLIENT INJECTION. (CADET DE GASSICOURT)

R. Decoction of Poppy, one pound.
Emulsion of Almonds, five ounces.
Fresh White of Egg, one dram.

Mix.—It is recommended in acute hemorrhagia.

DEMULCENT POTION. (SAINTE-MARIE)

R. Linseed, two drams.
Poppy Heads, one.
Water, a sufficient quantity
to obtain eight ounces of strained liquor
after a quarter of an hour's ebullition; add
Fresh Yolk of Egg, one.

Mix.—To be taken by spoonfuls, in twenty-four hours; in painful diarrhœa.

PECTORAL BROTH. (SAINTE-MARIE)

R. Flayed Chicken, one half.
Turnip, sliced, one.
Water, one pound and a half.

Boil down to three-fourths, adding towards the end

Poppy Heads,
two scruples or one dram.

Strain, and add
Sugar Candy, four ounces.

Mix.—Dose, two glassfuls, morning and evening.

ANTI-NEPHRITIC LIQUOR.

Liquor ad Nephritidem calculosam. (CADET DE GASSICOURT; SWEDIAUR)

R. Poppy Heads,
from four to eight ounces.
Boiling Water, two pounds.
Boil down to eight ounces, express, and add
Nitrate of Potass, one ounce.
Mix.

ANODYNE LOTION. (SAINTE-MARIE)

R. Poppy Heads, one ounce.
Water, three pounds.

Boil down to one half, and add to the cold strained liquor

Solution of Ammonia,
one or two ounces.

Mix.

SYRUP OF POPPY.

Syrupus Papaveris albi seu *Papaveris somniferi* seu *de Meconio* seu *diacodion* seu *de Capitis Papaveris* seu *Diacodii* seu *Diacodium liquidum Montani*. (LOND.; EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; SARD.; WIRTEM.; HERBIP.; COXE.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

Syrupus Papaveris. (LOND.)

R. Dried Poppy Heads, freed
from seeds and bruised,
fourteen ounces.

Refined Sugar, two pounds.
Boiling Water,
two gallons and a half.

Macerate the capsules in the water for twelve hours; then boil down to one gallon, in a water-bath, and express strongly: boil the strained liquor down to two pounds, and strain again while hot. Having set it apart for twelve hours that the fæces may subside, boil the clear liquor till it is reduced to one pint, and finally add the sugar and form a syrup.

Syrupus Papaveris Somniferi.
(EDINB.; COXE.)

R. Dried Poppy Heads, freed
from seeds, one part.
Boiling Water, fifteen parts.
Refined Sugar, two parts.

Macerate the sliced capsules in the water for twelve hours; then boil down to one third, and strain with strong expression. Boil the strained decoction to one half, and strain again; then add the sugar, and form a syrup.

R. Poppy Heads, gathered before
they are ripe, dried and freed
from seeds, seventeen ounces.
Boiling Water, two gallons.
Sugar, fifty-eight ounces.

Slice and bruise the capsules, pour the water over them, and macerate for twenty-four hours; then express the liquor, and evaporate it slowly to two pints: set the liquor apart that the fæces may subside, and finally add the sugar to the clear liquor, and form a syrup. (DUBL.)

R. Poppy Heads, freed from seeds, and
bruised, three parts.
Hot Water, fifteen parts.

Infuse for twelve hours at a temperature of from 30 to 40 degrees; then express strongly, and to one part of the strained liquor add

White Sugar, two parts.

Dissolve. (BAVAR.)

LIPP. directs one pound of capsules to be infused in five pounds of boiling water for twelve hours; and five pounds of sugar to be added to every two pounds of strained liquor.

R. Poppy Heads with the seeds,
eight ounces.
Spring Water, four pounds.

Boil down to one half; add to the strained liquor, depurated by repose,

White Sugar, three pounds.

Make a syrup. (BRUNS.; PALAT.; WIRTEM.)

SPIELMANN prescribes four pounds of sugar.—GENEV.—one pound of capsules, sixteen pounds of water, and four pounds of sugar.—FERR.—one pound of capsules, eight pounds of water, and four of sugar.

This formula presents numerous varieties:—GALL. directs one pound of capsules freed from seeds to be digested for twelve hours in eight pounds of boiling water; the liquor to be then evaporated in a water-bath to one half, set aside to depurate, and a syrup formed with four pounds of sugar.—BELG.—one pound of nearly dry capsules, freed from seeds, to be digested for twenty-four hours in thirty pounds of boiling water, and then boiled till only eight pounds of liquor remain: this is to be expressed, evaporated to one half, depurated, and a syrup then formed with four pounds of sugar.—HASS.; and PIDERIT.—one pound of capsules to be macerated for twenty-four hours in eight pounds of water; to be then boiled, and sixteen ounces of sugar added to every nine of the strained liquor, to make a syrup by boiling for a few instants.—HERBIP.—eight ounces of capsules, twelve pounds of water, and twelve hours' maceration; then reduction to one third, straining, evaporation to one half, and the addition of two pounds and a half of sugar.—VAN MONS—to macerate one

part of capsules in four parts of water for six hours; then to add eight of water, to boil down to six, and finally to make a syrup with the addition of six parts of sugar.

—SWEDIAUR—to macerate two pounds of capsules in thirty pounds of water for twelve hours; then to reduce it to one third, and having strained with expression, to evaporate the liquor to one half; to strain again, and make a syrup with four pounds of sugar.

—AMST.—to macerate one pound of capsules in twenty-four pounds of boiling water for twelve hours, then to boil down to one fourth, to strain with expression, to evaporate to one half, to strain again, to dehydrate by repose, and to make a syrup with three pounds of sugar.—HISP.—to macerate sixteen ounces of capsules in twenty pounds of warm water for twelve hours, to boil down to two thirds, to strain, to add four pounds of sugar with a sufficient quantity of water, then to clarify with white of egg, and to boil to a syrup.—BATAV.—to infuse one pound of capsules for twenty-four hours in thirty pounds of water, then to reduce to eight, to strain and boil down to one half; then to set apart, decant, and make the syrup with three pounds of sugar.

—LUSIT.—to macerate for twelve hours, two pounds of capsules in twenty-four of water; then to boil down to one third, strain, and evaporate to one half, and to make the syrup with four pounds of sugar.

—SARD.—to infuse for twenty-four hours, one pound of capsules in eight pounds of water; then to boil, express, and make the syrup by adding sixteen ounces of sugar to every nine of the strained liquor.—GENEV.

—to digest for twenty-four hours, two pounds of capsules in eight of water, to boil down to one half, and to make the syrup with four pounds of sugar.—FULD.—to macerate one pound of capsules in four pounds of water; then to boil down to one half, and add two pounds of sugar boiled *à la plume*.

R. Poppy Heads, without seeds and bruised, two pounds.
Rain Water, twenty pounds.

Macerate for twenty-four hours, gradually raising the temperature to boiling, and strain; boil the residue for a quarter of an hour with

Water, ten pounds.

Express strongly.—Mix the decoctions, evaporate to one half, and add

White Sugar, eight pounds.

Clarify, and boil to the proper consistence. (ANT.)

R. Dried Poppy Heads, fourteen ounces.
Root of Liquorice, half an ounce.
Boiling Water, eight pounds.

Macerate for one night; then boil down to one half, express strongly, and add

White Sugar, four pounds.

Boil to a syrup. (DAN.)

OLDEN. prescribes sixteen ounces of capsules, one ounce of liquorice, eight pounds of water, twelve hours' maceration, reduction to one half, and the addition of four pounds of sugar.

R. Poppy Heads, sliced, and freed from seeds, four ounces.
Water, eight pounds.

Boil down to three pounds, adding towards the end

Grated Liquorice Root, two ounces.

Strain with strong expression, and add

White Sugar, one pound and a half.

Clarify, and boil to the proper consistence. (AUSTR.)

R. Unripe Poppy Heads, with the seeds, two ounces.

Carob Beans, two ounces and a half.

Spring Water, one pound and a half.

Boil over a slow fire down to ten ounces, and add towards the end

Fresh Liquorice, sliced, one ounce and a half.

Dissolve in the strained liquor

White Sugar, sixteen ounces.

Skim, and strain. (HANN.; PALAT.; WIRTEM.; SPIELMANN)

CROCATED SYRUP OF POPPY.

Syrupus diacodium crocatum. (WIRTEM.)

R. Syrup of Poppy, one ounce.
Powdered Saffron, fifteen grains.
Mix.

ANISATED SYRUP OF POPPY. (VAN MONS)

R. Oil of Anise, one dram.
Mucilage of Gum Arabic, one dram and a half.
Syrup of Poppy, sixty-four ounces.
Mix.

PECTORAL SYRUP. (CADET DE GASSI-COURT)

R. Dates, two pounds.
Jujubes, one pound.
Root of Liquorice, half a pound.
—— Marshmallows, } of each,
Canadian Maidenhair, } four ounces.
Poppy Heads, }
Water, sixteen pounds.

Evaporate to the consistence of syrup.

SEDATIVE POTION. (GALL.; RATIER;
PIERQUIN)

R. Gum Potion, four ounces.
Syrup of Poppy, half an ounce.
Mix. (RATIER)

R. Cichory Water, three or four ounces.
Syrup of Poppy, four or six drams.
Mix. (PIERQUIN)

R. Infusion of Lime Flowers, } of each,
Pectoral Infusion, } two ounces.
Orange-flower Water, two drams.
Syrup of Poppy, half an ounce.
Mix. (RATIER)

R. Orange-flower Water, half an ounce.
Lettuce Water, three ounces.
Syrup of Poppy, two drams.
Mix. (GALL.)

R. Orange-flower Water, } of each, one
Lily Water, } ounce and a half.
Syrup of Poppy, } of each,
Borage, } half an ounce.

Mix. (PIERQUIN)

R. Orange-flower Water, one ounce.
Balm Water, three ounces.
Simple Syrup, } of each, one ounce,
Syrup of Poppy, }

Mix. (RATIER)

R. Water of Red Poppy, } of each,
Coltsfoot, } two ounces.
Water Lily, }
Prepared Crab's Eyes, twenty grains.

Syrup of Poppy, six drams.
Mix. (PIERQUIN)

R. Decoction of Lettuce, } of each,
Purslain, } three ounces.
Syrup of Poppy, two ounces.
Mix. (PIERQUIN)

DEMULCENT GARGLE. (RATIER)

R. Decoction of Starch, eight ounces.
Syrup of Poppy, one ounce.
Mix.

PECTORAL MIXTURE. (SAUNDERS)

R. Syrup of Poppy, two ounces.
Oil of Almonds, fourteen ounces.
Mix.

PECTORAL EMULSION. (SAUNDERS;
SAINTE-MARIE)

R. Melon Seeds, three ounces.
Barley Water, a sufficient quantity
to make one pound of emulsion. Add when
strained,

Syrup of Poppy, one ounce.

Mix. (SAUNDERS)

R. Olive Oil, one ounce.
Fresh Yolk of Egg, one.
Syrup of Poppy, six drams.
Water, six ounces.

Make an emulsion. (SAINTE-MARIE)

PARIETARIA OFFICINALIS, LINN.—PELLITORY OF THE WALL,
WALL PELLITORY.

SYNONIMES :—Parietaire, FR.; Nachtkraut, Glaskraut, GERM.; Den a noc, Sklenna bylina, BOHEM.; Murkurt, DAN.; Parietaria, ITAL.; PORT.; and SPAN.; Glaskruid, DUT.; Noc i dzien, POL.; Væggært, SWED.

AMST.; ANT.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant common throughout Europe. (*Polygama Monecia*, LINN.; *Urticea*, JUSS.; *Fig. Zorn, Ic. pl. t. 121*)

The herb (*Herba Parietariæ seu Helxines*) is employed. It consists of a round reddish stalk, and of oval, lanceolated, petiolated leaves, glabrous above and slightly hairy beneath.

It is feebly diuretic.

DECOCTION OF WALL PELLITORY. (RATIER)

R. Leaves of Wall Pellitory, one ounce.
Water, two pounds.

Boil.—It is usually employed as a vehicle for more active diuretics.

DISTILLED WATER OF WALL PELLITORY. (GALL.)

R. Fresh Herb of Wall Pellitory, five thousand parts.

Water, twelve thousand five hundred parts.

Distil ten thousand parts, and pour the product upon

Fresh Herb of Wall Pellitory, five thousand parts.

Add,

Water, ten thousand parts.

Distil ten thousand parts.—The same operation may be repeated a third time.

PARIS QUADRIFOLIA, LINN.—ONE-BERRY.

SYNONIMES :—Parisette, Raisin de Renard, FR.; Einbeere, Woolfsbeere, Fuchstraube, GERM.; Etbær, DAN.; Ubas de zorro, SPAN.; Woolfsbezie, DUT.; Uva di volpo, ITAL.; Wronie oko, POL.; Parisetta, PORT.; Trollbær, SWED.

GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of Europe. (*Octandria Tetragynia*, LINN.; *Asparagineæ*, JUSS.; Fig. Zorn, *Ic. pl. t. 10*)

The root, herb, and fruit are employed.

The root (*Radix Paradis seu Solani quadrifolii*) is fleshy, rather nodose, and furnished with a few fibres. It has a nauseous smell and taste.

It excites vomiting in the dose of from thirty to fifty grains. The ancients prescribed it in colic.

The herb consists of a long, slender, glabrous stem, naked on the lower part and furnished above with four oval, acute, verticillated leaves. Its odour is very disagreeable, and its taste sweetish.

It is purgative. It is also considered anodyne, sudorific, and alexipharmic. It has been principally employed in mania, spasms, hooping cough, poisoning from nuxvomica, &c.

The fruits (*Uvæ versæ seu vulpinæ*) are berries of a deep-purple or blackish-brown colour, of the size of cherries, and shining. They have a disagreeable odour and a virous taste. They are poisonous.

PARNASSIA PALUSTRIS, LINN.

SYNONIMES :—Parnassie des marais, FR.; Wasserleberblume, Sumpfparnassien gras, Sumpferzblume, Herzblümlein, GERM.

BRUNS.; BERGIUS; SPIELMANN.

A plant common in all parts of Europe. (*Pentandria Tetragynia*, LINN.; *Cappari-deæ*, Fig. Zorn, *Ic. pl. t. 124*)

The herb and flowers are employed.

The herb (*Herba Hepaticæ albæ*) consists of radical heart-shaped, very entire leaves, supported upon long foot-stalks.

The flowers are white and rather large.

The taste is rather bitter and acrid.

It is slightly astringent.

PARONYCHIA ARGENTEA, LMK.

SYNONIMES :—Paronique argentée, FR.; Nevadilla, SPAN.

HISP.

A plant of the south of France. (*Pentandria Monogynia*, LINN.; *Paronychicæ*, A. ST. HILL)

The flowering herb (*Herba Sanguinariæ minoris*) is employed. It consists of a branching stem bearing oval, acuminate, nearly glabrous leaves, with stipulæ shorter than they, broader at the base; the flowers are arranged in small terminal heads surrounded by large bractæ.

PASPALUM DACTYLUM, LMK.—CREEPING PANIC GRASS.

SYNONIMES :—Paspale, Pied de poule, FR.; Fingerfärmiges Hirsegras, GERM.; Grama, SPAN.; Gevingerd panik, DUT.; Grama digitate, PORT.

HISP.; GALL.; FERR.; GENEV.

A common plant of Europe. (*Triandria Digynia*, LINN.; *Gramineæ*, JUSS.; Fig. Schrad, *Fl. Germ. I. p. 165, t. 3. fig. 9*)

The radical offsets which are commonly employed in France under the name of *Chien dent* and erroneously looked upon as roots, are very long, of the diameter of a crow-

quill, cylindrical and knotty. Under a hard, yellow, shining epidermis it contains a white substance of a farinous and saccharine taste.

PASTINACA SATIVA, LINN.—GARDEN PARSNIP.

SYNONIMES :—Panais, FR. ; Pastinake, GERM. ; Passtrnak, BOHEM. ; Pastinak, DAN. ; Pastinaca, ITAL. ; PORT. ; and SPAN. ; Pinsternakel, DUT. ; Pasternak, POL. ; Palsternacka, SWED.

ANT. ; BATAV. ; GALL. ; FERR. ; SUEC. ; WIRTEM. ; BERGIUS ; BRUGNATELLI ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

This plant is cultivated in all parts of Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; *Fig. Zorn, Ic. pl. t. 568*)

The root and seeds are employed.

The root (*Radix Pastinacæ*) is an inch or more in diameter, fleshy, whitish or yellowish. It has a rather aromatic and saccharine taste.

The seeds are rounded, elliptical, nearly plane, surrounded with a narrow membranous border, striated, and of a rather aromatic and sweet taste.

It is excitant, and was formerly esteemed diuretic.

PEUCEDANUM.—SULPHURWORT.

Two species of this genus of plants are employed in medicine :

1. *Peucedanum Silaus*, LINN.—Meadow Sulphurwort.

SYNONIMES :—Seseli des pres, Saxifrage des Anglais, FR. ; Silau, Mattensteinbrech, GERM. ; Weld minum, DAN. ; Hairstreng, DUT.

GALL.

A plant of the south of Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; *Fig. Jacq. Fl. Austr. t. 15*)

The root and seeds are employed.

The root (*Radix Seselios pratensis* seu *Saxifragæ Anglicæ*) is large, thick, often bifurcated, provided below with a bundle of hairs.

The seeds are flat on one side, and bulged on the other, and marked with five striæ. Their odour and taste are aromatic and acrid.

2. *Peucedanum officinale*, LINN.—Common Sulphurwort.

SYNONIMES :—Peucedan, Fenouil de porc, Queue de porceau, FR. ; Saufenchel, Schwefelwurzelhaarstrang, GERM. ; Svinkenkel, DAN. ; Peucedan, ITAL. ; PORT. ;

and SPAN. ; Hairstreng, DUT. ; Weerpznicce, POL. ; Svinfinkel, SWED.

BRUNS. ; HISP. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of the north of Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; *Fig. Gmel. Fl. Sibir. I. t. 41*)

The root (*Radix Peucedani* seu *Feniculi porcini*) is employed. It is thick, long, round, fibrous, blackish without and whitish within. It has a viroous odour, and an acrid, bitter, and mucilaginous taste.

It is excitant, and is reputed anti-hysterical.

EXTRACT OF COMMON SULPHURWORT.
(SARD.)

R. Root of Sulphurwort, one pound.
White Wine, five or six pounds.

Digest for three days in a sand-bath ; then boil the expressed residue in a sufficient quantity of fresh wine. Mix the liquors and evaporate to the consistence of an extract.

SYRUP OF COMMON SULPHURWORT.
(SARD.)

R. Extract of Sulphurwort, } of each,
——— Liquorice, } one ounce.

Extract of Veronica, } of each,
 ————— Coltsfoot, } three drams.
 Honey, two ounces.
 Distilled Water, a sufficient quantity.

Mix over a slow fire, and make a syrup.

ESSENTIAL OIL OF COMMON SULPHUR-
 WORT. (SARD.)

R. Root of Sulphurwort, twenty-five parts.
 Water, seventy-five parts.
 Muriate of Soda, three parts.
 Distil and collect the supernatant oil.

PERSICA VULGARIS, CAND.—PEACH TREE.

SYNONIMES:—Pêcher, FR.; Pfirschenbaum, GERM.; Khowth, ARAB.; Breskew, BOHEM.; Persikketræ, DAN.; Prisco, SPAN.; Persikboom, DUT.; Persico, ITAL.; Schuftalu, PERS.; Broskwina drzewo, POL.; Pecegueira, PORT.; Persiketræd, SWED.

AUSTR.; ANT.; BELG.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; WIRTEM.; HERBIP.; AINSLIE; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A tree originally of Persia. (*Icosandria Monogynia*, LINN.; *Rosaceæ*, JUSS.; *Fig.* Zorn, *lc. pl. t.* 282)

The leaves, flowers, and seeds, are employed.

The leaves (*Folia Persicarum*) are simple, long, lanceolated, pointed, and finely indented on the borders. They have the taste and odour of bitter almonds. They are feebly laxative.

The flowers are reddish or rose-coloured, and possess the same property as the leaves.

The seed is a ligneous nut, very hard, and grooved; it encloses a bitter kernel containing a bland oil and hydrocyanic acid.

CONSERVE OF PEACH FLOWERS.

Conserva Florum Persicarum. (SARD.)

R. Fresh Peach Flowers, one part.
 Sugar, boiled *à la plume*, two parts.
 Mix.

JUICE OF PEACH FLOWERS.

Succus ex Floribus Persicarum. (SARD.)

R. Fresh Peach Flowers, freed from the calices, any quantity.

Sprinkle them with a little water, put them into a linen bag, and express and filter the juice. If it is intended to be preserved, it should be set apart to depurate, and after having added a little alcohol to the limped part, be kept in bottles under a stratum of oil.

INFUSION OF PEACH FLOWERS.

Infusum Persicarum. (HISP.; SARD.)

R. Peach Flowers, two pounds.
 Boiling Water, eight pounds.

Infuse for eight hours, and strain with slight expression; repeat the operation nine times successively with fresh flowers. (SARD.)

R. Peach Flowers, one pound.
 Boiling Water, two pounds.

Infuse for eight hours, and strain with expression. (HISP.)

DECOCTION OF PEACH LEAVES. (AUGUSTIN; SWEDIAUR)

R. Peach Leaves, one ounce.
 Water, two pounds.

Boil down to one pound and a half.

It is recommended in hæmaturia, dysuria and nephritis. Dose, one pound daily.

SYRUP OF PEACH LEAVES. (VAN MONS)

R. Fresh Peach Leaves, any quantity.

Bruise them and express the juice; coagulate it by boiling, and dissolve in five parts of the strained liquor.

White Sugar, nine parts.

SYRUP OF PEACH FLOWERS.

Syrupus Persicarum Florum seu de Amygdali Persicæ Floribus seu Amygdali Persicæ. (ANT.; BRUNS.; GALL.; FERR.; GENEV.; PALAT.; SARD.; WIRTEM.; HERBIP.; VAN MONS)

R. Fresh Peach Flowers, four pounds.
 Boiling Water, twelve pounds.

Infuse for twelve hours in a covered vessel placed in warm ashes, and strain with slight expression; set apart, decant, and add to the liquor.

White Sugar, seventeen pounds.

Boil to the consistence of syrup. (GALL.)

BRUNS.; PALAT.; and WIRTEM. direct one pound of flowers to be macerated for

twelve hours in one pound of boiling water; then to express, and to make a syrup by adding sixteen ounces of sugar to nine ounces of depurated juice.—FERR.—to macerate two pounds of flowers in three of boiling water for twelve hours; and to add two pounds of sugar to the strained liquor.—SARD.—to boil to a proper consistence a mixture of an equal part each, of infusion of peach flowers and of simple syrup.

R. Fresh Peach Flowers, two parts.

Bruise and pour upon them,

Hot Water, four parts,

in a covered vessel. When cold, strain with expression, and dissolve in four parts of the liquor,

White Sugar, seven parts.

Make a syrup. (VAN MONS)

R. Peach Flowers, eight ounces.

— Buds, two ounces.

Boiling Water, sixteen ounces.

Infuse in a covered vessel placed in a water-bath for twenty-four hours; then filter, and dissolve in five ounces of the infusion,

White Sugar, eight ounces.

Strain. (ANT.)

R. White Sugar, four pounds.

Spring Water, a sufficient quantity.

Boil to the consistence of thick syrup; pour this while hot, upon

Fresh Peach Flowers, two pounds.

Macerate for three days in a covered vessel, then boil a little and express strongly. (GENEV.)

SEDATIVE EMULSION.

Emulsio externa e Nucleis. (WIRTEM.)

R. Peach Kernels, one ounce.

Seeds of Hemp, } of each,

— Poppy, } half an ounce.

Water of Melilot, } of each,

— Water Lily, } four ounces.

Add to the emulsion,

Spirit of Camphor, half an ounce.

Mix.—It is applied, by means of compresses, to the forehead and temples, in cephalalgia.

PETROLEUM.

SYNONIMES :—*Oleum Petreæ*; *Bitumen Petroleum* (DUBL.; EDINB.): Barbadoes or Mineral Tar.—*Pétrole*, FR.; *Steinöl*, BERGEL, GERM.; *Neft*, ARAB.; *Muttie katali*, DUK.; *Petroleo*, SPAN.; *Mitti tel*, HIND.; *Petrolino*, ITAL.; *Keesa no abra*, JAP.; *Minnia tanna*, MAL.; *Oley skalny*, POL.; *Bluemi tailam*, SANS.; *Bergolja*, SWED.; *Ippu*, SUMAT.; *Mun tylum*, TAM.

LOND.; EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BRUGNATELLI.; COXE.; PIDERIT.; SPIELMANN.

A fluid bitumen, frequently of an oily consistence, of a blackish and sometimes reddish-brown colour, nearly opaque, lighter than water, unctuous to the touch, of a very strong smell and acrid taste. It is very combustible and burns with a thick black smoke.

It is stimulant, anti-spasmodic, diuretic, and anthelmintic.—Dose, from ten to fifteen drops, on sugar.

Vol. II.

ESSENTIAL OIL OF PETROLEUM.

Oleum volatile seu Naphtha Petrolei. (HERBIP.; SWEDIAUR)

R. Petroleum, two pounds.

Water, four pounds.

Distil over a slow fire, till limpid oil ceases to pass over.

It is stimulant and resolvent, and is principally employed externally, in arthrodynia, ecchymosis, and paralysis.

LINIMENT OF PETROLEUM.

R. Petroleum, } of each, one ounce.

Rectified Spirit, }

Mix. (SAUNDERS; SWEDIAUR)

R. Essence of Soap, eight drams.

Petroleum, one dram and a half.

Mix. (SAUNDERS)

ANTHELMINTIC MIXTURE. (AUGUSTIN; NIEMANN)

R. Petroleum, half an ounce.

Tincture of Assafoetida, six drams.

Mix.—Dose, forty drops, thrice a day.—It is recommended by SCHWARZ, against tænia.

2 Y

PHASEOLUS COMMUNIS.—FRENCH BEAN.

SYNONIMES:—Haricot commun, FR.;
Gemeine Schminkbohne, GERM.

GALL.; BERGIUS.

This is an original plant of the East Indies, but it is cultivated in every part of Europe, where it has produced numerous varieties. (*Diadelphia Decandria*, LINN.;

Leguminosæ, JUSS.; *Fig. Schk. Bot. Handb.* t. 199)

The seed (*Semen Phaseoli seu Fabarum*) is employed. It is more or less reniform, of a white, yellowish, red, violet, black or marbled colour, and farinaceous. The white beans (*Haricots blancs*) are those which are ordinarily used; they are smooth, shining, flattish, and rather long.

PHELLANDRIUM AQUATICUM, LINN.—WATER HEMLOCK.

SYNONIMES:—Phellandre aquatique, FR.;
Wasserfenchel, Peersaat, Pferdesaamen,
GERM.; Nandfennikel, DAN.; Water-
kervel, DUT.; Felandro, ITAL.; Koper-
wadny, Kruszykamien ziele, POL.;
Stækra, SWED.

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.;
BRUNS.; DAN.; GALL.; FERR.; HANN.;
HASS.; OLDEN.; POL.; BORUS.; ROSS.;
SAX.; SUEC.; WIRTEM.; HERBIP.;
WURCEB.; BERGIUS.; GUIBOURT.; MUR-
RAY and GMELIN.; SPIELMANN.

A plant common all over Europe. (*Pentandria Digynia*, LINN.; *Umbellifera*, JUSS.; *Fig. Cæd. Fl. Dan.* t. 1154)

The herb and seeds are employed.

The herb (*Herba Phellandrii seu Feniculi aquatici*) consists of a cylindrical, fistulous, striated, branched stem, and of trebly or quadruply-winged, glabrous leaves, with leaflets deeply incised by narrow linear notches.

The seed is oval, long, even on one side, and striated on the other, surmounted by the remains of the calyx and by the styles; they are of a brownish or yellowish-green colour.

The odour is disagreeable, strong, and penetrating; the taste aromatic and bitterish. It is narcotic, stimulant, and diuretic.—Dose, from ten to twenty grains of the powdered seeds, thrice a day.

RESOLVENT POWDER.

Pulvis ex Phellandro aquatico gummosus.
(BRERA)

R. Seeds of Water Hemlock, one scruple.
Gum Arabic, two drams.
Divide into six parcels.—Dose, one every

three hours, in scrofula and scrofulous phthisis.

ANTI-CATARRHAL POWDER. (AUGUSTIN;
SAINTE-MARIE)

R. Seeds of Water Hemlock, one dram.
Julep, } of each, one ounce.
Gum Arabic, }
Sugar of Milk, two ounces.

Mix.—Dose, a tea-spoonful in a glass of sweetened water three or four times a day; in chronic catarrhs. (SAINTE-MARIE)

AUGUSTIN prescribes five grains of water hemlock, eight of gum, six of nitrate of potass, and ten of sugar.—The dose to be repeated thrice a day.

ANTI-CATARRHAL EMULSION. (VAN
MONS)

R. Seeds of Water Hemlock, one ounce.
Gum Arabic, one dram.
Triturate together, and add gradually
Water, nine ounces.
Strain, set aside, decant, and add
Simple Syrup, one ounce.

Mix.

TINCTURE OF WATER HEMLOCK. (AUGUSTIN; NIEMANN)

R. Seeds of Water Hemlock, half an ounce.
Alcohol, six ounces.
Digest for twenty-four hours, and add
Muscadel Wine, six ounces.
Digest again for two days.

AUGUSTIN gives the same formula, but does not add the wine.

Dose, from ten to sixty drops, in pituitous phthisis.—MARTIUS considers it a very efficacious remedy.

PECTORAL DECOCTION.

Decoctum pectorale in phthisi. (BRERA)

- R. Bitter-sweet, two drams.
 Seeds of Water Hemlock, one dram.
 Water, a sufficient quantity

to obtain eight ounces of decoction. Add to it when strained,

- Almond Emulsion, two ounces.
 Oxy mel, one ounce.

Mix.—To be taken at twice, in the morning.

RESOLVENT PILLS.

Pilule in Bronchitide lenta. (BRERA)

- R. Seeds of Water Hemlock, three scruples.
 Proto-chloruret of Mercury, four grains.

Make fifteen pills.—Dose, three, every two hours in chronic catarrhs.

PHŒNIX DACTYLIFERA, LINN.—DATE TREE.

BAVAR.; BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A tree of the East Indies, of the north of Africa, and of the South of Europe. (*Diæcia Triandria*, LINN.; *Palmiers*, JUSS.; *Fig.* Flore Medic. III. 148)

The fruit, called *Dates* (*Dactyli*, *Palmulæ*.—Dattes, FR.; Dattel, GERM.; Datle, BOHEM.; Dadel, DUT.), are employed. They are oval longish drupes, of a yellow colour; under a thin smooth pellicle presenting a succulent pulp, enveloping a hard seed, one of the sides of which is marked with a longitudinal groove.

It is emollient and pectoral.

PULP OF DATES.

Pulpa Dactylorum. (GALL.)

- R. Dates, any quantity.
 Boiling Water, a sufficient quantity.

Boil, to soften the fruit; then remove the stones and pass the pulp through a hair-sieve, add to it the water of the decoction, and evaporate slowly to the proper consistence.

PASTE OF DATES. (GALL.)

- R. Dates, freed from their Stones, one pound and a half.

Cut them into small pieces, and boil them in Water, ten pounds, for one hour till they may easily be bruised between the fingers, and strain the decoction. Dissolve

Gum Senegal, six pounds, in twenty pounds of water, and strain the solution. Mix it with the other liquor, and add

- White Sugar, five pounds.
 Whites of Eggs, mixed with a little Water, five.

Boil, skimming, till reduced to one-third; then strain through a linen cloth, evaporate to the consistence of thick syrup, and add

- Orange-flower Water, nine ounces.

Evaporate to the consistence of soft extract, pour it into oiled tin moulds, and dry in a stove.

PHOSPHORUS.

1. *That which is met with in Commerce.*

AMER.; BATAV.; BAVAR.; BELG.; FENN.; SARD.; SUEC.; COXE.

2. *Purified Phosphorus of Commerce.*

BATAV.; BELG.

- R. Phosphorus of Commerce, any quantity.

Enclose it in a linen bag plunged in cold water; heat this gradually, and when the phosphorus is softened, press it gently through the cloth: then melt it in hot water, pour it into glass tubes, and keep them under water in a bottle covered with black paper.

3. *Prepared according to Formule.*

ANT.; AUSTR.; BAVAR.; BRUNS.; DAN.;

GALL.; FULD.; HANN.; HASS.; OLDEN.; POL.; BORUS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI.; SWEDIAUR.; VAN MONS.

A.

Phosphorus extracted from Urine.

BRUNS.; FULD.; WIRTEM.; HERBIP.

R. Muriate of Ammonia, in powder, two pounds.
Deutoxide of Lead, four pounds.

Distil in a glass retort, preserve the product, set apart the residuum, first putting three pounds of it into an iron vessel containing

Urine, putrified for eight weeks, and evaporated to the consistence of honey, ten pounds.
Charcoal, in powder, half a pound.

Stir the mixture till it is reduced to a black powder, which has ceased to exhale a urinous odour; fill with it to three-fourths a retort made of clay and charcoal (*de terre braquée*), adapt to this a receiver a little more than half filled with water, and distil with a gradually augmented heat. Collect the phosphorus which falls drop by drop into the water. (BRUNS.; HERBIP.)

R. Salt of Urine, obtained by crystallizing Urine evaporated to the consistence of Honey, eight parts.
Charcoal, in powder, one part.

Distil as before. (FULD.; WIRTEM.)

This process is abandoned.

B.

Phosphorus extracted from Bones.

ANT.; AUSTR.; BAVAR.; DAN.; GALL.; FERR.; HANN.; HASS.; OLDEN.; POL.; BORUS.; SAX.; BRUGNATELLI.; SWEDIAUR.; VAN MONS.

R. Bones of Mammiferous animals calcined to whiteness, and powdered, twelve parts.

Mix them in a wooden tub with a sufficient quantity of water to form a thick paste, and add gradually

Sulphuric Acid (66 degrees), ten parts, constantly stirring with a stick. Then pour more water upon the mass to reduce it to a thin paste, and leave it in repose for twenty-four hours: then put it upon a close cloth, and pour boiling water upon it till that passes through tasteless. Evaporate the liquor in a copper boiler; wash the precipitate which forms, and then, having mixed the water which is used in washing it with the former, evaporate the liquor to the consistence of syrup; add one fourth of char-

coal in powder, and heat nearly to redness in an iron stove to dry it completely. Fill a stone-ware retort to three fourths with this mixture; place it in a reverberatory furnace furnished with its dome; adapt to the neck a copper vessel prolonged into a tube which, passing through a cork, plunges into a glass bottle filled nearly to one half with water: then distil, regulating the fire in such a manner that the retort do not become red hot till at the expiration of an hour. Collect the phosphorus which passes into the receiver, and strain it through chamois leather under water heated to forty-five degrees.

Phosphorus is said to be aphrodisiac. It is one of the most violent excitants and most dangerous poisons. It should never be employed but with extreme circumspection, but it would be more prudent to banish it from the materia medica, at least as an internal medicament.—Dose, from the fourth of a grain to one grain daily.

PULVERIZED PHOSPHORUS. (HANN.)

R. Phosphorus, one dram.

Put it into a phial of the capacity of three ounces, and pour upon it

Water, one ounce.

Heat by plunging the phial into boiling water. When the phosphorus is liquified, shake the bottle strongly. Dry the powder upon a filter.

PHOSPHORATED EMULSION.

Emulsio phosphorata. (AUGUSTIN; NIEMANN)

R. Phosphorus, from two to four grains.
Oil of Almonds, one ounce.

Digest till dissolved; then make an emulsion with

Sweet Almonds, blanched, three.
Gum Arabic, half an ounce.
Water, three or four ounces.
Syrup of Marshmallows, half an ounce.

Mix.—Dose, half a spoonful or a spoonful, according to the quantity of the phosphorus.

PHOSPHORATED POTION. (AUGUSTIN; ELLIS; NIEMANN)

R. Phosphorus, two grains.
Mucilage of Gum Arabic, a sufficient quantity.

Spring Water, half a pound.

Make an emulsion, and add

Syrup of Almonds, one ounce.
HOFFMANN'S LIQUOR, thirty drops.

Mix.—Dose, a spoonful every two hours.

R. Spirit of Turpentine, half a dram.
Olive Oil, three drams.
Phosphorus, two grains.
Boiling Water, a sufficient quantity.
Gum Arabic, half an ounce.
Cinnamon Water, four ounces.
Syrup of Cinnamon, one ounce.

Mix.

R. Phosphorus, one grain.
DIPPEL's Animal Oil, one dram.
Oil of Cloves, one scruple.

Mix.—Dose, two or three drops, morning and evening, in paralytic amaurosis, tic doloieux and chronic rheumatism. (AUGUSTIN)

PHOSPHORATED ALCOHOL. (BRUGNATELLI)

R. Alcohol, eight ounces.
Phosphorus, cut small, eight grains.

Triturate in a glass mortar till the alcohol shall have lost its agreeable taste, and decant it.

PHOSPHORATED ÆTHER.

1. With Phosphoric Æther.
Æther Phosphoro-phosphoricus. (VAN MONS)

R. Phosphorus, one part.
Phosphoric Æther (60 degrees), sixty parts.

Shake strongly till solution is effected. (VAN MONS)

R. Phosphorus, four grains.
Phosphoric Æther, half an ounce.
Oil of Cloves, sixteen drops.

Dissolve at a moderate heat, in a covered vessel. (VAN MONS)

R. Phosphorus, one scruple.
Sub-carbonate of Ammonia, half a scruple.

Triturate together in a mortar moistened with phosphoric æther; put the mixture into a flask, and pour upon it

Phosphoric Æther,
two ounces and a half.

Shake till perfectly dissolved. (VAN MONS)

2. With Sulphuric Æther.

Æther sulphuricus phosphoratus, Naphtha Vitrioli phosphorata. (ANT.; DAN.; GALL.; HANN.; AUGUSTIN; BRUGNATELLI; CADET DE GASSICOURT; NIEMANN; PIERQUIN)

A.

Formula of BOUTTAZ or of PELLETIER.

R. Rectified Æther, four ounces.
Phosphorus, twenty-eight grains.

Dissolve. (ANT.; CADET DE GASSICOURT)

GALL. prescribes two drams and a half of phosphorus, and one pound of æther.—NIEMANN—four grains of phosphorus, and half an ounce of æther.—BRUGNATELLI—eight grains of phosphorus, and eight ounces of æther.—DAN.; and HANN.—sixteen grains of phosphorus, and two ounces of æther.—AUGUSTIN—two grains of phosphorus, and two drams of æther.

B.

Formula of FLITTNER.

R. Phosphorus, twenty-four grains.
Oil of Cloves, three drams.

Dissolve, and add

Sulphuric Æther, twenty-one drams.

Mix. (NIEMANN)

PIERQUIN prescribes three grains of phosphorus, twelve drops of oil of cloves, and half an ounce of æther.

C.

Formula of LÖBESTEIN LÜBEL.

R. Phosphorus, two grains.
Oil of Peppermint, from one scruple to half a dram.

Dissolve, and add

Sulphuric Æther, half an ounce.

Mix. (NIEMANN)

R. Phosphorus, three drops.
Sulphuric Æther, half an ounce.
Essential Oil of Valerian, ten drops.

Mix. (AUGUSTIN)

One ounce of pure æther dissolves six grains of phosphorus.—Dose, from two to four drops, on sugar, within twenty-four hours, increasing it daily by two or three drops.

PHOSPHORATED TINCTURE OF SUBCARBONATE OF POTASS. (VAN MONS)

R. Tincture of Subcarbonate of Potass, any quantity.

Heat it slowly in a sand-bath, and throw into it, by small pieces,

Phosphorus, a sufficient quantity, or till it ceases to be dissolved; set aside for some time, and decant.

EXCITANT POTION.

Mixtura Chinæ phosphorata. (AUGUSTIN)

R. Cinnamon Water, three ounces.
Extract of Cinchona, half an ounce.
Phosphorus, dissolved in Sulphuric Æther, three grains.
Syrup of Orange Peel, three ounces.

Mix.—Dose, one spoonful every two hours.

PHOSPHORATED LINIMENT. (FERR.; CADET DE GASSICOURT)

R. Phosphorus, cut, thirty grains.
Oil of Almonds, one pound.

Dissolve slowly, and strain when cold.
(CADET DE GASSICOURT)

FERR. prescribes twelve grains of phosphorus, and four ounces of oil.

It is applied in chronic rheumatism, paralysis, and atrophies.—From half an ounce to one ounce to be used at each application.

PHOSPHORATED LINIMENT OF CAMPHOR.

Linimentum phosphoratum camphoratum.
(HANN.; AUGUSTIN; CADET DE GASSICOURT; NIEMANN; VAN MONS)

R. Phosphorus, ten grains.
Camphor, two scruples.
Oil of Almonds, one ounce.

Dissolve. (HANN.; NIEMANN)

VAN MONS prescribes one scruple of phosphorus, four scruples of camphor, and one ounce of oil.—AUGUSTIN—three grains of phosphorus, six of camphor, and two drams of oil.

R. Phosphorus, six grains.
Oil of Almonds, one ounce.

Triturate together, digest in a water-bath, and add

Camphor, } of each,
Solution of Ammonia, } ten grains.

Mix. (HAMB.; AUGUSTIN)

R. Spirit of Turpentine, two ounces.
Camphor, two drams.
Pyro-oleose Subcarbonate of Ammonia, one dram.
Soap, two ounces.

Phosphorus, dissolved in twelve or fifteen drops of Oil of Cloves,
one grain.

Mix. (AUGUSTIN; NIEMANN)

PHOSPHORATED OINTMENT. (GALL.; CADET DE GASSICOURT)

R. Phosphorus, twenty parts.
Lard, one thousand parts.
Water, one hundred parts.

Boil till the phosphorus is dissolved and the water evaporated; then filter through bibulous paper, and add, *ad libitum*, one part of oil of lavender. (GALL.)

PHOSPHORATED OIL.

Oleum phosphoratum. (AUGUSTIN)

R. Phosphorus, two grains.
Oil of Henbane by infusion, half an ounce.
Extract of Chamomile, } of each,
Peppermint Water, } two ounces.

Mix.—Dose, a spoonful every two hours.

MAGENDIE prescribes, after LESCOT, an *Aromatic phosphorated oil*, of which the following is the formula:

R. Phosphorus, one ounce.
Olive Oil, one pound.

Digest at the ordinary temperature in a dark place for fifteen days, and flavour with oil of bergamotte.

Dose, from twenty-five to thirty drops, in twenty-four hours; in potion. (MAGENDIE)

PHYSALIS ALKEKENGİ, LINN.—WINTER CHERRY, ALKEKENGE.

SYNONIMES:—Alkekenge, Coqueret, FR.; Judenkirschen, GERM.; Jædekirsbær, DAN.; Alcúquerija, SPAN.; Kricken van over zee, DUT.; Alcachingi, ITAL.; Miechanki, POL.; Alcúqueriga, PORT.; Judekørsbær, SWED.

AMST.; ANT.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A plant of the South of Europe (*Pentandria Monogynia*, LINN.; *Solaneæ*, JUSS.; *Fig. Flore Medic. I.* 16)

The fruit (*Baccæ Alkekengi* seu *Solani vesicarii* seu *Halicacabi*) is employed. It

consists of globular, bilocular berries, as large as cherries, red, enclosing flat reniform seeds, inodorous, and of a sharpish yet sweetish taste, unless they may have been left in contact with the vesiculous calyx which encloses them, and which is extremely bitter.

They are diuretic.—Dose of the recently expressed juice, one ounce.

TROCHISCS OF WINTER CHERRY.

Trochisci Alkekengi. (SARD.; WIRTEM.)

R. Winter Cherries, ten drams.
Armenian Bole, } of each,
Gum Arabic, } five drams.
Mastic, }

Amber,	} of each,	five drams.
Seeds of Poppy,		
White Henbane,		
Opium,		one dram.
Mucilage of Tragacanth,		a sufficient quantity

to make a mass to be divided into trochiscs. (SARD.)

R. Winter Cherries,	} of each,	one ounce.
Seeds of White Poppy,		
Smallage,	} of each,	one ounce and a half.
Henbane,		
Gum Arabic,		
Starch,		
Gum Tragacanth,		
Root of Liquorice,		
Extract of Liquorice,		

Armenian Bole,	} of each,	eleven ounces.
Olibanum,		
Dragon's Blood,	} of each,	two drams.
Prepared Amber,		
Terra Sigillata,		

Moisten the powder with rose water, and make a mass to be divided into trochiscs. Two drams of opium may be added. (WIRTEM.)

Each dram contains one grain of opium.—These trochiscs were formerly extolled in the diseases of the urinary passages.—Dose, from two scruples to a dram.—The opium was not inserted when the patient had continued the use of them for some time.

PHYTOLACCA DECANDRA, LINN.—AMERICAN NIGHTSHADE.

SYNONIMES:—American Pukeweed, Gargel.—Phytolaque à dix étamines, Raisin d'Amerique, Morelle en grappes, Herbe de la laque, Méchoacan du Canada, FR.; Amerikanische Kermesbeere, GERM.; Fitolacca, Herba camin, SPAN.; Tienmannige luckplant, DUT.; Pianta lacca, ITAL.

AMER.; BRUNS.; HISP.; GALL.; ROSS.; WIRTEM.; COXE; MURRAY and GMELIN; SPIELMANN.

A plant originally of North America but naturalized in the south of Europe. (*Decandria Digynia*, LINN.; *Atriplicea*, JUSS.; *Fig. Zorn, Ic. pl. t. 164*)

The root and herb are employed.

The root (*Radix Phytolaccae* seu *Solani racemosi*) is thick, fleshy, and divided into several large fibres.

The herb consists of a cylindrical, branching stem, and of leaves of four or five

inches in length, alternate, glabrous, oval, lanceolated, and supported upon short footstalks. They have a very acrid taste.

HAYWARD says that the root may be substituted for ipecacuanha, in the dose of one scruple. The extract of the leaves has been extolled externally, in ulcerated cancer. The infusion has also been commended in rheumatism and piles.

OINTMENT OF AMERICAN NIGHTSHADE.

Unguentum Radicis Phytolaccae. (NIEMANN)

R. Powdered Root of American	
Nightshade,	one or two ounces.
Lard,	one pound.

Mix.

HAYWARD says it is serviceable in itch and obstinate tinea. It frequently excites heat and redness of the skin, when the ointment should be laid aside and a lotion used.

PIGMENTUM INDICUM.—INDIGO.

SYNONIMES:—*Indicus Color*.—Indigo, FR.; Nil, ARAB.; CYNG.; DUK.; and PERS.; Cham nho la, COCHIN.; Tarum, MAL.; Anil, PORT.; Nili, Nilini, SANS.; and TAM.; Nilum, TEL.

BRUNS.; WIRTEM.; AINSLIE; BERGIUS; GUIBOURT; SPIELMANN.

It is in small, solid, brittle masses, inodorous and insipid, of a deep azure blue co-

lour, assuming a cupreous lustre when rubbed.

It is obtained from the *Indigofera Anil, argentea*, and *tinctoria*, LINN. (*Diadelphia Decandria*, LINN.; *Leguminosae*, JUSS.), cultivated in South America. The *Isatis tinctoria* also yields it but in much less quantity.

It contains, with various other substances, a peculiar principle, *Indigotine*, which en-

ters, in the proportion of nearly one half, into the fine indigo of commerce.

It is sometimes, though rarely, employed to colour certain pharmaceutic preparations.

TINCTURE OF RESIN OF INDIGO. (BRUGNATELLI)

R. Resin of Indigo, one dram.
Alcohol, a sufficient quantity.

Dissolve and filter.—The resin of indigo

is obtained by distilling four parts of nitric acid with one part of indigo.

LIQUID SULPHATE OF INDIGO. (MAGENDIE)

R. Bengal Indigo, in powder, one part.
Concentrated Sulphuric Acid, six parts.

Digest with heat, and dilute it with

Distilled Water,
nine hundred and ninety-three parts.

This liquor is employed to ascertain the strength of the alkaline chlorurets, by the decoloration of it which these effect.

PIGMENTUM URUCU.

Urucu, Orleana, Orleana Terra.

BRUNS.; WIRTEM.; BERGIUS.; GUIBOUT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A colouring paste in cakes or balls of three or four pounds weight, brown with-

out and red within, of a soft consistence, of an animal odour and astringent taste.

This substance is prepared by collecting the red coating which covers the ripe fruit of the *Bixa orellana*, LINN.; (*Polyandria Monogynia*, LINN.; *Tiliaceae*, JUSS.; *Fig. Desc. Fl. Ant. I. 4*) a middling-sized tree of South America.

PILA MARINA.

SYNONIMES:—*Sphæra marina*, *Sphæra thalasia*, *Halcyonium rotundum*.—Pelote de mer, FR.

WIRTEM.; SPIELMANN.

By these names are designated certain light, round, depressed or oblong, inodorous, insipid masses, in which rushes, hair, and often fragments of shells are to be seen. They are found on the borders of the ocean and of the Baltic sea.

They are considered by SPIELMANN to be the agglomerations of the roots of the *Zostera marina*, LINN., a plant very common in our seas. (*Montecia Monandria*, LINN.; *Aroidæ*, JUSS.; *Fig. Ced. Fl. Dan. t. 15*)

CAVOLA assures us that he has found iodine in the ashes of the *Zostera*, which explains the reason of the *Pila Marina* having been classed among the substances which are useful in goitre.

PIMPINELLA.

Three species of this genus are employed in medicine:

1. *Pimpinella Anisum*, LINN.—Anise.

SYNONIMES:—Anis cultivé, FR.; Anis, GERM.; DAN.; SPAN.; and SWED.; Anison, ARAB.; Anyz, BOHEM.; and POL.; Kadis manis, BAL.; Souf, DUK.; Anisu, DUK.; Anys, DUT.; Mungfi, Adis manis, JAV.; Jera manis, MAND.;

Raryaneh rumie, PERS.; Herba doce, PORT.; Sataphuspha, SANS.; Sombu, TAM.; Sompu, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.;

AINSLIE ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOUT ; MURRAY and GME-LIN ; PARIS ; PIDERIT ; SPIELMANN ; ZARDA .

A plant originally of the Levant, and cul-tivated in Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; *Fig.* ZORN, *lc.* pl. t. 128)

The seed (*Semen Anisi vulgaris*) is em-ployed. It is long, oval, pediculated, flat on one side, bellied and grooved on the other, and of a greenish-yellow colour. It has an agreeable and aromatic odour, and an aromatic and sweet taste.

It contains an essential oil.

It is stimulant and carminative.—Dose, from fifteen grains to half a dram of the powder.

OLEOSACCHARUM OF ANISE.

Oleosaccharum Anisi. (AUSTR. ; BRUNS. ; DAN. ; GALL. ; HANN. ; OLDEN. ; PA-LAT. ; POL. ; BORUS. ; WIRTEM. ; SWE-DIAUR)

R. Essential Oil of Anise, one drop.
White Sugar, one dram.
Triturate for a long time. (BRUNS. ; DAN. ; GALL. ; PALAT. ; WIRTEM.)

AUSTR. ; HANN. ; OLDEN. ; POL. ; and BORUS. prescribe one dram of sugar, and three drops of oil.—SWEDIAUR—one dram of sugar and two drops of oil.

OPHTHALMIC POWDER.

Tragea ophthalmica. (BRUNS. ; SPIEL-MANN)

R. Seeds of Anise, } of each,
Fennel, } half an ounce.
Prepared Wood Lice, }
Root of Valerian, two drams.
Cinnamon, } of each, one dram.
Cubebs, }
White Sugar, two ounces.

Make a fine powder.

SPECIES DIANISI. (WIRTEM.)

R. Root of Pyrethrum, two drams.
Ginger, } of each,
Galanga, } five drams.
Spikenard, one dram and a half.
Liquorice, one ounce.
Calamint, half a dram.
Seeds of Anise, two ounces and a half.
Caraway, } of each,
Fennel, } five drams.
Lesser Cardamoms, } of each, one
Cloves, } dram and a half.
Cubebs, }

White Pepper, }
Black Pepper, } of each, two drams.
Pimento, }
Cinnamon, five drams.
Cassia Lignea, two drams.
Mace, five drams.
Saffron, one dram and a half.
Mastic, one ounce.

Make a powder.—Dose, from two scruples to one dram.

CARMINATIVE POWDER.

Pulvis ad flatum. (PALAT. ; WIRTEM. ; SPIELMANN)

R. Seeds of Anise, }
Caraway, } of each,
Coriander, } one ounce.
Fennel, }
Orange Peel, } of each, six drams.
Ginger, }
Prepared Crab's Eyes, half an ounce.
Mace, one dram.
White Sugar, six ounces.

Make a powder. (PALAT. ; WIRTEM.)

R. Seeds of Anise, }
Caraway, } of each,
Cumin, } three ounces.
Ginger, six drams.
Mace, three drams.
Saffron, one dram.

Make a powder. (SPIELMANN)

Dose, from a scruple to a dram.

DIGESTIVE POWDER. (PIERQUIN)

R. Anise, }
Juniper, } of each, three drams.
Root of Sweet Flag, }
Elicampane, } of each,
Florentine Iris, } two drams.
Ambergris, one dram
Sugar Candy, six drams.
Oil of Cinnamon, thirty drops.

Make a powder.—Dose, from a scruple to half a dram.

STOMACHIC POWDER. (SAINTE-MARIE)

R. Anise, }
Cinnamon, } of each, four scruples.
Ginger, }
Sugar of Milk, four ounces.

Make a powder.—Dose, a tea spoonful thrice a day.

CARMINATIVE BOLUSES. (CADET DE GAS-SICOURT)

R. Anise, sixteen drams.
Gentian, }
Angelica, } of each, twelve grains.
Castor, six grains.
Oil of Anise, twelve drops.

Oil of Cinnamon,	four drops.
Extract of Lesser	
Centauri,	of each,
Theriac,	thirty-five grains.
Syrup of Mint,	a sufficient quantity.

Make thirty boluses.—Dose, three, every four hours.

SPITZLAY'S LOZENGES. (HENRI)

R. Powdered Anise,	three parts.
Sugar,	five hundred parts.
Aqueous Extract of Opium,	one half part.
Gum Arabic,	sixty-four parts.
— Tragacanth,	of each,
Extract of Liquorice,	eight parts.

Make lozenges.

ANTI-HYSTERIC EPITHEM.

R. Anise, in powder,	two drams.
Theriac,	two ounces.
Oil of Cloves,	eight drops.
Spirit of Camphor,	a sufficient quantity.

Mix.—It was recommended by ZIMMERMANN.

PECTORAL LOHOCH.

Lohoch pulmonale seu de Pulmone Vulpis.
(HANN.; WIRTEM.; RATIER; SPIELMANN)

R. Powdered Anise,		
— Fennel,		
— Extract of Liquorice,	of each,	
— ice,	one	
— Canadian Maiden-hair,	ounce.	
— Fox Lung,		
Syrup,	fifteen ounces.	

Mix. (WIRTEM.; SPIELMANN)

R. Powdered Anise,		
— Fennel,		
— Extract of	of each,	
Liquorice,	one ounce.	
Syrup of Marshmallows,	twelve ounces.	

Mix. (HANN.)

R. Powdered Anise,	half a dram.
Water,	four ounces.
Sugar,	two drams.

Mix. (RATIER)

The two first formulæ give a medicament much celebrated formerly in cough and pulmonary phthisis; they used to be administered in the dose of two or three drams. The latter is well adapted to children, and may be given in tea spoonful doses.

OIL OF ANISE.

Oleum Anisi. (HISP.; GALL.)

R. Anise, bruised, any quantity.

Leave it exposed for eight minutes to the vapour of boiling water, upon a reversed sieve; then enclose it in a bag, and express. (GALL.)

HISP. directs the seeds to be bruised in a heated marble mortar, before expression.

ESSENTIAL OIL OF ANISE.

Oleum Anisi æthereum, Ætheroleum Anisi.
(LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FULD.; HANN.; HASS.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Oleum Anisi. (LOND.; DUBL.)

Ol. volatile Pimpinellæ Anisi.
(EDINB.)

R. Seeds of Anise, any quantity.
Pure Water,

a sufficient quantity
to prevent empyreuma. Distil, and collect the supernatant oil.

AMER.; AMST.; BATAV.; BELG.; FENN.; ROSS.; and COXE give the same formula.

ANT.; and BRUGNATELLI prescribe one part of seeds and three parts of water.—FULD.; and SWEDIAUR—one of seeds and four of water.—DAN.; and VAN MONS—one of seeds and six of water.—HISP.—one of anise and seven of water.—HANN.; OLDEN.; POL.; BORUS.; and SAX.—one of anise and eight of water.—BAVAR.—one of anise and nine of water.—GALL.—five of anise and seven of water.

R. Seeds of Anise,	four parts.
Water,	sixteen parts.
Muriate of Soda,	one part.

Distil. (BRUNS.; HANN.; PALAT.; SARD.; WIRTEM.; PIDERIT)

R. Seeds of Anise, any quantity.
Distilled Water of Anise,

a sufficient quantity.
Infuse for twelve hours, and distil. (HERBIP.)

R. Seeds of Anise,	one part.
Water,	twelve parts.

Macerate for twelve hours, and distil, with constant ebullition, till the water passes over insipid; re-distil this with an equal quantity of fresh seeds, adding spring water

till the desired quantity of oil be obtained. (AUSTR.)

Dose, from five to ten drops.

DISTILLED WATER OF ANISE.

Aqua Anisi. (BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FULD.; HASS.; SAX.; SARD.; PIDERIT; SWEDIAUR; VAN MONS)

R. Bruised Anise, one part.
Water, nine parts.

Macerate for twelve hours, and distil six parts. (DAN.)

BRUNS.; directs one part of anise to be macerated for several days in six parts of water, and three parts to be distilled.—BAVAR.—one of anise to be macerated for twenty-four hours in ten of water, and six to be distilled.—HERBIP.—one of anise, eight of water, and twenty-four hours' maceration.—SARD.—one of anise, six of salt water, and a few days' maceration.—GALL.—two of anise and fifteen of water; four to be distilled.—HASS.—one of anise, ten of water, and twenty-four hours' maceration; six to be distilled.—PIDERIT—one of anise, sixteen of water, and twenty-four hours' maceration; six to be distilled.—SAX.—one of anise, twenty-two of water, and twelve hours' maceration; two thirds to be distilled.—SWEDIAUR—one of anise and a sufficient quantity of water to obtain twenty parts of product.

R. Bruised Anise, one pound.
Subcarbonate of Potass, three ounces.
Water, ten pounds.

Distil slowly one half; and separate the oil. (FULD.)

R. Anise, three pounds.
Water, six pounds.
Alcohol, four ounces.

Macerate for two days, and distil three pounds in a water bath. (HISP.)

Dose, one or two ounces.

R. Oleosaccharum of Anise, one part.
Distilled Water, thirty-two parts.
Dissolve. (BAVAR.)

SPIRIT OF ANISE.

Alcohol seu Spiritus Anisi. (LOND.; ANT.; AUSTR.; BRUNS.; FULD.; HASS.; PALAT.; HERBIP.; WIRTEM.; BRUGNATELLI; COXE; PIDERIT)

Spiritus Anisi. (LOND.; COXE)

R. Seeds of Anise, bruised,
half a pound.

Proof Spirit, one gallon.
Water, a sufficient quantity
to prevent empyreuma. Macerate
for twenty-four hours, and distil one
gallon with a gentle heat.

AUSTR. prescribes one pound of anise, six pounds and a half of alcohol (0.910), and six pounds of water; to be digested for twenty-four hours, and six pounds of spirit marking 0.910, to be distilled in a water-bath.

R. Anise, bruised, one part.
Alcohol (20 degrees), nine parts.

Distil till the product marks 28 degrees. (ANT.; BRUNS.; PALAT.; WIRTEM.)

HERBIP. prescribes one part of anise and eight parts of alcohol.—FULD.; HASS.; BRUGNATELLI; and PIDERIT—one of anise and twelve of alcohol.

Dose, from sixty to one hundred and twenty drops.

COMPOUND SPIRIT OF ANISE.

Alcohol Anisi compositum, Spiritus Anisi compositus. (DUBL.; GENEV.; COXE; NIEMANN; SWEDIAUR)

Spiritus Anisi compositus. (DUBL.)

R. Seeds of Anise, } of each,
bruised, } half
——— Angelica, } a pound.
bruised,
Proof Spirit, one gallon.
Water, a sufficient quantity

to prevent empyreuma. Distil one gallon.

COXE; NIEMANN; and SWEDIAUR give the same formula.

GENEV. prescribes three ounces each of anise and of angelica, three pounds of proof spirit, and a sufficient quantity of water to prevent empyreuma; three pounds to be distilled.

It is carminative and stomachic.—Dose, from half a dram to four drams, in water or wine.

PHYSAGOGUE LIQUOR. (PALAT.; WIRTEM.)

R. Seeds of Anise, one pound.
——— Mountain Lasserwort,
one ounce and a half.
Cinnamon, }
Nutmegs, } two ounces.
Galanga, }
Mace, } of each, half an ounce
Cloves, }

White Wine, sixteen pounds.
 Alcohol, two pounds.
 Distil slowly one half. (WIRTEM.)

PALAT. prescribes two pounds of anise, one ounce of carrot seeds, two ounces of cinnamon, and two drams each, of the four latter substances.

It is stomachic and carminative.—Dose, from half an ounce to one ounce.

MYNSICHT'S TINCTURE.

Tinctura Proprietatis Mynsichti.
 (WIRTEM.)

R. Alkermes Confection, half an ounce.
 Flowers of Clove Pink, one dram.
 Spirit of Anise, sixteen ounces.
 Crocated Tincture of Aloes and
 Myrrh, one ounce.
 Digest, and filter.

It is excitant, and is reputed stomachic, carminative, and pectoral.—Dose, from one dram to one dram and a half.

INFUSION OF ANISE. (RATIER)

R. Anise, half an ounce.
 • Boiling Water, two pounds.
 Digest, and strain.

It is stimulant, tonic, and stomachic.

CARMINATIVE MIXTURE. (PIERQUIN)

R. Root of Marshmallows, two ounces.
 Water, two pounds.

Boil down to one half, and add

Seeds of Anise,	} of each,
— Fennel,	
— Coriander,	} two drams
— Rue Water,	
Pennyroyal Water,	} of each,
	} four ounces.

Macerate for four hours in a covered vessel, and add

Syrup of Poppy, four ounces.
 SYDENHAM'S Laudanum, forty drops.

Mix.

CARMINATIVE LAVEMENT. (SPIELMANN)

R. Seeds of Anise, }
 — Cumin, } of each, two drams.
 Juniper, }
 Water, sixteen ounces.
 Boil down to twelve ounces, and dissolve
 in the strained liquor
 Venice Soap, half an ounce.

AROMATIC ELIXIR.

Claretum aromaticum. (AUGUSTIN)

R. Seeds of Anise, half an ounce.

Seeds of Dill,	} of each,
— Fennel,	
— Coriander,	
— Caraway,	
Proof Spirit,	} half an ounce.
Infuse for a month, and add	
Sugar,	} of each, twenty ounces.
Water,	

Mix.—Dose, one or two spoonfuls, in dyspepsia.

2. *Pimpinella magna*, LINN.

SYNONIMES : — Boucage élevé, FR. ;
 Schwarze Bibernell, Blau Wurzel, GERM.
 GALL. ; WIRTEM. ; BERGIUS ; MURRAY
 and GMELIN ; SPIELMANN.

This plant grows all over Europe. (Fig. Jacq. Austr. 4. t. 396)

The root (*Radix Pimpinellæ nigra* seu *Dauci cyanopi* seu *Tragoselini majoris*) is employed. It is long, of the thickness of the finger, branching at the extremity, nearly black without, and bluish within. It has a strong, aromatic and agreeable odour, and an acrid, warm, and aromatic taste.

This plant may be used indiscriminately with the following :

3. *Pimpinella Saxifraga*, LINN.—Pimpinell Saxifrage, Burnet Saxifrage,

SYNONIMES : —Boucage à feuilles de pimprenelle, FR. ; Rossbibernelle, Steinbibernelle, Beckspeterlein, GERM. ; Bedznjk, BOHEM. ; Steenbrekke, Qvæseurt, Piergroed, Pimpinelle, DAN. ; Pimpinella blanca, SPAN. ; Kleine bevernell, DUT. ; Pimpinella bianca, ITAL. ; Piedrzyniec, POL. ; Pimpinella branca, PORT. ; Pækrot, SWED.

AMST. ; ANT. ; AUSTR. ; BRUNS. ; DAN. ; GALL. ; FENN. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; CAST. BORUS. ; BORUS. ; SARD. ; SAX. ; WIRTEM. ; HERBIP. ; BERGIUS ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN.

This plant grows in every part of Europe. (Fig. Jacq. Austr. IV. t. 395)

The root (*Radix Pimpinellæ albæ* seu *Nos-tratis* seu *Umbellifera* seu *hircinæ* seu *Tragoselini*) is employed. It is long, cylindrical, sub-fusiform, rather fibrous, striated circularly, of a yellowish-grey colour without, and white within. It has a strong goatish smell, provoking sneezing ; its taste is bitter, acrid, and burning.

It is excitant, diaphoretic, and resolvent.—Dose of the powder, from ten to twenty grains.

INFUSION OF BURNET SAXIFRAGE.
(CAST. BORUS.)

- R. Root of Burnet Saxifrage, one ounce and a half.
Boiling Water, one pound and a half.
Infuse for a quarter of an hour in a covered vessel, and strain.

EXTRACT OF BURNET SAXIFRAGE.

Extractum Pimpinellæ albæ. (PALAT.; POL.; BORUS.; WIRTEM.)

- R. Root of Burnet Saxifrage, one pound.
Boiling Water, six pounds.
Macerate for three days; then boil a little, strain with expression, and evaporate. (PALAT.)

- R. Root of Burnet Saxifrage, } of each,
Alcohol, } two pounds.
Water, } nine pounds.

Digest for twelve hours, express, distil off the alcohol, and evaporate the rest to the proper consistence. (POL.)

BORUS.; and SAX. prescribe two parts of root, three of alcohol, and nine of water.

- R. Root of Burnet Saxifrage, one pound.
Alcohol, five or six pounds.

Digest, and express; then digest and boil the residue in

Water, three pounds.

Clarify the decoction with white of egg. Mix the liquors, distil off the alcohol, and evaporate the rest to the proper consistence. (WIRTEM.)

TINCTURE OF BURNET SAXIFRAGE.

Tinctura seu Essentia Pimpinellæ albæ. (AUSTR.; BRUNS.; DAN.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; NIEMANN.; SWEDIAUR)

- R. Root of Burnet of Saxifrage, five ounces.
Rectified Spirit, two pounds.
Digest, express and filter. (HANN.; OLDEN.; POL.; BORUS.; NIEMANN)

AUSTR.; and SUEC. prescribe two ounces of root, and one pound of alcohol (0.910).—BRUNS.; DAN.; PALAT.; and WIRTEM.—one of root, and five of alcohol.—SWEDIAUR—two and a half of root, and six of alcohol.—FULD.; LIPP.; and SAX.—one of root, and four of alcohol.

It is excitant, diaphoretic, alexipharmic, diuretic, incisive, and alterative; and is principally recommended in the catarrhal affections of the fauces.—Dose, from thirty drops, to forty or more.

ALKALINE TINCTURE OF BURNET SAXIFRAGE.

Tinctura seu Essentia Pimpinellæ albæ kalica. (HASS.; PIDERIT)

- R. Root of Burnet Saxifrage, three ounces.
Subcarbonate of Potass, one dram.
Rectified Spirit, one pound.

Digest, express, and filter.

It is excitant, sudorific, diuretic, and deobstruent.—Dose, from twenty to sixty drops.

COMPOUND TINCTURE OF BURNET SAXIFRAGE.

Tinctura Pimpinellæ composita, Mixtura resolvers. (SWEDIAUR)

- R. Tincture of Burnet Saxifrage, } of each,
Alkalized Tincture of Antimony, } four ounces.

Digest for several days, frequently shaking. Add to the strained liquor

Æthereal Tincture of Amber, two ounces.

Medicinal Soap, one ounce.

Dissolve.

It is stimulant, and is extolled in chronic catarrhs, calculous nephritis, articular rheumatism, gout, and certain diseases of the skin.—Dose, from twenty drops to one dram, twice or thrice a day.

VULNERARY SPECIES.

Species pro Decocto vulnerario. (PALAT.; WIRTEM.; SPIELMANN)

- R. Root of Burnet Saxifrage, } of each,
Arum, } one ounce.
Herb of Sanicle, }
Wintergreen, } of each,
Ground Ivy, } half an
Speedwell, } ounce.
Tops of Lesser Centaury,
Cut and mix. (WIRTEM.)

- R. Herb of Ladies-mantle, }
Agrimony, }
Bugle, }
Great Comfrey, } of each,
St. John's-wort, } one
Thoroughwax, } ounce.
Burnet Saxifrage,
Plantain,
Tree Lungwort,
Golden Rod,
Nettle,
Root of Great Comfrey, } of each,
Pontic Rhubarb, } one pound.
Tormentil,

Cut and mix. (PALAT.)

- R. Root of Elicampane, } of each,
Great Comfrey, } half an
Tormentil, } ounce.

Root of Swallow-wort, half an ounce.
 ——— Liquorice, six drams.
 Herb of Thoroughwax, }
 ——— Sanicle, } of each,
 ——— Plantain, } half an
 ——— Mouse-ear, } ounce.
 ——— St. John's-wort,
 ——— Agrimony, }

Herb of BurnetSaxifrage, }
 ——— Bugle, } of each,
 ——— Golden Rod, } half an
 Flowers of Daisy, } ounce.
 Seeds of Fennel, six drams.
 Cut and bruise. (WIRTEM.; SPIEL-
 MANN)

PINUS.

The two following species of this genus are mentioned in pharmacopœias:

1. *Pinus Pineæ*, LINN.—Cultivated Pine Tree.

SYNONIMES:—Pin à pignons, FR.; Pinienbaum, GERM.; Domacy borowice, BOHEM.; Spansk syrt, DAN.; Pino alvo, SPAN.; Tamme pyomboom, DUT.; Pino sativo, ITAL.; Pinheiro manso, PORT.; Spansk tall, SWED.

BRUNS.; FERR.; GALL.; WIRTEM.; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A tree of the South of Europe. (*Monœcia Monadelphica*, LINN.; *Conifera*, JUSS.; Fig. Nouv. Duh. V. t. 72 bis, fig. 3. and t. 73)

The seeds (*Fircones*; *Nuces seu Nuclei Pineæ seu Pini seu Pincoli*) are employed. They are ligneous shells, sometimes hard and difficult to be broken, and sometimes soft and easily broken; they contain an oblong flattish kernel, larger at one end than at the other, about half an inch in length, and covered with a thin brown pellicle.

This kernel has an agreeable taste, and may be substituted for sweet almonds.

2. *Pinus sylvestris*, LINN.—Wild Pine,

SYNONIMES:—Pin sauvage, FR.; Gemeine Fichte, Kiefer, GERM.; Lesnj borowice, BOHEM.; Furs, DAN.; Pino, SPAN.; and ITAL.; Pyn, DUT.; Sosna parowa, POL.; Pinheiro, PORT.; Tall, FURN, SWED.

ANT.; AUSTR.; BRUNS.; DAN.; GALL.; FERR.; FORM. HÔP. FR.; FENN.; FULD.; GENEV.; HAMB.; HANN.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.;

BERGIUS; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A tree common in the greatest part of Europe, particularly in the north, and in mountainous regions. (Fig. Nouv. Duh. V. t. 66)

The young shoots (*Pini Turiones*, erroneously called *Strobili seu Coni Pini*) are employed. They appear in spring at the extremity of the branches. It consists usually of five or six lateral buds placed round the base of a larger terminal bud of from half an inch to an inch long. These buds are clothed with reddish scales, and are saturated with resin.

They have an agreeable and resinous odour, and a bitter and slightly aromatic taste.

It is excitant and anti-scorbutic.

DECOCTION OF PINE BUDS. (NIEMANN)

R. Pine Buds, three handfuls.
 Water, one pound and a half.

Boil for a quarter of an hour in a covered vessel; add to the liquor when cold, an equal quantity of white wine, and express at the end of twenty-four hours.

Dose, two ounces or more.

EMULSION OF FIR CONE KERNELS. (GALL.)

R. Fir cone Kernels, } of each,
 White Sugar, } one ounce.

Bruise for a long time in a marble mortar, adding water gradually, so as to form a soft paste. Then add,

Water (40 degrees, R.), one pound.
 Strain with expression, and add,

Orange-flower Water, half an ounce.
 Mix.

PIPER.—PEPPER.

This is the officinal name, applied in pharmacopœias, to six different plants.

1. *Piper Nigrum*, LINN.—Black Pepper

SYNONIMES:—Poivre commun, FR.; Pfeffer,

fer, GERM.; Filfil uswud, ARAB.; Czer-
ny pepr, BOHEM.; Micha, BAL.; Hu-
tseou, CHIN.; Gammiris, CYNG.; Peper,
DAN.; and DUT.; Kalimirchie, DUK.;
Pimienta, SPAN.; Miritch, Golmirch,
HIND.; Pepe, ITAL.; Maricha, JAV.;
Kaly mirchingay, MAH.; Lada, MAL.;
Saban, PALEM.; Filfil siab, PERS.;
Pieprz, POL.; Pimenta, PORT.; Perez,
RUSS.; Maricha, SANS.; Peppar, SWED.;
Mellaghu, TAM.; Mirialu, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.;
ANT.; BATAV.; BELG.; BRUNS.; DAN.;
HISP.; GALL.; FENN.; FERR.; FULD.;
HANN.; HASS.; LIPP.; OLDEN.; POL.;
BORUS.; ROSS.; SAX.; SUEC.; WIR-
TEM.; HERBIP.; AINSIE.; BERGIUS;
BRUGNATELLI.; COXE.; GUIBOURT;
MURRAY and GMELIN.; PARIS.; PIDE-
RIT.; SPIELMANN.; ZARDA.

A shrub of the East Indies. (*Fig. Flore
Medic. V. 276*)

The fruits (*Bacca Piperis* seu *Piper*) are
employed. They are round, hard berries,
of an aromatic and pungent odour, and of
an acrid, warm, pungent, and persistent
taste. They are black or brown externally
and white internally. Deprived of their
external pellicle they take the name of
White Pepper, *Piper Album*.

Their acridity is owing, according to
CERSTEDT, to a peculiar crystallizable prin-
ciple, similar to the resins, *Piperinum*, *Pi-
pera*.

2. *Capsicum Annuum*, LINN.—Capsicum, Guinea Pepper.

SYNONIMES:—Poivre de Guinée, Poivre
d'Espagne, Corail des Jardins, FR.; Spa-
nischer Pfeffer, GERM.; Spansk peber,
DAN.; Pimienta de Indias, SPAN.; Jaar-
lykse Spaansche peper, DUT.; Peperone,
ITAL.; Pieprz yca, POL.; Pimentao da
India, PORT.; Spansk peppar, SWED.

EDINB.; DUBL.; AMST.; ANT.; BATAV.;
BAVAR.; BRUNS.; DAN.; GALL.;
FERR.; FENN.; FORM. HÔP. FR.; GE-
NEV.; HASS.; POL.; BORUS.; SAX.;
WIRTEM.; HERBIP.; BERGIUS.; BRUG-
NATELLI.; COXE.; GUIBOURT.; MURRAY
and GMELIN.; PARIS.; SPIELMANN.

A plant of the Indies, from whence it has
been transported into America and Europe.
(*Pentandria Monogynia*, LINN.; *Solanææ*,
JUSS.; *Fig. Rheed. Malab. 2. tab. 35*)

The fruits (*Fructus Piperis Indici*, *Piper
Indicum* seu *Turcicum* seu *Hispanicum*) are
employed. These are dry, very smooth,
long berries, of a yellowish or vivid-red co-
lour, vesiculous, bilocular, of a very varia-

ble form, enclosing many flattish seeds.
These berries are destitute of odour; their
taste is very bitter, acrid, and burning.

Their acridity is attributed to a peculiar
alkaloid, *Capsinum*.

It is stimulant, but rather culinary than
medicinal.

AINSLIE does not say a word about this
species, but points out the *Capsicum frutes-
cens* (*Fig. Rumph. Amb. 5. tab. 88, f. 3.*),
also of the East Indies (Filfil, Achmar,
ARAB.; Tabia, BAL.; Meneschena, CAN.;
Lal Mirchie, HIND.; Lombok, JAV.; Tam-
bhudda mirchingay, MAH.; Lada mira,
MAL.; Fulfil, sarkh, PERS.; Brahn mari-
cha, SANS.; Mollaghai, TAM.; Merapa-
kaia, TEL.). This has the same properties
as the preceding.

3. *Myrtus Pimenta*, LINN.—Pimenta, Pi- mento, Allspice, Jamaica Pepper.

SYNONIMES:—Poivre de Jamaïque, Piment
des Anglais, Toute-Epice, FR.; Neue
Würze, Jamaikanischer Pfeffer, Engli-
scher Gewürz, Allerley Gewürz, Amœm-
lin, Nelkenpfeffer, Wunderpfeffer,
GERM.; Pimienta de Tabasco, Malag-
netta, SPAN.; Jamaica pepper, DUT.;
Angielski ziele, POL.; Kryddpepper,
SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.;
BATAV.; BELG.; BRUNS.; HISP.; GALL.;
FERR.; FENN.; GENEV.; HANN.; OL-
DEN.; POL.; BORUS.; SAX.; SUEC.;
BERGIUS.; BRUGNATELLI.; COXE.; GUI-
BOURT.; MURRAY and GMELIN.; PARIS.;
PIDERIT.; SPIELMANN.

A tree of the Antillas and East Indies.
(*Icosandria Monogynia*, LINN.; *Myrteæ*,
JUSS.; *Fig. Desc. Fl. Ant. III. 215*)

The fruits or berries (*Piper Jamaicaense*
seu *Pimenta* seu *Semen Amomi* seu *Piper
Tabasci*) are employed: they are dried be-
fore their maturity: they are wrinkled, a
little larger than the seeds of the black pep-
per, globular or oblong, umbilicated, and
of a deep brown colour. They have an aro-
matic odour and taste, resembling a mix-
ture of those of pepper, cinnamon, and
cloves.

4. *Piper longum*, LINN.—Long Pepper.

LOND.; EDINB.; DUBL.; ANT.; BRUNS.;
HISP.; FERR.; FORM. HÔP. FR.; WIR-
TEM.; GUIBOURT.; MURRAY and GME-
LIN.; SPIELMANN.

A sarmentose shrub of the East Indies.
(*Diandria Trigynia*, LINN.; *Urticææ*, JUSS.;
Fig. Blackw. Herb. t. 356)

The fruit (*Piper longum*) is a spike of several inches long, cylindrical, greyish and shagreened at the surface, of the diameter of a goose-quill, divided interiorly into several small cells, in each of which is a round seed. Its fracture is blackish and of a resinous appearance at the border and greyish and fibrous in the centre: its taste is acrid, hot, and bitterish. It contains, according to DULONG, the same principles as the black pepper.

5. *Piper Betel*, LINN.—Betel.

SYNONIMES:—Betel, Poivre des Indes, FR.; Tambul, ARAB.; PAN.; DUK.; and HIND.; Barg tambul, PERS.; Tambuli, SANS.; Vittiler, TAM.; Tamalapaku, TEL.

GALL.; AINSLIE.

A sarmentose shrub of the East Indies. (*Diandria Trigynia*, LINN.; *Urticeæ*, JUSS.; *Fig.* Flore Medic. II. 68)

The leaves are employed; they are rather large, subcordiform, acuminate, glabrous, and furnished with seven nervures of unequal length.

It is excitant, and irritant. It is considered febrifuge in the Indies.

6. *Piper Cubeba*, LINN.—Cubeb.

SYNONIMES:—Cubebes, Poivre à queue, FR.; Cubeber, Kubeber, Schwindelkerner, GERM.; Kebabeh, ARAB.; Kubeby, BOHEM.; Wal-gummeris, CYNG.; Cubeber, DAN.; Dumke mirchie, DUK.; Cubebas, SPAN.; Cubab chinie, HIND.; Koe beben, DUT.; Cubebi, ITAL.; Kumukus, JAV.; Komunkus, Lada barekor, MAL.; Kabeba, POL.; Cobebas, Pimenta rabuta, PORT.; Sugandha marichu, SANS.; Kobeber, SWED.; Val mellaghu, TAM.; Salavamirialu, TEL.

LOND.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; FENN.; FERR.; FULD.; GALL.; GENEV.; HAMB.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

A shrub of the East Indies. (*Diandria Trigynia*, LINN.; *Urticeæ*, JUSS.; *Fig.* Handb. der Pharm. Bot. II. p. 17)

The fruit (*Piper Cubeba* seu *caudatum*) consists of round, dry berries of the size of the seeds of black pepper, of a blackish grey colour, wrinkled, and supported upon a small peduncle. The external pellicle is easily broken, and covers another which is

blackish, and this encloses a white and oleaginous kernel. Its odour is agreeable and aromatic, and its taste is strongly aromatic and pungent.

It contains, according to VAUQUELIN, a resin similar to that of Copaiba, and which does not seem to differ much from Piperine, a small proportion of another coloured resin, gum, extractive, and salts.

It is stomachic, nerve, and strongly excitant.—Dose, from a scruple to two drams, twice or thrice a day, in syrup. It has been extolled in gonorrhoea; in its use in which affection the English have followed the Indians. These only employ it in the chronic stage of the disease, but in France it is given while there is yet acute inflammation. As it sometimes succeeds, though it fails or even aggravates the disease much more frequently, and always deranges more or less the digestive organs, it has been invested with a *specific* anti-gonorrhoeic virtue by some who prefer to believe blindly rather than give themselves the trouble of reasoning, and with whom the word specific serves as a rallying point. Fashion has brought cubeb to be employed in this affection as it did formerly in dyspepsia, and fashion will probably again leave it to neglect.

ERRHINE POWDER. (PIERQUIN)

R. Cubeb, } of each, one dram.
Valerian, }

Make a powder.

ANTI-GONORRHOEIC POTION. (PIERQUIN)

R. Cubeb, in powder, two drams.
Wine or Water, two or three ounces.
Essence of Bergamotte, one drop.

Mix.—To be taken every hour or two hours, till the discharge ceases. PIERQUIN has employed this potion since 1818.

TINCTURE OF BLACK PEPPER.

Essentia Piperis. (BRUNS.)

R. Black Pepper, one part.
Alcohol, six parts.
Digest and filter.—It is excitant.

TINCTURE OF CAPSICUM.

Tinctura Capsici seu *Capsici annui.* (LOND.; AMER.; HANN.; BORUS.; COXE.; ELLIS.; NIEMANN.; VAN MONS)

R. Capsicum, one ounce.
Proof Spirit, two pints.

Macerate for fourteen days, and filter. (LOND.)

AMER.; COXE.; and ELLIS give the same formula.—NIEMANN prescribes half an ounce

of capsicums and one pound of alcohol.—HANN.; and BORUS.—two ounces of pepper and one pound of alcohol.—VAN MONS—one part of pepper and ten parts of proof spirit.

It is excitant.—Dose, from one to three drams.—This tincture is sometimes employed as a gargle, diluted with two parts of water.

TINCTURE OF CAPSICUM AND OF CANTHARIDES.

Tinctura Capsici et Cantharidum. (AMER.; COXE; NIEMANN)

R. Cantharides, ten drams.
Capsicums, one dram.
Proof Spirit, one pound.

Digest for ten days, and filter.

This tincture is highly irritant, and is recommended in gangrenous affections; it may be given internally in the dose of ten drops in gum water or decoction of marsh-mallows.

STOMACHIC TINCTURE.

Essentia stomachica polychresta Gundelshheimeri. (WIRTEM.; SPIELMANN)

R. Capsicums, one ounce.
Black Pepper, }
White Pepper, } of each, two drams.
Long Pepper, }
Solution of Acetate of Potass, } of each,
Spirit of Ammonia, } six ounces.

Digest, and filter. (SPIELMANN)

WIRTEM. adds ten drams of lesser cardamoms.

It is stimulant, and is extolled as resolvent, in cachexia and dropsies.—Dose, from thirty to forty drops.

ERNSTING'S STOMACHIC TINCTURE.

Tinctura Stomachica acuta Ernstingii. (NIEMANN)

R. Capsicums, one ounce.
Black Pepper, }
White Pepper, }
Long Pepper, }
Grains of Paradise, } of each,
Cinnamon, } half an ounce.
Ginger, }
Root of Sweet Flag, }
Cretan Pennyroyal, }
Solution of Acetate of Potass, }
Alcohol, } twenty ounces.

Digest for several days in a warm place.

This tincture scarcely differs from the preceding.

Vol. II.

ÆTHEREAL TINCTURE OF PIMENTA.

Tinctura Pimentæ sulphurica. (SWEDIAUR)

R. Pimenta, three ounces.
Alcohol, fifteen ounces.

Infuse for fifteen days; add gradually to the strained tincture,

Sulphuric Acid, fifteen ounces.

It is stimulant, and is recommended in blennorrhœa and in affections attributed to atony of the stomach.—Dose, ten drops, twice a day in some proper vehicle.

VINEGAR OF PIMENTA. (VAN MONS)

R. Pimenta, one part.
Proof Spirit, four parts.
Vinegar, eight parts.

Infuse for eight days in a cool place, and filter.

VINEGAR OF CAPSICUM.

Acetum Capsici spirituosum. (VAN MONS)

R. Capsicums, two ounces.
Vinegar, } of each,
Proof Spirit, } twelve ounces.

Filter at the end of several days.

It is stimulant and antiseptic, and is recommended in laryngitis maligna, as a gargle with decoction of cinchona, in dropsy and cachexia.—Dose, two or three drams, several times a day, in any aromatic water.

DISTILLED WATER OF BLACK PEPPER. (VAN MONS)

R. Black Pepper, bruised, one part.
Water, fifteen parts.

Distil five parts.

DISTILLED WATER OF PIMENTA.

Aqua Pimentæ. (LOND.; DUBL.; EDINB.; BATAV.; BELG.; SWEDIAUR; VAN MONS)

R. Pimenta, half a pound.
Water, one pint.

Macerate for twenty-four hours; then add a sufficient quantity of water, and distil one gallon. (LOND.; DUBL.)

Aqua Myrti Pimentæ. (EDINB.)

R. Pimenta, half a pound.
Water, a sufficient quantity, to prevent empyreuma. Distil ten pounds.

BATAV.; BELG.; and SWEDIAUR give the same proportions as LOND.—VAN 3 A

MONS prescribes one pound of pimenta and eighteen pounds of water; six pounds to be distilled.

SPIRIT OF PIMENTA.

Spiritus Pimentæ; *Aqua Pimentæ vinosa*.
(LOND.; EDINB.; DUBL.; BELG.; FULD.;
GENEV.; HERBIP.; COXE.; SWEDIAUR)

Spiritus Pimentæ.

R. Pimenta, two ounces.
Proof Spirit, one gallon.
Water, a sufficient quantity,
to prevent empyreuma. Macerate for
twenty-four hours, and distil slowly
one gallon. (LOND.)

DUBL. gives the same formula but
requires three ounces of pimenta.

Spiritus Myrti Pimentæ. (EDINB.;
COXE)

R. Pimenta, half a pound.
Proof Spirit, nine pounds.

Macerate for two days in a cover-
ed vessel; then add a sufficient quan-
tity of water to prevent empyreuma,
and distil nine pounds.

R. Pimenta, one part.
Alcohol (20 degrees), sixteen parts.

Macerate, and distil a spirit marking 19
degrees. (BELG.)

SWEDIAUR prescribes one part of pi-
menta, four parts of proof spirit, and three
parts of water; four to be distilled.—GE-
NEV.—one ounce of pimenta, three pounds
of proof spirit, and a sufficient quantity of
water; three pounds to be distilled.

R. Pimenta, three ounces.
Subcarbonate of Potass, six drams.
Rhenish Wine, three pounds.

Digest for several days, and distil two
pounds. (FULD.)

R. Pimenta, one part.
Wine, eight parts.

Macerate for two days, and distil. (HERBIP.)

ESSENTIAL OIL OF CUBEBS.

Oleum Cubebæarum æthereum, *Ætheroleum Pi-
cubebæarum*. (BRUNS.; HANN.; SARD.)

R. Cubebs, one part.
Water, eight parts.

Distil. (HANN.)

BRUNS. prescribes four parts of cubebs,
sixteen of water, and one of muriate of
soda.—SARD.—one pound of cubebs, eight
pounds of water, and three ounces of mu-
riate.

ESSENTIAL OIL OF PIMENTA.

Oleum Pimentæ æthereum, *Ætheroleum Pi-
mentæ*. (LOND.; EDINB.; DUBL.; AMER.;
BRUNS.; GENEV.; PALAT.; SAX.;
SUEC.; HERBIP.; COXE)

Oleum Pimentæ. (LOND.)

Ol. Volatile Myrti Pimentæ. (EDINB.)

Ol. Baccarum Pimentæ. (DUBL.)

R. Pimenta, any quantity.
Water, a sufficient quantity.

Distil.

AMER.; GENEV.; and COXE give
the same formula.

SAX. prescribes one part of pimenta and
four parts of water.

R. Pimenta, one pound.
Water, sixteen pounds.
Muriate of Soda, two ounces.

Distil at the end of five or six days.
(BRUNS.; PALAT.)

R. Pimenta, any quantity.
Distilled Water of Pimenta,
a sufficient quantity.

Macerate for twelve hours, and distil.
(HERBIP.)

FEBRIFUGE POWDER. (NIEMANN)

R. Capsicums, six grains.
Bay Berries, two scruples.

Make a powder, and divide it into three
parcels. The first to be taken at the com-
mencement of the cold stage, the second on
the morrow at the same hour, and the last
on the following day. It is recommended
by BERGIUS.

STIMULANT PILLS. (ELLIS; NIEMANN)

R. Capsicums, one dram.
Rye Meal, five grains.
Water, a sufficient quantity.

Make twelve pills.—Dose, one every two
hours, in the black vomiting of yellow fever.
(NIEMANN)

R. Capsicums, one dram.
Crumb of Bread, } of each, a
Water, } sufficient quantity.

Make six pills.—Dose, one, three or four
times a day. (ELLIS)

FEBRIFUGE BOLUSES. (BRERA)

R. Piperine, one scruple.
Crumb of Bread, a sufficient quantity.

Make six boluses.—Dose, one every two
hours.—BRERA says these boluses fre-
quently fail.

STIMULANT PASTILES. (PIERQUIN)

R. Long Pepper, } of each, half a dram.
Nutmeg, }

Seeds of Stavesacre, } of each,
 Root of Pyrethrum, } two scruples.
 Mustard, } of each, one scruple.
 Origanum, }
 Honey, a sufficient quantity.

Make pastilles.

STIMULANT GARGLE. (NIEMANN)

R. Capsicums, half an ounce.
 Subcarbonate of Magnesia, one dram.
 Boiling Water, of each,
 Distilled Vinegar, } eight ounces.

Mix.—This formula is from HEADLY.—
 CURRIE employed an infusion of one grain
 of the capsicums in two ounces of water;
 ADAIR, the tincture of capsicums diluted
 with water.

OCYTIC POWDER. (PIERQUIN)

R. Seeds of Lavender, half a dram.
 ——— Plantain, } of each,
 ——— Endive, } two scruples.
 Pepper, one scruple.

Mix.—This, mixed with four ounces of
 endive water and four of chervil water, is
 administered to facilitate labour. It is also
 employed, rubbed upon the head, to destroy
 lice.

STOMACHIC ELECTUARY.

*Electuarium stomachicum alterans seu pi-
 perinum.* (SAUNDERS; SWEDIAUR)

R. Black Pepper, } of each,
 Conserve of Orange, } one ounce.
 Syrup of Orange Peel, a sufficient quantity.

Mix. (SAUNDERS)

R. Root of Sweet Flag, } of each,
 Galanga, } one ounce.
 Black Pepper, }
 Syrup of Orange Peel, a sufficient quantity.

Mix.—(SWEDIAUR) Half an ounce of
 black oxide of iron may be added.

It is recommended in gouty dyspepsia or
 cardialgia and in flatulent colic.—Dose, one
 dram twice a day.

ANTI-CACHECTIC ELECTUARY. (CADET
 DE GASSICOURT; SWEDIAUR)

R. Root of Elicampane, } of each,
 Fennel, } three ounces.
 Black Pepper, one ounce.
 White Sugar, } of each, two ounces.
 Honey, }

Mix.—Dose, one or two drams, thrice a
 day, in hæmorrhoidal cachexy, dyspepsia,
 and intestinal catarrh.

OINTMENT OF BLACK PEPPER.

Unguentum Pipiris nigri. (DUBL.; BORUS;
 COXE; CADET DE GASSICOURT; NIE-
 MANN)

R. Black Pepper, in powder, four ounces.
 Lard, one pound.

Triturate well together. (DUBL.;
 COXE)

R. Muriate of Soda, } of each,
 Black Pepper, } one pound.
 Soot, sifted, half a pound.
 Mustard, half an ounce.
 Yolks of Eggs, two.
 Lard, two pounds.

Make an ointment. (BORIES.)

R. White Soap, six ounces, two drams.
 Olibanum, five drams.
 Black Pepper, } of each,
 Muriate of Soda, } three drams.
 Alcohol, seven ounces.

Digest till the soap is dissolved; then
 boil for several minutes, stirring with a
 spatula, and spread it upon linen. (CADET
 DE GASSICOURT; NIEMANN)

All these ointments are rubefacient and
 more or less epispastic.

RUBEFACIENT CATAPLASM.

Cataplasma anti-pleuriticum. (GALL.;
 BORIES; CADET DE GASSICOURT; EL-
 LIS; PIERQUIN)

R. Long Pepper, } of each, half an ounce.
 Ginger, }
 White of Egg, a sufficient quantity.

Mix. (BORIES; CADET DE GASSICOURT;
 PIERQUIN)

R. Barley Meal, four ounces.
 Strong Vinegar, one ounce.
 Whites of Eggs, three.
 Water, a sufficient quantity.

Spread the whole upon tow, and sprinkle
 it with a mixture of

Powdered Black Pepper, } of each, half
 Fennel, } an ounce.

Apply it immediately. (GALL.; CADET
 DE GASSICOURT)

R. Mustard Flour, half a pound.
 Black Pepper, } of each, one dram.
 Ginger, }
 Simple Syrup, a sufficient quantity
 to make a soft paste. (ELLIS)

R. Powdered Capsicums, } of each,
 Cinnamon, } half an ounce.
 Cloves, }
 Flour, of each,
 Wine, } a sufficient quantity,
 to make a cataplasm. (ELLIS)

STIMULANT PLASTER.

Emplastrum piperitum, Dropax anti-paralyticum. (AUGUSTIN)

R. Pitch, two ounces.
Galbanum, } of each,
Black Pepper, } three drams.
Oil of Bay, a sufficient quantity.

Mix.—PLENK recommends it in paralysis.

INFUSION OF CAPSICUM. (ELLIS)

R. Capsicums, one dram and a half.

Boiling Water, half a pound.
Infuse.—To be taken by spoonfuls.

STIMULANT GARGLE.

Gargarisma piperitum. (AUGUSTIN)

R. Capsicums, half an ounce.
Subcarbonate of Magnesia, one dram.
Boiling Water, } of each.
Vinegar, } three ounces.

Infuse and Mix.—HEADLY recommends it in malignant sore throat.

PISTACIA VERA, LINN.—PISTACHIO TREE.

SYNONIMES :—Pistachier commun, FR.; Pistazienbaum, GERM.; Pistacy, Klockocka, BOHEM.; Pistaccira, SPAN.; Pistache boom, DUT.; Pistacchio, ITAL.; Fastuca, SIC.

BAVAR.; BRUNS.; GALL.; FERR.; GENEV.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMEIN; SPIELMANN; ZARDA.

An original tree of Asia and cultivated in the south of Europe. (*Diæcia Pentandria*, LINN.; *Terebinthaceæ*, JUSS.; *Fig.* Blackw. *Herb.* t. 46)

The seeds called *Pistachio Nuts* (*Pistaciæ*.—Pistaches, FR.; Pistazien, Syrische Nusschen, Pimperniisschen, GERM.), are employed. They are ovoid, oblong, reddish, wrinkled externally, and filled with an oleaginous kernel, of a pale green colour and agreeable taste.

PISTACHIO OIL.

Oleum Pistaciorum. (GALL.; SARD.)

R. Pistachio Nuts, any quantity.

Bruise them in a marble mortar with a wooden pestle; enclose the paste in a linen bag, and express the oil without heat.

PISTACHIO EMULSION. (GALL.)

R. Pistachio Nuts, } of each,
White Sugar, } one ounce.

Bruise together for a long time, adding water gradually; mix the paste with

Hot Water, one pound.

Strain with expression, and add

Orange-flower Water, half an ounce.

Mix,

PIX LIQUIDA.—TAR.

SYNONIMES :—*Pissa Cedria, Pix liquida nigra, Pix cedria, Resini Pini empyreumatica liquida, Terebinthina empyreumatica*.—Goudron, Poix liquide, FR.; Theer, Fliissiges Pech, GERM.; Tioere, DAN.; Zeer, DUT.; Zypera, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; BATAV.; BELG.; BRUNS.; DAN.; MIL. COPEN.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; ROSS.; SAX.; SUEC.; WIRTEM.; BERGIUS; COXE; GUIBOUT; PARIS; SPIELMANN.

A blackish-brown, tenacious, threading, semi-transparent liquid, of the consistence of very thick syrup, and of a resinous and em-

pyreumatic odour, with a bitter and resinous taste, and tinging the saliva with reddish-brown.

It is obtained by the slow, dry, and descending distillation of the chips of the wood of various coniferous trees, and consists of a half burnt resin, empyreumatic oil, and acetic acid.

Tar is a rather energetic stimulant which enters into the composition of several medicaments. It has also been prescribed alone, either with milk or beer, or in the form of pills, particularly in inveterate and obstinate itch. Its vapour has been recommended in the treatment of pulmonary phthisis. To reduce it to the pilular form, it is incorporated with hemlock powder,

and the mass thus obtained should be divided in pills of three grains each, of which the patient is first to take one, and augment the number daily till he takes twenty-four of them at once.

TAR WATER.

Aqua Picea seu Picis, Infusum Picis liquidæ seu Picis empyreumaticæ liquidæ, Potio picea. (DUBL.; ANT.; BAVAR.; BRUNS.; GALL.; FULD.; LIPP.; ROSS.; SAX.; SUEC.; WIRTEM.; CADET DE GASSICOURT; COXE; RATIER; SAUNDERS; NIEMANN; SPIELMANN; SWEDIAUR; PH. SYPH.; VAN MONS)

Aqua Picis liquidæ. (DUBL.)

R. Tar, two pints.
Water, one gallon.

Mix, stirring with a wooden rod for a quarter of an hour; then, after the tar shall have subsided, strain the liquor, and preserve it in well-corked bottles.

R. Tar, one part.
Water, four parts.

Stir well the mixture, and at the end of from a quarter of an hour to two days, pour off the supernatant liquor. (BRUNS.; SAX.; SUEC.; PH. SYPH.; COXE; SAUNDERS; SPIELMANN; SWEDIAUR)

NIEMANN prescribes one part of tar and five parts of water.—BAVAR.—one of tar and six of water; this is BERKELEY's formula.—FULD.; LIPP.; and WIRTEM.—one of tar and eight of water.—GALL.—five of tar and sixteen of water.—CADET DE GASSICOURT; and RATIER—one of tar and forty-eight of water; to be infused for one week.

R. Tar, one part.
Water, sixty-four parts.

Triturate together; then cover the vessel, leave it exposed to a gentle heat for a day, and then filter. (VAN MONS)

R. Water, twenty ounces.
Tincture of Tar, one scruple.

Instil the tincture into the water by three or four drops at a time, shaking the vessel quickly each time. (VAN MONS)

VAN MONS does not point out the manner of making this tincture; but it is known that tar is soluble in alcohol. His last formula differs probably from all the preceding by the presence, in the preparation, of a great quantity of empyreumatic oil; these, in fact, contain but a very little, which is rendered soluble in the water by the acetic acid.

This water was for some time held up as

a panacea on the credit of Bishop BERKELEY, but it is now nearly forgotten.—It was said to be anti-scorbutic, anti-variolous, anti-catarrhal, anthelmintic, alterative, diaphoretic, and diuretic. It is a rather strong stimulant to the stomach.—Dose, to children, to two or three ounces; to adults, about sixteen ounces.—To render it more agreeable, sugar may be added, with a few drops of oil of nutmeg or of sulphuric æther.

TAR OINTMENT.

Unguentum Picis Liquidæ. (LOND.; EDINB.; DUBL.; AMER.; MIL. COPEN.; SUEC.; COXE; SAUNDERS)

R. Tar, } of each,
Prepared Suet, } one pound.
Melt together, and strain through a linen cloth. (LOND.)

R. Tar, five parts.
Yellow Wax, two parts.
Melt the wax with a gentle heat; then add the tar and stir till cold. (EDINB.)

R. Tar, } of each,
Mutton Suet, } half a pound.
Melt together, and strain through a sieve. (DUBL.)

AMER.; and COXE give the same formula as EDINB.

R. Rye Meal, one pound.
Vinegar, a sufficient quantity.

Boil to a paste, and add,

Resin, } of each,
Tar, } half a pound.

Leave the mixture upon the fire, constantly stirring it till of a proper consistence. (SUEC.)

R. Tar, four ounces.
Rancid Salt Butter, two ounces.
Subcarbonate of Potass, one ounce.
Triturate together. (MIL. COPEN.)

R. Tar, one pound.
Sulphur, four ounces.
Melted Wax, one ounce.

Incorporate. (SAUNDERS)

All these ointments have been employed in cutaneous diseases but particularly in itch.

TAR LINIMENT.

Linimentum farinaceum cum Pice. (AUGUSTIN)

R. Rye Meal, four parts.
Gentian, in powder, one part.

Tar, ... a sufficient quantity.

Mix.—It is extolled in schirrus, conjointly with decoction of guaiac internally.

TAR PLASTER.

Emplastrum Picis liquidæ. (HANN.; SAX.; VAN MONS)

R. Yellow Wax, ... six parts.
Resin, ... three parts.

Melt together and add,

Tar, ... one part.

Strain the mixture, and stir till cold.
(VAN MONS)

HANN.; and SAX. prescribe eight parts of wax, one part of resin, and sixteen of tar.

PILLS OF TAR.

Pilulæ Picis. (AMER.; SARD.; COXE; NIEMANN)

R. Tar, ... any quantity.
Root of Elicampane, ... a sufficient quantity.

Make pills of six grains each. (AMER.; COXE; NIEMANN)

R. Tar, ... of each,
Balsam of Peru, } half an ounce.
Liquorice Powder, } one ounce.
Florentine Iris Powder, three drams.

Make a pillular mass. (SARD.)

Dose, half a dram, twice a day.—These are sometimes administered in pulmonary phthisis.

PLANTAGO.—PLANTAIN.

Five species of this genus are employed in medicine:

1. *Plantago major*, LINN.—Great Plantain.

SYNONIMES:—Plantain commun, Grand plantain, FR.; Grosser Wegerich, Wegetritt, GERM.; Vejbred, DAN.; Llantén major, SPAN.; Groote wegbblad, DUT.; Piantaggine maggiore, ITAL.; Kanasch, PERS.; Lubka, POL.; Tanchugem major, PORT.; Popuschnik, RUSS.; Grodblad, SWED.

AMST.; ANT.; BATAV.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; LIPP.; WIRTEM.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A plant which grows in all parts of Europe. (*Tetrandria Monogynia*, LINN.; *Plantagineæ*, JUSS.; *Fig. Flore Medic. V.* 275)

The root and leaves are employed.

The root (*Radix Plantaginis majoris* seu *latifolia* seu *Arnoglossi* seu *Septinervia*) is composed of a hard, thick, nearly ligneous stump, furnished below with cylindrical fibres.

The leaves are oval, large, obtuse, coriaceous, sinuated at the borders, and shining.

They are nearly inodorous, and have an herbaceous, bitterish, and slightly astringent taste.

2. *Plantago lanceolata*, LINN.—Small Plantain, Ribwort.

SYNONIMES:—Plantain lancéolé, Petit plantain, FR.; Spitzenwegerich, GERM.; Spids vejbred, DAN.; Smal-bladige wegbree, DUT.; Spets groblad, SWED.

GALL.; FERR.; WIRTEM.; SPIELMANN.

A plant growing all over Europe.—The root and herb are employed.

The root (*Radix Plantaginis lanceolata* seu *angustifolia* seu *quinquenervia minoris*) is sweetish and slightly astringent.

The herb consists of a bundle of lanceolated five-nerved leaves.

3. *Plantago media*, LINN.—Hoary Plantain.

SYNONIMES:—Plantain moyen, FR.; Mittelwegerich, GERM.; Gitrocel prostednj neb ssiroky, BOHEM.; Llantén medio, SPAN.; Middelbaare wegbree, DUT.; Piantaggine media, ITAL.; Tanchagem mediana, PORT.; Slosskæmper, SWED.

GALL.; FERR.; HERBIP.; ZARDA.

A plant of Europe. (*Fig. CEd. Fl. Dan. t.* 581)

The root and herb are employed.—The root is slightly astringent.

The herb consists of lanceolated leaves which have a styptic taste.

These three plants are slightly astringent.

CONSERVE OF PLANTAIN. (HERBIP.)

R. Fresh Herb of Plantain, ... one part.
Sugar, ... two parts.

Beat to a homogeneous mass.

JUICE OF PLANTAIN.

Succus Plantaginis. (SARD.; HERBIP.)

R. Fresh Herb of Plantain, ... any quantity.

Bruise it in a stone mortar, moistening it with a little water, and express strongly.

Set apart the juice, decant, and preserve it in bottles, adding one dram of alcohol to each pound of juice.

SARD. directs it to be clarified in a water-bath.—Dose, two or three ounces.

SYRUP OF PLANTAIN.

Syrupus Plantaginis seu Plantaginis majoris.
(HISP.; WIRTEM.)

R. Expressed Juice of Plantain, } of each,
 White Sugar, } four parts.
 Water, a sufficient quantity.

Clarify, and boil to the consistence of syrup. (HISP.)

R. Depurated Juice of Plantain,
 evaporated to one half, nine ounces.
 White Sugar, sixteen ounces.
Make a syrup. (WIRTEM.)

DISTILLED WATER OF PLANTAIN. (GALL.;
 (SARD.))

R. Fresh Herb of Plantain, five thousand parts.
 Water, twelve thousand five hundred parts.

Distil about ten thousand parts. Pour the product upon twice the quantity of fresh herb, add ten thousand parts of water, and again, distil ten thousand parts. The operation may be repeated a third time. (GALL.)

SARD. prescribes one part of herb and two parts of water; two-thirds to be distilled.

It enters into resolvent collyria, but it is perfectly inert.

EXTRACT OF PLANTAIN. (SARD.; WIRTEM.; HERBIP.)

R. Fresh Leaves of Plantain, any quantity.

Bruise them in a stone mortar, sprinkling them with a little water; express the juice and evaporate it immediately in a water-bath. (HERBIP.)

R. Leaves of Plantain, one pound.
 Spring Water, six pounds.

Digest for four days in a warm place, and then boil a little. Strain with expression and evaporate the liquor to the proper consistence. (SARD.; WIRTEM.)

ASTRINGENT FOMENTATION. (SAINTE-MARIE)

R. Leaves of Plantain, two handfuls.
 Petals of Red Roses, one pinch.
 Water, three pounds.

Boil down to one pound and a half, adding towards the end, the volume of a hen's

egg of quick lime. Set aside, decant, and add to it,

Proof Spirit, four ounces.

VULNERARY LIQUOR.

Aqua Calcis vulneraria. (SARD.)

R. Herb of Plantain, } of each,
 Sanicle, } one
 St. John's-wort, } handful.
 Wintergreen, }
 Quick Lime, two pounds.

Throw the powdered lime upon the chopped herbs, and leave them to macerate for several days; then pour over them

Pure Water, ten pounds.

Macerate for ten hours, and decant; pour over the residue

Plantain Water, ten pounds.

Macerate for twelve hours, and mix the two liquors. (SARD.)

4. *Plantago Psyllium*, LINN.—Clammy Plantain, Fleaseed.

SYNONIMES:—Herbe aux puces commune, FR.; Flohsaame, GERM.; Chmeljk, Blessnjik, BOHEM.; Psilleurt, DAN.; Zaragatena, SPAN.; Flookruid, DUT.; Pullicaria, ITAL.; Pulgueira, PORT.; Loppfrägras, SWED.

AMST.; ANT.; BATAV.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; FENN.; GENEV.; HANN.; OLDEN.; BORUS.; ROSS.; WIRTEM.; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of Egypt and of the temperate climates of Europe. (*Fig. Zorn, Ic. pl. t. 115*)

The seeds (*Semines Psyllii seu Pulicariæ*) are employed. They are small, long, of a deep brown colour, shining, smooth, soft to the touch, flat on one side, and bulged on the other. They are inodorous but have a mucilaginous, bitterish, and acrid taste.

It is emollient.

5. *Plantago Cynops*, LINN.

SYNONIMES:—Herbe aux puces vivaces, FR.; Strauchartizer Wegerich, GERM.

BAVAR.; BERGIUS.

A plant of Egypt and of the south of Europe.

Its seeds may be substituted for the preceding.

MUCILAGE OF FLEASEED.

Mucilago Psyllii. (GALL.; SARD.)

R. Seeds of Fleaseed Herb, two drams.
 Warm Water, one ounce and a half.

Digest for twenty-four hours in warm ashes, shaking the vessel from time to time, and strain through a linen cloth with expression. (GALL.)

SARD. directs one part of seeds to be macerated for twelve hours, and then boiled in ten parts of water; to be strained with expression.

PLATINA.

SYNONIMES :—Platine, Or blanc, Fr.

HANN.; HISP.

A solid metal, nearly as white as silver, very brilliant, ductile, malleable, and tenacious, infusible, and unalterable in the air. Its specific gravity is 20, 98.

CHLORURET OF PLATINA.

Chloruretum seu Murias Platinae. (NIEMANN)

R. Platina, any quantity.
Hydrochloro-nitric Acid, a sufficient quantity
to dissolve the metal. Evaporate and crystallize the solution.

It has been recommended in the treatment of syphilitic diseases.—Dose, from the fourth of a grain to one grain, to be rubbed upon the gums.

OXIDE OF PLATINA.

Oxydum Platinae. (NIEMANN)

R. Aqueous Solution of Chloruret of Platina, any quantity.
Pure Soda, a sufficient quantity,
or till it ceases to effect a precipitate: wash and dry this.

It is emetic and purgative, according to the dose.

PLUMBAGO EUROPEA, LINN.—LEADWORT.

SYNONIMES :—Dentelaire, Fr.; Bleywurz, Zahnwurz, GERM.; Velesa, SPAN.; Roodkruid, DUT.; Piombaggine, Crepanella, ITAL.

GALL.; FERR.; FULD.; HERBIP.; WIRTEM.; BRUGNATELLI; MURRAY and GMELIN.

A plant of the south of Europe. (*Pentandria Monogynia*, LINN.; *Plumbagineae*, JUSS.; *Fig. Flore Medic.* III. 149)

The root (*Radix Dentariae majoris seu Dentillariae seu Squamariae seu Ambluti*) is employed. It is straight, long, thick, scarcely branching, furnished with fibres. It is brownish without and white within, inodorous, and of an acrid and burning taste.

An infusion of it in oil has been employed against itch.

PLUMBUM.—LEAD.

SYNONIMES :—*Saturnus*; μόλυβδος.—Plomb, FR.; Blei, GERM.; Anuk, ARAB.; Hinyen, CHIN.; Blye, DAN.; Schisch, DUK.; Plomo, SPAN.; Sisa, HIND.; Bly, DUT.; Piombo, ITAL.; Tamaetam, MAL.; Surb, PERS.; Clow, POL.; Chumbo, PORT.; Swinez, RUSS.; Sisaka, SANS.; Bly, SWED.; Eium, TAM.; Schischum, TEL.

AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; GENEV.;

HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; COKE; GUIBOUT; PIDERIT; SPIELMANN.

A solid metal, of a bluish-white colour, soft, brilliant, insonorous, susceptible of being marked by the nail, malleable, not very tenacious, exhaling a sensible odour when rubbed. It is fusible at 322,22 degrees C., and volatilizable at a higher tem-

perature. Its specific gravity is 11,3523, according to BRISSON.

SULPHURET OF LEAD.

Sulphuretum Plumbi, Æs ustum. (AMST.; HISP.; FERR.; SARD.)

R. Lead, any quantity.

Melt it in an iron ladle, adding to it powdered sulphur by spoonfuls, till it is entirely converted into a black powder; porphyryze and preserve it. (AMST.; HISP.)

FERR.; and SARD. direct layers of lead and of sulphur to be stratified in an earthen pot and exposed to a violent heat; the residuum to be washed and porphyryzed.

AMST.; and SARD. erroneously designate the sulphuret of lead by the name of *Æs ustum*, which should be applied to the protoxide.

BLACK LEAD PLASTER.

Emplastrum Plumbi nigrum. (HISP.)

R. Sulphuret of Lead,	}	of each,
Protoxide of Lead,		two ounces.
Subcarbonate of Lead,		of each,
Sulphuret of Antimony,	}	one ounce.
Rose Oil,		three pounds.
White Wax,		nine ounces.

Melt the wax in the oil and add the other ingredients well porphyryzed; boil a little, and then stir till cold.

CHLORURET OF LEAD.

Magisterium Saturni. (SPIELMANN; VAN MONS)

R. Solution of Subacetate of Lead, one part.
Water, three parts.

Add to the mixture a solution of muriate of soda in six parts of water till it ceases to produce a precipitate: wash and dry this.

It is principally recommended as cosmetic, but it should not be used as such except with great circumspection.

MILK OF LEAD. (VAN MONS)

R. Muriate of Soda, one half part.

Dissolve it in the least possible quantity of water; add gradually to the solution

Solution of Acetate of Lead, one part.

Collect the precipitate, and having well washed it, mix it with

Almond Emulsion, twelve parts.

It is cosmetic.

PROTOXIDE OF LEAD.

I. That which is found in commerce.

Lithargyrum, Lithargyrium, Lythargyrus, Oxydum Plumbi fusum seu semifusum
Vol. II.

seu semivitreum, Oxydum plumbicum semivitreum.—Semivitreous Oxide of Lead, Litharge.

SYNONYMES:—Litharge, Oxide de plomb demi-vitreux, FR.; Bleiglatte, GERM.; Sølvglød, DAN.; Murdarzang, DUK.; HIND.; and PERS.; Lithargirio, Almagro, SPAN.; Lootglans, DUT.; Litar-girio, Piombo semivitreo, ITAL.; Gleyta, POL.; Silfwerlitt, SWED.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HER-BIP.; WURCEB.; AINSIE.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.

In small scales, greasy to the touch, produced by the crystallization of the melted and slowly-cooled protoxide of lead. According as the colour may be white or yellow, it is called *Silver Litharge* (*Lithargyrus*; Litharge d'argent, FR.; Silberglätte, GERM.), or *Gold Litharge* (*Lithargyrium*; Litharge d'or, FR.; Goldglätte, GERM.)

BRUGNATELLI describes the process employed in commerce in manufacturing litharge.

II. Prepared according to various formula.

1. *Æs ustum, Oxodes Plumbi nigrum, Oxydum Plumbi.*—Black or Grey Oxide of Lead.

BATAV.; BELG.; BRUNS.; FENN.; FULD.; HANN.; PALAT.; SAX.; WIRTEM.; SPIELMANN.; ZARDA.

R. Lead, any quantity.

Keep it in fusion in an earthen dish, stirring it frequently, and remove the grey stratum which covers it in proportion as it appears. When all the metal is converted into oxide, reduce it to powder, sift, porphyryze, and wash it in a large quantity of water.

2. *Cerussa citrina.*—Yellow Oxide of Lead, Yellow Ceruss.

VAN MONS.

R. Subcarbonate of Lead, any quantity.

Put it into a Hessian crucible covered with a tile, and apply a sufficient heat to expel all the carbonic acid, but not to fuse the oxide; when cold, triturate it with water, and form it into trochiscs.

WIRTEM. requires the yellow oxide of commerce.

It is a mixture of protoxide of lead with a small proportion of metal.

POWDER AGAINST ERYSIPELAS.

Pulvis erysipelatodes. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

R. Flour, } six ounces.
Black Oxide of Lead, } of each,
Armenian Bole, } two ounces.
Mastic, } of each,
Olibanum, } one ounce.
Subcarbonate of Lead, }

Make a very fine powder.

SATURNINE INJECTION. (SWEDIAUR)

R. Black Oxide of Lead, twenty grains.
Sulphate of Zinc, six grains.
Rose Water, four ounces.

Mix.—It is extolled by CLARE in gonorrhœa, even in the acute stage.

DEUTOXIDE OF LEAD.

Minium, Oxydum Plumbi rubrum, Deutoxydum Plumbi rubrum, Super-oxydum plumbosum, Plumbum oxydatum rubrum.—Red Lead, Minium, Red Oxide of Lead.

SYNONIMES:—Deutoxide de plomb, Oxide rouge de plomb, FR.; Menning, GERM.; Isrenj, ARAB.; Yuen-tan, CHIN.; Mænnie, DAN.; Sendur, DUK.; Minio, ITAL.; and SPAN.; Sindur, HIND.; Menie, DUT.; Temamera, MAL.; Minia, POL.; Sindura, SANS.; Alænja, SWED.; Segapu senduerum, Jum sindurum, TAM.; Yerra sinduriam, TEL.

EDINE.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BRUGNATELLI; COXE; PIERIT; SPIELMANN.

A very fine, very heavy, red powder.

BRUGNATELLI describes the process which is followed in preparing the red-lead or minium of commerce.

ASTRINGENT ERRHINE. (PIERQUIN)

R. Aloes, } of each,
Myrrh, } one ounce and a half.
Quick Lime, } one dram.
Deutoxide of Lead, half a dram.
Alum, four scruples.
Pomegranate Rind, } of each,
Flowers, } two drams.
Oil of Myrtle, } of each,
Rose Water, } two ounces and a half.
Plantain Water, a sufficient quantity.

Mix.—Against ulcers of the nostrils.

OINTMENT OF DEUTOXIDE OF LEAD.

Unguentum seu Ceratum Minii. (VAN MONS)

R. White Wax, one part.
Oil, two parts.

Incorporate with these melted together
Deutoxide of Lead, triturated with
a little Oil, one half part.

PLASTER OF RED OXIDE OF LEAD.

Emplâtre défensif rouge de Wepfer. (WIRTEM.)

R. Fresh Butter, } of each,
Linseed Oil, } six ounces.
Green Bark of Elder, four ounces.

Boil slowly to the consumption of the humidity; and add to the expressed mixture

Yellow Wax, } of each,
Deutoxide of Lead, } six ounces.

Mix.—It is recommended to be applied to chronic ulcers, and also to open cancers, to appease pain.

RED CAMPHORATED CERATE. (BRUNS.; PALAT.; WIRTEM.)

R. Yellow Wax, } of each,
Stag's Suet, } half a pound.

Melt over a slow fire, and incorporate in the nearly cold mass,

Deutoxide of Lead, triturated with
half a pound of Rose Oil, six ounces.

Add lastly

Camphor, one dram and a half.

CAMPHORATED OINTMENT OF DEUTOXIDE OF LEAD. (WIRTEM.; SPIELMANN)

R. Rose Oil, one pound and a half.
Crystallized Protoxide of Lead, two ounces.

Deutoxide of Lead, three ounces.
Subcarbonate of Lead, one ounce and a half.

Tutty, } of each, three drams.
Camphor, }
White Wax, two ounces.
Distilled Vinegar, three ounces.

Melt the wax over a slow fire, add the oil, then the other substances in powder, then the vinegar, and lastly the camphor triturated with a little alcohol. (WIRTEM.)

R. Ointment of Litharge, fifteen ounces.
Camphor, two scruples.

Triturate together, and add

Deutoxide of Lead, one ounce and a half.

Tutty, two drams.

Mix. (SPIELMANN)

This ointment, as the cerate which pre-

cedes it, is reputed desiccative and anodyne. It is applied to burns.

RED DEFENSIVE PLASTER. (HANN.; PALAT.; WIRTEM.; VAN MONS)

- R. Yellow Wax, two pounds.
 Resin, one ounce.
 Rose Oil, four ounces.
 Quick Lime, washed with
 Rose Water, and then } of each,
 dried, ten drams.
 Pounded Brick,
 Armenian Bole,
 Deutoxide of Lead, half an ounce.
 Camphor, dissolved in Oil, two drams.

Melt together the wax, resin, and oil; then add the dry ingredients, and finally the camphor. (WIRTEM.; PALAT.)

- R. Yellow Wax, } of each,
 Mutton Suet, } six ounces.
 Spermaceti, half an ounce.

Melt together, and add to the mass when half cold

- Deutoxide of Lead, three ounces.
 Subcarbonate of Lead, } of each,
 Camphor, dissolved in } six ounces.
 three ounces of Oil,

Mix. (HANN.)

- R. Deutoxide of Lead, two parts.
 Subcarbonate of Lead, four parts.

Triturate together, and incorporate with a melted mixture of

- Mutton Suet, eight parts.
 Olive Oil, six parts.
 Yellow Wax, sixteen parts.

Then add

- Camphor, triturated with a little Oil,
 one part.

Mix. (VAN MONS)

RED SPARADRAP.

Emplastrum ad Fonticulos. (PALAT.; VAN MONS)

- R. Turpentine, one part.
 Mutton Suet, two parts.
 White Wax, six parts.

Melt, and add

- Deutoxide of Lead, triturated with
 a little Suet, one part.

Dip strips of linen in the melted mixture. (VAN MONS)

PALAT. prescribes six ounces of yellow wax, two of stag's suet, one and a half each, of turpentine and of oil of lily, and four ounces of deutoxide.

RESOLVENT PLASTER. (CADET DE GAS-SICOURT; PIERQUIN)

- R. Diapalm Plaster, four ounces.

- Lard, } of each,
 White Wax, } two ounces.
 Melt together, and add

- Deutoxide of Lead, two ounces.

- R. Olive Oil, one pound twelve ounces.
 Deutoxide of Lead, one pound.
 Yellow Wax, four ounces.

- Venice Turpentine, } of each,
 Pulverized Loadstone, } one ounce and
 a half.

- Olibanum, } of each,
 Mastic, } three drams.
 Myrrh,
 Alum,

- Camphor, two drams.

Mix.

TRITOXIDE OF LEAD.

Oxydum Plumbi purum, Super-oxydum seu Peroxydum Plumbi. (VAN MONS)

- R. Deutoxide of Lead, one part.
 Nitric Acid, three parts.

Dilute the acid with its own weight of water, and pour it upon the pulverized oxide, stirring well; heat the mixture, decant the liquor, and well wash and dry the powder.

OINTMENT OF TRITOXIDE OF LEAD. (VAN MONS)

- R. Tritoxide of Lead, one part.
 Lard, twelve parts.
 Yellow Wax, two parts.

Make an ointment.

BROWN CERATE. (VAN MONS)

- R. Tritoxide of Lead, one part.
 Simple Cerate, eight parts.

Triturate in a glass mortar.

SUB-PROTO-CARBONATE OF LEAD.

Oxydum Plumbi carbonatum per Acidum acetosum, Oxydum Plumbi album, Cerussa alba Norica seu Hollandica, Subcarbonas Plumbi, Carbonas Plumbi seu plumbicum seu plumbicus, Subacetas Plumbi.—Subcarbonate of Lead, White Lead, Ceruss.

SYNONIMES:—Sous-carbonate de protoxide de plomb, Blanc d'argent ou de plomb, Blanc de Krems, Craie de plomb, Plomb-spathique, Méphite de plomb, Blanc de Cérusse, Oxide de plomb blanc, FR.; Bleiweiss, Schieferweiss, GERM.; Asfidaj, ARAB.; Blegghvidt, DAN.; Suffidah, DUK.; HIND.; and PERS.; Albayalde, SPAN.; Tootwit, DUT.; Cerussa, ITAL.; Bleywas, POL.; Blyhwitt, Kiferhwitt, SWED.; Vullay, Muthu vullay, TAM.; Sibaydu, TEL.

I. *That which is met with in Commerce.* Cerussa.—Ceruss.

LOND.; DUBL.; EDINB.; AMER.; AMST.;

ANT.; AUSTR.; BATAV.; BAYAR.;
BELG.; BRUNS.; DAN.; HISP.; GALL.;
FENN.; FERR.; GENEV.; HAMB.;
HANN.; HASS.; LIPP.; OLDEN.; LUSIT.;
POL.; CAST. BORUS.; BORUS.; ROSS.;
SAX.; SUEC.; WIRTEM.; HERBIP.;
WURCEB.; AINSLIE.; BRUGNATELLI.;
COXE.; GUIBOUT.; PARIS.; PIDERIT.;
SPIELMANN.

A white and very heavy powder.

VAN MONS indicates the following method of purifying the ceruss of commerce:

R. Ceruss, any quantity.

Triturate it in a porcelain mortar, adding gradually a weak solution of acetate of lead. After a short time, decant, and add more solution. Continue thus till the decanted liquor is precipitated by a drop of hydro-sulphate of potass in the same manner as pure water is. Wash the residue largely with water, and form it into trochiscs.

II. Prepared according to various Formule.

BORUS.; FULD.; WIRTEM.; BRUGNATELLI.; PIDERIT.; VAN MONS.

R. Thin Sheets of Lead, any quantity.

Suspend a certain number of them in a vessel containing good vinegar, and apply heat that this may be slowly evaporated; remove from time to time the white crust with which the sheets become covered, and continue the operation till nothing remains but these. (BRUGNATELLI.; PIDERIT.; VAN MONS)

R. Neutral Acetate of Lead, eighteen parts.
Effloresced Subcarbonate of Soda, five parts.

Dissolve the salts, each separately in an excess of water, and mix the solutions, strongly agitating; set aside, decant, wash the precipitate, and form it into trochiscs. (VAN MONS)

BRUNS.; FULD.; and WIRTEM. prescribe the subcarbonate of potass, and call the product *Magistery of Lead*, *Magisterium Plumbi*.

A current of carbonic acid gas may be passed through a solution of subacetate of lead till no more precipitate is formed. This process is indicated by THENARD, and is followed at Clichy.

RHAZES' WHITE TROCHISCS.

Trochisci Rhazis seu albi seu albi ex Rhazis seu Carbonati Plumbi camphorati, Pulvis e Plumbo seu e Cerussa compositus, Sief album. (AMST.; BRUNS.; HISP.; LUSIT.; PALAT.; SARD.; WIRTEM.; CA-

DÉT. DE GASSICOURT; PIERQUIN; PIDERIT; SPIELMANN; SWEDIAUR)

R. Subcarbonate of Lead, five drams.
Gum Tragacanth, three drams.

Triturate together. (SWEDIAUR)

LUSIT. prescribes two parts of subcarbonate, and one part of gum.

R. Subcarbonate of Lead, one ounce and a half.
Gum Ammoniac, } of each,
Starch, } two drams.
Camphor, half a dram.

Pulverize each separately, mix the powders, and make trochiscs with water.

R. Subcarbonate of Lead, five ounces.
Sarcocolla, one ounce and a half.
Gum Tragacanth, half an ounce.

Make a powder. (SPIELMANN)

One twenty-eighth of opium may be added.

R. Subcarbonate of Lead, six drams.
Gum Arabic, } of each,
Tragacanth, } half an ounce.
Sarcocolla, three ounces.

Make a powder. (BRUNS.; WIRTEM.)

Half a grain of opium may be added to each scruple, which gives the *Sief cum Opio*.

R. Subcarbonate of Lead, ten drams.
Starch, half an ounce.
Gum Arabic, } of each,
Tragacanth, } two drams.
Camphor, in powder, half a dram.

Make a fine powder. (AMST.)

R. Subcarbonate of Lead, ten drams.
Sarcocolla, three drams.
Starch, two drams.
Gum Arabic, } of each,
Tragacanth, } one dram.
Opium, half a dram.

Make a powder. (PIERQUIN)

CADET DE GASSICOURT prescribes half a dram of camphor instead of the opium.

R. Subcarbonate of Lead, six drams.
Starch, half an ounce.
Isinglass, three drams.

Make a powder. (PIDERIT)

One or two grains of opium may be added to each dram.

This powder, mixed with rose water, was formerly celebrated in ophthalmia, burns, and other inflammations arising from external causes.

COMPOUND SATURNINE POWDER. (SWEDIAUR)

R. Subcarbonate of Lead, one ounce.
Oak Bark, one pound.

Mix.—This is proposed to be sprinkled upon the surface of scrofulous ulcers.

BALLS AGAINST ERYSIPELAS.

Globuli ad Erysipelas. (SPIELMANN)

R. Subcarbonate of Lead, } of each,
Chalk, } one ounce.
White Bole, } two ounces.
Spirit of Camphor, a sufficient quantity
to make a paste to be formed into balls.

ANTI-OPHTHALMIC LINIMENT—(PIERQUIN)

R. Subcarbonate of Lead, four ounces.
Starch, three drams.
Gum Arabic, one dram.
— Tragacanth, dissolved in
Rose Water, one ounce and a half.

Mix.—This liniment scarcely differs from some of the formulæ of the *Sief album*, except by its liquidity.

OINTMENT AGAINST TIC DOLOREUX.
(RATIER)

R. Subcarbonate of Lead, one ounce.
Opium, one dram.
Lard, half an ounce.
Balsamum Tranquillum,
a sufficient quantity.

Mix.—It is recommended by FOUQUIER.

OINTMENT OF SUBCARBONATE OF LEAD.

Ceratum de Cerussa, Unguentum album seu album simplex seu Carbonatis plumbici seu Cerussæ seu Oxydi Plumbi Carbonati seu Subcarbonatis Plumbi. (EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; CADET DE GASSICOURT; PIDERIT; PIERQUIN; SPIELMANN; SWEDIAUR; VAN MONS)

Unguentum Carbonatis Plumbi.

R. Simple Ointment, five parts.
Carbonate of Lead, one part.

Mix. (EDINB.)

R. Ointment of White Wax,
one pound.
Carbonate of Lead, in powder,
two ounces.

Make an ointment. (DUBL.)

R. Subcarbonate of Lead, one part.
Lard, one part and a half.

Triturate together. (ANT.; BRUNS.; DAN.; GALL.; FULD.; HASS.; LIPP.; PALAT.; SUEC.; WIRTEM.; BRUGNATELLI; PIDERIT; SPIELMANN)

SAX. prescribes three parts of subcarbonate, and two of lard.

R. Lard, two parts.
Mutton Suet, one part.

Melt together, and add

Subcarbonate of Lead, two parts.

Triturate well. (BAVAR.; HANN.; OLDEN.; POL.; BORUS.)

FERR. prescribes two pounds of olive oil, two and a half of mutton suet, and two of subcarbonate.

R. Olive Oil, three parts.
White Wax, } of each,
Subcarbonate of Lead, } one part.

Make an ointment. (CADET DE GASSICOURT; PIERQUIN)

LUSIT.; and WIRTEM. prescribes four parts of oil, and one part each, of wax and of subcarbonate.—SWEDIAUR—thirteen of oil, two and a half of wax, and ten of subcarbonate; or, three of oil, one of wax, and one and a half of subcarbonate.—VAN MONS—directs twelve parts of subcarbonate to be triturated with a little oil, and then to add a mixture of five parts of white wax, and eighteen of olive oil.

R. White Wax, three parts.
Rose Oil, twelve parts.
Subcarbonate of Lead, four parts.

Make an ointment. (SARD.)

HISP. prescribes four parts of wax, eighteen of rose oil, and twelve of subcarbonate.

R. White Cerate, six parts.
Subcarbonate of Lead, one part.

Mix. (AMER.; and COXE)

R. Simple Ointment, three parts.
Melt over a slow fire, and add
Subcarbonate of Lead, one part.

Mix. (AMST.)

R. Simple Ointment, five parts.
Olive Oil, one half part.
Subcarbonate of Lead, one part.

Make an ointment. (BATAV.; BELG.)

This ointment is principally employed as a dressing for burns.

CAMPHORATED OINTMENT OF SUBCARBONATE OF LEAD.

Unguentum album camphoratum. (BATAV.; BELG.; BRUNS.; DAN.; FULD.; LIPP.; OLDEN.; PALAT.; POL.; SARD.; SAX.; WIRTEM.; HERBIP.; BRUGNA-

TELLI; CADET DE GASSICOURT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Ointment of Subcarbonate of Lead,
twenty-four parts.

Camphor, triturated with a little

Oil of Almonds, one part.

Mix by trituration. (BATAV.; BELG.; BRUNS.; OLDEN.; PALAT.; POL.; SAX.; WIRTEM.; BRUGNATELLI)

LIPP. prescribes one thirtieth of camphor.—SPIELMANN—one thirty-second.—DAN.—one fortieth.—FULD.—one forty-eighth.—CADET DE GASSICOURT; and SWEDIAUR—one sixty-sixth.—SARD.; and HERBIP.—one seventy-second.—VAN MONS—one ninety-sixth.

OINTMENT AGAINST BURNS. (VAN MONS)

R. Camphorated Ointment of } of each,
Subcarbonate of Lead, } any
Ointment of Marshmallows, } quantity.

Mix.

WHITE SPARADRAP. (VAN MONS)

R. Turpentine, one part.
Mutton Suet, two parts.
White Wax, six parts.
Subcarbonate of Lead, triturated
with a little suet, one part.

Melt and incorporate. Dip strips of linen in the mixture.

DESICCATIVE SPARADRAP. (CADET DE GASSICOURT; VAN MONS)

R. Olive Oil, one part.
Beef Marrow, one part and a half.
Yellow Wax, two parts.
Suet, three parts.

Melt together, and add.

Powdered Calamine, one part.
Subcarbonate of Lead,
one part and a half.

Dip strips of linen in the mixture. (VAN MONS)

CADET DE GASSICOURT prescribes six ounces of oil, four of wax, and one and a half each, of subcarbonate of lead and of calamine.

OINTMENT AGAINST BURNS. (PALAT.; WIRTEM.)

R. Camphor, one ounce.
Rose Oil, three ounces.
Subcarbonate of Lead, six ounces.
Powdered Olibanum, two ounces.
Whites of Eggs, seven.

Triturate together in a leaden mortar. (PALAT.)

R. Inner Bark of Elder, } of each, one
Lime, } ounce and a half.

Juice of Plantain, two ounces.
Great Houseleek, one ounce.
Linseed Oil, six ounces.
Stag's Suet, one ounce.

Boil over a slow fire to the consumption of humidity; then strain, and triturate it in a leaden mortar with

Subcarbonate of Lead, one ounce.
Sugar, two drams.
Oil of Eggs, half an ounce.

Make an ointment. (WIRTEM.)

DESICCATIVE OINTMENT. (SWEDIAUR)

R. Olive Oil, one pound.
White Wax, half a pound.

Add to the melted mixture

Calamine, half a pound.
Subcarbonate of Lead, nine ounces.
Protoxide of Lead, three ounces.

Stir till cold.

COMPOUND OINTMENT OF LIQUORICE.

Unguentum de Liquiritia. (PALAT.; WIRTEM.; SPIELMANN)

R. Pulp of Fresh Liquorice } of each,
Root, } sixteen
Fresh Butter, } ounces.

Boil to the consumption of the humidity; then express, and add to the liquid mass

Oxide of Zinc, six drams.
Subcarbonate of Lead, three ounces.
Camphor, in powder, two scruples.

Mix.—It is recommended in excoriations and fissures of the breast.

ANTI-HÆMORRHOIDAL OINTMENT.
(PALAT.)

R. Starch, }
Subcarbonate of Lead, } of each,
Protoxide of Lead, } six drams.
Grey Oxide of Lead, }
Gum Tragacanth, }
Opium, } of each, four scruples.
Camphor, }
Whites of Eggs, two.
Mucilage of Fleaseed, } of each,
Rose Oil, } half a pound.
Oil of Lesser Mullein, }
Yellow Wax, three ounces.

Melt the wax in the oils, then add the camphor, the powders, and lastly the whites of eggs.

COMPOUND SATURNINE OINTMENT.
(SWEDIAUR)

R. Subcarbonate of Lead, one ounce.
Protoxide of Lead, two drams.
Armenian Bole, one ounce

Triturate these with

Honey of Roses, half an ounce.
Lard, a sufficient quantity

to make an ointment.

This preparation was for a long time kept secret. It was extolled as an application to scrofulous, phagedenic, and cancerous ulcers, particularly of the penis.

It is exceedingly stimulant, as is proved by the acute pain which it occasions for some hours after its application. Two drams of tritoxide of iron are sometimes added, and then it resembles the red defensive plaster, particularly that which is prepared according to the formula of HISP.

OPHTHALMIC LIQUOR. (SAUNDERS)

R. RHazes' White Trochiscs, one dram.
Rose Water, four ounces.
Mix.

ASTRINGENT OINTMENT. (PIERQUIN)

R. Juice of Plantain, } of each,
——— Shepherd's Purse, } two
——— Lesser Mullein, } ounces.

Boil slightly, and add

Powdered Dragon's Blood, one ounce.
——— Armenian Bole, } of each,
——— Lemnian Bole, } one dram
——— Root of Bistort, } and a half.
Subcarbonate of Lead, two scruples.

Mix.

SUB-PROTO-SULPHATE OF LEAD.

This salt is never prepared separately, but it is incidentally formed when solution of acetate of lead is mixed with a selenitious water. It forms the base of the following preparation :

VIRGIN'S MILK. (BRUNS.; WIRTEM.)

R. Semi-vitreous Protoxide of Lead, half a pound.
Distilled Vinegar, one pound and a half.
Boil together to two-thirds. Strain, and add

Alum, four ounces.
Water, two pounds,
mixed together. Preserve the lactescent liquor which results.

It is a dangerous cosmetic.

PROTO-NITRATE OF LEAD.

This salt is only mentioned as one of the constituent elements of the following preparation :

BALSAM OF LEAD. (VAN MONS)

R. Lead, one part.

Diluted Nitric Acid,

a sufficient quantity
to dissolve the metal. Pour the solution while hot, continually agitating, into

Melted Lard, sixteen parts.

Pour the mass into paper moulds, and cut it when cold into convenient forms.

SUB-PROTO-ACETATE OF LEAD.

Acetas Plumbi liquidus, Acetas super-plumbicus aquosus, Aqua Lythargyri Acetatis, Decoctum saturninum, Liquor Acetatis Plumbi basica seu Plumbi subacetati seu Subacetatis Lythargyri, Solutio Plumbi in Acido acetoso, Subacetas Plumbi liquidus, Extractum seu Balsamum Saturni.—Solution of Subacetate of Lead, Extract of Lead.—(LOND.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; CAST. BORUS.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI.; COXE.; PIDERIT.; SAUNDERS.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

Liquor Plumbi Subacetatis. (LOND.)

R. Semi-vitrified Oxide of Lead, two pounds.

Dilute Acetic Acid, one gallon.

Mix, and boil down to six pints, constantly stirring; then set apart the solution that the impurities may subside, and strain.

Plumbi Subacetatis Liquor. (DUBL.)

R. Semi-vitreous Oxide of Lead, one part.

Distilled Vinegar, twelve parts.

Put them into a glass vessel, and boil, constantly stirring, till eleven parts remain; then set apart the solution till the fæces shall have subsided, and strain.

R. Semi-vitreous Oxide of Lead, one part.
Vinegar, four parts.

Digest for twenty-four hours in a glass matrass, occasionally stirring; then apply a moderate heat for three or four hours, till turnsol paper is no longer reddened by the liquor: then let it cool and preserve it. It should be of the specific gravity of 1.250: this may be concentrated till two ounces and two drams are of the same volume as one ounce of water.

VAN MONS directs twenty-one and a half parts of semi-vitreous oxide to be dissolved, by the heat of a sand-bath, in six parts of concentrated vinegar, and the solution to be evaporated to from 29 to 48 degrees.—PLANCHE—two pounds of the oxide and eight and a half of vinegar to be boiled in a leaden vessel till the mass has acquired the consistence of honey; nine pounds of water to be then added, the ebullition to be continued for a quarter of an hour, and the liquor evaporated to 27 degrees.—BRUGNATELLI—one part of the oxide to be dissolved in four parts of distilled vinegar, and the filtered liquor to be preserved.—AMER.; and COXE give the same process as LOND.—LIPP.—one pound of the oxide to be boiled in six pounds of vinegar till, on the one hand, ten drams occupy the volume of one ounce of water; and on the other, till two ounces of the liquor occupy the volume of one ounce of water.—FENN.; and SUEC.—one part of the oxide to be boiled in six parts of vinegar till the solution weighs 1,14.—AMST.—one part of the oxide to be dissolved in four parts of vinegar, the solution to be filtered and evaporated to the consistence of honey.—ANT.—one pound of the oxide to be boiled in eight ounces of vinegar till the liquor marks 27 degrees.—PALAT.; SARD.; WIRTEM.; and SAUNDERS—one part of the oxide to be boiled in four parts of vinegar for an hour, and the liquor to be filtered.—LUSIT.—four ounces of the oxide to be boiled in two pounds of vinegar till one pound and a half of solution remain.—GENEV.; HERBIP.; and SPIELMANN—one part of the oxide to be boiled in three parts of vinegar to the consumption of one half of the liquid.—OLDEN.—one of the oxide to be boiled in five of vinegar till this is perfectly saturated, and the liquor thus obtained to be evaporated till six drams of the liquor occupy the volume of an ounce of water.—BRUNS.—one pound of the oxide to be boiled in four pounds of vinegar to the consumption of one half of the liquid, and this to be further evaporated till only nine ounces remain.—HISP.; and SARD.—one part of the oxide to be digested in four parts of vinegar for several days, and the liquor to be filtered.—FULD.—one of the oxide to be boiled with two of concentrated vinegar.

SPIELMANN also directs one part of deutoxide of lead to be boiled in eighteen parts of vinegar till the solution has acquired the consistence of honey.—BORUS.—one of deutoxide and nine of distilled vinegar to be boiled together, and the liquor to be evaporated to one third.—DAN.—one of deutoxide to be boiled in four of vinegar, and the liquor to be evaporated to one half, and also to only three-fourths.

ROSS.; and SAX. direct one part of sub-

carbonate of lead and eight parts of vinegar to be boiled till reduced to one half.—HASS.; and PIDERIT—one of subcarbonate and eight of vinegar; either to be only boiled for a quarter of an hour, or to be evaporated to the consistence of honey.

ROSS. allows either the deutoxide or the subcarbonate to be used, and PIDERIT—either the deutoxide, subcarbonate, or protoxide.

R. Crystallized Acetate of Lead,

	five parts.
Distilled Water,	nine parts.

Dissolve by the aid of a gentle heat, and add,

Semi-vitreous Oxide of Lead, one part.

Boil, constantly stirring, till the oxide is dissolved and the liquor marks thirty degrees by the areometer. Let it cool, strain, and preserve it in a well-stopped bottle. (GALL.; FORM. HÔP. FR.)

SAX. prescribes one part of the acetate, from twenty to twenty-five of water, and one and a half of the oxide; to be boiled together till the liquor renders the syrup of violets green, and a phial capable of containing one ounce of water contains one ounce and a half of it.—HANN.; and NIEMANN—five of the oxide and eight of the acetate to be dissolved in forty of water, and boiled down to twenty-four, when the specific gravity of the liquor will be 1.420.—VAN MONS—one hundred seven and a half of the oxide, one hundred eighty-four and a half of the acetate, and two and a half of water; to be boiled together till the liquor marks 48 or 29 degrees.—BAVAR.—one of oxide and three of acetate to be boiled in nine of water till the liquor weighs from 1.360 to 1.365.

There enters thrice as much oxide, for the same quantity of acid, in the sub-acetate as in the neutral acetate.

With regard to the semi-vitreous oxide, which is preferable to the deutoxide as well as to the subcarbonate. THENARD observes that it should be calcined before it is put into re-action with the neutral acetate, in order to destroy the carbonate of lead which it always contains in a greater or less quantity, and which the neutral acetate would not combine with.

Several pharmacopœias indicate solutions of subacetate of lead of two different densities; the weaker is called *Lithargyrated* or *Saturnine Vinegar*; *Acetum Plumbi* seu *Lithargyri* seu *Saturni* seu *Lithargyrites* seu *saturninum*, and the more concentrated GOULARD's *Extract*, *Extract of Lead*, *Extractum Saturni* seu *Plumbi* seu *Goulardi*, *Balsamum Saturni*, *Acetum Plumbi concentratum*. The concentration of the latter generally varies between 27 and 30 degrees

of REAUMUR's areometer.—GALL.; and FORM. HÔP. FR. require it at 30 degrees. As the salt can be obtained crystallized, it would be well if the solution, from its constant variability, were entirely suppressed as an official preparation.

AUSTR.; and CAST. BORUS. give the name of extract of lead or of saturnine vinegar to a solution of the neutral acetate in water. This is an abuse, or rather a reprehensible error.

The medicinal properties and chemical incompatibilities of the subacetate of lead are the same as those of the following salt.

I. *Preparations in which the salt has watery liquids for excipient.*

OPHTHALMIC LIQUOR.

Collyrium plumbatum, Aqua saturnina seu ophthalmica. (AMER.; FULD.; HAMB.; BORUS.; COXE; DISP. CHA. PARIS; HUFELAND; RATIER; SWEDIAUR)

R. Solution of Subacetate of Lead,
half an ounce.
Distilled Water, two pounds.
Mix. (BORUS.)

AMER.; and COXE prescribe a scruple of solution and half a pound of water.—DISP. CHA. PARIS—two drops of solution and one ounce of water.—RATIER—ten drops of solution and four ounces of water.

R. Solution of Subacetate of Lead,
ten drops.
Decoction of Marshmallows,
two ounces.

Mix. (HAMB.)

R. Mucilage of Quince, one ounce.
Distilled Water, three ounces.
Solution of Subacetate of Lead,
twelve drops.

Mix. (FULD.)

HUFELAND prescribes two drops of solution, one ounce of rose water, and half a dram of mucilage of quince.

R. Flowers of Melilot, half a dram.
Water, one pound.
Infuse, and add when strained,
Solution of Subacetate of Lead,
one dram.

Mix. (RATIER)

R. Solution of Subacetate of Lead,
eight drops.
Rose Water, four ounces.

Mix. (SWEDIAUR)

SATURNINE COLLUTORIUM. (FORM. HÔP. FR.)

R. Solution of Subacetate of Lead,
one part.

Despumated Honey, twenty parts.
Mix.

ASTRINGENT INJECTION.

Injectio ad Leucorrhæam. (AUGUSTIN; CADET DE GASSICOURT; SWEDIAUR; PH. SYPH.)

R. Solution of Subacetate of Lead,
two drams.
Distilled Vingar, half a pound.
Rose Water, one pound and a half.

Mix.—It is recommended by YOUNG, in leucorrhœa. (CADET DE GASSICOURT; SWEDIAUR; PH. SYPH.)

R. Solution of Subacetate of Lead,
from nine to eighteen drops.
Water, nine ounces.
Opium,

from twelve to twenty-four grains.

Mix.—It is recommended by GIRTANNER in painful gonorrhœa. (AUGUSTIN)

SEDATIVE INJECTION. (CADET DE GASSICOURT; SWEDIAUR; PH. SYPH.)

R. Aqueous Extract of Opium,
from one to three drams.
Warm Water, one pound.
Dissolve, and add,

Solution of Subacetate of Lead,
from one to three drams.

Mix. (CADET DE GASSICOURT; SWEDIAUR)

PH. SYPH. prescribes one dram of extract of opium, two pounds of rose water, and sixty drops of solution; or, three drams of opium, twelve ounces of distilled water, and from one to three drams of solution.

It was extolled by HAMILTON, especially in gonorrhœa.

OPIATE SATURNINE COLLYRIUM. (AMER.; WURCEB.; COXE; ELLIS)

R. Solution of Subacetate of Lead,
two drops.
Rose Water, one ounce.
Aqueous Extract of Opium, one grain.
Mix. (WURCEB.)

R. Solution of Subacetate of Lead,
one scruple.
Water, half a pound.
Tincture of Opium, one dram.

Mix. (AMER.; and COXE)

R. Solution of Subacetate of Lead,
twelve drops.
Wine of Opium, forty drops.
Rose Water, three ounces.

Mix. (ELLIS)

When the diluted solution is prepared with common water, it is lactescent from a part of the salt becoming decomposed, producing subcarbonate or sulphate of lead, which is precipitated, and acetate of lime which remains in solution. But this is of no importance medicinally, except, that it slightly diminishes the strength of the liquor; but the effect of this result may be obviated, by shaking it immediately before employing it, in order to mix the precipitate which has formed at the bottom.

The diluted solution of subacetate of lead is employed in fomentations and lotions, and to moisten cataplasms.

SATURNINE FOMENTATION.

Fomentatio refrigerans, Lotio saturnina seu plumbata. (FULD.; HAMB.; CADET DE GASSICOURT; PIERQUIN; SWEDIAUR; PH. SYPH.)

R. Diluted Solution of Subacetate of Lead, one pound and a half.
Alcohol, three ounces.

Mix. (HAMB.)

R. Solution of Subacetate of Lead, } of each,
Proof Spirit, } one ounce.
Rose Water, three pounds.

Mix. (CADET DE GASSICOURT; PIERQUIN)

SWEDIAUR prescribes from two to four drams of solution, four of alcohol, and two pounds of water.—PH. SYPH.—from two to four drams of solution, one ounce of alcohol, and two pounds of water.

Three or four drams of spirit of camphor may be added.

R. Solution of Subacetate of Lead, } of each,
Proof Spirit, } half an ounce.
Plantain Water, one pound.

Mix. (PIERQUIN)

R. Diluted Solution of Subacetate of Lead, two pounds.
Alcohol, three ounces.
Sulphuric Acid, one ounce.

Mix. (FULD.)

RESOLVENT CATAPLASM. (FORM. HÔP. FR.; RATIER; SPIELMANN; PH. SYPH.)

R. Resolvent Meals, one part.
Diluted Solution of Subacetate of Lead, eight parts.

Boil to the consistence of soft paste. (SPIELMANN)

LIQUOR AGAINST CHILBLAINS.

Liquor ad Perniones. (SWEDIAUR)

R. Solution of Subacetate of Lead, one part.

Spirit of Rosemary, four parts.
Mix.

RESOLVENT FOMENTATION. (FULD.)

R. Muriate of Ammonia, half an ounce.
Water, twelve ounces.

Dissolve, and add

Alcohol, two ounces.
Solution of Subacetate of Lead, two drams.

Mix.

SATURNINE COLLYRIUM. (AUGUSTIN; CADET DE GASSICOURT)

R. Solution of Subacetate of Lead, } of each,
Spirit of Camphor, } six drops.
Mucilage of Tragacanth, half an ounce.
Plantain Water, six ounces.

Mix. (CADET DE GASSICOURT)

R. Diluted Solution of Subacetate of Lead, } of each,
Rose Water, } two ounces.
Opium, eight grains.

Mix. (AUGUSTIN)

R. Solution of Subacetate of Lead, two drams.
Infusion of Elder, six ounces.
Spirit of Camphor, one dram.

Mix. (AUGUSTIN)

COROBORANT LOTION. (CADET DE GASSICOURT)

R. Solution of Subacetate of Lead, } of each,
Spirit of Camphor, } two drams.
Sulphate of Zinc, one dram.
Chamomile Water, half a pound.

Mix.—It is recommended by SELLE, in atonic ophthalmia.

ASTRINGENT FOMENTATION.

Fomentum saturninum cum opio. (AUGUSTIN)

R. Water, ten ounces.
Extract of Opium, ten grains.
Solution of Subacetate of Lead, half a dram.

Spirit of Camphor, twenty drops.

Mix.—VOGEL recommends it in strangulated hernia.

III. Preparations in which the Subacetate of Lead has fatty substances for the excipient.

SATURNINE BOUGIES.

Cereoli saturnini seu plumbici, Candela chirurgica seu resolventes seu mitigantes, Cerei medicati. (BRUNS.; FULD.; POL.;

BORUS.; ROSS.; SAX.; SUEC.; BRUGNATELLI; PIDERIT; PIERQUIN; SPIELMANN; SWEDIAUR)

- R. Yellow Wax, two pounds.
Melt over a slow fire, and add
Solution of Subacetate of Lead,
one ounce.

Mix with constant stirring; withdraw the mass from the fire, and form bougies. (BRUNS.; FULD.; POL.; BORUS.; ROSS.; SAX.; BRUGNATELLI; PIDERIT; PIERQUIN; SPIELMANN; SWEDIAUR)

- R. Yellow Wax, one pound.
Spermaceti, three drams.
Solution of Subacetate of Lead,
two drams.

Add the solution to the two first melted together. (SUEC.)

OINTMENT OF SUBACETATE OF LEAD.

Balsamum universale seu saturninum, Butyrum Saturni, Ceratum Saturni seu Goulardi seu Lithargyri seu Plumbi Superacetatis seu Plumbi seu e Plumbo, Linimentum plumbatum, Pomatum Saturni, Unguentum Acetatis Plumbi seu Acetatis superplumbici seu Lithargyri seu de Lithargyro seu nutritum seu tripharmacum seu saturninum seu Subacetatis plumbici seu Oxydi Plumbi acetatum. (AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; CAST. BORUS.; BORUS.; SARD.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BRUGNATELLI; CADET DE GASSICOURT; PIDERIT; PIERQUIN; SAINTE-MARIE; SPIELMANN; SWEDIAUR; PH. SYPH.; VAN MONS)

- R. Olive Oil, two parts.
Solution of Subacetate of Lead,
one part.

Triturate together. (AMST.)

HISP.; and SWEDIAUR prescribe an equal part of each.—FULD.; LIPP.; and PIDERIT—one and a half of oil and one of solution.—PIERQUIN—eight of oil and one of solution.

- R. Lard, four parts.
Solution of Subacetate of Lead,
one part.

Triturate together. (SAX.)

- R. Olive Oil, }
Lard, } of each,
Solution of Subacetate }
of Lead, } one part.

Triturate together in a glass mortar. (BELG.)

- R. Rose Oil, one part.
Solution of Subacetate of Lead,
two parts.

Triturate together. (WIRTEM.)

- R. Semi-vitreous Oxide of Lead, one part.
Vinegar, two parts.
Rose Oil, four parts.

Triturate the oxide in a leaden mortar, adding alternately the oil and vinegar by small portions, till the whole is well incorporated. (BRUNS.)

- R. Semi-vitreous Oxide of }
Lead, } of each,
Subcarbonate of Lead, } three ounces.
Vinegar, eight ounces.
Oil, one pound and a half.

Triturate together, adding the vinegar and oil alternately till they are absorbed. (SARD.)

WIRTEM. prescribes one half part of the oxide, one and a half of the subcarbonate, one half pint of vinegar, and one of rose oil.—PALAT.—two of the oxide, one half of the subcarbonate, one of vinegar, and two and a half of rose oil.

- R. Semi-vitreous Oxide of Lead, one pound.
Olive Oil, three pounds.
Vinegar, sixteen ounces.

Boil the oxide with one pound of the oil, and water gradually till the mass has acquired the consistence of cerate; then triturate it with the rest of the oil and with the vinegar. (HISP.)

SPIELMANN prescribes sixteen parts of the oxide, eighteen of oil, and three of vinegar.—SAINTE-MARIE—one of oil, two of vinegar, and two of the oxide, to be boiled till reduced to one half.

- R. Olive Oil, four parts.
White Wax, one half part.

Melt together, and add gradually when the mass begins to cool, constantly stirring,

Solution of Subacetate of Lead,
one part.

Mix well. (OLDEN.; BRUGNATELLI)

DAN.; FULD.; LIPP.; and VAN MONS prescribe one part of solution, two parts of wax, and six of oil.—HERBIP.—one of solution, one and a half of wax, and four of oil.—PH. SYPH.—one of solution, six of wax, and sixteen of oil.—SWEDIAUR—one each of solution and of wax, and three of oil.—SPIELMANN—one of solution, six of wax, and twenty-four of oil; or four of solution, eight of wax, and eighteen of oil; or, one half of solution, nine of wax, and eight of oil.—FENN. and SUEC.—four of solution, four of wax, and nine of oil.—HASS.—two

and a half of solution, four of wax, and twelve of oil.—ROSS.—four of solution, five of wax, and nine of oil.—LUSIT.; and BRUGNATELLI—two and a half of solution, four of wax, and nine of oil.

- R. Lard, one part.
White Wax, one half part.

Melt together. Add when nearly cold,

Solution of Subacetate of Lead,
one part.

Triturate together. (AUSTR.)

CAST. BORUS. prescribes two parts of solution, one part of wax, and three of lard.

- R. Walnut Oil, twelve parts.
Yellow Wax, four parts.

Melt over a slow fire, and add gradually

Spring Water, twelve parts.
Solution of Subacetate of Lead,
one part.

Mix. (DAN.; PIDERIT)

POL.; and VAN MONS prescribe one part of solution, two parts of water, one of wax, and four of oil.—BORUS.—one of solution, two of rose water, two of wax, and eight of oil.—WIRTEM.—one of solution, twenty-four of water, one of wax, and four of oil. HANN.; and OLDEN.—one of solution, four of rose water, two of wax, and eight of oil.—HANN.—two and a half each of solution, of water, and of wax, and eight of oil.—SAX.—one half of extract, three and a half of water, one of wax, and four of oil.

- R. Wax, } of each, two parts.
Mutton Suet, }
Olive Oil, one part.

Melt over a slow fire, and add when nearly cold

Solution of Subacetate of Lead,
one part.

Triturate well together. (PIDERIT)

- R. Cerate,
one hundred and twenty-five parts.
Solution of Subacetate of Lead,
one part.

Mix carefully in a mortar. (GALL.)

CADET DE GASSICOURT; and PIERQUIN prescribe one part of solution and ninety-six of cerate.—FORM. HÔP. FR.—one of solution and twenty-five of cerate.—BATAV.—one of solution and twenty of cerate.—BAVAR.—one of solution and ten of cerate.—ANT.; MIL. COPEN.; and SWEDIAUR—one of solution and eight of cerate.—WURCEB.—one of solution and four of cerate.—FERR.—two pounds of olive oil, three ounces of white wax, and six ounces of solution.

These preparations only differ from each other in their degree of consistence; they ought not therefore to be separated. They are all considered sedative, soothing, resolvent,

and healing. They are in rather extensive use in surgical practice; indeed they are used perhaps more frequently than is necessary, though they have lost much of the reputation they possessed when the treatment of wounds and ulcers consisted nearly entirely in the application of ointments and plasters.

ANTI-HÆMORRHOIDAL OINTMENT.

Unguentum ad Varices Ani. (HAMB.; SWEDIAUR)

- R. Spermaceti Ointment, eight parts.
Solution of Subacetate of Lead,
one part.

Triturate together. (SWEDIAUR)

- R. Ointment of Toad-flax, } of each,
Mucilage of Quince Seeds, } two parts.
Solution of Subacetate of Lead,
one part.

Triturate together. (HAMB.)

OINTMENT AGAINST BURNS. (FULD.)

- R. Ointment of Subacetate of Lead,
one ounce.
Resin, } of each,
Compound Ointment of } one dram.
Elemi,

Mix.

KIRKLAND'S DESICCATIVE CERATE. (CADET DE GASSICOURT; PIERQUIN)

- R. Prepared Chalk, } of each,
Distilled Vinegar, } four ounces.

After the cessation of the effervescence, add

Olive Oil, four ounces.
Simple Plaster, half a pound.

Melt over a slow fire, and incorporate with the mass when half cold,

Solution of Subacetate of Lead,
fourteen ounces.

Mix.

OPIATE OINTMENT OF SUBACETATE OF LEAD. (RATIER)

- R. Solution of Subacetate of Lead,
half an ounce.
SYDENHAM'S Laudanum, two drams.
Oil, two ounces.
Shake well together.

CAMPHORATED OINTMENT OF SUBACETATE OF LEAD.

Baume de Saturne de Goulard, Pommade anti-herpétique de Goulard. (LOND.; AMER.; AMST.; BRUNS.; FULD.; LUSIT.; HERBIP.; COXE; PIERQUIN; SPIELMANN; SWEDIAUR)

Ceratum Plumbi compositum. (LOND.)

- R. Solution of Subacetate of Lead,
two fluid ounces and a half.
Yellow Wax, four ounces.
Olive Oil, nine fluid ounces.
Camphor, half a dram.

Melt the wax, and mix it with eight ounces of the oil; then take the mixture from the fire, and as soon as it begins to thicken, add gradually the solution, stirring assiduously with a wooden spatula till cold: finally add the camphor dissolved in the remainder of the oil.

The other pharmacopœias prepare it by adding camphor to the simple ointment of the subacetate at the moment of its becoming cold.

AMST. requires one thirty-third of camphor triturated with a little oil.—HERBIP.—one fifty-fourth.—FULD.—one sixty-sixth.—BRUNS.—one one hundred and sixty-eighth.—SPIELMANN—one two hundred and tenth.—PIERQUIN; and SPIELMANN—one two hundred and fortieth.—AMER.; LUSIT.; and COXE—one two hundred and forty-eighth.—SPIELMANN—one eight hundred and fortieth.

The enormous diversity which reigns among the relative proportions in these formulæ, shows how much better it would be, were it left entirely to the judgment of the practitioner prescribing it,

ANTI-HERPETIC OINTMENT. (SAINTE-MARIE)

- R. Solution of Subacetate of } of each,
Lead, } one
Unguentum Populeum, } ounce.
Camphor, } two drams.

Mix.—To be applied to inflamed tetter.

ANTI-HYSTERIC LINIMENT. (SAINTE-MARIE)

- R. Rose Oil, two ounces.
Unguentum Populeum, four drams.
Camphor, two drams.
Solution of Subacetate of Lead, one dram.

Mix.—To be rubbed on the abdomen every four hours.

GOULARD'S SPARADRAP.

Peau de Goulard. (PIERQUIN; SPIELMANN; VAN MONS)

- R. Olive Oil, fourteen ounces.

Yellow Wax, twenty-four ounces.
Melt together, and add

Solution of Subacetate of Lead, one ounce.
Camphor, triturated with a little oil, half a dram.

Dip a moderately fine cloth into the mixture, and smooth its surface. (VAN MONS)

PIERQUIN prescribes twelve pounds of wax, three of oil, eight ounces of solution, and two drams of camphor.—SPIELMANN—ninety-six ounces of wax, twenty-eight of oil, four of solution, and one dram of camphor.

This sparadrap has been extolled as an application to atonic ulcers of the leg, as also in cases of rheumatic pains.

SAPONACEOUS OINTMENT OF SUBACETATE OF LEAD.

Unguentum Saturni saponaceum; Pommade ou Onguent de Goulard. (BRUNS.; FULD.; LUSIT.; PIERQUIN)

- R. Common Soap, eight ounces.
Spring Water, two pounds.

Dissolve over a slow fire, constantly stirring; then add

Solution of Subacetate of Lead, two ounces.
Camphor, one dram.

Mix. (BRUNS.; PIERQUIN)

- R. Unguentum Nutritum, eight ounces.
White Soap, four ounces.

Boil slowly, to evaporate the humidity, add

Camphor, one dram.

Mix. (FULD.)

LUSIT. directs three ounces of soap to be added to its simple ointment of subacetate of lead.

This ointment is reputed more resolvent and discutient than the simple. It has been extolled particularly in ankylosis.

SATURNINE BOUGIES. (FERR.)

- R. Yellow Wax, one pound.
Solution of Subacetate of Lead, half an ounce.

Boil together till the liquor is evaporated, and make bougies with strips of linen dipped in the mixture.

PROTO-ACETATE OF LEAD.

Acetas Plumbi seu Plumbi acidulus siccus, Acetas plumbicum crystallisatum, Acetis Plumbi, Cerussa acetata, Plumbum aceticum seu acetatum, Superacetas Plumbi, Saccharum saturninum, Sal Saturni, Sal Plumbi.—Acetate of Lead, Neutral Acetate of Lead, Salt of Lead, Sugar of Lead.

SYNONIMES:—Acetate de protoxide de plomb, Acétate de plomb, Acetate de plomb neutre, Sel de saturne, Sucre de plomb, Sucre de saturne, FR.; Bleizucker, Essigsäures Bley, Bleysalz, GERM.; Blyesukker, DAN.; Lootzuiker, DUT.; Zuccherò di saturno, ITAL.; Cukior Ołowiany, Przyocian ołowin nieczysty, POL.; Blysocker, SWED.

I. *That which is found in Commerce.*

BATAV.; BAVAR.; BELG.; MIL. COPEN.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; SAX.; SUEC.; WURCEB.; COXE; GUIBOURT; PARIS; PIDERIT.

Before this salt is used it should be purified:

R. Acetate of Lead, two parts.
Boiling Water, three parts.

Add a little concentrated acetic acid to the solution to render it perfect; filter it while hot, and set it apart in a cool place that crystals may form. (BAVAR.)

FENN. prescribes one part of acetate, four parts of water, and one of distilled vinegar.—SAX. directs three parts of acetate to be dissolved in three of water, and one part of vinegar to be added; the mixture to be then boiled, strained, and set aside to crystallize.

R. Acetate of Lead, } of each,
Water, } one part.

Dissolve, strain the solution through a linen cloth, add one part of water, apply heat, filter, and set aside to crystallize slowly. (HANN.; HASS.; OLDEN.; BORUS.; SUEC.; SWEDIAUR; VAN MONS)

The salt may also be dissolved in the smallest possible quantity of water, and a current of carbonic acid gas passed into the solution till precipitate ceases to be formed. (VAN MONS)

II. *Manufactured according to various Formule.*

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BRUNS.; DAN.; HISP.; GALL.; FULD.; GENEV.; HASS.; LUSIT.; PALAT.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS.

Plumbi Acetas.

R. Subcarbonate of Lead,
one pound.

Strong Acetic Acid, one pint.
Boiling Distilled Water,
one pint and a half.

Mix the acid with the water, and add the subcarbonate gradually; then filter the solution through paper, and evaporate till a pellicle appears on its surface. Set it then apart that crystals may form; pour off the fluid, and dry the crystals on bibulous paper. (LOND.)

R. Carbonate of Lead, called
Cerussa, any quantity.
Distilled Vinegar,
ten times its weight.

Digest in a glass vessel till the vinegar has become sweet, and having poured this off, add more till it no longer comes off sweet. Filter the liquor, and crystallize by alternately evaporating and cooling it: dry the crystals in the shade. (DUBL.)

Acetas Plumbi. (EDINB.)

R. White Oxide of Lead,
any quantity.
Weaker Acetic Acid,
a sufficient quantity.

Put the oxide into a cucurbit, and pour upon it ten times its weight of the acid. Stand the mixture in a warm sand-bath till the acid becomes sweet; then pour it off and add fresh portions of acid successively till it ceases to acquire any sweetness. Evaporate the mixed liquors, freed from impurities, in a glass vessel to the consistence of thin honey, and set it aside in a cool place that crystals may form, which should be dried in the shade. Evaporate the residue again that more crystals may be obtained, repeating the evaporation till they cease to be formed.

R. Subcarbonate of Lead, one part.
Distilled Vinegar, ten parts.

Digest in a glass vessel till the acid is saturated. Decant and filter the liquor, set it apart to crystallize, and dry the crystals. (AMER.; AMST.; GENEV.; HASS.; LUSIT.; HERBIP; COXE; PIDERIT; VAN MONS)

The operatory process is the same in the other pharmacopœias; there are a few slight differences in the respective proportions of the subcarbonate and of the acid, similar to those which the preceding present, but they are unnecessary to be particularized: but PALAT. prescribes the oxide of lead obtained by the calcination of the metal at the open fire.—AUSTR.; DAN.; and FERR. require the deutoxide.—HISP.; and VAN MONS allow either the subcarbonate or the semi-vitreous oxide.—FULD.; SPIELMANN; and SWEDIAUR—either the subcarbonate or the deutoxide.—WIRTEM.—either the deutoxide or the semi-vitreous oxide.—BRUNS.; and GALL. prescribe the semi-vitreous oxide.

ANT.; SARD.; and BRUGNATELLI simply require the solution of the subacetate to be evaporated slowly, and set aside in a cool place to crystallize.

It is considered astringent, desiccative, and repperussive, and is employed externally to dissipate or resolve certain phlegmonous and erysipelatous inflammations. It is also given internally to diminish the colliquative sweats of consumptive patients, to arrest certain diarrhœas, in passive hæmorrhage, and in various cases of chronic catarrh.

It is incompatible with lime water, the alkalis and their subcarbonates, the sulphuric and hydro-chloric acids, the alkaline soaps, alum, the sulphates of iron, of copper, and of zinc, borax, gum, and the greater number of animal substances.

A.

Preparations into which the salt enters in the solid state.

TEMPERANT POWDER. (PIERQUIN)

- R. Acetate of Lead, one dram.
Nitrate of Potass, one ounce.

Make a powder.—Dose, from two scruples to a dram, in a glass of water, two or three times a week.

WIRTEM. gives this preparation, under the name of *Nitrum saturninum*, with the following formula.

- R. Nitrate of Potass, three ounces.
Acetate of Lead, two ounces.

Dissolve them in a sufficient quantity of rose water; then evaporate it to the consistence of thin paste, and finally dry it completely with a moderate heat.

This saturnine nitre used formerly to be employed externally, in collyriums, gargles, and injections.

ANTI-PHTHISICAL POWDER. (BORIES)

- R. Acetate of Lead, } of each, six grains.
Opium, }
Sugar, one scruple.

Mix, and divide into twelve parcels.—Dose, one, morning and evening, in the colliquative sweats and diarrhœas of consumptive patients.

STYPTIC POWDER. (CADET DE GASSI-COURT)

- R. Saturated Solution of Hydrochlorate of Iron, four ounces.
Evaporate it to dryness, and add to it,
Acetate of Lead, an equal weight.

Continue the evaporation, at a moderate heat, to dryness; and pulverize and sift the residuum.

It is employed internally and externally, in hæmorrhages.—Dose, internally, from four to twelve grains.

ANTI-PSORIC POWDER. (RATIER)

- R. Acetate of Lead, } of each, two parts.
Sublimed Sulphur, }
Sulphate of Zinc, one part.

Mix.—CHAUSSIER has proposed this powder. A pinch of it is to be mixed with a few drops of oil, and rubbed upon the palms of the hands.

ESCHAROTIC POWDER. (PIERQUIN)

- R. Acetate of Lead, } of each,
Subcarbonate of Lead, } one pound.
Alum, }

Mix.—It is applied to quicken the cicatrization of atonic ulcers.

PILLS OF ACETATE OF LEAD. (PIERQUIN; RATIER)

- R. Acetate of Lead, } of each,
Powdered Marshmallows, } one dram.
Simple Syrup, a sufficient quantity

to make thirty-six pills.—Dose, from four to twelve daily, to moderate the sweats of consumptive patients. Eight or ten are usually sufficient. They should be taken at the time when the sweating is expected to come on.

ANTI-PHLOGISTIC PILLS. (PIERQUIN)

- R. Acetate of Lead, six grains.
Camphor, } of each,
Nitrate of Potass, } three grains.
Conserve of Roses, a sufficient quantity

to make six pills.—To be taken in the course of the day, against the excessive sweats of consumptive patients and in certain inflammations of the chest.

STYPTIC PILLS. (ELLIS; RATIER)

- R. Acetate of Lead, four grains.
 Purified Opium, one grain.
 Extract of Liquorice, one scruple.

Make sixteen pills.—DUMÉRIEUX employs them as astringent in chronic and colliquative diarrhoea. (RATIER)

- R. Acetate of Lead, twelve grains.
 Opium, six grains.
 Conserve of Roses, a sufficient quantity.

Make twelve pills.—They are useful in hæmoptysis and other hæmorrhages. (ELLIS)

PILLS OF ACETATE OF LEAD AND OF IPECACUANHA. (NIEMANN)

- R. Acetate of Lead, } of each, one dram.
 Ipecacuanha, }
 Opium, ten grains.
 Simple Syrup, a sufficient quantity

to make forty pills.

THACHER praises them in passive uterine hæmorrhages.—One to be taken every three hours.

B.

Preparations in which the Acetate of Lead has fatty substances for the excipient.

SATURNINE CERATE.

Ceratum saturninum seu Goulardi. (GENÈVE.)

- R. Acetate of Lead, half an ounce.
 GALEN'S Cerate, twelve ounces.
 Triturate together.

SATURNINE BALSAM.

Oleum seu Balsamum Saturni. (HASS.; CADET DE GASSICOURT; PIDERIT; PIERQUIN; VAN MONS)

- R. Spirit of Turpentine, any quantity.
 Heat it in a glass vessel, and add by portions,

Powdered Acetate of Lead, a sufficient quantity, constantly stirring, till no more of it is dissolved; then set it apart and decant the liquor while hot. (VAN MONS)

HASS.; CADET DE GASSICOURT; PIDERIT; and PIERQUIN direct eight ounces of spirit of turpentine, to be poured upon four ounces of dried acetate, and left to digest in a sand-bath for twenty-four hours.

It is stimulant, and is extolled as an application to spreading and painful ulcers, as well as to gangrene; but it should not be employed in such cases but with great circumspection.

Vol. II.

ANTI-HERPETIC OINTMENT. (PIERQUIN)

- R. RHAZES' White Trochiscs, half an ounce.
 Acetate of Lead, } of each,
 Proto-chloruret of } half a dram.
 Mercury,
 Lard, a sufficient quantity,
 to form an ointment. (PIERQUIN)

SATURNINE OINTMENT. (BORIES)

- R. Acetate of Lead, one ounce.
 Distilled Vinegar, eight ounces.

Dissolve, evaporate to dryness, pulverize, and add

- Lard, two ounces.
 Extract of Hemlock, two drams.

Mix.—BARTHEZ calls it discutient and resolvent.

ANTI-HÆMORRHOIDAL OINTMENT. (PIERQUIN)

- R. Ointment of Snails, one ounce.
 Subcarbonate of Lead, } of each,
 Tincture of Opium, } half a dram.
 Acetate of Lead, two drams.
 Powdered Marshmallows Root,
 a sufficient quantity.

Mix.

RESOLVENT OINTMENT. (BORIES; PIERQUIN; SAINTE-MARIE)

- R. Acetate of Lead, three drams.
 Camphor, two drams.
 White Wax, one ounce.
 Olive Oil, four ounces.

Mix.—BARTHEZ recommends it for scorbutic ulcers.

CERATE OF ACETATE OF LEAD. (LOND.; DAN.; COXE; SWEDIAUR)

Ceratum Plumbi Acetatis. (LOND.; COXE)

- R. Acetate of Lead, in powder, two drams.
 White Wax, two ounces.
 Olive Oil, half a pint.

Melt the wax in seven ounces of the oil; then gradually add the acetate triturated with the remainder of the oil, and stir with a wooden spatula till thoroughly incorporated.

DAN. prescribes one ounce and a half of the acetate, twelve ounces of oil, and three of wax.—SWEDIAUR—one dram of acetate, eight ounces of oil, and one ounce and a half of wax.

ANTI-HÆMORRHOIDAL OINTMENT. (BORIES)

- R. Acetate of Lead, fifteen grains.
 Burnt Cork, in powder, half an ounce.
 Fresh Butter, two ounces.
- Triturate together.

OPHTHALMIC OINTMENT.

Unguentum oculare. (SPIELMANN)

- R. Acetate of Lead, one scruple.
 Spermaceti Ointment, five drams.
 Compound Tincture of Benzoin, two scruples.

Mix.

POTT'S CERATE.

Ceratum saponaceum. (CADET DE GASSI-COURT; SWEDIAUR; VAN MONS)

- R. Acetate of Lead, in powder, four parts.
 White Soap, scraped, one part and a half.

Mix intimately in a mortar heated in a water-bath; after the evaporation of the water, add,

White Wax, } of each,
 Olive Oil, } three parts,

previously melted together, and mix. (VAN MONS)

- R. Semi-vitreous Oxide of Lead, one pound.
 Vinegar, eight pounds.
 Soap, half a pound.

Boil over a slow fire till all humidity is dissipated; then add,

Olive Oil, } of each, one pound.
 Yellow Wax, }

Mix by trituration. (CADET DE GASSI-COURT; SWEDIAUR)

It is desiccative and is employed in the treatment of ulcers. It does not differ in its mode of action, from the ointment of subcarbonate of lead.

RESOLVENT SPARADRAP. (PIERQUIN)

- R. White Wax, eight ounces.
 Olive Oil, one pound.

Melt together, and add,

Camphor, } of each,
 Muriate of Ammonia, } one dram.
 Acetate of Lead, one ounce.

Dip strips of linen into the mixture.

SEDATIVE OINTMENT. (PIERQUIN)

- R. Sagapenum, } of each, six drams.
 Gum Ammoniac, }
 Turpentine, } of each, eight ounces.
 Resin, }

Proof Spirit, a sufficient quantity.
 Dissolve, and add,

Acetate of Lead, } of each,
 Subcarbonate of } one ounce and
 Lead, } a half.
 Olibanum, ten drams.
 Euphorbium, three drams.
 Lard, one pound.
 Oil of Lily, } of each,
 ——— Lavender, } a sufficient
 White Wax, } quantity.

Make an ointment.—This is stimulant, and was formerly considered resolvent, but is now disused.

RESOLVENT PLASTER. (BORIES; PIERQUIN)

- R. Acetate of Lead, one dram.
 Sublimed Sulphur, half an ounce.
 White Resin, two ounces.

Make a plaster.—It is applied to syphilitic buboes, and is renewed every two days.

C.

Preparations in which the Acetate of Lead has liquids for the excipient.

AQUEOUS SOLUTION OF ACETATE OF LEAD.

Acetas Plumbi solutus seu liquidus, Aquacum Saccharo Saturni. (AUSTR.; BAVAR.; MIL. COPEN.; HAMB.; LIPP.; CAST. BORUS.; RATIER)

- R. Acetate of Lead, one part.
 Distilled Water, six parts.

Dissolve, and filter. (AUSTR.)

AUSTR. gives this solution, very improperly, under the name of *Acetum Lithargyri*.—CAST. BORUS.—calls a solution of one part of the salt in three parts of water, by the name of *Extractum Saturni*.—The *Aqua Saturnina* of MIL. COPEN. is a solution of one part of the salt in ninety-six parts of water.

BAVAR. prescribes one part of the salt and nine parts of water.—HAMB.; and LIPP.—one of the salt and forty-eight of water.

RATIER gives a *Resolvent Injection* composed of one part of the salt, and from twenty-four to forty-eight of water.

OPIATE AQUEOUS SOLUTION OF ACETATE OF LEAD. (AUGUSTIN)

- R. Acetate of Lead, two grains.
 Tincture of Opium, half a dram.
 Water, six ounces.

Mix.—Dose, a spoonful every three hours, against the immoderate sweats of consumptive patients.

ASTRINGENT TINCTURE.

Tinctura Saturnino-martialis. (AUGUSTIN)

- R. Acetate of Lead, one ounce.
 Sulphate of Iron, half an ounce.
 Alcohol, eight ounces.

Mix.—Dose, ten drops to children, twenty to adults; in nasal hæmorrhage.

OPHTHALMIC LIQUOR. (NIEMANN)

- R. Acetate of Lead, fifteen grains.
 Rose Water, six ounces.

Dissolve.

ANTISEPTIC LOTION. (PIERQUIN)

- R. Acetate of Lead, one dram.
 Plantain Water, two ounces.

Dissolve.

RESOLVENT COLLYRIUM. (GENEV.; LIPP.; AUGUSTIN; PIERQUIN; RATIER)

- R. Acetate of Lead, half a scruple.
 Water, two ounces.

Dissolve. (GENEV.)

- R. Acetate of Lead, six grains.
 Infusion of Elder Flowers, four ounces.

Dissolve, and add,

Vulnery Spirit, two drams.

Mix. (RATIER)

- R. Acetate of Lead, dissolved in water, three parts.
 Mucilage of Quince, one part.

Mix. (LIPP.)

- R. Acetate of Lead, half a dram.
 Plantain Water, } of each,
 Rose Water, } two ounces.

Mix. (PIERQUIN)

- R. Acetate of Lead, three grains.
 Opium, four grains.
 Quince Seeds, six grains.
 Rose Water, two or three ounces.

Mix. (AUGUSTIN)

- R. Acetate of Lead, ten grains.
 Alum, one dram.
 Rose Water, four ounces.

Mix. (AUGUSTIN)

- R. Acetate of Lead, three grains.
 Quince Seeds, six grains.
 Rose Water, } of each, two ounces.
 Fennel Water, }
 SYDENHAM'S Laudanum, one dram and a half.

Mix. (AUGUSTIN)

SEDATIVE JULEP. (PIERQUIN)

- R. Acetate of Lead, eight grains.
 Water of Bugloss, } of each,
 ——— Water-Lily, } three ounces.

Syrup of Poppy, one ounce.

Mix.

SEDATIVE GARGLE. (PIERQUIN)

- R. Acetate of Lead, sixty grains.
 Aqueous Extract of Opium, six grains.
 Rose Water, eight ounces.
 Syrup of Mulberries, one ounce.

Mix.

ASTRINGENT GARGLE.

Collutorium Catechu saturninum. (AUGUSTIN)

- R. Catechu, three drams.
 Lime Water, twelve ounces.
 Acetate of Lead, one scruple.
 Honey of Roses, two ounces.

Mix.

RESOLVENT FOMENTATION.

Fomentum saturninum. (AUGUSTIN)

- R. Acetate of Lead, two drams.
 Vinegar, three ounces.
 Water, twenty-four ounces.

Mix.—It is recommended by GIRTANNER, against phymosis.

ASTRINGENT GARGLE.

Gargarisma saturninum. (AUGUSTIN)

- R. Herb of Sage, half an ounce.
 Water, a sufficient quantity
 to obtain four ounces of decoction. Add
 when strained,

Alum, one dram.
 Saturnine Liquor, two ounces.

Mix.—It is recommended in mercurial salivation.

ASTRINGENT INJECTION. (AUGUSTIN)

- R. Acetate of Lead, two grains.
 Water, one ounce.

Dissolve.—It is recommended at the commencement of gonorrhœa.

- R. Acetate of Lead, one dram.
 Spirit of Turpentine, five ounces.

Mix.—A little camphor may be added. This remedy, employed in chronic gonorrhœa, was called *divine* by HOFFMANN.

- R. Acetate of Lead, one dram.
 Water, six ounces.
 Solution of Myrrh, two ounces.
 Tincture of Opium, one ounce.

Mix.—For chronic gonorrhœa.

POTION AGAINST HOOPING-COUGH. (CADET DE GASSICOURT)

- R. Acetate of Lead, from two to four grains

Rose Water, two ounces.
Syrup of Violets, two drams.

Mix.—It is recommended by ROBERT THOMAS, in the dose of a tea-spoonful, every four hours.

MYRRHOUS SATURNINE DECOCTION.
(NIEMANN)

R. Acetate of Lead, one dram.
Distilled Water, six ounces.

Dissolve, and add,

Aqueous Tincture of Myrrh, two ounces.
Tincture of Opium, one ounce.

Mix.—FRANK extolled it in chronic menorrhœa. — It may be compared with PLENK'S *Vulnerary Liquor* (BORIES), of which this is the formula.

R. Lime Water, one pound.
Burning Soot from an oven, one ounce.
Subcarbonate of Lead, half an ounce.

Boil the whole together for half an hour; then add,

Oil of Myrrh, half an ounce.

Mix.—It is probable that a little pyro-acetate of lime and pyro-acetate of lead is formed during the operation.

HEISTER'S DECOCTION.

Decoctum ad Fungos Articulorum seu ad Lupias synoviales Heisteri. (BRUNS.; SPIELMANN)

R. Semi-vitreous Oxide of Lead, six ounces.
Armenian Bole, one ounce.
Mastic, } of each, half an ounce.
Myrrh, }
Vinegar, two pounds.

Boil over a slow fire for half an hour.

It is applied as a fomentation.

BELL'S VEGETO-MINERAL WATER. (NIEMANN)

R. Acetate of Lead, half an ounce.
Vinegar, four ounces.
Distilled Water, two pounds.

Dissolve, and mix.

ASTRINGENT INJECTION. (SAUNDERS)

R. Acetate of Lead, half a dram.
Distilled Vinegar, one pound.
Rose Water, one pound and a half.

Dissolve.—To be used in uterine hæmorrhages.

SEDATIVE FOMENTATION. (PIERQUIN)

R. Acetate of Lead, } of each,
Camphor, } two ounces.

Oxyerat, six ounces.
Very cold Water, a sufficient quantity.
Dissolve and mix.

COMPOUND ALUMINOUS INJECTION. (PH. SYPH.)

R. Acetate of Lead, one dram.
Alum, half a dram.
Sulphate of Zinc, eighteen grains.
Distilled Water, sixteen ounces.

CAMPHORATED VULNERARY LIQUOR.
(HAMB.; HERBIP.; PIERQUIN; SPIELMANN)

R. Acetate of Lead, two drams.
Distilled Water, one pound.
Spirit of Camphor, two ounces.

Dissolve and mix. (HAMB.)

R. Acetate of Lead, half a dram.
Camphor, dissolved in one ounce of alcohol, three ounces.
Add gradually shaking the vessel,
Spring Water, four pounds.

Preserve. (HERBIP.; PIERQUIN; SPIELMANN)

VIRGIN MILK. (PIERQUIN)

R. Semi-vitreous Oxide of Lead, one ounce.
Subcarbonate of Lead, three drams.
Camphor, two scruples.

Dissolve these in

Vinegar, two ounces,
and add to the solution

Distilled Water of Solanum	} of each, three ounces.
Nigrum,	
————— Lettuce,	
————— Water-	
Lily,	

Mix.—To be applied as a cosmetic, three or four times a day.

SEDATIVE ELECTUARY. (PIERQUIN)

R. Acetate of Lead, half a dram.
Nitrate of Potass, six drams.
Conserve of Dog-rose, two ounces.
Syrup of Maidenhair, a sufficient quantity.

Mix.—It is extolled against salacity in both sexes.—Dose, half a dram, morning and evening, for eight days.—Laudanum and camphor may be added.

GARMANN'S ANTI-PHTHISICAL TINCTURE.
(WIRTEM.; NIEMANN)

R. Acetate of Lead, half an ounce.
Sulphate of Iron, three drams.

Vinegar, } of each, two ounces.
 Alcohol, }
 Rose Water, six drams.

Dissolve the acetate in the vinegar at a moderate heat, and add the pulverized sulphate; when the solution is effected, add the alcohol mixed with the rose water, and set it aside in a moderate temperature.

It is extolled by GARMANN, WEDEL, VOGEL, and others against the colliquative sweats and diarrhœas of consumptive patients; it has also been recommended to arrest gonorrhœa and nocturnal emissions.—Dose, from twenty to thirty drops. The compilers of WIRTEM. say of it, *caute tamen ubique, vel nunquam, adhibendum esse, censemus*. Some practitioners of the present day are far from sharing in this proper mistrust of it.

PROTO-GALLATE OF LEAD.

This salt is never prepared expressly, but it makes, at least most probably, the basis of the two following preparations

QUERCICO-SATURNINE LINIMENT.

Magna scytodepsicum ex Quercu, Linimentum saturnino-quercicum. (AUGUSTIN; NIEMANN)

R. Oak Bark, in powder, one ounce.
 Spring Water, six ounces.

Boil down to one half; instil into the strained decoction.

Solution of Subacetate of Lead,
 half an ounce.

Filter, and preserve the precipitate under alcohol, without drying it.

AUTENRIETH extolled this substance as an application to excoriations produced by long lying in bed.

ASTRINGENT LOTION.

Lotio Quercus saturnina. (AUGUSTIN)

R. Decoction of Oak Bark, } of each,
 Saturnine Liquor, } six ounces.
 Whites of Eggs, two.
 Solution of Myrrh, two ounces.

Mix.—It is recommended to excoriations which threaten gangrene.

PROTO-OLEO-MARGARATE OF LEAD.

This name, under which we place all the plasters in which an intimate combination is effected between the protoxide of lead and the acid products produced by the re-action which the elements of fatty substances mutually undergo under the influence of heat, is not perfectly exact. Indeed the greater number of the plasters in question contain,

besides oleate and margarate, stearate of lead, some of them caproate, perhaps also butyrate, and some sulphate and even acetate of lead. However it seems better that we should adopt it on account of its shortness, and because the two salts which it expresses are nearly always by far the most abundant.

The saturnine plasters are sometimes prepared with the semi-vitreous oxide, sometimes with the deutoxide, and sometimes with the subcarbonate of lead, which has determined us to rank them in three different divisions, to one or the other of which we have thought proper to refer those in which there enters two or even three of those saturnine compounds. It will be useful to bear in mind that, according to HAGEN, one part of oil is required to dissolve one of the subcarbonate, one and a half to dissolve one of the deutoxide, and two parts to dissolve one of the semi-vitreous oxide. The investigations of HENRI have shown us, 1st with respect to the fat bodies, that the oils which are naturally mucilaginous or which are rendered so artificially give plasters which are scarcely consistent, that of the olive deserves the preference over every other, and that lard does not seem to differ much from it in this respect; 2nd with regard to the oxides, that the deutoxide combines slowly and imperfectly, that the yellow oxide gives a mass without coherence, that the semi-vitreous oxide is the best adapted of all for the making of plasters, and that the English is better for use than that from Hamburg.

I. Prepared with Protoxide of Lead.

SIMPLE PLASTER.

Emplastrum commune seu diachylum seu Lithargyri seu Lithargyrii simplex seu Oxydi Plumbi semi vitrei seu Oxydi plumbici seu Plumbi.

1. Simple Diachylon Plaster; *Emplastrum diachylum simplex seu Oxydi Plumbi mucilaginem.* (BRUNS.; HASS.; LIPP.; PALAT.; SARD.; WIRTEM.; HERBIP.; SPIELMANN)

R. Mucilage of Fenugreek, }
 Linseed, } of each.
 Marshmal- } one pound.
 lows Root, }
 Olive Oil, three pounds.
 Semi-vitreous Oxide of Lead, two pounds.

Boil the oil and oxide over a slow fire occasionally adding one or two ounces of spring water, and constantly stirring; then add the mucilages, and continue the slow ebullition till the mass has acquired the

proper consistence. (BRUNS.; SARD.; WIRTEM.; HERBIP.; SPIELMANN)

PALAT. gives the same formula, but requires only one pound and a half of the oxide.

R. Root of Marshmal- } of each,
lows, } half a pound.
Linseed, }
Water, } eight pounds.

Infuse, boil, and express. Then

R. Of this mucilage, three pounds.
Semi-vitreous Oxide of Lead,

Olive Oil, two pounds.
four pounds.

Boil the oil and oxide together, adding the mucilage gradually till a plaster is formed. (HISP.)

R. White Plaster, } of each,
Emollient Plaster, } two pounds.

Melt together over a slow fire, and mix carefully. (LIPP.)

This variety of the simple plaster owes its Greek name of *diachylon* and its latin epithet of *mucilagineum*, to the mucilages which enter into its composition. It is fallen totally into disuse, but, by a very common abuse, the name of *diachylon* remains attached to the following variety, the only one now in use, although it does not contain a particle of mucilage.

2. Simple or common plaster; *Emplastrum simplex seu commune*. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; MIL. COPEN.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; GENEV.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; BRUGNATELLI; CADET DE GASSICOURT; COXE; SWEDIAUR; VAN MONS)

Emplastrum Plumbi. (LOND.)

R. Semi-vitreous Oxide of Lead,
in fine powder, five pounds.
Olive Oil, one gallon.
Water, two pints.

Boil together over a slow fire, constantly stirring, till the oil and oxide unite and become of the consistence of plaster. It will be necessary, however, to add a little boiling water, if that which was employed at the beginning is consumed before the end of the operation.

Emplastrum Lithargyri. (DUBL.)

R. Litharge, in powder, five pounds.
Olive Oil, nine pounds.
Boiling Water, two pints.

Mix at a high temperature, constantly stirring, till the oil and litharge unite and form a plaster, supplying from time to time any waste of water that may take place.

Emplastrum Oxydi Plumbi Semi-vitrei. (EDINB.)

R. Semi-vitreous Oxide of Lead, one part.
Olive Oil, two parts.
Water, a sufficient quantity.

Boil, continually stirring, till the oil and oxide unite and form a plaster.

AMER.; ANT.; AUSTR.; BELG.; DAN.; MIL. COPEN.; FERR.; ROSS.; BRUGNATELLI; COXE; and VAN MONS give the same formula as EDINB.

SWEDIAUR prescribes one part of the oxide and four parts of the oil.—HASS.—four of oxide and seven of oil.—GENEV.—four of oxide and two and a half of oil.—VAN MONS—two of oxide and three of oil.—LUSIT.—five of oxide and eight of oil.—FENN.—four of oxide and three of oil.—BAVAR.; HANN.; OLDEN.; BORUS.; SAX.; and SUEC.—five of oxide and nine of oil.

R. Semi-vitreous Oxide of Lead, six parts.
Lard, fifteen parts.

Proceed as above. (POL.; VAN MONS)

R. Semi-vitreous Oxide of Lead, } of each,
Olive Oil, } one
Lard, } part.

Heat slightly together, constantly stirring with a spatula; pour water upon the mass, and boil, constantly stirring, and replacing the water in proportion as it becomes evaporated, till no trace of the oxide is perceptible; then remove it from the fire, and knead it in fresh water. (GALL.; FORM. HÔP. FR.; CADET DE GASSICOURT)

AMST.; and BATAV. prescribe two parts of lard, and three each, of oil and of oxide.

This plaster is very much used in surgical practice, principally for protecting from the action of the air, wounds and ulcers whose surfaces are not very irritable. It also enters into the composition of a great many other plasters.

SATURNINE BOUGIES.

Cereoli exploratorii, Candelæ cereæ, Specilla cerea. (LIPP.)

R. Simple Plaster, four ounces.
Melt over a slow fire, and add to it
Wax, one ounce and a half.
Olive Oil, three ounces.

Dip strips of linen into the mixture, and form bougies. (LIPP.; VAN MONS)

- R. Turpentine, one part.
 White Wax, eight parts.
 Simple Plaster, four parts.
 Make bougies. (VAN MONS)

SATURNINE CERATE. (VAN MONS)

- R. Simple Plaster, six parts.
 White Wax, one half part.
 Melt separately, mix, and add
 Olive Oil, two parts and a half.
 Mix.

ANODYNE PLASTER.

Emplastrum Sebi anodynum. (HISP.)

- R. Simple Plaster, one pound.
 White Wax, six ounces.
 Lard, four ounces.
 Goat's Suet, two ounces.
 Melt together.

By adding four ounces of olive oil we obtain what HISP. calls *Ceratum anodynum*.

EPULOTIC PLASTER.

Emplastrum Oxydi Plumbi epuloticum. (HISP.)

- R. Semi-vitreous Oxide of Lead, } of each,
 Olive Oil, } three
 Lard, } pounds.

Boil with a sufficient quantity of water, constantly stirring, till the mass no longer adheres to the fingers; then add

- White Wax, nine ounces.
 Resin, six ounces.

Boil to the consistence of a plaster.

This scarcely differs from the simple plaster.

EPULOTIC SPARADRAP.

Tela emplastica epulotica, Sparadrapus palidus. (HISP.)

- R. Epulotic Plaster, two pounds.
 Yellow Wax, half a pound.
 Melt, and spread the mass upon strips of linen.

ANTI-HEMORRHOIDAL OINTMENT.

Unguentum ad Hemorrhoides seu ad Varices Ani. (SARD.)

- R. Fresh Root of Scrofularia, three ounces.
 Leaves of White Henbane, three handfuls.
 Fresh Butter, one pound and a half.

Semi-vitreous Oxide of Lead, two ounces.

Yolks of Eggs, two.

Boil the root and leaves slowly in the butter; after the consumption of the humidity, express, add the oxide, stir till perfectly cold, and finally incorporate the yolks of eggs.

ASTRINGENT PLASTER.

Emplastrum roborans seu adstringens. (COXE; SWEDIAUR)

- R. Simple Plaster, two pounds.
 Powdered Dragon's Blood, three ounces.
 ———— Olibanum, half a pound.
 Mix.

BROWN SATURNINE CERATE.

Ceratum fuscum, Unguentum fuscum seu Oxydi Plumbi fuscum seu Oxyduli Plumbi semi-vitrei adustum seu Acetatis Plumbi carbonatum seu matris seu dippurans. (ANT.; AUSTR.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; LIPP.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; NIEMANN; SPIELMANN; SWEDIAUR; VAN MONS)

- R. Olive Oil, five hundred parts.
 Lard, } of each, two
 Fresh Butter, } hundred and fifty parts.
 Mutton Suet, }

Melt together, and add gradually

Semi-vitreous Oxide of Lead, two hundred and fifty parts.

Boil till the mass becomes of a blackish-brown colour; then add a melted mixture of

- Yellow Wax, one hundred and eighty parts.
 Black Resin, eighty parts.

Pour the mass into paper moulds. (GALL.; NIEMANN)

FORM. HÔP. FR. suppresses the butter, and doubles the quantity of the lard.

- R. Lard, } of each, one part.
 Fresh Butter, }
 Mutton Suet, }

Boil together, and add

Semi-vitreous Oxide of Lead, one part.

Boil, constantly stirring, till the mass becomes of a brown colour; then add

- Yellow Wax, one part.

Preserve when cold. (BRUNS.; SAX.; NIEMANN; SPIELMANN)

- R. Lard, } of each, two pounds.
 Olive Oil, }

Fresh Butter, } of each, one pound.
Yellow Wax, }

Apply heat till the mixture begins to be carbonized, and add to it

Semi-vitreous Oxide of Lead, one pound,
constantly stirring with a wooden spatula;
continue the application of heat till the
mass becomes of a blackish-brown colour.
(ANT.; BELG.)

R. Lard, } of each,
Yellow Wax, } four ounces.
Fresh Butter, }
Mutton Suet, }
Olive Oil, one pound.
Semi-vitreous Oxide of Lead, four ounces.

Melt the fat substances over a slow fire;
then add the oxide, and boil, constantly
stirring, till the plaster becomes of a black
colour. (WIRTEM.)

HISP.; and GENEV. prescribe one pound
of oil, and half a pound each, of lard, of
butter, of wax, of goat's suet, and of the
oxide. BRUGNATELLI—fifteen ounces of
oil, and half a pound each, of lard, of
butter, of wax, of mutton suet, and of the
oxide.—LIPP.—six ounces of oil, and four
each, of lard, of butter, of wax, of stag's
suet, and of the oxide.

R. Lard, fourteen ounces.
Mutton Suet, }
Yellow Wax, } of each,
Semi-vitreous Oxide of } four ounces.
Lead, }

Boil till the mixture acquires a brown
colour. (AUSTR.)

R. Lard, } of each, six ounces.
Olive Oil, }
Yellow Wax, two ounces.

Add gradually to the melted and boiling
mass

Powdered Semi-vitreous Oxide of
Lead, two ounces.

Stop the boiling when the mass has be-
come black. (FULD.)

R. Simple Plaster, } of each, three parts.
Lard, }
Yellow Wax, one part.

Melt together, and encrease the heat till
the mass blackens. (VAN MONS)

R. Simple Plaster, fourteen parts.
Boil it, constantly stirring, till it has ac-
quired a brown tint, then add

Black Pitch, eight parts.
Mutton Suet, five parts.

Preserve. (BAVAR.)

R. Nuremburgh Plaster, eight parts.
Olive Oil, five parts.

Heat till the mixture becomes brown.
(HANN.)

This ointment is considered to be com-
posed of oleate, margarate, stearate, ace-
tate, and sometimes of caproate of pro-
tioxide of lead, incorporated with a very
carbonated fat.

It is reputed emollient and maturative.

HJERNE'S PLASTER.

*Emplastrum Hjernerii seu saponaceo-plum-
bicum.* (SUEC.)

R. Hard Potass-soap, scraped, ten parts.
Olive Oil, forty-five parts.

Melt the soap in the oil, and add

Powdered Semi-vitreous Oxide of
Lead, eighteen parts.
Subcarbonate of Lead, nine parts.

Boil till the mass has acquired a brown
colour; then add to it

Turpentine, one part and a half.
Mix.

DIAPALM PLASTER.

Emplastrum diapalmæ seu diaphænicon.
(BRUNS.; PALAT.; SARD.; WIRTEM.;
PIDERIT; SPIELMANN; VAN MONS)

R. Palm Oil, six parts.
Olive Oil, sixty-six parts.
Young Leaves of Oak, eight parts.

Boil to the consumption of the humidity.
Add to the expressed liquid

Semi-vitreous Oxide of Lead,
forty-eight parts.

Boil to the consistence of plaster, gra-
dually adding water, holding in solution

Sulphate of Zinc, four parts.

Preserve. (SPIELMANN)

BRUNS.; SARD.; and WIRTEM. prescribe
eight parts each, of lard, of olive oil, of oak
leaves, and of the oxide, and one part of
the sulphate dissolved in twelve of water.
—PALAT.—six of old lard, nine of palm or
olive oil, nine of the oxide, and one of the
sulphate dissolved in water.

R. Semi-vitreous Oxide of Lead,
two pounds.
Lard, three pounds.

Boil, adding gradually,

Water, a sufficient quantity,
of which sixteen ounces contain one dram
of alum in solution. (PIDERIT)

R. Olive Oil, nine parts.
Lard, six parts.
Semi-vitreous Oxide of Lead, nine parts.

Sulphate of Zinc, one part.
Make a plaster. (VAN MONS)

The following, by SOUBEIRAN, is a more simple formula :

- R. Simple Plaster, three thousand parts.
Sulphate of Zinc,
one hundred and twenty-eight parts.

Liquify the plaster, add the sulphate dissolved in water, and expose to a moderate heat, constantly stirring, till the water is totally evaporated.

It may be presumed that the object of the inventors of this plaster was to render it styptic: whence the precept originally given to stir it with a spatula of palm wood, esteemed astringent, then to put into it a decoction of palm fruit, and lastly a decoction of oak leaves: whence also the addition of sulphate of zinc, for which PIDERIT substitutes alum. However it may be, if the diapalm plaster really possesses any astringency, it must be in a very trifling degree. Its composition varies according to the formula which is followed in preparing it; but, with the exception of PIDERIT's, it contains sulphate of lead, with a variable quantity of oleate, margarate, and stearate of lead and of zinc. SARD.; and SPIELMANN erroneously give as an equivalent to this plaster, that which bore the name of *diachalciteos*, for in this latter the sulphate of zinc was replaced by an equal quantity of chalcitis, or of sulphate of iron calcined to redness.

GAULTHIER'S SPARADRAP.

Sparadrapum album, Tela Gualtheri.
(FORM. HÔP. FR.; WIRTEM.; SPIELMANN)

- R. Diapalm Plaster, } of each,
Simple Plaster, } one pound.
Ceruss Plaster, } half a pound.

Melt, and add

Powdered Florentine Iris,
one ounce and a half.

Spread the mass upon strips of linen.
(WIRTEM.; SPIELMANN)

- R. Simple Plaster, } of each,
Gummy Diachylon } fifteen parts.
Plaster, }
Yellow Wax, } five parts.
Turpentine, } three parts.

Melt in a water-bath, and spread upon linen. (FORM. HÔP. FR.)

RED STRENGTHENING PLASTER.

Emplastrum confortativum Vigonis seu defensionum rubrum seu Ferri seu Oxydi Plumbi rubrum seu Oxydi Plumbi semi-vitrei cum Sulphate Ferri calcinato seu Oxydi Ferri rubri seu sticticum seu roborans seu stypticum seu Siliquarum.
(EDINB.; DUBL.; BATAV.; BRUNS.; VOL. II.

HISF.; FERR.; PALAT.; SARD.; WIRTEM.; COXE; HENRI; SPIELMANN; VAN MONS)

Emplastrum Oxydi Ferri rubri. (EDINB.; COXE)

- R. Plaster of Semi-vitreous Oxide of Lead, twenty-four parts.
White Resin, six parts.
Yellow Wax, } of each,
Olive Oil, } three parts.
Red Oxide of Iron, eight parts.

Triturate the red oxide of iron with the oil; then add the other ingredients melted, and mix.

Emplastrum Thuris. (DUBL.)

- R. Litharge Plaster, two pounds.
Olibanum, half a pound.
Red Oxide of Iron, three ounces.

Melt the plaster and olibanum together, add the oxide, and stir to form a plaster.

- R. Semi-vitreous Oxide of }
Lead, } of each,
Deutoxide of Lead, } half a pound.
Calamine, }
Linseed Oil, } of each, one pound.
Olive Oil, }

Boil over a very slow fire, constantly stirring, and adding gradually one pound of water holding in solution

Sulphate of Zinc, one ounce.

When the plaster has arrived at the required consistence, add

- Resin, } of each, one pound.
Yellow Wax, }
Turpentine, } of each,
Oil of Bay, } half a pound.
Gum Ammoniac, }
Galbanum, }
Bdellium, }
Myrrh, } of each,
Hepatic Aloes, } one ounce.
Long Birthwort, }
Dragon's Blood, }
Hæmatitis, }

Mix. (BRUNS.; PALAT.; WIRTEM.)

- R. Semi-vitreous Oxide of Lead, } of
Deutoxide of Lead, } each,
Linseed Oil, } twelve
Olive Oil, } ounces.

Boil to the consistence of a plaster, adding

Sulphate of Zinc, four ounces,

dissolved in water. When the boiling is finished, add,

Resin, } of each,
Wax, } twenty-four ounces.
Turpentine, } twelve ounces.

Add to the mass when strained,

Powdered Gum Ammoniac, }
Galbanum, }
Bdellium, } of
Myrrh, } each,
Hepatic Aloes, } two
Mummy, } ounces.
Root of Long }
Birthwort, }
Dragon's Blood, }
Hæmatitis, }
Oil of Bay, } twelve ounces.

Mix. (SPIELMANN)

R. Mucilage of Comfrey Root, six pounds.
Walnut Oil, four pounds.
Deutoxide of Lead, } of each,
Semi-vitreous Oxide of } one pound.
Lead, }
Calamine, } four ounces.
Dragon's Blood, }
Colcothar, } of each, one ounce.
Alum, }
Root of Round Birthwort, } one ounce and a half.

Boil the oxides and the calamine with the oil, gradually adding the mucilage, and incorporate in powder, when the mass is nearly cold. (SARD.)

R. Mucilage of Marshmallows } of each,
Root, } two
Comfrey Root, } pounds.
Oil of Myrrh, }
Goat's Suet, } of each,
Semi-vitreous Oxide of } one pound.
Lead, }
Deutoxide of Lead, } six ounces.
Turpentine, }
Mastic, } of each, four ounces.
Armenian Bole, }

Boil the oxides with the mucilages, suet, and oil; then add the turpentine, the bole, and finally the mastic. (SARD.)

R. Root and Leaves of Great } of each,
Comfrey, } four
Myrtle Berries, } ounces.
Petals of Red Roses, }
Tops of St. John's-wort, }
Water, } nine pounds.

Boil to the reduction of one third. Then

R. Olive Oil, four ounces.
Goat's Suet, one pound.
Semi-vitreous Oxide of Lead, two pounds.

Boil to the consistence of plaster, gradually adding the preceding decoction. Then melt in the mass,

Yellow Wax, one pound.
Turpentine, three ounces.

Boil sufficiently; and when the mass is half cold, add a mixture, in powder, of

Olibanum, } of each, two ounces.
Myrrh, }
Mastic, }
Dragon's Blood, three ounces.
Armenian Bole, four ounces.
Deutoxide of Lead, six ounces.

Mix well. (HISP.)

R. Resin, eight ounces.
Yellow Wax, six ounces.

Add to the melted and strained mass,

Extract of Carob Bean, reduced to the consistence of honey, one pound.
Plaster of Deutoxide of Lead, four ounces.

Subcarbonate of Iron, six drams.

Mix. (HISP.)

R. Semi-vitreous Oxide of Lead, twelve ounces.
Olive Oil, twenty-four ounces.
Boil to the consistence of plaster, and melt in the mass,
Yellow Wax, } of each,
Venice Turpentine, } two ounces.
Olibanum, }

When liquified, add,

Colcothar, three ounces.

Mix well. (SPIELMANN)

R. Yellow Wax, } of each, one part.
Resin, }

Melt, and add,

Simple Plaster, sixteen parts.

At the moment of its cooling, add,

Turpentine, one part

trituated, hot, with

Washed Colcothar, one part and a half.

Mix. (VAN MONS)

R. Simple Plaster, two pounds.
Resin, half a pound.
Yellow Wax, } of each, three ounces.
Olive Oil, }
Colcothar, } eight ounces.

Add the oxide trituated with the oil to the two other substances melted together over a slow fire. (BATAV.)

R. Simple Plaster, } of each,
Gummy Diachylon Plaster, } an equal
Yellow Wax, } part.
Olive Oil, }
Colcothar, }

Mix the oxide with a little oil, and incorporate it with the other ingredients melted together. (HENRI)

R. Simple Plaster, two pounds.
Mutton Suet, sixteen ounces.

Melt and add,

Colcothar, two ounces.
Mix. (FERR.)

IMPERIAL PLASTER.

Emplastrum Siliquarum imperiale. (HISP.)

R. Plaster of Black Pitch, } of
 ——— Deutoxide of Lead, } each,
 ——— Opodeldoc (Form. } four
 of HISP.), } ounces.
 Sulphuretted Plaster of } of each,
 Resin, } two
 Red Strengthening Plaster, } ounces.
 Acetate of Lead, } of each, two drams.
 Camphor, }
 Red Oxide of Iron, half an ounce.

Melt the plasters together over a slow fire, and, before the mass becomes cold, add to it the three other ingredients in powder.

This farrago scarcely differs in effect, from the red strengthening plaster.

OPODELDOC PLASTER.

Emplastrum Collæ ad Herniam; Emplâtre fondant de Rustaing. (HISP.; WIRTEM.; BORIES; PIERQUIN)

R. Semi-vitreous Oxide of Lead, two pounds.
 Olive Oil, two pounds and a half.
 Yellow Wax, one pound.
 Chio Turpentine, } of each,
 Oil of Bay, } four ounces.
 Opoponax, two ounces and a half.
 Bdellium, }
 Gum Ammoniac, } of each,
 Sarcocolla, } two ounces.
 Olibanum, }
 Mastic, }
 Myrrh, }
 Socotrine Aloes, one ounce.
 Root of Birthwort, two ounces.
 Camphor, three ounces.

Make a plaster. (BORIES; PIERQUIN)

R. Semi-vitreous Oxide of Lead, one pound and a half.
 Olive Oil, two pounds.
 Boil over a slow fire, adding gradually,
 Vinegar, } of each, four ounces,
 Spring Water, }
 till the mass has acquired a light brown colour. Add,

Yellow Wax, half a pound.
 Turpentine, four ounces.

When cold, add a mixture, dissolved in vinegar and reduced to the consistence of honey, of

Gum Ammoniac, }
 Bdellium, } of each, one ounce.
 Galbanum, }
 Opoponax, }

Then incorporate,

Powdered Round Birthwort, }
 ——— Long Birthwort, } of
 ——— Calamine, } each,
 ——— Olibanum, } one
 ——— Myrrh, } ounce.
 ——— Mastic, }

Oil of Bay, one ounce and a half.
 Stir till perfectly cold. (WIRTEM.)

R. Myrtle Berries, bruised, } of each,
 Earth Worms, washed } six ounces.
 in wine,
 Water, eight pounds.

Boil down to one half. Then

R. Of this Decoction, three pounds.
 Olive Oil, two pounds.
 Semi-vitreous Oxide of Lead, one pound.

Boil to the consumption of the humidity and to the consistence of a plaster; then add a melted and strained mixture of

Yellow Wax, one pound.
 Black Pitch, nine ounces.
 Resin, eighteen ounces.

Add further, a mixture, dissolved in vinegar and evaporated to the consistence of honey, of

Galbanum, } of each,
 Gum Ammoniac, } four ounces.

When the mass is nearly cold, incorporate a mixture, in powder, of

Myrrh, }
 Olibanum, }
 Mastic, }
 Goat's Blood, }
 Glue, }
 Root of Round Birthwort, } of each,
 ——— Long Birthwort, } three
 ——— Great Comfrey, } ounces.
 ——— Lesser Comfrey, }

Nutgalls,
 Armenian Bole,
 Gypsum,
 Mummy,

Mix. (HISP.)

This incongruous mixture, which resembles some of the formulæ for the red strengthening plaster, was considered discutient, resolvent, cicatrizing, and astringent.

OINTMENT OF THE TWELVE APOSTLES.

Unguentum Apostolorum seu dodecapharmacum. (BRUNS.; PALAT.; WIRTEM.; PIERQUIN; SPIELMANN)

R. Turpentine, } of each,
 Resin, } two ounces.
 Yellow Wax, }
 Long Birthwort, }
 Olibanum, }
 Bdellium, } of each,
 Myrrh, } half an ounce.
 Galbanum, }

Oil of Lavender, two ounces.
 ——— Linseed, one ounce and a half.

Mix. (PALAT.)

R. Live Frogs, ten.
 ——— Earth Worms, four ounces.
 Herb of Lemon-grass, } of each,
 French Lavender, } half an ounce.
 Mother-wort, }
 Root of Elicampne, } of each,
 ——— Dwarf Elder, } three ounces.
 White Wine, two pounds.
 Vinegar, one pound and a half.

Boil slowly to the reduction of one half, and strain; add gradually to the liquor, a mass prepared with

Olive Oil, } of each,
 Semi-vitreous Oxide } one pound.
 of Lead, }

Having removed the plaster from the fire, add,

Oil of Bay, } of each, two ounces.
 Turpentine, }
 Yellow Wax, }
 Powdered Olibanum, six drams.
 ——— Euphorbium, five drams.
 Storax, } of each, half an ounce.
 Saffron, }

Mix. (WIRTEM.; SPIELMANN)

R. Live Frogs, eighty.
 Fresh Root of Dwarf } of each,
 Elder, } one pound
 ——— Burdock, } and a half.
 Vinegar, three pounds.
 Water, nine pounds.

Macerate for two days; then boil to the reduction of one half. Then

R. Of the strained Decoction, five pounds.
 Olive Oil, nine pounds.

Boil to the consumption of the humidity; remove from the fire, and add,

Lard, three pounds.
 Semi-vitreous Oxide of Lead, six pounds.
 Of the preceding Decoction,
 a sufficient quantity.

Boil till the mass no longer adheres to the finger. Then add,

Yellow Wax, three pounds.
 Turpentine, two ounces.

Finally add to the mass when nearly cold,

Powdered Euphorbium, one ounce.

Mix. (HISP.)

This plaster, once so celebrated, is now entirely disused. More complicated than the ancient simple diachylon, it does not differ from it in action, except by being more excitant.

DIABOTANUM PLASTER.

Emplastrum Oxydi Plumbi diabatanum seu

resolutivum. (HISP.; SARD.; WIRTEM.; SPIELMANN)

R. The Roots and Fresh Herbs of
 Burdock, }
 Ground Pine, }
 Valerian, } of each,
 Angelica, } three
 Elicampne, } ounces.
 Round Birthwort, }
 Horse-radish, }
 Wild Cucumber, }
 Scrofularia, }
 Sowbread, }
 Lesser Celandine, }
 Juice of Hemlock, } of each,
 ——— Ormin, } four
 ——— Great Celandine, } pounds.
 Semi-vitreous Oxide of Lead, two pounds and a half.

Olive Oil, four pounds.
 Yellow Wax, }
 Turpentine, } of each, one pound.
 Resin, }
 Tacamahac, }
 Sulphur, }
 Asphaltum, }
 Olibanum, } of each, six ounces.
 Mastic, }
 Myrrh, }
 Euphorbium, } of each,
 Camphor, } one ounce and a half.
 Oil of Cloves, half an ounce.

Bruise the herbs and roots, pour the juices over them, and leave to macerate for two days; then, having boiled to the consumption of one half, express: boil the strained liquor to the proper consistence with the oil and oxide, constantly stirring; add the wax, resin, and turpentine, then the tacamahac, mastic, and euphorbium, and, finally, when cold, the other substances reduced to powder, and triturated with the oil of cloves. (SARD.)

R. Semi-vitreous Oxide of Lead, two pounds.
 Olive Oil, four pounds.
 Vinegar, a sufficient quantity.

Boil to the consistence of a plaster, adding towards the end, a liquified mixture of

Inspissated Juice of Hemlock, } of each,
 ——— Great Ce- }
 landine, } two
 ——— Wild Cu- } ounces.
 ——— cumber, }
 Extract of Root of Eli- } of each,
 ——— campne, } half
 ——— Valerian, } an ounce.

Boil for a few moments; then add a melted and strained mixture of

Yellow Wax, }
 Resin, } of each, half a pound.
 Turpentine, }
 Storax, three ounces.

Incorporate in mass when nearly cold,

Powdered Gum Ammoniac,	} of each, three ounces.
Galbanum,	
Olibanum,	} two ounces.
Mastic,	
Tacamahac,	} of each, one ounce.
Bay Berries,	
Root of Iris,	} of each, six drams.
Black Hellebore,	
Seeds of Cumin,	} three ounces. half an ounce.
Olive Oil,	
Camphor,	

Mix. (Hisp.)

R. Fresh Herbs and Roots of	} of each, three ounces.
Burdock,	
Butterbur,	
Marigold,	
Ground Pine,	
Lovage,	
Valerian,	
Angelica,	
Hemlock,	
Elicampne,	
Horse-radish,	
Wild Cucumber,	
Scrofularia,	
Lesser Houseleek,	
Great Celandine,	
Lesser Celandine,	
Gratiola,	
Water,	twenty pounds.

Boil to the consumption of one half; express, and add to the liquor,

Juice of Hemlock,	} of each, two pounds.
Great Celandine,	
Ormin,	
Lesser Houseleek,	half a pound.

Evaporate to the consistence of extract. Then

R. Of this Extract,	half a pound.
Galbanum,	} of each, four ounces,
Gum Ammoniac,	
Oponox,	
Sagapenum,	

the whole dissolved in vinegar of squill, and evaporated to the consistence of extract. Then

R. Semi-vitreous Oxide of Lead,	one pound.
Oil of Earth worms,	} of each, four ounces.
Puppies [<i>Petits chiens</i>],	
Mellilot,	
Mucilage,	
Spring Water,	ten ounces.

Boil to the consistence of plaster; then gradually add to it, the extract mixed with the gums, and

Sublimed Sulphur,	seven ounces and a half.
-------------------	--------------------------

Powdered Roots of	} of each, three drams.
Florentine Iris,	
Sowbread,	
Ranunculus,	
Imperial Crown,	
Serpentaria,	
White Hellebore,	} of each, half an ounce.
Solomon's Seal,	
Asarum,	
Long Birthwort,	} of each, one dram.
Round Birthwort,	
Clematis,	
Bay Berries,	two drams.
Seeds of Angelica,	} of each, three drams.
Cress,	
Cumin,	one ounce and a half.
Pigeon's Dung,	half an ounce.
Powdered Asphaltum,	} of each, four ounces.
Olibanum,	
Mastic,	} six ounces.
Tacamahac,	
Bdellium,	} of each, one ounce and a half.
Myrrh,	

Then add gradually a melted mixture of
Yellow Wax, } of each,
Resin, } half a pound.

Finally incorporate,

Storax,	six ounces.
Camphor, dissolved in half an ounce of Oil of Cloves,	six drams.
Oil of Bricks,	ten drams.

Preserve. (WIRTEM.; SPIELMANN)

This monstrous farrago, which owes its name to the great number of plants which enter into it, and which formerly equalled VIGO's plaster in reputation, was preferred by some to that as resolvent and discutient. It is now forgotten every where except in Spain.

II. Prepared with the Deutoxide of Lead.

ONIMENT OF DEUTOXIDE OF LEAD.

Unguentum magistrale seu de Minio seu Oxydi Plumbi rubrum. (HISP.)

R. Deutoxide of Lead,	five ounces.
Subcarbonate of Lead,	} of each, one ounce
Semi-vitreous Oxide of Lead,	
Rose Oil,	one pound and a half.

Boil a little, constantly stirring. Add
White Wax, four ounces.

Stir till perfectly cold.

OPIATE RESOLVENT PLASTER. (BORIES)

R. Linseed Oil,	two pounds.
Deutoxide of Lead,	} of each, eight ounces.
Subcarbonate of Lead,	
Yellow Wax,	} three ounces.
Turpentine,	
Opium,	one ounce.

Make a plaster.—It is recommended for incipient scirrhus tumours.—It is to be renewed every eight days.

NUREMBURG PLASTER.

Emplastrum seu Ceratum Noribergense seu Noricum seu tripharmicum seu Minii adustum seu Minii camphoratum. (BRUNS.; DAN.; GALL.; HANN.; SARD.; SAX.; WIRTEM.; CADET DE GASSICOURT; NIEMANN; SPIELMANN; VAN MONS)

R. Deutoxide of Lead, eight ounces.
Olive Oil, one pound.

Boil, constantly stirring, till the mass assumes a brown colour; add to this when half cold,

Yellow Wax, one ounce and a half,
and after the liquefaction of this

Camphor, triturated with a little oil,
two drams.

Mix. (HANN.)

SPIELMANN prescribes twelve ounces of deutoxide, twenty-four ounces of rose oil, three ounces of wax, and two drams of camphor.—VAN MONS—sixteen ounces of deutoxide, twenty-four of olive oil, three of wax, and three drams of camphor.—CADET DE GASSICOURT—one pound three ounces and one dram of olive oil, four ounces three drams of deutoxide, one pound of wax, and six drams of camphor.—GALL.—six hundred parts of oil, three hundred of deutoxide, five hundred of wax, and twenty-four of camphor.—The two latter require a certain quantity of water to be added to the oil and oxide before boiling.

R. Deutoxide of Lead, one pound.
Oil of Rape, two pounds.

Boil constantly stirring, till the plaster becomes of a brown colour, and add

Camphor, triturated with alcohol,
two drams.

Mix. (SAX.; NIEMANN)

This formula is by TROMMSDORF.

R. Deutoxide of Lead, eight ounces.
Vinegar, four ounces.
Rose Oil, one pound.

Boil to the consistence of a soft plaster, and add

Yellow Wax, one ounce and a half.
Camphor, two drams.

Mix. (SARD.; WIRTEM.)

DAN. prescribes one pound of deutoxide, half a pound of vinegar, two pounds of oil, one ounce and a half of wax, and two drams of camphor.—BRUNS.; and NIEMANN—one pound each of deutoxide and of vinegar, two pounds of oil, and one ounce and a half of camphor, with no wax.

This plaster is considered soothing and defensive. It is employed principally to

accelerate the cicatrization of ulcers, and to protect parts irritated by long recumbence.

STAHL'S CAMPHORATED PLASTER. (SPIELMANN)

R. Olive Oil, twelve ounces.
Deutoxide of Lead, eight ounces.

Boil to the required consistence, and add to the mass when nearly cold,

Camphor, dissolved in a little oil,
two ounces.

This plaster should be preferred to the preceding, as being more simple.

LEEK PLASTER. (BORIES)

R. Deutoxide of Lead, two pounds.
Olive Oil, four pounds.
Juice of Leeks, six pounds.

Boil to the required consistence, and add

Yellow Wax, one pound.
White Soap, two pounds.

Stir till perfectly cold.

ANGLO-SAXON PLASTER. (NIEMANN)

R. White Wax, } of each,
Deutoxide of Lead, } one pound.
Powdered Amber, } of each,
———— Camphor, } two drams.
———— Calcined Alum, }
Olive Oil, one pound.

Boil the oil till it acquires a reddish-brown colour; then add the deutoxide, continue the boiling, and when the mass has acquired the consistence of plaster, add the amber, and when cold, the camphor and alum.

It is extolled as an application to foul ulcers.

MAMMILLARY PLASTER. (PALAT.)

R. Plaster of Deutoxide of Lead, } of each,
———— Mucilage, } half
Powdered Deutoxide of Lead, } an ounce.
———— Semi-vitreous Oxide of Lead, }
Oil of Lily, } of each,
———— Eggs, } one ounce.
———— Wild Cucumber, }
———— Sesame, }
Fresh Butter, }
Spermaceti, }
Camphor, half an ounce.

Make a plaster.—It is esteemed resolvent in engorgements of the breasts of nurses, and prevents fissures of the nipples.

DOYEN'S PLASTER. (CADET DE GASSI-COURT)

- R. Deutoxide of Lead, } of each
 Oil of St. John's-wort, } one pound.
 Resin, } four ounces.
 Olibanum, } two ounces.
 White Soap, } half an ounce.
 Make a plaster.

ANTI-ARTHRITIC PLASTER. (WIRTEM.)

- R. Henbane Oil, } of each, four ounces.
 Poppy Oil, }
 Linseed Oil, }
 Deutoxide of Lead, } one pound.
 Elder Water, } half a pound.
 Boil slowly to the consistence of plaster,
 and add
 Venice Soap, three ounces.
 Yellow Wax, two ounces.

It used formerly to be recommended to be applied to parts affected with pains and spasms.

MIRACULOUS PLASTER. (WIRTEM.)

- R. Deutoxide of Lead, } of each,
 Semi-vitreous Oxide } half a pound.
 of Lead, }
 Subcarbonate of Lead, } two pounds.
 Olive Oil, } three pounds.
 Boil slowly and add
 Venice Soap, scraped, four ounces.
 When the mass has become of the con-
 sistence of a plaster, add further.
 Powdered Bay Berries, half an ounce.
 ——— Camphor, one ounce.

It was formerly considered,—and hence its name,—a sovereign remedy against old ulcers, and even against recent wounds. It does not sensibly differ from the following.

BARBETTE'S SAPONACEOUS PLASTER. (BRUNS.; HISP.; PALAT.; WIRTEM.)

- R. Deutoxide of Lead } of each,
 Subcarbonate of Lead, } four ounces.
 Rose Oil, } one pound.
 Boil, constantly stirring, and adding a few ounces of spring water, to the consistence of a plaster; then throw into the mass,
 Venice Soap, scraped, four ounces.
 and when melted,
 Camphor, triturated with alcohol, half an ounce.

Mix. (BRUNS.; PALAT.; WIRTEM.)

HISP. directs twelve ounces of deutoxide and six of subcarbonate to be mixed with thirty of oil, three of soap, dissolved in a sufficient quantity of water, to be added, and the mixture shaken so as to give it the appearance of a liniment; it is then to be

boiled slowly, continually stirring and adding a little water from time to time; the humidity to be then entirely evaporated, one ounce and a half of wax to be added, and finally, when nearly cold, one ounce of camphor.

This plaster was considered discutient, resolvent, and anodyne; it used also to be applied to the abdomen, as a relaxant to children.

BALSAMIC PLASTER. (NIEMANN)

- R. Olive Oil, three pounds.
 Venice Soap, } of each,
 Deutoxide of Lead, } one pound.
 Subcarbonate of Lead, }
 Castor, one ounce and a half.
 Camphor, three ounces.

Mix.—For the secret of this plaster Catharine II. gave thirty thousand rubles. It was extolled in head-ache, blindness, deafness, rheumatism, tooth-ach, goitre, palsy, swelled breasts, cancer of the face, and a host of other affections. The advisers of the empress might have given a better direction to her munificence.

III. Prepared with the Subcarbonate of Lead.**PLASTER OF SUBCARBONATE OF LEAD.**

Emplastrum Cerussæ seu album coctum, seu *e Plumbo compositum seu Subcarbonatis Plumbi*. (BAVAR.; BELG.; BRUNS.; DAN.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SARD.; SAX.; WIRTEM.; HERBIP.; NIEMANN; SPIELMANN; SWEDIAUR; VAN MONS)

- R. Subcarbonate of Lead, four parts.
 Olive Oil, three parts.

Boil, constantly stirring and adding hot water from time to time, till the subcarbonate is dissolved. (SPIELMANN)

BAVAR.; LIPP.; and SAX. prescribe two parts of subcarbonate and one part of oil.—VAN MONS—one of subcarbonate and one of oil.—POL.—nine of subcarbonate and six of lard.

- R. Subcarbonate of Lead, one part.
 Olive Oil, twelve parts.

Boil, constantly stirring, and adding water gradually, till the mass assumes the consistence of plaster; then add

- White Wax, three parts.
 Preserve when cold. (SWEDIAUR)

HANN.; OLDEN.; BORUS.; and NIEMANN prescribe eight parts of subcarbonate, one part of wax, and eight parts and a half of oil.—HERBIP.—eight of subcarbonate four of wax, and one of oil.—BRUNS.; PALAT.; and WIRTEM.—eight of subcarbonate, one of wax, and six of oil.—DAN.—eight of subcarbonate, one of wax, and four of oil.—SARD.—four of subcarbonate, one of wax,

and four of oil.—FULD. — twelve of subcarbonate, one of wax, and ten of oil.—
VAN MONS—four of subcarbonate, one of wax, and six of oil.

- R. Subcarbonate of Lead, one part.
Semi-vitreous Oxide of Lead,

two parts.
Olive Oil, four parts.

Boil with a sufficient quantity of water, and add

White Wax, melted, one part.

Mix. (VAN MONS)

- R. Simple Ointment, five parts.
Olive Oil, one half part.

Soften over a slow fire, and add

Subcarbonate of Lead, one part.

Stir till cold. (BELG.)

This plaster is considered desiccative and sedative.

PLASTER OF FROG'S SPAWN.

Emplastrum de Spermate Ranarum. (PALAT.; WIRTEM.; SPIELMANN)

- R. Frog's Spawn, } of each, three pounds.
Olive Oil, }
Subcarbonate of Lead, }
four pounds and a half.

White Wax, one pound.
Alum, four ounces.
Turpentine, two drams.
Camphor, one ounce.

Make a plaster. (PALAT.)

- R. Oil of Frog's Spawn, } of each,
Subcarbonate of Lead, } five pounds.

Boil till the subcarbonate is dissolved, adding a little water and vinegar to prevent the mass from burning. Withdraw it from the fire, and add

Alum, five ounces.
Camphor, triturated with Alcohol, one ounce.

Mix well. (WIRTEM.)

SPIELMANN prescribes sixty parts of the oil of frog's spawn, eighteen of subcarbonate, six of alum, and two of camphor.

This plaster differs but little from the preceding in its action: if that is nearly disused now, this is still more completely forgotten.

COMPOUND PLASTER OF SUBCARBONATE OF LEAD.

Emplastrum Plumbi Subcarbonatis compositum. (AMER.; COXE)

- R. Subcarbonate of Lead, one pound.
Olive Oil, two pounds.
Yellow Wax, four ounces.

Vol. II.

Simple Diachylon Plaster,

one pound and a half.

Powdered Iris, nine ounces.

Boil the oil and the subcarbonate in a little water, constantly stirring, till perfectly incorporated; then add the wax and plaster, and, when melted, stir in the powder.

COMPOUND CERATE OF SUBCARBONATE OF LEAD. (AMER.; COXE)

- R. Compound Plaster of Subcarbonate of Lead, five parts.
Olive Oil, one part.

Melt the plaster, and add the oil, constantly stirring till the mass is cold.

OINTMENT OF SUBCARBONATE OF LEAD.

Unguentum Carbonatis Plumbi.

- R. Simple Ointment, five parts.
Carbonate of Lead, one part.

Mix. (EDINB.)

- R. Ointment of White Wax, one pound.
Carbonate of Lead, in fine powder, two ounces.
Make an ointment. (DUBL.)

RICOUR'S OINTMENT. (CADET DE GASSI-COURT)

- R. Rose Oil, twelve ounces.
White Wax, eight ounces.
Subcarbonate of Lead, four ounces.
Semi-vitreous Oxide of Lead, two ounces.

Boil, and add

Balsam of Peru, half an ounce.

Mix.—It is recommended to indolent ulcers.

EMPLASTRUM BENEDICTUM.

Emplastrum Carbonatis Plumbi ustum.
(HISP.; PALAT.)

- R. Olive Oil, one pound and a half.
Root of Sweet Flag, one ounce.
Iris, } of each,
Marshmallows, } six drams.

When the oil begins to boil, throw into it the roots cleaned and sliced, and boil till the liquor becomes black; then remove it from the fire, and add

Subcarbonate of Lead, half an ounce.
Prepared Tutty, six drams.

Boil again to the consistence of plaster, and add

Yellow Wax, half an ounce.
Balsam of Peru, one dram and a half

Mix. (PALAT.)

3 F

HISP. prescribes three pounds of oil, two ounces each, of marshmallows root and of root of sweet flag, three ounces of root of iris, six ounces of subcarbonate, three

ounces of semi-vitreous oxide of lead, no tutty, four ounces of yellow wax, and half an ounce of balsam.

POCGEREBÆ CORTEX.

BRUNS.

An American bark, in quilled pieces the size of which varies from that of a goose-quill to that of the finger, hard, compact,

solid, straight or curved, rugose, and of a deep brown externally, brown internally, inodorous, and of a weak and astringent taste.

PODOPHYLLUM PELTATUM, WILLD.—MAY APPLE.

AMER.; COXE.

A plant of North America. (*Polyandria Polygynia*, LINN.; *Podophyllaceæ*, CAND.)

The root is employed. It is of the diameter of the finger.—It is an excellent purgative in the dose of twenty grains.

EXTRACT OF MAY APPLE.

R. Root of May Apple, one pound.

Alcohol,
Water,

four pounds.
ten pints.

Macerate the root for four days in the alcohol; then boil it in the water till this is reduced to two pints: evaporate separately the two liquors, then mix them, and evaporate slowly to the proper consistence.

POLANISIA DODECANDRA, CAND.—BASTARD MUSTARD.

SYNONIME :—Mosambe dodécandre, FR.

COXE.

A plant of the East Indies. (*Tetrad-*

namia Siligiosa, LINN.; *Capparideæ*, JUSS.; *Fig. Bart. Flor. N. Amer. t. l. XXII*)

The root is reputed anthelmintic.

POLYGALA.

Four species of this genus of plants are indicated in pharmacopœias.

1. *Polygala amara*, LINN.—Milkwort.

SYNONIMES :—*Polygala amer*, FR.; *Bittere Kreuzwurz*, *Kreutzbleumenkraut*, *Milchblume*, GERM.; *Vstawaæ*, *Zezhulka*, *Mudatka*, *BOHEM.*; *Korsblomster*, DAN.; *Poligala lechera*, SPAN.; *Kruisbloem*, DUT.; *Poligala*, ITAL.; *Wyczka*, POL.; *Poligala leitera*, PORT.; *Jungfrumariælin*, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.;

BELG.; BRUNS.; DAN.; HISP.; GALL.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of Europe. (*Diadelphina Octandria*, LINN.; *Polygaleæ*, JUSS.; *Fig. Jacq. Fl. Austr. V. t. 412*)

The root (*Radix Polygalæ amaræ*) is employed. It is hard, almost ligneous, and filamentous. It is yellowish without, whitish within, inodorous, and of a taste at once bitter and sweetish.

The entire plant is usually gathered on account of the smallness of the roots, but it is seldom met with in commerce, the *P. vulgaris* being almost always substituted for it.

It is tonic, and is reputed pectoral. It has been much extolled in pulmonary phthisis.—Dose of the powder, half a dram.

2. *Polygala vulgaris*, LINN.

SYNONIMES : — *Polygala commun*, FR. ; *Kreutzb Blumen*, GERM.

AMST. ; GALL. ; WIRTEM. ; HERBIP. ; GUIBOURT ; MURRAY and GMELIN.

A plant of Europe. (*Diadelphia Octandria*, LINN. ; *Polygaleæ*, JUSS. ; *Fig. Jacq. Fl. Austr. V. 413*)

The root and herb (*Radix et Herba Polygalæ vulgaris* seu *Ambarvalis*) are employed.

They are generally gathered together. They are said to be less active than those of the *P. amara*, though that appears to be merely a variety of it, and is smaller than it by one half in all its parts.

3. *Polygala Seneka*, LINN.—Seneka, Rattle Snake Root.

SYNONIMES : — *Polygala de Virginie*, Seneka, Seneka, FR. ; *Klapperschlangenzurzel*, Senegawurz, GERM. ; *Indyczk Wirginiansky*, POL.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; MIL. COPEN. ; GALL. ; FERR. ; FENN. ; GENEV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY and GMELIN ; PARIS ; SPIELMANN.

A plant of North America. (*Diadelphia Octandria*, LINN. ; *Polygaleæ*, JUSS. ; *Fig. LINN. Amen. acad. 2. p. 139. t. 2*)

The root (*Radix Senegæ* seu *Senekæ* seu *Polygalæ Virginianæ*) is employed. It is contorted, nodose, branching, furnished with a greyish and resinous bark covering a whitish parenchyma. It has a feeble and nauseous odour, and a taste at first sweet, then bitter and acrid, exciting a flow of saliva.

It contains, according to PESCHIER, three peculiar principles, *Polygaline* or *Senegine* (GEHLEN), *Isolusine*, and *Polygalic Acid*.

It is excitant, diuretic, deobstruent, and emetic and purgative in a large dose.—Dose of the powder, from ten grains to half a dram.

4. The last species is the *Polygala rubelia*, WILLD.

AMER. ; COXE.

PECTORAL SPECIES. (HANN.)

R. Root of Polygala, half an ounce.
Flowers of Leopard's Bane, two drams.
Cut, and mix.

DEOBRUENT POWDER.

Pulvis resolvens anti-phlogisticus. (FULD.)

R. Root of Seneka, } of each,
Nitrate of Potass, } one dram.

Mix.—Dose, from a scruple to half a dram.

RESOLVENT PILLS. (NIEMANN)

R. Root of Seneka, any quantity.
Mucilage of Tragacanth, a sufficient quantity.

Make pills of two grains each.—Dose, one pill every two hours.—WENDT extols them in procuring the resolution of hydropyon.

BITTER PECTORAL POTION. (RATIER)

R. Root of Polygala, two drams.
Simple Syrup, two ounces.
Water, four ounces.

Infuse, and mix.

PECTORAL ELECTUARY.

Electuarium Polygalæ amaræ. (AUGUSTIN)

R. Powdered Root of Polygala amara, half an ounce.
Honey, a sufficient quantity.

Mix.—Dose, a tea spoonful daily, in chronic affections of the bronchiæ.

EXTRACT OF POLYGALA AMARA. (HERBIP.)

R. Polygala, any quantity.
Boil in a sufficient quantity of water for an hour, and express ; having set aside and decanted the liquor, evaporate it to the consistence of thick honey.

EXTRACT OF SENEKA.

Extractum Senegæ Radicis. (FULD. ; BORUS. ; SAX.)

R. Root of Seneka, one pound.
Proof Spirit, six pounds.

Digest for a few days in a moderate heat ; then filter the tincture, and evaporate it to the consistence of honey : boil the residue with

Water, three pounds.

Having expressed and evaporated the decoction, mix the two extracts, and continue to evaporate them to the proper consistence. (FULD.)

- R. Root of Seneka, two parts.
 Alcohol, three parts.
 Water, nine parts.

Digest for twenty-four hours; then distil off the alcohol, and evaporate the remainder to the consistence of extract. (BORUS.; SAX.)

Dose, from six to fifteen grains.

TINCTURE OF SENEKA.

Tinctura seu Essentia Senegæ. (HANN.; VAN MONS)

- R. Root of Seneka, one part.
 Proof Spirit, six parts.

Infuse for several days without heat. (VAN MONS)

HANN. prescribes five ounces of root and two pounds of alcohol.—Dose, thirty drops, several times a day.

INFUSION OF POLYGALA AMARA. (RATIER)

- R. Root of Polygala amara, half an ounce.
 Boiling Water, two pounds.

Infuse, and strain.

It is tonic, and is recommended as drink, in low fevers.

DECOCTION OF POLYGALA AMARA. (HISP.; FULD.; NIEMANN)

- R. Root of Polygala amara, one ounce.
 Water, a sufficient quantity
 to obtain two pounds of strained decoction. (HISP.)

NIEMANN prescribes two ounces of polygala and three pounds of water, reduced to one half.

- R. Root of Polygala amara, one ounce and a half.
 Water, eighteen ounces.

Boil down to two thirds, and add to the strained decoction

Oxymel, two ounces.

Mix. (FULD.)

It is recommended in pulmonary phthisis.—Dose, from four to six ounces, several times a day.

DECOCTION OF SENEKA.

Decoctum Senegæ seu Polygalæ Senegæ. (LOND.; EDINB.; AMER.; FULD.; GENEV.; HERBIP.; WURCEB.; BRUGNATELLI.; COXE.; ELLIS.; NIEMANN.; SAUNDERS.; SWEDIAUR)

- R. Root of Seneka, one ounce.

Water, two pints.

Boil down to one pint. (LOND.; EDINB.)

AMER.; HERBIP.; WURCEB.; BRUGNATELLI.; and COXE give the same formula.

NIEMANN prescribes three ounces of the root and two pounds of water, reduced to one-third.—FULD.; and GENEV.—one ounce of root, and one pound and a half of water, reduced to one pound.

- R. Root of Seneka, one ounce.
 Water, two pounds.

Boil down to one pound; then strain, and add

Simple Syrup, one ounce.

Mix. (SAUNDERS.; SWEDIAUR)

- R. Root of Seneka, one ounce.
 Liquorice, half an ounce.
 Water, one pound and a half.

Boil down to one pound. (ELLIS)

It is excitant, and is recommended in catarrhal affections of the lungs, arthrodynia, and dropsies.—Dose, two ounces, twice a day; or half an ounce may be given every hour till stools are obtained.

SYRUP OF SENEKA.

Syrupus Senegæ. (AMER.; FENN.; HANN.; POL.; BORUS.; SUEC.; COXE.; NIEMANN.; VAN MONS)

- R. Root of Seneka, one ounce.
 Boiling Water, one pound and a half.

Boil down to ten ounces; strain, and add White Sugar, one pound and a half.

Make a syrup. (FENN.; HANN.; POL.; BORUS.; SUEC.; NIEMANN)

- R. Root of Seneka, one part.
 Water, twelve parts.

Infuse in a moderate heat, in a covered vessel, for several hours; then add to the slightly expressed liquor

White Sugar, eighteen parts.

Make a syrup. (VAN MONS)

- R. Root of Seneka, four ounces.
 Water, one pound.

Boil to the consumption of one half, and add to the decanted liquor,

Sugar, one pound.

Make a syrup. (AMER.; COXE)

GLYCYRRHIZED SYRUP OF POLYGALA.

Syrupus Senegæ cum Liquiritia. (AUGUSTIN)

- R. Tincture of Extract of Liquorice, one dram.
 Syrup of Seneka, one ounce.

Mix.—Dose, a tea-spoonful, every hour.

MOSCHATED SYRUP OF SENEKA.

Syrupus Senegæ moschatus. (AUGUSTIN)

- R. Syrup of Seneka, three ounces.
 Gum Ammoniac, triturated with
 mucilage, one scruple.
 Musk, ten grains.
 Tincture of Extract of Liquorice,
 from fifteen to twenty drops.

Mix.—A spoonful every hour.

PECTORAL DECOCTION. (HAMB.; BRERA;
RATIER)

- R. Root of Seneka, } of each,
 Iceland Moss, } half an ounce.
 Spring Water, } sixteen ounces.

Boil over a slow fire, till reduced to ten ounces. Add to the strained liquor,

Glycyrrhized Honey, one ounce.

Mix. (HAMB.)

- R. Root of Polygala amara, three ounces.
 Herb of Coltsfoot, } of each,
 ——— Veronica, } one ounce.
 Water, three pounds.

Boil down to two thirds. (RATIER)

Dose, three or four ounces, several times a day.—It may be sweetened with sugar or honey.

DIURETIC DECOCTION. (HAMB.)

- R. Root of Seneka, half an ounce.
 Squill, one scruple.
 Spring Water, twelve ounces.

Boil down to two thirds; strain, and add,

Nitric Spirit of Æther, two drams.

Tincture of Opium, two scruples.

Glycyrrhized Honey, one ounce.

Mix.

EMMENAGOGUE DECOCTION. (ELLIS)

- R. Root of Seneka, one ounce.
 Orange Peel, two drams.
 Water, one pound.

Boil down to two thirds with the root, then add the peel, and strain after a short time.—Dose, four ounces daily, in amenorrhœa.

PECTORAL LOTION.

Decoction pectorale corroborans, Haustus pectoralis incitans. (AUGUSTIN; BRERA)

- R. Root of Seneka, two drams.
 Water, a sufficient quantity

to obtain six ounces of decoction. Strain, and add,

Oxymel of Squill, one ounce.

Mix. (BRERA)

- R. Root of Seneka, two drams.
 Water, a sufficient quantity

to obtain eight ounces of decoction. Strain and add,

Camphor, triturated with mucilage,
one dram.

Mix. (BRERA)

- R. Root of Seneka, two drams.
 Boiling Decoction of Cinchona,

a sufficient quantity
to obtain seven ounces of infusion. Add to it when strained,Camphor, triturated with mucilage,
half a dram.Sulphuric Æther, thirty drops.
Syrup of Orange Peel, } of each,
———— Marshmallows, } one ounce.

Mix. (BRERA)

- R. Root of Seneka,
 from half a dram to one dram.
 ——— Liquorice, two drams.
 Boiling Water, three ounces.

Infuse, and add when strained,

Paregoric Elixir, one dram.
Syrup of Marshmallows, half an ounce.

Mix. (AUGUSTIN)

Dose, one spoonful.

POLYGONATUM ANGULOSUM, DESF.—SOLOMON'S SEAL.

SYNONIMES:—Sceau de Salomon, FR.; Weisswurz, Weisswurzthal blume, GERM.; Salomon's segel, DAN.; Selli de Salomon, SPAN.; Salomons zege, DUT.; Sigillo de Salomone, ITAL.; Korkoryczka, POL.; Scella di Salomee, PORT.; Salomon's Sigil, SWED.

BRUNS.; HISP.; GALL.; WIRTEM.; BERGIUS.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN.

A plant which is met with in nearly every

part of Europe. (*Hexandria Monogynia*, LINN.; *Asparagineæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 171*)

The root (*Radix Sigilli Salominis* seu *Polygonati majoris* seu *latifolii* seu *vulgaris* seu *Geniculatæ* seu *Genicellæ*) is employed. It is articulated, long, rather fibrillous, whitish without, white within, and of about the diameter of the little finger. It presents a kind of seal-print on its joints. Its odour, rather disagreeable, disappears by desiccation. Its taste is sweetish, mucilaginous, and rather acrid.

POLYGONUM.

Five species of this genus are employed in medicine :

1. *Polygonum aviculare*, LINN.—Knotgrass.

SYNONIMES :—Renouée, Centinode, Frainasse, FR. ; Vogelknöterich, Vogelwegtritt, Wegrass, GERM. ; Varkengross, DUT.

BRUNS. ; HISP. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A common plant of Europe. (*Octandria Trigynia*, LINN. ; *Polygonæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 468*)

The herb (*Herba Centumnodiæ seu Centinodiæ seu Polygoni seu Sanguinariæ seu Sanguinalis*) is employed. It consists of a nodose, branching, smooth stem, having alternate, lanceolate, nearly sessile, entire, smooth leaves. It is destitute of odour, but has a sharp and astringent taste.

2. *Polygonum Persicaria*, LINN.—Spotted Persicaria.

SYNONIMES :—Persicaire douce, FR. ; Gemeiner Knöterich, GERM. ; Loppegræs, DAN. ; Persicaria monchada, SPAN. ; Persenkruid, DUT. ; Pessigueira, PORT. ; Røknesva, SWED.

HISP. ; GALL. ; SPIELMANN.

A plant of Europe. (*Fig. CEd. Fl. Dan. t. 702*)

The herb (*Herba Persicariæ minoris*) is employed. It consists of a round, fistulous, reddish, nodose, branching stem, and of alternate, lanceolate, petiolated leaves. Its taste is slightly acid and rather saline.

It was formerly esteemed vulnerary.

3. *Polygonum Hydropiper*, LINN.

SYNONIMES :—Persicaire poivrée, P. brulante, P. âcre, Poivre d'eau, Piment brulant, FR. ; Bitterknöterich, Wasserpfefferknöterich, GERM.

GALL. ; GENEV. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of Europe. (*Fig. Blackw. Herb. t. 119.*)

The (*Herba Hydropiperis seu Persicariæ urentis seu Mercurius terrestris*) is employed. It consists of a smooth, round,

nodose, branching stem, and of alternate lanceolate leaves. It is destitute of odour, but has an exceedingly acrid, burning, and peppery taste.

When fresh, it acts as a rubefacient, applied to the skin.

DIURETIC SPECIES.

Species pro Infuso diuretico. (AUGUSTIN)

R. Leaves of Spotted Persicaria,	} of each, half an ounce.
Root of Wild Rush,	
Leaves of Parsley,	} of each, one dram.
Herb of Hyssop,	
Cinnamon,	
Orange Peel,	
Anise,	

Cut and mix.

4. *Polygonum Bistorta*, LINN.—Bistort, Snakeweed.

SYNONIMES :—Bistorte, FR. ; Natterknöterich, wiesknöterich, Natterwurzknoeterich, GERM. ; Slangeurt, DAN. ; Bistorta, ITAL. ; PORT. ; and SPAN. ; Natterwortel, Slangenwortel, Hartstongue, DUT. ; Wezownik, POL. ; Ormrot, SWED.

DUBL. ; EDINB. ; AMST. ; BATAV. ; BELG. ; BRUNS. ; HISP. ; GALL. ; FENN. ; FERR. ; FORM. HÔP. FR. ; GENEV. ; HANN. ; HASS. ; LIPP. ; LUSIT. ; ROSS. ; SAX. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; COKE ; GUIBOUT ; MURRAY and GMELIN ; PARIS ; PIDERIT ; SPIELMANN.

A plant common to nearly all Europe. (*Fig. Flore Medic. II. 71.*)

The root (*Radix Bistortæ seu Colubrinæ seu Serpentinæ vulgaris seu rubræ*) is employed. It is of about the length of the finger, hard, subcylindrical, twice or thrice twisted, marked with annular intersections, furnished with numerous fibrillæ, of a blackish-brown colour without, and reddish or rose coloured within. It is destitute of odour or nearly so, but its taste is austere and very astringent, particularly in the fresh state.

It contains much tannin and starch, gallic acid, and oxalic acid.

It is powerfully astringent.—Dose, of the powder, from half a dram to one dram ; of the juice, from two to three ounces.

EXTRACT OF BISTORT.

Extractum Bistortæ. (HISP. ; PALAT. ; SARD.)

R. Root of Bistort, sliced, one pound.
Boiling Water, six pounds.

Digest for three days ; then boil slightly ; evaporate the strained decoction to the consistency of an extract. (PALAT.)

SARD. directs the infusion to be reduced to two thirds before straining and evaporating it.

R. Root of Bistort, sliced, two pounds.
Water, twenty pounds.

Infuse for twenty-four hours in a warm place, then boil a little, and strain ; boil the residue with a fresh quantity of water, till reduced to eight pounds, and strain. Mix, set aside, and decant the liquors ; then evaporate it, first over a slow fire, and then in a water-bath. (HISP.)

Dose, from a scruple to a dram.

DECOCTION OF BISTORT. (RATIER)

R. Root of Bistort, one ounce.
Water, two pounds.

Boil, and strain.

It is prescribed as an astringent in internal passive hæmorrhages.

ASTRINGENT LAVEMENT. (RATIER)

R. Root of Bistort, one ounce.

Poppy Heads, half an ounce.
Water, two pounds.

Infuse.—It is used in chronic diarrhœa and dysentery.

STOMACHIC POTION. (RATIER)

R. Bistort, one dram.
Rob of Elder, three drams.
Simple Syrup, one ounce.
Water, four ounces.

Infuse, and strain.

ASTRINGENT POTION. (RATIER)

R. Root of Bistort, two drams.
Syrup of Quince, one ounce.
Tincture of Catechu, two drams.
Water, four ounces.

Infuse, and strain.

5. *Polygonum Fagopyrum*, LINN.—Buck-wheat.

SYNONIMES : — Sarrazin, Blé noir, FR. ; Heydenkorn, Gemeiner Buchweizen, GERM.

GALL. ; WIRTEM. ; BERGIUS ; SPIELMANN.

A plant of Asia Minor. (*Fig. Mart. Fl. Russ. t. 46*)

The seed (*Semen Fagopyri* seu *Tritici fagini*) is employed. It is triangular, smooth, of the size of a hemp-seed, of a blackish-brown colour, and ferruginous taste.

POLYPODIUM VULGARE, LINN.—COMMON POLYPODY.

SYNONIMES : — Polypode commun, FR. ; Engelsüsstapfelfarren, Engelsuss, Süss-faren, Tufelfarren, GERM. ; Osladyc, BOHEM. ; Engelsædl, DAN. ; Polipodio, SPAN. ; Engelzæd, DUT. ; Felce quercina, ITAL. ; Raprotka, POL. ; Stensœta, SWED.

AMST. ; ANT. ; AUSTR. ; BAVAR. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A common plant of Europe. (*Crypto-*

gamia, LINN. ; *Filices*, JUSS. ; *Fig. Flore Medic. V. 278*)

The stock, improperly called the root (*Radix Polypodii* seu *Filiculæ dulcis*) is employed. It is hard, thick, ligneous, covered with numerous reddish, membranous scales, and furnished with blackish fibres. It is nearly inodorous ; its taste, at first saccharine, is bitter, nauseous, and slightly astringent.

It contains, according to BUCHOLZ, saccharine matter, gum, resin, oil, and starch.

It is feebly astringent, and is especially recommended in catarrhal disorders. It is also employed against worms, and forms the basis of HERRENSCHWAND'S remedy. —Dose, one or two drams of the powder.

POLYPORUS OFFICINALIS, Mic.—AGARIC, FUNGUS OF THE LARCH.

SYNONIMES:—Agaric blanc, Fr.; Lærchen-schwamm, GERM.; Bjla drjnowa hauba, BOHEM.; Agarico del alcerce, SPAN.; Lorkenswam, DUT.; Agarico del tarice, ITAL.; Gebka modrzewowa, POL.; Lærkeswamp, SWED.

AMST.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; PIDERIT; SAUNDERS; SPIELMANN; ZARDA.

A parasitical mushroom (*Agaricus albus* seu *Fungus Laricis*) which grows at the base and middle of the trunk of the larch. (*Fig. Flore Medic. 1. 6*)

It forms irregular masses, usually conical, triangular; slightly convex, marked with fawn-coloured zones, and brown or orange coloured above, of a yellowish-white and furnished with very small pores underneath. Its substance is yellowish-white and tenacious.

It is prepared for use by removing its exterior, which is very bitter, exposing it to the sun for several weeks, and then beating it with a mallet. It then appears white, light, of a homogeneous texture, and friable. It is without odour; its taste is first vapid and farinaceous, but soon becomes bitter, acrid, and nauseous.

In one thousand parts it contains, according to BUCHOLZ, five hundred of resin, thirty of a bitter principle, sixty of gummo-mucous matter, and three hundred and six of fibrine. The resin constitutes its active principle.

It is purgative, but not much used now.—Dose, ten grains.

TROCHISCS OF AGARIC.

Agaricus trochiscatus, Trochisci de Agarico. (BRUNS.; HISP.; GALL.; PALAT.; SARD.; WIRTEM.)

R. Agaric, eight parts.
Gum Tragacanth, one part.

Reduce the gum to thick mucilage with a sufficient quantity of water, then add the agaric, and make a soft mass, to be divided into trochiscs divisible into a very fine powder. (GALL.)

R. Agaric, four ounces.

White Wine, a sufficient quantity.

Triturate together, and form the paste into trochiscs. (HISP.)

BRUNS.; PALAT.; and WIRTEM. prescribe wine in which ginger has been infused.—WIRTEM. adds besides, a little gum tragacanth, and recommends the trochiscs to be made in winter, that they may be the whiter.—SARD. prescribes three ounces of agaric, one dram of ginger, and a sufficient quantity of mucilage of tragacanth.

DRASTIC POTION. (BORIES; PIERQUIN)

R. Trochiscs of Agaric,	}	of each,
Senna,		one dram.
Indian Jalap,		
Supertartrate of		
Potass,	}	of each,
Socotrine Aloes,		fifteen grains.
Cinnamon Powder,		twelve grains.
Boiling Water,		six ounces.

Infuse for twelve hours; then strain, and add,

Syrup of Buckthorn, one ounce.

Mix.

AQUEOUS EXTRACT OF AGARIC. (GALL.)

R. Agaric, one pound.
Cold Water, four pounds.

Macerate for twenty-four hours, occasionally shaking, and strain; macerate the residue in fresh water, and strain again. Mix the liquors, and evaporate in a water-bath, to the consistence of extract.

ALCOHOLIC EXTRACT OF AGARIC. (HISP.; WIRTEM.)

R. Agaric, one pound.
Alcohol, a sufficient quantity.

Digest for three days in a covered vessel and in a warm place; then express, and distil off the alcohol slowly by the heat of a water-bath, (WIRTEM.)

HISP. directs one pound of agaric to be digested for twelve hours in eight pounds of boiling water and four ounces of alcohol; then to boil for two hours, to express the liquor when half cold, to set it aside to depurate, and finally to evaporate it, first at the open fire, and then in a water-bath.

This extract is preferable to the preceding and is much more active.—Dose, from fifteen grains to a scruple. It is seldom given alone, but combined with other substances.

POLYTRICHUM COMMUNE, LINN.—GREAT GOLDEN MAIDENHAIR,

SYNONIMES : — Polytric commun, FR. ; Goldner Wiederthorn, Gùldenwiederthorn, Gùldenhaarmoss, GERM. ; Gùldhaar, DAN. ; Golden Wederhood, DUT. ; Bærmossa, SWED.

BRUNS. ; GALL. ; WIRTEM. ; BERGIUS ; GUIBOUT ; MURRAY and GMELIN ; SPIELMANN.

A plant common in all Europe. (*Cryp-*

togamia, LINN. ; *Musci*, JUSS. ; *Fig. Zorn*, *Ic. pl.* t. 415)

The herb (*Herba Adianthi aurei* seu *Polytrichi aurei* seu *Musci capillacei major*) is employed. It consists of a simple, proliferous stem, and of linear, lanceolated, toothed leaves. It has neither odour, nor taste.

It is feebly astringent.

POPULUS.

Three species of this genus of plants are mentioned in pharmacopœias.

1. *Populus balsamifera*, LINN.—Balsam Tree.

SYNONIMES : — Peuplier baumier, FR. ; Balsamespe, GERM. ; Balsam Popelier, DUT.

GALL. ; FORM. HÔP. FR. ; LIPP. ; SAX. ; BERGIUS ; GUIBOUT ; MURRAY and GMELIN ; PIDERIT.

A tree of North America and Siberia. (*Diœcia Octandria*, LINN. ; *Amentaceæ*, JUSS. ; *Fig. Cat. Car. I.* p. 34. t. 34)

The buds are employed ; they are rather large, oblong, acute, imbricated, with firm, oblong, concave scales. Their surface is covered with a limpid resinous juice of a golden colour, of an aromatic, warm, and bitterish taste.

2. *Populus Nigra*, LINN.—Poplar Tree.

SYNONIMES : — Peuplier noir, Peuplier franc, FR. ; Pappel, GERM. ; Topol, BOHEM. ; Poppel, DAN. ; Chopo, SPAN. and PORT. ; Popelier, DUT. ; Pioppo, ITAL. ; Topola Czarna, POL. ; Asp, SWED.

DUBL. ; AMST. ; ANT. ; BATAV. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FORM. HÔP. FR. ; GENEV. ; HANN. ; SAX. ; SUEC. ; WIRTEM. ; BRUGNATELLI ; GUIBOUT ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

A common tree of Europe. (*Fig. Zorn*, *Ic. pl.* 577)

The buds (*Oculi* seu *Gemmæ Populi*.—Pappelknospen, Pappelknœpfe, GERM.) are

employed. They are rounded, long, pointed, of a yellowish-brown colour, composed of scales of various sizes, lying one upon the other, shining at the surface, covered with a viscous juice of a golden-yellow colour and transparent. They have a penetrating, balsamic, and very agreeable odour, and a bitter, warm, and aromatic taste.

3. *Populus tremula*, LINN.—Trembling Poplar Tree.

SYNONIMES : — Tremble, FR. ; Zitternde Pappel, Gemeine Espe, GERM.

SUEC.

A tree of Europe. (*Fig. Nouv. Duh. II.* t. 87)

The twigs of the root are employed ; they are astringent.

TINCTURE OF POPLAR.

Tinctura seu *Essentia Populi* seu *Oculorum Populi*. (LIPP. ; VAN MONS)

R. Fresh Poplar Buds, one part.
Alcohol, six parts.

Infuse without heat, and preserve the filtered liquor. (VAN MONS)

LIPP. prescribes one part of buds and two parts of alcohol.

VULNERARY ESSENCE OF POPLAR.

Essentia Populi vulneraria. (WIRTEM.)

R. Fresh Poplar Buds, half a pound.
Concentrated Alcohol, one pound and a half.

Digest for several days, and express. Add to the filtered liquor,
3 G

Essence of the Flowers of St.
John's-wort, half a pound.

Digest for some time; then distil off one half of the alcohol, and preserve the remainder.

It is excitant, and is recommended externally to atonic ulcers, and internally, in gonorrhœa, leucorrhœa, diarrhœa, and phthisis pulmonalis.—Dose, from thirty to fifty or more drops.

POPLAR OINTMENT.

Unguentum Populeum. (FERR.; LIPP.; BRUGNATELLI)

R. Dried Buds of Balsam Tree,
half a pound.

Fresh Butter, one pound.

Boil slowly to the consumption of the humidity, and express. (LIPP.)

BRUGNATELLI directs eight ounces of dried buds to be bruised, two pounds of melted lard to be added; the mixture to be kept fluid in a water-bath for three hours, and then decanted.—FERR.—two pounds of fresh buds, bruised, to be digested for eight days in four pounds of lard.

This is the only ointment which really deserves the term populeum.

PORTULACA OLERACEA, LINN.—PURSLAIN.

SYNONIMES:—Pourpier commun, FR.; Portulak, DAN.; GERM.; and SWED.; Berdolaga, SPAN.; Porselein, DUT.; Porcellana, ITAL.; Kurza noga, POL.; Beldroega, PORT.

AMST.; BRUNS.; HISP.; GALL.; WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A cultivated plant of Europe. (*Dodecandria Monogynia*, LINN.; *Portulacæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 489*)

The herb and seeds are employed.

The herb (*Herba Portulacæ*) consists of a rounded, smooth, shining, succulent stem, and of oblong, cuneiform, fleshy, smooth leaves, of a deep green colour. Its taste is watery and feebly aromatic.

The seeds are small, reniform, curved, rather wrinkled, black, and tasteless.

The seeds are classed among the four minor cold seeds. As to the herb, which is

rather culinary than medicinal, it purges when a large quantity is taken at once.

CONSERVE OF PURSLAIN.

Conserva Portulacæ. (VAN MONS)

R. Leaves of Purslain, one part.
White Sugar, two parts.

Bruise the leaves, add the sugar, and heat slightly.

WATER OF PURSLAIN. (GALL.; PALAT.)

R. Herb of Purslain, five parts.
Water, twelve parts and a half.

Distil about ten parts; pour the product upon ten parts of herb, add ten parts of water, and distil ten parts again. The operation may be repeated a third time. (GALL.)

PALAT. prescribes one part of herb and four parts of water; one and a half to be distilled.

POTASSIUM.

This metal is not used in medicine, but its numerous compounds are applied to many purposes.

PROTOXIDE OF POTASSIUM.

Potassa caustica seu pura, Kali vegetabile purum seu causticum.

I. In the liquid state.

Aqua Potassæ seu Potassæ puræ seu Kali puri Lixivia caustica, Liquor Potassæ seu Kali caustici seu Potassæ causticæ, Lixivium causticum seu magistrale seu saponarium seu causticum vegetabile.—Solution of Potass, Solution of caustic Potass, Liquid Potass, Soap ley. (LOND.; DUBL.; EDINB.; AMER.; AMST.; BATAV.; BAVAR.; BELG.; GALL.; FERR.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Liquor Potassæ. (LOND.)

R. Subcarbonate of Potass,
one pound.

Fresh-burnt Lime, half a pound.
Boiling Distilled Water, one gallon.

Dissolve the subcarbonate in two pints of the water. Pour the remainder of the water upon the lime, and add the solution. Set apart the mixture in a covered vessel, and when cold, strain it through a cotton bag. If, by adding any diluted acid, effervescence is excited, more lime must be added, and the filtration repeated. A pint of this solution should weigh sixteen ounces.

Aqua Potassæ. (EDINB.)

R. Fresh-burnt Lime, eight ounces.
Subcarbonate of Potass, six ounces.
Boiling Water, twenty-eight ounces.

Put the lime into an earthen or iron vessel with twenty ounces of the water. When the ebullition has ceased, add the subcarbonate dissolved in the remainder of the water, and having thoroughly mixed the whole, cover the vessel till the mixture has become cold; then agitate it well, and pour it into a glass funnel the tube of which is obstructed by a piece of clean linen. Cover the upper opening of the funnel, while its tube is inserted into another glass vessel, that the liquor may gradually drop through the linen into the lower vessel. When it first ceases to drop, pour a few ounces of water into the funnel, but cautiously, so that the matter may be disturbed as little as possible, continue to pour on water thus till three pounds of liquor have filtered through, which will be in the space of two or three days; then shake the solution and preserve it in a well stopped vessel.

Aqua Potassæ causticæ. (DUBL.)

R. Fresh burnt Lime, } of each,
Carbonate of Potass, } two parts.

Water, fifteen parts.

Pour upon the lime, in an earthen vessel, one part of the water made hot, and when it is slaked, add the subcarbonate and the remainder of the water. Pour the mixture when cold into a well stopped bottle, and keep it for three days, frequently shaking it. When the carbonate of lime shall have subsided, decant the supernatant liquor and preserve it in well stopped green glass bottles.

The specific gravity of this solution is to that of distilled water, as 1080 to 1000.

There are nearly as many variations in this process, but none of much importance, as there are pharmacopœias.—GALL. says mix together two parts of subcarbonate, one part of lime, and twelve parts of water; strain the liquor, concentrate it till it marks 36 degrees by the areometer, set it aside, and then decant it: its weight will be 1.334, and a phial of the capacity of three ounces will contain four ounces of it.—FERR.; GENEV.; COXE; and SWEDIAUR give the same as EDINB.—AMST.; and BELG. say pour four ounces of hot water upon nine of quick lime, and add when it is slaked, six ounces of subcarbonate; pour the whole into a funnel stopped with cotton, and when no more passes through, add boiling water so as to obtain three pounds of liquor.—ROSS.—extinguish gradually four ounces of lime with a little hot water, add eighteen ounces of hot water and then three of subcarbonate of potass, agitate strongly, and when cold add a sufficient quantity of water to obtain eighteen ounces of liquor.—SAX.—dissolve subcarbonate of potass in six times its weight of water; then apply heat and add lime gradually, ascertaining from time to time if the liquor dropped into an acid produce effervescence: cease to add lime when this no longer takes place, and strain through linen with expression: wash the residuum with a little hot water, express again, set aside, pour off, and concentrate till the liquor marks 36 degrees by the areometer.—LIPP.—extinguish four pounds of lime with a sufficient quantity of boiling water, and then add sixteen ounces of subcarbonate dissolved in nine pounds of water; set apart for twenty-four hours, test the liquor alternately with lime and sulphuric acid; then, when it is no longer rendered turbid by either of them, filter it in a funnel furnished with cotton, and, if required, evaporate it till a vessel of the capacity of one pound contains sixteen ounces of it.—FULD.—extinguish six ounces of lime

with a sufficient quantity of water, add three ounces of subcarbonate dissolved in one pound and a half of water, and filter after some time in a funnel furnished with cotton, adding to the liquor which passes a sufficient quantity of water to raise its weight to one pound and a half.

R. Subcarbonate of Potass, two parts.
Lime, three parts.

Mix together, and add

Rain water, sixteen parts.

Boil for half an hour, frequently stirring; then strain through a double linen cloth, expressing slightly towards the end. Pour upon the residuum,

Water, four parts.

Evaporate the two liquors mixed together till only eight parts remain; set this aside in a large well stopped bottle, decant the clear liquor, and preserve it in small phials. (BATAV.)

HASS.; and PIDERIT direct six pounds of spring water to be poured upon one pound of quick lime, and when this is slaked, half a pound of subcarbonate of potass to be added; the liquor to be then boiled till a goose feather thrown into it becomes corroded and dissolved, then to be diluted with a sufficient quantity of water, and filtered. —FENN.; HANN.; OLDEN.; POL.; BORUS.; SUEC.; and VAN MONS—two pounds of subcarbonate to be dissolved in ten of water, three of powdered lime to be gradually added to the solution, and the mixture boiled for a quarter of an hour, stirring it with a spatula; then to set it aside for some time, and the liquor to be strained and evaporated till it has acquired the specific gravity of 1.337. —BAVAR.—one part of subcarbonate to be dissolved in ten parts of water, the solution to be boiled, and two thirds of a part of pure lime to be added, constantly stirring; after a quarter of an hour's boiling, the liquor is to be strained through linen, and the residuum to be boiled with five parts of water; the two liquors to be mixed together and set apart for twenty-four hours, and then to be decanted and evaporated till of the specific gravity of 1.33.

BATAV. also proposes to dissolve one part of fused potass in two parts of distilled water; which is more simple and expeditious, and is also preferable with respect to the quality of the product.

II. In the solid state.

Alkali causticum, Causticum alkalinum seu commune seu commune fortius seu potentiale seu salinum, Cauterium potentiale, Kali causticum siccum, Hydras kalicum seu Potassa pura fusa, Lapis causticus seu causticus Chirurgorum seu infernalis

alkalinus seu septicus, Potassa fusa seu pura seu caustica seu caustica fusa, Ruptorium commune.—Fused Potass, Caustic Potass, Hydrate of Protoxide of Potassium.—(LOND.; DUBL.; EDINB.; AMER.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Potassa fusa. (LOND.)

R. Solution of Potass, one gallon.

Evaporate it in a clean iron vessel over the fire, till the ebullition has ceased and the potass has become melted; then pour it out upon a clean iron plate, and cut it into pieces of a proper form.

Potassa, olim Causticum commune acerrimum. (EDINB.)

R. Solution of Potass, any quantity.

Evaporate it in a very clean covered iron vessel, till, the ebullition having ceased, the saline matter flows smoothly like oil, which happens before the vessel becomes red hot. Then pour it upon a clean iron plate, cut it into small masses before it hardens, and preserve them in well-stopped phials.

Potassa caustica. (DUBL.)

R. Solution of Potass, any quantity.

Evaporate it over the fire in a clean iron vessel, till, the ebullition having ceased, the saline matter, on the heat being increased, remains quiescent in the vessel. Pour out the fused salt upon a clean iron plate, and while it is consolidating, cut it into proper pieces, and enclose them immediately in a well-stopped bottle. During the evaporation the operator should avoid the particles which may be thrown from the vessel.

R. Solution of Potass, any quantity.

Evaporate it till a drop let fall upon a cold stone immediately hardens; then put

the mass into a crucible, and heat it till it flows like oil; then pour it into moulds of the size of a goose-quill. (AMER.; BATAV.; BRUNS.; DAN.; FERR.; FENN.; FULD.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR)

R. Quick Lime, three parts.

Having slaked it with a sufficient quantity of water, add

Pure Subcarbonate of Potass, one part.

Diffuse the whole in much water, stir well, filter, and evaporate to dryness; then fuse it, and pour it either into greased iron moulds, or upon a warm slab: break it into pieces when cold. (HERBIP.)

HISP. directs two pounds of quick lime, one pound of subcarbonate of potass, and sixteen pounds of water, to be digested and shaken together; the liquor to be filtered, and evaporated to dryness, and the residuum fused and poured upon a hot slab.

R. Subcarbonate of Potass, two parts.
Lime, in coarse powder, one part.
Water, twelve parts.

Boil for half an hour; then strain, and evaporate to dryness: fuse the residuum in a crucible, and pour it upon a hot slab. (GALL.; FORM. HÔP. FR.)

BRUGNATELLI directs one pound of subcarbonate, three pounds of lime, and sixteen of water to be boiled together for a quarter of an hour, the liquor to be evaporated to dryness, the residuum to be fused and poured upon an oiled stone.—AUSTR.—one pound of subcarbonate to be dissolved in ten pounds of boiling water, and two pounds of quick lime to be added, constantly stirring; the mixture to be boiled for an hour, then set aside, decanted, evaporated till the matter flows like oil, and then poured either into moulds or upon a hot iron plate.—SARD.—four pounds of calcined ashes of vine twigs, and two of quick lime, to be boiled with ten times the quantity of water till reduced to one half; the liquor to be strained, and evaporated to dryness, and the residuum melted in a crucible and poured upon a warm oiled stone.—SPIELMANN—one part of subcarbonate, four parts of quick lime, and twenty times the weight of water to be boiled together for several hours; the liquor to be filtered, evaporated to dryness, and the residuum slowly fused, and poured into moulds.

R. Crystallized Caustic Potass, any quantity.

Put it by portions into a silver crucible; after the dissipation of the water of crystallization, when it is equally fused and tran-

quil, pour it into a warmed iron mould (VAN MONS)

VAN MONS recommends that the moulds be not oiled into which the fused potass is poured.

Some chemists prefer to pour the potass, by drops, upon an oiled slab. (PLANCHE)

III. In a state of perfect purity.

ALCOHOLATED POTASS.

Potasse à l'alcohol. (GALL.)

R. Solution of Potass, any quantity.

Having evaporated it to dryness, dissolve it in alcohol (40 degrees), and then remove part of this alcohol by distillation; evaporate the residue in a silver pan, remove with a spoon the blackish pellicle which covers the liquor, pour what remains upon a silver plate, and, when cold, break it into pieces.

The product is pure hydrate of protoxide of potassium. The fused potass contains salts of potass, sulphates, hydro-chlorates, &c. The pure solution of potass is corrosive and caustic in its concentrated state, but is only strongly stimulant when diluted with a large quantity of water. It has been employed in this form as an antacid, diuretic, and lithontriptic, in the dose of from ten to fifteen drops in a proper vehicle. The fused potass is used to open issues.

CAUSTIC POTASS WITH LIME.

Potassa cum Calce, Causticum commune mitius. (LOND.; DUBL.; EDINB.; AMER.; GENEV.; COXE; HENRI; SPIELMANN)

Potassa cum Calce. (LOND.)

R. Solution of Potass, three pints.
Fresh-burnt Lime, one pound.

Boil the solution down to a pint; then add the lime slaked with water, and mix intimately.

Potassa cum Calce, olim Causticum commune mitius. (EDINB.)

R. Solution of Potass, any quantity.

Evaporate it to one third in a covered iron vessel; then mix with it as much recently slaked lime as will bring it to the consistence of a solid paste, which is to be preserved in a well-stopped vessel.

Potassa caustica cum Calce. (DUBL.)

Evaporate solution of caustic potass to one fourth; then add as much fresh-

burnt lime, in powder, as will form a mass of a proper consistence, which is to be preserved in a well-stopped bottle.

AMER.; GENEV.; and COXE direct the solution to be evaporated to one third, and a sufficient quantity of quick lime added to obtain a firm paste.—SPIELMANN—one part of caustic potass to be dissolved in three parts of water, and a sufficient quantity of quick lime added as will form a paste.

HENRI says that before pouring the ordinary fused potass, a little powdered quick lime should be added, which prevents it from attracting the humidity of the atmosphere so readily, and brings it near to the present preparation.

This caustic does not act so strongly as the preceding.

ELSE'S CAUSTIC PASTE. (WIRTEM.; NIEMANN)

R. Solution of Potass, eight ounces.

Boil down to two ounces, and add to the boiling liquor,

Extract of Opium, two drams.
Quick Lime, six drams,

or, a sufficient quantity to absorb all the liquid.

SAUNDERS' LITHONTRIPTIC LIQUOR.
(CADET DE GASSICOURT)

R. Calcined Potass of Commerce, half a pound.

——— Oyster Shells, one pound and a half.

Boiling Water, eight pounds and a half.

Set aside for twenty-four hours, and filter.

Three drams to be given daily at three doses, in an infusion of linseed, to which fifteen grains of calcined magnesia are added.

BRUNNINGHAUSEN'S FOMENTATION.
(NIEMANN)

R. Fused Potass, one ounce.
Water, forty-eight ounces.

Dissolve.—This solution has been applied, it is said, with success, to the abdomen in a case of convulsions in a pregnant woman.

GIRTANNER'S INJECTION. (NIEMANN)

R. Caustic Potass, half a dram.
Distilled Water, six ounces.

Dissolve.—It is recommended by GIRTANNER, at the commencement of gonorrhœa, to arrest it.

GIMBERNAT'S COLLYRIUM. (CADET DE GASSICOURT; NIEMANN; PIERQUIN; SWEDIAUR)

R. Caustic Potass, one grain.
Distilled Water, one ounce.

Dissolve. (NIEMANN; SWEDIAUR)

CADET DE GASSICOURT; and PIERQUIN double the quantity of the potass.

A few drops are inserted into the eye affected with speck of the cornea, and the organ then washed with a mucilaginous decoction.

COHEN'S SOLUTION. (AUGUSTIN; NIEMANN)

R. Solution of Potass, from ten to twenty drops.
Decoction of Juniper, one pound.

Mix.—To be taken in the course of the day.

RESOLVENT OPHTHALMIC LIQUOR.
(HAMB.)

R. Caustic Potass, one grain.
Distilled Water, four ounces.
Extract of Opium, four grains.

Dissolve, and filter.

SAVIARD'S SOLUTION. (NIEMANN)

R. Caustic Potass, two drams.
Camphor, two scruples.
White Sugar, two ounces.
Spring Water, two pounds.

Mix.—As a lotion to indolent ulcers and filled with luxuriant growths.

BOUGIES OF CAUSTIC POTASS.

Cereoli cum Kali caustico. (AUGUSTIN)

R. Caustic Potass, four grains.
Extract of Opium, one dram.
Water, two ounces.

Gum Arabic, a sufficient quantity to thicken the liquor. Dip threads into this, dry them, and continue to do so till the bougies are sufficiently large.

HECKER recommended these soluble bougies in chronic gonorrhœa.

ALCOHOLIC SOLUTION OF POTASS.

Tinctura Metallorum seu Antimonii tartarizata seu Stibii alkalizata seu Antimonii acris seu Antimonii alkalina seu Antimonii nigra seu Antimonii balsamica seu Salis Tartari seu Tartari seu alkalina seu kalina seu alkalina acris, Alcohol Potasse seu alkalisatum, Spiritus potassatus. (BAVAR.; BRUNS.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.;

SARD.; SAX.; WIRTEM.; HERBIP.;
SPIELMANN; SWEDIAUR; VAN MONS)

R. Martial Regulus of Antimony,
two ounces.

Copper, } of each, one ounce.
Tin, }

Fuse together, pulverize the amalgam, and detonate it by small portions with twelve ounces of nitre; increase the heat so that the residuum may be reduced to greenish scorïæ: then pulverize it, and add

Alcohol, one pound and a half.

Filter the liquor when it is of a fine red colour. (BRUNS.; SPIELMANN)

R. Martial Regulus of Antimony,
three ounces.

Nitrate of Potass, eighteen ounces.

Project the mixture by portions into a red hot crucible, and after the deflagration, calcine for an hour; then fuse the mass, pour it into an iron mortar rubbed with chalk, and rub it to powder. Put this while hot into a matrass, and pour upon it

Concentrated Alcohol,
forty-eight ounces.

Digest for three days; then decant and filter. (FULD.)

LIPP. prescribes one ounce and a half of regulus, nine ounces of nitre, and two pounds of alcohol.—WIRTEM.—four ounces of regulus, twenty ounces of nitre, and one pound and a half of alcohol.—BRUNS.—one pound of regulus, one of nitre, and one and a half of alcohol.—SPIELMANN—four ounces of regulus, eight of nitre, and twelve of alcohol.

R. Sulphuret of Antimony, half a pound.
Subcarbonate of Potass, one pound.

Fuse the mixture in a crucible, and reduce it to yellowish scorïæ; then bruise it in a heated iron mortar, and pour upon the powder

Alcohol, three pounds.

Strain at the expiration of three days. (BRUNS.; PALAT.; SARD.; WIRTEM.; HERBIP.; SPIELMANN; SWEDIAUR)

R. Sulphuret of Antimony, six ounces.

Fuse it in a crucible, and add

Nitrate of Potass, one ounce and a half.

Having kept it on the fire for a quarter of an hour, pulverize the mass, and boil it with

Fixed Nitre, eight ounces,

dissolved in two pounds of water, till of the consistence of syrup; then add

Alcohol, one pound.

Digest, and decant. (BRUNS.)

R. Medicinal Regulus of Antimony,
while hot, and in powder,
four ounces.

Pour upon it a solution prepared with
Subcarbonate of Potass, six ounces.
Water, eight ounces.

Boil to the consistence of syrup, and add
Concentrated Alcohol, ten ounces.

Digest at a very moderate heat, frequently shaking, till the liquor becomes of a blackish-red colour. (FULD.)

R. Subcarbonate of Potass, six ounces.

Calcine it strongly in a crucible, and, having powdered it, pour upon it while yet hot,

Concentrated Alcohol, twelve ounces.

Leave it to digest in a sand-bath till the liquor becomes of a red colour. (BRUNS.; FULD.; PALAT.; SARD.; WIRTEM.; HERBIP.; SWEDIAUR)

SPIELMANN directs eight ounces of subcarbonate of potass to be calcined with three ounces of quick lime for an hour, and the residuum to be digested with twenty-four ounces of concentrated alcohol.—WIRTEM.—six ounces of subcarbonate to be strongly calcined with one ounce and a half of charcoal powder, and the residuum to be digested with one pound of alcohol.

R. Caustic Potass, one part.
Concentrated Alcohol, eight parts.

Digest for several days in a sand-bath. (BAVAR.; HANN.; OLDEN.; POL.; BORUS.; SAX.; VAN MONS)

This preparation is no longer used; the older practitioners used to prescribe it as stimulant, deobstruent, diuretic, and diaphoretic.—Dose, from twenty to sixty drops.

STAHL'S TONICO-NERVINE MIXTURE.

Mixtura tonico-nervina Stahlîi. (BRUNS.; FULD.; PALAT.; SAX.; WIRTEM.; NIEMANN; SPIELMANN)

R. Alcoholic Solution of Potass,
two ounces.
Spirit of Hartshorn, one ounce.

Mix.

It is excitant, resolvent, nervine, antispasmodic, diaphoretic, and diuretic.—Dose, from thirty to fifty drops.

SALINE TINCTURE. (SARD.)

R. Purified Potass, six ounces.
Fresh Orange Peel, } of each, one
Root of Gentian, } ounce and a half.
Distilled Water, one pound and a half.

Digest for four days, and add

Rectified Alcohol, eight ounces.

Digest again for twenty-four hours, and filter.—Dose, from one to two drams.

SPIRITUOUS RESOLVENT MIXTURE.

Mixtura resolvens spirituosā. (FULD.)

- R. Black Tincture of Antimony, one ounce.
 Tincture of Burnet Saxifrage, six drams.
 ————— Amber, three drams.
 Spanish Soap, two drams.

Mix, and dissolve.

This tincture is reputed stimulant, tonic, balsamic, resolvent, anti-spasmodic, nervine, diuretic, and diaphoretic.—Dose, from forty to seventy drops.

DIPPEL'S CHEMICAL SOAP. (WIRTEM.)

- R. Subcarbonate of Potass, one pound.
 Sulphuret of Antimony, half a pound.
 Copper Filings, three ounces.

Fuse the mixture, and powder it before it is cold; then mix it with

Oil of Juniper or of Turpentine,
 nine ounces.

Pour over the mass

Concentrated Alcohol,

a sufficient quantity

to cover it by several fingers' breadth. Let it digest for several days, and then decant the coloured liquor. Repeat the digestion two or three times with fresh alcohol; then, having mixed the tinctures, distil off part of the spirit, and add Venice soap and a little balsam of Peru.

This rather absurd preparation was recommended by DIPPEL, applied externally, as resolvent and anodyne, in gout and chilblains.

PROTOXI-SULPHURET OF POTASSIUM.

Hepar Sulphuris seu Sulphuris salinum seu Sulphuris vulgare, Kali sulphuratum, Sulphuretum Lixiviæ seu Potassæ seu Kalicum.—Sulphuret of Protoxide of Potassium, Sulphuret of Potass, Liver of Sulphur.—(LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; DAN.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS)

I. In the solid state.

Sulphuretum Potassæ.

- R. Washed Sulphur, one ounce.
 Subcarbonate of Potass,
 two ounces

Rub them together, and keep them

over the fire in a covered crucible till they unite. (LOND.)

- R. Subcarbonate of Potass,
 two parts.
 Sublimed Sulphur, one part.

Rub them together, and put them into a large covered crucible, to which apply heat cautiously till the mixture becomes fused. Preserve the mass in a well-stopped vessel. (EDINB.)

- R. Carbonate of Potass, four parts.
 Sublimed Sulphur, one part.

Mix, and put them into a crucible, then having adapted a cover to it, expose it to a gradually raised heat till they unite. (DUBL.)

- R. Sublimed Sulphur, one part.
 Subcarbonate of Potass, two parts.

Fuse together in a matrass, at a gradually augmented heat; and when the vessel is cold, break it, and immediately transfer its contents to a bottle. (AUSTR.; BATAV.; BRUNS.; DAN.; GALL.; FENN.; FORM. HÔP. FR.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; WIRTEM.; HERBIP.; COXE; SWEDIAUR)

It is necessary to heat the mass till it flows tranquilly like water, for so long as it swells it may be known to contain carbonic acid. The bottles into which it is to be put should be previously warmed in order to deprive them of air and moisture.

VAN MONS directs six parts and a half of subcarbonate of potass, in powder, to be triturated with three of sublimed sulphur, and the mixture fused in a crucible.—BAVAR.—four of subcarbonate and three of sulphur.—FERR.; LIPP.; SAX.; SUEC.; and BRUGNATELLI—an equal part of each. The latter proportions, also recommended by VAUQUELIN, give a mixture of sulphate of potass and of sulphuret of potassium, with an excess of sulphur.—BERAL employs fifteen parts of pearlsh and forty of sulphur: a principal inconvenience of which is that it furnishes a product which contains sulphuret of iron, which gives a greenish tinge to the skin.—HENRI has proposed to use two parts of subcarbonate of potass obtained by the combustion of tartar and one part of pulverized roll-sulphur; to fill a matrass with this mixture to two-thirds, to place it in a sand-bath, with at first a moderate heat, to be gradually raised till the mass becomes liquified.

- R. Roll-Sulphur,
 one part and three-fourths.

Fused Caustic Potass, five parts and a half.

Mix together, and fuse in a matrass placed in a sand-bath. (VAN MONS)

VAN MONS calls this *Simple Sulphuret of Potass*, which name has also been applied to the sulphuret obtained by the decomposition of the sulphate by charcoal at a high temperature. The preparation which he calls *Supersulphuret of Potass*, is obtained according to the following formula :

R. Roll-Sulphur, five parts and a half.
Fused Caustic Potass, three parts.

Fuse together in a glass matrass, placed in a sand-bath, to be gradually heated.

II. In the liquid state.

SOLUTION OF SULPHURET OF POTASSIUM.

Potassæ Sulphureti Aqua. (DUBL.)

R. Sublimed Sulphur, one part.
Solution of Caustic Potass, eleven parts.

Boil together for ten minutes, and filter through paper. Preserve the liquor in well-stopped phials. Its specific gravity is to that of distilled water as 1117 to 1000.

R. Sublimed Sulphur, half an ounce.
Solution of Caustic Potass, nine ounces.

Boil for ten minutes, and filter. (COXE)

The specific gravity of the liquor is 1.120.

The sulphuret of potass is excitant, stimulant, irritant, poisonous in a large dose, expectorant and diaphoretic in many cases, internally; and detergent, externally. It has been prescribed in asthma and croup, but without much advantage, in chronic catarrhs, chronic rheumatism, gout, cancer, tetters, itch, tinea, and many other cutaneous disorders. It has been represented as an antidote to the arsenical and saturnine poisons.—Dose, from five to fifteen or twenty grains.

The acids and the acidulous, metallic, and earthy salts are incompatible with it.

POWDER AGAINST SALIVATION. (BORIES; SAINTE-MARIE)

R. Sulphuret of Potass, four scruples.
Supertartrate of Potass, four drams.
Divide into twenty-four parcels.

HAHNEMANN has recommended this powder against mercurial salivation. The patient should take three parcels in the day, in a little sweetened water. Notwithstanding the praises which have been bestowed

on it, it is, as are all other derivatives, frequently inefficacious.

ALKALINE CHARCOAL PILLS.

Pilule Carbonum Kalinæ. (AUGUSTIN)

R. Sulphuret of Potass, one dram.
Extract of Blessed Thistle, } of each,
Charcoal, } two drams.

Mix.—Dose, from thirty to forty grains, four times a day, in phthisis pulmonalis.

SINGER'S ANTI-SIALAGOGUE SOAP.

Sapo sulphurato-crocatus. (BAVAR.)

R. Sulphuret of Potass, one part.
Boiling Water, two parts.

Dissolve, and add

Yellow Wax, one part.

Evaporate, constantly stirring, till the mass no longer adheres to the fingers; then pour it out into paper moulds.

JADELOT'S OINTMENT.

Linimentum saponaceum hydrosulphuretum.
(GALL; AUGUSTIN; CADET DE GASSICOURT; NIEMANN)

R. Sulphuret of Potass, six ounces.

Dissolve it, one day in advance, in one-third of its weight of water; then triturate it with a solution of

White Soap, two pounds,

in Oil of Clove-pink, four pounds.

Add

Essential Oil of Thyme, two drams.

Mix.—It is recommended in itch. One ounce to be rubbed in each time. (AUGUSTIN; CADET DE GASSICOURT; NIEMANN)

R. Common Soap, fifty parts.

Melt in a glass vessel placed in a water-bath, and add

Oil of Clove-pink, twenty-five parts.

Triturate intimately together, and add

Sulphuret of Potass, ten parts.

Mix well, and add

Oil of Clove-pink, seventy-five parts.

Incorporate. (GALL.)

RATIER prescribes six ounces of sulphuret, two ounces of water, and two pounds each of soap and of olive oil.

HENRI recommends the soap to be softened in a small quantity of water, which is to be again evaporated by the heat of a water-bath; the oil to be then added by portions, and the mixture stirred, still in the bath, till it is perfectly homogeneous; it is

then to be poured into a glazed pan, and the sulphuret gradually incorporated with it, constantly stirring with a wooden spatula.

This preparation may be resembled to ALIBERT'S *Sulphuro-alkaline Ointment*, which is also used in the treatment of itch, and which is prepared with two parts of sublimed sulphur, one part of potass, and eight parts of lard. (RATIER)

MARTIAL SULPHURET OF POTASS. (FERR.)

- R. Subcarbonate of Potass, } of each,
Sulphur, } one ounce.
Deutoxide of Iron, } two drams.

Fuse in a crucible, pour the mass upon a marble slab, and break it into pieces when cold.

To be taken in pills of three or four grains each; or, in the dose of a scruple in ten ounces of water; to be taken by spoonfuls.

RESOLVENT AND ANTI-CANCEROUS PLASTER. (BORIES)

- R. Sulphuret of Potass, } of each,
Powdered Hemlock, } two drams.
Camphor, } of each, four drams.
Turpentine, }
White Soap, half a dram.
Yellow Wax, one ounce.
Simple Plaster, four ounces.

Make a plaster.—It is recommended to enlarged lymphatic glands.

SOLUTION OF LIVER OF SULPHUR. (HAMB.)

- R. Sulphuret of Potass, half an ounce.
Distilled Water, three ounces.

Dissolve, and filter.

SULPHUREOUS POTION.

Aqua Kali sulphurati. (AUGUSTIN)

- R. Sulphuret of Potass, one dram.
Water, one pound.
Sugar, one dram.

Mix.—It is recommended, as an antidote, in poisoning by arsenic.

ANTI-HERPETIC POTION. (CADET DE GASSICOURT)

- R. Sulphuret of Potass, forty grains.
Water, one pound.

Dissolve, and filter.

SULPHUREOUS BATH.

Balneum sulphuratum. (BRERA; RATIER)

- R. Sulphuret of Potass, two ounces.
Water, a sufficient quantity
for a bath; about a hundred pounds.

DUPUYTREN'S SULPHUREOUS AND GELATINOUS BATH. (RATIER)

- R. Sulphuret of Potass, four ounces.
Water, two hundred pounds.

Pour into the solution

- White Flemish Glue, two pounds,
dissolved in

Boiling Water, ten pounds.

Mix.

RESOLVENT BATH. (SAINTE-MARIE)

- R. Sulphuret of Potass, one ounce.
Muriate of Soda, two ounces.
Soda, four drams.
Leaves of Sage, one or two handfuls.

Boil the whole, for twenty-four hours, in Water, nine pounds.

It is employed in lotions, douches, or fomentations, in spina ventosa, and scrofula. When immersion is possible, it should continue half an hour, and be repeated three or four times a day.

SYRUP OF SULPHURET OF POTASS.

Syrupus de Sulphureto Potassæ seu Sulphuris seu Kali sulphurati, Arcanum bechicum Willisii. (GALL.; WIRTEM.; AUGUSTIN; CADET DE GASSICOURT; NIEMANN; PLANCHE)

- R. Sulphuret of Potass, half an ounce.
Hyssop or Fennel Water, half a pound.
White Sugar, fifteen ounces.

Dissolve the sulphuret without heat; then add the sugar, and form a syrup in a water-bath. (GALL.)

PLANCHE prescribes two drams of sulphuret, eight ounces of fennel water, and fifteen of sugar.—CADET DE GASSICOURT—two drams of sulphuret, half a pound of fennel water, and one pound of sugar.

- R. Sulphuret of Potass, half an ounce.
Spring Water, } of each, four
Fennel Water, } ounces and a half.

Dissolve, and add

White Sugar, sixteen ounces.

Boil to the consistence of syrup, and strain. (WIRTEM.; AUGUSTIN; NIEMANN)

WILLIS directed two ounces of sulphuret to be digested in three pounds of Spanish wine for twenty-four hours; the liquor to be filtered, and two pounds of sugar dissolved in it. (CADET DE GASSICOURT)

BOULLAY and PLANCHE have proposed the following formula:

- R. Pure Soda, prepared with alcohol, one dram.
Distilled Water, five drams.

Apply heat, and add as much pure sulphur as it will dissolve. Take one ounce of

the liquid, mix it with thirty-one ounces of syrup, and agitate the mixture in a well stopp'd bottle.

It is recommended by CHAUSIER in catarrhal affections of the lungs and in croup. It excites the mucous secretions, sometimes vomits and purges, and disposes to perspiration.—Dose, a tea spoonful to children, and one or two table spoonfuls to old persons.

ALCOHOLIC SOLUTION OF SULPHURET OF POTASS. (VAN MONS)

R. Dry Sulphuret of Potass, any quantity.
Alcohol, a sufficient quantity.

Dissolve at a moderate heat. When cold, filter the solution, and preserve it in a bottle turned upside down.

VAN MONS again brings this preparation forward under the title of *Tinctura Sulphuris*, which he directs to be prepared by dissolving one part of sulphuret of potass in twelve parts of proof spirit.—SARD. directs two ounces of sulphur to be fused with one ounce of subcarbonate of potass, and the product to be digested with one pound of alcohol.—SPIELMANN requires the same product to be digested for several days with eight ounces of alcohol.

This tincture used formerly to be recommended against the itch.—Dose, sixty drops.

BOLUSES AGAINST SALIVATION.

Boli in Ptyalismo mercuriali. (BRERA)

R. Sulphuret of Potass, three grains.
Rob of Elder, a sufficient quantity.

Make six boluses.—Dose, one every three hours.

SUB-PROTO-CARBONATE OF POTASSIUM.

I. In the solid state.

A.

Impure.

Carbonas Potassæ venale seu impurum, Hypercarbonas Potassæ venale, Kali carbonicum seu carbonicum crudum seu subcarbonicum crudum seu carbonicum impurum, Alkali fixum vegetabili impurum, Potassa impura seu venalis, Subcarbonas Potassæ impurus seu kalicum impurum.—Potass of Commerce, Pearl-ash. — (LOND.; EDINB.; DUBL.; AMER.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENÈV.; HAMB.; HASS.; LIPP.; OLDEN.;

POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; COXE; PARIS; PIDERIT; SPIELMANN)

This is a mixture of subcarbonate of potass, of sulphates and hydrochlorates of potass, lime, and magnesia, of silica, of oxide of iron, and sometimes of oxide of manganese, in very variable proportions.

B.

Pure.

1. Purification of the Potass of Commerce.

Lixivia purificata, Kali præparatum seu carbonicum, Cineres clavellati depurati, Sal Cinerum Clavellatorum depuratus, Subcarbonas kalicum depuratum, Alkali Lignorum depuratum, Carbonas superpotassicus purus seu depuratus.—Purified Potass, Prepared Pearl-ash. — (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; FENN.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; BORUS.; SAX.; SUEC.; HERBIP.; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Potassæ Subcarbonas. (LOND.)

R. Impure Potass (Pearl-ash), in powder, three pounds.
Boiling Water, three pints and a half.

Dissolve, and filter; pour the solution into a clean iron pot, and evaporate with a gentle heat till the liquor has become thick; then take it from the fire, and stir assiduously with an iron spatula till the salt concretes into small grains.

A purer subcarbonate of potass may be prepared in a similar manner from tartar, previously burnt till of an ash colour.

Subcarbonas Potassæ. (EDINB.)

Put impure subcarbonate of potass into a crucible, and expose it to a red heat; then triturate it with an equal weight of water. Pour the solution, when the impurities have subsided, into a clean iron pot, and boil it to dryness, stirring the salt constantly towards the end to prevent it from adhering to the vessel.

Potassæ Carbonas e Lixivæ Cinere.
(DUBL.)

R. Pearl-ash, in coarse powder, } of each,
Cold Water, } one part.

Triturate together, and macerate in an open vessel for the space of a week, frequently stirring; then filter the solution, and evaporate it to dryness, assiduously stirring towards the end with an iron spatula. Having in this manner reduced to coarse powder, preserve it in well stopped bottles. If the pearl-ash is very impure, it should be previously roasted in a crucible till it becomes white.

ANT.; BELG.; FULD.; HANN.; HASS.; LIPP.; ROSS.; SAX.; PIDERIT; SPIELMANN; and VAN MONS give the same process as DUBL.—COXE; and SWEDIAUR the same as EDINB.—FENN.; and SWEDIAUR the same process, but with one part of pearl-ash and one and a half of water.

R. Pearl-ash, one part.
Boiling Water, three parts.

Boil for a quarter of an hour; decant the liquor while hot, and set it apart for twenty-four hours in an earthen pan that the extraneous salts may subside: then strain, evaporate to one half, set it apart in a cool place, and finally evaporate to dryness. (BAVAR.; BORUS.)

R. Pearl-ash, one part.

Calcine it strongly in an iron crucible; then infuse it for three days in

Water, three parts,
stirring it from time to time. Decant, wash the residue till the water ceases to render blue vegetable colours green, and evaporate the united liquors; then set aside for several days, decant, separate the extraneous salts, and evaporate the rest to dryness. (AMST.; BELG.; FORM. HÔP. FR.; HERBIP.)

R. Pearl-ash, } of each, one part.
Cold Water, }

Leave the mixture for twenty-four hours, occasionally stirring, then filter, and evaporate to dryness. Bring the residue to a white heat in a crucible, then dissolve in one part and a half of water; filter the solution and let it cool slowly; on the following day separate the crystals which are formed and dry them. (VAN MONS)

VAN MONS says that when the potass of commerce is purified by evaporating the solution to a pellicle in order to free it from crystallizable salts, so much of these pretended extraneous salts may be removed as

to leave at last no alkali. It was by following the process just described, that he ascertained the crystallizability of the subcarbonate of the potass.

Under the name of *Gallato-carbonate of Potass*, and as being a neutral carbonate or bicarbonate, a preparation has been given which is really nothing more than a subcarbonate; the following is the formula.

R. Subcarbonate of Potass, any quantity.

Dissolve it in the least possible quantity of water, filter the solution, and convert it into a dry paste with tan; expose this to a strong red heat in a crucible for half an hour: when cold, dissolve this mass in water, filter the liquor, and evaporate it to dryness. (VAN MONS)

2. *Formation of the salt by the combustion of Tartar.*

Alkali Tartari, Carbonas Potassæ purissimus, Kali subcarbonicum purissimum, Sal Tartari.—Salt of Tartar, (LOND.; DUBL.; EDINB.; AMST.; AUSTR.; BAVAR.; DAN.; HISP.; GALL.; FENN.; FERR.; GENEV.; HANN.; HASS.; OLDEN.; PALAT.; POL.; LUSIT.; BORUS.; SARD.; SAX.; SUEG.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; SPIELMANN; VAN MONS)

LOND. directs this to be prepared from tartar, previously burnt till of an ash colour, in the same manner as the subcarbonate from pearl-ash.

Subcarbonas Potassæ purissimus.
(EDINB.)

R. Impure Supertartrate of Potass, any quantity.

Wrap it up in moist bibulous paper or put it into a crucible, and having placed it among live coals, let it be burnt to a black mass; reduce this to powder, and expose it in an open crucible to a moderate fire, till it becomes white or at least ash coloured, taking care that it do not fuse. Then dissolve it in warm water; strain the solution through a linen cloth, and evaporate it in a clean iron vessel, stirring constantly towards the end with an iron spoon, lest any of it should adhere to the bottom of the vessel. A very white salt will remain, which is to be left a little longer on the fire till the bottom of the vessel becomes red-hot. Fi-

nally, when cold, enclose it in well-stopped glass vessels.

Potassæ Carbonas e Tartari Crystallis. (DUBL.)

R. Crystals of Tartar, any quantity.

Heat it to redness in a silver crucible lightly covered, till fumes cease to be emitted. Let the residue be reduced to a coarse powder, and then, in the same crucible, uncovered, roast it for two hours, stirring frequently. Then boil it in twice its weight of water, for a quarter of an hour, and when the impurities have subsided pour off the clear liquor. Repeat this part of the process thrice. Filter the mixed liquors and evaporate in a silver vessel; then, while the saline residue is drying, granulate it by fresh agitation, and expose to a dull red heat. Take it out of the vessel before it is quite cold, and preserve it in well stopped bottles.

BAVAR.; FENN.; and SAX. also prescribe the crystals of tartar. All the other pharmacopœias prescribe either the crude tartar, or the residuum of the distillation of tartar, and require the first to be calcined to whiteness in a crucible or in an iron boiler, or finally upon burning coals, enclosed in folds of paper, till it has ceased to exhale fumes.

HENRI observes that it is better to mix the tartar with one fourth of pulverized vegetable charcoal, and to put the mixture into small crucibles disposed in layers with charcoal.

3. *Formation of the salt by the deflagration of nitre with charcoal.*

Nitrum fixum. — Fixed Nitre. (GALL.; SARD.; SPIELMANN; SWEDIAUR; VAN MONS)

R. Nitrate of Potass, any quantity.

Having fused it in a crucible, throw into it powdered charcoal, by portions, till it ceases to raise any flame; then calcine the residuum for a quarter of an hour, dissolve it in water, filter the solution, and evaporate it to dryness.

HENRI recommends it to be calcined strongly, to prevent hyponitrite being left, which is frequently found in the product.

4. *Formation of the salt by the deflagration of nitre and of tartar.*

Nitre fixed by Tartar. (GALL.; FERR.; VAN MONS)

R. Nitrate of Potass,
Supertartrate of Potass, } of each,
an equal part

Put the mixture into an iron mortar, and inflame it with lighted charcoal. Dissolve the residuum in a sufficient quantity of water, strain the solution, and evaporate it to dryness. (FERR.; GALL.)

VAN MONS prescribes one part of tartar and two parts of nitre. These proportions are better than those of the preceding. If, instead of proceeding thus, the deflagration was effected at a high heat, and the subcarbonate brought to the state of pasty fusion, we should obtain this latter mixed with cyanuret of potassium, according to the important remark made by GUIBOURT.

5. *Formation of the salt by double decomposition.* (VAN MONS)

R. Carbonate of Barytes, twelve ounces.
Sulphate of Potass, fifteen ounces.
Pulverize the two salts, and pour upon them Boiling Water, twenty-four ounces.

Stir the mixture, filter the solution when cold, and evaporate it to dryness.

6. *Formation of the salt by the combustion of vegetables.*

Sal Tachenianus.—Tachenius, Lixivial Salt. (BRUNS.; HISP.; GALL.; FULD.; PALAT.; SARD.; WIRTEM.; HERBIP.; SPIELMANN)

R. Dried Herb of Wormwood, any quantity.

Put it into a covered iron pot placed over a moderate fire, and let it be slowly burnt and carbonized, producing much smoke but not throwing out flame; then remove the lid, and pulverize the mass by stirring it with a rod; boil the ashes with six times its weight of water, strain the boiling liquor and evaporate it to dryness.

In the same manner as the *Salt of Wormwood*, are prepared those of

Blessed Thistle. (HISP.; SARD.)

Lesser Centaury. (AUSTR.; GALL.; SARD.)

Juniper Wood. (SARD.)

Rest-harrow. (SARD.)

Tamarisk. (SARD.)

Bean-stalks. (SARD.)

In these cases the alkali obtained bears the name of the plant which has furnished

it; but it is in every instance precisely the same.

FULD. directs the charcoal produced by the combustion of one pound of wormwood, to be boiled with twelve ounces of purified potass dissolved in a sufficient quantity of water, and the decoction to be evaporated to dryness. This process is absurd.

The subcarbonate of potass is incompatible with the mineral acids, the sub-borate of soda, the muriate and acetate of ammonia, alum, the sulphate of magnesia, lime, lime-water, and all the metallic salts.

VEGETABLE SOAP.

Poudre gommeuse alcaline. (GALL.)

- R. Crystallized Subcarbonate of Potass, four parts.
Gum Arabic, thirty-two parts.
Triturate together for a long time.

POWDER AGAINST COLIC.

Pulvis anti-colicus Camerarii. (WIRTEM.)

- R. Salt of Wormwood, } of each,
——— Chamomile, } three drams.
Prepared Rock Crystal, half an ounce.
——— Hartshorn, six drams.
Ivory Shavings, two ounces.

Make a fine powder.

CAMERARIUS extolled it in disorders attributed to the presence of acids in the primæ viæ.—Dose, one or two scruples to adults, and half a scruple to children.

DIGESTIVE POWDER. (PALAT.)

- R. Salt of Wormwood, ten drams.
Supertartrate of Potass, six drams.
Sulphate of Potass, of each,
Sulphuretted Scammony, } two scruples five grains.
Oleo-saccharum of Lemon, one ounce.

Make a powder.

It is recommended in abdominal obstructions.—Dose, from a scruple to one dram.

AQUEOUS SOLUTION OF SUBCARBONATE OF POTASS.

Liquor Salis Tartari seu Kali carbonici seu Kali Subcarbonici, Liguamen Salis Tartari, Hypocarbonas Potassæ liquidum, Lixivium Tartari seu Alkali vegetabilis, Aqua Kali, Alcahest Glauberi, Carbonas Lixiviæ alkalinus solutus seu Potassæ alkalescens liquidus.

1. *Spontaneous Solution.* *Oleum Tartari per deliquium, Sal Tartari per se liquidus; Deliquesced Oil of Tartar.* (BRUNS;

HASS.; PALAT.; SARD.; WIRTEM.; HERBIP.; PIDERIT; SPIELMANN; SWEDIAUR)

It is obtained by putting the crystallized subcarbonate in a damp place, that it may deliquesce, and then filtering the liquor.

2. *Solution expressly prepared.*

The degree of saturation of the deliquesced solution varying to infinity, as may be easily conceived, it is better prepared with fixed proportions.

Liquor Potassæ Subcarbonatis. (LOND.)

- R. Subcarbonate of Potass, one pound.
Distilled Water, twelve fluid ounces.
Dissolve, and filter through paper.

Potassæ Carbonatis Aqua. (DUBL.)

- R. Carbonate of Potass, from crystals of tartar, one part.
Distilled Water, two parts.
Dissolve, and filter.

HISP. prescribes one part of salt and one half part of water.—AMER.; ANT.; and COXE give the same proportions as LOND.—VAN MONS—one of salt and one and a half of water.—AUSTR.; BAVAR.; DAN.; FENN.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; and SUEC.—the same proportions as DUBL.—LUSIT.; and ROSS.—one of salt and three of water.—WIRTEM.—one of salt and four or five of water.—FULD.—one of salt and a sufficient quantity of water.

This solution is employed externally as cosmetic; internally as absorbent, stimulant, incisive, alterative, and diuretic.

LITHONTRIPTIC SOLUTION. (AUGUSTIN; NIEMANN)

- R. Subcarbonate of Potass, nineteen drams twelve grains.
Water, fifteen ounces.
Dissolve.—Dose, one or two scruples thrice a day.

AUGUSTIN directs a solution of three drams of subcarbonate in two pounds of water, to be taken daily.

BONHOMME'S ANTI-RACHITIC LOTION. (PIERQUIN)

- R. Calcined Subcarbonate of Potass, from half an ounce to one ounce.
Water, one pound.
Dissolve.

ALKALINE PEDILUVIUM. (RATIER)

- R. Subcarbonate of Potass, eight ounces.
Hot Water, a sufficient quantity.

Mix.—It is irritant and revulsive.

MASCAGNI'S ALKALINE PTISAN. (SAINTE-MARIE)

- R. Subcarbonate of Potass, two drams.
Water, two pounds.

Dissolve.—To be taken by spoonfuls within twenty-four hours; it should be mixed when taken with an equal quantity of mucilaginous syrup. MASCAGNI recommended it in chronic pneumonia.

LITHONTRIPTIC POTION. (ELLIS)

- R. Subcarbonate of Potass, two drams.
Lime Water, two pounds.

Mix.—Dose, a glassful every two hours, with milk.

JUNIPER LEY. (SAUNDERS)

- R. Ashes of Juniper Wood, three ounces.
Water, two pounds.

Leave the mixture to digest for twelve hours in warm ashes; then filter.

In the same manner are prepared the leys of wormwood, of butcher's broom, and of tamarisk.

ANTI-DYSENTERIC DRINK. (SAINTE-MARIE)

- R. Ashes of Vine Twigs, four drams.
Water, two pounds.

Boil down to one third.

To be taken within twenty-four hours, mixed with an equal quantity of milk.

ALKALINE COLLUTORIUM. (FORM. HÔP. FR.)

- R. Subcarbonate of Potass, one part.
Despumated Honey, twenty parts.
Triturate together in a glass mortar.

ALKALINE SYRUP. (GENEV.)

- R. Subcarbonate of Potass, three drams.
Simple Syrup, six ounces.
Dissolve.

LOTION AGAINST CHILBLAINS. (NIEMANN)

- R. Subcarbonate of Potass, one ounce.
Rose Water, one pound.

Filter the solution.—BRUGNATELLI recommends rags to be wetted with it and applied to the chilblains.

ANTI-STRUMOUS POTION. (SAINTE-MARIE)

- R. Subcarbonate of Potass, one dram and a half.

- Ipecacuanha, } of each,
Protochloruret of Mer- } one dram
cury, } and a half,
Water, one pound and a half.

Boil down to one half. Dose, two spoonfuls in the course of the day, mixed with a pint of ordinary drink.

ABSORBENT POTION. (AUGUSTIN; ELLIS)

- R. Subcarbonate of Potass, } of each,
White Sugar, } one dram.
Compound Tincture of Lavender, }
two drams.
Tincture of Opium, } forty drops.
Mint Water, } four ounces.

Mix.—Dose, a spoonful every hour or two. (ELLIS)

- R. Subcarbonate of Magnesia, one dram.
Water, five ounces and a half.
Tincture of Cinnamon, three drams.
Solution of Subcarbonate of Potass, one dram.

Mix.—Dose, two or three spoonfuls after dinner. (AUGUSTIN)

ANTI-SPASMODIC POTION. (AUGUSTIN)

- R. Subcarbonate of Potass, one dram.
Water, four ounces.

Dissolve.—Dose, a spoonful every two hours, giving, in the intervals, ten drops of SYDENHAM'S laudanum.—Many German practitioners have found this remedy useful in convulsions, employed in concert with lavements prepared with two scruples of subcarbonate of potass and infusion of chamomile, and with epithems made with one ounce of caustic potass and two pounds of water.

ANTI-ASTHMATIC POTION.

Aqua kalino-ætherea. (AUGUSTIN)

- R. Subcarbonate of Potass, one dram.
Peppermint Water, four ounces.
HOFFMANN'S Liquor, two scruples.
Simple Syrup, half an ounce.

Mix.—Dose, a spoonful, in asthma.

SAPONACEOUS POTION.

Mixtura saponacea. (AUGUSTIN)

- R. Solution of Subcarbonate of Potass, half an ounce.
Oil of Almonds, one ounce and a half
Water, ten ounces.
Amygdaline Syrup, one ounce.

Mix.—To be taken by spoonfuls every half hour, in poisoning by arsenic.

EXPECTORANT POTION. (ELLIS)

- R. Subcarbonate of Potass, one scruple.
Cochineal, half a scruple.
Sugar, one dram.

Water, four ounces.
 Mix.—Dose, a tea-spoonful, every two or three hours; in hooping cough.

FEBRIFUGE DECOCTION.

Decoctum Carbonatis Potassæ Absinthii; Febrifugum salsum. (HISP.)

R. Salt of Wormwood, one dram.
 Honey, four ounces.
 Water, two pounds.

Boil, remove the scum, and strain.

ALKALINE MIXTURE. (SAINTE-MARIE)

R. Fresh Yolks of Eggs, four.
 Triturate them, adding gradually,
 Boiling Water, one pound.

Add,

Deliquesced Subcarbonate of Potass,
 from ten grains to a scruple.

Mix.—To be taken by spoonfuls every quarter of an hour or more frequently.—It has been recommended in poisoning by corrosive sublimate.

ANTI-LACTEAL POTION. (SAINTE-MARIE)

R. Subcarbonate of Potass, one scruple.
 Lily Water, six ounces.
 Orange-flower Water, one ounce.
 Syrup of Marshmallows, three ounces.

Mix.—To be taken by spoonfuls in the course of twenty-four hours, fomenting the breasts at the same time with a strong solution of soap in decoction of poppy heads.

ANTI-SCORBUTIC POTION. (BRERA)

R. Subcarbonate of Potass, two drams.
 Scurvy-grass Water, six ounces.
 Orange-flower Water, half an ounce.
 Syrup of Wild Mustard, one ounce.

Mix.

DIURETIC POTION. (SAINTE-MARIE; SWEDIAUR)

R. Salt of Butcher's Broom, twenty-four grains.
 Depurated Juice of Chervil, twenty-four spoonfuls.
 Juice of Wood-lice, one hundred.

Mix. (SAINTE-MARIE)

R. Subcarbonate of Potass, ten grains.
 Oxymel of Squill, one dram and a half.
 Compound Tincture of Lavender, half a dram.
 Cinnamon Water, six drams.

Mix. (SWEDIAUR)

ALKALINE LINIMENT. (HERBIP.; SPIELMANN)

R. Solution of Subcarbonate of Potass, two ounces.

Olive Oil, four ounces.
 Yolks of Eggs, two.
 Mix well.—PLENK recommends it in rhagades.

SEDATIVE POTION. (PIERQUIN)

R. Subcarbonate of Potass, two drams.
 Lime-flower Water, } of each,
 Rose Water, } four ounces.

Dissolve, and add

Syrup of Orange Peel, one ounce.

Mix.—To be taken at three or four doses in the course of the twenty-four hours, in an appropriate vehicle.

DIURETIC PTISAN. (RATIER)

R. Gum Arabic, five drams.
 Medicinal Soap, half a dram.
 Subcarbonate of Potass, one dram and a half.

Nitrate of Potass, twenty-four grains.
 Infusion of Juniper, two pounds.

Mix.—RECAMIER has found it serviceable in various cases of dropsy.

EMULSION AGAINST LUMBAGO. (CADET DE GASSICOURT)

R. Subcarbonate of Potass, half a dram.
 Water, ten ounces.

Dissolve, and add

Yolk of Egg, a sufficient quantity.
 Oil of Almonds, one ounce and a half.
 Syrup of Marshmallows, two ounces.

Mix.—To be taken by glassfuls, every half hour.

PURGATIVE AND FEBRIFUGE POTION. (CADET DE GASSICOURT)

R. Subcarbonate of Potass, one scruple.
 Castor Oil, two ounces.

Triturate in a glass mortar, and add

Syrup of Maidenhair, half an ounce.
 Peppermint Water, three ounces.

Mix.—To be taken at twice, half an hour elapsing between each dose.

RESOLVENT FOMENTATION. (CADET DE GASSICOURT)

R. Subcarbonate of Potass, } of each,
 Soap, } one ounce.
 Elder Water, two pounds.

Filter, and add to the solution

Muriate of Ammonia, two drams.

ANTI-PSORIC OINTMENT. (ELLIS)

R. Subcarbonate of Potass, half an ounce.
 Rose Water, } of each, one ounce.
 Cinnabar, }
 Oil of Bergamotte, half an ounce.

Sublimed Sulphur, } of each,
Lard, } nine ounces.

Triturate together.

ANTI-ARTHRITIC TINCTURE. (BRUNS.)

R. Earth-worms reduced to a pulp
with deliquesced Subcarbonate
of Potass, } one ounce.
Root of Round Birth- } of each,
wort, } half an ounce.
——— Juniper, }
Extract of Ground Pine, three drams.
Castor, } of each,
Saffron, } two drams.
Subcarbonate of Potass, }
Spirit of Scurvy Grass, twelve ounces.
Infuse, and express.

RED COLLYRIUM. (BORIES)

R. Subcarbonate of Potass, }
twenty-five grains.
Camphor, } ten grains.
Water of Great Celandine, two ounces.
Digest for twenty-four hours in a sand-
bath; then filter, and add to the liquor,
Tincture of Aloes, twenty-four drops.
Mix.—It is recommended against specks
of the cornea. A few drops to be instilled
into the eye.

STOMACHIC MIXTURE.

*Julapium stomachicum, Mixtura Carbonatis
Potassæ Absinthii stomachica.* (HISP.)

R. Salt of Wormwood, } four scruples.
Water of Lemon Peel, } of each,
——— Wormwood, } four ounces.
——— Mint, }
——— Cinnamon, } two ounces.
Tincture of Aloe Wood, } two drams.
Syrup of Lemon Peel, } one ounce.

Mix.—It is stimulant.—To be taken by
spoonfuls.

BESNARD'S ANTI-SYPHILITIC TINCTURE.

Tinctura anti-syphilitica Besnardi. (AU-
GUSTIN; BORIES; NIEMANN; VAN
MONS)

R. Opium, } two parts.
Vinous Cinnamon Water, twenty parts.
Macerate, and add
Gum Arabic, } two parts.
Subcarbonate of Ammonia, } one part.
——— Potass, } four parts.

Filter after one day's digestion in a mo-
derate heat. (VAN MONS)

R. Subcarbonate of Potass, } of each,
Simple Cinnamon Water, } one pound.
Opium, } two ounces.
Vinous Cinnamon Water, four ounces.

VOL. II.

Digest for three weeks in a water-bath,
frequently stirring, and add when strained,
Gum Arabic, } two ounces.
Subcarbonate of Ammonia, } one ounce.
Cinnamon Water, } six ounces.

Filter at the end of several days. (BO-
RIES; NIEMANN)

R. Subcarbonate of Potass, three ounces.
Opium, } of each,
Gum Arabic, } half an ounce.

Triturate together, and dissolve in
Chamomile Water, } six ounces.

Then add
Subcarbonate of Am- } of each,
monia, } two drams.
Proof Spirit, }

Digest in a water-bath, and filter when
cold. (VAN MONS)

AUGUSTIN prescribes half an ounce of
subcarbonate of potass, four ounces of cin-
namon water, two grains of opium, and one
dram of subcarbonate of ammonia.

Dose, from half a scruple to four scru-
ples in a proper vehicle.

TALBOT'S ANTI-SYPHILITIC TINCTURE.
(AUGUSTIN)

R. Subcarbonate of Potass, } two ounces.
Balm Water, } three ounces and a half.
Extract of Opium, } of each,
——— Celandine, } one scruple.

Digest for twenty-four hours, and add
Solution of Ammonia, } two drams.

Mix.—This tincture, which has been
lately extolled by EICHHEIMER, very much
resembles the preceding.

INCISIVE POTION. (BORIES)

R. Subcarbonate of Potass, } one scruple.
Cochineal, } half a dram.
Hyssop Water, } four ounces.
Syrup of Poppies, } of each,
——— Tolu, } half an ounce.
——— Ipecacuanha, } of each,
Orange-flower Water, } one ounce.

Triturate together the cochineal and sub-
carbonate, then add the hyssop water hot;
when cold, filter, and add the other sub-
stances.—Dose, a tea-spoonful or dessert
spoonful, according to the age of the chil-
dren; in hooping cough.

REDFEARN'S ANTI-DIABETIC REMEDY.
(AUGUSTIN)

R. Red Cinchona, } one ounce.
Calcined Alum, } three drams.
Subcarbonate of Potass, } two drams.
Petroleum, } a sufficient quantity
to make an electuary.—Dose, the volume of
a nutmeg, thrice a day. At the same time

a solution of three drams of sulphuret of potass in sixteen ounces of lime water should be taken in the course of the day, together with four of the following pills:

- R. Sulphate of Soda, one dram.
Medicinal Soap, four scruples.
Mucilage of Gum Arabic,
a sufficient quantity.

Make thirty pills.

ANTI-PSORIC OINTMENT.

Unguentum Burdenii. (BRERA)

- R. Subcarbonate of Potass, one ounce.
Sublimed Sulphur, two ounces.
Lard, four ounces.

Mix.—To be rubbed in, at four times, in the course of the day.

PROTO-CARBONATE OF POTASSIUM.

Carbonas Protoxidi Potassii.—Carbonate of Potass, Bicarbonate of Potass. (LOND.; EDINB.; DUBL.; ANT.; BATAV.; BAVAR.; BELG.; DAN.; HISP.; GALL.; FENN.; FERR.; HANN.; POL.; SUEC.; COXE; SWEDIAUR; VAN MONS)

Potassæ Carbonas. (LOND.)

- R. Solution of Subcarbonate of Potass, one gallon.

Transmit carbonic acid through it by means of a proper apparatus, till saturated, and filter it. Evaporate the solution till crystals form, being careful not to raise the heat above 120 degrees. Separate the crystals from the liquor, and dry them in bibulous paper.

The carbonic acid may be easily obtained from white marble and dilute sulphuric acid.

Carbonas Potassæ. (EDINB.)

- R. Pure Subcarbonate of Potass, two parts.
Water, three parts.

Dissolve the subcarbonate in the water, and pass a stream of carbonic acid gas into the solution by means of a proper apparatus. When the solution has ceased to absorb the acid, filter it, and evaporate with a heat not exceeding 180 degrees, that crystals may form. The carbonic acid may be easily obtained by pouring

dilute sulphuric acid on powdered carbonate of lime.

Potassæ Bicarbonas. (DUBL.)

- R. Carbonate of Potass, one part.
Distilled Water, two parts.

Dissolve. Expose this solution in a proper apparatus, to a stream of carbonic acid gas, evolved from white marble by diluted muriatic acid, till the liquor becomes turbid; then strain, and expose it again to the stream of carbonic acid till the alkali is saturated: finally, put the solution in a cool place that it may crystallize; dry these without heat, and preserve them in well-stopped bottles.

- R. Subcarbonate of Potass, two parts.
Water, three parts.

Dissolve, filter, and pass through the solution a current of carbonic acid gas. At the end of twenty-four hours, evaporate and set aside that crystals may form. (ANT.; BATAV.; BAVAR.; BELG.; GALL.; FERR.; FENN.; HANN.; POL.; SUEC.; COXE)

- R. Subcarbonate of Potass, one pound.
Ammonia, three ounces.
Water, one pound.

Add the ammoniacal salt to the subcarbonate of potass dissolved in the water, and expose the mixture for three hours to a temperature of 380 degrees, F., in a sand-bath, till all the ammonia is expelled; then set the liquor apart that it may crystallize. (HENRI)

This process is due to BERTHOLLET.

- R. Alcoholated Potass, one part.
Distilled Water, three parts.

Pass into the liquor a current of carbonic acid gas, till it is perfectly saturated; then evaporate slowly, and set apart to crystallize. (SWEDIAUR)

HENRI recommends, to obtain fine crystals, the solution, well charged with gas, and in which crystals have begun to be formed, to be evaporated to two-thirds, and then left to repose for two days.

GALL. requires the tube to be plunged into the solution.—ANT.; BATAV.; BAVAR.; BELG.; and VAN MONS—the surface of it to be scarcely touched by the tube.

SOLUTION OF SUPER-CARBONATE OF POTASS.

Aqua Carbonatis Potassæ acidula seu Su-

per-carbonatis Potassæ seu aërata alkalina seu Super-carbonatis kalici, Aqua lithontriptica, Potio alkalina.—Alkaline Potion, Alkaline mephitic Water. (EDINB.; AMER.; DAN.; HISP.; FENN.; POL.; BORUS.; BRUGNATELLI; COXE; HENRI; SWEDIAUR; VAN MONS)

Aqua Supercarbonatis Potassæ.
(EDINB.)

R. Water, ten pounds.
Pure Subcarbonate of Potass, one ounce.

Dissolve, and expose the solution to a stream of carbonic acid gas obtained from

Carbonate of Lime,	} of each,	
in powder,		three
Sulphuric Acid,		ounces.
Water,		three pounds,

gradually and cautiously mixed together.

The chemical apparatus of DR. NOOTH is well adapted for this preparation; but if a larger quantity of the solution is required, an apparatus should be used which will allow of a sufficiently great pressure. The solution should be kept in well stopped vessels.

AMER.; FENN.; BRUGNATELLI; COXE; and SWEDIAUR give the same formula.

R. Subcarbonate of Potass, two parts.
Water, three parts.

Pour the solution into a bottle filled with carbonic acid, and agitate for a long time. (POL.; BORUS.; HENRI)

R. Subcarbonate of Potass, two drams.
Water charged with Carbonic Acid, one pound.

Dissolve, and preserve in a well-stopped bottle. (DAN.)

SUEC. prescribes three ounces of the salt, and fifty-six pounds of saturated gaseous water.

R. Subcarbonate of Potass, one part.
Water, one hundred and twenty parts.

Charge the solution with the desired number of atmospheres of carbonic acid gas. (VAN MONS)

R. Neutral Carbonate of Potass, three ounces.
Water, twelve ounces.

Dissolve, and filter. (VAN MONS)

R. Neutral Carbonate of Potass, one part.
Water charged with Carbonic Acid, one hundred and twenty parts.

Dissolvè. (VAN MONS)

HISP. prescribes three drams of neutral carbonate, and two pounds of gaseous water.

It is excitant, and is considered digestive and lithontriptic. It has also been extolled in what are called lymphatic disorders.—Dose, twelve ounces in the course of the day, to be augmented gradually.—A little brandy may sometimes be added to it.

LITHONTRIPTIC POTION. (ELLIS)

R. Neutral Carbonate of Potass, from ten to fifteen grains.
Seltz Water, six ounces.

Dissolve.—To be taken three or four times a day.

PROTO-CARBONATE OF POTASSIUM AND OF AMMONIA.

Carbonas Protoxidi Potassii et Ammoniae seu Potassæ et Ammoniae. (SWEDIAUR)

R. Crystallized Subcarbonate of Potass, four parts.
Ammonia, one part.

Triturate together, and pour upon the mixture

Cold Water, a sufficient quantity to merely dissolve it. Filter the liquor, and pass through it a current of carbonic acid till saturated; apply a little heat, that the salt which was precipitated may be redissolved, and crystallize.

It is recommended in diabetes, dyspepsia, and in cases of urinary calculi.—Dose, half a dram, in half an ounce of juniper or mint water, four times a day.

HYPOPHOSPHATE OF POTASS. (VAN MONS)

R. Tincture of Salt of Tartar, any quantity.
Granulated Phosphorus, a sufficient quantity

to saturate it without heat. Decant, and preserve.

Granulated phosphorus may also be put into a bottle with concentrated solution of potass, and rectified alcohol added; the mixture left in re-action for several days, and then decanted.

PROTO-SULPHATE OF POTASSIUM.

Alkali vegetabile vitriolatum, Arcanum duplicatum Mynsichti, Kali sulphuricum seu vitriolatum, Lixivia vitriolata, Nitrum

vitriolatum Schræderi, Panacea duplicata, Sal de duobus seu polychrestum Glaseri seu Lemery seu Sapientiæ seu polychrestus Parisiensis seu Tartari sulphuratum, Specificum Paracelsi, Spiritus Vitrioli coagulatus Mynsichti, Sulphas Potassæ, Tartarus vitriolatus.—Sulphate of Potass, Vitriolated Tartar, Vitriol of Potass.

I. Sulphate furnished by commerce.

EDINB.; BAVAR.; BELG.; HISP.; FORM.
HÔP. FR.; HASS.; LIPP.; SAX.; COXE;
PARIS.

II. Purification of the Sulphate of commerce.

BATAV.; BAVAR.; BELG.; HASS.; LIPP.;
SAX.

R. Sulphate of Potass, one part.
Water, eight parts.

Dissolve; instil into the liquor an aqueous solution of the subcarbonate of potass till it is no longer rendered turbid; then filter, evaporate till a slight pellicle is formed, and crystallize. (BATAV.; BELG.; HASS.; SAX.)

LIPP. doubles the quantity of water.—BAVAR. directs one part of the salt to be dissolved in five parts of boiling water, and the solution to be set apart in a cool place to crystallize.

III. Prepared according to Formula.

LOND.; DUBL.; EDINB.; AMER.; AMST.;
ANT.; AUSTR.; BATAV.; BELG.;
BRUNS.; DAN.; HISP.; GALL.; FENN.;
FULD.; GENEV.; HANN.; HASS.; LIPP.;
OLDEN.; LUSIT.; PALAT.; POL.; BORUS.;
ROSS.; SARD.; SUEC.; WIRTEM.;
HERBIP.; BRUGNATELLI; COXE; SPIELMANN; VAN MONS.

Potassæ Sulphas. (LOND.)

R. Residuary Salt after the distillation of Nitric Acid,
two pounds.
Boiling Water, two gallons.

Dissolve, and add as much subcarbonate of potass as may suffice to saturate the excess of acid; then boil till a pellicle forms on the surface of the liquor, and set it apart to crystallize after having filtered it. Pour off the water, and dry the crystals on bibulous paper.

Sulphas Potassæ.

Dissolve the salt which remains

after the distillation of nitric acid in a sufficient quantity of warm water, and add as much common carbonate of potass as will saturate the superfluous acid. Evaporate the liquor, and crystallize. (DUBL.)

Dissolve the acidulous salt remaining after the distillation of nitrous acid, in hot water, and add as much powdered carbonate of lime as will be sufficient to saturate the superfluous acid; then set apart till the fæces have subsided. Finally, pour off the liquor, filter it, and evaporate till crystals form. (EDINB.)

BATAV.; DAN.; FERR.; FULD.; HANN.; OLDEN.; POL.; BORUS.; SUEC.; BRUGNATELLI; COXE; SPIELMANN and VAN MONS give the same process as DUBL.—AUSTR.; and GALL.—the same as EDINB.—SARD. recommends the solution to be evaporated to two thirds, filtered, and evaporated to dryness; the residuum to be calcined for several hours with a violent heat, then dissolved in water, filtered, evaporated to dryness, and calcined again; it is finally to be dissolved in water, the filtered solution evaporated to one half, and set apart to crystallize in a cool place.

FENN. also directs the excess of acid to be removed by the application of a strong heat. CHEVALLIER and IDT remark that the salt thus obtained has seemed to them to contain sometimes a certain quantity of nitrate of potass.

R. Subcarbonate of Potass, one part.
Water, twelve parts.

Dissolve, and add gradually to the liquor,
Sulphuric Acid (20 degrees),
a sufficient quantity

to saturate the alkali. Strain, evaporate to a pellicle, and let it cool slowly. (AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HASS.; LIPP.; LUSIT.; PALAT.; ROSS.; SARD.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT)

R. Sulphate of Iron, two parts.
Boiling Water, eight parts.

Dissolve. On the other hand,

R. Subcarbonate of Potass,
Water, a sufficient quantity,
five parts.

Dissolve.

Mix these to perfect saturation; then filter, and wash the precipitate well; mix the water used in washing it with the filtered liquor, evaporate to a pellicle, and crystal-

lize. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

R. Nitrate of Potass, } of each,
Sulphate of Iron calcined } an equal
to whiteness, } part.

Fill an open crucible to one half with the mixture, and calcine till smoke is no longer disengaged. Dissolve the residuum in boiling water, evaporate to dryness, and calcine the product for several hours; dissolve it again in water, and finally, filter, evaporate, and crystallize. (SARD.)

Sulphas Potassæ cum Sulphure.
(EDINB.)

R. Nitrate of Potass, } of each,
in powder, } an equal
Sublimed Sulphur, } weight.

Mix well together, and throw the mixture by small portions into a red hot crucible. When the deflagration is finished, let the salt cool, and preserve it in a well-stopped glass vessel.

R. Nitrate of Potass, two parts.
Sublimed Sulphur, one part.

Project the mixture by portions into a red hot iron vessel. When flame ceases to rise keep the mass over the fire for two hours, bringing the whole vessel to a red heat. Pulverize the residuum when cold, dissolve it in boiling water, and strain; evaporate the solution to a pellicle, and set it aside to crystallize in a cool place. (SARD.)

AMST.; BATAV.; BRUNS.; WIRTEM.; HERBIP.; COKE; and SPIELMANN prescribe an equal part of each of the two ingredients.

LIPP. substitutes the subcarbonate of potass for the nitrate, and prescribes an equal part each, of that and of sulphur.

The product appears to be a mixture of sulphate and of sulphite of potass. It is very certain, as DUNCAN has remarked, that it does not contain free sulphur, and therefore the name which is imposed on it by EDINB. does not accord with its composition.

R. Urine of a healthy man, any quantity.

Expose it to cold so that half of it may be converted to ice; filter the rest, pour it into plates, wash the crystals which form spontaneously, and dry them. (HERBIP.)

This salt was formerly called *Native Salt*, *Fusible Salt*, or *Microcosmic Salt of Urine*. It is a mixture of the phosphates of soda and of ammonia, the hydrochlorates of soda and of ammonia, and the sulphates

of potass and of soda; the first of these salts entering in the proportion of 2,94, the second, 1,65, the third, 4,45, the fourth, 1,50, the fifth, 3,71, and the sixth, 3,16, in 100 parts, according to BERZELIUS. To render it colourless and perfectly pure, it should be first treated with alcohol, then re-dissolved in water, and again crystallized.

It is excitant, alterative, and purgative. —Dose from a scruple to a dram as an alterative, and from one dram to six, as a purgative.

It is incompatible with the hydrochloric and nitric acids, the chloruret of calcium, mercurial salts, and nitrate of silver.

DIGESTIVE POWDER.

Pulvis digestivus. (LIPP.; WIRTEM.; SWEDIAUR)

R. Sulphate of Potass, two parts.
Prepared Oyster Shells, one part.

Mix. (WIRTEM.)

R. Sulphate of Potass, two parts.
Root of Arum, one part.

Mix. (LIPP.)

R. Sulphate of Potass, eight parts.
Citrate of Lime, four parts.
Root of Arum, one part.

Mix. (SWEDIAUR)

The usual dose is half a dram.

COMPOUND DIGESTIVE POWDER.

Pulvis digestivus cum Rheo. (MIL. COPEN.)

R. Sulphate, four parts.
Rhubarb, one part.

Mix.—Dose, half a dram.—The sulphate of soda or of magnesium may be substituted for that of potass.

ANTI-PHLOGISTIC POWDER. (HUFELAND)

R. Sulphate of Potass, } of each,
Nitrate of Potass, } an equal part.
Sulphate of Magnesia, }

Mix.

ALTERATIVE AND RESOLVENT POWDER.
(BORIES)

R. Sulphate of Potass, } of each,
Tartarized Iron, } two drams.
Salt of Butcher's-broom, } one dram.
Nitrate of Potass, half a dram.
Wood-lice, thirty grains.

Mix carefully.

GRIMAUD recommended it in obstructions of the abdominal viscera, cedema, infiltration, suppression, or diminution of the urine.—Dose, fifteen or twenty grains, at bed time.

LOZENGES AGAINST THIRST. (PIERQUIN)

R. Sulphate of Potass, three drams.
 Sugar, one pound.
 Oil of Lemon, eight drops.
 Gum Tragacanth, a sufficient quantity
 to make lozenges.

There is evidently a mistake in this formula which is presented as offering dry lemonade in the form of lozenges. It ought from its title to contain tartaric or citric acid. However, it may be kept, such as it is. They are excitant and alternative.—Dose, half an ounce or more, in water.

SPAGYRIC CONSERVE. (WIRTEM.; SPIELMANN)

R. Salt of Wormwood, two ounces.
 ——— Balm, } of each, one ounce.
 ——— Nettle, }
 Sulphate of Potass,
 two ounces and a half.

Make a powder, and add to it,

Sulphurous Acid, two drams.

Add, when the effervescence has ceased,

Conserve of Red Roses, } of each,
 ——— Clove Pink, } one pound.

Mix. (WIRTEM.)

R. Salt of Wormwood, three ounces six drams.
 Powdered Cinnamon, two drams.
 ——— Mace, } of each, one dram.
 ——— Cloves, }
 ——— Ginger, }
 Sulphate of Potass, one ounce six drams.
 White Sugar, eight ounces.
 Conserve of Red Roses, sixteen ounces.
 ——— Clove Pink, eight ounces.
 Sulphuric Acid, two drams.
 Oil of Anise, } of each, one dram.
 ——— Lemon, }
 ——— Fennel, half a dram.
 ——— Cinnamon, one scruple.
 ——— Mint, four drops.

Mix carefully. (SPIELMANN)

This inconsistent preparation was formerly considered resolvent in abdominal obstructions, jaundice, obstinate quartan fevers, amenorrhœa, &c.—Dose, from a dram to half an ounce.

SALT OF GLASS.

Fel Vitri, Recrementa Vitri, Sal Vitri, Anatron. (BRUNS.; WIRTEM.; SPIELMANN)

In white, light, spongy masses.

It is a mixture of chloruret of potassium or of sodium and of sulphate of potass or of soda, which is separated during the fusion of glass; and floats on the surface of the vitreous mass.

It was formerly employed as incisive and laxative.

AQUEOUS SOLUTION OF SULPHATE OF POTASS. (BAVAR.)

R. Sulphate of Potass, one part.
 Distilled Water, nineteen parts.

Dissolve.

It should be remembered when this salt is prescribed in solution, that, according to GAY LUSSAC, 100 parts of water dissolve 10 57.100 of it at 12 degrees 72, and 26 33.100 at 101 degrees, 50. C.

LAXATIVE POTION. (SAUNDERS)

R. Sulphate of Potass, six drams.
 Balm Water, four ounces.

Dissolve and add,

Syrup of Raspberries, one ounce.

Mix.

REFRESHING EMULSION. (SAUNDERS)

R. Sulphate of Potass, one dram.

Dissolve it in one pound and a half of an emulsion made with

Seeds of Mallow, three ounces.
 Barley Water, a sufficient quantity.

Then add,

Syrup of Marshmallows,
 one ounce and a half.

Mix.—To be taken by glassfuls.

ACID MIXTURE. (SAUNDERS)

R. Sulphate of Potass, half an ounce.
 Nitrate of Potass, one dram.
 Spring Water, ten ounces.

Dissolve and add,

Oxymel, one ounce.
 Vinegar, one ounce and a half.

Mix.

VINOUS MIXTURE. (SAUNDERS)

R. Sulphate of Potass, half a dram.
 Decoction of Barley, two pounds.

Dissolve, and add,

Rob of Goose-berries, } of each,
 Wine, } one ounce and
 Vinegar, } a half.
 Syrup of Raspberries, one ounce.
 two ounces.

Mix.

RESOLVENT POTION. (BRERA)

R. Sulphate of Potass, three drams.
 Infusion of Rhubarb, six ounces.

Dissolve.—To be taken gradually, in lacteal engorgements and metastasis.

SUPER-PROTO-SULPHATE OF POTASSIUM.

Potassæ Supersulphas seu Bisulphas.
(LOND.; DUBL.; FERR.; SUEC.; COXE;
PALAT.; VAN MONS)

Potassæ Supersulphas. (LOND.)

- R. Residuary Salt after the distillation of nitric acid,
two pounds.
Boiling Water, four pints.

Dissolve, and filter. Then boil the solution till one half is dissipated and set aside to crystallize. Pour off the water, and dry the crystals on bibulous paper.

Potassæ Bisulphas. (DUBL.)

- R. Sulphuric Acid of commerce,
two parts.
Common Carbonate of Potass,
a sufficient quantity.
Water, six parts.

Mix one part of the acid with the water, and saturate it with the carbonate; then add the other part of the acid, and evaporate the liquor till crystals form on cooling.

COXE; and VAN MONS give the same formula as LOND.

- R. Sulphuric Acid, } of each, seven parts.
Water, }
Subcarbonate of Potass, four parts.
Dissolve, mix, and evaporate. (VAN MONS)

- R. Sulphuric Acid, ten parts.
Subcarbonate of Potass,
six parts and a half.
Water, twenty parts.

Mix the acid with the water, and add the salt to the mixture while hot, as rapidly as the effervescence will permit; then filter, evaporate slightly, and crystallize. (VAN MONS)

- R. Powdered Sulphate of Potass,
eight parts and a half.
Sulphuric Acid, five parts.

Dissolve, evaporate to dryness, and heat the residuum to redness. Dissolve it in one part and a half of water, filter, and crystallize. (VAN MONS)

FERR. prescribes two ounces of sulphate and three of acid.

- R. Sulphate of Potass, eight parts.
Tartaric Acid, seven parts and a half.
Triturate together, and wash it with a

little cold water; filter the liquor, concentrate it, and set it apart to crystallize. (VAN MONS)

- R. Sulphate of Potass, two parts.
Sulphuric Acid, one part.
Water, three parts.

Mix, evaporate to dryness, and pulverize. (SUEC.)

It is purgative, and considered anti-pu-
rescent; it is said to have been given with
advantage in gastric and adynamic diseases,
and is sometimes combined with decoction
of cinchona.

SULPHURICO-SALINE POTION. (VAN MONS)

- R. Super-sulphate of Potass, six drams.
Water, six pounds.
Proof Spirit, } of each,
Syrup of Violets, } one ounce and a half.
Dissolve, and mix.

It is excitant, slightly laxative, and con-
sidered antiseptic; and is prescribed in bi-
lious and adynamic fevers, hæmatemesis,
and menorrhagia.—Dose, a glassful, more
or less frequently.

PROTO-SULPHATE OF POTASSIUM AND OF AMMONIA.

Sulphas Potassæ et Ammoniæ. (VAN MONS)

- R. Supersulphate of Potass, one part.
Hot Water, two parts.

Pour into the solution
Solution of Ammonia,
a sufficient quantity,
to saturate the excess of acid, and crys-
tallize.

PROTO-SULPHATE OF POTASSIUM AND OF MAGNESIUM.

Sulphas Potassæ et Magnesiæ. (VAN MONS)

- R. Sulphate of Potass,
three parts and a half.
Magnesia, ten parts.

Mix, dissolve in a sufficient quantity of
hot water, filter, and crystallize.

- R. Sulphate of Magnesia, ten parts.
Calcined Subcarbonate of Potass,
three parts and one fourth.

Triturate, adding gradually,
Warm Water, a sufficient quantity,
to effect the solution. Boil for a few mo-
ments, filter, and crystallize.

- R. Supersulphate of Potass, one part.
Water, eighteen parts.
Saturate the solution, applying heat, with
Magnesia, a sufficient quantity.
Filter and crystallize.

PROTO-HYPOSULPHATE OF POTASSIUM.
(VAN MONS)

- R. Sulphate of Potass, }
seven parts and a half.
Fused Caustic Potass, }
five parts and a half.
Powdered Sulphur, }
one part and a half.

Dissolve the salt in the smallest quantity of water required, add the potass, then the sulphur, and apply heat for half an hour in a covered vessel; then filter and crystallize.

- R. Sulphuret of Potass, } of each,
Sulphate of Potass, } an equal part.

Triturate together for some time; then dissolve the mixture in the least possible quantity of water, filter, and crystallize.

- R. Hydrosulphate of Potass, any quantity.

Dissolve it in water, and pass a current of sulphurous acid gas into the solution till no more of it is absorbed; then heat slightly, filter, and crystallize.

A current of hydrosulphuric acid gas may also be passed into a solution of sulphate of potass; the liquor then heated, filtered, and crystallized.

This salt may be substituted for the sulphuretted sulphite of soda.

PROTO-SULPHITE OF POTASSIUM. (VAN MONS)

- R. Subcarbonate of Potass, } of each,
Water, } one part.

Dissolve, and filter. Saturate with sulphurous acid gas; then heat in a warm water-bath, and set apart the liquor that crystals may form.

PROTO-NITRATE OF POTASSIUM.

Kali nitricum, Nitras kalicum seu Lixivia seu Potassæ, Nitrum, Nitras Protoxidi Potassii, Nitras Potassæ.—Nitre, Salt-petre.

I. Nitre of commerce.

Nitrum crudum seu venale.—Crude Nitre.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BRUGNATELLI; COXE; PARIS; PIDERIT; SPIELMANN; SAINTE-MARIE; VAN MONS.

II. Purified Nitre of commerce.

Nitrum depuratum seu purificatum seu purum, Nitras Potassæ depuratus seu purus.

AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FENN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; PARIS; PIDERIT; SPIELMANN; SAINTE-MARIE; VAN MONS.

- R. Nitre of Commerce, } two parts.
Boiling Water, } one part.

Dissolve, strain, and crystallize. (AMST.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; LIPP.; OLDEN.; LUSIT.; PALAT.; ROSS.; WIRTEM.; HERBIP.; BRUGNATELLI; PARIS; PIDERIT; SPIELMANN)

FENN.; and SUEC. direct the salt to be fused at a slow heat in a crucible, the mass to be pulverized when cold, and dissolved in four times its weight of water, the solution to be strained, evaporated to a pellicle, and rapidly cooled, constantly stirring; the crystals which form to be washed with a little water, and dried.—ANT.; BATAV.; BELG.; POL.; and BORUS.—the salt to be washed in a sufficient quantity of boiling water, a solution of subcarbonate of potass instilled into the liquor till it ceases to trouble it; then to be set aside, filtered, evaporated, and crystallized.—HASS.; SAX.; and SWEDIAUR—the same process to be followed, but, besides, the filtered liquor to be boiled with charcoal.—HANN.—solution of nitrate of silver to be instilled into it till it ceases to render it turbid.

VAN MONS requires the nitre to be boiled with one fourth of its weight of boiling water, the solution to be decanted, one fourth part of cold water added to it, and the mixture shaken till perfectly cold; the liquor to be then decanted, a little cold water poured upon the crystallized salt, which is then to be put into funnels, and washed without heat, first, at several times, with the same saturated solution of nitrate, depurated and afterwards with a solution which has not yet been used: the crystals having been left to drip, are then to be dissolved in five parts of boiling water, the solution filtered while hot, and set apart to crystallize.

III. Regenerated Nitre.

Nitrum regeneratum.

BORUS.

- R. Nitric Acid, } any quantity.
Solution of Subcarbonate of Potass, } a sufficient quantity,
or, till it ceases to excite effervescence. Filter the liquor, evaporate it to a pellicle, and crystallize.

IV. Nitre impregnated with foreign substances.

Nitre which has been crystallized after

having been dissolved in infusion of poppy or of red roses. It is called when prepared with the former *Nitrum papaveratum* (PALAT.), and when with the latter *Nitrum rosatum* (WIRTEM.).

V. Nitre mixed with Sulphate of Potass.

Sal seu Lapis Prunellæ, Nitras Potassæ cum Sulphure fusus, Crystallum minerale, Crystallus mineralis, Nitrum tabulatum.—*Sal Prunella.*

AMST.; ANT.; BRUNS.; HISP.; GALL.; PALAT.; SARD.; WIRTEM.; SPIELMANN; VAN MONS.

R. Nitre, two pounds and a half.

Fuse it in a crucible and add to it gradually

Sublimed Sulphur, half an ounce.

After the deflagration and the vessel has become a little cooled, pour the liquid mass into a pan, and move it about in such a manner that the salt may be spread over the sides in a thin and uniform layer: take this out when cold, and break in pieces. (HISP.)

ANT. prescribes two pounds of nitre and half an ounce of sulphur.—SARD.—one pound of nitre and two drams of sulphur.—AMST.—one part of sulphur and twenty parts of nitre.—GALL.—one of sulphur and one hundred and twenty-eight of nitre.—VAN MONS; BRUNS.; PALAT.; and WIRTEM.—one of sulphur and twenty-four of nitre. The operatory process is the same in every instance.

It is excitant, acting as purgative in large doses; it is however classed among the refrigerants or antiphlogistics, certainly only on account of the revulsion which it determines by irritating the primæ viæ; it is also considered to have a specific diuretic property. It is prescribed in dropsies, hæmorrhages, tetters, gonorrhœa, &c.—The dose is usually from ten grains to half a dram, but it has been carried much higher by some modern experimentalists, by which we have learnt that nitre is not so poisonous as FODERE and ORFILA would have us believe.

It is incompatable with sulphuric acid, the sulphates of soda and of magnesia, alum, and the metallic sulphates.

A

Preparations in which the action of nitre is not sensibly modified by that of other substances.

PEARLED NITRE.

Nitrum perlatum. (WIRTEM.)

R. Nitre, four ounces.
Prepared Mother of Pearl,
half an ounce.

VOL. II.

Mix.—This powder used formerly to be prescribed, in the dose of from five grains to a scruple, in fevers of a bad character; its effects were chiefly attributed to the inert substance accompanying the nitre.

AUGUSTIN gives an analogous preparation under the name of *Pulvis nitrosus*, composed of nitre and of prepared oyster shells, at an equal part each.

ANTI-CATARRHAL POWDER.

Pulvis Iridis cum Nitro. (AUGUSTIN)

R. Root of Florentine Iris, one ounce.
Nitre, one dram.
Spermaceti, two drams.
Sugar, one ounce.

Mix.—Dose, a tea spoonful, in catarrhal affections.

APERITIVE POWDER.

Pulvis aperitivus. (GENEV.; VAN MONS)

R. Nitre, one ounce.
Gum Arabic, one ounce and a half.
Root of Liquorice, } of each,
Marshmallows, } three ounces.

Mix.—It is diuretic and emollient, and is recommended in the acute stage of urethritis.—Dose, one dram and a half or two drams thrice a day, in water. (VAN MONS)

R. Nitre, three drams.
Magnesia, half an ounce.

Divide into twelve parcels. (GENEV.)

APERITIVE PILLS.

Pilulæ aperitivæ. (VAN MONS)

R. Nitre, six drams.
Gum Arabic, three drams.
Root of Liquorice, } of each,
Marshmallows, } three ounces.
Simple Syrup, a sufficient quantity.

Make pills of five grains each.

R. Nitre, one part and a half.
Prepared Oyster Shells, one part.
Gum Arabic, two parts.
Root of Liquorice, } of each,
Marshmallows, } four parts.
Simple Syrup, a sufficient quantity

to form a pilular mass. (VAN MONS)

They are diuretic, emollient, and anodyne, and are recommended in urethritis and dysury.—Dose, five or six, thrice a day.

It is rather singular that VAN MONS prescribes the addition of the oyster shells in gonorrhœa only when it is provoked by the use of beer.

TROCHISCS OF NITRE.

Trochisci de Nitro seu Nitratis Potassæ.
(EDINB.; HERBIP.; COXE; SPIELMANN)

Trochisci Nitratis Potassæ (EDINB.;
HERBIP.)

R. Nitre, one part.
Sugar, three parts.
Mucilage of Tragacanth,
a sufficient quantity
to form a mass, to be divided into
trochiscs.

SPIELMANN prescribes one part of nitre,
six parts of sugar, one part of lemon juice,
and a sufficient quantity of mucilage.

TEMPERANT ELECTUARY. (GENEV.;
HAMB.)

R. Nitre, two drams.
Conserve of Roses, } of each,
Oxymel, } two ounces.

Mix. (HAMB.)

GENEV. prescribes half an ounce of nitre
and four ounces of conserve of roses, with-
out oxymel.

NITRATED POTION. (PIERQUIN)

R. Nitre, half a dram.
Common Drink, four pounds.

Dissolve.—To be taken by glassfuls.

DIURETIC POTION. (SAINTE-MARIE)

R. Nitre, half a dram.
Juice of Chervil,
twenty-four spoonfuls.

Dissolve.—To be taken by spoonfuls.

FEBRIFUGE POTION. (NIEMANN)

R. Nitre, one ounce.
Cranberry Juice, one pound.

Dissolve.

It is employed in the north of Europe
against intermittent fevers.—Dose, from an
ounce to half a pound.—The juice of bil-
berry or of raspberry may be used instead
of that of the cranberry.

NITROUS MIXTURE.

Mixtura nitrosa. (CAST. BORUS.)

R. Nitre, two drams.
Water, eight ounces.
Simple Syrup, one ounce.

Mix.

COMMON GARGLE. (FULD.; LIPP.)

R. Nitre, one dram.

Water, eight ounces.
Syrup of Vinegar, one ounce.
Mix.—It is refreshing, stimulant, and re-
solvent.

REFRIGERANT MIXTURE.

*Mixtura e Kali nitrico seu nitrata seu re-
frigerans seu sedativo-refrigerans.* (WUR-
CEB.; SAUNDERS; SAINTE-MARIE)

R. Nitre, one dram.
Water, eight ounces.
Simple Oxymel, one ounce.

Mix.—Two table spoonfuls to be taken
every hour. (WURCEB.)

R. Nitre, half an ounce.
Water, six ounces.
Vinegar, } of each,
Syrup of Raspberry } one ounce.
Vinegar, }

Mix.—One or two spoonfuls to be taken
every two hours. (SAINTE-MARIE)

R. Nitre, one dram.
Elder-flower Water, eight ounces.
Syrup of Mulberries,
one ounce and a half.

Mix.—Dose, two table spoonfuls every
hour. (SAUNDERS)

REFRIGERANT LINCTUS. (SAUNDERS)

R. Nitre, half a dram.
Syrup of Raspberries, one ounce.
Gooseberries, three ounces.

Mix.

LITHONTRIPTIC INJECTION. (BORIES)

R. Nitre, two ounces.
Nitric Acid, two drams.
Flemish Glue, one ounce.
Water, one pound.

Mix.—This injection has been proposed
as a solvent for calculi formed of phosphate
and oxalate of lime.

APERITIVE POTION. (RATIER)

R. Nitre, one scruple.
Decoction of the Five Roots,
five ounces.
Syrup of the Five Roots, two ounces.

Mix.—The syrup may be replaced by
oxymel, and half an ounce of solution of
acetate of ammonia may be added if there be
occasion.—It is reputed useful in dropsies.

RATIER quotes, under the name of *Ni-
trous Potion*, a similar mixture, composed
of eighteen grains of nitre, four ounces of
decoction of dog-grass, and one ounce of
the syrup of the five roots. They are both
diuretic.

TEMPERANT POTION. (SAINTE-MARIE)

R. Nitre, fifteen grains.

Lettuce Water, six ounces.
 Orange-flower Water, two drams.
 Syrup of Water Lily, two ounces.

Mix.—To be taken at twice or thrice, during the night, at the commencement of acute diseases.

REFRIGERANT MIXTURE. (PIERQUIN)

R. Nitre, half a dram.
 Infusion of Maidenhair, five ounces.
 Orange Juice, two ounces.
 Syrup of Pomegranate, one ounce.

Mix.—To be taken at four doses.

REFRIGERANT GARGLE.

Gargarisma refrigerans seu nitrosus.
 (SAUNDERS; SAINTE-MARIE)

R. Nitre, six drams.
 Decoction of Barley, one pound.
 Oxydel, two ounces.

Mix. (SAINTE-MARIE)

SAUNDERS prescribes three ounces of honey of roses instead of the oxydel.

ANTI-PHLOGISTIC POTION. (HAMB.)

R. Nitre, three drams.
 Decoction of Marshmallows, eight ounces.
 Oxydel, two ounces.

Mix.—To be taken by spoonfuls.

ANTI-PHLOGISTIC DRINK. (AUGUSTIN; CADET DE GASSICOURT; ELLIS)

R. Pearl Barley, twelve ounces.
 Water, three pounds.

Boil till the grain is burst; then add to the decoction

Nitre, one dram and a half.
 Syrup of Vinegar, two ounces.

A small cupful every hour in inflammatory fevers. (CADET DE GASSICOURT)

AUGUSTIN prescribes two pounds of decoction of barley, one dram of nitre, and one ounce of oxydel.—ELLIS.—one pound of decoction of barley, two drams of nitre, and one or two ounces of lemon juice.

DIURETIC POTION. (CADET DE GASSICOURT)

R. Nitre, half an ounce.
 Decoction of Barley, two pounds.
 Oil of Almonds, four ounces.
 Syrup of Marshmallows, six ounces.

Mix.—To be taken by glassfuls, every four hours; in dysury and strangury.

RESOLVENT STOMACHIC POTION.

Decoctum stomachicum rosolvens. (BRERA)

R. Bitter Decoction, one pound.

Extract of Dandelion, one dram.
 Nitre, one dram and a half.

Mix.—To be taken gradually in engorgements of the abdominal viscera with languid digestion.

R. Bitter Decoction, one pound.
 Nitre, two drams.

Mix.—To be taken in the course of the day.

REFRIGERANT LAVEMENT. (AUGUSTIN; SAUNDERS)

R. Nitre, half a dram.
 Decoction of Linseed, five ounces.
 Linseed Oil, } of each, one ounce.
 Oxydel, }

Mix. (SAUNDERS)

R. Decoction of Barley, ten ounces.
 Nitre, one dram.
 Honey, one ounce.

Mix. (AUGUSTIN)

R. Whey, four ounces.
 Nitre, one dram.
 Oxydel, one ounce.

Mix. (AUGUSTIN)

NITROUS EMULSION. (FORM. HÔP. FR.; AUGUSTIN; BRERA; SAUNDERS)

R. Nitre, one scruple.

Dissolve it in six ounces of an emulsion prepared with

Sweet Almonds, blanched, six drams.
 Seeds of White Poppy, one dram.
 Lime-flower Water, a sufficient quantity.

Add to the solution

Syrup of Lemon Juice, one ounce.

Mix.—A spoonful to be taken every hour. (SAUNDERS)

R. Sweet Almonds, blanched, half an ounce.
 Water, one pound.
 Nitre, half an ounce.
 Syrup of Marshmallows, one ounce.

Make an emulsion. (AUGUSTIN)

R. Almond Emulsion, one pound.
 Nitre, one dram.
 Despumated Honey, one ounce.

Mix. (BRERA)

FORM. HÔP. FR. simply prescribes three-tenths of nitre to be added to simple emulsion.

BARTHEZ' EMULSION. (BORIES)

R. Nitre, ten grains.

Dissolve it in an emulsion prepared with

Major Cold Seeds, half an ounce.
 Decoction of two drams of Cinchona, eight ounces.

Add to the solution

Syrup of Maidenhair, one ounce.

Mix.—To be taken at twice in hæmoptysis and other affections of the chest.

VINOUS POTION. (PIERQUIN)

R. Nitre, half a dram.
Water, two pounds.
White or Red Wine, five ounces.

Mix.—As a stimulant to the alimentary canal.—To be taken by glassfuls.

REFRIGERANT POTION. (SAUNDERS)

R. Nitre, } of each,
White Sugar, } two scruples.
Simple Alexitereous Water, two ounces.
Essence of Pimenta, two drams.

Mix.—The name of this potion is misapplied; it is stimulant.

SEDATIVE AND DIURETIC POTION. (PIERQUIN)

R. Nitre, twenty grains.
Distilled Water of Wall
Pellitory, } of each,
Parsley, } one
Bean- } ounce.
skins,
Alcoholized Nitric Acid, one scruple.
Syrup of Lemon, one ounce.

Mix.—For one dose, to be repeated every four hours, if the pain does not abate, adding, if necessary, one ounce of syrup of poppy.

SEDATIVE AND PURGATIVE JULEP. (PIERQUIN)

R. Nitre, three drams.
Lettuce Water, }
Rose Water, } of each,
Red Poppy Water, } three ounces.
Purslain Water, }
Syrup of Violets, } of each,
Pomegranate, } one ounce
and a half.

Mix.—To be taken in the course of the day, at three doses.

TEMPERANT JULEP. (SAINTE-MARIE)

R. Nitre, twelve grains.
Lettuce Water, }
Purslain Water, } of each, two ounces.
Syrup of Violets, }
Lemon, one ounce.
Orange-flower Water, half an ounce.

Mix.—To be taken in the course of the night, at two or three doses.

B.

Preparations in which the action of Nitre is more or less modified by that of other substances.

1. By Camphor.

NITRO-CAMPHORATED POWDER.

Pulvis nitroso-camphoratus. (PH. SYPH.)

R. Nitre, ten grains.
Camphor, from four to eight grains.
Gum Arabic, one scruple.

Mix.—To be taken at twice or thrice.

R. Nitre, } of each, fifteen grains.
Sugar, }
Camphor, from two to four grains.

Mix.—To be taken at twice or thrice.

CAMPHORATED TEMPERANT POWDER.

Pulvis temperans camphoratus. (DAN.; MIL. COPEN.)

R. Refrigerant Powder, eighteen parts.
Camphor, one part.

Mix.

ALEXIPHARMIC POWDER. (WIRTEM.)

R. Nitre, one ounce and a half.
Crab's Eyes saturated with Citric
Acid, one ounce.
Armenian Bole, one ounce and a half.
Camphor, one dram.

Pulverize, and mix.—Dose, from one to two scruples.

CAMPHORATED DIURETIC LIQUOR. (CADET DE GASSICOURT)

R. Nitre, two ounces.
Camphor, in powder, one dram.
Water, two pounds.

Triturate, and mix.—To be taken by spoonfuls.

R. Nitre, four ounces.
Water of Wall Pellitory, two pounds.

Dissolve, and add

Acetic Acid, four ounces.
Alcohol, half a pound.
Camphor, six drams.

Agitate for a long time, and filter.—Dose, a spoonful every hour.

REFRIGERANT EMULSION. (SAINTE-MARIE)

R. Nitre, ten grains.

Dissolve it, when the liquor is cold, in an emulsion prepared with

Hemp Seed, a pinch.
Boiling Water, two pounds.

Then add to the solution

Camphor, dissolved in æther, six grains.

Syrup of Water Lily, two ounces.

Mix.—A fourth part to be taken every hour, in the evening; in gonorrhœa with chordee.

CAMPHORATED NITROUS EMULSION.
(SAINTE-MARIE)

R. Nitre, one scruple.
Camphor, eight grains.
Melon Seeds, two ounces.
Spring Water, two pounds.
Sugar, two ounces.

Make an emulsion.—Dose, a glassful every two hours, adding to each, a few drops of lemon juice.

DIURETIC AND REFRIGERANT MIXTURE.
(PIERQUIN)

R. Nitre, fifty grains.
Camphor, forty grains.
Honey, one ounce.

Triturate for a long time, and add Water, a sufficient quantity.

Mix.—It is recommended at the commencement of urethritis.

CAMPHORATED NITROUS MIXTURE.

Mixtura nitrato-camphorica. (VAN MONS)

R. Nitre, one dram and a half.
White Sugar, three drams.
Camphorated Oil, from half a dram to one dram.

Mix exactly together, and triturate with Honey, one ounce and a half.
Sage Water, ten ounces.

Mix.—It is excitant, and is esteemed anti-spasmodic and sudorific, and is recommended in exanthematic fevers, and especially in pleurodynia.—Dose, one or two spoonfuls every two hours.

2. *By Iron.*

FERRUGINOUS DIURETIC POWDER.
(HAMB.)

R. Diuretic Powder, any quantity.
Levigated Iron Filings, four scruples.
Mix, and divide into eight doses.

3. *By Hydrochlorate of Ammonia.*

REFRIGERANT POWDER. (VAN MONS)

R. Nitre, two parts.
Hydrochlorate of Ammonia, one part.
Triturate together.

It is to be enclosed in a bladder and moistened with water or mixed with pounded ice, and applied as a refrigerant to the surface of the body; it is principally used in cases of strangulated hernia.

SCHMUCKER'S REFRIGERANT LOTION.

Fotus refrigerans seu frigidus Schmuckeri, Mixtura frigorifica seu anti-phlogistica.
(FULD.; HANN.; AUGUSTIN; BRERA; BRUGNATELLI; CADET DE GASSICOURT; NIEMANN; PIERQUIN; SPIELMANN; SWEDIAUR)

R. Nitre, eight ounces.
Hydrochlorate of Ammonia, four ounces.
Water, twenty pounds.

Dissolve and add

Vinegar, two pounds.

Mix. (CADET DE GASSICOURT)

NIEMANN prescribes one pound of nitre, one pound of hydrochlorate of ammonia, six hundred and forty ounces of water, and ninety-six of vinegar.—PIERQUIN—one pound of nitre, eight ounces of the hydrochlorate, twelve pounds of water, and four of vinegar.—BRUGNATELLI—two ounces of nitre, one ounce of hydrochlorate, and two pounds of spring water acidulated with acetic acid.—SPIELMANN—half an ounce of nitre, two ounces of hydrochlorate, one hundred and twenty ounces of water, and twelve ounces of vinegar.—HANN.—one ounce of hydrochlorate, one ounce of nitre, sixteen ounces of water, and sixteen ounces of vinegar.—FULD.—two ounces of nitre, one ounce of hydrochlorate, half a pound of vinegar, and six pounds of water.—AUGUSTIN—four ounces of nitre, two of hydrochlorate, one pound of vinegar, and eight pounds of water.—BRERA—three drams of nitre, two of hydrochlorate, one pound of vinegar, and four pounds of water.

This liquor, which is applied cold by means of linen four or six times folded, and renewed every hour, is useful in contusions and ecchymoses.

ANTI-GONORRHOIC TINCTURE. (PIERQUIN)

R. Nitre, two ounces.
Hydrochlorate of Ammonia, two drams.
Water, one pound and a half.

Dissolve at a moderate heat, and add

Camphor, two drams.
Alcohol, a sufficient quantity.

Macerate till perfectly cold and filter.—Dose, a spoonful in a glass of water.

4. *By Opium.*

BOLUSES FOR COUGH. (PIERQUIN; SAINTE-MARIE)

- R. Nitre, twelve grains.
 Pill of Hounds-tongue, four grains.
 Conserve of Red Roses, two scruples.

Make two boluses, to be taken in the evening at an interval of two or three hours.

ANTI-NEPHRITIC POTION. (FERR.)

- R. Poppy Heads, six ounces.
 Water, three pounds.

Boil down to eight ounces, and add when strained,

- Nitre, one ounce.

Dissolve.—Dose, one or two drams daily, in eight ounces of decoction of linseed or of marshmallows.

5. *By Squill.*

DIURETIC POWDER.

Pulvis diureticus. (PIERQUIN; SWEDIAUR)

- R. Calcined Nitre, eight parts.
 Root of Swallow-wort, six parts.
 Squill, one part.

Mix. (PIERQUIN; SWEDIAUR)

SWEDIAUR also prescribes fifteen grains of nitre, ten of squill, and ten of pimenta.

- R. Nitre, } of each,
 Supertartrate of Potass, } four scruples.
 Squill, } eight grains.
 Ginger, } sixteen grains.

Mix, and divide into eight parcels. (HAMB.)

It is recommended in dropsies, particularly in ascites, in asthma, and in chronic diseases of the bladder, as diuretic and sudorific.—The first formula is from VAN HELMONT.—Dose, from ten to twenty grains, two, three, or four times a day.

DIURETIC MIXTURE. (SWEDIAUR)

- R. Nitre, two drams.
 Water, eight ounces.
 Powdered Squill, half a dram.
 Oxymel of Squill, one ounce.

Mix.—It is recommended in dropsies and in affections of the mucous membrane of the lungs.—Dose, two or three spoonfuls, thrice a day.

DIURETIC AND RESOLVENT POTION.

Decoctum resolvens et diureticum. (BRERA)

- R. Decoction of Dog-grass, one pound.
 Extract of Buckbean, } of each,
 Nitre, } one dram.
 Oxymel of Squill, one ounce.

Mix.—To be taken gradually, in engorgements of the liver and of the spleen.

DIURETIC POTION.

Potus anti-phlogisticus diureticus. (BRERA)

- R. Decoction of Asparagus Roots, two pounds.
 Nitre, two drams.
 Oxymel of Squill, half an ounce.

Mix.—To be taken gradually.

6. *By acidulous or neutral salts.*

REFRIGERANT MIXTURE. (AUGUSTIN; SWEDIAUR)

- R. Nitre, } of each,
 Supertartrate of Potass, } two drams.
 Water, } eight ounces.
 Syrup of Raspberry Vinegar, one ounce.

Mix.—A spoonful to be taken every two hours. (SWEDIAUR)

- R. Nitre, } of each,
 Supertartrate of Potass, } three drams.
 Water, } six ounces.
 Lemon Juice, } of each,
 Syrup of Lemon, } one ounce.

Mix.—Dose, a spoonful every two hours. (AUGUSTIN)

SIMPLE SALINE MIXTURE. (SAUNDERS)

- R. Nitre, } of each,
 Supertartrate of Potass, } six drams.
 Fennel Water, } five ounces.
 Sulphuric Ether, } one dram.
 Syrup of Poppy, } two ounces.

Mix.—To be administered in the same manner as the preceding.

REFRIGERANT AND DIURETIC DRINK.

Potus anti-phlogisticus et diureticus. (BRERA)

- R. Decoction of Dog-grass, two pounds.
 Supertartrate of Potass, one ounce.
 Nitre, three drams.
 Sugar, one ounce.

Mix.—To be taken gradually in the course of the day.

NITRATED LOZENGES.

Tabellæ de tribus, Saccharum nitratum. (SAUNDERS)

- R. Nitre, one ounce.
 Supertartrate of Potass, half an ounce.
 White Sugar, one pound.

DIURETIC TEMPERANT POWDER.

Pulvis temperans diuretica, Pulvis temperans seu temperans albus seu nitrosus seu Nitri

tartaricus. (FENN.; HANN.; LIPP.; BRERA)

R. Nitre, three drams.
Supertartrate of Potass, half an ounce.
Mix, and divide into eight parcels.—Dose, one every two hours.

R. Nitre, one part.
Supertartrate of Potass, three parts.

Mix. (HAMB.)

LIPP. prescribes an equal part of each.

R. Nitre, } of each,
Supertartrate of Potass, } one ounce.
White Sugar, three ounces.

Pulverize each separately, and mix.

(FENN.)

Dose, half a dram to adults.

TARTARIZED NITROUS POWDER.

Pulvis Nitri tartaricus. (SUEC.)

R. Nitre, ten parts.
Tartaric Acid, two parts.
Subcarbonate of Magnesia, one part.
White Sugar, thirty parts.

Triturate the acid with the subcarbonate, then add the sugar, and lastly the nitre.

REFRIGERANT MIXTURE. (SWEDIAUR)

R. Nitre, three drams.
Tartaric Acid, one scruple.
White Sugar, one ounce.
Water, eight ounces.

Dissolve.—One or two ounces of oil of almonds are sometimes added.—Dose, a spoonful every hour.

BOUILLON LAGRANGE'S TEMPERANT POWDER. (CADET DE GASSICOURT)

R. Nitre, three pounds.
Supertartrate of Potass, four pounds and a half.
Boracic Acid, one pound.
Pulverize and sift.

It is temperant and laxative, and is recommended in diseases of the skin, in some catarrhal affections, and in certain retentions of urine.—Dose, half a dram in a pound and a half of water, to which two or four drams of effloresced sulphate of soda may be added, and a glassful taken every morning.

ROSENSTEIN'S TEMPERANT POWDER.

Pulvis Nitri oxalicus seu nitrosus. (SUEC.; NIEMANN; VAN MONS)

R. Nitre, one ounce and a half.
Superoxalate of Potass, half an ounce.
Lemon Lozenges, seven ounces and a half.

Pulverize and mix. (NIEMANN)

R. Nitre, three ounces.
Superoxalate of Potass, one ounce.
White Sugar, ten ounces.

Pulverize separately and mix. (SUEC.)

VAN MONS prescribes one ounce of superoxalate, two ounces of nitre, and three of sugar.

DIAPHORETIC POWDER. (ELLIS)

R. Nitre, one dram.
Tartarized Antimony, one grain.
Proto-chloruret of Mercury, six grains.

Mix, and divide into six doses; one to be taken every two hours in syrup.

REFRIGERANT TROCHISCS.

Trochisci refrigerantes seu temperantes.
(FULD.; SPIELMANN)

R. Nitre, six drams.
Superoxalate of Potass, two drams.
White Sugar, six ounces.
Mucilage of Tragacanth, a sufficient quantity
to make a mass for trochiscs. (FULD.)

SPIELMANN prescribes one ounce of nitre, four drams of the superoxalate, six ounces of sugar, half a scruple of essence of lemon, and mucilage of tragacanth.

Dose, from one to four drams.

SALINE FEBRIFUGE POWDER. (VAN MONS)

R. Nitre, two parts and a half.
Subcarbonate of Potass, one part and a half.
Supertartrate of Potass, two parts.
Sulphate of Potass, three parts.
Nutmeg, one part.
Pulverize.

TEMPERANT POWDER. (HANN.; OLDEN.; CAST. BORUS.; BORUS.; ROSS.; VAN MONS)

R. Nitre, } of each,
Sulphate of Potass, } an equal part.
Pulverize.

UNZER'S TEMPERANT POWDER. (BRUNS.; PIDERIT; SPIELMANN)

R. Nitre, } of each,
Sulphate of Potass, } one ounce.
Compound Powder of Tragacanth, two drams.

Mix. (PIDERIT)

R. Nitre, } of each,
Sulphate of Potass, } an
Subcarbonate of Magnesia, } equal part.
Mix. (BRUNS.)

SPIELMANN gives the same formula, but substitutes prepared crab's eyes for the subcarbonate of magnesia.

7. By Sulphur.

STRUENSEE'S PYRIC POWDER.

Pulvis pyricus. (NIEMANN)

- R. Nitre, six parts.
Sulphur, }
Charcoal, } of each, one part.

Pulverize and mix.

It is recommended to cauterize the bites of rabid animals.—It is a means not to be recommended, as the cauterization is never but superficial.

PROTO-CHLORATE OF POTASSIUM.

Alkali vegetabile salito - dephlogisticatum, Chloruretum Potassæ oxydatum, Haloïdum oxygenatum, Murias Potassæ hyperoxygenatum seu oxygenatum, Oxygenochloruretum Potassii, Potassæ Euchloras. (GALL.; FERR.; BRUGNATELLI; NIEMANN; SWEDIAUR; VAN MONS)

- R. Muriate of Soda, } of each, five parts.
Sulphuric Acid, }
Fused Potass, five parts and a half.
Oxide of Manganese, eight parts and a half.

Add water to the potass till the solution marks 30 degrees B.; filter the solution and pour it into a long, narrow, glass cylinder: pass into it a current of chlorine gas produced by the mixture of the three other substances, collect the crystals which are deposited, and wash them with a little cold water. (GALL.; FERR.; BRUGNATELLI; NIEMANN; SWEDIAUR; VAN MONS)

HENRI; and SWEDIAUR prescribe subcarbonate.—HENRI requires the solution to mark 20 degrees, and the crystals to be redissolved in boiling water.

It is excitant, and recommended in scurvy, tetters, chronic hepatitis, and syphilitic diseases.—Dose, from fifteen to thirty grains.

SOLUTION OF CHLORATE OF POTASS.

Solutio Muriatæ Potassæ oxygenatæ, Liquor ad Ulcera atonica. (NIEMANN; SWEDIAUR; PH. SYPH.)

- R. Chlorate of Potass, one dram.
Distilled Water, twelve ounces.

Dissolve.

It is to be applied, with a brush or by means of lint, as a stimulant to indolent ulcers.

OXYMURIATED ALKALINE LIQUOR.

Aqua alkalina oxymuriatica. (NIEMANN)

- R. Decrepitated Muriate of Soda, two pounds.
Oxide of Manganese, one pound.
Water, }
Sulphuric Acid, } of each, two pounds.

Mix the muriate and oxide together, put them into a matrass, and add the water; then, having gradually added the acid, pass the current of gas which is disengaged, into a solution prepared with

Subcarbonate of Potass, four ounces.
Water, twenty-nine ounces.

The saturated liquor weighs 1.087.

OXYGENATED POTION.

Potus oxygenata. (SWEDIAUR)

- R. Solution of Chlorate of Potass, two pounds.
Water, four pounds.

Mix.

It is excitant, and has been extolled in cases of general debility.—Dose, one or two pounds daily.

PROTO-HYDROCHLORATE OF POTASSIUM.

Murias seu Hydrochloras potassæ, Sal digestivus seu febrifugus Sylvii, Spiritus Salis marini coagulatus. (BAVAR.; BRUNS.; HISP.; GALL.; SARD.; WIRTEM.; NIEMANN; SPIELMANN; VAN MONS)

- R. Subcarbonate of Potass, any quantity.
Water, a sufficient quantity
to effect the solution. Filter, and add to it,
Hydrochloric Acid, a sufficient quantity
to saturate the potass. Evaporate the liquor till it marks 30 degrees, and set it aside that crystals may form. (BAVAR.; HISP.; GALL.; WIRTEM.)

- R. Muriate of Soda, five parts and a half.
Calcined Subcarbonate of Soda, six parts and a half.
Water, sixteen parts.

Boil, filter, and crystallize, keeping the liquor hot till crystals cease to be formed. (VAN MONS)

SARD.; NIEMANN; and SPIELMANN direct the residuum of the preparation of the subcarbonate of ammonia to be used.

Sulphate or supertartrate of potass may also be dissolved in distilled water, and solution of hydrochlorate of lime poured into the liquor till no more precipitate is formed. This process is recommended by CHEVALIER and IDT.

When in the crystallized state this salt is called *Chloruret of Potassium*.

It is excitant, and is classed among the resolvents and deobstruents, and is recommended in quartan and tertian fevers as sudorific and diuretic.—Dose, one or two scruples, in four times as much water.

HYDROSULPHATE OF POTASS. (VAN MONS)

R. Calcined Caustic Potass, one part.
Water, six parts.

Pass a current of hydrosulphuric acid gas into the solution, set it apart that it may become clear, and preserve it.

PROTO-ACETATE OF POTASSIUM.

Acetis seu Acetos Potassæ seu Lixivie seu kalicum, Alkali Tartari Aceto saturatum, Alkali vegetabile cum Aceto, Arcanum Tartari, Kali acetatum seu aceticum Lixivie acetata, Magisterium Tartari purgans, Oxytartarus, Sal diureticus, Tartarus acetosus, Terra foliata Tartari.—Acetate of Potass, Acetate of Protoxide of Potassium. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COHEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

1. In the solid state.

Potassæ Acetas.

R. Subcarbonate of Potass, one pound and a half.
Strong Acetic Acid, }
Boiling Distilled } of each,
Water, } two pints.

Add the subcarbonate to the acid and water mixed together till all effervescence has ceased, and filter the liquor. Evaporate it then in a water-bath till no more bubbles rise; then expose it to a gradually augmented heat, and continue the evaporation till a pellicle forms; then remove this, and dry it in bibulous paper. Repeat the evaporation again and again, removing the pellicles as they form, and drying them as the first. (LOND.)

Vol. II.

R. Pure Subcarbonate of Potass, one pound.
Weaker Acetic Acid, a sufficient quantity.

Boil the subcarbonate in five pounds of the acid, and add more acid at different times, till the watery part of the former portion being nearly dissipated by evaporation, the recently added acid occasions no effervescence, which will be when about twenty pounds of it are consumed; then evaporate slowly to dryness. Liquefy this impure salt with a moderate heat for a short time, then dissolve it in water and filter the solution through paper. If the liquefaction has been properly effected, the filtered liquor will be limpid, if not, it will be of a brown colour. Evaporate this liquor in a shallow glass vessel, so that when removed from the fire it may form into a crystalline mass, which should be preserved in closely-stopped vessels. (EDINB.)

R. Carbonate of Potass from
Crystals of Tartar, any quantity.

Add to it at different times, about five times its weight of distilled vinegar moderately heated. When the effervescence has ceased and the fluid has been evaporated a little, add, at intervals, more distilled vinegar till it entirely ceases to excite effervescence; then evaporate to dryness, and having encreased the heat a little, liquify cautiously the mass. Dissolve the salt in water when cold; filter the solution, and boil it till, on removing it from the fire, it concretes into a crystalline mass, which should be very white. Enclose it immediately in well-stopped bottles. (DUBL.)

BATAV.; BELG.; BRUNS.; DAN.; HISP.; FERR.; FENN.; FULD.; GENEV.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SUEC.; WIRTEM.; COXE; PIDERIT; SPIELMANN; and SWEDIAUR give the same process as DUBL.

R. Dry and pulverized Subcarbonate of Potass, one pound.
Strong Vinegar, two or three pounds.

3 L

Heat the salt in a glass capsule, and gradually add the vinegar to it; when the effervescence has ceased distil the liquor in a water-bath to dryness; add fresh vinegar to the residuum, and distil again to dryness: continue thus till the vinegar passes over unaltered. Then dry the salt at a moderate temperature, dissolve it in thrice its weight of alcohol, filter the liquor, distil it in a water-bath, and preserve the dry residuum. (PALAT.; SARD.)

R. Subcarbonate of Potass, one part.

Dissolve it in a small quantity of water, and add to the solution

Distilled Vinegar, eight parts.

Boil to the consistence of syrup, and add more vinegar to perfect saturation; evaporate the liquor slowly, constantly stirring it: dissolve the residuum in distilled water, filter the solution, and finally evaporate it with a gentle heat. (AMST.; ANT.; HERBIP.; BRUGNATELLI)

R. Subcarbonate of Potass, any quantity.

Throw it by portions into

Distilled Vinegar, a sufficient quantity to dissolve it completely. Strain the liquor, and evaporate it in a silver capsule; when it is reduced to about one fourth, let it cool; then set it apart, and having decanted it, add vinegar gradually to it; lastly, evaporate it to dryness. (GALL.)

VAN MONS, who also indicates this process, requires the acid to be slightly heated before throwing in the salt, and the liquor to be rapidly dried by the aid of a brisk heat, constantly stirring.

To render the salt the whiter it has been recommended to pass the liquor through a charcoal filter, or to treat it without heat with animal charcoal, or to subject the salt to several successive solutions and evaporations.

R. Dried Acetate of Lime, seven parts and a half.

—— Subcarbonate of Potass, eleven parts and three fourths.

Mix with trituration, adding the quantity of water which is strictly necessary to dissolve the acetate produced; pour this liquid at several times, then express, and dry immediately as above. (VAN MONS)

R. Sulphate of Potass, eight parts and a fourth.

Crystallized Acetate of Barytes, thirteen parts.

Mix together, add water, and proceed as above. (VAN MONS)

VAN MONS also recommends the tartrate of potass to be decomposed by the acetate of lime mixed in equal proportions and dissolved in water to saturation; the tartrate

is to be poured into the acetate, the liquor expressed, the residuum washed, and mixed liquors evaporated.

II. In the liquid state.

Acetas Potassæ liquidus seu aquosus, Liquor Ferræ foliatæ seu Tartari vegetabilis seu Kali acetici seu digestivus Boerhaavii seu Alkali acetici nigri seu Terræ foliatæ Tartari seu Salis Alkali vegetabilis acetati, Salsilago Oxytartari, Acetas Lixivie solutus, Mixtura salina pauperum. (AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; MIL. COPEN.; FENN.; FORM. HÔP. FR.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

R. Subcarbonate of Potass, any quantity.
Concentrated Acetic Acid, a sufficient quantity

to saturate the salt. Add the acid gradually, stirring the mixture with a glass tube. The filtered liquor should weigh from 1.230 to 1.235. (BAVAR.)

R. Subcarbonate of Potass, twelve ounces.
Vinegar, a sufficient quantity

for perfect saturation. Mix, digest for several days, with occasional agitation; add more vinegar, if necessary, and evaporate till only eighty-eight ounces remain—BATAV.; or, till the liquor marks seven degrees. (BELG.)

This formula presents numerous variations:—HASS.—Mix together one pound of subcarbonate of potass and a sufficient quantity of vinegar to saturate the salt; then evaporate the liquor to three pounds, and filter it.—AUSTR.—Saturate subcarbonate with distilled vinegar (1,005), and evaporate till the liquor weighs 1,200.—FERR.; HENRI; and FEE—Saturate subcarbonate with vinegar, and add a sufficient quantity of water.—MIL. COPEN.; FULD.; CAST. BORUS.; SPIELMANN.; and VAN MONS—Dissolve in vinegar at six degrees as much subcarbonate as possible; filter and preserve the liquor.—HAMB.—Saturate one ounce of subcarbonate with a sufficient quantity of vinegar, and evaporate to six ounces.—FORM. HÔP. FR.—Mix together one part of subcarbonate and sixteen parts of acid at thirteen degrees; heat to perfect saturation, and preserve.

R. Acetate of Potass, one part.
Distilled Water, two parts.

Dissolve. (AMST.; BRUNS.; DAN.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SWEDIAUR.)

HASS.; and PIDERIT recommend the salt to be left in the open air to deliquesce, and the liquor to be filtered.

R. Subcarbonate of Potass, two drams.
Vinegar, a sufficient quantity
to saturate the salt. Add to the liquor
Water, from two to four ounces.
Simple Oxy mel, two ounces.

Mix. (NIEMANN)

R. Subcarbonate of Potass, one pound.
Add to it by portions

Vinegar, a sufficient quantity
to saturate it. At the end of three days,
filter the liquor, and evaporate it in a sand-
bath, till reduced to three pounds; then add

Alcohol, two ounces.

Preserve it in a well-stopped bottle.
(LIPP.)

The acetate of potass is an excitant which
acts, according to doses and circumstances,
as resolvent, diuretic, and laxative. It is
employed in dropsies, jaundice, and in ob-
structions of the abdominal viscera. The
dose is from a scruple to a dram or more.
Of the solution from forty to eighty drops
are usually given, in an appropriate vehicle.

It is incompatible with the mineral acids,
the decoction of tamarinds, the deuto-chlo-
ruret of mercury, the nitrate of silver, the
sulphates of soda and of magnesia, the hy-
dro-chlorate of ammonia, and the supertar-
trate of potass.

A.

*Preparations in which the action of the Salt
is not modified, at least sensibly, by that
of other substances.*

ANTI-LACTEAL BOLUS. (SAINTE-MARIE)

R. Acetate of Potass, one scruple.
Conserve of Marigold, a sufficient quantity.

Mix.

To be taken in the morning, for twenty
days; it may be repeated in the evening, if
necessary. A cupful of a decoction of cher-
vil and celery should be taken with each
bolus.

DIURETIC POTION. (RATIER; SAUNDERS;
SAINTE-MARIE)

R. Acetate of Potass, half a dram.
Juice of Chervil, twelve ounces.
Compound Syrup of Smalage, two ounces.

Mix.

R. Acetate of Potass, two drams.
Infusion of Lime-flowers, four ounces.
Cinnamon Water, one dram.
Syrup of Vinegar, one ounce.

Mix. (RATIER)

R. Prepared Crab's Eyes, three drams.
Vinegar, one ounce and half.

After the cessation of the effervescence,
add

Spring Water, eight ounces.
Acetate of Potass, one dram.
Syrup of Fennel, one ounce.

Mix. (SAUNDERS)

To be taken by spoonfuls.—In the last
formula the diuretic effect must be referred
principally to the action of the acetate of
lime.

APERITIVE POTION. (SAINTE-MARIE)

R. Acetate of Potass, two drams.
Infusion of Chamomile, six ounces.
Hordeated Cinnamon Water, two ounces.
Extract of Milfoil, four drams.

Mix.—Dose, a spoonful thrice a day; at
the same time taking a broth made with
four ounces of neck of mutton, and a hand-
ful of leaves of cichory, of dandelion, and
of water cress.

ANTI-ICTERIC MIXTURE. (CADET DE
GASSICOURT)

R. Acetate of Potass, } of each,
Liquid Extract of Dan- } half an
delion, } ounce.
Hemlock, from ten to twenty grains.
Fennel Water, six ounces.
Compound Syrup of Smalage, one ounce.

Mix.—Dose, two spoonfuls every four
hours.

APERITIVE POTION. (SAINTE-MARIE)

R. Acetate of Potass, half a dram.
Nitre, one scruple.
Syrup of Borage, two ounces.
Clarified Whey, one pound and a half.

Dissolve, and mix.—DESBOIS DE ROCHE-
FORT directed it to be taken in the course
of the day, but the greater part in the morn-
ing fasting.

SALINE MIXTURE. (HASS.; SUEC.; SAUN-
DERS)

R. Subcarbonate of Potass, three ounces.
Vinegar, nine ounces,
or a sufficient quantity to saturate the al-
kali. Add

River Water, one pound.
Simple Syrup, two ounces.

Mix and filter. (SUEC.)

HASS. directs the subcarbonate to be sa-
turated with vinegar, four or five ounces of
the liquor diluted with an equal quantity of
distilled water, and one ounce of syrup
added.—SAUNDERS—nine ounces of vine-

gar, one pound of water, and two ounces of sugar, to be poured into a phial; then three drams of subcarbonate added, and the bottle well-stopped and kept in a cool place.

Dose, two spoonfuls, every two or three hours.

RESOLVENT AND ALTERNATIVE CONSERVE. (BORIES)

R. Acetate of Potass, half an ounce.
Sulphate of Soda, one dram.
Juice of Cress, }
Fumitory, } of each,
Herb Mercury, } two ounces.
Dandelion, }
Sugar, a sufficient quantity.

Mix.—Dose, a spoonful, three or four times a day.

ANTI-SCROFULOUS ELECTUARY. (SAINTE-MARIE)

R. Acetate of Potass, six drams.
Conserve of Scurvy-grass, two ounces.
Extract of Dog-grass, } of each,
Dandelion, } one ounce.
Compound Syrup of Cichory, a sufficient quantity.

Mix.—Dose, a tea-spoonful, four times a day, in a glassful of weak decoction of hop.

DIURETIC POTION.

Potus diureticus simplex. (AUGUSTIN ; BRERA)

R. Decoction of Dog-grass, one pound.
Solution of Acetate of Potass, two ounces.

Mix.—To be taken gradually.

R. Subcarbonate of Potass, two drams.
Vinegar, a sufficient quantity
to saturate the alkali. Add

Water, four ounces.
Syrup of Lemon, half an ounce.

Mix.—To be taken gradually.

R. Subcarbonate of Potass, two drams.
Vinegar, a sufficient quantity.
Water, four ounces.
Syrup of Raspberries, two ounces.

Mix.—To be taken gradually.

R. Subcarbonate of Potass, two drams.
Aromatic Vinegar, a sufficient quantity.
Cinnamon Water, one ounce.
White Wine, four ounces.

Mix.—Dose, a spoonful every hour.

RESOLVENT AND DIURETIC POTION.

Decoctum resolvens et diureticum. (BRERA)

R. Decoction of Dog-grass, two pounds.
Solution of Acetate of Potass, two ounces.

Mix.—To be taken gradually, in dropsies.

R. Bitter Decoction, one pound.
Aqueous Tincture of Rhubarb, four ounces.
Acetate of Potass, two drams.

Mix.—To be taken in engorgements of the abdominal viscera.

R. Infusion of Orange Peel, one pound.
Acetate of Potass, two drams.
Sugar, one ounce.

Mix.—To be taken in asthenic inflammations.

B.

Preparations in which the action of Acetate of Potass is more or less modified by that of other substances.

1. By Alcohol.

Besides the following formulæ, the solution of acetate of potass of LIPP., previously given, should be placed here.

DIURETIC POTION. (RATIER)

R. Acetate of Potass, one ounce.
White Wine, one pound.

Dissolve.—It is extolled in dropsies which are designated passive; that is, when there is no symptom of much irritation of the intestinal canal.

ALCOHOLIC SOLUTION OF ACETATE OF POTASS. (BRUNS.; PALAT.; WIRTEM.)

R. Acetate of Potass, very dry, two ounces.
Concentrated Alcohol, eight ounces.

Digest for several days, and filter.—It used formerly to be extolled as an excellent remedy against hypochondria, gout, quartan fever, and all disorders caused by obstructions.—Dose, from thirty to sixty drops.

DELLA LENA'S LITHONTRIPTIC SOLUTION. (NIEMANN)

R. Solution of Acetate of Potass, two ounces and a half.
Subcarbonate of Potass, one ounce.
Distilled Water, four ounces.
Alcohol, two ounces.

Digest till the salt is perfectly dissolved.—From ten to thirty drops are to be given in any proper vehicle.

ESSENTIA DULCIS. (WIRTEM.; CADET DE GASSICOURT)

R. Acetate of Potass, two drams.
White Sugar, two ounces.
Water, one ounce and a half.

Boil in a porcelain capsule till the mass has acquired a black tinge from the sugar

becoming candied. When cold pour upon it

Alcohol (30 degrees), twelve ounces.

Dissolve, constantly stirring, and filter.
(CADET DE GASSICOURT)

WIRTEM. directs the mass to be dissolved in the first three pounds of liquor obtained by distilling together four pounds of proof spirit, eight ounces of alum, and one pound of water.

It is stimulant, tonic, alterative, balsamic, and diuretic.—Dose, from thirty to sixty drops, in proper drink, to adults; and from three to twelve, to children.

2. By Antimony.

DEOBSTRUENT MIXTURE.

Liquor Kali acetici cum Rheo. (AUGUSTIN; PIDERIT)

R. Subcarbonate of Potass, one dram.
Vinegar, a sufficient quantity
to saturate the alkali. Add to the liquor
Aqueous Tincture of Rhubarb, one ounce.
HUXHAM'S Antimonial Wine, one dram.

Mix.

ANTI-SCROFULOUS ELECTUARY. (BORIES)

R. Acetate of Potass, } of each,
Burnt Sponge, } two drams.
Proto-chloruret of Mercury, }
twelve grains.
Sulphuret of Antimony, one dram and a half.
Jalap, sixteen grains.
Sublimed Sulphur, one dram.
Simple Syrup, a sufficient quantity.

Mix.—It is recommended by BAUME in scrofulous engorgements of the lymphatic glands.—Dose, from twelve to twenty-four grains, twice a day.

3. By Æther.

SALINE MIXTURE. (MIL. COPEN.)

R. Solution of Acetate of Potass, two ounces.
Sulphuric Æther, one dram.
Infusion of Peppermint, six ounces.
Mix well.

RESOLVENT LIQUOR. (SAUNDERS)

R. Solution of Acetate of Potass, four ounces.
Sulphuric Æther, twenty drops.
Extract of Lesser Centaury, two drams.
Alkermes Syrup, two ounces.
Mix.

4. By Cinchona.

RESOLVENT MIXTURE. (BORIES; PIERQUIN)

R. Acetate of Potass, ten grains.
Extract of Cinchona, three grains.
Hemlock, two grains.
Infusion of Heart's-ease Leaves, three ounces.

Mix.—It is recommended by BAUME in tabes mesenterica.—Dose, a spoonful, three or four times a day.

FEBRIFUGE AND APERITIVE ELECTUARY. (BORIES)

R. Solution of Acetate of Potass, two ounces.
Extract of Cinchona, two drams.
Cascarilla, one dram.
Compound Tincture of Aloes, one ounce.

Mix.—GAUBIUS extolled it in intermittent fevers complicated with obstructions or dropsy.—Dose, thirty drops, in a glass of water, every two hours, on the days intervening between the paroxysms.

5. By Squill.

SCILITIC LIQUOR. (HAMB.)

R. Subcarbonate of Potass, a sufficient quantity.
Vinegar of Squill, one ounce.
Mix to saturation.

KEUP'S SCILITIC ESSENCE. (BRUGNATELLI; CADET DE GASSICOURT)

R. Subcarbonate of Potass, half an ounce.
Vinegar of Squill prepared with distilled Vinegar, two ounces.
Boil to the consistence of honey, and add, Alcohol (36 degrees), six ounces.
Digest for several days, and decant.

This liquor has been recommended in asthma and dropsy.—Dose, from forty to sixty drops in a proper vehicle.

MILMANN'S DIURETIC POTION.

Haustus diureticus Milmanni. (NIEMANN)

R. Subcarbonate of Potass, fifteen grains.
Distilled Vinegar, a sufficient quantity
to dissolve the alkali. Add to the liquor,
Water, half an ounce.
Peppermint Water, one ounce.
Pimenta Water, } of each, one dram.
Vinegar of Squill, }
Simple Syrup, two drams.

Mix.—Dose, from half a dram to one dram, every six hours, in dropsy.

RESOLVENT MIXTURE. (SAUNDERS)

- R. Acetate of Potass, two drams.
 Oxymel of Squill, two ounces.
 Fennel Water, }
 Parsley Water, } of each, six ounces.
 Juniper Water, }

Dissolve and mix.

- R. Solution of Acetate of Potass, half an ounce.
 Extract of Elicampane, two drams.
 Oxymel of Squill, one ounce.
 Hyssop Water, six ounces.
 Mix.

DIURETIC AND SEDATIVE POTION. (PIERQUIN)

- R. Acetate of Potass, } of each, one dram.
 Oxymel of Squill, }
 Lime-flower Water, four ounces.
 SYDENHAM'S Laudanum, fifteen drops.

Compound Syrup of Smallage, one ounce.

Mix.—To be taken by spoonfuls.

RESOLVENT ELIXIR. (CADET DE GASSICOURT)

- R. Subcarbonate of Potass, two ounces.
 Vinegar of Squill, a sufficient quantity.

Saturate, and add,

- Chamomile Water, four pounds.
 Tincture of JAMES'S } of each,
 Antimony, } one
 Tartarized Tincture of Iron, } ounce.
 Extract of Fumitory, } of each,
 ———— Buckbean, } half
 ———— Milfoil, } an ounce.
 ———— Blessed Thistle, }

Mix and filter.—To be taken by teaspoonfuls, every two hours, as a stimulant to the alimentary canal.—It has been recommended in dropsy.

SUPER-PROTOXALATE OF POTASSIUM.

Oxalas acidulum Potassæ seu Potassæ acidulus seu basicus, Superoxalas kalicum seu Potassæ, Hyperoxalas Potassæ, Acetoselæ Sal, Oxalium, Kali oxalicum acidulum, Oxalas subpotassicus, Binoxalas Potassæ.
 —Super-oxalate of Potass, Salt of Sorrel.

I. Superoxalate of Potass of commerce.

ANT.; BATAV.; BAVAR.; BELG.; DAN.; FENN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; COXE.

II. Superoxalate of Potass prepared expressly.

BRUNS.; FERR.; FULD.; PALAT.; SARD.;

WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SPIELMANN; VAN MONS.

- R. Fresh Leaves of Wood Sorrel, any quantity.

Bruise them in a stone mortar with a wooden pestle; express the juice, heat it in a water-bath, and add to it white of egg beaten up. When clarified, evaporate the liquor in a water-bath, and crystallize in a cool place. (BRUNS.; FERR.; FULD.; PALAT.; SARD.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SPIELMANN; VAN MONS)

- R. Oxalic Acid, twelve parts.
 Calcined Subcarbonate of Potass, six parts and a half.

Mix, and dissolve in the least quantity of water possible; filter, and crystallize. (VAN MONS)

It is anti-phlogistic, refrigerant, and astringent.—It is not much used.

POWDER OF SUPEROXALATE OF POTASS. (NIEMANN)

- R. Superoxalate of Potass, two scruples.
 Oleosaccharum of Lemon, two ounces and a half.

Mix.

TEMPERANT TROCHISCS. (WIRTEM.; PIDERIT)

- R. Superoxalate of Potass, two drams.
 White Sugar, eight ounces.
 Oil of Lemon, ten drops.
 Mucilage of Tragacanth, a sufficient quantity.

Make trochiscs.

PROTO-CITRATE OF POTASSIUM.

Citras Potassæ seu kalicum seu kalinum, Kali citratum, Sal Absinthii citratum. (BAVAR.; HANN.; OLDEN.; BORUS.; BRUGNATELLI; SWEDIAUR; VAN MONS)

- R. Subcarbonate of Potass, any quantity.
 Fresh Lemon Juice, a sufficient quantity to saturate the alkali. Add the juice gradually, testing the liquor from time to time to ascertain when it is perfectly saturated; then set apart for twenty-four hours, filter, and evaporate to dryness.

SIMPLE SALINE MIXTURE.

Citras Potassæ agnosus, Mixtura salina, Solutio Citratis kalici. (FENN.; LUSIT.; SUEC.; AUGUSTIN; BRUGNATELLI; NIEMANN; SWEDIAUR)

- R. Subcarbonate of Potass, three drams.
 Lemon Juice, a sufficient quantity.
 Simple Syrup, one ounce.

Spring Water, a sufficient quantity to give one pound of liquor. Mix well. (FENN.; LUSIT.; SUEC.; BRUGNATELLI; NIEMANN)

R. Subcarbonate of Potass, one dram.
Lemon Juice, three ounces.
Cinnamon Water, six ounces.
White Sugar, half an ounce.

Mix. (SWEDIAUR)

Dose, three spoonfuls every five or six hours.

COMPOUND SALINE MIXTURE.

Mixtura salina composita seu Macbridii.
(LUSIT.)

R. Lemon Juice, one ounce.
Subcarbonate of Potass, a sufficient quantity.
Mint Water, seven ounces.
Tartarized Antimony, one grain.
Simple Syrup, half an ounce.

Mix.

TONIC MIXTURE. (ELLIS)

R. Lemon Juice, one ounce and a half.
Subcarbonate of Potass, one dram.
Tincture of Cinchona, one ounce.
Cinnamon Water, three ounces.

Mix.—A spoonful to be taken every two hours.

PROTO-TARTRATE OF POTASSIUM.

Alkali vegetabile tartarisatum seu fixum tartarisatum, Kali tartaricum seu tartarisatum, Lixivia tartarisatus, Sal vegetabile, Sal panchrestum seu diureticum vegetabile, Tartarus tartarisatus seu solubilis, Tartras kalinus seu Lixiviæ seu Potassæ.
—Tartrate of Potass, Neutral Tartrate of Potass, Soluble Tartar. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; COKE.; PARIS; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

1. In the solid state.

Potassæ tartras.

R. Subcarbonate of Potass, sixteen ounces.
Supertartrate of Potass, three pounds.
Boiling Water, one gallon.

Dissolve the subcarbonate in the water and add the supertartrate in powder till effervescence ceases. Filter the solution, boil till a pellicle ap-

pears on the surface, and set it apart to crystallize. Pour off the water from the crystals, and dry them on bibulous paper. (LOND.)

R. Subcarbonate of Potass, one part.
Boiling Water, fifteen parts.
Supertartrate of Potass, three parts, or, a sufficient quantity.

To the subcarbonate dissolved in the water, add, by portions, the supertartrate in fine powder so long as it excites effervescence, which gradually ceases before thrice the weight of the subcarbonate is added. Filter the liquor when cold, and, after due evaporation, set it aside that crystals may form. (EDINB.)

R. Common Carbonate of Potass, five parts.
Bitartrate of Potass, fifteen parts.
Boiling Water, forty-five parts.

Dissolve the carbonate in the water, and gradually add the bitartrate, reduced to fine powder. Filter the liquor, evaporate, and set it aside that crystals may form as it cools. (DUBL.)

A similar formula is indicated by all the other pharmacopœias. — GALL.; FULD.; LIPP.; SARD.; HERBIP.; SPIELMANN; and VAN MONS direct the solution of subcarbonate to be added to that of the supertartrate.

2. In the Liquid state. *Liquor spleneticus ex Tartara.*

R. White Tartar, } of each, one pound.
Nitre, }

Detonate together in a crucible, and dissolve the residue in

Boiling Water, three pounds.
Strain, and add,

White Tartar, one pound.
Digest for eight days, and decant. (PALAT.; WIRTEM.; SPIELMANN)

It is excitant, diuretic, resolvent, and laxative.—Dose, from a scruple to a dram, as an alterative; and about an ounce, as a purgative.

It is incompatible with the sulphuric, nitric, and hydrochloric acids, lime-water, hydrochlorate of lime, sulphate of soda, and the vegetable acids.

KLEIN'S DIGESTIVE POWDER.

Pulvis leniens anti-hypochondriacus Kleinii, Solamen hypochondriacorum. (HAMB.; NIEMANN; SAINTE-MARIE)

R. Tartrate of Potass, } of each,
Rhubarb, } an equal part.
Orange Peel, }

Make a powder. (HAMB.; NIEMANN)

R. Tartrate of Potass, two drams.
Rhubarb, } of each, one dram.
Orange Peel, }
Cajeput Oil, one scruple.

Mix, and divide into twenty-four parcels. (SAINTE-MARIE)

It is highly extolled by KLEIN and BALDINGER; it is stimulant and sometimes produces a slight purgative effect.—Dose, two or three parcels every two days.

RESOLVENT POWDER.

Pulvis incidens. (SAUNDERS)

R. Tartrate of Potass, } of each,
Muriate of Ammonia, } one ounce.
Subcarbonate of Magnesia, }
Root of Arum, two drams.

Mix.

AQUEOUS SOLUTION OF TARTRATE OF POTASS.

Liquor Tartari tartarisati, Solutio Tartari solubilis. (BRUNS.; CAST. BORUS.; RATIER)

R. Tartrate of Potass, half an ounce.
Boiling Water, two pounds.

Dissolve. (RATIER)

CAST. BORUS. prescribes one ounce of tartrate and twelve ounces of water, and half an ounce of honey to be added.

It is laxative; to be taken by glassfuls.

The preparation of BRUNS., composed of one pound each of the salt and of water, is only intended to be kept in the shops for use.

STIMULANT FOMENTATION. (RATIER)

R. Tartrate of Potass, one ounce.
Water, one pound.

Dissolve.—To be applied to atonic and stationary ulcers.

GRANT'S PURGATIVE. (SAINTE-MARIE)

R. Tartrate of Potass, six or eight drams.
Infusion of Cichory, twenty ounces.
Manna, two ounces.

Dissolve.—To be taken every day or two days for a fortnight at four doses, in the

morning, drinking veal broth during the effect of the remedy. It has been extolled in hypochondria and chronic affections of the liver.

LAXATIVE DRINK. (CADET DE GASSICOURT)

R. Tartrate of Potass, half an ounce.
Infusion of Cichory Leaves, two pounds.

Dissolve and add

Honey, a sufficient quantity.

Mix.—It is mildly laxative; to be taken at several doses in the course of the day.

MUTZEL'S RESOLVENT MIXTURE.

Potio ad Atrabilem. (FULD.; NIEMANN; SAINTE-MARIE)

R. Tartrate of Potass, } of each,
Honey, } half an ounce.
Water, } eight ounces.

Dissolve.—One or two spoonfuls to be taken every hour in melancholy and mania. (FULD.; NIEMANN)

SAINTE-MARIE gives the following, after BALDINGER:

R. Tartrate of Potass, one ounce.
Honey, four ounces.

Mix, and add a few drops of essence of anise.

To be taken by tea spoonfuls, drinking afterwards a glassful of whey.

The formula which CADET DE GASSICOURT gives under this name varies still more from the original:

R. Tartrate of Potass, half an ounce.
Extract of Gentian, } of each,
Lesser Centaury, } two drams.
taury, }
Water, } eight ounces.

Dissolve, and mix.—Dose, half an ounce every hour, in obstructions of the liver.

It resembles the following, which is the same in its mode of action.

R. Tartrate of Potass, half an ounce.
Extract of Lesser Centaury, two drams.
Water, eight ounces.

Dissolve and mix.—To be taken in the same manner. (SWEDIAUR)

SALINE EMULSION. (SUEC.)

R. Tartrate of Potass, three drams.
Honey, one ounce and a half.
Spring Water, one pound.
Dissolve, and triturate the liquor, in a stone mortar with

Sweet Almonds blanched, one ounce.
Strain the emulsion.

It is a very mild laxative; to be taken by glassfuls.

PURGATIVE POTION FOR CHILDREN.
(HAMB.)

R. Tartrate of Potass, two drams.
Laxative Infusion, } of each,
Spring Water, } one ounce.
Glycyrrhized Honey, half an ounce.
Mix.

PURGATIVE DRINK. (BORIES)

R. Tartrate of Potass, four drams.
Jalap, in powder, eight grains.
Beer, two glassfuls.
Infuse for twenty-four hours; filter, and add
Sulphuric Æther, twenty drops.
Mix.—To be taken at twice, in the morn-
ing, at three hours interval.

ALCOHOLIC SOLUTION OF TARTRATE OF
POTASS. (PIDERIT)

R. Tartrate of Potass, four ounces.
Distilled Water, eight ounces.
Alcohol, two drams.
Dissolve, mix, and filter.

This liquor is intended to be put in po-
tions, and mixtures.—Dose, from one to
three ounces.

SALINE LIQUOR. (LIPP.)

R. Tartrate of Potass, one pound.
Hydrochlorate of Ammonia, two ounces.
Orange Peel, one ounce.
Distilled Water, three pounds.

Digest for four days at a moderate tem-
perature, shaking frequently; then filter.

BITTER ELIXIR.

Elixir amarum seu aperiens seu resolvens.
(BRUNS.; FULD.; HAMB.)

R. Tartrate of Potass, half an ounce.
Extract of Buckbean, one ounce.
Peppermint Water, eight ounces.
Mix. (HAMB.)
R. Tartrate of Potass, } of each,
Extract of Dandelion, } half an ounce.
Infusion of Rhubarb, } of each,
Peppermint Water, } three ounces.

Mix well. (HAMB.)

R. Tartrate of Potass, half an ounce.
Extract of Gentian, one ounce.
Distilled Water, six ounces.
VOL. II.

Peppermint Water, two ounces.
Mix. (FULD.)

R. Tartrate of Potass, two ounces.
Water, twelve ounces.
Extract of Buckbean, } of each,
Blessed Thistle, } one
Lesser Centaury, } ounce.
Cinnamon Water, two ounces.

Mix. (BRUNS.)

SAUNDERS gives the following similar
formula under the title of *Mixtura resolvens*
roboreans:

R. Tartrate of Potass, three drams.
Extract of Fumitory, } of each; one
Buckbean, } dram and a half.
Water of Blessed Thistle, six ounces.
Pepper- } of each,
mint, } one ounce and a half.
Syrup of Tolu, }

Mix.—It is excitant, and is considered
resolvent and aperitive.—To be taken by
spoonfuls.

RESOLVENT MIXTURE.

Mixtura deobstruens. (SWEDIAUR)

R. Tartrate of Potass, } of each,
Honey, } half an ounce.
Cinnamon Water, eight ounces.
Alcoholized Æther, one dram.

Dissolve, and mix.—Dose, at first half
an ounce every two hours, and afterwards
thrice a day.

SUPER-PROTOTARTRATE OF POTASSIUM.

*Hypertartras seu Supertartras seu Bitar-
tras Potassæ, Tartras superpotassicus seu
subpotassicus, Tartras acidulus Potassæ,
Supertartras kalicum.*—Supertartrate of
Potass.

I. *That which is met with in commerce.*

1. Impure. *Tartarus, Tartarus crudus seu
impurus.*—Crude Tartar, Essential Salt
of Tartar.

LOND.; DUBL.; EDINB.; ANT.; AUSTR.;
BATAV.; BAVAR.; BELG.; BRUNS.;
DAN.; MIL. COPEN.; HISP.; GALL.;
FENN.; FERR.; FULD.; FORM. HÔP.
FR.; GENEV.; HANN.; HASS.; LIPP.;
OLDEN.; LUSIT.; POL.; CAST. BORUS.;
BORUS.; ROSS.; SAX.; SUEC.; WIR-
TEM.; HERBIP.; BERGIUS; BRUGNA-
TELLI; COKE; MURRAY and GMELIN;
PARIS; PIDERIT; SPIELMANN.

The saline crust which is deposited on the
inner surface of the barrels in which wine
is kept, formed of an assemblage of nume-
rous crystalline spangles. It is white or
red according to the colour of the wine

which furnishes it. It consists of supertartrate of potass combined with a certain quantity of tartrate of lime and colouring matter.

2. Purified. *Tartari Crystalli, Cremor Tartari*.—Crystals of Tartar, Cream of Tartar.

These names are given to the following salt, which is presented either crystallized in quadrangular prisms, or in the form of powder. (LOND.; EDINB.; AMER.; ANT.; AUSTR.; BATAV.; BAVAR.; MIL. COPEN.; FORM. HÔP. FR.; LIPP.; SAX.; SUEC.; COXE)

II. Purification of the salt furnished by commerce.

BATAV.; BELG.; BRUNS.; HISP.; GALL.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; BORUS.; SARD.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS.

R. Tartar, any quantity.
Water, a sufficient quantity

to dissolve the salt. Strain the solution, clarify it with white of egg, remove the scum, strain through flannel, and crystallize. (GALL.; BRUGNATELLI)

BATAV.; BELG.; BRUNS.; HASS.; LUSIT.; SARD.; WIRTEM.; HERBIP.; PIDERIT; SPIELMANN; and VAN MONS give the same process, but do not require the solution to be clarified with the white of egg.

HANN.; OLDEN.; POL.; and BORUS. simply require the extraneous substances to be removed, the tartar to be washed with cold water, and then dried and pulverized.

R. Tartar, ten pounds.
Pure Argil, one pound.
Water, eighteen pounds.

Boil together for two hours, strain the liquor while hot, and set it aside in a cool place that crystals may form. Wash these with cold water, then dissolve them in water, and crystallize again. (HISP.; SWEDIAUR)

R. White Tartar, washed with warm water, and powdered, one part.
Water, twenty parts.

Boil till the tartar is dissolved and a thick pellicle formed, and remove it: continue thus till a pellicle ceases to be formed. (HERBIP.)

This process is the only one which really merits the name of cream of tartar.

It is refrigerant, diuretic, and purgative. —Dose, from one dram to two ounces in from eight ounces to two pounds of water.

RED HEPATIC POWDER.

Pulvis hepaticus ruber. (WIRTEM.; SPIELMANN)

R. Supertartrate of Potass, four ounces.
Red Sanders Wood, } of each,
Petals of Red Roses, } half an ounce.
Diluted Sulphuric Acid, two drams.

Bruse the petals, sprinkling them with the acid; then add the wood, and finally the salt.

It is cooling.—Dose, from two scruples to a dram.

TARTAREOUS MIXTURE.

Mixtura Kali tartarici. (AUGUSTIN)

R. Supertartrate of Potass, } of each,
Extract of White Hore- } two drams.
hound, }
Balm Water, three ounces.

Dissolve, and mix.—To be taken at twice, morning and evening, in piles with constipation.

ELECTUARY OF SUPERTARTRATE OF POTASS. (NIEMANN; SAINTE-MARIE)

R. Supertartrate of Potass, one part.
Rob of Prunes, two parts.

Mix.—Dose, half a spoonful every two hours.

TARTAREOUS OXYMEL. (SWEDIAUR)

R. Supertartrate of Potass, one part.
Despumated Honey, two parts.

Mix.—It is recommended in bilious fevers and in dropsies.

DENTIFRICE POWDER.

Pulvis dentrificus seu Tartari compositus.
(BRUNS.; DAN.; FENN.; FERR.; HANN.; OLDEN.; PALAT.; POL.; BORUS.; SARD.; SUEC.; WIRTEM.; CADET DE GASSICOURT; VAN MONS)

R. Supertartrate of Potass, four ounces.
Root of Florentine Iris, two ounces.
Myrrh, } of each,
Dragon's Blood, } one ounce.
Oil of Cloves, forty drops.

Pulverize and mix. (DAN.; HANN.; OLDEN.; BORUS.)

R. Supertartrate of Potass, } of each, an
Ratanhy Root, } equal part.
Root of Florentine Iris, a sufficient quantity
to give it an agreeable odour. Mix. (FERR.)

R. Ratanhy Root, six ounces.
Cuttle Fish Bone, ten ounces.

Supertartrate of Potass, } of each,
Root of Iris, } four drams.

Mix. (FERR.)

R. Supertartrate of Potass, four ounces.
Root of Iris, two ounces.
Mastic, }
Myrrh, } of each, one ounce.
Catechu, }
Oil of Cloves, forty drops.

Make a fine powder. (POL.)

R. Supertartrate of Potass, twenty-four parts.

Root of Iris, }
Myrrh, } of each, eight parts.
Kino, }
Cochineal, triturated with a few drops
of lemon juice, one part.

Make a powder, and add to each ounce
four drops of oil of cloves. (FENN.; SUEC.)

R. Supertartrate of Potass, thirty-six parts.

Cuttle Fish Bone, } of each,
Armenian Bole, } twenty-four
Prepared Red Coral, } parts.
Dragon's Blood, twelve parts.
Cinnamon, six parts.
Cochineal, three parts.
Cloves, one part.

Make a fine powder. (FULD.)

R. Cuttle Fish Bone, } of each,
Root of Iris, } one ounce.
Dragon's Blood, } of each, half
Supertartrate of Potass, } an ounce.
Mastic, two drams.
Cloves, one dram.

Make a powder. (SARD.)

R. Chalk, }
Lac, } of each, sixteen parts.

Supertartrate of Potass, forty-eight parts.

Cinnamon, }
Cloves, } of each, one part.

Mix. (VAN MONS)

R. Seed Lac, }
Cuttle Fish Bone, } of each, one ounce.
Root of Iris, }
Supertartrate of Potass, six drams.
Cloves, }
Myrrh, } of each, two drams.
Mastic, }

Mix. (BRUNS.; PALAT.; WIRTEM.)

R. Prepared Red Coral, } of each,
Cuttle Fish Bone, } six drams.
Armenian Bole, }
Supertartrate of Potass, nine drams.
Dragon's Blood, three drams.
Cinnamon, one dram and a half.
Cochineal, one ounce one scruple.
Cloves, twenty grains.

Mix. (CADET DE GASSICOURT)

DENTIFRICE ELECTUARY.

Electuarium dentifricium seu gingivale.
(BRUNS.; HANN.)

R. Supertartrate of Potass, } of each,
Pumice Stone, } six ounces.
Myrrh, }
Crab's Eyes, } of each, two ounces.
Root of Iris, }
Cochineal, six drams.
Honey of Roses, three pounds.
Oil of Cloves, one dram.

Pulverize very finely, and mix. (HANN.)

R. Supertartrate of Potass, one dram and a half.
Cochineal, one dram.
Saturated Tincture of Myrrh, dis-
tilled to one half, three ounces.
Honey of Roses, four ounces.
Tincture of Lac, one ounce.

Mix well together by trituration, digest
for twenty-four hours, and add to the
strained liquor while hot,

Oil of Cinnamon, seven drops.

Preserve. (BRUNS.)

BALDINGER'S OPHTHALMIC POWDER.

Pulvis ophthalmicus seu Tartari argillatus
seu ad Maculas Corneæ. (FULD.; HASS.;
ROSS.; SUEC.; HERBIP.; VAN MONS)

R. Supertartrate of Potass, } of each, half
White Sugar, } an ounce.
Gum Arabic, } two drams.

Make a very fine powder. (HASS.)

R. Supertartrate of Potass, } of each,
Armenian Bole, } an
White Sugar, } equal part.

Pulverize, and pass through a doubled
linen cloth. (FULD.; ROSS.; SUEC.;
HERBIP.; VAN MONS)

TROCHISCS OF SUPERTARTRATE OF POTASS.

Trochisci de Cremore Tartari. (WIRTEM.)

R. Supertartrate of Potass, one dram and a half.
White Sugar, six ounces.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs.—They are recommended
to abate thirst in fever.

TARTARIZED MANNA.

Manna tabulata seu tartarisata. (SPIEL-
MANN)

R. Supertartrate of Potass, half an ounce.
Manna, four ounces.
Water, ten ounces.

Boil to the proper consistence, and form tablets.

They form an agreeable laxative.

TRONCHIN'S LAXATIVE AND TEMPERANT POWDER.

Pulvis temperans seu refrigerans. (DAN.; MIL. COPEN.; HASS.; SAINTE-MARIE)

R. Supertartrate of Potass, three ounces.
Nitrate of Potass, three drams.
White Sugar, four ounces.

Mix. (SAINTE-MARIE)

HASS. prescribes one pound of supertartrate, and three ounces of nitrate, without sugar.—MIL. COPEN.—two ounces of supertartrate, and half an ounce each, of nitrate and of sugar.—DAN.—three ounces of supertartrate, and half an ounce each, of nitrate and of oleosaccharum of lemon.—Dose, one or two drams, in the morning, in a glass of whey.

REFRIGERANT ELECTUARY.

Electuarium lenitivum. (AUSTR.; SAINTE-MARIE)

R. Supertartrate of Potass, one ounce.
Nitrate of Potass, half an ounce.
White Sugar, ten drams.
Syrup of Lemon, a sufficient quantity.

Mix.—HERRENSCHWAND prescribed it in the dose of a tea spoonful every two hours in the morning, with a glass of whey after each dose. (SAINTE-MARIE)

R. Pulp of Prunes, three pounds.
Rob of Elder, one pound.
Supertartrate of Potass, half a pound.
Despumated Honey, a sufficient quantity.

Make an electuary. (AUSTR.)

ECCROFOTIC POWDER. (FULD.)

R. Supertartrate of Potass, one ounce.
Subcarbonate of Magnesia, } of each,
Sublimed Sulphur, } half an ounce.
Nitrate of Potass, two drams.

Mix.—It is incisive, diuretic, resolvent, and laxative.—Dose, from one to three drams.

ECPHRATIC POWDER. (CADET DE GASSI-COURT)

R. Supertartrate of Potass, }
Magnesia, } of each,
Sublimed Sulphur, } half
Rhubarb, } an ounce.
Chamomile Flowers,
Oleosaccharum of Fennel,

Mix.—It is recommended by SELLE in obstructions of the abdominal viscera.—

Dose, a spoonful in the course of the day, at several times.

SALINE POWDER.

Pulvis salinus seu Tartari compositus. (HERBIP.; WURCEB.)

R. Supertartrate of Potass, half an ounce.
Sulphate of Potass, two drams.

Mix. (HERBIP.)

R. Supertartrate of Potass, five drams.
Sulphate of Potass, three drams.
Tartarized Antimony, one grain.

Mix, and divide into eight parcels. (WURCEB.)

PURGATIVE ELECTUARY. (SAUNDERS)

R. Supertartrate of Potass, five drams.
Tartarized Soda, half an ounce.
Manna, six drams.
Pulp of Tamarinds, three ounces.
Syrup of Pale Roses, a sufficient quantity.

Make a thick electuary.

SOLUTION OF SUPERTARTRATE OF POTASS.

Solutio Tartari depurati, Decoctum Crystallorum Tartari. (HANN.)

R. Supertartrate of Potass, three drams.
Hot Distilled Water, thirty-two ounces.
Syrup of Raspberries, two ounces.

Dissolve, and mix.

It is laxative, diuretic, and cooling.—Dose, two pounds daily.

SAX.; and NIEMANN indicate two similar preparations: NIEMANN, under the title of *Cream of Tartar Drink*, prescribes half an ounce of the supertartrate, one pound of boiling water, and one ounce of syrup of raspberries:—SAX., under that of *Decoctum Tartari citratum*,—one part of supertartrate, twenty parts of water, to be reduced to fifteen by boiling with the salt, and one half part of lemon juice.

HYDRAGOGUE POTION. (ELLIS; SWEDIAUR)

R. Supertartrate of Potass, two drams.
Water, five ounces.

Dissolve.—To be taken, three, four, five, and six times a day.—It is recommended in ascites and anasarca. (SWEDIAUR)

ELLIS prescribes one ounce of supertartrate and two pounds of water.

MILMAN'S DRINK.

R. Supertartrate of Potass, half an ounce.
Decoction of Barley, two pounds.
Brandy, one or two ounces.

Dissolve, and mix.—This drink has been extolled in dropsies.

DIURETIC POTION. (SWEDIAUR ; NIEMANN)

R. Supertartrate of Potass, }
from half an ounce to one ounce and a half.
Decoction of Barley, }
a sufficient quantity.

Dissolve.—To be taken by spoonfuls in the course of the day. (SWEDIAUR)

R. Supertartrate of Potass, } of each,
Raisins, } two ounces.
Boiling Water, } four pounds.

Infuse for some hours, then boil a little, and strain.—To be taken by glassfuls.

IMPERIAL DRINK. (NIEMANN)

R. Supertartrate of Potass, half an ounce.
Orange Peel, three ounces.
Boiling Water, four pounds.

Infuse, and strain.

ACIDULOUS PURGATIVE MIXTURE. (SAINTE-MARIE)

R. Supertartrate of Potass, } of each,
White Sugar, } four drams.
Syrup of Gooseberries, three ounces.
Essence of Anise, one drop.

Mix.—It is recommended by STOLL in bilious fevers.—To be taken in the morning fasting, at four doses, with lemonade or veal broth.

FEBRIFUGE AND LAXATIVE POTION. (BORIES)

R. Supertartrate of Potass, half a pound.
Strong Infusion of Chamomile, one pound and a half.

Dissolve.—It is extolled in obstinate intermittent fevers.

PURGATIVE POTION. (BRERA ; PIERQUIN)

R. Supertartrate of Potass, one dram.
Syrup of Lemon, } of each,
Cichory, } one ounce.
Infusion of Chamomile, two ounces.

Mix. (PIERQUIN)

R. Pulp of Tamarinds, one ounce.
Water, eight ounces.
Supertartrate of Potass, one ounce.

Dissolve, and mix.—To be taken at twice. (BRERA)

R. Pulp of Tamarinds, one ounce.
Water, eight ounces.
Manna, } of each,
Supertartrate of Potass, } one ounce.

Dissolve, and mix.—To be taken at twice. (BRERA)

LAXATIVE MIXTURE. (SAINTE-MARIE)

R. Supertartrate of Potass, four drams.
Sulphate of Potass, two drams.
Lettuce Water, six ounces.
Syrup of Mulberries, two ounces.

Dissolve, and mix.—Dose, a table spoonful, three or four times a day.

COMPOUND TARTARIC POTION. (VAN MONS)

R. Supertartrate of Potass, one ounce.
White Sugar, two ounces.
Boiling Water, three pounds.

When cold, decant, and add,
Tartarized Antimony, half a grain.

LAXATIVE LIQUOR.

Liquor aperitivus seu salinus laxativus.
(SPIELMANN)

R. Sulphate of Magnesia, } of each,
Supertartrate of Potass, } one ounce.
Sulphate of Potass, half an ounce.
Boiling water, ten ounces.

Dissolve, and filter.

SYDENHAM'S DIURETIC WINE.

Vinum Genistæ Sydenhami. (BORIES ; NIEMANN ; PIERQUIN)

R. Ashes of Butcher's-broom, two ounces.
White Wine, one pound and a half.

Infuse for twenty-four hours, and filter. (BORIES ; PIERQUIN)

NIEMANN prescribes one pound of ashes, half a dram of wormwood leaves, and two pounds of Rhenish wine.

Dose, four ounces thrice a day ; in dropsies.

HYDRAGOGUE WINE. (SAUNDERS)

R. Rhenish Wine, one pound.
Subcarbonate of Potass, half an ounce.

After the cessation of the effervescence, add,
Syrup of Buckthorn, one ounce and a half.

Mix.—To be taken by glassfuls.

ALKALIZED WINE OF JUNIPER. (HERBIP.)

R. Subcarbonate of Potass, half an ounce.
Juniper, one ounce.
White Wine, two pounds and a half.

Infuse for twelve hours, and strain.

EXCITANT WINE. (HERBIP.)

R. Subcarbonate of Potass, } of each,
Generous Wine, } half an ounce.

Dissolve, and add,

Oxymel of Colchicum, two ounces.
Sulphuric Aether, one scruple.

Mix.

DIURETIC WINE.

Vin diuretique Anglais. (CADET DE GASSICOURT)

R. Subcarbonate of Potass, one dram and a half.
Squill, }
Rhubarb, } of each, one dram.
Juniper, }
Root of Zedoary, two drams.
Cinnamon, three drams.
White Wine, one pound and a half.

Macerate, and filter.

Dose, three or four glassfuls daily, in dropsy.

BORATED SUPER-PROTO-TARTRATE OF POTASSIUM.

Cremor Tartari solubilis.—Borated Tartar. (BAVAR. ; BELG. ; GALL. ; FULD. ; HANN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; VAN MONS)

R. Sub-borate of Soda, one part.
Water, ten parts.

Boil till dissolved, and add,

Supertartrate of Potass, three parts ;
or, as much as the liquor will dissolve. Set aside, filter, and evaporate to dryness. (BAVAR. ; BRUNS. ; HANN. ; BORUS. ; SAX. ; SUEC. ; WIRTEM. ; VAN MONS)

FULD. prescribes four parts of supertartrate and one part of sub-borate.

R. Supertartrate of Potass, three parts.
Sub-borate of Soda, one part.

Pulverize and mix. (POL.)

ROSS. prescribes one part of sub-borate and four parts of supertartrate. — VAN MONS—nine of sub-borate and nine and a half of supertartrate.

R. Supertartrate of Potass, two hundred and ten parts.
Boracic Acid, thirty parts.
Distilled Water, twenty-parts.

Dissolve the acid in the water with heat, and while the solution is hot, add the supertartrate gradually, constantly stirring. Evaporate to dryness, and pulverize. (GALL.)

VAN MONS recommends eight parts of super-tartrate and one part of boracic acid to be separately powdered, and then mixed together.

SOUBEIRAN directs four parts of supertartrate and one part of the acid to be dissolved over the fire in twenty-four parts of water ; the solution to be filtered, evapo-

rated to dryness with a gentle heat, and the residuum dried and powdered. (HENRI)

VAN MONS has proposed another method of increasing the solubility of the supertartrate of potass ; it differs from all the preceding :

R. Supertartrate of Potass, one part.
Water, eighteen parts.

Dissolve by boiling in a covered vessel ; add tartaric acid gradually, till a few drops of the liquor being placed upon a cold body no longer deposit supertartrate of potass ; then filter, and evaporate to dryness.

LAXATIVE LEMONADE. (PIERQUIN)

R. Supertartrate of Potass, one ounce.
Sub-borate of Soda, three drams.
White Sugar, two ounces.

Mix.—To be taken in three glassfuls of water, at intervals of half an hour.

TARTRO-BORATED DRINK.

Eau saline fondant de Lassone. (FORM. HÔP. FR. ; HAMB. ; BORIES)

R. Supertartrate of Potass, one ounce.
Sub-borate of Soda, three drams.
Boiling Water, one pound.

Dissolve. (HANN.)

BORIES prescribes one dram each, of the salts, and one pound of water.

R. Supertartrate of Potass, twenty parts.
Sub-borate of Soda, five parts.
Despumated Honey, sixty parts.
Water, a sufficient quantity

to make a pound and a half of liquor.

Triturate the two salts together, add the water by small portions, and then mix in the honey. (FORM. HÔP. FR.)

It is mildly laxative, taken by glassfuls, and might advantageously replace the decoction of tamarinds.—The bottle should be shaken each time the liquor is used.

DIURETIC POTION. (AUGUSTIN ; NIEMANN)

R. Borated Tartar, from four to six drams.
Juniper Water, } of each, three ounces.
Parsley Water, }
Nitric Aether, two drams.
Syrup of Poppies, one ounce.

Dissolve and mix.—Dose, two spoonfuls every two hours. (NIEMANN)

R. Borated Tartar, half an ounce.
Rob of Dwarf Elder, two ounces.
Juniper Water, three ounces.
Syrup of Buckthorn, } of each,
Violets, } half an ounce.

Dissolve and mix.—To be taken by spoonfuls. (AUGUSTIN)

HYDRAGOGUE PILLS.

Pilulæ hydragoguæ seu e Tartaro boraxato.
(AUGUSTIN)

- R. Extract of Buckbean, two ounces.
Borated Tartar, one dram and a half.
Root of Seneka, two drams.
—— Colchicum,
one dram and a half.

Make pills.—Dose, from twelve to twenty-four grains, every two hours.

PROTO-TARTRATE OF POTASSIUM AND OF SODIUM.

Alkali minerale tartarizatum, Sal polychrestus Seignetti seu Rupellensis, Soda tartarizata, Natrum tartarizatum, Tartarus natronatus, Tartras kalico-natricum seu Potassæ et Sodæ seu Potassæ sodicus seu Sodæ et Kali.—Tartrate of Potass and of Soda, Tartarized Soda, Soluble Sal polychrest, Rochelle Salt. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; GENEV.; HANN.; HASS.; OLDEN.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; PIDERIT; SPIELMANN; VAN MONS)

Soda Tartarizata. (LOND.)

- R. Subcarbonate of Soda,
twenty ounces.
Supertartrate of Potass,
two pounds.
Boiling Water, ten pints.

Dissolve the subcarbonate in the water, and gradually add the supertartrate. Filter the liquor, boil till a pellicle forms on its surface, and set aside to crystallize. Pour off the water, and dry the crystals in bibulous paper.

Tartras Sodæ et Potassæ. (EDINB.)

- R. Subcarbonate of Soda, one part.
Boiling Water, fifteen parts.
Supertartrate of Potass,
three parts; or,
a sufficient quantity.

To the subcarbonate dissolved in the water, gradually add the supertartrate, in fine powder, so long as effervescence is excited, which usually ceases before thrice the weight of the

subcarbonate is used. When the liquor is cold, filter, evaporate, and set it apart to crystallize.

Sodæ et Potassæ Tartras. (DUBL.)

- R. Carbonate of Soda, five parts.
Bitartrate of Potass, seven parts.
Boiling Water, fifty parts.

Dissolve the carbonate in the water, and gradually add the bitartrate in very fine powder. Filter, evaporate, and let cool that crystals may form.

- R. Subcarbonate of Soda, any quantity.
Boiling Water, a sufficient quantity to dissolve the salt. Add to the liquor,
Supertartrate of Potass,

a sufficient quantity;

or, as much as it will dissolve. Evaporate the liquor to 35 degrees of the areometer, and set it aside to crystallize. (AMER.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; FENN.; FERR.; GENEV.; HANN.; HASS.; OLDEN.; POL.; BORUS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; PIDERIT; VAN MONS)

GALL.; SPIELMANN; and VAN MONS direct the subcarbonate to be added to the solution of the supertartrate.

HENRI observes that the mother liquor is found to be almost entirely formed of tartrate of soda, and that it is sufficient to obtain more Rochelle salt, for a certain quantity of tartrate of potass to be added to it.

- R. Tartrate of Potass,
eleven parts and a half.
Dried Subcarbonate of Soda,
two parts and a half.
Water, twenty-four parts.
Set aside, decant, and crystallize. (VAN MONS)

- R. Supertartrate of Potass, two pounds.
Water, a sufficient quantity to dissolve the salt. Saturate this, while boiling, with potass, and add
Hydrochlorate of Soda,
five ounces two drams.

When the solution is cold, and the hydrochlorate of potass is deposited, evaporate the liquor that it may crystallize. (VAN MONS)

- R. Tartrate of Potass,
eleven parts and a half.
Sulphate of Soda,
three parts and four fifths.
Warm Water, twenty-four parts.

Heat together, add a little caustic soda to the liquor, filter, and crystallize. (VAN MONS)

It is excitant, diuretic, and purgative.—Dose, from two drams to one ounce in a pound and a half of water.

FORDYCE'S POWDER. (SAINTE-MARIE)

R. Tartarized Soda, ten grains.
Rhubarb, from three to seven grains.

Mix.—It is recommended in tabes mesenterica.—It is to be given every morning for a fortnight.

RESOLVENT MIXTURE. (SAINTE-MARIE)

R. Tartarized Soda, half an ounce.
Sulphate of Potass, two drams.
Infusion of Cichory, one pound.
Oxymel of Squill, four ounces.

Dissolve, and mix.—BOERHAAVE recommended it in abdominal obstructions.—Dose, four table-spoonfuls, in the morning, at intervals of an hour.

VAN SWIETER'S WHEY. (SAINTE-MARIE)

R. Root and Leaves of
Dandelion, } of each,
Herb of Fumitory, one
Leaves of Water Cress, handful.
Chervil,

Chop the herbs and boil them for five or six minutes with

Clarified Whey, one pound.

Strain with strong expression, and add
Tartarized Soda, one or two drams.
Honey, six drams.

Mix.—To be taken every morning, fasting, for a month, in four doses at intervals of an hour. It is recommended in abdominal engorgements arising from intermittent fevers, or inveterate hypochondriacal affections.

PROTO-TARTRATE OF POTASSIUM AND OF AMMONIA.

Ammonium tartaricum, Tartarus ammoniacus seu ammoniacalis seu solubilis ammoniacalis, Sal ammoniacum tartareum, Alkali volatile tartaricatum.

I. In the solid state.

DAN.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; WIRTEM.; NIEMANN.; VAN MONS.

R. Supertartrate of Potass, any quantity.
Boiling Water, a sufficient quantity
to dissolve the salt; add gradually to the liquor

Solution of Subcarbonate of
Ammonia, a sufficient quantity

to saturate the excess of acid. Filter the liquor, and evaporate it to dryness over a slow fire.

HASS.; LIPP.; and SAX. prescribe the caustic ammonia.—HANN.; and VAN MONS direct it to be crystallized, and the crystals reduced to powder.—NIEMANN recommends, before crystallizing it, a little solution of ammonia to be added to re-dissolve the crystals of tartar which may have been precipitated during the evaporation.

II. In the liquid state.

Liquor Tartari solubilis.

LIPP.

R. Solution of Subcarbonate of Ammonia, six ounces.
Supertartrate of Potass, a sufficient quantity

to saturate it. Set aside for a few days, and filter.

It is excitant, diuretic, and slightly laxative, and has been employed, though rarely, in active fevers, dropsies, and engorgements of the abdominal viscera—Dose, one dram, several times a day.

PROTO-OLEO-MARGARATE OF POTASSIUM.

We propose to place in this article all the soaps properly so called, even those which have soda for their base, because, however they may differ in their composition, they are all precisely the same in their mode of action.

I. SOAPS OF COMMERCE.

1. *Sapo Alicantinus seu Aloniensis seu Hispanicus albus, Sapo albus.*—Spanish Soap.

LOND.; EDINB.; DUBL.; AMER.; AMST.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; BORUS.; GALL.; FORM. HÔP. FR.; GENEV.; HAMB.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; SAX.; WIRTEM.; WURCEB.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; SAUNDERS; SPIELMANN.

It is prepared with olive oil and soda. It is white, hard, and homogeneous.

2. *Sapo Venetus.*—Venice Soap.

BELG.; BRUNS.; FENN.; FERR.; PALAT.; SUEC.; PIDERIT.

It is prepared as the preceding. It is hard and marbled.

3. *Sapo domesticus seu nostras.*

AUSTR.; BAVAR.; MIL. COPEN.; GALL.; FERR.; FORM. HÔP. FR.; HANN.; HASS.; POL.; CAST. BORUS.; BORUS.;

ROSS.; SAX.; WURCEE.; MURRAY and GMELIN; PARIS.

It is prepared in the same manner as the preceding; it presents blue marblings, from the hydro-sulphate of iron and alumine combined with the soda.

4. *Sapo mollis seu niger seu viridis seu kalicus*.—Soft Soap.

LOND.; EDINB.; MIL. COPEN.; HASS.; SUEC.; COXE; PARIS; PIDERIT.

Prepared with Oil and Potass.

5. *Sapo Wologdensis*.—Wologd Soap.

FENN.; ROSS.

Prepared with Suet and Potass, at Wologd and other places.

II. PHARMACEUTIC SOAPS.

A.

Sapo medicinalis seu medicatus seu amygdalinus.—Medicinal Soap, Almond Soap.

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; CADET DE GASSICOURT; SWEDIAUR; VAN MONS.

R. Subcarbonate of Soda, } of each,
Quick Lime, } four pounds.
Water, } twenty pounds.

Boil together, strain through linen, and evaporate the liquor till eight ounces of it fill a vessel capable of containing six ounces of water. Add to it when cold

Olive Oil, twice the quantity.

Boil, constantly stirring till a thick homogeneous mass is formed. Pour it into moulds, and dry it. (ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; HISP.; FERR.; FENN.; FORM. HÔP. FR.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; HERBIP.; BRUGNATELLI; SWEDIAUR)

GALL. prescribes one hundred parts of caustic ley (36 degrees), and two hundred and ten parts of oil of almonds.—VAN MONS—one part of ley, and two parts and a fourth of oil of almonds.—CADET DE GASSICOURT—two pounds of ley, and four pounds three ounces of oil of almonds.

BRUNS.; SARD.; WIRTEM.; HERBIP.; and VAN MONS also give a similar formula, but require a potass ley instead of one of soda.

VOL. II.

B.

Sapo cacaotinus.—Cocoa Soap.

BATAV.; FERR.; HANN.; LIPP.; SPIELMANN; VAN MONS.

R. Soda Ley, one part.
Melted Cocoa Nut Oil, two parts.
Boil together till incorporated.

BEEF MARROW SOAP.

Sapo Wologdensis. (GALL.; CADET DE GASSICOURT; SWEDIAUR; VAN MONS)

R. Prepared Beef Marrow, five hundred parts.

Solution of Potass, two hundred and fifty parts.

Melt over a slow fire, constantly stirring with a spatula, till the mixture has become totally soluble in water. Then dissolve it in

Boiling Water, two thousand parts.

Add to the liquor

Muriate of Soda, one hundred and eighty parts

dissolved in

Distilled Water, one thousand parts.

Set aside to cool, express through linen, and evaporate. (GALL.; CADET DE GASSICOURT)

SWEDIAUR prescribes two parts of solution of potass, and four and a half of beef marrow.—VAN MONS—one and a half of caustic soda, and two of marrow.

All these soaps are more or less stimulant. They have been extolled as resolvent, diuretic, and lithontriptic. They are administered with advantage in cases of poisoning by acids. Externally they are resolvent or maturative.—Dose, from ten grains to one dram twice or thrice a day.

AROMATIC ANIMAL SOAP. (CADET DE GASSICOURT; VAN MONS)

R. Expressed Oil of Nutmeg, one part.
Beef Marrow, five parts.

Melt together, and incorporate

Solution of Caustic Soda, seven parts, adding it gradually, and constantly stirring till perfectly incorporated. (VAN MONS)

CADET DE GASSICOURT prescribes seven parts of marrow, one part of oil of nutmeg, one of spermaceti, and a sufficient quantity of solution of caustic soda.

CERATE FOR EXAMINATION DURING LABOUR.

Cerat pour le Toucher. (HENRI)

R. Olive Oil, six thousand parts.

3 N

Spermaceti, } of each,
White Wax, } four hundred parts.

Melt together, strain through linen, and add gradually, constantly stirring,

Caustic Soda, }
four hundred and eighty parts.

This cerate should not be employed till after it has been prepared for some time.

COSMETIC POWDER. (NIEMANN)

R. Spanish Soap, twelve ounces.
Subcarbonate of Potass, two ounces.
Root of Iris, three ounces.
Horse-chesnuds, two pounds.
Oil of Lavender, } of each,
— Bergamotte, } forty drops.
— Lemon, thirty drops.
— Cloves, ten drops.
Sugar, half an ounce.

Pulverize and mix.

R. Spanish Soap, twelve ounces.
Subcarbonate of Soda, one ounce and a half.
Root of Iris, one ounce.
Lesser Cardamoms, one dram.
Oil of Lavender, } of each,
— Bergamotte, } thirty drops.
Lemon, twenty drops.
Cinnamon, three drops.
Sugar, half an ounce.

Make a powder.

SAPONACEOUS BOLUSSES.

Boli saponacei. (SAUNDERS; SWEDIAUR)

R. Medicinal Soap, two scruples.
Oil of Caraway, two or three drops.
Syrup of Tolu, a sufficient quantity.

Mix.—Four grains of aloes or seven of rhubarb are sometimes added. They are purgative.

LITHONTRIPTIC BOLUSSES.

Boli lithontriptici. (SPIELMANN)

R. Spanish Soap, eight ounces.
Calcined Egg shells, } of each,
Pulp of Hips, } one ounce.
Subcarbonate of Potass, one dram.
Proof Spirit, a sufficient quantity.

Make sixty boluses.

SAPONACEOUS PILLS.

Pilule saponaceæ seu resolventes. (GALL.; FORM. HÔP. FR.; HAMB.; HUFELAND.; NIEMANN; SAINTE-MARIE; SPIELMANN; SWEDIAUR; VAN MONS)

R. Medicinal Soap, half a pound.
Root of Marshmallows, one ounce.
Nitrate of Potass, two drams.

Oil of Almonds, a sufficient quantity.
Make a pilular mass. (GALL.; FORM. HÔP. FR.)

As the oil soon becomes rancid, HENRI proposes to substitute simple syrup for it.

R. Medicinal Soap, two drams.
Rhubarb, } of each, one dram.
Myrrh, }

Make pills of four grains each. (SWEDIAUR)

R. White Soap, one ounce.
Extract of Juniper, } of each,
Powdered Gentian, } four drams.

Make pills of four grains each. (SAINTE-MARIE)

R. Venice Soap, } of each,
Galbanum, } two drams.
Extract of Celandine, }

Liquorice Powder, a sufficient quantity.

Make pills of two grains each. (HAMB.)

R. Medicinal Soap, one dram and a half.
Gum Ammoniac, one dram.
Extract of Dog-grass, four scruples.

Make pills of one grain each. (NIEMANN)

HUFELAND; and NIEMANN prescribe an equal part each, of soap, of gum ammoniac, and of extract of celandine; to be made into pills of two grains each.

R. Official Soap, one ounce.
Gum Ammoniac, } of each,
Extract of Lesser } half an ounce.
Centaurry, }

Oil of Amber, ten drops.

Make a pilular mass. (SPIELMANN)

R. Acetate of Potass, } of each, one part.
Gum Ammoniac, }
Pills of Myrrh and }
of Aloes, }
White Soap, two parts.
Simple Syrup, a sufficient quantity

to make a mass. (VAN MONS)

LAXATIVE SUPPOSITORY.

Suppositorium ad Alvum sollicitandum. (PALAT.; NIEMANN; SPIELMANN)

R. Soap, two ounces.
Muriate of Soda, one ounce.
Inspissated Honey,

a sufficient quantity.

Make cones of various sizes, to be rubbed with oil of wormwood before using them.

MATURATIVE CATAPLASM.

Cataplasma saponaceum seu resolvens. (FULD.; HENRI; NIEMANN; RATIER; SPIELMANN)

R. Crumb of Bread, eight ounces.
Venice Soap, one ounce.

Cow's Milk, one pound.
Boil together. (FULD.; HENRI; NIEMANN)

R. Barley Meal, eight ounces.
White Soap, four ounces.
Water, a sufficient quantity.

Make a cataplasm. (RATIER)

SPIELMANN prescribes sixteen ounces of bread poultice and two ounces of scraped soap.

SAPONACEOUS HONEY. (NIEMANN)

R. Soap, } of each, four ounces.
Honey, }
Subcarbonate of Potass, half an ounce.

Fumitory Water, two drams.

Mix.—It is recommended as resolvent to indolent tumours.

SAPONACEOUS LOHOCH.

Linctus saponatus seu saponaceus; Lohoch saponaceum seu saponatum. (FULD.; LIPP.; WIRTEM.; HERBIP.; CADET DE GASSICOURT; SAUNDERS; SPIELMANN; SWEDIAUR)

R. Soap, one part.
Oil of Almonds, eight parts.
Syrup of Manna, twelve parts.

Triturate together. (LIPP.)

HERBIP. prescribes three parts of soap, sixteen of oil, and twenty-four of syrup of manna.

R. Soap, one part.
Oil of Almonds, eight parts.
Syrup of Maidenhair, twelve parts.

Triturate together. (WIRTEM.; SPIELMANN)

SAUNDERS prescribes one part of soap, sixteen parts of oil, and twenty-four parts of syrup of maidenhair.

R. Soap, one part.
Oil of Almonds, twenty-four parts.
Syrup of Rhubarb, thirty-six parts.

Triturate together. (FULD.)

R. Soap, one part.
Oil of Almonds, eight parts.
Syrup of Lemon, one part and a half.

Triturate together. (CADET DE GASSICOURT)

R. Soap, four parts.
Oil of Almonds, eighteen parts.
Syrup of Poppy, forty-eight parts.

Triturate together. (SWEDIAUR)

SAPONACEOUS OINTMENT. (RATIER)

R. White Soap, one pound.
Water, one pound and a half.

Soften the soap in the water, and pass it through a sieve. It is employed against the itch.

AROMATIC SOAP. (NIEMANN)

R. Concentrated Aqueous Solution
of Soap, six ounces.
Oil of Bergamotte, } of each,
Lavender, } two drams.
Rosemary, }

Mix.—It is employed in baths, as nervine and tonic.

SAPONACEOUS LAVEMENT.

Clysmma resolvens. (BRERA; SPIELMANN)

R. Leaves of Mallow, one handful.
Water, sixteen ounces.

Boil down to twelve ounces, and dissolve in the strained liquor.

Venice Soap, two drams.

Mix. (SPIELMANN)

R. Decoction of Barley, six ounces.
Venice Soap, one dram.
Honey, six drams.

Dissolve. (BRERA)

LAXATIVE AND ANTI-SPASMODIC FOMENTATION. (BORIES)

R. Flowers of Chamomile, } of each,
Melilot, } one handful.
Bay Berries, } of each,
Caraway Seeds, } one ounce.
Milk, a sufficient quantity,

to obtain three pounds of strained decoction after a quarter of an hour's boiling. Add

Venice Soap, one ounce and a half.
Saffron, two drams.

Mix.

SAPONACEOUS POTION. (CADET DE GASSICOURT; NIEMANN; RATIER)

R. Medicinal Soap, three drams.
Water, four ounces.
Syrup of Marshmallows, one ounce.
Magnesia, one dram.

Dissolve, and mix.—It is recommended against poisoning by mineral acids. (CADET DE GASSICOURT)

R. Soap, six drams.
Infusion of Balm, four ounces.
Syrup of Maidenhair, one ounce.

Dissolve.—To be taken at three or four doses in some cases of abdominal flatulency. (RATIER)

R. Soap, two drams.
Powdered Pearls, one dram.
Crab's Eyes, one dram and a half.

Syrup of Marshmallows, half an ounce.
 Mint Water, } of each,
 Water of Lemon Peel, } two ounces.
 Fennel, }

Dissolve, and mix.—To be taken by spoonfuls.—BOERHAAVE employed it against acidity of the primæ viæ of children. (NIEMANN)

ALCOHOLIC SOLUTION OF SOAP.

Alcohol saponatum, Spiritus saponatus, Tinctura Saponis. (BAVAR.; FERR.; FORM. HÔP. FR.; CAST. BORUS.; RATIER)

R. Soap, one part.
 Alcohol (0.900) four parts.
 Dissolve, and filter. (BAVAR.)

R. Spanish Soap, one pound.
 Alcohol, three pounds.
 Distilled Water, one pound.
 Dissolve, and filter. (CAST. BORUS.)

FORM. HÔP. FR. prescribes three parts of alcohol (23 degrees), and one part of soap.—RATIER—one pound of alcohol (26 degrees), and four ounces of soap; or, one pound of alcohol and two ounces of soap; or, one pound of alcohol and half an ounce of soap; or lastly, two pounds of alcohol, two pounds of water, and ten ounces of soap.

R. Spanish Soap, eight ounces.
 Subcarbonate of Potass, four ounces.

Dissolve the salt in sixteen ounces of water, add the soap, and boil to the consistence of honey; then pour upon the mass

Rectified Alcohol, sixteen ounces.
 Digest for a week, and filter. (FULD.)

This solution, the concentration of which should of course be varied according to circumstances, furnishes a resolvent which may be employed with success in cases of sprains when the inflammation has subsided. LUGOL has also applied it with advantage in the treatment of itch.

ANTI-PHLOGISTIC FOMENTATION. (HASS.; PIDERIT)

R. Venice Soap, three ounces.
 Lime Water, two pounds.
 Add to the solution
 Alcohol, five ounces.
 RHAZES White Trochiscs, half an ounce.

Mix.—It is resolvent.

AROMATIC ALCOHOLIC SOLUTION OF SOAP.

Spiritus Saponis seu saponatus seu Saponaceus seu resolvens Tinctura Saponis. (ANT.; AUSTR.; HANN.; LIPP.; OL-

DEN.; BORUS.; ROSS.; SAX.; NIEMANN; RATIER; SPIELMANN; VAN MONS)

R. White Soap, one part.
 Spirit of Lavender, eight parts.

Dissolve, and filter. (VAN MONS)

R. White Soap, one part.
 Alcohol, three parts.
 Rose Water, one half part.

Dissolve, and filter. (HANN.; OLDEN.; BORUS.; SAX.; NIEMANN)

POL. prescribes three parts and a half of alcohol.

R. White Soap, six ounces.
 Subcarbonate of Potass, one dram.
 Alcohol (0.910), one pound and a half.
 Lavender Water, half a pound.

Filter the solution. (AUSTR.)

SPIELMANN directs four ounces of Venice soap to be triturated with two of subcarbonate of potass, and the mass to be digested in a mixture of twelve ounces of alcohol and half a dram of essence of bergamotte.—ANT.—four ounces of soap to be dissolved in twenty-four of proof spirit, and two ounces each, of spirit of lavender and of spirit of rosemary, to be added to the solution.

R. Spanish Soap, four ounces.
 Subcarbonate of Potass, two ounces.

Boil in one pound of water to a thickish consistence. Pour upon this

Spirit of Lavender, sixteen ounces.
 Digest over a slow fire for four days, and filter. (ROSS.)

BATAV. prescribes eight ounces of soap, one pound of subcarbonate, and sixteen ounces of spirit of rosemary.

R. Olive Oil, two ounces.
 Soap, two drams.
 Vulnerary Spirit, one pound.

Dissolve, and filter. (RATIER)

It is resolvent, and is applied to indolent chronic tumours, rheumatic pains, and contractions of the joints.

EMOLLIENT AND ANTI-SEPTIC BROTH.

R. Leaves of Rue, } of each,
 Scordium, } two ounces.
 Mallow, }
 Water, a sufficient quantity,

to obtain two pounds of decoction. Add to it when strained,

Venice Soap, one dram,
 dissolved in

Alcohol, six drams.
 Mix.

CAMPHORATED ALCOHOLIC SOLUTION OF SOAP.

Linimentum Saponis seu saponaceum seu

Saponis compositum, Alcohol Camphora saponaceum, Aqua Vitæ saponacea. (HISP.; LUSIT.; NIEMANN; SPIELMANN; VAN MONS)

R. White Soap, three ounces.
Alcohol, sixteen ounces.

Dissolve and add
Camphor, one ounce.

Filter. (LUSIT.)

HISP. prescribes two ounces of soap, six drams of camphor, and one pound of alcohol.—SPIELMANN—one ounce of soap, and sixteen ounces of spirit of camphor.—NIEMANN—an arbitrary quantity of soap, one ounce of hot water, and one ounce of spirit of camphor.

R. Spirit of Camphor, sixteen parts.
White Soap, one part.

Macerate till dissolved and filter. (VAN MONS)

KEMPF'S SAPONACEOUS LINIMENT. (CADET DE GASSICOURT; PIDERIT)

R. Venice Soap, one ounce.
Alcohol, four ounces.

Set light to the alcohol, and stir it with a rod till it ceases to burn. Add to the liquor when cold,

Camphor, two scruples.
Filter.—It is resolvent.

AROMATIC AND CAMPHORATED ALCOHOLIC SOLUTION OF SOAP.

Balsamum saponaceum seu Saponis, Linimentum Saponis seu saponaceum seu Saponis compositum seu Saponis camphoratum. (LOND.; DUBL.; EDINB.; AMER.; ANT.; DAN.; FENN.; FULD.; SUEC.; WIRTEM.; COXE; SPIELMANN; SWEDIAUR; VAN MONS)

Linimentum Saponis compositum.
(LOND.; DUBL.)

R. Hard Soap, three ounces.
Camphor, one ounce.
Spirit of Rosemary, one pint.

Dissolve the camphor in the spirit; then add the soap, and macerate in the heat of a sand-bath till perfectly dissolved.

Tinctura Saponis camphorata, vulgo Linimentum Saponaceum. (EDINB.)

R. Hard Soap, four ounces.
Camphor, two ounces.
Oil of Rosemary, half an ounce.

Alcohol, two pounds.

Digest the soap in the alcohol for three days, then add the camphor and the oil, frequently shaking the mixture.

WIRTEM. prescribes six ounces of soap, two pounds of spirit of rosemary, and half an ounce of camphor.—ANT.—three ounces of soap, one pound of spirit of rosemary, and one ounce of camphor.—SWEDIAUR; and VAN MONS give the same formula as LOND.—AMER.; and COXE prescribe twelve ounces of soap, two ounces of camphor, two ounces of oil of rosemary, and eight pints of alcohol.—FULD.—five ounces of soap, two ounces of camphor, half an ounce of oil of rosemary, and two pounds of alcohol.

R. Hard Soap, two ounces.
Spirit of Spruce Fir Buds, one pound.

Dissolve, and add
Camphor, half an ounce,

and, if required,
Oil of Rosemary, one dram.
Filter. (SUEC.)

R. White Soap, two ounces.
Compound Spirit of Milfoil, one pound.

Dissolve, and add
Camphor, half an ounce.
Preserve. (FENN.)

R. Spanish Soap, ten ounces.
Alcohol, four pounds.

Dissolve at a moderate heat, and add
Camphor, two ounces.
Oil of Rosemary, } of each,
Thyme, } half an ounce.

Filter the solution. (DAN; SPIELMANN)
It is resolvent.

CAMPHORATED OPIATE ALCOHOLIC SOLUTION OF SOAP.

Balsamum anodynum, Linimentum Saponis cum Opio seu anodynum seu opiatum. (EDINB.; AMER.; GALL.; LUSIT.; SUEC.; BORIES; CADET DE GASSICOURT)

Tinctura Saponis et Opii, vulgo Linimentum anodynum. (EDINB.)

R. Hard Soap, four ounces.
Opium, one ounce.
Camphor, two ounces.
Oil of Rosemary, half an ounce.
Alcohol, two pounds.

Digest the soap and opium in the

alcohol for three days; then strain the liquor, and add the camphor and oil, frequently shaking the mixture.

R. White Soap, three ounces.
Opium, six drams.
Alcohol, sixteen ounces.

Dissolve, and add

Camphor, one ounce.

Preserve. (LUSIT.)

BORIES; and CADET DE GASSICOURT prescribe one pound and a half of alcohol, five ounces of soap, one ounce of opium, half an ounce of oil of rosemary, and three drams of camphor.—AMER. directs one ounce of opium to be added to the aromatic and camphorated alcoholic solution of soap.

R. White Soap, two ounces.
Opium, half an ounce.
Spirit of Spruce Fir Buds, one pound.

Dissolve, and add

Camphor, half an ounce.

It is employed externally, in cases of chronic pain, and internally, in what is called nervous colic.—Dose, from fifteen to twenty drops in an infusion of balm.

ANODYNE BALSAM. (GALL.)

R. Almond Soap, half an ounce.
Tincture of Opium, one ounce.
Triturate the solution with
Oil of Almonds, two ounces.

ÆTHEREAL SOLUTION OF SOAP.

Linimentum æthericum; Liniment acétique, Liniment de Savon éthereux, Savon d'Ether acétique, Savon acétique éthere. (BORIES; CADET DE GASSICOURT; HENRI; PIERQUIN; VAN MONS)

R. Marrow Soap, one part.
Acetic Æther, five parts.

Dissolve in a covered vessel placed in a warm bath, and pour off the liquor. (VAN MONS)

VAN MONS also prescribes one part and a half of soap and one of acetic æther; and also, three of soap and two of æther.—CADET DE GASSICOURT; and PIERQUIN—one of soap and eight of æther.—HENRI—three of soap and sixteen of æther.

It is employed in frictions, in chronic rheumatism.

CAMPHORATED ACETIC BALSAM. (BORIES; CADET DE GASSICOURT; HENRI; PIERQUIN)

R. Animal Soap, } of each, one dram.
Camphor, }
Essence of Thyme, ten drops.

Dissolve with the heat of a water-bath, and add

Acetic Æther, one ounce.

Filter. (BORIES; CADET DE GASSICOURT; PIERQUIN)

HENRI directs one dram each, of soap and of camphor, to be dissolved in two ounces of acetic æther, and ten drops of spirit of turpentine to be added to the solution.

It is recommended as a liniment in rheumatism and sciatica.

ANTI-ARTHRITIC BALSAM. (CADET DE GASSICOURT)

R. Spirit of Lavender, four ounces.
Aromatic Animal Soap, } of each,
Acetic Æther, } one ounce.
Camphor, two drams.
Oil of Peppermint, }
Cinnamon, } of each,
Lavender, } fifteen drops.
Nutmeg, }
Cloves, }
Sassafras, }

Mix.—For the same uses as the preceding.

ÆTHEREOUS LINIMENT. (SAUNDERS)

R. Soft Soap, } of each,
Rectified Alcohol, } one ounce.
Dissolve, and add
Camphor, half an ounce,
dissolved in
Sulphuric Æther, one ounce.

SAAPONACEOUS OINTMENT.

Unguentum saponaceum, Linimentum saponaceum camphoratum. (HAMB.; CADET DE GASSICOURT; SWEDIAUR; VAN MONS)

R. White Soap, one part.
Olive Oil, five parts.

Triturate the soap in a mortar, then dissolve it in the oil, and decant after sufficient repose. (VAN MONS)

R. Camphor, two drams.
Olive Oil, two ounces.

Dissolve, and add

Venice Soap, one ounce.

Mix. (HAMB.)

R. Soft Soap, half an ounce.
Camphor, two drams.
ARCEUS' Balsam, one ounce.

Mix. (CADET DE GASSICOURT; SWEDIAUR)

It is recommended in rheumatism.

ANTI-PLEURITIC LINIMENT.

Linimentum anti-pleuriticum seu Saponis.
(WIRTEM.; ELLIS; SPIELMANN)

- R. Venice Soap, two ounces.
Oil of Bay, } of each,
—— Earth Worms, } one ounce.

Add to the solution

Solution of Ammonia, two drams.

Mix. (WIRTEM.)

- R. Venice Soap, two ounces.
Oil of Bay, one ounce.

Melt with a gentle heat, and add, when cold,

Balsamum Tranquillum, one ounce.

Solution of Ammonia, two drams.

Mix. (SPIELMANN)

- R. Liniment of Soap, two ounces.
Solution of Ammonia, one ounce.
Tincture of Opium, half an ounce.

Mix. (ELLIS)

ANTI-PARALYTIC OINTMENT.

Unguentum Paralyticum. (BRUNS.; HERBIP.; NIEMANN; SPIELMANN)

- R. Venice Soap, six ounces.
Oil of Juniper, one ounce.
Seeds of Rocket, six drams.

Triturate together.

BALSAM OF AMBER. (VAN MONS)

- R. Insipissated Oil of Amber, one part.
Spanish Soap, two parts.

Melt together.

RESOLVENT OINTMENT. (SAINTE-MARIE)

- R. Olive Oil, }
Lime Water, } of each, two ounces.
White Soap, }
White of Egg, }

Shake well together.

SAPONACEOUS PLASTER OF CUMIN.
(SAUNDERS)

- R. Hard Soap, } of each, one ounce.
Yellow Wax, }
Cumin Plaster, three ounces.

Mix.

SAPONACEOUS PLASTER.

Emplastrum de Sapone seu Saponis seu saponatum seu saponaceum seu mamillare, Ceratum Saponis. (LOND.; DUBL., EDINB.; AMER.; ANT.; GALL.; FORM. HÔP. FR.; LIPP.; LUSIT.; SARD.; SUEC.; CADET DE GASSICOURT; COXE; VAN MONS)

Ceratum Saponis. (LOND.; AMER.; COXE)

- R. Hard Soap, eight ounces.
Yellow Wax, ten ounces.
Semi-vitreous Oxide of Lead, one pound.
Olive Oil, one pint.
Vinegar, one gallon.

Boil the vinegar with the oxide over a slow fire, constantly stirring, till they incorporate; then add the soap, and boil again in the same manner till the humidity is entirely evaporated: finally add the wax melted with the oil.

- R. Subcarbonate of Lead, } of each,
Deutoxide of Lead, } one pound.
Olive Oil, three pounds.
Soap, eight ounces.

Boil together, constantly stirring, adding water gradually, till the mass assumes the consistence of a plaster. (SARD.)

- R. Diachylon Plaster, two hundred parts.
Deutoxide of Lead, fifty parts.
Subcarbonate of Lead, twenty-five parts.

White Wax, ninety-six parts.
Water, a sufficient quantity.

Boil, constantly stirring; and, having removed it from the fire and separated the water, add

White Soap, one hundred and twenty-five parts.

Boil slowly. (GALL.; CADET DE GASSICOURT)

- R. Simple Diachylon Plaster, two pounds.
White Wax, three ounces.

Liquify over a slow fire, and add

White Soap, half a pound.

Make a plaster. (ANT.)

JEROMEL has proposed to melt together one pound of diachylon, and one ounce of white wax, and to incorporate with the mass when cold, by small portions, three ounces of medicinal soap.—FORM. HÔP. FR. prescribes eight parts of diachylon, two of wax, and four of soap.

Emplastrum Saponis. (LOND.; DUBL.)

- R. Hard Soap, sliced, half a pound.
Lead Plaster, three pounds.

Melt the plaster, add the soap, and boil to a proper consistence.

Emplastrum saponaceum. (EDINB.)

R. Semivitreous Oxide of Lead, four parts.
 Gum Plaster, two parts.
 Soap, one part.

Mix the soap with the plasters melted together, and boil so as to form a plaster.

SUEC. prescribes eight parts of diachylon plaster, and one part of soap.—LUST.—one of plaster, and three of soap.—COXE—six of plaster, and one of soap.—VAN MONS—ten parts of gum plaster, one half part of olive oil, one part of white wax, and four parts of white soap.—LIPP.—two parts of gum plaster, and one part of Spanish soap.

CAMPHORATED SAPONACEOUS PLASTER.

Emplastrum saponatum seu Saponis camphoratum seu resolvens cum Sapone et Camphora. (AMST.; ANT.; BATAV.; BAVAR.; BELG.; DAN.; GALL.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; BRERA; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Oil of Roses, twelve ounces.
 Deutoxide of Lead, } of each,
 Subcarbonate of Lead, } six ounces.

Boil, and add

Soap, four ounces.

Stir till cold, then add

Camphor triturated with a little Oil,
 half an ounce.

Mix. (SPIELMANN)

R. Simple Diachylon Plaster, three pounds.
 Yellow Wax, half a pound.
 Soap, three ounces.

Melt together, and add

Camphor, half an ounce.

Mix. (BATAV.; BELG.; DAN.; HANN.; OLDEN.; POL.; BORUS.)

ANT. directs ten grains of camphor to be added to each ounce of simple saponaceous plaster.—LIPP.—one twenty-fourth.—FORM. HÔP. FR.—one twenty-fifth and one fiftieth.—VAN MONS—one forty-eighth.—SAX.—one part of camphor, two parts of soap, four of wax, and twenty-four of simple plaster.

R. Simple Diachylon Plaster, one pound and a half.

Melt it over a slow fire, and add, constantly stirring,

Spanish Soap, scraped, four ounces.
 Camphor, triturated with a little Oil,
 half an ounce.

Mix. (HASS.; PIDERIT)

AMST. prescribes three pounds of plaster, half a pound of soap, and six drams of camphor.—ROSS.—three pounds of plaster, half a pound of soap, and one ounce of camphor.—FULD.—two ounces of plaster, one ounce of soap, and one dram of camphor.—BRUGNATELLI—six ounces of plaster, one ounce of soap, and one dram of camphor to each pound.—SWEDIAUR—one pound of plaster, one ounce of soap, and one dram of camphor.—GALL.—one pound of plaster, five ounces of wax, and one ounce of camphor.—BRERA—half a pound of plaster, four ounces of soap, and half an ounce of camphor.

R. Yellow Wax, } of each, four parts.
 Resin, }

Melt, and add

Soap, four parts.

Add to the mass when nearly cold

Camphor, triturated with an equal
 quantity of Oil,
 one ninety-sixth part.

Mix. (BAVAR.)

BALSAMIC PLASTER. (SWEDIAUR)

R. Diachylon Plaster, five pounds.

Melt, and add

White Soap, one pound.
 Camphor, three ounces.
 Castor, triturated with Alcohol,
 one ounce and a half.

Mix.

OIL OF SOAP.

Oleum Saponis. (BRUNS.; HISP.; FULD.; PALAT.; SARD.; WIRTEM.)

R. Soap, one part.
 Washed Sand, three parts.

Distil from a retort placed in a sand-bath, and rectify the oil upon ashes of vine twigs. (BRUNS.; FULD.; PALAT.; SARD.; WIRTEM.)

HISP. directs one part of soap to be distilled with one of pounded brick, and the oil to be re-distilled twice or thrice to obtain it limpid.

It is excitant, and was formerly extolled as a resolvent to ecchymoses, to appease pain, and as a detergent to sordid ulcers.

POTENTILLA.

Two species of this genus are used in medicine :

1. *Potentilla Anserina*, LINN.—Silverweed.

SYNONIMES :—Anserine vulgaire, FR. ; Gänserich, Fingerkraut, Silberkraut, GERM.

BRUNS. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A common plant of Europe. (*Icosandria Polygynia*, LINN. ; *Rosaceæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 15*)

The root and herb are employed.

The root (*Radix Anserinæ* seu *Argentina* seu *Argentariæ*) is fibrous and blackish. It has a sweetish taste.

The herb consists of a very slender stem, furnished with pinnate leaves with toothed, hairy leaflets, brilliant and silvery underneath. It is without odour, but of an acerb and rather styptic taste.

It is astringent.

2. *Potentilla reptans*, LINN.—Cinquefoil, Five-leaved Grass.

SYNONIMES :—Quintefeuille, FR. ; Fünffingerkraut, Fünfblatt, Handblatt, GERM. ; Femfingerert, DAN. ; Cinco en rama, PORT. ; and SPAN. ; Vyfvingerkruid, DUT. ; Cinquefoglio, ITAL. ; Pieciornik, POL. ; Femfingerært, SWED.

AMST. ; BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; SPIELMANN.

A common plant of Europe. (*Icosandria Polygynia*, LINN. ; *Rosaceæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 302*)

The root and herb are employed.

The root (*Radix Potentillæ* seu *Pentaphylli* seu *Quinquefolii majoris*) is cylindrical, of a blackish-brown colour without, whitish within, and of a feebly styptic taste.

The herb consists of a filiform stem, and of petiolated, quinate leaves with retuse, serrated leaflets of a light green colour. Its taste is bitterish and astringent.

POTERIUM SANGUISORBA, LINN.—PIMPERNEL, SMALL BURNET.

SYNONIMES :—Pimprenelle commune, FR. ; Pibernelle, GERM. ; Pimpinelle, DAN. ; Pimpinella, ITAL. ; PORT. ; and SPAN. ; Pimpernelle, DUT. ; Pimpernella, SWED.

BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; MURRAY and GMELIN.

A plant of the south of Europe. (*Monœcia Polyandria*, LINN. ; *Rosaceæ*, JUSS.)

The root and herb are employed.

The root (*Radix Pimpinellæ minoris* seu *hortensis* seu *rubræ* seu *Italica*) is thick, long, reddish, and divided into several fibres.

The herb consists of an angular stem, rather branched, and furnished, particularly at the base, with winged leaves having slightly hairy footstalks ; they are composed of from eleven to twenty-one leaflets nearly equal, round or oval, glabrous, and rather deeply toothed. Its odour is feeble and aromatic, and its taste astringent and bitterish.

PRIMULA OFFICINALIS, LINN.—COWSLIP.

SYNONIMES :—Primevère, FR. ; Schlüsselblume, GERM. ; Oxedrif, DAN. ; Primula veris, SPAN. ; Sleutelbloem, DUT. ; Primavera, ITAL. ; Bakwiza, POL. ; Primevera, PORT. ; Oxelægg, SWED.

AMST. ; BRUNS. ; HISP. ; GALL. ; GENEV. ; WIRTEM. ; HERBIP. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of Europe (*Pentandria Mono-* Vol. II.

gynia, LINN. ; *Primulaceæ*, JUSS. ; *Fig. Flore Medic. V. 284*)

The root, herb, and flowers are employed.

The root (*Radix Primulæ veris* seu *paræzeos* seu *Verbasculi pratensis*) is a large, somewhat reddish, stock, furnished with long, white, fleshy fibres. It has an agreeable and anisated odour, and a slightly astringent and bitterish taste.

The herb consists of oblong, furrowed, thick, obtuse, pubescent, dentated leaves, whitish underneath. It is nearly inodorous, and of a purely herbaceous taste.

The flowers, the corollæ of which are only used, are monopetalous, cylindrical, five-lobed, of a yellow colour, of a pleasant odour and taste.

CONSERVE OF COWSLIP.

Conserva Primulæ veris. (PALAT.; WIRTEM.)

R. Flowers of Cowslip, one part.
White Sugar, two parts.

Triturate together.

WATER OF COWSLIP.

Aqua Paralyseos. (PALAT.)

R. Flowers of Cowslip, one part.
Water, three parts.

Distil one part and a half.

SYRUP OF COWSLIP.

Syrupus Florum Paralyseos seu Primulæ veris. (PALAT.; WIRTEM.)

R. Fresh Flowers of Cowslip,
three pounds and a half.

Boiling Water, four pounds.

Infuse for twenty-four hours, and express; dissolve in eighteen ounces of the strained liquor,

White Sugar, thirty-two ounces.

VINEGAR OF COWSLIP.

Acetum Primulæ veris. (HERBIP.)

R. Root of Cowslip, half dried, and bruised, one ounce and a half.
Vinegar, one pound.

Macerate for fifteen days, frequently agitating; then express slightly, and add,

Alcohol, one ounce.

Decant at the end of several days.

OXYMEL OF COWSLIP.

Oxymel Primulæ veris. (HERBIP.)

R. Vinegar of Cowslip, one part.
Honey, two parts.

Boil slowly to the consistence of syrup.

PRUNELLA VULGARIS, LINN.—SELFHEAL.

SYNONIMES: — Prunelle, FR.; Gottheil, Braunhëil, Selbstheil, GERM.; Czernoklawek, Swalnjk menssj, BOHEM.; Prunelle, DAN.; Brunella, SPAN.; Bruinella, DUT.; Zwokest modry, POL.; Prunella, PORT.

BRUNS.; HISP.; GALL.; WIRTEM.; HERBIP.; BERGIUS; MURRAY and GMELIN; SPIELMANN; ZARDA.

A common plant of Europe. (*Didymia Gymnospermia*, LINN.; *Labiata*, JUSS.; Fig. Zorn, Ic. pl. t. 156)

The flowering herb (*Herba Prunellæ seu Brunellæ seu Consolida minoris*) is employed. It consists of a square, rather branching stem, furnished with petiolated, oblong, oval, serrated leaves; the flowers are arranged in a terminal spike, and are formed of an assemblage of close verticilli. It is destitute of odour, but its taste is bitterish and mucilaginous.

It was formerly esteemed vulnerary.

JUICE OF SELFHEAL.

Succus Prunellæ (HERBIP.)

R. Fresh Herb of Selfheal, any quantity.

Bruise it in a wooden mortar, sprinkling it with a little water; express strongly, set aside the juice, decant, and preserve it in bottles, adding to each pound of it, one dram of alcohol.

SYRUP OF SELFHEAL.

Syrupus Prunellæ. (WIRTEM.)

R. Juice of Selfheal, depurated and reduced to one half by slow evaporation, nine ounces
White Sugar, sixteen ounces.

Dissolve.

PRINOS VERTICILLATUS, LINN.—DECIDUOUS WINTERBERRY.

SYNONIMES:—Prinos, Apalachine à feuilles de prunier, FR.

AMER.; COXE.

A shrub of North America. (*Hexandria*

Monogynia, LINN.; *Rhamnea*, JUSS.; *Fig. Duham. I. t. 23*)

The bark is employed, it is bitter and sub-acrid.

PROPOLIS.

SYNONIMES:—Propolis, FR.; Stopfwachs, Bienenwachs, GERM.

GENEV.; and SPIELMANN.

A resinous matter elaborated by the domestic bee, *Apis mellifica*, LINN.

To obtain this substance pure, it is put into a pan with about twice its weight of water; when this is boiling and the propolis is melted, the whole is to be poured upon a new cloth, strained with expression, and the resin received into cold water; when it has become congealed it is rolled into cylinders.

OINTMENT OF PROPOLIS.

Unguentum propolicum. (CADET DE GASSICOURT; SWEDIAUR)

R. Purified Propolis, four ounces.
Olive Oil, one pound and a half.

Dissolve at a moderate heat, and strain while hot. (CADET DE GASSICOURT)

SWEDIAUR prescribes one part of propolis and two parts of oil.

It is applied to old ulcers and to piles.

PRUNUS.

Two species of this genus of plants are indicated in pharmacopœias:

1. *Prunus domestica*, LINN.—Plum Tree.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; WIRTEM.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A tree which is met with in every orchard. (*Icosandria Monogynia*, LINN.; *Rosaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 493*)

The fruit, called Plum (*Prunus*), is employed.

SYNONIMES:—Prune, FR.; Pflaume, GERM.; Sswestka, Ssjwa, BOHEM.; Plom, DAN.; Cirnelo, SPAN.; Pruin, DUT.; Prugno, ITAL.; Stiwinia, POL.; Ameixa, PORT.; Plom, SWED.

This fruit varies to infinity in size, form, and colour. It has a sweet and generally subacidulous taste. When dried, either by the sun or in an oven, they are called *Prunes* (*Pruneaux*, FR.), when the damask plum (*Prune de Damas*, FR.) is usually employed.

2. *Prunus spinosa*, LINN.—Sloe.

SYNONIMES:—Prunier sauvage, FR.; Steckdorn, Schlehenbaum, GERM.; Slaetorn, Kreygebærtorn, DAN.

AMST.; AUSTR.; BAVAR.; DAN.; HISP.; GALL.; FERR.; FULD.; LIPP.; ROSS.; SAX.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A shrub common to all parts of Europe. (*Icosandria Monogynia*, LINN.; *Rosaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 4*)

The bark, flowers, and fruit are employed.

The bark (*Cortex Acaciæ nostratis seu vulgaris seu Germanicæ seu Pruneoli sylvestris*) is of a brown-grey colour, and of a bitter and astringent taste.

The flowers, which are white, have an agreeable odour which is not entirely dissipated by desiccation, and a bitter and rather styptic taste.

The fruit, called *Sloes* (Prunelles, Fr.), are round, black drupes, covered with a blue powder; under a greenish-yellow flesh, each contains a round smooth stone. Their taste is acerb and astringent, but becomes sweet after frost.

The kernel has the taste of bitter almonds.

The whole plant is astringent. The inspissated juice of the fruit constitutes the *Acacia nostras*. The flowers constitute a domestic aperient, in the dose of half an ounce in the fresh state, or of two drams in the dried, infused in six ounces of water.

PULP OF PRUNES.

Pulpa Prunorum. (ANT.; BAVAR.; DAN.; GALL.; HISP.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; HERBIP.; PIDERIT; SAINTE-MARIE; VAN MONS)

R. Tours Prunes, any quantity.

Soften them in hot water; pour off the liquor, remove the stones and pass the pulp through a hair sieve, pressing it with a wooden spoon: preserve it in a covered pot. (LIPP.; VAN MONS)

GALL.; and LUSIT.; direct the pulp to be mixed with the water of the infusion, and the mixture to be evaporated with a gentle heat, to a proper consistence.

R. Prunes, one part.
Water, two parts.

Boil till the fruit becomes soft; then remove the stones, strain the pulp through a sieve, and evaporate it, adding one sixth of sugar. (BAVAR.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SAINTE-MARIE)

HISP.; LUSIT.; and PIDERIT prescribe one part of sugar and two parts of pulp.—ANT.; DAN.; ROSS.; and HERBIP.—one of sugar and three of pulp.

It is a mild laxative.

LAXATIVE DECOCTION.

Decoctum purgans anti-phlogisticum. (BRERA; SPIELMANN)

R. Prunes, two ounces.

Bruise them with the stones, and add,

Water, three pounds.

Boil down to one third, and add to the strained liquor,

Simple Syrup, one ounce.

Mix.—To be taken at twice. (BRERA)

R. Prunes, four ounces.

Water, three pounds.

Boil down to one-fifth, and add towards the end,

Liquorice, four drams.

Strain when cold. (SPIELMANN)

CONSERVE OF SLOES. (SARD.; BRUGNATELLI)

R. Sloes, any quantity.

Digest them in water over a slow fire till they become soft; then pass the pulp through a silk sieve, add three parts of sugar, and evaporate to the proper consistence, constantly stirring. (SARD.)

BRUGNATELLI directs one part of sloes to be bruised with two parts of sugar, so as to form a homogeneous paste.

ROB OF SLOES.

Acacia nostras, Rob Prunorum sylvestrium seu Fructuum Acacia Germanica. (AUSTR.; SARD.)

R. Sloes, nearly ripe, any quantity.

Bruise them, express the juice, and evaporate it in a water-bath, constantly stirring with a spatula, till it is of the consistence of an electuary.

WATER OF SLOE-FLOWERS.

Aqua Florum Acaciae. (FULD.; LIPP.; PALAT.; SAX.; HERBIP.; SAINTE-MARIE)

R. Dried Sloe-flowers, two parts.
Water, a sufficient quantity.

Distil twenty parts. (SAINTE-MARIE)

PALAT. prescribes one part of flowers and three parts of water; one half to be distilled.—LIPP.; and HERBIP.—one of flowers and six of water.—SAX.—one of flowers and eight of water; one half to be distilled.

R. Dried Sloe-flowers, three pounds.

Suspend them, in a bag, in the middle of a cucurbit containing,

Water, twelve pounds.

Distil one pound and a half without making the water boil. (FULD.)

SYRUP OF SLOE-FLOWERS.

Syrupus Acaciarum seu Aciarum Florum. (PALAT.; WIRTEM.)

R. Fresh Sloe-flowers, } of each,
Boiling Water, } two pounds.

Digest for one night in a covered vessel. Add to nineteen ounces of the strained liquor,

White Sugar, thirty-two ounces.

Boil, and strain. (WIRTEM.)

PALAT. directs two pounds of flowers to be infused in four of boiling water for twelve hours ; the liquor to be poured upon two pounds of fresh flowers : this to be repeated a third time, and the syrup to be

finally made with the strained liquor and four pounds of sugar.

It is laxative, anthelmintic, and anti-nephritic.—Dose, from one dram to three drams, to children ; and from six drams to one ounce, to adults.

PSORALEA GLANDULOSA, LINN.—STRIPED FLOWERED PSORALEA.

SYNONIMES :—Thé du Paraguay, Thé de la mer du Sud, FR. ; Drusige Psorbalea, GERM. ; Culen, SPAN.

ELLIS.

A plant of Peru and of Brasil, (*Diadel-*

phia Decandria, LINN. ; *Leguminosæ*, JUSS. ; *Fig. Feuill. Per. 7. t. 3*)

The herb (*Herba Culen*) is employed. It consists of ternate leaves with lanceolated leaflets.

It is stomachic, and anthelmintic.

PTERIS AQUILINA, LINN.—COMMON BRAKE.

SYNONIMES :—Fougère femelle, Pteride, FR. ; Flügelfarren, Adlersaumfarren, Farrenkrautweiblein, GERM.

FULD. ; **ROSS.** ; **WIRTEM.** ; **HERBIP.** ; **MURRAY** and **GMELIN.**

A plant which grows in every part of Europe. (*Cryptogamia*, LINN. ; *Filices*, JUSS. ; *Fig. Bull. Herb. t. 207*)

The stock, improperly called the root

(*Radix Pteridis aquilinæ*) is employed. It is of the thickness of the finger, long, black without, and yellow within. When cut obliquely it presents two blackish-crossed lines, rudely representing an eagle with two heads. It has a very strong odour, and a taste at first mucilaginous, then bitterish, nauseous, and slightly styptic.

It is anthelmintic.—Dose, thirty grains to children, and three drams to adults.

PULMONARIA OFFICINALIS, LINN.—LUNGWORT.

SYNONIMES :—Grande pulmonaire, FR. ; Lungenkraut, GERM. ; Spettel lungeurt, DAN. ; Pulmonaria, PORT. ; and SPAN. ; Gewoon lungekruid, DUT. ; Polmonaria, ITAL. ; Placknik, POL. ; Meduniza, RUSS. ; Lungært, SWED. ; Koiwiek, TAR.

AMST. ; **ANT.** ; **BRUNS.** ; **HISP.** ; **GALL.** ; **FERR.** ; **WIRTEM.** ; **BERGIUS** ; **BRUGNATELLI** ; **GUIBOURT** ; **MURRAY** and **GMELIN** ; **SPIELMANN.**

A plant of Europe. (*Pentandria Monogymia*, LINN. ; *Borraginææ*, JUSS. ; *Fig. Flore Medic. V. 286*)

The herb (*Herba Pulmonariæ maculosæ seu Symphyti maculosi*) is employed. It consists of radical, oval, petiolated, bristled leaves sprinkled with white spots, and of

stems the leaves of which are alternate, sessile, longer, narrower, and entire. It is nearly destitute of odour, and has an herbaceous, mucilaginous, and rather astringent taste.

It used formerly to be esteemed in pulmonary catarrh. It is now nearly disused.

JUICE OF LUNGWORT.

Succus Pulmonariæ. (GALL.)

R. Leaves of Lungwort, sixteen parts.

Bruise them in a stone mortar, adding gradually,

Water, one part.

Express the juice, set aside, and filter without heat through bibulous paper.

PUMEX, LAPIS PUMICIS.—PUMICE STONE.

SYNONIMES :—Ponce, Pierre Ponce, FR. ; Bimstein, Bimsenstein, GERM. ; Piedra pomez, SPAN. ; Puimsteen, DUT. ; Pietra pomice, ITAL. ; Zazel kamienna, POL.

BRUNS. ; HISP. ; GALL. ; HANN. ; WIRTEM. ; HERBIP. ; SPIELMANN.

A volcanic stone, of a whitish-grey or yellowish colour, very porous, light, tenacious, fragile, fibrous, shining or even silky in the course of its fibres, presenting a grained and unequal fracture ; it is dry to

the touch, lighter than water, and fusible into a white glass.

It enters into several dentifrice powders.

WHITE DENTIFRICE POWDER. (BRUNS. ; SPIELMANN)

R. Pumice Stone,	two ounces.
Cuttle Fish Bone,	one ounce.
Supertartrate of Potass,	of each,
Root of Iris,	halfan ounce.
Oil of Cloves,	four drops.

Make a very fine powder.

PUNICA GRANATUM, LINN.—POMEGRANATE TREE.

SYNONIMES :—Grenadier, Balaustier, FR. ; Granatbaum, GERM. ; Ranua, Roman, ARAB. ; Pomapran, Granatowe gabto, BOHEM. ; Delunghedie, CYNG. ; Anar, HIND. ; and PERS. ; Balaustries, SPAN. ; Rodia, MOD. GR. ; Granatboom, DUT. ; Granato, ITAL. ; Gangsalan, JAV. ; Dailimb, MAH. ; Dalema, Daime, MAL. ; Drzewo granatowe, POL. ; Romeira, PORT. ; Granatoschnoe derewo, RUSS. ; Dadima, SANS. ; Darim, SWED. ; Magilam palam, TAM. ; Dadima pundu, TEL. ; Nar, TUR.

LOND. ; DUBL. ; AMER. ; AMST. ; ANT. ; BATAV. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; SAX. ; WIRTEM. ; HERBIP. ; AINSLIE. ; BERGIUS. ; BRUGNATELLI. ; COXE. ; GUIBOURT. ; MAGENDIE. ; PARIS. ; SPIELMANN. ; ZARDA.

A tree of the south of Europe. (*Icosandria Monogynia*, LINN. ; *Myrtæa*, JUSS. ; *Fig. Zorn, Ic. pl. t. 270*)

The bark of the root, the flowers, and the fruit are employed.

The bark of the root is yellowish-grey externally, and greyish-yellow or slightly reddish internally, inodorous, and of a feebly styptic and bitterish taste. Its powder is greenish-yellow. It has been represented as an infallible remedy for the expulsion of tænia.

The flowers (*Balaustia*, *Flores Balaustiorum*) are hemispherical, of a fine red colour, nearly inodorous, and a slightly styptic taste. They are astringent, but less so than the rind of the fruit.

The fruit, called Pomegranates (*Mala*

Punica), are round berries, terminated by the sepals of the calyx and divided by a transverse partition into two multilocular cells, filled with angular seeds surrounded by a pulpy arillus. The exterior pellicle (*Malacorium*, *Malicorium*) is greyish-yellow or reddish, thick, hard, and coriaceous.

It has an astringent taste. It has been employed in every case in which an astringent is indicated.

It is incompatible with the salts of iron. REHMANN has recommended the rind of the fruit in intermittent fevers, from six to twelve doses of from two to six scruples each, to be taken during the apyrexia.

ASTRINGENT SPECIES.

Species Decocti pro Gargarismate adstringente seu adstringentes pro Gargarismate. (GALL. ; FORM. HÔP. FR. ; HANN. ; PALAT. ; NIEMANN. ; PIDERIT)

R. Root of Bistort ;	} of each,
Pomegranate Flowers,	
Hollyhock Flowers,	

Cut, and mix. (HANN.)

R. Herb of Plantain,	} of each,
Selfheal,	
Sage,	} four ounces.
Flowers of Red Roses,	
Ligustrum,	
Pomegranate,	} of each,

Cut, and mix. (PALAT. ; WIRTEM.)

BRUNS. gives the same formula but without the ligustrum.

R. Root of Bistort,	} of each,
Herb of Agrimony,	
Periwinkle,	
Flowers of Hollyhock,	
Pomegranate,	two ounces.

Cut, and mix. (PIDERIT)

R. Pomegranate Rind, } of each,
 Root of Bistort, } an equal part.
 ——— Tormentil, }

Cut, and mix. (GALL.; FORM. HÔP. FR.)

ANTI-HÆMORRHOIDAL SPECIES.

Species pro Fotu ad hæmorrhoidum fluxum immodicum. (PALAT.)

R. Herb of Milfoil, } of each,
 ———— Plantain, } four handfuls.
 ———— Lesser Mullein, }
 Flowers of Elder, }
 Pomegranate Rind, one ounce.

Mix.

JUICE OF POMEGRANATE.

Succus Granatorum. (HISP.; GALL.; SWEDIAUR)

R. Unripe Pomegranates, any quantity.

Remove the external rind, cut the rest, and triturate with a wooden spatula in order to separate the seeds; then bruise in a stone mortar, and express the juice either in a bag or between two layers of rye straw chopped and washed with warm water. Set the juice aside, decant, and preserve it in bottles under a stratum of oil.

SYRUP OF POMEGRANATE.

Syrupus Granatorum seu de Granatis. (HISP.; GALL.; PALAT.; WIRTEM.)

R. Depurated and filtered juice of Pomegranate, two pounds.
 White Sugar, three pounds and a half.

Dissolve at a moderate heat. (GALL.)

HISP.; and LUSIT. prescribe one part of juice and two parts of sugar.—WIRTEM.—ten ounces of the former and sixteen of the latter, to be boiled to a syrup.

DECOCTION OF THE BARK OF POMEGRANATE ROOT. (ELLIS; FEE; MAGENDIE; NIEMANN)

R. Bark of Pomegranate Root, two ounces.
 Water, two pounds.

Boil down to three-fourths.

It is recommended against tænia.—Dose,

two ounces every half hour.—Four doses usually suffice for the expulsion of the worm.

DECOCTION OF POMEGRANATE RIND. (FORM. HÔP. FR.)

R. Pomegranate Rind, one ounce.
 Water, a sufficient quantity,
 to obtain two pounds of strained decoction.
 Boil for a quarter of an hour, and add
 Liquorice, two drams.
 It is astringent.

ASTRINGENT FOMENTATION. (RATIER)

R. Pomegranate Rind, } of each,
 Root of Bistort, } two ounces.
 Red Wine, one pound.
 Boil and add when strained,
 Muriate of Ammonia, two drams.
 Dissolve.

ASTRINGENT LAVEMENT. (HISP.)

R. Pomegranate Rind, one ounce.
 Water, a sufficient quantity,
 to obtain two pounds of strained decoction.

ASTRINGENT GARGLE. (CADET DE GASCICOURT)

R. Infusion of Agrimony, } of each,
 ———— Cypress Nuts, } four
 ———— Pomegranate } ounces.
 Rind, }
 ———— Elder Flowers, }
 Honey of Roses, } of each,
 Syrup of Mulberries, } three drams.
 Sulphuric Acid, } twenty drops.

Mix.

REHMANN'S DECOCTION OF POMEGRANATE RIND. (NIEMANN)

R. Pomegranate Rind, two ounces.
 Gum Arabic, one dram.
 Spring Water, two pounds.
 Boil down to one half, and to each pound of the decoction add

Extract of Buckbean, half a dram.
 Alcoholized Sulphuric Æther, one dram.

Mix.—It is astringent.—Dose, half a glassful every hour and a half.

PYROLA ROTUNDIFOLIA, LINN.—WINTERGREEN.

SYNONIMES:—Pyroleà feuilles rondes, FR.; Waldmangold, Rundblättriges Wintergrün, GERM.; Vintergræn, DAN.

BRUNS.; HISP.; GALL.; LIPP.; SAX.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of the north of Europe. (*Decandria Dggyinia*, LINN.; *Ericineæ*, JUSS.; *Fig. CEd. Fl. Dan. t. 110*)

The herb (*Herba Pyrolæ*) is employed. It consists of a bundle of radical leaves,

round, green, and coriaceous, having long footstalks. It is without odour, but has a bitter and somewhat styptic taste.

It was formerly esteemed vulnerary.

QUASSIA AMARA, LINN.—QUASSIA.

SYNONIMES:—Quassie, Bois de Surinam, FR.; Quassienholz, Americanisches Bitterholz, GERM.; Drzewo kwassya, POL.

LOND.; EDINE.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; LUSIT.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.

A tree of Surinam (*Decandria Monogynia*, LINN.; *Simaroubeæ*, JUSS.; *Fig. Desc. Fl. Ant. I. 5*)

The wood of the trunk and especially of the root is employed. It comes to us, in its bark, in pieces of various dimensions, sometimes scarcely of the thickness of the little finger, sometimes of the thickness of the arm, several feet long, cylindrical and smooth, or cloven lengthwise, usually straight, seldom curved, sometimes knotty and branched. The bark is of a yellowish-grey colour, not thick, smooth to the touch, and slightly adherent to the wood. The latter, is of a yellowish-white colour, light, soft in the interior. It is free from odour, but is extremely bitter.

Its bitterness is owing to a peculiar principle, quassine.

This wood is also attributed to the *Quassia excelsa*, Sw.; a tree of Jamaica. (*Fig. Comm. Hort. I. p. 149. t. 942*)

It is a tonic, and stomachic, and is used in dyspepsia, diarrhœa, and hæmorrhages.—Dose, of the powder, from a scruple to half a dram.

AQUEOUS EXTRACT OF QUASSIA.

Extractum Quassie. (AMER.; AMST.; ANT.; BAVAR.; BRUNS.; HANN.; HASS.; OLDEN.; POL.; LUSIT.; ROSS.; SAX.; HERBIP.; COXE; PIDERIT)

R. Quassia Shavings, one pound.
Cold Water, ten pounds.

Macerate for two days, occasionally agitating; then strain, set aside and decant the liquor, and finally evaporate it over a slow fire. (POL.)

R. Quassia Shavings, any quantity.

Boil it in a sufficient quantity of water for an hour; then express, set aside, and decant the liquor, and finally evaporate it to the consistence of thick honey. (HERBIP.)

LUSIT. prescribes one pound of wood and six pounds of water.

R. Quassia Shavings, one pound.
Spring Water, six pounds.

Digest for four days in a warm place, then boil a little; express the decoction, and evaporate it to the consistence of an extract. (BRUNS.)

AMER.; HASS.; COXE; and PIDERIT direct one pound of wood to be boiled in eight pounds of water till reduced to one half.

R. Quassia Shavings, one pound.
Boiling Water, eight pounds.

Digest for twenty-four hours, then boil for a quarter of an hour, and strain; boil the residue for a quarter of an hour with four pounds of water, and strain again. Mix the decoctions, set apart, decant, and evaporate. (ANT.)

AMST.; DAN.; HANN.; OLDEN.; POL.; BORUS.; ROSS.; and SAX. direct one part of wood to be boiled for several times successively in ten parts of water, and the mixed decoctions to be evaporated.—BAVAR.—one part of wood to be first boiled with five and then with fifteen parts of water, and the mixed liquors evaporated.

Dose, from ten to fifteen grains.

ALCOHOLIC EXTRACT OF QUASSIA.

Extractum Quassie alcoholicum. (FULD.; LIPP.)

R. Quassia Shavings, one pound.
Proof Spirit, six pounds.

Digest for several days at a moderate heat; then express and filter the tincture, and evaporate it to the consistence of honey; boil the residue with three pounds of water, and express, filter, and evaporate in the same manner. Mix the liquors, and finally evaporate to the consistence of extract. (FULD.)

LIPP. directs one pound and a half of wood to be digested, first in six pounds of alcohol for two days, then in nine pounds of boiling water for twenty-four hours; the liquors to be then mixed and evaporated.

TONIC PILLS. (AUGUSTIN; ELLIS)

- R. Extract of Quassia, one dram.
Powdered Gentian, a sufficient quantity.

Make twenty pills.—Dose, one, three or four times a day. (ELLIS)

AUGUSTIN prescribes one dram of the extract and a sufficient quantity of powdered ginger to make pills of two grains each, ten of which form a dose.

ANTI-ACID ELIXIR.

Elixir cum Aqua Calcarie usta. (AUGUSTIN)

- R. Extract of Quassia, one dram.
Lime Water, eight ounces.
Dissolve.—Dose, a tea spoonful.

INFUSION OF QUASSIÆ.

Infusum Quassiæ. (LOND.; DUBL.; EDINB.; AMER.; LUSIT.; COXE; ELLIS; NIEMANN; SWEDIAUR)

- R. Quassia Chips, one scruple.
Boiling Water, half a pint.

Macerate for two hours in a lightly covered vessel, and strain. (LOND.; DUBL.)

Infusum Quassiæ Excelsæ. (EDINB.)

- R. Quassia Shavings, half a dram.
Boiling Water, eight ounces.

Macerate for two hours in a lightly covered vessel, and strain.

AMER.; and COXE give the same proportions as EDINB.—NIEMANN—the same as LOND.—SWEDIAUR prescribes two drams of wood and one pound of water.—LUSIT.—three drams of wood and one pound of water.

- R. Quassia Shavings, } of each,
Serpentaria, } half an ounce.
Orange Peel, }
Boiling Water, two pounds.

Infuse, and strain. (ELLIS)

It is prescribed as a bitter.—Dose, one or two ounces several times a day.

INFUSION OF QUASSIA WITH SULPHATE OF ZINC.

Infusum Quassiæ cum Sulphate Zinci.
(AMER.; COXE)

- R. Quassia Shavings, one dram.
VOL. II.

Sulphate of Zinc, eight grains.
Cold Water, half a pound.
Macerate for two hours, and strain.

STOMACHIC POTION.

Haustus stomachicus. (BRERA)

- R. Calumba, one dram.
Quassia, one dram and a half.
Water, a sufficient quantity
to obtain eight ounces of strained infusion.

Add

WHYTT's Stomachic Elixir,
two drams.

Mix.—To be taken gradually.

DECOCTION OF QUASSIA.

Decoctum Ligni Quassiæ. (FULD.; HAMB.)

- R. Quassia Shavings, one ounce.
Spring Water, two pounds.

Boil till reduced to a pound and a half. (HAMB.)

- R. Quassia Shavings, one ounce and a half.
Water, four pounds.

Boil down to three pounds, adding towards the end

Chamomile Flowers,
one ounce and a half.

Set aside and decant. (FULD.)

Dose, a spoonful every two or three hours.

COMPOUND DECOCTION OF QUASSIA.

Decoctum Quassiæ Ligni compositum, Decoctum amarum. (HERBP.)

- R. Quassia Shavings, one ounce.
Chamomile Flowers, one ounce and a half.
Subcarbonate of Potass, two drams.
Spring Water, four pounds.

Boil down to three-fourths, and strain.

VINOUS TINCTURE OF QUASSIA. (HANN.)

- R. Quassia Shavings, four ounces.
Malaga Wine, two pounds.
Digest, and filter.

It is tonic.—Dose, three drams or half an ounce, twice a day.

ALCOHOLIC TINCTURE OF QUASSIA.

Tinctura Quassiæ. (DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BELG.; DAN.; GENEV.; HANN.; OLDEN.; SAX.; COXE; SWEDIAUR; VAN MONS)

- R. Quassia Chips, one ounce.
Proof Spirit, two pints.

Digest for seven days, and strain.
(DUBL.)

Tinctura Quassiae excelsæ. (EDINB.)

R. Quassia Shavings, one ounce.
Proof Spirit,

two pounds and a half.

Digest for seven days, and filter.

AMER.: and COXE give the same proportions as DUBL.—BATAV.; and BELG. prescribe one part of wood and six parts of alcohol (15 degrees).—AMST.—one of wood and six of alcohol (0.917).—SWEDIAUR—two ounces of wood and two pounds and a half of alcohol (0.935).—VAN MONS—one part of wood and six parts of proof spirit.

—HANN.; and OLDEN.—five ounces of wood and two pounds of alcohol.—DAN.—one part of wood and five parts of alcohol. SAX.—one of wood and ten of alcohol.

R. Quassia Shavings, one ounce.
Alcohol (20 degrees), four ounces.

Digest for four days in a water-bath, and strain; pour upon the residue

Alcohol (20 degrees), two ounces.

Strain at the end of two days. Mix the filtered tinctures. (ANT.)

R. Quassia Shavings, two ounces.
Orange Peel, one ounce.
Lesser Cardamoms, half an ounce.
Proof Spirit, two pounds.

Macerate for four days, and strain.
(GENEV.)

QUERCUS.

Several species of this genus are mentioned in pharmacopœias:

1. *Quercus alba*, LINN.—White Oak.

SYNONIMES:—Chêne blanc, FR.; Weisse Eiche, GERM.

AMER.; COXE.

A tree of the United States. (*Monœcia Polyandria*, LINN.; *Amentaceæ*, JUSS.; *Fig. Mich. Arb. Amer.* II. t. 1)

The bark is employed; it is astringent.

2. *Quercus tinctoria*, MICH.—Black Oak, Quercitron Oak.

SYNONIMES:—Chêne Quercitron, FR.; Quercitroneneiche, GERM.

AMER.; COXE; GUIBOURT.

A tree of the United States. (*Fig. Mich. Arb. Amer.* II. t. 22)

The bark is employed; it is very astringent.

3. *Quercus Robur*, LINN.—Oak.

SYNONIMES:—Chêne Rouvre, FR.; Eiche, GERM.; Dub, BOHEM.; Eeg, DAN.; Majup'hal, HIND.; and SANS.; Eik, DUT.; Quercia, ITAL.; Kora debowa, PORT.; Ek, SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.;

LIPP.; OLDEN.; LUSIT.; CAST. BORUS.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

A large tree common to all parts of Europe. (*Fig. Nouv. Duh.* VII. 52)

The bark and fruit are employed.

The bark of the young branches (*Cortex Quercus seu quercinus*) is glabrous, ash coloured externally, and reddish internally. It has a peculiar odour, and a bitter and very astringent taste.

The fruit, acorns (*Glandes seu Nuces Quercus*;—Glands, FR.; Eichel, GERM.; Zoladz, POL.), are ovoid or oval, and prolonged; they are inodorous, but of an acerb and bitter taste.

Their cups (*Cupulæ seu Calyculæ Glandium Quercus*) are covered with greyish scales closely imbricated. (BATAV.; BAVAR.; BRUNS.; DAN.; FENN.; HANN.; HASS.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; HERBIP.; BERGIUS; MURRAY and GMELIN)

The *Quercus pedunculata*, WILLD. is also indicated by BAVAR.

4. *Quercus Ilex*, LINN.

SYNONIME:—Chêne Yeuse, FR.

HISP.

A tree of the south of Europe and north of Africa. (*Fig. Zorn, Ic. pl.* t. 518)

Its acorns are employed; they are sometimes sweet and good for eating. The cups are furnished with very small scales, minutely imbricated, and cottony.

The barks of the oaks, especially of the *Q. Robur* contain much tannin with an extractive matter. They possess a very decided astringency, so that when administered in a large dose, they irritate the digestive organs, and yet, at the same time, do not determine a general reaction. They have been proposed as a substitute for cinchona, but they seem to be very inferior to it as a febrifuge.—Dose, from a dram to one ounce, of the powder.

As to the acorns, those of them which are bland are used as food: the others, when roasted, have been recommended as a useful excitant in glandular affections, particularly in scrofula.

5. *Quercus Suber*, LINN.—Cork Tree.

SYNONIMES :—Liege, FR.; Korcheiche, GERM.; Korketræ, DAN.; Alcorhogre, SPAN.; Korkboom, DUT.; Saverio, ITAL.; Savereiro, PORT.; Korktræd, SWED.

GALL.; SUEC.; WIRTEM.; BERGIUS.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

A tree of the south and east of Europe. (*Fig. Nouv. Duh. VII. 45*)

The bark and fruit are employed.

The bark (*Cortex Suberi seu Suber*) is flexible, light, cracked, spongy, elastic, and of a reddish-yellow colour. It used formerly to be given internally as an astringent.

It has for its base, according to CHEVREUIL, a peculiar principle called *Suberine*.

Its acorns have generally a sweet and agreeable taste.

DECOCTION OF CORK. (BORIES; PIERQUIN)

R. Cork, rasped, two scruples.
Water, three pounds.

Boil down to one half. (PIERQUIN)

BORIES directs two ounces of syrup of maidenhair to be added.

Dose, eight ounces, thrice a day, for eight or ten days.—Syrup of water-lily may be added, and its action aided by a lavement made with a strong decoction of cork or of parsley, administered every morning.

COMPOUND POWDER OF OAK BARK. (CAST. BORUS.; AUGUSTIN)

R. Oak Bark, one scruple.
Root of Sweet Flag, } of each,
Gentian, } five grains.

Make a powder. (CAST. BORUS.)

R. Oak Bark, five grains.
Orange Peel, } of each, ten grains.
Olibanum, }
Make a powder. (AUGUSTIN)

ANTI-SEPTIC CATAPLASM.

Cataplasm e tanniné. (FORM. HÔP. FR.; LIPP.)

R. Oak Bark, in powder, } of each,
Willow Bark, } one part.
Camphorated Vinegar, } six parts.
Mix. (LIPP.)

R. Barley Meal, } of each,
Rye Meal, } an equal part.
Oak Bark, in powder, }
Water, a sufficient quantity.

Boil to a proper consistence. (FORM. HÔP. FR.)

By adding to the mass when cold a fiftieth or twenty-fifth of camphor, the *Camphorated anti-septic cataplasm* is obtained. (FORM. HÔP. FR.)

CORROBORANT SUPPOSITORY. (CADET DE GASSICOURT)

R. Oak Bark, } of each, one part.
Root of Tormantil, }
Honey, two parts.
Pulverize, and mix.

EXTRACT OF OAK BARK.

Extractum Corticis Quercus. (DUBL.; CAST. BORUS.; BRUGNATELLI)

R. Oak Bark, one pound.
Pure Water, eight pounds.

Boil down to one half, and express; evaporate the strained decoction slowly to the proper consistence. (DUBL.)

R. Oak Bark, one part.
Water, six parts.

Boil for two hours, and strain through a sieve; boil the residue with fresh water, mix the strained decoctions, and evaporate. (BRUGNATELLI)

R. Oak Bark, in powder, one pound.
Boiling Water, twelve pounds.

Digest for twenty-four hours, frequently stirring; then evaporate the liquor slowly to the reduction of one third: filter when cold, and evaporate it to the consistence of an extract. (CAST. BORUS.)

TINCTURE OF OAK BARK.

Tinctura Corticis Quercus. (AUSTR.)

R. Oak Bark, in coarse powder, two ounces.
Alcohol (0.910), one pound.

Digest, and filter.—It is astringent.

DECOCTION OF OAK BARK.

Decoctum Quercūs seu Quercūs Roboris seu Corticis Quercūs. (LOND.; EDINB.; CAST. BORUS.; NIEMANN)

Decoctum Quercūs. (LOND.; NIEMANN)

R. Oak Bark, one ounce.
Water, two pints.

Boil down to one pint, and strain.

Decoctum Quercūs Roboris. (EDINB.)

R. Oak Bark, one ounce.
Water, two pounds and a half.

Boil down to one pound, and strain.

CAST. BORUS. prescribes two ounces of bark, and sixteen ounces of water; to be boiled down to twelve.

It is astringent, tonic, and febrifuge, and is recommended in intermittent fevers.—Dose, a cupful, every three hours.

ASTRINGENT FOMENTATION. (FORM. HÔP. FR.)

R. Oak Bark, one ounce.
Water, a sufficient quantity
to obtain a pound and a half of strained decoction.

ASTRINGENT INJECTION. (SWEDIAUR)

R. Oak Bark or Leaves, one or two ounces.
Water, one pound and a half.

Boil down to one pound.

To be injected into the urethra three or four times a day, in obstinate chronic gonorrhœa.

It may also be prepared with half an ounce of nutgalls in powder.

ASTRINGENT FOMENTATION.

Fomentum quercinum. (AUGUSTIN; RATIER)

R. Oak Bark, half an ounce.
Water, three pounds.

Boil down to one half, and add

Chamomile Flowers, one ounce.

Infuse for half an hour, and add when strained

Vinegar, four ounces.

Mix. (AUGUSTIN)

R. Tops of Milfoil, one ounce.
Root of Avens, } of each, half an ounce.
Oak Bark, }
Lime Water, twenty-four ounces.

Boil down to twelve ounces, and add

Alcohol, three ounces.

Mix. (AUGUSTIN)

R. Root of Cinquefoil, one ounce.
Oak Bark, in powder, } of each,
Red Roses, } six drams.
Water, one pound.

Boil, and strain. (RATIER)

ASTRINGENT MILK. (SAUNDERS)

R. Oak Bark, } of each,
Pomegranate Bark, } two drams.
Root of Tormentil, }
Spring Water, } of each, one pound.
Cow's Milk, }

Boil down to one half, adding towards the end,

Cinnamon, in powder, two drams.

Strain.

DECOCTION OF OAK BARK WITH ALUM.

Decoctum Corticis Quercūs cum Alumine, Fotus adstringens. (MIL. COPEN.; SAUNDERS; SWEDIAUR)

R. Oak Bark, two ounces.
Water, thirty-six ounces.

Boil down to one half; then add to the expressed liquor

Alum, three drams.

Dissolve. (MIL. COPEN.)

R. Oak Bark, one pound.
Pomegranate Flowers, four ounces.
Spring Water, six pounds.

Boil down to one half, and add to the strained liquor,

Alum, six drams.

Dissolve. (SAUNDERS)

DECOCTION OF OAK BARK WITH VINEGAR.

Decoctum Corticis Quercūs cum Aceto, Fotus anti-septicus. (MIL. COPEN.; FULD.; HAMB.)

R. Oak Bark, one ounce.
Herb of Wormwood, two ounces.
Water, thirty-six ounces.

Boil till reduced to twenty ounces; add to the expressed liquor,

Vinegar, four ounces.

Mix. (MIL. COPEN.)

R. Oak Bark, two ounces.
Vinegar, thirty-six ounces.

Boil, adding towards the end,

Herb of Rue, } of each,
Scordium, } three drams.
Muriate of Ammonia, half an ounce.

Strain. (FULD.)

HAMB. prescribes one ounce of oak bark, two pounds of vinegar, one handful each, of rue and of scordium, and half an ounce of muriate of ammonia to be dissolved in the pound and a half of strained liquor.

ASTRINGENT BEER.

Cerevisia quercina. (VAN MONS)

- R. Acorns, one part.
Strong Beer, sixteen parts.

Infuse in a water-bath for several days, and strain.

ASTRINGENT EMULSION. (BORIES)

- R. Acorns, peeled, } of each, six.
Sweet Almonds, }
Infusion of Lesser Mullein, six ounces.
Syrup of Ground Ivy, one ounce.

It is recommended in diabetes.—To be taken in the evening at bed time.

ASTRINGENT ELECTUARY. (BORIES)

- R. Conserve of Hips, six ounces.
Acorns, in powder, three ounces.
Prepared Red Coral, } of each,
Catechu, } one ounce and
a half.
Cinnamon, three drams.
Syrup of Red Roses, a sufficient quantity.

Mix.—It is recommended in diarrhoea and chronic dysentery.—Dose, one dram, every four hours.

TORRIFIED ACORNS.

Glandes seu Nuces Quercus tostæ. (BAVAR. ; LIPP. ; SAX.)

- R. Acorns, any quantity.

Remove from them the cups and rind, cut the kernels each into four parts, and dry them on a sieve placed before a fire; then torrify them in an iron pan, constantly stirring, till the pieces become brittle and of a brown-yellow colour. When cold, reduce them to powder by grinding.

INFUSION OF ACORNS.

Infusum Glandium Quercus tostorum.
(NIEMANN)

- R. Torrified Acorns, half an ounce.
Boiling Water, six or eight ounces.

Infuse, and strain.—It is recommended in obstructions of the mesentery, and in atrophy.—The infusion may be rendered more agreeable by the addition of a dram of chocolate.

ACORN COFFEE.

Coffea Glandium. (AUGUSTIN)

- R. Torrified Acorns, six drams.
Burnt Coffee, two drams.
Water, a sufficient quantity.

Boil for a short time, and strain.

This drink, taken with milk and sugar, is reputed useful in scrofula and rickets.—Dose, two or three cupfuls daily.

QUINA, CHININA, KININUM.—QUININE.

FERR. ; HENRI ; MAGENDIE.

- R. Yellow Cinchona, any quantity.

Treat it with alcohol; wash the extract obtained, with a solution of potass, till this comes off colourless: wash the residue with distilled water to free it from potass, and dissolve it in hydrochloric acid; filter the liquor, and boil it in thrice its weight of magnesia; filter again, and evaporate to dryness. Treat the residue with alcohol, evaporate the tincture, and preserve the residue. (MAGENDIE)

- R. Yellow Cinchona, one hundred parts.
Water, five hundred parts.
Sulphuric Acid, fifty parts.

Boil for two hours, and express. Repeat this operation twice or thrice; then mix the liquors, and when the mixture is cold, throw

slaked lime into it, by portions and in fine powder, till it has become sensibly alkalized and a kind of reddish coagulum is formed in the liquor. Let the deposit drip on a linen cloth, dry it in a stove, and having reduced it to powder, digest it at several times in alcohol (32 degrees); decant, or filter, and distil the tinctures, treat the residue with hot acidulated water and with a small quantity of animal charcoal: filter the liquor, and add to it when cold, a slight excess of ammonia. Collect, wash, and slowly dry the precipitate. (HENRI)

Quinine, which was discovered by PELLÉTIER and CAVENTOU, exists only in the yellow cinchona; it is mixed with cinchonine in the red, but the grey contains none of it. Seven hundred parts of water are required to dissolve it. It is not employed in medicine.

SULPHATE OF QUININE. (FERR.; BORIES;
FEE; HENRI)

1. *Process by Pelletier and Caventou.*

R. Alcoholic Extract of Yellow Cin-
chona, any quantity.

Treat it, with heat, with water acidulated with hydrochloric acid; boil the acid liquor with an excess of subcarbonate of magnesia to perfect discolouration; then filter, collect the deposit, wash it with cold water, and dry it; then digest it at three or four times with alcohol (36 degrees), distil the tinctures, and combine the residue with sulphuric acid. (FEE)

2. *Process by Henri*

R. Yellow Cinchona, in powder, two pounds.
Water, sixteen pounds.
Sulphuric Acid, two ounces.

Boil for an hour, and strain; add a fresh quantity of water and acid to the residue, and boil, and strain again: repeat this operation a third time. Then mix the liquors together, and saturate, without heat, the acid of the mixture with eight ounces of quick lime in powder. Strain through a silk sieve, set aside, dry the deposit, treat it at thrice with alcohol (36 degrees), and filter. Distil the mixed tinctures to three-fourths, add to the residue water weakly acidulated with sulphuric acid, and filter, with heat, adding a certain quantity of animal charcoal, and collect the crystals which are deposited on cooling. (FERR.; BORIES; FEE; HENRI)

BERNARDET has modified this process in the following manner. Boil the cinchona in the acidulated water, and repeat the operation with the residue and fresh acidulated water for seven successive times. Treat the boiling mixed decoctions with lime recently slaked and sifted. When cold, wash the deposit, dry it in a stove, pulverize it, and treat it at several times, in a water bath, with alcohol (36 degrees), which is then to be entirely re-distilled. Then heat the residue, dissolve it in alcohol, and mix the solution with water strongly acidulated with sulphuric acid, applying the heat of a water-bath. Finally filter, decolorize with animal charcoal, filter again, and evaporate.

PELLETIER and CAVENTOU found that a hundred weight of cinchona produced two pounds one dram and thirty grains of sulphate of quinine; according to which proportion, one pound yields two drams sixty-six grains and one-sixteenth, and one ounce, thirteen grains and two-sixteenths. But, as the sulphate of quinine is composed of nine parts and nine-tenths of sulphuric

acid and ninety parts and one tenth of quinine, it therefore results, the ordinary dose of cinchona being two drams, that a patient takes three grains and nine thirty-seconds of sulphate of quinine or three grains and a half of quinine, taking into account a small quantity of cinchonine which may be mixed with it. Four grains of the salt are therefore more than the equivalent of the dose which is usually prescribed when the cinchona is used. Consequently, those who have given thirty-six grains and seventy-two grains of the sulphate, have given what is equal to twenty-seven drams of cinchona in the first case, and fifty-four in the second. It is difficult to imagine that any one could order such enormous doses of cinchona, much more think of repeating them several times in succession.

The sulphate of quinine is given in moistened water; the patient should drink a cup of chamomile tea afterwards.

ANTI-PERIODIC POWDER. (PIERQUIN)

R. Sulphate of Quinine, six grains.
Subcarbonate of Magnesia, thirty-two grains.

Mix.

PELLETIER'S DENTIFRICE POWDER. (FEE; HENRI)

R. Sulphate of Quinine, four grains.
Prepared Coral, one ounce.
Carmine Lac, eight grains.
Essence of Myrrh, two drops.

Mix.

FEBRIFUGE POTION. (BORIES; ELLIS)

R. Sulphate of Quinine, ten grains.
Simple Syrup, one ounce.
Water, three ounces,

acidulated with

Sulphuric Acid, half a dram.

Mix. (BORIES)

R. Sulphate of Quinine, twelve grains.
Sugar, } of each, one dram.
Gum Arabic, }
Cinnamon Water, two ounces and a half.

Mix.—Dose, a tea spoonful, every hour. (ELLIS)

R. Sulphate of Quinine, twelve grains.
Aromatic Sulphuric Acid, ten drops.
Sugar, one dram.
Cinnamon Water, two ounces and a half.

Mix.—To be taken as the preceding. (ELLIS)

SYRUP OF SULPHATE OF QUININE.

Sirop de Quinine. (FERR.; BORIES; CADET DE GASSICOURT; FEE; HENRI; MAGENDIE; RATIER)

R. Sulphate of Quinine, sixty-four grains.
Dissolve it in a little water acidulated with a few drops of sulphuric acid, and add
Warm Simple Syrup, two pounds.
Mix.

COMPOUND SYRUP OF SULPHATE OF QUININE. (BORIES)

R. Coffee, roasted and ground, four ounces.
Boiling Water, two pounds four ounces.
Infuse in a covered vessel; strain when cold, and add
Sugar, four pounds.
Dissolve in a water-bath, and add further
Sulphate of Quinine, dissolved in a little water slightly acidulated with sulphuric acid, sixty grains.
Mix.

TINCTURE OF SULPHATE OF QUININE. (FERR.; BORIES; FEE; MAGENDIE; PIERQUIN; RATIER)

R. Sulphate of Quinine, six grains.
Alcohol (34 degrees), one ounce.
Dissolve. (FERR.; BORIES; FEE; MAGENDIE; PIERQUIN)
RATIER prescribes two scruples of sulphate and one pound of alcohol.

WINE OF SULPHATE OF QUININE. (FERR.; BORIES; CADET DE GASSICOURT; FEE; MAGENDIE; PIERQUIN; RATIER)

R. Sulphate of Quinine, twelve grains.
Madeira Wine, two pounds.

Dissolve. (FERR.; BORIES; FEE; MAGENDIE; PIERQUIN; RATIER)

CADET DE GASSICOURT prescribes eight grains of sulphate.

R. Wine, one pound and a half.
Tincture of Quinine, two ounces.
Mix. (MAGENDIE)

EMMENAGOGUE BOLUSES.

Boli incitantes emmenagogi. (BRERA)

R. Black Oxide of Iron, thirty grains.
Sulphate of Quinine, from six to twelve grains.
Powdered Roses, } of each,
Extract of Horehound, } one dram and a half.
Make eight boluses.—Dose, one every two, three or four hours.

FEBRIFUGE BOLUSES.

Boli anti-febriles. (BRERA)

R. Sulphate of Quinine, three grains.
Crumb of Bread, a sufficient quantity to make one bolus.—Dose, four, six, or eight daily, at intervals of two hours.
R. Sulphate of Quinine, twelve grains.
Opium, two grains.
Honey, a sufficient quantity.
Make four boluses.—Dose, one every four hours.

TONIC PILLS. (ELLIS)

R. Sulphate of Quinine, ten grains.
Conserve of Roses, a sufficient quantity.
Make ten pills.—Dose, one every hour.

RADIX LOPEZIANA.—LOPEZ ROOT.

SYNONIME :—Racine de Jean Lopez.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; GALL.; FERR.; FULD.; WIRTEM.; AINSIE; MURRAY and GMELIN; SPIELMANN.

This root is in rather large, ligneous pieces, the wood of which is of a yellowish-white colour, compact, veined, inodorous, and very bitter. The bark consists of an inner stratum, which is reddish, bitter, and styptic, and of an outer one which is grain-

ed, pulverulent, and of a pale yellow colour; the latter is probably derived from some cryptogamic growth.

This bark comes from Africa, but we are ignorant of the plant which furnishes it. Is it the *Morus Indica*? By a singular error, FERR. assigns it to the *Lopezia racemosa*, CAV., of Mexico.

It is excitant and tonic, and is prescribed in chronic diarrhoea.

Dose, from six to twenty grains of the powder, three or four times a day.

ASTRINGENT SYRUP. (SAINTE-MARIE)

R. Lopez Root, cut small, one ounce.
Boiling Water, two pounds.

Digest in hot ashes for forty-eight hours; then boil slowly till reduced to one half. Strain this and dissolve in it,

White Sugar, one pound,
and boil to a syrup.—Dose, six spoonfuls, in the course of the day.

ASTRINGENT ELECTUARY. (SAINTE-MARIE)

R. Lopez Root, two drams.

Conserve of Red Roses, } of each,
Hips, } two ounces.
Syrup of Gum, a sufficient quantity.

Mix.—Dose, a tea-spoonful, twice or thrice a day.

ASTRINGENT TINCTURE. (SAINTE-MARIE)

R. Lopez Root, one dram and a half.
Alcohol, two ounces and a half.

Digest and filter.

Dose, a tea-spoonful, thrice a day, in any proper vehicle.

RANUNCULUS.

Six species of this genus of plants are used in medicine :

1. *Ranunculus acris*, LINN. — Upright Crow-foot.

SYNONIMES : — Renoncule âcre, FR. ; Scharfer Hahnenfuss, GERM.

BRUNS. ; HISP. ; GALL. ; GUIBOUT ; MURRAY and GMELIN ; SPIELMANN.

A very common plant of Europe. (*Polyandria Polygynia*, LINN. ; *Ranunculaceæ*, JUSS. ; *Fig. Flore Medic. VI. 294*)

The herb and flowers are employed.

The herb (*Herba Ranunculi seu Ranunculi acris seu pratensis*) consists of leaves divided into three lobes, and these again much subdivided ; the superior are linear.

The flowers are yellow.

2. *Ranunculus aquatilis*, LINN.

SYNONIMES : — Renoncule aquatique, FR. ; Wasserhahnenfuss, Wasserhähnlein, GERM.

HISP.

A plant of Europe. (*Fig. CEd. Fl. Dan. t. 376*)

The herb and flowers are employed.

The herb consists of leaves, some finely cut into capillary segments, others entire and peltate. It has an acrid taste.

The flowers are white.

3. *Ranunculus bulbosus*, LINN. — Bulbous Crow-foot ; Butter Cup.

SYNONIMES : — Renoncule bulbeuse, FR. ; Knolliger Hahnenfuss, GERM. ; Knollartige ranonkel, DUT. ; Solvifear, SWED.

GALL. ; GUIBOUT ; MURRAY and GMELIN.

A common plant of Europe. (*Fig. CEd. Fl. Dan. t. 351*)

The root and herb are employed.

The root (*Bulbus Ranunculi bulbosi*) is simple, round, tuberculous, and white. The herb consists of a straight stem and of compound leaves.

The whole plant is extremely acrid.

4. *Ranunculus sceleratus*, LINN. — Marsh Crow-foot.

SYNONIMES : — Renoncule des marais, FR. ; Giftiger Hahnenfuss, Froscheppich, Feigblatterneppich, GERM. ; Puggepeber, DAN. ; Ranunculo mulvado, SPAN. ; Waterhaanevoet, DUT.

GALL. ; COXE ; GUIBOUT ; MURRAY and GMELIN.

A plant of Europe, common in marshy places. (*Fig. Blackw. Herb. t. 259*)

The herb (*Herba Ranunculi palustris seu aquatici*) is employed. It consists of a glabrous, fistulous, and very branching stem, and of caulinary, petiolated, rounded glabrous leaves, half cut into three incised or crenelated lobes, smooth, and of a light-green colour, and into caulinary leaves cut more deeply, narrower, linear, and digitate or palmate.

This species is one of the most acrid and irritant poisons ; it is considered by some authors as the *Herba Sardoa* of SALLUST, from the name of which we derive the designation of *sardonic* given to the apparent laughter which is produced by the spasmodic contraction of the muscles of the mouth and cheeks, which the use of this plant excites.

5. *Ranunculus Flammula*, LINN.—Lesser Spearwort.

SYNONIMES :—Pétite douve, FR.; Kleiner Sumpfhahnenfuss, GERM.; Liden gaul hanefod, DAN.; Kleen egeikolen, DUT.; Ält gräs, SWED.

GALL.; GUIBOURT; MURRAY and GMELIN.

A common plant of Europe. (*Fig. Zorn, Ic. pl. t. 326*)

The herb (*Herba Flammula*) is employed. It consists of a smooth stem, furnished with very entire oval, lanceolated, petiolated leaves. Its taste is extremely acrid and burning.

It is irritant. The Swedish peasants apply it to the inner side of the wrist in intermittent fevers.

6. *Ranunculus Ficaria*, LINN.—Pilewort.

SYNONIMES :—Pétite Chélidoine, FR.; Feigwarzenkraut, Scharbockskraut, Schmalzsternblume, GERM.; Speenkruid, DUT.

BRUNS.; GALL.; GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant which grows in all parts of Europe. (*Fig. Céd. Fl. Dan. t. 499*)

The root (*Radix Ficariae seu Chelidonii minoris*) is employed. It consists of small oblong, and fleshy tubercles, adherent by fibres. It has an acrid and biting taste.

It is irritant; vesicant externally, and sometimes internally.

RAPHANUS.—RADISH.

Two plants of this name are used in medicine:

1. *Raphanus sativus*, LINN.—Turnip Radish.

SYNONIMES :—Raifort cultivé, Radis noir, FR.; Rettig, Ratich, Reddick, GERM.; Reddike, DAN.; Rabano, SPAN.; Tamme radys, DUT.; Rafano, ITAL.; Rabao, PORT.; Rættika, SWED.

AMST.; BRUNS.; DAN.; HISP.; GALL.; FERR.; PALAT.; WIRTEM.; HERBIP.; BERGIUS; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

This plant is cultivated every where. It is said to be originally from China. (*Tetradynamia Siliquosa*, LINN.; *Cruciferae*, JUSS.; *Fig. Flore Medic. IV. 292*)

The root (*Radix Raphani nigri seu hor-tensis*) is large, orbicular, black without and white within; its substance is firm and succulent. It has a strong, peculiar odour, and a very acrid taste, which is particularly distinct in the rind.

There are several varieties of this species produced by culture.

2. *Raphanus rusticus* (*Cochlearia Armoracia*), LINN.—Broad Horse-radish.

SYNONIMES :—Raifort sauvage, Cran, FR.; Meerrettig, Meerrettigsscharbockheil, Kreen, Kren, GERM.; Kren, BOHEM.; Peberrod, DAN.; Rabano, Marvisiro, SPAN.; Meeradys, DUT.; Armorraccio, Peperello bobrasso, ITAL.; Chrzan, POL.; Rabao rusticano, PORT.; Chren, RUSS.; Pepparrot, SWED.

VOL. II.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; SAUNDERS.; SPIELMANN; ZARDA.

A plant common to various parts of Europe. (*Tetradynamia Siliquosa*, LINN.; *Cruciferae*, JUSS.; *Fig. Zorn, Ic. pl. t. 457*)

The root (*Radix Armoraciae seu Raphani rusticani seu rustici seu marini*) is employed. It is long, cylindrical, thick, yellow without, white and fibrous within.

It has a very penetrating odour, and an acrid, pungent, and bitter taste.

It contains starch and a volatile oil, of which sulphur appears to be one of the principal constituents.

It is an energetic stimulant, and is stomachic, and anti-scorbutic.—Dose, of the juice, half a dram, in milk or wine.

IRRITANT PEDILUVIUM.

Pediluvium ex Armoracia. (AUGUSTIN)

R. Hot Water, four pounds.
Horse-radish, bruised,
from two to four ounces.

Mix.—SCHLEGEL recommends it to restore suppressed perspiration of the feet.

3 Q

ANTI-SCORBUTIC SPECIES.

Species contra Scorbutum. (PALAT.)

- R. Herb of Scurvy-grass, }
 ——— Water-cress, } of each,
 ——— Brooklime, } half an
 ——— White Horehound, } ounce.
 ——— Fumitory,
 ——— Buckbean,
 Root of Horse-radish, } half a pound.
 ——— Polypody, } of each,
 ——— Sarsaparilla, } three ounces.
 Seeds of Anise, } six drams.
- Cut, bruise, and mix.

DIURETIC JUICE.

Succus Armoracæ compositus. (SUEC.; NIEMANN.)

- R. Juice of Horse-radish, } four ounces.
 Oxymel of Squill, } one ounce.
- Mix. (NIEMANN)
- R. Juice of Horse-radish, } one ounce.
 Spirit from grain, } six ounces.
 Vinegar, } two ounces.
- Mix. (SUEC.)
- Dose, two spoonfuls.

ANTI-CATARRHAL DRINK. (PIERQUIN; SAINTE-MARIE)

- R. Expressed Juice of boiled } of each,
 Radishes, } eight
 Water, } ounces.
 Honey, } two ounces.
- Mix.—To be taken in the course of the day, by cupfuls.

HONEY OF HORSE-RADISH.

Mel Armoracæ. (SUEC.; NIEMANN; VAN MONS)

- R. Juice of Horse-radish, } three ounces.
 Honey, } one pound.
- Mix without heat. (SUEC.)

VAN MONS prescribes one part of juice and four parts of honey.—NIEMANN—one of juice and two of honey, evaporated to a proper consistence over a slow fire.

WATER OF HORSE-RADISH.

Aqua Armoracæ. (HENRI)

- R. Fresh Root of Horse-radish, } one part.
 Water, } five parts.
- Distil two parts.

DIURETIC WATER. (SARD.; CADET-DE GASSICOURT; PIERQUIN)

- R. Root of Horse-radish, }
 ——— Parsley, } of each,
 ——— Saxifrage, } one pound.
 Juniper, } six ounces.

- Lemon Juice, } of each,
 Water of Rest-harrow, } three pounds.
- Macerate for twenty-four hours, and distil in a water-bath. (SARD.)
- R. Juice of Leek, }
 ——— Onion, } of each,
 ——— Horse-radish, } two pounds.
 ——— Wall Pellitory, } of each,
 ——— Lemon, } half a pound.
- Digest for three days, and distil slowly. (CADET DE GASSICOURT; PIERQUIN)
- Dose, one ounce, morning and evening.

SPIRIT OF HORSE-RADISH.

Spiritus Raphani rusticani. (HASS.; SARD.)

- R. Fresh Root of Horse-radish, } of each,
 Juice of Horse-radish, } six pounds.
 Water, }
 Alcohol, } two pounds.
- Distil one half. Add to the product
 Fresh Root of Horse-radish, } three pounds.
- Distil in a water-bath. (SARD.)
- R. Root of Horse-radish, } of each,
 Water, } six parts.
 Alcohol, } four parts.
- Distil two parts. (HASS.)

COMPOUND SPIRIT OF HORSE-RADISH.

Spiritus Armoracæ compositus. (LOND.; DUBL.; FULD.; NIEMANN)

- R. Fresh Horse-radish, } of each,
 sliced, } one
 Dried Orange Peel, } pound.
 Nutmegs, bruised, } half an ounce.
 Proof Spirit, } one gallon.
 Water, } a sufficient quantity

to prevent empyreuma. Macerate for twenty-four hours, and distil one gallon slowly. (LOND.; DUBL.; NIEMANN)

- R. Root of Horse-radish, } three pounds.
 Alcohol, } four pounds.

Distil four pounds by the heat of a water-bath, and add

- Water of Pimenta, } eight pounds.

Mix. (FULD.)

Dose, a spoonful.

ANTI-SCORBUTIC PTISAN. (DISP. CHA. PARIS)

- R. Compound Spirit of Horse-radish, } one ounce.
 Infusion of Cichory, }
 one pound and a half.
- Mix.

ANTI-SCORBUTIC BEER.

Cerevisia Armoracæ. (CADET DE GASSICOURT; HUFELAND; VAN MONS)

- R. Horse-radish, five ounces.
Beer, two pounds.

Infuse for twenty-four hours, and add to the strained liquor,

Simple Syrup, one ounce.

Mix. (HUFELAND)

- R. New Beer, twenty pounds.
Suspend in it, during its fermentation, a bag containing

Betony,	}	of each, one handful.
Sage,		
Agrimony,		
Scurvy-grass,		
Lesser Wormwood,	}	of each, four ounces.
Root of Elicampane,		
—— Horse-radish,		

Strain at the end of two or three days. (CADET DE GASSICOURT)

- R. Oranges, three.

Express the juice, and pour it into

Strong Beer, sixty-four parts.

Macerate, without heat, in the mixture

Fresh Leaves of Scurvy-	}	of each, one part.
grass,		
—— Cress,		
—— Sage,	}	of each, one half part.
Seeds of Mustard,		
Root of Horse-radish,		

At the end of several days, strain with slight expression. (VAN MONS)

Dose, two or three glassfuls, daily, in scurvy and disorders of the loins, bladder, and liver.

COMPOUND SPRUCE BEER.

Bièrre Sapinette. (CADET DE GASSICOURT)

- R. Beer, four pounds.
Root of Horse-radish, two ounces.
Leaves of Scurvy-grass, one ounce and a half.

Buds of Spruce Fir, one ounce.

Infuse, and express.—Dose, from two to four ounces daily.

VINEGAR OF HORSE-RADISH.

Acetum Cochleariæ Armoracæ. (SWEDIAUR)

- R. Root of Horse-radish, one ounce.
Vinegar, one pound.

Macerate for eight days, and filter.

It is principally employed as a cosmetic for the removal of freckles; it is also used against scorbutic ulcers of the mouth.

EMETIC LIQUOR. (PIERQUIN)

- R. Green Walnuts, } of each,
Root of Horse-radish, } two pounds.
Vinegar, three pounds.

Digest in a moderate heat for five or six days, and strain.—Dose, from half an ounce to three ounces.

COMPOUND TINCTURE OF HORSE-RADISH.

Tinctura Raphani composita. (FORM. HÔP. FR.; NIEMANN)

- R. Fresh Root of Horse-radish, two ounces.
Dried Orange Peel, one ounce.
Spirit of Scurvy-grass, four pounds.

Digest for several days, and express.

It is excitant, and is recommended in scurvy.—Dose, a spoonful.

ANTI-SCORBUTIC GARGLE. (RATIER)

- R. Bitter Infusion, six ounces.
Tincture of Horse-radish, half an ounce.
Honey of Roses, two ounces.

Mix.

WINE OF HORSE-RADISH.

Vinum Armoracæ. (SUEC.)

- R. Root of Horse-radish, half a pound.
White Wine, eight pounds.

Digest for four days, and strain.

It is excitant and anti-scorbutic.

COMPOUND WINE OF HORSE-RADISH.

Vinum Cochleariæ compositum. (FORM. HÔP. FR.; NIEMANN; SWEDIAUR)

- R. Root of Horse-radish, one ounce.
Cinnamon, } of each,
Seeds of Black Mus- } one ounce and
tard, a half.
White Wine, two pounds and a half.

Macerate for three days, frequently agitating, and afterwards decant the clear part. (SWEDIAUR)

- R. Compound Tincture of Horse-radish, one ounce.
White Wine, two pounds.

Mix. (FORM. HÔP. FR.; NIEMANN)

It is excitant, and is recommended in scurvy, derangement of the digestive functions from weakness of the stomach, and in paralysis.—Dose, five or six spoonfuls twice a day. It may be mixed with water when taken.

ANTI-SCORBUTIC WINE. (GALL.; SARD.; SPIELMANN; VAN MONS)

- R. Root of Horse-radish, three parts.
Seeds of Mustard, one part and a half.

Herb of Water Cress, } of each,
 — Scurvy-grass, } one part.
 — Brooklime, }

Spirit of Scurvy-grass, one part and a half.
 Rhenish Wine, ninety-six parts.

Infuse for eight days; then express, and filter. (VAN MONS)

R. Fresh Root of Horse-radish, one ounce.

— Leaves of Scurvy-grass, } of each,
 — Water } half an ounce.
 — Cress, }
 — Buckbean, }

Seeds of Mustard, }
 Muriate of Ammonia, } two drams.
 White Wine } two pounds.

Digest for thirty-six hours, occasionally agitating; strain with slight expression, and add to the liquor

Spirit of Scurvy-grass, half an ounce.

Mix. (GALL.)

R. Fresh Root of Horse-radish, twelve ounces.

— Burdock, } five ounces.
 — Leaves of Scurvy-grass, }
 — Cress, } of each,
 — Brooklime, } six ounces.
 — Fumitory, }

Seeds of Mustard, }
 Muriate of Ammonia, } three ounces.

White Wine, three hundred and eighty ounces.

Infuse for eight days; then express strongly, and filter. (SPELMANN)

R. Fresh Root of Horse-radish, four ounces.

— Leaves of Scurvy-grass, } of each,
 — Buckbean, } two handfuls.
 — Sage, } one handful.
 White Wine, } six pounds.

Digest for twenty-four hours at a moderate heat, and strain. (SAUNDERS)

Dose, one ounce, and more.

DIURETIC WINE. (SAUNDERS)

R. Root of Horse-radish, } of each,
 — Iris, } six ounces.
 White Wine, a sufficient quantity

to obtain, after half an hour's infusion, two pounds of strained liquor; add to this

Syrup of Fennel, } of each,
 — Juniper, } two ounces.

Mix.

ANTI-SCORBUTIC SYRUP.

Syrupus anti-scorbuticus seu anti-scorbuticus compositus seu Cochleariae officinalis com-

positus seu de Armoracia compositus.
 (ANT.; BELG.; GALL.; SPIELMANN;
 SWEDIAUR; VAN MONS)

R. Bruised Cinnamon, one part.

Fresh Leaves of Scurvy-grass, } of each,
 — Brooklime, } thirty parts.
 — Cress, }

Seeds of Mustard, thirty-five parts.

Fresh Root of Horse-radish, forty-five parts.

Orange Juice, ninety parts.

Red Wine, two hundred and sixty parts.

Infuse for twenty-four hours at a moderate temperature; when cold, express the liquor, and dissolve in it, at a gentle heat,

White Sugar, four hundred and eighty parts.

Strain the syrup through a sieve. (VAN MONS)

R. Fresh Leaves of Scurvy-grass, } of each,
 — Buckbean, } one pound.
 — Water }
 — Cress, }
 Root of Horse-radish, }
 Bitter Oranges, sliced, }
 Cinnamon, bruised, }

White Wine, one ounce and a half.
 four pounds.

Macerate for two days in a tin cucurbit covered with its capital and well luted. Distil in a water-bath one pound of liquor, with which and

White Sugar, two pounds,
 make a syrup in a covered vessel placed in a water-bath. Strain, without expression, what remains in the cucurbit. Set apart the liquor, and finally decant, and make a syrup with

White Sugar, two pounds.
 Clarify the syrup, and when it is only warm mix it with the preceding. (GALL.)

R. Fresh Herb of Scurvy-grass, } of each,
 — Pepper-wort, } five pounds.
 — Water Cress, }
 Root of Horse-radish, ten pounds.
 Cinnamon, in powder, six ounces.
 Bitter Oranges, sliced, one pound.
 Alcohol (20 degrees), two pounds.
 Distilled Water, eight pounds.

Mix in an alembic, and distil slowly five pounds of liquor, with which and

White Sugar, ten pounds,
 make a syrup by the heat of a water-bath. Express strongly what remains in the cucurbit. Add to the strained liquor

White Sugar, ten pounds.

Clarify, boil slowly to the consistence of syrup, and when cold, mix it with the preceding. (ANT.; BELG.)

SWEDIAUR prescribes, instead of the mixture of alcohol and water, ten pounds of white wine. He also recommends the second syrup to be made with the liquor expressed from the residue, ten pounds of water, and twenty pounds of sugar.

R. Juice of Scurvy-grass, } of each,
 ——— Brooklime, } twelve
 ——— Water Cress, } ounces.
 ——— Horse-radish Root, }
 ——— Bitter Oranges, twenty ounces.
 Fresh Peel of Bitter Orange, one ounce.

Digest without heat in a covered vessel, occasionally agitating, for about twelve hours. Filter, and add to the liquor

White Sugar, twice the quantity.

When dissolved, add further,

Spirit of Scurvy-grass, } of each,
 Cinnamon Water, } one ounce.

Set aside, and decant. (SPIELMANN)

BITTER ALTERATIVE SYRUP. (CADET DE GASSICOURT)

R. White Wine, sixteen pounds.
 Leaves of Scurvy-grass, three pounds.
 ——— Buckbean, } of each,
 Fresh Root of Horse- } two pounds.
 radish, }
 Dried Root of Gentian, one pound.
 Macerate for several days; then

R. Of this Wine, one pound.
 Sugar, one pound twelve ounces.

Dissolve slowly, and to each pound of this syrup add one ounce of another syrup prepared with spirit of scurvy-grass.

INFUSION OF HORSE-RADISH.

Infusum Armoracæ seu Armoracæ compositum seu Seminum Sinapeos seu Raphani. (LOND.; DUBL.; AMER.; GENEV.; HERBIP.; COXE; NIEMANN)

Infusum Armoracæ Compositum. (LOND.; DUBL.)

R. Fresh Root of Horse- } of each,
 radish, sliced, } one
 Mustard Seeds, bruised, } ounce.
 Boiling Water, one pint.

Macerate for two hours in a lightly covered vessel, and strain. Add

Compound Spirit of Horse-radish, one fluid ounce.

Mix.

NIEMANN gives the same formula.—AMER.; and COXE also give the same, omitting the spirit.

R. Root of Horse-radish, }
 Seeds of Mustard, } of each,
 Conserve of Scurvy- } one ounce.
 grass, }
 Boiling Water, two pounds.

Infuse for twenty-four hours in a covered vessel, and strain. (HERBIP.; NIEMANN)

R. Root of Horse-radish, } of each,
 Seeds of Mustard, } one ounce.
 Cannella Alba, three drams.
 Boiling Water, one pound and a half.

Infuse for two hours; then strain, and add

Water of Anise, two ounces.

Mix. (GENEV.)

It is excitant, and is recommended in scurvy, palsy, dropsy, and intermittent fevers.—Dose, from one to three ounces.

SYRUP OF HORSE-RADISH.

Syrupus Armoracæ seu Raphani rusticani. (FULD.; LIPP.; SUEC.; WIRTEM.; NIEMANN)

R. Recently expressed Juice of Horse-radish, any quantity.
 White Sugar, a sufficient quantity.

Make a syrup. (SUEC.)

FULD.; and LIPP. prescribe an equal part of each.

R. Root of Radish, } of each,
 Water, } an equal part.

Triturate together, and express. Add to nine parts of the juice

White Sugar, seven parts.
 Make a syrup in a close vessel. (VAN MONS)

CULLEN recommended one dram of horse-radish root, rasped, to be infused in two ounces of hot water, and twice the quantity of sugar to be added to the strained liquor.—Dose, from one to three drams.

ANTI-PARALYTIC INFUSION.

Infusum paralyticum seu ad Paralysis. (SAUNDERS; SWEDIAUR)

R. Root of Horse-radish, } of each,
 Seeds of Mustard, } one ounce and
 a half.
 Boiling Water, two pounds.

Macerate for two hours; then strain, and add

Alcohol, two ounces.

Mix.

It is excitant, and is recommended in paralysis.—Dose, four ounces, twice a day.

ANTI-HYDROPIC INFUSION.

Infusum anti-hydropicum seu ad Hydricos.
(SAUNDERS; SWEDIAUR)

- R. Bitter Infusion, } of each,
Anti-paralytic Infusion, } one pound.
Mix.—Dose, four ounces, twice a day.

ANTI-SCORBUTIC DECOCTION.

Decoctum anti-scorbuticum. (GALL.; BORRIES; CADET DE GASSICOURT; SPIELMANN)

- R. Root of Horse-radish, two ounces.
Water, eighty ounces.

Boil for a quarter of an hour, and add

- Herb of Brooklime, }
Cress, } of each,
Scurvy-grass, } two handfuls.
Sorrel, }

Boil a little, strain, and add

- Vinegar, four ounces.

Mix. (SPIELMANN)

- R. Root of Horse-radish, } of each,
Parelle, } half an ounce.
Buckbean, }
Water Cress, } of each, one pinch.
Scurvy-grass, }
Brooklime, }
Water, a sufficient quantity

to produce three glassfuls of decoction.
Add to it when strained

PORTAL'S Anti-scorbutic Syrup
without Sublimate, one ounce.

Mix. (BORIES)

- R. Root of Patience, } of each,
Burdock, } one ounce.
Water, four pounds.

Boil for twenty minutes, and add

- Root of Horse-radish, one ounce.
Leaves of Scurvy-grass, } of each,
Buckbean, } six drams.
Water Cress, }

Infuse for ten minutes, and strain.
(GALL.; CADET DE GASSICOURT)

It is recommended in tetters, scurvy, and disorders of the lymphatic system.—Dose, several glassfuls in the course of the day.

DIURETIC APOZEM. (PIERQUIN; SAINTE-MARIE)

- R. Root of Horse-radish, from two to four drams.
Juniper, half a pinch.
White Horehound, a large pinch.
Boiling Water, three large glassfuls.

Macerate for an hour, and strain when cold.

To be taken at thrice, in the morning, fasting; adding to each dose, a spoonful of a mixture of four ounces of syrup of squill with four ounces of vinous syrup of seneka.

RESINA ELEMI.—ELEMI.

SYNONIMES:—Elemi, FR.; and DAN.; Elemiharz, Wildäelbaumharz, GERM.; Goma de lemon, SPAN.

LOND.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY; GMELIN; PARIS; FIDERIT; SPIELMANN; ZARDA.

There are two sorts of resins which bear this name.

1. *Elemi orientale*; Oriental or Ethiopian Elemi.—This is of a yellowish colour approaching to green, hard externally, soft internally, of a strong and disagreeable odour resembling that of fennel, and of an aromatic taste.

2. *Elemi occidentale*; Occidental or Spurious Elemi.—This is in round cakes, of a pale yellow colour, semi-transparent, some-

times variegated with white and yellow grains, of a strong and agreeable odour, and of a bitter taste.

The first was attributed to the *Gardenia gummifera*, LINN., a tree of Ceylon (*Fig. Thunb. Diss. No. 4, tab. 2*), and the second to the *Amyris Elemifera*, LINN. (*Fig. Plum. Ic. I. 100*), a tree of New Spain. GUIBOUT has ascertained that both are originally from America, that the resin of the amyris is now no longer met with in commerce, and, that what is, is derived from the *Iceia Icacariba*, CAND.

It is excitant.

OINTMENT OF ELEMI.

Unguentum Elemi. (DUBL.)

- R. Elemi Resin, one pound.
White Wax, half a pound.
Prepared Lard, four pounds.

Form an ointment, and strain through a sieve while hot.

COMPOUND OINTMENT OF ELEMI.

Unguentum Elemi compositum.
(LOND.)

R.	Elemi,	one pound.
	Turpentine,	ten ounces.
	Prepared Suet,	two pounds.

Olive Oil, two fluid ounces.

Melt the resin with the suet; then remove the mixture from the fire, add the turpentine and oil immediately, and strain the whole through a linen cloth.

RHAMNUS.

Two species of this genus are employed :

1. *Rhamnus Frangula*, LINN.—Black Alder Tree.

SYNONIMES :—Bourgene, FR.; Faulbaum, GERM.; Tærsttræ, DAN.; Arraclan, SPAN.; Verelboom, DUT.; Frangula, ITAL.; Kruszyna, POL.; Frangulina, PORT.; Brakvæd, SWED.

GALL.; FORM. HÔP. FR.; GENEV.; WIRTEM.; HERBIP.; BERGIUS; MURRAY and GMELIN; SAUNDERS; SPIELMANN.

A shrub of the north of Europe. (*Pentandria Monogynia*, LINN.; *Rhamneæ*, JUSS.; *Fig. Nouv. Duh.* III. t. 15)

The bark (*Cortex Frangule* seu *Alni nigre baceifera*) is employed. It is of a deep green colour spotted with white. It appears yellow when the epidermis is removed, and of a brownish-red after desiccation. It is without odour, but has a bitterish taste.

It is laxative.—The berries have also the same property.

2. *Rhamnus cathartica*, LINN.—Buckthorn.

SYNONIMES :—Nerprun, FR.; Gemeine Kreuzdorn, Færbebeere, GERM.; Psygahody, Rzesetlak, BOHEM.; Korsbær, Vrietorn, DAN.; Ramno catartico, SPAN.; Purgeerende wegedoorn, DUT.; Spino cervino, ITAL.; Szaklak kzewia, POL.; Espinha cervina, PORT.; Getappel, Saftrænt, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HANN.; LIPP.; LUSIT.; SAX.; SUEC.; WIRTEM.; BORIES; COXE; GUIBOURT; MURRAY and GMELIN; SAUNDERS; SPIELMANN; ZARDA.

A shrub which grows in every part of Europe. (*Pentandria Monogynia*, LINN.; *Rhamneæ*, JUSS.; *Fig. Nouv. Duh.* III. t. 10)

The fruit (*Baccæ Spinæ cervinæ* seu *domesticæ*) is employed. It consists of berries of the size of a pea, round, shining, and blackish, enclosing four grains in the middle of a succulent pulp. They have a disagreeable odour, and a bitter, nauseous, and acrid taste.

They contain, according to VOGEL, a peculiar colouring principle, *Rhamnium*.

It is a very strong purgative, and has been much extolled in dropsy.

JUICE OF BUCKTHORN.

Succus Rhamni cathartici seu *Spinæ cervinæ*.
(GALL.; SARD.)

- R. Ripe Buckthorn Berries, any quantity.

Bruise them with the fingers, and leave the mass to ferment for three or four days; then express, and preserve the juice in bottles, under a stratum of oil.

Dose, from two to four drams.

ROB OF BUCKTHORN.

Rob Spinæ cervinæ. (AUSTR.; GALL.; HERBIP.)

- R. Ripe Buckthorn Berries, any quantity.

Bruise them, and leave the mass for three days till fermentation has commenced; then express strongly, strain the juice through flannel, decant at the end of three or four hours, and evaporate slowly to the consistence of an extract. (GALL.)

- R. Juice of Buckthorn Berries, one pound.

Boil it to the consistence of honey, adding towards the end

Sugar, two ounces.

AUSTR. prescribes one part of sugar and nine parts of juice.

Dose, from a scruple to a dram and a half.

SYRUP OF BUCKTHORN.

Syrupus Rhamni seu *Rhamni cathartici* seu *domesticus* seu *de Spina cervina*. (LOND.;

DUBL.; EDINB.; AMER.; AMST.; ANT.;
BELG.; BRUNS.; GALL.; FORM. HOP.
FR.; GENEV.; HANN.; LIPP.; LUSIT.;
PALAT.; SARD.; WIRTEM.; HERBIP.;
COXE; SPIELMANN; SWEDIAUR; VAN
MONS)

Syrupus Rhamni cathartici. (EDINB.)

- R. Clarified Juice of ripe Buck-
thorn Berries, two parts.
Refined Sugar, one part.

Boil to a syrup.

GALL. prescribes an equal part of each.—
AMER.; ANT.; LUSIT.; and COXE—four
parts of juice and two of sugar.—HERBIP.
—six of juice and four of sugar.—AMST.;
and BELG.—sixteen ounces of juice and two
pounds and a half of sugar.—FORM. HOP.
FR.—three parts of juice and four of sugar.
—SWEDIAUR—seven pounds and a half of
juice and three and a half of sugar.—GE-
NEV.—six pounds of juice and three of sugar.
—LIPP.—one part of juice and one and a
half of sugar.

SWEDIAUR allows a pound of tincture of
pimenta to be added.

Syrupus Rhamni.

- R. Fresh Juice of Buckthorn
Berries, four pints
Ginger, sliced, } of each,
Pimenta Berries, } half
bruised, } an ounce.
Refined Sugar,
three pounds and a half.

Set apart the juice that the fecu-
lencies may subside, and strain it.
To one pint of this juice add the
ginger and pimenta, macerate in a
gentle heat for four hours, and strain;
boil the remainder of the juice down
to a pint and a half; then mix the
liquors, add the sugar, and boil to a
syrup. (LOND.)

- R. Fresh Juice of Buckthorn
Berries, two pints and a half.
Ginger, sliced, } of each,
Pimenta Berries, } two drams.
bruised, }

Refined Sugar,
three pounds and a half.

Set apart the juice for three days
that the fæces may subside, and strain.
Add to one pint and a half of it the
ginger and pimenta, and strain after
four hours' maceration in a gentle
heat: then add the remainder of the
juice and the sugar, and boil to a
syrup. (DUBL.)

- R. Ripe Buckthorn Berries,
sixteen pounds.

Express the juice; cover the residue with
boiling water, macerate it for twenty-four
hours, and strain with expression. Mix the
liquors, and add

White Sugar, eight pounds.

Clarify with white of egg, and boil to the
consistence of syrup. When this is cold,
add to it, triturated with sugar,

Oil of Cinnamon, }
Cloves, } of each, one scruple,
Mace, }

Mix well. (VAN MONS)

- R. Depurated Juice of Buckthorn Berries,
twenty ounces.
White Sugar, thirty-two ounces.

Boil to the proper consistence, adding
towards the end, contained in a bag,

Anise, }
Mastic, } of each, three drams.
Ginger, }
Cinnamon, } of each,
Cloves, } one dram and a half.

Preserve. (BRUNS.; HANN.; WIRTEM.
SPIELMANN)

PALAT. prescribes one pound of juice,
two pounds of sugar, and one dram each,
of the aromatics.—SARD.—an equal part
each, of the juice and of sugar, two drams
each, of anise and of mastic, and one dram
each, of ginger, of cinnamon, and of
cloves.

Dose, from half an ounce to one ounce.

PURGATIVE POTION. (DISP. CHA. PARIS)

- R. Decoction of Prunes, four ounces.
Syrup of Buckthorn, one ounce.
Mix.

RHEUM.—RHUBARB.

SYNONIMES :—*Rhabarbarum*.—Rhubarbe, FR.; Rhabarber, DAN.; GERM.; DUT.; and SWED.; Rawend, ARAB.; Rabarbara, BOHEM.; Tayhuam, Hay-hoang, CHIN.; Ta hoam, COCHIN.; Rewund chini, DUK.; Ruibarbo, PORT.; and SPAN.; Reywan chinie, HIND.; Rabarbaro, ITAL.; Schara modo, MO.; Reywand, PERS.; Korzen Rabarbarowy, POL.; Variatu kalung, TAM.; Dschumza, THIB.

The pharmacopœias principally designate under this name the two following substances :

A.

Official Rhubarb.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

There are several kinds of official rhubarb :

1. Indian or Chinese Rhubarb; *Rheum seu Rhabarbarum verum seu Indicum seu Sinense seu Chinense seu Danicum*.—This is in round pieces, of a dull yellow colour externally, of a compact texture, brownish-yellow and veined internally, and of a bitter taste. It grits between the teeth, and tinges the saliva with orange-yellow. The powder is between a fawn and an orange colour. It comes from China by Canton.

2. Russian, Tartarian, or Bucharian Rhubarb; *Rheum seu Rhabarbarum Tartaricum seu Moscoviticum seu Russicum*. This is in flattish, irregular, angular pieces, pierced with large holes, of a pure yellow colour externally, less compact than the preceding, presenting red and white veins, very distinct and irregular. It has a very decided odour, and a bitter and astringent taste. It tinges the saliva with saffron-yellow, and grits between the teeth. Its powder is of a purer yellow than that of the preceding. This is the kind which is most esteemed, though the Chinese is also valued highly when perfectly sound. This also comes from China, but by Kiachta in Siberia.

VOL. II.

GUIBOURT thinks these two varieties are furnished by the *Rheum palmatum*, LINN. (*Fig. Flore Medic. VI. 297*), which grows in the environs of Kian-Sin and Chan-Sin in Chinese Tartary. PFAFF does not reject this origin, but still thinks it doubtful.

Another kind is also met with in commerce :

3. European Rhubarb; *Rheum seu Rhubarbarum Britannicum seu Gallicum seu nostras*; it comes from the *Rheum undulatum*, *compactum*, and also *palmatum*, cultivated in various countries of Europe, in England, France, and Germany.—This differs from the two preceding, by its characters both physical and chemical, and presents generally, with a few slight modifications, those of the second variety of the *Rheum Rhaponticum*, which is equally indigenous.

Oriental rhubarb contains, according to HENRI, a peculiar colouring principle, *Rhabarbarine* or *Caphopieicrite*, a fixed oil, supermalate of lime, gum, starch, oxalate of lime, which makes one third of its weight, &c. It is from the presence of this last mentioned salt that it grits between the teeth. HORNEMANN considers *rhabarbarine* to be an impure substance; he attributes another peculiar principle to rhubarb, which he calls *Rheumine*, and which approximates to the acids although it is very different to the *Rheumatic acid* admitted by HENDERSON, but of which the existence has not been fully established. He has also discovered in it a saccharine substance the presence of which has been also ascertained by MEISSNER.

Rhubarb is tonic or purgative, according to the dose; it is tonic in that of from five to ten grains, and purgative in that of from eighteen to twenty-four or more.

B.

Pontic Rhubarb; Rheum Rhaponticum,—LINN.

SYNONIMES :—*Rhapontic*, FR.; and DUT.; *Pontischer Rhabarber*, GERM.; *Reupontykum*, Zemezluc wetssj, BOHEM.; *Rapontico*, ITAL.; PORT.; and SPAN.

BRUNS.; HISP.; FERR.; FORM. HÔP. FR.; WIRTEM.; HERBIP.; BELG.; BRUNS.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant originally of the south of Russia, and cultivated in France. (*Enneandria Trigynia*, LINN.; *Polygoneæ*, JUSS.; *Fig. Flore Medic. VI. 296*)

3 R

The root (*Radix Rhapontici veri seu Rhabarbari Rhapontici*) is employed. It is presented in two different forms; one of the varieties is as large as the first, or less, of a ligneous appearance, of a reddish-grey colour externally, veined, when broken transversely, with red and white, forming very close striae, radiating from the centre to the circumference; it has a very astringent and mucilaginous taste, and tinges the saliva yellow, but does not grit between the teeth; its odour is analogous to that of rhubarb but is more disagreeable; its powder is yellow with a reddish tinge. The other variety is about three or four inches in length and two or three in thickness, of a less ligneous appearance, of a light yellow colour, purer or less reddish; its fracture presents a radiated appearance, its taste is astringent, and mucilaginous, its odour is similar to that of rhubarb, and it does not grit between the teeth.

It contains, according to HENRI, much more starch, and less oxalate of lime than rhubarb, and also more colouring principle, which in this is red. HORNEMANN has discovered a peculiar substance in it which he calls *Rhaponticine*.

It is tonic and laxative.

The following is the only formula of the Pontic rhubarb:

EXTRACT OF PONTIC RHUBARB.

Extractum Rhapontici. (FORM. HÔP. FR.)

R. Dried Root of Pontic Rhubarb, one part.
Cold Water, four parts.

Infuse for twenty-four hours, then strain with slight expression, and infuse the residue in

Cold Water, two parts.

Strain. Mix the infusions, and evaporate in a water-bath to the consistence of an extract.

I. Preparations which contain Rhubarb in substance.

TORRIFIED RHUBARB.

Rhabarbarum tostum. (HISP.; LUSIT.; SARD.)

R. Rhubarb, in powder, any quantity.

Expose it in a shallow iron pan to the heat of a moderate fire, constantly stirring it with a spatula, till it has acquired a brown colour; then let it cool, and preserve it.

DIATURBITH SPECIES WITH RHUBARB. (PALAT.; WIRTEM.; VAN MONS)

R. Rhubarb, } of each, two ounces.
Indian Jalap, }

Hermodactyls, one ounce.
Sulphuretted Scammony, } of each,
Supertartrate of Potass, } half an
Seeds of Violet, } ounce.

Make a powder, and add,

Oil of Cinnamon, ten drops.

Preserve. (PALAT.; WIRTEM.)

R. Cinnamon, one part.
Ginger, } of each, three parts.
Seeds of Anise, }
Scammony, four parts.
Hermodactyls, } of each, eight parts.
Indian Jalap, }
Rhubarb, ten parts.

Make a powder. (VAN MONS)

Dose, from one to two scruples. It is purgative and anthelmintic.

ABSORBENT POWDER. (ELLIS)

R. Prepared Crab's Claws, one dram.
Rhubarb, two scruples.

Mix, and divide into four parcels.—Dose, one or two in the day.

STIMULANT POWDER. (ELLIS)

R. Rhubarb, } of each, one dram.
Ipecacuanha, }

Mix.—It is applied with advantage to indolent ulcers.

POWDER OF RHUBARB AND OF SUPERTARTRATE OF POTASS.

Pulvis Rhei cum Tartaro, Pulvis catharticus seu Rhei compositus seu lenitivus seu lenitivus tartarisatus. (MIL. COPEN.; FULD.; HANN.; SAINTE-MARIE; PH. SYPH.)

R. Rhubarb, } of each,
Supertartrate of Potass, } half an ounce.

Make a powder. (MIL. COPEN.)

Dose, a scruple.

PH. SYPH. prescribes twenty grains of rhubarb, and ten of supertartrate, for one dose.—SAINTE-MARIE—an equal part each, of rhubarb and of tartrate of potass.

R. Rhubarb, } of each,
Supertartrate of Potass, } one ounce.
Orange Peel, }

Make a powder. (HANN.)

FULD. prescribes two drams of rhubarb, half an ounce of supertartrate and four scruples of orange peel.

Dose, from half a dram to one dram.

POWDER OF RHUBARB AND OF SULPHATE OF POTASS.

Pulvis Rhei compositus seu anti-hecticus seu purgans. (CAST. BORUS.; SAUNDERS; SWEDIAUR)

R. Rhubarb, six grains.
Sulphate of Potass, twelve grains.

Make a powder. (SPIELMANN)

CAST. BORUS. prescribes one part of rhubarb and three parts of sulphate.

R. Rhubarb, half a dram.
Sulphate of Potass, ten grains.
Scammony, eight grains.
Oil of Fennel, one drop.

Make a powder. (SARD.)

POWDER OF RHUBARB AND OF MAGNESIA.

Pulvis Puerorum epilepticus cum Rheo seu catharticus absorbens seu laxans seu Infantum cum Rheo seu Magnesia compositus seu Rhei salinus seu Rhei cum Magnesia seu digestivus seu e Rheo. (BRUNS; DAN.; MIL. COPEN.; FERR.; GENEV.; LIPP.; SUEC.; ELLIS; NIEMANN; SAUNDERS; SWEDIAUR)

R. Rhubarb, one part.
Subcarbonate of Magnesia, three parts.

Make a powder. (SAUNDERS; SWEDIAUR)

SUEC. prescribes an equal part of each.—GENEV.—one of rhubarb and four of magnesia.—FERR.—one of rhubarb and eight of subcarbonate of magnesia.

R. Rhubarb, one scruple.
Magnesia, half a dram.
Oil of Cinnamon, one drop.

Mix.—To be taken in syrup. (ELLIS)

R. Rhubarb, half an ounce.
Subcarbonate of Magnesia, } of each,
Fennel, } two
White Sugar, } drams.

Mix. (SUEC.)

BRUNS. prescribes one ounce each, of rhubarb and of magnesia, two drams of fennel, and half an ounce of sugar.

R. Rhubarb, } of each,
Subcarbonate of Magnesia, } an equal
Oleosaccharum of Fennel, } part.

Make a powder. (DAN.; MIL. COPEN.)

Dose, one scruple.

R. Rhubarb, half an ounce.
Miseltote, } of each, six drams.
Liquorice, }
Subcarbonate of Magnesia, two ounces and a half.

Make a powder. (LIPP.; NIEMANN)

Dose, from a scruple to half a dram.

SAPONACEOUS POWDER OF RHUBARB.

Pulvis ad Tormina Infantum. (AUGUSTIN; SPIELMANN)

R. Rhubarb, half a dram.
Soap, one dram.

Crab's eyes, half an ounce.
Anise, } of each, one scruple.
Cumin, }
Saffron, four grains.
White Sugar, three drams.

Mix. (SPIELMANN)

R. Medicinal Soap, from ten to twenty grains.
Subcarbonate of Mag- } of each,
nesia, } half a dram.
Sugar, }
Rhubarb, }

Mix. (AUGUSTIN)

BOLUSES OF RHUBARB.

Boli e Rheo. (FORM. HÔP. FR.; SAUNDERS; SWEDIAUR)

R. Rhubarb, half a dram.
Simple Syrup, a sufficient quantity.

Mix. (SAUNDERS; SWEDIAUR)

Five grains of protochloruret of mercury or ten grains of jalap may be added if required.

R. Rhubarb, one part.
Sulphate of Soda, two parts.

Make a bolus with a little water. (FORM. HÔP. FR.)

ASTRINGENT BOLUS.

Bolus ad Diarrhœam seu e Rheo sedativus. (SAUNDERS; SWEDIAUR)

R. Rhubarb, fifteen grains.
Diascordium, half a dram.
Simple Syrup, a sufficient quantity.

Make a bolus. (SAUNDERS)

R. Rhubarb, twelve grains.
Aromatic Powder, three grains.
Tincture of Opium, twenty drops.
Syrup, a sufficient quantity.

Mix. (SWEDIAUR)

RESOLVENT BOLUSES.

Boli resolventes et purgantes. (BRERA)

R. Rhubarb, one dram.
Venice Soap, one dram and a half.
Rob of Elder, a sufficient quantity,
to make eight boluses.—Dose, one every two hours.

PILLS OF RHUBARB.

Pilulæ Rhei. (LIPP.; SAUNDERS)

R. Rhubarb, any quantity.
Infusion of Rhubarb, a sufficient quantity.

Make a pilular mass.—Dose, from ten to thirty grains.

RESOLVENT PILLS.

Pilule resolventes cum Rheo seu a Rheo acetico seu ephracticæ. (LIPP.; CAST. BORUS.; WURCEE.; ELLIS; SAINTE-MARIE)

- R. Rhubarb, } of each,
 Acetate of Soda, } two drams.
 Inspissated Ox-gall, }
 Gum Arabic, half a dram.

Make pills of two grains each. (CAST. BORUS.; WURCEE.)

- R. Rhubarb, one dram and a half.
 Venice Soap, four drams.
 Spirit of Turpentine, a sufficient quantity.

Make pills of five grains each. (SAINTE-MARIE)

- R. Inspissated Ox-gall, } of each,
 Gum Ammoniac, } one ounce.
 Rhubarb, }

Make pills of two grains each. (LIPP.)

- R. Rhubarb, one dram,
 White Soap, ten grains.
 Water, a sufficient quantity.

Make fifteen pills.—Dose, four. (ELLIS)

- R. Rhubarb, one dram and a half.
 Sulphate of Iron, half a dram.
 White Soap, two scruples.
 Water, a sufficient quantity.

Make forty pills. (ELLIS)

BITTER PILLS. (VAN MONS)

- R. Extract of Gentian, } of each,
 White Soap, } one part.
 Rhubarb, one part and a half.

Make a pilular mass.

ABSORBENT PILLS.

Pilule Rhubarbarinæ antacidæ. (SWEDIAUR)

- R. Rhubarb, } of each, two drams.
 Magnesia, }
 Extract of Gentian, a sufficient quantity.

Make pills of five grains each.

STIMULANT PILLS.

Pilule roborantes ad Blennorrhæam. (PH. SYPH.; SAUNDERS)

- R. Rhubarb, one dram.
 Turpentine, two drams.

Make thirty-six pills. (SAUNDERS)

- R. Rhubarb, } of each, one ounce.
 Gum Ammoniac, }
 Copaiba, a sufficient quantity.

Make pills of four grains each. (PH. SYPH.)

STOMACHIC ELECTUARY.

Electuarium stomachicum. (SWEDIAUR)

- R. Rhubarb, one dram and a half.
 Root of Arum, } of each,
 Aromatic Powder, } one dram.
 Black Oxide of Iron, one scruple.
 Conserve of Orange Peel, two ounces.
 Syrup of Orange Peel, a sufficient quantity.

Mix.—Dose, the volume of a nutmeg, twice a day.

ANTI-DYSENTERIC ELECTUARY. (CADET DE GASSICOURT)

- R. Confection of Catechu, two ounces.
 LOCATELLI's Balsam, one ounce.
 Rhubarb, half an ounce.
 Syrup of Marshmallows, a sufficient quantity.

Mix.—Dose, the same as the preceding.

PURGATIVE MARMALADE.

Conditum purgativum, Pullicula saponacea. (SAUNDERS; SAINTE-MARIE; SPIELMANN)

- R. Sulphate of Potass, one dram.
 Supertartrate of Potass, half an ounce.
 Rhubarb, one dram and a half.
 Pulp of Tamarinds, two ounces.

Mix. (SAUNDERS).—Dose, a tea spoonful.

- R. Soap, half a dram.
 Mint Water, three drams.

Dissolve, and add

- Yolk of Egg, one dram.
 Subcarbonate of Magnesia,

- Rhubarb, two scruples.
 Syrup of Marshmallows, } of each,
 Chamomile, } two drams.

Mix. (SPIELMANN)

It is recommended in infants' colic.

- R. Rhubarb, } of each,
 Supertartrate of Potass, } two drams.
 Pulp of Prunes, eight ounces.
 Syrup of Quince, a sufficient quantity.

Mix. (SAINTE-MARIE)

Dose, four tea spoonfuls, daily.

VERMIFUGE RAISINS.

Passula anthelmintica. (PARIS)

- R. Corinth Raisins, half a pound.
 Rhubarb, in a bag, half an ounce.
 Corsican Moss, one ounce.
 Purslain Water, four ounces.

Put the mixture into an oven after the bread has been taken out, and leave the vessel in it till the raisins are swollen and soft.

TROCHISCS OF RHUBARB.

Tabellæ de Rheo seu Diarrhodon cum Rhubarbaro. (GALL.; SARD.)

- R. Rhubarb, half an ounce.
Mucilage of Tragacanth, made
with cinnamon water,
a sufficient quantity.

Make trochiscs of twelve grains each.
(GALL.)

- R. Diarrhodon Powder, } of each, one
Rhubarb, } ounce and a half.
White Sugar, one pound.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs. (SARD.)

- R. Rhubarb, } of each,
Supertartrate of Potass, } two drams.
Fresh Orange Peel, half a dram.
Sugar, dissolved in Orange-flower
Water, four ounces.

Make trochiscs. (BRUNS.)

PURGATIVE POTION.

Haustus catharticus pro Infantibus. (PIERQUIN; SWEDIAUR)

- R. Rhubarb, twelve grains.
Cichory Water, three ounces.
Syrup of Cichory, one ounce.

Mix.—To be taken by spoonfuls; for a
child of nine or ten years old. (PIERQUIN)

- R. Rhubarb, eight grains.
Subcarbonate of Lime, ten grains.
Water, three drams.
Cinnamon Water, two drams.
Simple Syrup, one dram.

Mix.—To be taken as the preceding.
(SWEDIAUR)

PURGATIVE AND VERMIFUGE POTION.
(PIERQUIN)

- R. Cinnabar, ten grains.
Resin of Guaiac, } of each, five grains.
Jalap, }
Rhubarb, one dram.
Deutoxide of Antimony, fifteen grains.
Compound Syrup of Cichory,
six drams.
Water of Cichory, two ounces.

Mix.

BITTER MIXTURE.

Mixtura amara cum Rheo. (SWEDIAUR)

- R. Sulphate of Potass, half an ounce.
Rhubarb, } of each,
Extract of Gentian, } two drams.
Peppermint Water, eight ounces.

Dissolve, and mix. To be taken by
spoonfuls.

II. Preparations which contain the active
principle of Rhubarb without the vehicle
employed to extract it.

AQUEOUS EXTRACT OF RHUBARB.

Extractum Rhei seu Rhubarbari. (AMST.;
BAVAR.; BRUNS.; DAN.; HISP.; GALL.;
FERR.; GENEV.; HANN.; LIPP.; OL-
DEN.; PALAT.; BORUS.; SARD.; SUEC.;
WIRTEM.; VAN MONS)

- R. Rhubarb, sliced, any quantity.
Cold Water, a sufficient quantity,

to cover it. At the end of twenty-four
hours, add more water so as to make up
thrice the weight of the rhubarb, and strain
without expression; pour upon the residue
an equal weight of water, and express
slightly. Mix the two strained liquors, cla-
rify with white of egg, and evaporate to the
consistence of thick extract. (VAN MONS)

BAVAR. directs one part of rhubarb to
be macerated in five parts of cold water, at
twice, and each time for twenty-four hours.

- R. Rhubarb, one pound.
Spring Water, six pounds.

Macerate for four hours in a warm place,
and then boil a little. Express, and eva-
porate the liquor. (BRUNS.; HISP.; PA-
LAT.; WIRTEM.; FERR.)

- R. Rhubarb, one pound.
Boiling Water, sixteen pounds.

Digest for twenty-four hours; then set
aside and strain; pour upon the residue

Boiling Water, eight pounds,
and strain again at the end of twenty-four
hours. Mix, and evaporate the liquors to
the proper consistence. (AMST.; SARD.)

- R. Rhubarb, one part.
Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and
express; boil the residue with

Water, four parts,
and strain again. Mix and decant the li-
quors, and finally evaporate. (DAN.;
HANN.; LIPP.; OLDEN.; BORUS.)

SUEC. prescribes three or four successive
ebullitions with three parts of water each
time.

- R. Rhubarb, one pound.
Cold Water, four pounds.

Macerate for twenty-four hours, occasi-
onally stirring, and strain; pour fresh wa-
ter upon the rhubarb, macerate in the same
manner, and express lightly. Mix the li-
quors, strain again, and evaporate to the
proper consistence in a water-bath. (GALL.)

- R. Rhubarb, two pounds
Cold Water, twelve pounds.

Macerate for twenty-four hours, occasi-
onally stirring, and strain; macerate the
residue in

Cold Water, eight pounds,
and strain again. Mix the liquor, and evaporate in a water-bath to the consistence of syrup; then pour the liquor into dishes to about the depth of two lines, and evaporate to dryness, in an oven or water-bath, then detach the scales. (GALL.)

This is the *Essential Salt of Rhubarb*.
Dose, from fifteen to twenty grains.

ALCOHOLIC EXTRACT OF RHUBARB.

Extractum Rhei seu Rhubarbari alcoholicum. (LOND.; DUBL.; FULD.; LUSIT.; POL.; SAX.; HERBIP.; COXE)

Extractum Rhei. (LOND.; DUBL.)

R. Rhubarb, in Powder, one pound.
Proof Spirit, one pint.
Water, seven pints.

Macerate for four days in a moderate heat; then strain the liquor, and set it apart that the feculencies may subside. Pour off the clear liquor, and evaporate it to the proper consistence.

R. Rhubarb, sliced, } of each,
Alcohol, } two pounds.
Water, } nine pounds.

Digest for twelve hours, and express; distil off the alcohol, and evaporate the residue to the consistence of an extract. (POL.)

FULD.; and HERBIP. prescribe one pound of rhubarb, one of alcohol, and eight pounds of water; to be digested for three days.—SAX.—two parts of rhubarb, three of alcohol, and nine of water; to be digested for twenty-four hours.—COXE gives the same formula as LOND.

R. Rhubarb, in powder, one pound.
Alcohol, four pounds.

Digest for four days, and decant; boil the residue with

Water, ten pounds,
till reduced to two: strain the decoction and evaporate; then distil the tincture to the consistence of honey, mix it with the evaporated decoction, and evaporate the mixture, constantly stirring, to the proper consistence. (LUSIT.)

It is to be taken at the same dose as the preceding.

COMPOUND EXTRACT OF RHUBARB.

Extractum Rhei compositum seu catholicum seu panchymagogum seu cnico-pharmacum.

(DAN.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; NIEMANN; PIDERIT)

R. Extract of Rhubarb, three ounces.
Aloes, one ounce.

Soften them in alcohol in a vessel placed in a water-bath, and evaporate to the consistence of pilular mass. (POL.)

R. Extract of Rhubarb, three ounces.
Aloes, one ounce.

Mix, and add
Soap of Jalap, dissolved in alcohol, one ounce.

Evaporate in a water-bath to a proper consistence. (OLDEN.; BORUS.; SAX.; NIEMANN)

DAN. prescribes two ounces of extract of rhubarb, five drams of extract of aloes, and one ounce of soap of jalap, dissolved in alcohol.

R. Extract of Rhubarb, two ounces.
Aqueous Extract of Aloes, half an ounce.

Extract of Black Hellebore, one dram thirty-six grains.

Mix in a heated mortar with
Resin of Jalap, in powder, two drams twenty-four grains.

Preserve for use. (PIDERIT)

R. Rhubarb, one ounce.
Seeds of Distaff-thistle, four ounces.
Polypody of the Oak, two ounces.
Ginger, two drams.
Senna, } of each, two
Indian Jalap, } ounces and a half.
Agaric, six drams.
Scammony, ten drams.
Manna, two ounces and a half.

Infuse the agaric, scammony, and jalap, at several times in two pounds of alcohol, in order to exhaust them; then macerate the others in three pounds of water, and boil to the consumption of one third, adding towards the end the residue of the spirituous extraction: then express, dissolve the manna in the liquor, clarify the solution, mix it with the spirituous extract, dissipate the alcohol with a gentle heat, and evaporate to the consistence of soft extract. (PALAT.)

It is purgative.—The dose of the last is from a scruple to half a dram, without manna, and from a dram to four scruples with manna.

III. *Preparations which contain the active principle of Rhubarb with the vehicle employed to extract it.*

A.

Extraction by water.

INFUSION OF RHUBARB.

Infusum Rhei seu Rhubarbari, Tinctura

Rhei aquosa, Anima Rhei. (LOND.; DUBL.; EDINB.; AUSTR.; BAVAR.; GENEV.; HAMB.; LUSIT.; HERBIP.; BRERA; BRUGNATELLI; COKE; DISP. CHA. PARIS; NIEMANN; RATIER)

Infusum Rhei.

- R. Rhubarb, sliced, one dram.
Boiling Water, half a pint.

Macerate for two hours in a lightly covered vessel, and strain. (LOND.; DUBL.)

- R. Rhubarb, sliced, half an ounce.
Boiling Water, eight ounces.
Spirit of Cinnamon, one ounce.

Macerate the rhubarb with the water for twelve hours in a lightly covered vessel; then add the spirit, and strain. (EDINB.)

DISP. CHA. PARIS; and NIEMANN give the same formula as LOND.—BAVAR. prescribes one part of rhubarb and twelve parts of water.—AUSTR.—half an ounce of rhubarb and half a pound of water.—BRUGNATELLI—three ounces of rhubarb and two pounds and a half of water.—RATIER—one ounce of rhubarb and two pounds of water.—HAMB.—one ounce of rhubarb, and a sufficient quantity of water to obtain eight ounces of strained infusion.—BRERA—three drams of rhubarb and six ounces of water. LUSIT.; and COKE give the same formula as EDINB.—HERBIP. prescribes three drams of rhubarb, half a pound of water, and six drams of spirit of cinnamon.

- R. Rhubarb, in powder, one ounce.
Peppermint Water, twelve ounces.
Digest in a gentle heat, and strain. (GENEV.)

It is tonic and slightly purgative; and is recommended in some chronic diseases, and in convalescence from acute affections of which the digestive organs have not been the seat.

TONIC POTION.

Infusum Rhei amarum, Decoctum amaro-aromaticum laxativum. (ANT.; BRERA)

- R. Bitter Decoction, one pound.
Infusion of Rhubarb, four ounces.
Peppermint Water, one ounce.

To be taken gradually. (BRERA)

- R. Rhubarb, three drams.
Orange Peel, half a dram.
Peppermint Water, four ounces.

Mix with the strained liquor,
Extract of Lesser Centaury,
two drams.

To be taken in the course of the day.
(AUGUSTIN)

EXCITANT AND RESOLVENT POTION.

Haustus incitans et resolvens. (BRERA)

- R. Rhubarb, one dram.
Decoction of Cinchona, eight ounces.
Infuse, and add when strained,
Camphor, triturated with Mucilage
of Gum Arabic, half a dram.
Sulphuric Ether, thirty drops.
Syrup of Orange Peel, one ounce.
Mix.—Dose, two spoonfuls every two hours.

LAXATIVE AND SEDATIVE POTION.

Infusio laxativa sedans. (BRERA)

- R. Infusion of Rhubarb, six ounces.
Syrup of Acetate of Morphine, one ounce.

Mix.—To be taken at twice; for irritable subjects.

- R. Infusion of Rhubarb, six ounces.
Peppermint Water, one ounce.
HOFFMANN'S LIQUOR, one scruple.

Mix.—To be taken at four doses.

LAXATIVE AND RESOLVENT POTION.

Potio resolvens et laxans. (BRERA)

- R. Infusion of Rhubarb, eight ounces.
Subcarbonate of Potass, three drams.
Compound Syrup of Cichory, one ounce.

Mix.—To be taken at twice or thrice.

DECOCTION OF RHUBARB.

Decoctum Rhabarbari, Anima Rhabarbari.
(HISP.)

- R. Rhubarb, bruised, two drams.
Water, eight ounces.
Boil a little, and strain.—For one dose.

SYRUP OF RHUBARB.

Syrupus Rhei seu Rhei palmati seu de Rheo seu de Rhabarbaro. (AMER.; ANT.; BATAV.; BAVAR.; BELG.; FENN.; HASS.; LUSIT.; SUEC.; COKE; SWEDIAUR; VAN MONS)

- R. Infusion of Rhubarb, one part.
White Sugar, two parts.

Make a syrup. (FENN.; HASS.; LUSIT.; SUEC.)

- R. Rhubarb, sliced, one part.
Boiling Water, twelve parts.

Infuse for twelve hours at the temperature of from 30 to 40 degrees, and add to one part of strained liquor,

White Sugar, two parts.
 Make a syrup. (AMER.; BAVAR.; COXE)
 R. Rhubarb, sliced, one part.
 Water, eight parts.

Infuse; repeat the infusion in the same quantity of water, mix the strained liquors, and add

White Sugar, fourteen parts.

Clarify with white of egg, and boil to the consistence of syrup. (SWEDIAUR; VAN MONS)

R. Rhubarb, sliced, eight ounces.
 Water, four pounds.

Infuse for twenty-four hours, and to the liquor depurated by repose, add

White Sugar, four pounds.

Add to the residue

Water, four pounds.

Infuse, gradually increasing the heat to ebullition; strain with expression, and add to the liquor

White Sugar, four pounds.

Clarify, evaporate, and mix the syrup, when half cold, with the former. (ANT.)

R. Rhubarb, sliced, three ounces.
 Boiling Water, two pounds.

Macerate for twelve hours, and then boil a little; express, filter, set aside, decant, and add

White Sugar, two pounds.

Make a syrup with a gentle heat. (BATAV.; BELG.)

AROMATIC SYRUP OF RHUBARB.

Syrupus Rhei aromaticus. (AMER.; GALL.; POL.; COXE)

R. Rhubarb, three ounces.
 Cinnamon, six drams.
 Boiling Water, two pounds.

Digest for one night; strain in the morning, and add to twenty ounces of strained infusion

White Sugar, three pounds.

Make a syrup. (POL.)

R. Rhubarb, in powder, ten drams.
 Seeds of Coriander, of each,
 ——— Fennel, } half a dram.
 Water, twelve ounces and a half.

Put the mixture into a bottle which it fills to the neck, plunge it into a water-bath, and let this boil for twenty minutes; then remove the bath from the fire, let it gradually cool, withdraw the bottle, decant the liquor at the end of twenty-four hours, strain, and add to it

White Sugar, twenty-five ounces.

Dissolve in a water-bath. (GALL.)

R. Rhubarb, five drams.

Cloves, } of each, half an ounce.
 Cinnamon, }
 Nutmegs, two.
 Water, one pint.

Digest; then evaporate to one half, and add

Sugar, one pound.
 Proof Spirit, half a pint.

Boil a little. (AMER.; COXE)

COMPOUND SYRUP OF RHUBARB.

Syrupus de Rheo seu de Cichorio compositus.
 (AMER.; AMST.; AUSTR.; BRUNS.;
 HISP.; GALL.; GENEV.; HANN.; PA-
 LAT.; SARD.; WIRTEM.; HERBIP.; RO-
 RIES; PIDERIT)

R. Fresh Herb of Cichory, four ounces.
 Water, four pounds.

Boil to the consumption of one third, adding towards the end

Rhubarb, sliced, four ounces.

Express, and add to the liquor

White Sugar, four pounds.

Clarify, and boil to the consistence of syrup. (HISP.)

R. Fresh Herb and Root of Cichory, two ounces.
 Water, three pounds.

Boil to the consumption of one third, and add

Rhubarb, sliced, three ounces.

Digest for twelve hours, and strain after a slight boiling; to the strained liquor, depurated by repose, add

White Sugar, three pounds.

Boil to a syrup. (AMST.)

R. Depurated Juice of Cichory, two pounds.
 Rhubarb, sliced, two ounces.

Digest for one night in warm ashes; express in the morning, and digest the residue with

Boiling Water, two pounds.

After a short boiling, express; mix the liquors together, and add

Simple Syrup, three pounds.

Make a syrup. (SAUNDERS)

R. Root of Cichory, six ounces.
 Leaves of Cichory, nine ounces.
 ——— Fumitory, } of each,
 ——— Hounds- } three ounces.
 tongue,
 Winter Cherries, two ounces.
 Water, fourteen ounces.

Boil down to twelve pounds; strain, and add to the liquor

White Sugar, five pounds.

Clarify, and make a thick syrup over a slow fire. On the other hand,

R. Rhubarb, bruised, six ounces.

Yellow Sanders, } of each,
Cinnamon, } half an ounce.
Water, } eight pounds.

Infuse for twenty-four hours in a covered vessel, and express slightly. Mix the liquor with the preceding syrup, boil for a few moments, and pour the syrup while hot upon a bag containing

Cinnamon, } of each,
Yellow Sanders, } half an ounce.

Infuse for six hours, and decant. (GALL.)

Bosson recommends the leaves and woods to be boiled for a quarter of an hour, and then infused for two hours, in ten pounds of water; the liquor to be strained and decanted, and with five pounds of sugar, a syrup made at 25 degrees; on the other hand, an infusion to be made on the previous evening with the rhubarb, the aromatic substances, three pounds of water, and two drams of subcarbonate of potass; this to be strained without expression, and the liquor mixed with the syrup, which is to be boiled again, if necessary, and the boiling mixture poured upon the sanders and cinnamon.

R. Root of Cichory, } of each,
Dandelion, } four ounces.
Spring Water, a sufficient quantity

to leave three pounds after proper boiling. Then

R. Rhubarb, sliced, six ounces.
Boiling Water, three pounds.

Macerate for twenty-four hours; express, mix the liquor with the preceding decoction, and boil with

White Sugar, six pounds.

Pour the boiling syrup upon

Yellow Sanders, } of each,
Cinnamon, } half an ounce.

Express when cold, and strain through flannel. (GENEV.)

R. Root of Butcher's-broom, } of each,
Asparagus, } three ounces.

Leaves of Cichory, }
Pimpernel, } of each,
Maidenhair, } one handful.
Agrimony, }
Chervil, }

Water, three pounds.

Boil, and infuse in the decoction

Senna, three ounces.
Rhubarb, bruised, one ounce.

Indian Jalap, } of each,
Hermodyctyls, } six drams.

Flowers of Marigold, } of each,
Tamarisk, } half a handful.
Elder, }

Add to the decanted liquor

White Sugar, one pound and a half.

Boil to a syrup. (BRERA)

VOL. II.

R. Herb of Cichory, one ounce.
Root of Cichory, three ounces.
Rhubarb, four ounces.
Subcarbonate of Potass, half a dram.
Spring Water, eight pounds.

Boil down to six pounds; express strongly, and add to the liquor

White Sugar, six pounds.

Clarify, and boil to the consistence of syrup. (AUST.)

R. Root of Cichory, two ounces and a half.
Spring Water, one pound and a half.

Boil down to one pound, and add

Subcarbonate of Potass, twenty grains.

Boil slowly, and add to the expressed liquor

White Sugar, sixteen ounces

Clarify, strain, and add

Spirituous Cinnamon Water, one ounce.

Mix. (BRUNS.; HANN.; PALAT.; WIRTEM.)

R. Root of Cichory, three ounces.
Spring Water, three pounds.

Boil to the consumption of one third.

Add

Rhubarb, sliced, two ounces.
Subcarbonate of Potass, half a dram.

Digest for several hours in a warm place; then boil for a few moments. Add to the expressed liquor

Sugar, thirty-two ounces.

Strain the syrup. (HERBIP.)

PIDERIT, who gives the same formula, adds

Spirituous Cinnamon Water, two ounces.

SPIELMANN prescribes two ounces and a half of cichory root, eighteen ounces of water boiled down to twelve, one ounce of rhubarb, one scruple of subcarbonate, and twenty-two ounces of sugar.

ALKALINE AQUEOUS TINCTURE OF RHUBARB.

Tinctura Rhei seu Rhabarbari, Anima Rhei, Infusum Rhei alkalisatum seu cum Subcarbonate Potassae. (AMST.; ANT.; AUST.; BAVAR.; BRUNS.; DAN.; MIL. COPEN.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; BORUS.; SAX.; WIRTEM.; NIEMANN.; SWEDIAUR.; VAN MONS)

R. Simple Infusion of Rhubarb, eleven parts.
Subcarbonate of Potass, one part.

Dissolve. (BAVAR.)

R. Rhubarb, sliced, three parts.
Water, fifty parts.

2 S

Heat to ebullition; then withdraw it from the fire, and add

Subcarbonate of Potass, one part.
Strain when cold. (VAN MONS)

R. Rhubarb, half an ounce.
Subcarbonate of Potass, one dram.
Water, six ounces.

Infuse for two hours at a gradually augmented heat till the liquor has begun to boil; then strain, set aside, and decant. (ANT.; AUSTR.)

DAN. prescribes one ounce of rhubarb, two drams of subcarbonate, and ten ounces of boiling water.—FULD.—one ounce of rhubarb, four scruples of subcarbonate, eight ounces of water, and twenty-four hours' digestion.

R. Rhubarb, two drams.
Subcarbonate of Potass, half a dram.
Water, a sufficient quantity

to obtain four ounces of strained liquor. When it has commenced boiling, add

Simple Syrup, one ounce.

Mix. (SWEDIAUR)

R. Rhubarb, two parts.
Boiling Water, sixteen parts.

Digest for two hours; express slightly, and add

Subcarbonate of Potass, one half part.
Alcohol, one part.

Mix. (LIPP.)

SWEDIAUR prescribes one ounce of rhubarb, three drams of subcarbonate, two ounces of water, twenty-four hours' maceration, and two ounces of spirit of cinnamon.—AMST.—one ounce of rhubarb, six ounces of boiling water, six hours' digestion, four ounces of cinnamon water, and two drams of subcarbonate.—HANN.; OLDEN.; BORUS.; SAX.; and NIEMANN— one ounce and a half of rhubarb, three drams of subcarbonate, twelve ounces of water, twelve hours' maceration, and two ounces of vinous cinnamon water.—MIL. COPEN.—half an ounce of rhubarb, one dram of subcarbonate, four ounces of boiling water, twelve hours' digestion, and one ounce of fennel water.—WIRTEM.—one ounce of rhubarb, two drams of subcarbonate, six ounces of cichory water, six ounces of buglossated cinnamon water, and twenty-four hours' digestion.—PALAT.—one ounce of rhubarb, two drams of subcarbonate, sixteen ounces of cichory water, and twenty-four hours' digestion.—BRUNS.—one ounce of rhubarb, one dram and a half of subcarbonate, three ounces of pure water, and six ounces of cinnamon water.

It is tonic and absorbent; and is recommended in jaundice, hypochondria, and acidity of the primæ viæ.—Dose, from two

drams to one ounce, two or three times a day.

SALINE AQUEOUS TINCTURE OF RHUBARB.

Infusum Rhei salinum seu boraxatum.
(HASS.; POL.; HERBIP.; CADET DE GASSICOURT; FEE; NIEMANN; PIDERIT; SPIELMANN)

R. Rhubarb, bruised, one ounce.
Distilled Water, nine ounces.
Acetate of Potass, one dram.

Infuse for four hours, and filter. (CADET DE GASSICOURT)

R. Vinegar, nine ounces.
Subcarbonate of Potass, a sufficient quantity

to saturate the acid. Digest in the liquor for twelve hours,

Rhubarb, sliced, one ounce.

Filter. (SPIELMANN)

R. Rhubarb, half an ounce.
Tartrate of Potass, two drams.
Boiling Water, half a pound.

Digest for twenty-four hours, and filter. (PIDERIT)

HERBIP. gives the same formula, adding half a scruple of cochineal.—HASS. leaves the choice between tartrate of potass and borated tartar.

R. Rhubarb, one ounce and a half.
Sub-borate of Soda, two drams.
Boiling Water, one pound.

Macerate for twelve hours, and add to ten ounces of the strained liquor

Vinous Cinnamon Water, two ounces.

Mix. (POL.; NIEMANN; FEE)

Dose, one or two spoonfuls, three or four times a day.

ALTERATIVE DECOCTION. (BRUNS.)

R. Root of Rhubarb, } of each,
—— Madder, } three drams.
Subcarbonate of Potass, two drams.
Spring Water, eight pounds.

Boil for an hour. Add to the strained decoction

Decoction of Ginger, three ounces.

Mix.

ROSENSTEIN'S VISCERAL ELIXIR.

Elixir viscerale Rosensteini. (NIEMANN)

R. Rhubarb, sliced, eight ounces.
Subcarbonate of Potass, three ounces.
Cinnamon Water, forty-eight ounces.

Digest for fifteen days in a well-stopped

flask, in a cool place, frequently agitating ; then add

Fresh Orange Peel,	two ounces.
Lesser Cardamoms, }	of each,
Seeds of Fennel, }	half an ounce.

Digest for eight days, with frequent agitation, and express ; pour upon the residue

Decoction of Cichory, six ounces.

Agitate, express, mix the liquor with the former, and dissolve in the mixture

Sugar, twenty-four ounces.

It is tonic.—Dose, from half a spoonful to one spoonful.

ALKALINE SYRUP OF RHUBARB.

Syrupus Rhei alka-lisatus. (DAN. ; FERR. ; FULD. ; HANN. ; LIPP. ; OLDEN. ; BORUS. ; ROSS. ; WIRTEM. ; SWEDIAUR)

R. Alkaline Aqueous Tincture of Rhubarb, one pound.
White Sugar, two pounds.

Boil to a syrup. (FULD.)

LIPP. prescribes ten ounces of the tincture and sixteen of sugar.

R. Rhubarb, four ounces.
Subcarbonate of Potass, two drams.
Boiling Water, two pounds.

Boil over a slow fire, and express. Add to twenty ounces of the liquor

White Sugar, three pounds.
Make a syrup. (DAN.)

R. Rhubarb, three ounces.
Subcarbonate of Potass, two drams.
Boiling Water, two pounds.

Digest for twelve hours, and add to twenty ounces of the strained infusion

White Sugar, three pounds.
Make a syrup. (OLDEN.)

ROSS. prescribes two ounces of rhubarb, one dram and a half of the salt, eighteen ounces of water, and two pounds of sugar.

R. Rhubarb, three ounces.
Cinnamon, three drams.
Subcarbonate of Potass, two drams.
Boiling Water, two pounds.

Digest for twelve hours, and strain. Add to twenty ounces of the strained liquor

White Sugar, three pounds.
Make a syrup. (SWEDIAUR)

HANN. ; and BORUS. prescribe six drams of cinnamon.

R. Rhubarb, two ounces.
Subcarbonate of Potass, one dram.
Juice of Cichory, boiling, three pounds.

Macerate for twelve hours, express, and add to the strained liquor

Sugar, two pounds.
Dissolve. (FERR.)

R. Rhubarb, } of each,
Senna, } one ounce and a half.
Cinnamon, } one dram and a half.
Subcarbonate of Potass, two scruples.
Ginger, half a dram.
Cichory Water, ten ounces.
Rose Water, four ounces.

Digest for three days in a warm place, then boil for an instant, and express. Clarify the liquor with white of egg, and add

Sugar, sixteen ounces.

Make a syrup. (WIRTEM.)

MANNA AND RHUBARB POTION. (FORM. HÔP. FR. ; RATIER)

R. Rhubarb, half an ounce.
Water, four ounces.

Boil for a few minutes ; then dissolve in the strained decoction

Manna, two ounces.

PURGATIVE POTION. (PIERQUIN)

R. Rhubarb, } of each,
Subcarbonate of Potass, } one dram.
Decoction of Wall Pellitory, a sufficient quantity.

Boil, and dissolve in one ounce of the liquor,

Manna, two ounces.

Strain, and add

Compound Syrup of Rhubarb, one ounce.

Mix.

PURGATIVE AND TONIC POTION. (PIERQUIN)

R. Myrobalani Citrini, two drams.
Yellow Sanders, one dram.
Rhubarb, bruised, half a dram.
Manna, two ounces.
Boiling Water, a sufficient quantity.

Strain, and add

Confection of Hyacinth, one dram.

Mix.

CARMINATIVE MIXTURE. (SAINTE-MARIE)

R. Aqueous Tincture of Rhubarb, six ounces.

Mineral Anodyne Liquor, one dram.

Mix.—Dose, a spoonful, three or four times a day.

APERITIVE TINCTURE. (HAMB.)

R. Aqueous Infusion of }
Rhubarb, } of each,
Solution of Acetate of } half an ounce.
Potass, }
HUXHAM'S Antimonial Wine, one dram.

Mix.

B.

Extraction by Wine.

WINE OF RHUBARB.

Tinctura Rhei seu Rhabarbari vinosa, Vinum Rhei seu Rhei palmati aromaticum seu compositum, Elixir Rhei dulce seu compositum. (EDINB.; AMER.; ANT.; BATAV.; BELG.; FENN.; GENEV.; OLDEN.; LUSIT.; SUEC.; CADET DE GASSICOURT; COXE; VAN MONS)

Vinum Rhei. (EDINB.)

R. Rhubarb, sliced, two ounces.
 Canella Bark, bruised, one dram.
 Proof Spirit, two ounces.
 Spanish White Wine, fifteen ounces.

Macerate for seven days, and filter.

AMER.; AMST.; BATAV.; CADET DE GASSICOURT; and COXE give the same formula.

R. Rhubarb, two ounces.
 Cinnamon, one dram.
 White Wine, one pound and a half.

Macerate for eight days, and strain. (LUSIT.)

R. Rhubarb, eight parts.
 Indian Anise, one part.
 Malaga Wine, ninety-six parts.

Digest without heat for eight days, then express lightly, and filter. (VAN MONS)

R. Rhubarb, two ounces.
 Lesser Cardamoms, one dram.
 Alcohol (15 degrees), two ounces.
 Spanish White Wine, one pound.

Macerate for eight days, and strain. (BERGIUS)

R. Rhubarb, two ounces.
 Lesser Cardamoms, two drams.
 Spanish Wine, one pound.

Digest for four days, and strain with expression. (FENN.; SUEC.)

R. Rhubarb, two ounces.
 Bitter Orange Peel, half an ounce.
 Lesser Cardamoms, two drams.
 White Wine, two pounds.
 Alcohol, two ounces.

Macerate for several days, and strain. (OLDEN.; NIEMANN)

R. Rhubarb, one ounce.
 Lesser Cardamoms, two drams.
 Saffron, one dram.
 Spanish Wine, eight ounces.

Digest for six days, and strain; pour upon the residue

Spanish Wine, four ounces.

Digest for twenty-four hours, and mix the strained liquors together. (ANT.)

GENEV. prescribes two ounces of rhubarb, half an ounce of cardamoms, two drams of saffron, and two pounds of Spanish wine.

It is excitant and slightly laxative, and is extolled in hypochondria, hysteria, scrofula, and tabes mesenterica.—Dose, one or two spoonfuls to adults, and from thirty to sixty drops to children.

BITTER WINE OF RHUBARB.

Vinum Rhei seu Rhabarbari amarum, Elixir Rhei amarum. (DAN.; FENN.; SUEC.; VAN MONS)

R. Rhubarb, two ounces.
 Gentian, six drams.
 Lesser Cardamoms, half an ounce.
 White Wine, two pounds.

Digest in a gentle heat, and strain. (DAN.)

R. Rhubarb, one ounce.
 Gentian, two drams.
 Canella Alba, one dram.
 Spanish Wine, one pound.

Digest for three days, and express. (FENN.; and SUEC.)

R. Cinnamon, one part.
 Madeira Wine, ninety-six parts.

Infuse without heat for two days, and add, Rhubarb, eight parts.

Continue the digestion for eight days, then add,

Gentian, one part.

Express at the expiration of several days, and filter. (VAN MONS)

It is excitant and tonic.—To be taken in the same cases and at the same doses as the preceding.

DAREL'S TINCTURE.

Tinctura Rhei Darelli seu dulcis. (BAVAR.; BRUNS.; HANN.; HASS.; POL.; BORUS.; SAX.; SUEC.; NIEMANN; PIERQUIN; VAN MONS)

R. Rhubarb, eight parts.
 Bitter Orange Peel, two parts.
 Lesser Cardamoms, one part.
 Root of Elicampagne, four parts.
 Madeira Wine, one hundred and twenty-eight parts.

Macerate without heat for several days; then express slightly, and filter. (VAN MONS)

R. Rhubarb, sliced, two ounces.
 Orange Peel, half an ounce.
 Lesser Cardamoms, two drams.
 Malaga Wine, two pounds.

Digest for four days at a moderate heat; then express, and add,

Extract of Elicampane, half an ounce.
White Sugar, three ounces.

Mix. (BAVAR.; POL.; BORUS.; SAX.)

- R. Rhubarb, in powder, two ounces.
Corinth Raisins, one ounce.
Lemon Peel, } of each, half an ounce.
Liquorice, }
Lesser Cardamoms, two drams.
Oporto Wine, two pounds.

Digest for two days, and add to the expressed liquor,

Extract of Elicampane, half an ounce.
White Sugar, three ounces.

Mix. (BRUNS.; HANN.; HASS.; NIEMANN; PIERQUIN)

- R. Rhubarb, one ounce and a half.
Lesser Cardamoms, half an ounce.
Raisins, stoned, three ounces.
Spanish Wine, one pound and a half.

Macerate for four days in a warm place; then add to the expressed liquor,

White Sugar, four ounces.
Extract of Elicampane, half an ounce.
Mace, half a dram.
Subcarbonate of Potass, one dram.

Preserve. (SUEC.)

The properties and uses of this are the same as with the preceding preparation. This has also been recommended against worms, but then the wine should be replaced by hydromel, and after four days' digestion, two ounces of extract of tansey should be mixed with the tincture instead of the extract of elicampane.

C.

Extraction by Alcohol.

ALCOHOLIC TINCTURE OF RHUBARB.

Alcohol cum Rheo. (AMST.; AUSTR.; BRUNS.)

- R. Rhubarb, sliced, one ounce.
Alcohol, one pound.

Macerate for one day, and filter. (BRUNS.)

AMST. prescribes one part of rhubarb, eight parts of alcohol (0.907), and eight days' maceration.—AUST.—two ounces of rhubarb and one pound of alcohol (0.910).

It is excitant and purgative, and is recommended in diarrhœa, colic, diseases attributed to weakness of the intestines, and in engorgements of the abdominal viscera.—It is used in chemistry for ascertaining the presence of alkalis which turn it red.

AROMATIC ALCOHOLIC TINCTURE OF RHUBARB.

Tinctura Rhei spiritiuosa seu Rhabarbari.
(LOND.; DUBL.; EDINB.; AMER.; FULD.; HASS.; LIPP.; LUSIT.; CADET

DE GASSICOURT; COXE; PIDERIT;
SWEDIAUR; VAN MONS)

Tinctura Rhei.

- R. Rhubarb, sliced, two ounces.
Cardamom Seeds, bruised, one ounce and a half.
Saffron, two drams.
Proof Spirit, two pints.

Macerate for fourteen days in a gentle heat, and filter. (LOND.)

- R. Rhubarb, sliced, two ounces.
Lesser Cardamoms, } of each,
bruised, } half
Liquorice, bruised, } an ounce.
Saffron, two drams.
Proof Spirit, two pints.

Digest for seven days, and filter. (DUBL.)

- R. Rhubarb, sliced, three ounces.
Lesser Cardamoms, bruised, half an ounce.

Proof Spirit,
two pounds and a half.

Digest for several days, and filter. (EDINB.)

AMER.; CADET DE GASSICOURT; COXE; and SWEDIAUR give the same formula as EDINB.—LUSIT. prescribes two ounces of rhubarb.

- R. Rhubarb, two ounces.
Lesser Cardamoms, half an ounce.
French Brandy, two pounds.

Digest for three days. Add to the filtered tincture,

White Sugar, four ounces.

Digest till the sugar is dissolved. (HASS.; PIDERIT)

This formula gives the *Tinctura Rhei spiritiuosa dulcis*.

Tinctura Rhei composita. (LOND.; DUBL.; COXE)

- R. Rhubarb, sliced, two ounces.
Liquorice, bruised, half an ounce.
Ginger, sliced, } of each,
Saffron, } two drams.
Proof Spirit, one pint.
Water, twelve fluid ounces.

Macerate for fourteen days in a gentle heat, and filter.

AMER. directs two ounces of rhubarb, one ounce each, of liquorice and of anise,

and two ounces of sugar, to be digested for ten days in two pints and a half of proof spirit. It is excitant and has been particularly recommended as carminative and digestive.—Dose, from half an ounce to one ounce in an appropriate vehicle.

BITTER TINCTURE OF RHUBARB.

Tinctura Rhei amara seu composita, Tinctura Rhei seu Rhubarbari et Gentianæ. (EDINB.; AMER.; BRUNS.; HAMB.; HANN.; WIRTEM.; COXE; VAN MONS)

Tinctura Rhei et Gentianæ. (EDINB.)

R. Rhubarb, sliced, two ounces.
Gentian, sliced, half an ounce.
Proof Spirit,

two pounds and a half.

Digest for seven days, and filter.

AMER.; and COXE give the same formula.

R. Tincture of Rhubarb,
one pound and a half.
Compound Tincture of Gentian,
half a pound.

Mix. (LUSIT.)

R. Rhubarb, two ounces.
Gentian, one ounce.
Orange Peel, half an ounce.
Alcohol, two pounds.

Digest for forty-eight hours, and filter. (HAMB.)

R. Rhubarb, eight parts.
Gentian, one part and a half.
Serpentaria, one part.
Proof Spirit, forty-eight parts.

Infuse without heat for several days; then express and filter. (VAN MONS)

BRUNS. directs one ounce of rhubarb, one dram and a half of gentian, one dram of serpentaria, and one scruple of cochineal; to be digested for three days in one pound of proof spirit.

R. Rhubarb, one ounce.
Gentian, one dram and a half.
Serpentaria, one dram.
Proof Spirit, } of each, half a pound.
White Wine, }

Digest, and filter. (HANN.; WIRTEM.)

R. Rhubarb, one ounce.
Gentian, three drams.
WINTER'S Bark, one dram.
Proof Spirit, sixteen ounces.

Infuse without heat for several days; then express lightly, and filter: (VAN MONS)

It is excitant, and was formerly extolled in bilious affections, fevers, and dysentery. —Dose, from fifty to eighty drops.

TINCTURE OF RHUBARB WITH ALOES.

Tinctura Rhei et Aloes. (EDINB.)

R. Rhubarb, sliced, ten drams.
Socotrine Aloes, powdered, six drams.
Lesser Cardamoms, bruised, half an ounce.

Proof Spirit,
two pounds and a half.

Digest for seven days, and filter.

RHODODENDRUM CHRYSANTHUM, LINN.—YELLOW-FLOWERED RHODODENDRON.

SYNONIMES:—Rosage à fleurs blanches, FR.; Sibirische Schneerose, GERM.; Roze Sybiryjskie, POL.

EDINB.; FERR.; HANN.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; COXE; MURRAY and GMELIN.

A shrub of the Alps and of Siberia. (*Decandria Monogynia*, LINN.; *Rhodbraceæ*, JUSS.; *Fig. Pall. Fl. Ross.* I. t. 30)

The stems and leaves (*Stipites et Folia Rhododendri chrysanthi*) are employed. They consist of branches of the thickness of goose-quills, long, and of a brownish colour; its leaves are oblong, rough, shining, and have short peduncles.

It is excitant, narcotic, sudorific, and diuretic.

CONSERVE OF RHODODENDRON.

Conserba Rhododendri chrysanthi. (SWEDIAUR)

R. Fresh Leaves of Rhododendron, one part.
White Sugar, two parts.
Beat together, and form a homogeneous paste.

ANTI-ARTHRITIC POWDER. (CADET DE GASSICOURT)

R. Liquorice Root, two drams.

Leaves of *Rhododendron*, ten grains.
Pulverize, mix, and divide into ten doses.
—Dose, one morning and evening, increasing by one daily till the patient experiences an abundant diaphoresis.

INFUSION OF RHODODENDRON.

Infusum Rhododendri chrysanthi. (AUGUSTIN; NIEMANN)

R. Leaves of *Rhododendron*,
from two drams to half an ounce.
Spring Water, ten ounces.
Digest for twenty-four hours in a covered vessel, and strain. (NIEMANN)

R. Leaves of *Rhododendron*,
half an ounce.
Boiling Water, eight ounces.
Infuse, and add to the strained liquor,

Aromatic Liquor, two drams.
Mix. (AUGUSTIN)

It is recommended by KÆLPIN, in chronic rheumatism. — Dose, a coffee cupful twice or thrice a day.

TINCTURE OF RHODODENDRON.

Tinctura Rhododendri chrysanthi. (NIEMANN)

R. Leaves of *Rhododendron*, two ounces.
French Brandy, } of each, half a pound.
Spanish Wine, }

Digest for fifteen days.

It is excitant and slightly narcotic, and is also considered diuretic. — Dose, five spoonfuls, in a decoction of the residue of the tincture with one pound of water.

RHUS.—SUMACH.

Three species of this genus of plants are employed in medicine.

1. *Rhus Coriaria*, LINN. — Carriers' Sumach.

SYNONIMES: — Sumac des Corroyeurs, FR.; Sumach, Gerberbaum, GERM.; Tuntumtum, ARAB.; Zumague, SPAN.; Sumack, DUT.; Sommaco, ITAL.; Sumak, PERS.; Sumagro, PORT.

BRUNS.; GALL.; HISP.; FERR.; WIRTEM.; HERBIP.; AINSLIE; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

A large shrub of the south of Europe. (*Pentandria Digynia*, LINN.; *Terebinthaceae*, JUSS.; Fig. Zorn, *l.c.* pl. t. 510)

The leaves and fruit are employed.

The leaves (*Folia Sumac*) are winged and composed of fifteen or seventeen oval, sessile leaflets, hairy above, and bordered with obtuse teeth. They have a very astringent taste.

The berries are of a beautiful red colour, of the size of a lentil, and of an astringent and acid taste.

It is astringent, but is not much used.

2. *Rhus cotinus*, LINN.

SYNONIMES: — Fustet, FR.; Perückensumach, Venussumach, GERM.; Coccigria, ITAL.

AUSTRI.; FERR.

A tree of the south of Europe. (*Pentandria Digynia*, LINN.; *Terebinthaceae*, JUSS.; Fig. Jacq. *Fl. Austr.* t. 210)

The bark and leaves are employed.

The bark (*Cortex Cotini*) is smooth and whitish without.

The leaves are of a reversed oval figure, entire, and smooth on both sides. They have a styptic, balsamic, and resinous taste.

It is astringent and tonic.

3. *Rhus Toxicodendrum et radicans*, LINN. — Common Upright Sumach, Poison Oak, Toxicodendron.

SYNONIMES: — Sumac vénéneux, FR.; Giftbaum, Giftsumach, Wurzelsumach, GERM.; Vergiftboom, DUT.; Sommaco radicante, ITAL.

LOND.; EDINB.; AMER.; ANT.; BAVAR.; BELG.; GALL.; FERR.; SAX.; BRUGNATELLI; COXE; GUIBOUT; PARIS.

A shrub of North America. (*Pentandria Digynia*, LINN.; *Terebinthaceae*, JUSS.; Fig. Dill. *Elth.* 389. t. 291. f. 375)

The leaves (*Folia Rhus* seu *Rhois Toxicodendri*) are employed. They are ternate, dentated angularly or entire, downy, and have petiolated leaflets. They possess such acridity that by simply touching them the skin becomes violently irritated; some say the exhalation from them is sufficient to do so.

It is stimulant and poisonous, and has been recommended in tetters and palsy.

EXTRACT OF TOXICODENDRON.

Extractum Rhois radicanis. (ANT. ; BELG. ; GALL. ; HANN. ; SAX. ; NIEMANN ; VAN MONS)

R. Fresh Leaves of Toxicodendron, any quantity.

Bruise them in a marble mortar, sprinkling them with a little water; and express the juice; evaporate it immediately in a water-bath, constantly stirring towards the end, with a spatula. (GALL. ; HANN. ; NIEMANN)

R. Fresh Leaves of Toxicodendron, any quantity.

Bruise them rapidly in a stone mortar, adding one half of their weight of water, and express the juice; set it aside for a moment; strain without expression, and evaporate to the consistence of an extract, constantly stirring. (VAN MONS)

R. Fresh Leaves of Toxicodendron, } of each,
Cold Water, } one part.

Macerate for twelve hours, and add,
Boiling Water, four parts.

Strain when cold, clarify with white of egg, and set the liquor upon the fire; when it has begun to thicken, strain it through a sieve, and evaporate it constantly stirring. (ANT. ; VAN MONS)

R. Leaves of Toxicodendron, two pounds.
Boiling Water, ten pounds.

Digest for twenty-four hours in a covered vessel, and then boil for one hour; decant and express when cold, and after the liquor is depurated by repose, evaporate it to a proper consistence. (BELG.)

VAN MONS directs one part of dried leaves to be boiled twice successively in ten parts of water, the liquor to be strained through a silk-sieve, and evaporated.—SAX.—one part of dried leaves to be boiled in thirty parts of water till reduced to fifteen, and the liquor to be strained and evaporated.

Dose, half a dram four times a day, gradually increasing to six drams and even to an ounce; mixed with sweetened water. Several ounces have been given in the course of the day.

EXCITANT POTION.

Mixtura Extracti Rhois radicanis.
(BRERA)

R. Extract of Toxicodendron, one dram.
Water, one ounce.

Mix.—Dose, ten drops, gradually increasing it to a tea-spoonful, twice a day; in palsy of the extremities.

PILLS OF TOXICODENDRON.

Pilulae Toxicodendri. (FERR. ; BRERA ; VAN MONS)

R. Chloruret of Barium, one part.
Extract of Toxicodendron, six parts.
Leaves of Toxicodendron, in powder, a sufficient quantity.

Make pills of four grains each.—Dose, three, twice or thrice a day, in tetters. (VAN MONS)

R. Leaves of Toxicodendron, one grain.
Liquorice Powder, one scruple.
Rob of Elder, a sufficient quantity.

Make four pills, to be taken in the course of the day; in paralysis consecutive to apoplexy. (FERR. ; BRERA)

TINCTURE OF TOXICODENDRON.

Tinctura Rhois radicanis. (SAX. ; HUFELAND)

R. Juice of Toxicodendron, } of each,
recently expressed, } one part.
Concentrated Alcohol, }

Infuse without heat for several days, and filter. (SAX.)

HUFELAND prescribes two ounces of leaves and one pound of alcohol; to be infused for three days.

RIBES.

Three species of this genus are employed:

1. *Ribes Nigrum*, LINN.—Black Currant.

SYNONIMES: — Cassis, Groseillier noir, FR. ; Schwarze Johannisbeere, Gichtbeere, GERM. ; Rybes neb wjno sjana cerne, BOHEM. ; Sæller, DAN. ; Swarta winbær, SWED.

AMST. ; BATAV. ; BELG. ; BRUNS. ; DAN. ; GALL. ; FENN. ; FULD. ; GENEV. ; HASS. ; LIPP. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A shrub of the northern parts of both continents. (*Pentandria Monogynia*, LINN. ; *Grossularia*, CAND. ; Fig. Zorn, Ic. pl. t. 305)

The fruit is employed; it consists of globular black berries, disposed in bunches, and rather aromatic than acid.

CONSERVE OF BLACK CURRANT. (VAN MONS)

R. Leaves of Black Currant, beaten
to a fine pulp, one part.
White Sugar, two parts.

Beat together, and apply the heat of a water-bath to the mass placed in a covered vessel till the sugar is dissolved.

JUICE OF BLACK CURRANTS.

Succus Ribesiorum nigrorum. (FULD.)

R. Ripe Black Currants, any quantity.

Bruise them in a stone mortar, and set aside the mass in a cellar for several days; then express strongly, strain the juice through flannel, and preserve it in long-necked bottles.

ROB OF BLACK CURRANTS.

Rob seu Gelatina Ribesiorum nigrorum.
(AMST.; BATAV.; BELG.; LIPP.; ROSS.; SAX.; SWEDIAUR)

R. Ripe Black Currants, any quantity.

Boil with a little water in a covered tin vessel, constantly stirring, till the berries burst; strain the liquor through a sieve, and express the residue strongly: add half a pound of sugar to each pound of the juice, and evaporate to the proper consistence, removing the scum as it is formed. (AMST.)

BATAV.; and BELG. prescribe four parts of sugar, and ten of juice.—SAX.—one of sugar, and eight of juice.—ROSS.—one of sugar, and six of juice.—LIPP.; and SWEDIAUR—one of sugar, and four of juice.

It is extolled in sore throats.—It is added to gargles.

SYRUP OF BLACK CURRANTS.

Syrupus Ribis nigri seu Ribesia nigræ.
(FENN.; SUEC.; COXE; VAN MONS)

R. Depurated Juice of Black Currants,
one pound.
White Sugar, two pounds.

Make a syrup at a gentle heat. (FENN.; SUEC.)

Coxe prescribes one pint of juice, and two pounds of sugar.

R. Residue of the expressed Juice of
Black Currants, one part.
Boiling Water, one part and a half.

Digest for twenty-four hours, and add to the liquor,

White Sugar, an equal part.

VOL. II.

Clarify, and boil to the consistence of a syrup. (VAN MONS)

SPIRIT OF BLACK CURRANTS. (VAN MONS)

R. Ripe Black Currants, two parts.

Bruise them after having infused them in warm water; leave the mass to ferment, and then distil with

Fresh Leaves of Black Currants,
one fourth part.
Proof Spirit, one part and a half.
Water, twelve parts.

Distil three parts.

2. *Ribes rubrum*, LINN.—Red Currant.

SYNONIMES:—Groseillier rouge, FR.; Rothes Johannisbeere, GERM.; Rybes neb wjno, sjana cerwene, BOHEM.; Ræde ribs, DAN.; Agracejo encarnada, SPAN.; Rothe aalbezie, DUT.; Uva de'prati, ITAL.; Porzeczki, POL.; Smorodina krasnaja, PORT.; Ræda winbær, SWED.

AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; HISP.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A shrub which grows in every part of Europe. (*Pentandria Monogynia*, LINN.; *Grossulariæ*, CAND.; *Fig. Flore Medic. IV.* 189)

The fruit (*Baccæ Ribesiorum seu Ribium rubrorum*) is employed; it consists of small, very juicy, globular berries, disposed in bunches, red or yellowish-white, inodorous, and of an acidulous and sweet taste.

JUICE OF RED CURRANTS.

Succus Ribium rubrorum. (GALL.; HISP.; FULD.; HASS.; SAX.; SARD.; PIDERIT; VAN MONS)

R. Red Currants, any quantity.

Bruise them with the fingers, and leave the mass in a cellar till there appears on the surface a clear liquor; then express the juice, set it aside in a cool place for several days, and preserve it, when strained, in long necked bottles under a stratum of oil. (GALL.; HISP.; FULD.; SARD.; SAX.)

R. Red Currants, any quantity.

Bruise them in a copper pan, and strain the juice through a sieve; having set it aside to depurate by repose, strain it through flannel: add one sixteenth of sugar to the liquor, and boil for an instant. Pour the juice into bottles, filling them only to the

shoulders, and cork them after having dipped a lighted match into the neck. (VAN MONS)

HASS.; and PIDERIT add one fourth of sugar to the juice.

ROB OR JELLY OF RED CURRANTS.

Roob Ribium seu Ribesiorum rubrum, Gelatina Ribium. (AMST.; ANT.; BELG.; BATAV.; BRUNS.; GALL.; FULD.; HASS.; ROSS.; SARD.; SAX.; WIRTEM.; HERBIP.; PIDERIT; SWEDIAUR; VAN MONS)

R. Ripe Red Currants, any quantity.

Boil them with a little water in a tin vessel, constantly stirring, till they are burst; then strain through a sieve, and express strongly. Add four parts of sugar to ten of juice, and evaporate to the consistence of honey, constantly stirring. (BATAV.; BELG.)

AMST. prescribes one part of sugar, and five parts of juice.—ANT.; BRUNS.; FULD.; HASS.; WIRTEM.; HERBIP.; PIDERIT, and SWEDIAUR—one of sugar, and four of juice.—ROSS.—one of sugar, and six of juice.—SAX.—one of sugar, and sixteen of juice.

R. Recently expressed Juice of Red Currants, any quantity.

Strain, and boil it over a moderate fire to the consistence of honey. (GALL.; SARD.)

R. Ripe Red Currants, fifteen pounds.
White Sugar, twelve pounds.

Boil together over a slow fire, constantly stirring, till one fourth of the humidity is evaporated; strain while hot, and leave the mass to gelatinize in a cool place. (WIRTEM.)

VAN MONS directs two pounds of sugar to be added to each pound and a half of expressed juice, and the mixture evaporated to the consistence of jelly.

R. Syrup, boiled à la grande plume, two pounds.
Juice of Red Currants, } of each,
Water, } one pound.

Boil to the consistence of jelly, and strain. (HISP.)

It is acidulous and refreshing, and is principally employed to render drinks grateful to patients.

SYRUP OF RED CURRANTS.

Syrupus Ribium seu Ribesiorum seu Ribis rubri seu de Ribesiis. (BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; WIRTEM.; HERBIP.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

R. Juice of Red Currants, depurated and filtered, two pounds.
White Sugar, three pounds and a half.

Dissolve at a moderate heat. (GALL.)

BRUNS.; DAN.; FULD.; PALAT.; WIRTEM.; and HERBIP. prescribe ten ounces of juice, and sixteen of sugar.—HANN.; OLDEN.; POL.; and BORUS.—twenty ounces of juice, and three pounds of sugar.—HASS.; and PIDERIT—ten ounces of juice, and thirteen and a half of sugar.—SWEDIAUR—sixteen ounces of juice, and twenty-eight of sugar.—BELG.—sixteen ounces of juice, and two pounds and a half of sugar.—HISP.—two pounds two ounces of juice, and four pounds of sugar.—BAVAR.—one part of juice, and two parts of sugar.—VAN MONS—two parts of juice, and three and a half of sugar.—LIPP.—three of juice, and four of sugar.—FERR.; and COXE—an equal part of each.

R. Red Currants, one hundred pounds.

Bruise them, put them into a pan, and apply heat, constantly stirring, till the seeds burst; then put the mass upon a sieve, express, and add to the juice obtained

Juice of Sour Cherries, five pounds.

Leave the mixture in a cool cellar, and at the end of thirty-six hours, break down the coagulum with an osier broom; strain and dissolve twenty-eight ounces of sugar in each pound of the juice obtained.

ACIDULOUS POTION.

Potus acidus. (AUGUSTIN)

R. Syrup of Red Currants, } of each,
Raspberries, } one
Barberries, } ounce.
Boiling Water, two pounds.
Mix.

3. *Ribes grossularia et Uva crispæ*, LINN.—Gooseberry.

SYNONIMES:—Grosseillier à maquereau, FR.; Stachelbeere, GERM.; Stikkelbær, DAN.; Grossellero, SPAN.; Kruisbezie, DUT.; Grispignolo, ITAL.; Agrest, POL.; Groselheira espin, PORT.; Steckelbær, SWED.

HISP.; GALL.; FERR.; BERGIUS; BRUGNATELLI; GUIBOURT.

A shrub of the north of Europe. (*Fig. Blackw. Herb. t. 277*)

The fruit is employed; it consists of globular or ovoid berries, greenish or reddish, of the size of a filbert, of an acid and astringent taste before maturity, and vapid and saccharine after.

The same preparations of this fruit are required by the above pharmacopœias as of the preceding species.

RICINUS COMMUNIS, LINN.—RICINUS, PALMA CHRISTI.

SYNONIMES:—Ricin, FR.; Wunderbaum, GERM.; Shocœwetssj, BOHEM.; Purgeerkorn, DAN.; Ricino, ITAL.; and SPAN.; Wonderboom, DUT.; Mamoma, POL.; Erando, SANS.; Undertræd, SWED.; Vullak unnay, TAM.; Amidum, TEL.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GME-LIN.; PARIS.; SPIELMANN.; ZARDA.

A plant of the south of Europe and of the north of Africa. (*Monœcia Monadelphica*, LINN.; *Euphorbeaceæ*, JUSS.; *Fig. Flore Medic. VI. 298*)

The seed (*Semen Ricini*) is employed. It is oval, prolonged, and flat on both sides. Under a thin, shining, brittle pellicle, veined with grey and white, the eye of which is yellowish and projecting, it contains a white and oleaginous kernel.

OIL OF RICINUS; CASTOR OIL.

Oleum Ricini seu Palmæ Christi.

1. *That which is met with in commerce.* (LOND.; DUBL.; EDINB.; DAN.; HASS.; OLDEN.; POL.; BORUS.; SAUNDERS)
2. *Prepared expressly.* (LOND.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; LIPP.; LUSIT.; ROSS.; SAX.; SUEC.; HERBIP.; BRUGNATELLI.; COXE.; SWEDIAUR.; VAN MONS)

Oleum Ricini. (LOND.)

R. Seeds of Ricinus, any quantity.

Bruise them after having freed them from the husks, and express the oil without heat.

R. Seeds of Ricinus freed from their envelope and of their embryo, any quantity.

Bruise them in a marble mortar with a wooden pestle; enclose the paste in a close cloth, and express slowly. Filter the oil by the heat of a warm-bath.

FIGUIER has recommended the seeds to be bruised with alcohol of commerce, the

paste to be submitted to the press, and the oil which flows to be distilled, drawing off two thirds of the alcoholic vehicle; the residue to be then boiled till all the humidity is dissipated.

This process is bad.

HENRI recommends the seeds to be tor-rified in an iron stove, to be bruised when cold, and the paste mixed with a certain quantity of water; the mixture to be then put over the fire, constantly stirring, till it boils: then the supernatant and swollen oily matter to be removed, heated apart that the water may be evaporated, and when the oil is isolated, to be strained through a close flannel, and finally set aside and decanted.

Castor oil is frequently acrid. This acridity is by some attributed to the pellicle and by others to the embryo of the seeds.—LECANU and BUSSY have ascertained that when subjected to distillation, the oil yields two acids, the *Ricinic* and the *Oleo-Ricinic*, which are formed, as well as a third, *Stearo-ricinic*, by the re-action of alkalis; that the two first are present in old oil, and it is these which render it acrid.

When new, castor oil is a mild purgative and is frequently prescribed as anthelmintic.—Dose, from two drams to three ounces. in broth; or a spoonful every hour till it produces the desired effect.

PURGATIVE EMULSION. (GALL.; AUGUSTIN.; BRERA.; CADET DE GASSICOURT.; ELLIS.; PIERQUIN.; SAINTE-MARIE.; VAN MONS)

R. Castor Oil, one ounce.

Triturate it with yolk of egg and one bitter almond, and dissolve it in

Broth, three ounces.

To be taken immediately. (BRERA)

R. Castor Oil,	} of each,
Emulsion of Bitter	
Almonds,	
Broth,	two ounces.

Mix.—To be taken by spoonfuls.

R. Castor Oil, two ounces.

Oil of Almonds, one ounce and a half.
Emulsion of Gum Arabic, four ounces.

Mix.—To be taken gradually. (BRERA)

R. Castor Oil,	three ounces.
Mucilage of Gum Arabic,	two drams.
Mint Water,	eight ounces.

Mix.—Dose, two spoonfuls every half hour. (AUGUSTIN)

R. Castor Oil, one ounce.
 Gum Arabic, one dram.
 Syrup of Marshmallows, half an ounce.
 Elder-flower Water,

one ounce and a half.
 Nitric Æther, one dram.
 SYDENHAM'S Laudanum, one scruple.

Mix.—Dose, a spoonful every hour. (AUGUSTIN)

R. Castor Oil, one ounce and a half.
 Gum Arabic, two drams.
 White Sugar, one dram.
 Mint Water, four ounces.
 Tincture of Opium, forty or fifty drops.

Mix.—Dose, a spoonful every hour. (ELLIS)

R. Castor Oil, three ounces.
 Yolk of Egg, one.
 Syrup of Marshmallows, two ounces.
 Water, ten ounces.

Mix.—Dose, two spoonfuls every half hour. (AUGUSTIN)

R. Castor Oil, three ounces.
 White Sugar, half an ounce.
 Yolk of Egg, one.
 Rose Water, ten ounces.

Mix.—Dose, two spoonfuls every half hour. (AUGUSTIN)

R. Castor Oil, two ounces.
 Yolk of Egg, one.
 Syrup of Marshmallows, one ounce.

Triturate together, adding gradually

Pure Water, eight ounces.
 Hordeated Cinnamon Water, two drams.

Mix. (SAINTE-MARIE)

GALL prescribes one ounce of oil, one half of a yolk of egg, two ounces of water, and half an ounce each, of orange-flower water and of simple syrup.—CADET DE GASSICOURT—one ounce of oil, two ounces of water, a sufficient quantity of yolk of egg, and half an ounce each, of compound syrup of cichory and of orange-flower water.—VAN MONS—two parts of oil, one part of yolk of egg, one of syrup, one of orange-flower water, and four of pure water.

R. Castor Oil, one ounce.
 Yolk of Egg, one.
 Hyssop Water, eight ounces.
 Subcarbonate of } of each,
 Magnesia, } two drams.
 Manna, }

Mix.

PURGATIVE POTION.

Haustus catharticus oleosus. (AUGUSTIN; BRERA; BORIES; DISP. CHA. PARIS; RATIER; SWEDIAUR)

R. Castor Oil, two ounces.

Syrup of Lemon, one ounce.

Mix. (RATIER)

RATIER also prescribes an equal part each, of the oil and of compound syrup of cichory.—BORIES.—an equal part each, of the oil and of syrup of peach flowers.

R. Castor Oil, one or two ounces.
 Simple Syrup, one ounce.
 Gum Arabic, two drams.
 Water, four ounces.

Mix. (RATIER)

R. Castor Oil, one or two ounces.
 Tartaric Syrup, one ounce.
 Water of Cichory, two ounces.

Mix. (RATIER)

RATIER also prescribes two ounces of oil, two of syrup of buckthorn, and one ounce of mint water; or, two ounces of oil, two of compound syrup of cichory, and two drams of mint water; or, six drams of oil, half an ounce of oxymel of squill, and one ounce of syrup of buckthorn.

R. Castor Oil, one ounce.
 Proof Spirit, } of each, half an ounce.
 Sugar, }

Mix. (SWEDIAUR)

R. Castor Oil, two ounces.
 Broth, three ounces.

Mix.—For one dose. (BRERA)

R. Castor Oil, one ounce.
 Decoction of Herbs, four ounces.

Mix. (SPELMANN)

R. Castor Oil, three ounces.
 Soap, half an ounce.
 Syrup of Orange Peel, three ounces.

Mix.—Dose, a spoonful every half hour. (AUGUSTIN)

R. Castor Oil, three ounces.
 Subcarbonate of Potass, one dram.
 Honey, one ounce.
 Lime-flower Water, ten ounces.

Mix.—Dose, two spoonfuls every two hours. (AUGUSTIN)

ANTHELMINTIC POTION.

Potio anthelmintica oleoso-aromatica.
 (BRERA)

R. Castor Oil, } of each,
 Peppermint Water, } two ounces.

Mix.—To be taken at one dose.

VERMIFUGE LAVEMENT.

Clyisma emolliens anthelminticum. (BRERA)

R. Castor Oil, three ounces.
 Decoction of Linseed, four ounces.

Mix.

ROBINIA PSEUDO-ACACIA, LINN.—FALSE ACACIA.

SYNONIMES :—Robinier, Faux Acacia, FR.;
Unächte Acacienbaum, GERM.

GALL.

An original tree of America, but natu-

ralized in Europe. (*Diadelphii Decandria*,
LINN.; *Leguminosæ*, JUSS.; *Fig. Nouv.*
Duh. II. t. 16)

Its bark is purgative and emetic.

ROCELLA, ORCELLA, ORCHILLA.—LITMUS.

SYNONIMES :—Argol, Archil.—Orseille,
Orchel, FR.

DUBL.; BAVAR.; GALL.; BERGIUS;
COKE; GUIBOUT; MURRAY.

This is a solid colouring paste, of a very
deep violet-red colour, and of a strong and
disagreeable smell.

It is prepared from the *Rocella tinctoria*,
CAND., a cryptogamous plant which grows
on the rocks of the Canaries, of Cape Verd,
and of the Archipelago. (*Fig. Pluk. Alm.*
t. 255, f. 6)—The *Variolaria orcina*, ACH.
also yields it, but this is of an inferior
quality.

ROSA.—ROSE.

Five species of this genus of plants are
mentioned in pharmacopœias :

1. *Rosa centifolia*, LINN.—Hundred-leaved
Rose, Damask Rose.

SYNONIMES :—Rosier à cent feuilles, FR.;
Centifolienrose, GERM.; Ruze cerwena,
BOHEM.; Fyld rose, Ræd rose, DAN.;
Rosa de Alexandria, SPAN.; Damask ro-
sen, DUT.; Roze centyfoliie, POL.; Rosa,
RUSS.; Roser, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.;
ANT.; AUSTR.; BATAV.; BAVAR.;
BELG.; BRUNS.; DAN.; HISP.; GALL.;
FENN.; FERR.; FULD.; GENEV.; HANN.;
HASS.; LIPP.; OLDEN.; LUSIT.; POL.;
BORUS.; ROSS.; SAX.; SUEC.; WIR-
TEM.; HERBIP.; BERGIUS; BRUGNA-
TELLI; COKE; GUIBOUT; MURRAY
and GMELIN; PIDERIT; SPIELMANN;
ZARDA.

A shrub cultivated in every garden of
Europe. (*Icosandria Polygynia*, LINN.,
Rosaceæ, JUSS.; *Fig. Zorn, Ic. pl. t. 600*)

The flowers (*Flores Rosarum incarnata-*
rum seu pallidarum) are employed. They
consist of oval, entire petals, of a pale red-
dish colour. Their odour is very agree-
able, but is almost entirely removed by de-
siccation; their taste is bitterish, acrid,
and styptic.

2. *Rosa moschata*, AIT.—Musk Rose.

GALL.

An original shrub of the north of Africa.
The flowers, according to WILLDENOW,
are used in preparing the otto of roses.

3. *Rosa alba*, LINN.—White Rose.

BRUNS.; GALL.; WIRTEM.; BERGIUS;
MURRAY and GMELIN; SPIELMANN;
ZARDA.

A shrub generally spread over Europe.
(*Fig. CEd. Fl. Dan. t. 1215*)

Its flowers (*Flores Rosæ albe*) have an
agreeable odour, and a bitterish and styptic
taste.

4. *Rosa Gallica*.—Red Rose, Provence Rose.

SYNONIMES :—Rose rouge, Rose de Pro-
vence, FR.; Essigrose, GERM.; Edike
rose, DAN.; Rosa rubra, Rosa Castella-
na, SPAN.; Fransche rosen, DUT.; Roza
Damascenskie, POL.; Rosa vermalha,
PORT.; Rättikeroser, SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.;
BATAV.; BELG.; BRUNS.; DAN.; HISP.;
GALL.; FERR.; FENN.; FORM. HÔP. FR.;
GENEV.; HANN.; HASS.; OLDEN.;
LUSIT.; POL.; BORUS.; ROSS.; SAX.;
SUEC.; WIRTEM.; WURCEB.; BERGIUS;

BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; SPIELMANN.

A shrub of the south of Europe. (*Fig. Blackw. Herb. t. 82*)

The flowers (*Flores Rosæ rubræ*) are employed. They are used before the petals are expanded, and the claws should be cut off.

5. *Rosa canina*, LINN.—Dog Rose.

SYNONIMES:—Rosier sauvage, Eglantier, FR.; Hundsrose, Hahnbuttenstrauch, Hagebutten, GERM.; Ssypec, Plana ruze, BOHEM.; Hybentorn, DAN.; Rosa silvestre, SPAN.; Hondsroos, DUT.; Rosa selvatica, ITAL.; Roza pulna, POL.; Rosa brava, PORT.; Neupon, SWED.

LOND.; EDINB.; AMST.; ANT.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; WIRTEM.; BERGIUS; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; ZARDA.

A shrub which grows in every part of Europe. (*Fig. Zorn, Ic. pl. t. 529*)

The flowers, fruit, and seeds are employed.

The flowers (*Flores Rosæ sylvestris seu rubræ vulgaris*) consist of reddish or nearly white petals, of a sweet and agreeable odour. They are slightly laxative.

The fruit, called *Hips* (*Fructus Cynosbati, Cynorrhodon*), consists of oval, smooth, scarlet berries; they are dried after the seeds and the hairs which adhere to them are removed. Its taste is rather styptic.

It is reputed diuretic and anti-spasmodic; but it is more certainly a powerful astringent and irritant, the abuse of which occasions cough, vomiting, and tenesmus.

The seeds (*Semina Cynosbati*) are long, angular, hard, and hairy. They are now no longer used.

All these plants are more or less astringent and tonic.

JUICE OF PALE ROSES.

Succus Rosarum pallidarum. (HISP.; GALL.; SARD.)

R. Petals of Pale Roses, any quantity.

Bruise them in a marble or wooden mortar, and strain with strong expression. Filter the juice, and preserve it in bottles under a stratum of oil. (HISP.; GALL.)

SARD. recommends the flowers to be put into a bag, plunged into boiling water, and hung up; first a greenish liquor drops from it, and afterwards a reddish, when it should be expressed.

CONSERVE OF RED ROSES.

Conserva seu Confectio Rosæ seu Rosarum. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; CADET DE GAS-SICOURT; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Confectio Rosæ Gallicæ. (LOND.; EDINB.)

R. Petals of the Red Rose, one pound.

Refined Sugar, three pounds.

Beat the petals in a stone mortar; then add the sugar, and continue to beat till they are incorporated.

Conserva Rosæ. (DUBL.)

R. Petals of the Red Rose, one part.

Refined Sugar, three parts.

Beat together till incorporated.

AMER.; AMST.; LIPP.; and COXE give the same formula.—ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; FULD.; HANN.; HASS.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; VAN MONS; and SWEDIAUR prescribe one part of petals and two parts of sugar.—FORM. HÔP. FR.; and SUEC.—an equal part of each.

R. Fresh Petals of Roses, fifteen parts.
Sugar, thirty parts.

Triturate together, strain the pulp through a hair sieve, and add

Simple Syrup, boiled to the consistence of an electuary,
one hundred and twenty parts.

Mix. (GALL.)

R. Fresh Petals of Roses, one pound.

Bruise in a glass mortar, adding gradually

Thick Simple Syrup, two pounds.

Evaporate slowly to the proper consistence. (HISP.)

R. Red Roses, one pound.

Put them into a bag, and keep it for some time in boiling water; then hang up the bag till a red liquid begins to drip from it, then express strongly, pound the residue in a mortar, and boil it with

Thick Simple Syrup, three pounds,
gradually adding the expressed juice, till it
acquires the consistence of a conserve.
(SARD.)

R. Dried Petals of Red Roses, ninety parts.
Rose Water, a sufficient quantity
to form a pulp. Having let it macerate for
six hours, add

Sugar, one thousand parts,
dissolved in rose water, and boiled to the
consistence which would be necessary for
forming it into trochiscs, and mix inti-
mately in a marble mortar, with a wooden
pestle. (GALL.)

FORM. HÔP. FR. prescribes pure water,
six hours' digestion at a moderate heat,
and the mixture of the pulp with an equal
quantity of sugar.—LUSIT.—three ounces
of powdered roses, eight ounces of rose
water, six hours' infusion, and one pound
and a half of sugar.—CADET DE GASSI-
COURT—six ounces of powder, a sufficient
quantity of rose water, and half a pound of
sugar.—FERR.—three ounces of powder,
eight ounces of rose water, and two pounds
of sugar.

R. Powdered Rose Petals, three ounces.
Rose Water, eight ounces.

Macerate for several hours, and add

White Sugar, one pound and a half,
previously formed into syrup with a suffi-
cient quantity of water, and mix. (GENEV.)

Dose, from half a dram to one dram.

VITRIOLATED CONSERVE OF ROSES.

Conserva Rosarum vitriolata. (BRUNS.;
SARD.; WIRTEM.)

R. Conserve of Roses, one pound.
Sulphuric Acid, one dram.

Mix well. (BRUNS.; SARD.)

WIRTEM. prescribes two drams of acid.

DRY CONSERVE OF ROSES.

*Tabellæ rosatæ seu Manus Christi, Conserva
Rosarum sicca, Saccharum rosatum.*
(BRUNS.; PALAT.; SARD.; WIRTEM.;
SPIELMANN)

R. Dried Petals of Red Roses, one ounce.
Simple Syrup, boiled to the proper
consistence, sixteen ounces.

Mix, and make trochiscs. (WIRTEM.;
SPIELMANN)

BRUNS. directs the powdered petals to be
first sprinkled with twelve drops of sul-
phuric acid.—PALAT. prescribes six drams
of petals, and four ounces of sugar boiled.

R. Red Roses, one ounce.
Simple Syrup, boiled to the proper
consistence, one pound and a half.
Lemon Juice, two drams.

Pulverize the roses, sprinkle them with
the juice, mix in the boiling syrup, and
form trochiscs. (SARD.)

R. White Sugar, one pound.
Rose Water, a sufficient quantity.

Boil to the proper consistence, and form
trochiscs. (PALAT.)

By adding towards the end, half an ounce
of prepared pearls, the preparation formerly
called *Saccharum perlatum* is obtained.

DIARRHODON TROCHISCS.

Tabellæ diarrhodon. (SARD.)

R. Diarrhodon Powder, one ounce and a half.
White Sugar, one pound.
Mucilage of Tragacanth, a sufficient quantity.

Make trochiscs.

PULP OF HIPS.

Pulpa Cynorrhodontos. (GALL.)

R. Hips, red but not perfectly ripe,
sixteen parts.

Cut off the crowns and pedicles, remove
the seeds, and macerate the rest for three
or four days, in

White Wine, two parts,
till the fruit becomes perfectly soft; then
bruise it in a marble mortar with a wooden
pestle, and pass the pulp through a close
silk-sieve.

CONSERVE OF HIPS.

*Conserva seu Confectio seu Rob Cynosbatos
seu Fructus Rosæ Caninæ.* (LOND.;
EDINB.; GALL.; FULD.; GENEV.; LU-
SIT.; PALAT.; SARD.; WIRTEM.; NIE-
MANN; SWEDIAUR; VAN MONS)

Confectio Rosæ Caninæ.

R. Pulp of Hips, one pound.
Refined Sugar, in powder,
twenty ounces.

Beat together till well incorpo-
rated. (LOND.)

R. Fresh Hips, freed from seeds
and hairs, one part.
Refined Sugar, three parts.

Reduce the fruit to a pulp, then
add the sugar, and beat till tho-
roughly incorporated. (EDINB.)

- R. Pulp of Hips, } of each,
White Sugar, } three pounds.

Evaporate slowly to the proper consistence. (GENEV.; SWEDIAUR)

FULD.; and PALAT. prescribe two pounds of pulp and one pound of sugar.—WIRTEM.—three of pulp and two of sugar.—LUSIT.; and VAN MONS—one of pulp and two of sugar.—NIEMANN.; and SARD.—one of pulp and three of sugar.—GALL.—one of pulp and one and a half of syrup boiled to the consistence of an electuary.

It is slightly astringent, diuretic, antispasmodic, and stomachic.—Dose, from four to twelve drams.

ASTRINGENT ELECTUARY.

Electuarium adstringens Fulleri, Conserva Rosarum rubrarum adstringens. (HISP.; BORIES; BRERA; PIERQUIN; SAINTE-MARIE)

- R. Conserve of Red Roses, two ounces.
Diascordium, half an ounce.
Armenian Bole, two drams.
Dragon's Blood, one dram and a half.
Pomegranate Flowers, half a dram.
Oil of Nutmeg, } of each,
Cinnamon, } two drops.
Syrup of Red Roses, one ounce.

Mix. (HISP.)

- R. Conserve of Red Roses, four ounces.
Nitrate of Potass, one dram.
Syrup of Poppy, one ounce.
Orange-flower Water, two drams.

Mix. (BORIES; PIERQUIN; SAINTE-MARIE)

- R. Conserve of Red Roses, four ounces.
Syrup of Tolu, one ounce.

Mix. (BRERA)

- R. Conserve of Hips, four ounces.
Syrup of Pomegranate, two drams.
Poppy, one ounce.
Nitrate of Potass, two scruples.

Mix. (SAINTE-MARIE)

It is prescribed in spitting of blood.—Dose, a tea spoonful every two or three hours.

TRAUMATIC ELECTUARY.

- R. Conserve of Red Roses, two ounces.
LOCATELLI's Balsam, one ounce.
Syrup of Tolu, half an ounce.

Mix.

ASTRINGENT MIXTURE.

Linctus communis. (SAUNDERS; SAINTE-MARIE)

- R. Olive Oil, two ounces and a half.
Conserve of Hips, half an ounce.

- Syrup of Poppy, one ounce.
Mix. (SAUNDERS)

- R. Conserve of Red Roses, two ounces.
Rosemary, } of each,
Elecampane, } one ounce.
Syrup of Amber,

Mix intimately. (SAINTE-MARIE)

Dose, a tea spoonful, thrice a day.—It is prescribed in inveterate leucorrhœa.

ASTRINGENT POTION. (BORIES; PIERQUIN; SAINTE-MARIE)

- R. Flowers of Lesser Mullein, one handful.
Water, one pound and a half.

Boil down to one half, and add to the strained liquor

Conserve of Red Roses, one ounce.

Mix.—It is recommended by FOUQUET, in dysentery.

INFUSION OF ROSES.

Infusum seu Mucharum seu Brodium Rosarum. (HISP.; FERR.; HASS.; LIPP.; SARD.; WIRTEM.; PIDERIT)

- R. Fresh Petals of Roses, two pounds.
Boiling Water, eight pounds.

Digest for eight hours, express, and pour the strained liquor upon a similar quantity of fresh petals; having repeated this operation a third time, preserve the liquor in a long-necked bottle, under a stratum of oil. (SARD.)

- R. Fresh Petals of Roses, } of each,
Boiling Water, } eighteen pounds.

Macerate for twenty-four hours, and strain; preserve the liquor in a bottle under a stratum of olive oil. (WIRTEM.)

HISP. prescribes one pound of roses and two pounds of water; to be infused for eight hours.—HASS.; and PIDERIT—one pound of red roses and one pound and a half of water; to be infused for several days.—LIPP.—an equal part of each; to be digested for three days.—FERR.—an equal part of each; to be digested for twelve hours.

This infusion is principally used in the preparation of the syrup and honey of roses.

ACIDULOUS INFUSION OF ROSES.

Infusum Rosæ acidulum seu compositum. (LOND.; EDINB.; DUBL.; AMER.; HISP.; GENEV.; SUEC.; BRUGNATELLI; COXE; NIEMANN; RATIER)

Infusum Rosæ compositum.

- R. Dried Petals of Red Roses, half an ounce.

Boiling Water, two pints and a half.
 Dilute Sulphuric Acid, three fluid drams.
 Refined Sugar, one ounce and a half.

Pour the water upon the petals in a covered glass vessel; then instil the acid, and macerate for half an hour; finally strain the liquor, and add the sugar to it. (LOND.)

DUBL. prescribes half an ounce of petals, three pints of water, three drams, by weight, of acid, and one ounce and a half of sugar.

Infusum Rosæ Gallicæ. (EDINB.)

R. Dried Petals of Red Roses, one ounce.

Boiling Water, two pounds and a half.
 Dilute Sulphuric Acid, half an ounce.
 Refined Sugar, one ounce.

Macerate the petals in an earthen vessel, which has not been glazed with lead, for four hours; then instil the acid, strain, and finally add the sugar.

AMER.; SUEC.; COXE; and NIEMANN give the same formula as LOND.—BRUGNATELI—one ounce of roses, five pounds of water, one dram of sulphuric acid, twenty-four hours' infusion, and two ounces of sugar.—HISP.—half an ounce of roses, three pounds of water, a quarter of an hour's infusion, one scruple of acid, and no sugar.—RATIER—half an ounce of roses, one pound of water, one ounce and a half of sugar, and half a dram of acid.—GENEV.—half an ounce of roses, two pounds and a half of water, one scruple of acid, and one ounce and a half of sugar.

It is slightly astringent, but is rarely prescribed alone. It is principally employed as a vehicle for various medicaments, and in gargles. It is sometimes given in passive hæmorrhages.

HÆMASTATIC TEA. (PIERQUIN)

R. Dried Petals of Roses, half an ounce.
 Vinegar, three spoonfuls.
 Sugar, one ounce.
 Boiling Water, two pounds.

Infuse for two hours, and strain.—To be taken by glassfuls in the course of the day.

VOL. II.

It is recommended in diarrhœa and hæmorrhages.

ASTRINGENT INFUSION. (BORIES)

R. Red Roses, } of each,
 The Five Myrobalani, } half an ounce.
 bruised,
 Water, one pound.

Digest in warm ashes. Add to the strained liquor

Syrup of Pomegranate, half an ounce.

Mix.—It is astringent, and is recommended in diarrhœa and hæmorrhages.—To be taken at three or four doses, in the course of the day.

SYRUP OF RED ROSES.

Syrupus Rosarum rubrarum seu Rosæ Gallicæ seu e Rosis siccis, Rhodosaccharum. (EDINB.; AMST.; ANT.; HISP.; GALL.; FERR.; GENEV.; SARD.; WIRTEM.; COXE; VAN MONS)

Syrupus Rosæ Gallicæ. (EDINB.; COXE)

R. Dried Petals of Red Roses, one part.

Boiling Water, nine parts.
 Refined Sugar, ten parts.

Macerate the petals in the water for twelve hours; then boil a little, and strain. Add the sugar to the liquor, and boil to a syrup.

R. Dried Petals of Red Roses, eight ounces.

Boiling Water, five pounds and a half.

Digest for twelve hours; then boil down to one half, and express: set the liquor apart to depurate by repose, and evaporate, so that on the addition of

Sugar, sixteen ounces,
 a syrup may be formed without ebullition. (AMST.)

R. Depurated Juice of Red Roses, two pounds.

White Sugar, four pounds.
 Water, a sufficient quantity.

Boil to the consistence of syrup. (HISP.)

R. Fresh Petals of Red Roses,

four pounds.
 Boiling Water, eight pounds.

Infuse for twelve hours in a covered vessel; express slightly, set aside the juice for several hours, decant, and add to it

White Sugar, twice the quantity.

Make a syrup in a water-bath. (GALL.)

WIRTEM. prescribes half a pound of dried roses, one pound and a half of boiling

3 U

water, twenty-four hours' maceration, and sixteen ounces of sugar to every nine ounces of the strained liquor.—ANT.; and BELG.—one pound of petals, three pounds of water, twelve hours' digestion, and four pounds of sugar.

- R. Fresh Petals of Red Roses, one part.
Cold Water, three parts.
Boiling Water, seven parts.

Infuse for six hours, strain through a sieve with slight expression, and add

- White Sugar, four parts.
Clarify, and make a syrup. (VAN MONS)

SYRUP OF PALE ROSES.

Syrupus Rosæ seu de Rosis pallidis seu Rosarum pallidarum seu rosatus simplex seu Rosarum solutivarum seu rosatus solutivus. (LOND.; EDINB.; DUBL.; AMST.; ANT.; HISP.; GALL.; FERR.; GENEV.; SARD.; WIRTEM.; COXE; VAN MONS)

Syrupus Rosæ. (LOND.; DUBL.)

- R. Dried Petals of the hundred-leaved Rose, seven ounces.
Refined Sugar, six pounds.
Boiling Water, four pints.

Macerate the petals in the water for twelve hours, and strain; then evaporate the liquor in a water-bath till reduced to two pints and a half, add the sugar, and form a syrup.

Syrupus Rosæ centifoliæ. (EDINB.)

- R. Fresh Petals of the hundred-leaved Rose, one part.
Boiling Water, four parts.
Refined Sugar, three parts.

Macerate the petals in the water for twelve hours; then add the sugar to the strained infusion, and boil to a syrup.

HISP. prescribes three parts of petals, six of boiling water, twelve hours' infusion, and two parts of sugar.—FERR.—one of petals, eight of boiling water, twenty-four hours' infusion, and six parts of sugar.

- R. Expressed Juice of }
Pale Roses, } of each,
White Sugar, } an equal part.

Boil slowly to the consistence of syrup. (GALL.)

SARD. prescribes simple syrup instead of sugar.

- R. Fresh Petals of Red Roses, one part.
Hot Water, two parts and a half.

Having reduced the petals to a pulp, macerate them in the water for twelve hours; then strain with slight expression, and add to the liquor

- White Sugar, an equal weight.
Clarify, and evaporate to a syrup. (VAN MONS)

- R. Fresh Petals of Pale Roses, twelve pounds.

Bruise them in a marble mortar; then infuse for twelve hours, in

- Water, four pounds.
Add to the expressed liquor

White Sugar, four pounds.
Clarify, and boil slowly to a syrup. To each pound of this, add another cold syrup, prepared with

- Water of Pale Roses, ten drams.
White Sugar, two ounces.
Mix. (ANT.)

- R. Sugar, two pounds.
Spring Water, a sufficient quantity.

Boil to the consistence of syrup; pour this, while boiling, upon

- Fresh Petals of Pale Roses, two pounds.

Macerate for three days; then boil a little, and express strongly. (GENEV.)

- R. Fresh Petals of Pale Roses, one pound.
Boiling Water, four pounds.

Macerate for twelve hours; then evaporate in a water-bath, constantly stirring, till the liquor is reduced to one half. Add to the expressed liquor, depurated by repose,

- White Sugar, two pounds.
Boil to the required consistence. (AMST.)

- R. Petals of Pale Roses, four pounds.
Boiling Water, eight pounds.

Infuse for twenty-four hours, and express strongly. Digest in the liquor a fresh quantity of roses. Repeat this operation a third time, and finally add to every nine ounces of the infusion,

- White Sugar, sixteen ounces,
and boil to a syrup. (BRUNS.; PALAT.; WIRTEM.)

This syrup is called *Solutive Syrup of Roses*, to distinguish it from the *Simple Syrup of Roses*, prepared with the same triple infusion, but with only thirty-two ounces of sugar to one pound and a half of liquid.

COMPOUND SYRUP OF ROSES.

Syrupus Rosarum solutivus cum Senna seu de Senna rosaceus. (AMST.; BAVAR.; HERBIP.)

- R. Fresh Petals of Pale Roses, one pound.
Boiling Water, six pounds.

Macerate for twelve hours; then boil down to four pounds, and express: macerate in the liquid for two hours

Senna, eight ounces.

Express. To the liquor, depurated by repose, add

White Sugar, two pounds.

Boil to the consistence of syrup, and add to it when cold,

Oil of Anise, }
Fennel, } of each, five drops.

Mix.

HERBIP. directs two pounds of infusion of pale roses to be poured upon one ounce of senna, and, after several hours' digestion, twice the quantity of sugar to be added to the strained liquor, and boiled to a syrup.—BAVAR.—one part of petals and two parts of senna to be infused for half an hour in eighty-eight of hot water, and two parts of sugar to be added to one part of the strained infusion.

HONEY OF ROSES.

Mel rosatum seu rosaceum seu de Rosis seu Rose Gallica, Rhodomet. (LOND.; DUBL.; AMST.; ANT.; AUST.; BATAV.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI.; COXE.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

Mel Rosæ. (LOND.; DUBL.)

R. Dried Petals of Red Roses, four ounces.

Boiling Water, three pints.

Despumated Honey, five pounds.

Macerate the petals in the water for six hours; then mix the honey with the strained liquor, and boil to the consistence of syrup, removing the scum.

R. Dried Petals of Red Roses, ten ounces.
Boiling Water, six pounds.

Macerate for twelve hours; then boil over a slow fire, and express: add to the decanted liquor

Despumated Honey, four pounds.

Evaporate slowly to the consistence of syrup. (BATAV.)

ANT.; and FORM. HÔP. FR. prescribe one part of dried petals, four parts of water, twelve hours' digestion, and six parts of honey.—BRUNS.; and WIRTEM.—six of fresh petals, six of water, twenty hours' digestion, and six parts of honey.—PALAT.—

nine of fresh petals, twelve of water boiled till reduced to five, and four of honey.—HANN.; OLDEN.; POL.; and SWEDIAUR.—eight of dried petals, four of water, twelve hours' digestion, and eight parts of honey.—FENN.; LUSIT.; and SUEC.—one half part of dried petals, three parts of water, twelve hours' digestion, and six parts of honey.—BORUS.—four of dried petals, eight of water, twelve hours' digestion, and eight of honey.—BAVAR.—one of dried petals, eight of water, three hours' digestion, and twice the weight of honey.—AUST.; and ROSS.—three parts of dried petals, twelve of water, three hours' digestion, and twenty-four parts of honey.—BRUGNATELLI.—one of dried petals, four of water, six hours' digestion, and five parts of honey.—SPIELMANN.—six of petals, six of water, twenty-four hours' digestion, and eight parts of honey.—SAX.—one of dried petals, six of water, twelve hours' digestion, and twice the weight of honey.—COXE.—four ounces of roses, three pounds of water, six hours' digestion, and five pounds of honey.—GENEV.—three ounces of petals, two pounds of water, several hours' digestion, and two pounds and a half of honey.—AMST.—eight ounces of dried petals, six pounds of water, twelve hours' maceration, and two pounds of despumated honey.

R. Petals of Red Roses, one part.
Water, eight parts.

Infuse at twice, straining each time without expression; shake the white of one or two eggs with the liquor, boil for an instant, strain through flannel, and add

White Sugar, one part.
Honey, five parts.

Evaporate to the consistence of syrup. (VAN MONS)

R. Dried Petals of Red Roses, one part.
Decoction of the Calices of Red Roses, four parts.

Infuse for two hours. Add to the strained liquor

Honey, six parts.

Clarify with white of egg, and evaporate to the proper consistence. (GALL.)

FORM. HÔP. FR. gives the same process, but substitutes water for the decoction of the calices, which introduces into the compound a great proportion of tannin and of gallic acid. HENRI also observes that the albumen of the white of egg ordered for the clarification, combines with much tannin, and thus destroys a part of the active principle of this syrup.

R. Depurated Juice, or Saturated Infusion of Red Roses, one half part.
Honey, one part.

Mix well. (HASS.)

PIDERIT directs nine ounces of saturated infusion of roses and one pound of despumated honey to be boiled to the consistence of syrup.—**HISP.**—an equal part each, of depurated juice of roses and of honey.—**SARD.**—one of depurated juice, and two of despumated honey.—**LIPP.**—one of infusion, and four of honey.—**HERBIP.**—one of infusion, and four of honey.—**FERR.**—an equal part each, of infusion and of honey.

R. Rose Water, one ounce.
Despumated Honey, four ounces.

Mix. (**DAN.**; **FERR.**; **FULD.**)

The great differences which result from the infinite diversity of these processes are of but little importance, as the syrup is scarcely ever employed except externally. It is used as an astringent in gargles and injections.

SOLUTIVE HONEY OF ROSES.

Mel rosatum solutivum. (**SARD.**; **WIRTEM.**)

R. Honey, two pounds.
Juice of Pale Roses, three pounds.

Boil together, skim, and evaporate slowly to the required consistence. (**WIRTEM.**)

SARD. prescribes an equal part of each.

It is rarely administered internally, but sometimes enters into the composition of lavements.—Dose, from one ounce and a half to two ounces.

DETERGENT GARGLE.

Gargarisma commune. (**FORM.** **HÔP. FR.**; **GENEV.**; **RATIER**; **SAINTE-MARIE**; **SPIELMANN**)

R. Acidulous Infusion of Roses, half a pound.
Honey of Roses, one ounce.

Mix. (**GENEV.**)

R. Honey of Roses, fifteen parts.
Decoction of Barley, one hundred parts.

Mix. (**FORM.** **HÔP. FR.**)

RATIER prescribes one ounce of honey of roses, and one pound of decoction of barley.

R. Leaves of Agrimony, one handful.
Sage, one pinch.
Water, one pound and a half.

Boil for a quarter of an hour; then strain and add

Honey of Roses, two ounces.

Mix. (**SAINTE-MARIE**)

R. Barley, one ounce.
Water, thirty-two ounces.

Boil down to one half, and add

Leaves of Agrimony, } of each,
Bramble, } two pinches.

Strain, and dissolve in the liquor,

Nitre, one dram.
Honey of Roses, one ounce.

Mix. (**SPIELMANN**)

DETERGENT GARGLE.

Collutorium detergens seu adstringens.
(**BRERA**)

R. Honey of Roses, one ounce.
Decoction of Barley, one pound.
Tincture of Lac, twenty drops.

Mix.

R. Honey of Roses, one ounce.
Spirit of Scurvy-grass, } of each,
HOFFMANN'S LIQUOR, } thirty drops.
Decoction of Barley, two pounds.

Mix.

R. Honey of Roses, one ounce.
Decoction of Barley, one pound.
HOFFMANN'S LIQUOR, thirty drops.

Mix.

TINCTURE OF ROSES. (**VAN MONS**)

R. Fresh Petals of Red Roses, one part.
Proof Spirit, four parts.

Macerate without heat for two days; then express and filter.

VINEGAR OF ROSES.

Acetum Rosæ seu Rosarum. (**BRUNS.**; **DAN.**; **HISP.**; **FERR.**; **HANN.**; **OLDEN.**; **LUSIT.**; **PALAT.**; **POL.**; **BORUS.**; **SARD.**; **SAX.**; **SUEC.**; **WIRTEM.**; **HERBIP.**; **VAN MONS**)

R. Petals of Red Roses, one part.
Vinegar, sixteen parts.

Macerate for fifteen days in a covered vessel, occasionally shaking; then strain and filter.

SARD. prescribes one part of roses, four parts of vinegar, six days' maceration, and a repetition of the infusion once or twice with fresh flowers.—**DAN.**; **FERR.**; **HANN.**; **OLDEN.**; **POL.**; **BORUS.**; **SAX.**; and **VAN MONS** prescribe one part of roses, eight parts of vinegar, and ten days' maceration without heat.—**HERBIP.**—one of roses, nine of vinegar, and three weeks' maceration in the sun.—**SUEC.**—one of roses, twelve of vinegar, and four days' maceration.—**BRUNS.**; **PALAT.**; and **WIRTEM.**—one of roses, and six of vinegar.

R. Dried Petals of Red Roses, one pound.
Vinegar, eight pounds.

Macerate in the sun for several days, or in a water-bath for several hours; then express, and add.

Proof Spirit, two ounces.

Mix. (**HISP.**)

LUSIT. directs one pound of roses to be

infused in six pounds of vinegar, then to be boiled a little, and half a pound of alcohol added.

COMPOUND VINEGAR OF ROSES.

Acetum Rosarum compositum. (VAN MONS)

R. Dried Flowers of Red Roses,	one part and a half.
Sumach,	one part.
Quince Wine,	four parts.
Vinegar,	ten parts.

Macerate without heat for ten days ; then express lightly and filter.

VAN MONS also prescribes one part of roses, one of sumach, and twelve of vinegar; to be infused in a moderate heat.

It is astringent and antiseptic, and is recommended externally in hæmorrhages and ecchymoses.

ROSE OIL.

Oleum Rosarum seu *rosatum*. (BRUNS.;
HISP.; GALL.; SARD.; WIRTEM.)

R. Fresh Roses, one part.
Olive Oil, two parts.

Digest for two days in a water-bath, and express when cold. Repeat the operation a second time. (SARD.)

R. Roses, pounded,	four parts.
Olive Oil,	five parts.

Expose the mixture to the sun for eight days, then boil to the consumption of the humidity, and express. (BRUNS.; HISP.; WIRTEM.)

R. Roses, pounded,	one part.
Olive Oil,	four parts.

Expose to the sun for three days, and express. Repeat the operation twice again, prolonging the last digestion for three months.

It is extolled against chilblains.

OIL OF ROSES.

Oleum Rosarum æthereum, Ætheroleum Rosarum. (GALL. ; SARD.)

R. Fresh Petals of Roses, five parts.
Water, seven parts.
Distil. (GALL.)

SARD. prescribes twenty-five parts of roses, seventy-five of water, and three of muriate of soda.

ROSE WATER.

Agua Rosæ seu *Rosarum*. (LOND.; EDINB.;
DUBL.; AMER.; AMST.; ANT.; AUSTR.;
BATAV.; BAVAR.; BELG.; DAN.; GALL.;
FERR.; FENN.; FULD.; GENEV.; HANN.;

HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.;
POL.; ROSS.; SAX.; SUEC.; WIRTEM.;
HERBIP.; BRUGNATELLI; COXE; PI-
DERIT; SWEDIAUR; VAN MONS)

Aqua Rosæ. (LOND.; DUBL.)

R. Petals of the hundred-leaved Rose, freed from their claws, eight pounds,
Water, a sufficient quantity,
to prevent empyreuma. Distil a gallon.

Aqua Rosæ Centifoliæ. (EDINB.)

R. Fresh Petals of the hundred-leaved Rose, six pounds.
Water, a sufficient quantity,
to prevent empyreuma. Distil ten pounds.

R. Fresh Rose Petals, three parts.
Water, a sufficient quantity,
to prevent empyreuma. Distil eight parts.
(FERR. ; HASS. ; PIDERIT)

LUSIT. prescribes six parts of petals and a sufficient quantity of water ; eight parts to be distilled.—FENN. ; and SUEC.—three of petals and a sufficient quantity of water ; six to be distilled.—GENEV.—four of petals and a sufficient quantity of water ; six to be distilled.—AMST. ; BATAV. ; and BELG.—one of petals and a sufficient quantity of water ; ten to be distilled.—HANN. ; OLDEN. ; and POL.—one of flowers and a sufficient quantity of water ; five to be distilled.—AMER.—eight of petals and a sufficient quantity of water ; eight to be distilled.—ANT.—one of petals and five of water ; four to be distilled.—BRUGNATELI.—three of flowers and fifteen of water, eight to be distilled.—SARD. ; and HERBIP.—one of petals and four of water ; two to be distilled.—GALL.—one of petals and four of water ; one to be distilled.—ROSS.—ten of petals and forty of water ; fifteen to be distilled.—BAVAR.—one of petals and nine of water ; three to be distilled.—SAX.—one of petals and ten of water ; four to be distilled.—PALAT. ; and WIRTEM.—one of petals and three of water ; one half to be distilled.

R. Fresh Rose Petals, three parts.
Water, twelve parts.

Enclose the petals in a linen bag suspended in the middle of a cucurbit in such a manner that it may not touch the water, and distil six parts without bringing this to ebullition. (FULD.; LIPP.)

R. Fresh Rose Petals, four parts.
Water, a sufficient quantity.

Muriate of Soda, one part.

Distil twenty parts. (SWEDIAUR)

AUSTR. prescribes one part of salted petals and eight parts of water; three parts to be distilled.—DAN.—one of salted flowers and six of water; three to be distilled.

R. Fresh Rose Petals, ten parts.
Proof Spirit, one half part.
Water, one hundred and sixty parts.

Distil forty parts. (VAN MONS)

R. Fresh Rose Petals, one pound.
Water, a sufficient quantity,
to prevent empyreuma. Macerate for two days; then distil ten pounds, and add to the product

Proof Spirit, five ounces.

Mix. (COXE)

Dose, one or two ounces.

SPIRIT OF ROSES.

Spiritus Rosarum. (BRUNS.; PALAT.; SAX.; WIRTEM.)

R. Fresh Roses, two parts.
Alcohol, } of each, one part.
Water, }

Distil one part. (SAX.)

R. Petals of Roses, one pound.
White Wine, ten pounds.
Alcohol, one pound and a half.

Distil one half, and having poured the product upon fresh petals, distil again one half. (BRUNS.; WIRTEM.)

R. Petals of Pale Roses, twelve pounds,
White Sugar, two pounds.
Yeast, half a pound.
Warm Water, thirty pounds.

Leave the mixture to ferment till a vinous odour is exhaled; then distil, and let the product be rectified. (PALAT.; WIRTEM.)

It is cordial and nerveine, but is used more externally than internally.

ACID TINCTURE OF ROSES.

Tinctura Florum Rosarum seu Rosarum acidula. (BRUNS.; OLDEN.; POL.; BORUS.; SAX.; WIRTEM.; VAN MONS)

R. Petals of Pale Roses, half an ounce.
Sulphuric Acid, one dram.
Rose Water, four ounces.

Digest for twenty-four hours; then express and filter.

HANN.; OLDEN.; BORUS.; and SAX. prescribe one ounce and a half of petals, half an ounce of acid, and twelve ounces of boiling water; to be macerated for twelve hours.—POL.—one dram of petals, one

dram of acid, and one ounce of water; to be macerated for three hours.

It is temperant.—Dose, from half an ounce to one ounce.

ROSE JULEP.

Julapium rosatum seu Rosarum seu regium, Syrupus regius seu alexandrinus. (BRUNS.; HASS.; PALAT.; WIRTEM.; PIDERIT; SPIELMANN)

R. Rose Water, twenty ounces.
Sugar, two pounds and a half.

Boil for a few moments, and strain. (PALAT.; WIRTEM.; and SPIELMANN)

BRUNS. prescribes sixteen ounces of water, and two pounds and a half of sugar.—HASS.; and PIDERIT—nine ounces of water and sixteen of sugar.

ASTRINGENT COLLYRIUM. (RATIER)

R. Rose Water, } of each,
Infusion of Melilot, } four ounces.

Mix.

DETERGENT MIXTURE. (SAINTE-MARIE)

R. Rose Water, three ounces.
Cream, } of each, one ounce.
White of Egg, }
Syrup of Violets, }

Mix.—It is recommended by BOERHAAVE to be applied to aphthæ and swelled gums.

ROSE OINTMENT.

Unguentum rosatum seu Rosarum seu pomadinum seu refrigerans seu Aqua Rose. (AMER.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SARD.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Lard, any quantity.

Triturate with a wooden pestle, gradually adding,

Rose Water, a sufficient quantity; or, till the lard becomes of a snowy whiteness. (SAX.; SUEC.)

FENN.; and LUSIT. prescribe eight parts of lard and one part of rose water.

R. Lard, thirty-two ounces.

Add to it by drops,

Rose Water, four ounces.

When the mixture is very white, add some essential oil, such as that of lavender, to it. (ROSS.)

BRUNS.; WIRTEM.; and HERBIP. pre-

scribe one pound of lard washed with rose water, and twelve drops of oil of rhodium.—SPIELMANN—twelve ounces of lard washed with rose water, and sixteen drops of oil of rhodium.—SWEDIAUR—one pound of lard washed with rose water, and one dram of oil of rhodium.

R. Lard, two pounds.
Mutton Suet, three ounces.

Melt together, and pour the mass into rose water; stir well, let the ointment cool, and dry it. One dram of oil of rhodium may be added to each pound. (BRUNS.)

AMER.; and COXE direct half an ounce of spermaceti and one dram of white wax to be melted in two ounces of oil of almonds, and two ounces of rose water to be stirred with the mixture till cold.

R. Lard, washed with rose water, eight ounces.
White Wax, two ounces.

Melt slowly together, and when cold, add gradually,

Rose Water, three ounces.
Oil of Lemon, one dram.

Mix well. (HANN.; POL.; and BORUS.)

VAN MONS directs one part of white wax to be melted and added to four parts of lard, and one ounce and a half of rose water to be gradually incorporated in the mixture.—FULD.—thirty-two ounces of lard to be triturated with four ounces of rose water, the mixture then melted, the water separated, and the lard triturated with thirty drops of oil of lemon.—LIPP.—one pound of lard to be triturated with two ounces of rose water and two ounces of orange-flower water, the mixture to be then melted, the water separated, and the lard incorporated with twenty-four drops of oil of bergamotte and fifteen of oil of lavender.

R. Lard, one pound.
Rose Water, } of each,
Orange-flower Water, } two ounces.
Oil of Lemon, twenty-four drops.
— Lavender, fifteen drops.
— Cloves, twelve drops.

Mix. (HASS.; OLDEN.; PIDERIT)

SWEDIAUR prescribes one pound of lard, half an ounce of orange-flower water, three ounces of rose water, twelve drops of oil of lemon, and twelve of oil of cloves.

R. Lard, ten ounces.
White Wax, two ounces.

Melt together, and add,

Rose Water, three ounces.
Orange-flower Water, one ounce.

Set aside for twenty-four hours; then pour off the water, and triturate the lard with

Oil of Lavender, } of each, half a dram.
— Lemon, }

Mix. (NIEMANN)

R. Lard, washed with rose water, twelve ounces.
Rose Water, three ounces.
Orange-flower Water, } of each,
Spirit of Roses, } half an ounce.
Oil of Lavender, half a dram.
— Rhodium, } of each,
— Lemon, } sixteen drops.

Mix. (SPIELMANN)

R. Rose Oil, one pound.
White Wax, three ounces.

Melt together, wash the mixture in water, and agitate it well that it become white. (WIRTEM.)

SARD. prescribes one pound and a half of rose oil and six ounces of wax.

R. Lard, three pounds.
Storax, } of each, one ounce.
Benzoin, }
Iris, }
Lemons, } of each, two.
Oranges, }
Rennet Apples, four.

Boil to the consumption of the humidity, and add,

Oil of Cinnamon, } of each,
— Cloves, } twenty-five drops.

Beat the whole together with spirit of roses. (PALAT.)

R. Cloves, two drams.
Cinnamon, } of each, three drams.
Benzoin, }
Storax, }
Mace, two scruples.
Lavender, }
Rhodium Wood, } of each, one dram.
Aloe Wood, }
Sassafras Wood, half a dram.
Florentine Iris, one ounce.
Rose Water, half a pound.
Lard, one pound.
Pulp of Rennet Apples, six ounces.
Goat's Suet, four ounces.

Boil slowly, and then digest for three days; then boil again till all the water is evaporated: agitate the strained liquid with rose water till it becomes white, and add,

Oil of Rhodium, twenty drops.

Mix. (NIEMANN)

This formula is by CARTHEUSER. It is to this preparation that the term pomatum or pommadine ointment (*pommade ou onguent pommadin*) is applied.

R. Lard, washed with rose water, } of each,
Pale Roses, bruised, } an equal part.

Leave together for two days, then melt the lard, and macerate it with an equal quantity

of roses for two other days; then liquify again in a water-bath, and set it aside. (GALL.)

BRUGNATELLI directs two pounds of lard and three ounces of suet to be melted together, two ounces of red roses and half an ounce of alkanet to be added, the mass to be strained, and stirring till it becomes cold. —SARD.—two pounds each, of red roses and of pale roses to be digested in six pounds of lard for three days in a water-bath, the liquid to be expressed, six ounces of wax added, melted, and the mixture set apart, decanted, and finally cooled.

R. Lard, twelve pounds.
Petals of Red Roses, six pounds.

Macerate for three days, then boil to the consumption of the humidity, and express. (HISP.)

PALAT.; and WIRTEM. direct one pound of lard to be boiled with four ounces of roses to the consumption of the humidity; the liquid to be then expressed, and three grains of essence of roses added.

R. Lard, nine pounds.
Mutton Fat, one pound.
Fresh Rose Petals, nine pounds.

Digest the petals for one day in the fats melted together, and express. (FERR.)

SPECIES OF HYACINTH. (BRUNS.; PALAT.; WIRTEM.)

R. Prepared Hyacinth,	} of each, one dram and a half.
Armenian Bole,	
Lemnian Bole,	
Prepared Red Coral,	} of each, one dram.
———— Pearls,	
———— Emeralds,	
———— Hartshorn,	} of each, four scrup- les.
———— Ivory Shavings,	
———— Bone of Ox-heart,	
Root of Fraxinella,	
———— Tormentil,	
White Sanders,	
Yellow Sanders,	
Red Sanders,	
Petals of Red Roses,	
Kermes Grains,	
Seeds of Sorrel,	} of each, half an ounce.
———— Lemon,	
———— Purslain,	
Saffron,	
Myrrh,	} of each, two drams.
———— Sweet Basil,	
Cinnamon,	
Spikenard,	
Rhubarb,	} of each, one dram.
Saffron,	
Ivory Shavings,	
Prepared Hartshorn,	
Mastic,	} of each, one scruple.
Prepared Mother of Pearl,	

These are called *completa*, when the addition is made of

Ambergris, } of each, four scruples.
Musk, }

Make a powder.

It is slightly astringent. — Dose, from half a scruple to one scruple.

POWDER AGAINST ABORTION.

Pulvis contra abortum. (PALAT.; WIRTEM.)

R. Kermes Grains, half an ounce.
Red Sanders, }
Petals of Red Roses, } of each,
Prepared Red Coral, } three drams.
Ivory Shavings, }
Mastic, } of each, one dram.
Nutmegs, }
White Sugar, }
an equal weight of the whole.

Mix. (WIRTEM.)

PALAT. prescribes one dram of kermes, half a dram of red sanders, one scruple of cloves, two scruples of nutmeg, half a dram of tormentil, one scruple each, of pearls, of red coral, and of white coral, and three drams of sugar. — Dose, one dram.

In common with all stimulants, this powder would be more likely to occasion than to prevent abortion.

RED CEPHALIC POWDER. (VAN MONS)

R. Red Roses, six ounces.
Tartaric Acid, half an ounce.
Florentine Iris, six ounces.

Pulverize, and moisten the mixed powders with rose water.

ODORIFEROUS POWDER.

Pulvis nasalis odoratus. (PALAT.)

R. Petals of Red Roses, six drams.
Flowers of Star-thistle, } of each,
———— Marigold, } one dram.
———— Wallflower, }
Florentine Iris, two drams.
Marjoram, } of each, one scruple.
Wild Thyme, }
Thyme, }

Add to the powder,

Oil of Rhodium Wood, four drops.
———— Cloves, six drops.

Mix.

DIARRHODON POWDER.

Pulvis seu Species Diarrhodon. (PALAT.; SARD.; WIRTEM.)

R. Petals of Red Roses, two ounces.
White Sanders, } of each, half an ounce.
Red Sanders, }
Liquorice, } of each, two drams.
Seeds of Anise, }
———— Sweet Basil, }
Cinnamon, }
Spikenard, }
Rhubarb, } of each,
Saffron, } one dram.
Ivory Shavings, }
Prepared Hartshorn, }
Mastic, }
Prepared Mother of Pearl, }

Make a powder. (PALAT.; WIRTEM.)

- | | | |
|-------------------|-------------|-----------|
| R. Red Roses, | two ounces. | |
| Yellow Sanders, | one ounce. | |
| Lesser Cardamoms, | | |
| Cinnamon, | | |
| Saffron, | | |
| Mastic, | | |
| Spikenard, | | |
| Fennel, | | |
| | | of each, |
| | | one dram. |

Make a powder. (SARD.)

It is stomachic.—Dose, from half a dram to two scruples.

SPECIES FOR AN EPITHEM.

Species pro Epithemate Cordis et Pulsuum. (PALAT.; WIRTEM.; SPIELMANN)

- | | | |
|-------------------------------|--|----------------|
| R. Herb of Rosemary, | | |
| — Marjoram, | | |
| — Balm, | | |
| Flowers of Yellow Wallflower, | | |
| — Clove-pink, | | |
| — Lavender, | | |
| — Roses, | | |
| Mace, | | |
| Nutmegs, | | |
| Aloe Wood, | | |
| Rhodium Wood, | | |
| Cinnamon, | | |
| Lemon Peel, | | |
| Orange Peel, | | |
| | | of each, |
| | | half an ounce. |
| | | of each, |
| | | six drams. |
| | | of each, |
| | | three drams. |

Cut, bruise, and mix. (WIRTEM.)

SPIELMANN directs the mixture to be powdered.

- | | | |
|------------------------|--|----------------------|
| R. Flowers of Bugloss, | | |
| — Borage, | | |
| — Balm, | | |
| White Sanders, | | |
| Red Sanders, | | |
| Yellow Sanders, | | |
| Petals of Red Roses, | | |
| Lemon Peel, | | |
| Seeds of Sorrel, | | |
| Aloe Wood, | | |
| Cloves, | | |
| | | of each, |
| | | half a handful. |
| | | of each, |
| | | two drams. |
| | | of each, |
| | | one dram and a half. |

- | | | |
|--------------|--|----------------|
| Doronica, | | |
| Red Behen, | | |
| White Behen, | | |
| Amber, | | |
| | | of each, |
| | | four scruples. |

Make a powder. (PALAT.)

SPECIES AGAINST GANGRENE.

Species pro Decocto ad Gangrænam. (PALAT.; WIRTEM.; VAN MONS)

- | | | |
|-------------------------|--|--------------------|
| R. Leaves of Horehound, | | |
| — Pennyroyal, | | |
| — Rue, | | |
| — Rosemary, | | |
| — Sage, | | |
| — Scordium, | | |
| Flowers of Chamomile, | | |
| — Red Roses, | | |
| — Elder, | | |
| — Marshmallows, | | |
| Seeds of Fenugreek, | | |
| — Flax, | | |
| | | of each, |
| | | one part. |
| | | of each, |
| | | two parts. |
| | | six parts. |
| | | eight parts. |
| | | twenty-four parts. |

Cut, bruise, and mix. (VAN MONS)

- | | | |
|---------------------|--|----------------|
| R. Herb of Milfoil, | | |
| — Wormwood, | | |
| — Scordium, | | |
| — Sage, | | |
| — Hyssop, | | |
| — Rue, | | |
| Flowers of Elder, | | |
| — St. John's-wort, | | |
| — Roses, | | |
| Myrrh, | | |
| Aloes, | | |
| Assafoetida, | | |
| Litharge, | | |
| | | of each, |
| | | six drams. |
| | | of each, |
| | | half an ounce. |
| | | one ounce. |
| | | six ounces. |

Cut, bruise, and mix. (PALAT.; WIRTEM.)

These species are boiled in five pounds of water and two of vinegar, and the decoction applied by means of flannel.—From four to six ounces of camphorated spirit may be added.—At the present day emollients, instead of stimulants, are preferred to be employed against gangrene.

ROSMARINUS OFFICINALIS, LINN.—ROSEMARY.

SYNONIMES:—Romarin, FR.; Rosmarin, GERM.; DAN.; and SWED.; Hasalban achsir, ARAB.; Rosmaryn, BOHEM.; DUT.; and POL.; Yong tsao, CHIN.; Tay-duong-choi, COCHIN.; Romero, SPAN.; Ramerino, Rosmarino, ITAL.; Alecrim, PORT.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; VOL. II.

FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of the south of Europe. (*Dian-dria Monogynia*, LINN.; *Labiata*, JUSS.; *Fig. Flore Medic. VI. 300*)

The flowering herb (*Herba et Flores An-3 X*)

thos seu Rosmarini seu Rorismarini hortensis) is employed. It consists of slender, long, ash-coloured branches, bearing sessile, opposite, rather hard, narrow, linear leaves, green above, whitish beneath, rolled at the edges, and furnished with pale blue flowers disposed in small tufted racemes, axillary, opposite. It has a very penetrating, agreeable odour, and an aromatic, camphorous, and subacid taste.

Internally it is stimulant, externally resolvent.—Dose, from ten grains to two scruples of the powder.

CONSERVE OF ROSEMARY.

Conserva Anthos. (AMST. ; BATAV. ; BELG. ; SARD. ; WIRTEM. ; CADET DE GASSICOURT)

R. Flowers of Rosemary, bruised, one part.
White Sugar, three parts.

Mix. (AMST.)

BATAV. ; BELG. ; and WIRTEM. prescribe one part of flowers and two parts of white sugar.—SARD.—one of flowers and three of sugar.—CADET DE GASSICOURT—one of flowers in powder, four of sugar, and a sufficient quantity of water.

AROMATIC POWDER. (RATIER)

R. Rosemary, } of each, four ounces.
Sage, }
Thyme, }
Muriate of Ammonia, } of each,
Camphor, } twenty grains.

Pulverize, and mix.—DUPUYTREN applies it, instead of cinchona, in gangrene and hospital gangrene.

AROMATIC SPECIES. (FULD.)

R. Root of Florentine Iris, }
Zedoary, } of each,
Herb of Balm, } six drams.
Rosemary, }
Wild Thyme, }
Flowers of Lavender, }
Cloves, } of each, two drams.
Nutmeg, }
Cinnamon, }

Mix.—They are resolvent, nervine, and corroborant. To be applied in bags.

DIANTHOS SPECIES. (WIRTEM.)

R. Root of Liquorice, six drams.
Ginger, }
Zedoary, } of each,
Galanga, } four scruples.
Aloe Wood, }

Cinnamon,	}	of each, four scruples.
Cloves,		
Lesser Cardamoms,		
Mace,		
Nutmegs,	}	one ounce.
Spikenard,		
Flowers of Rosemary,	}	of each, six drams.
— Roses,		
— Violets,		
Seeds of Anise,	}	of each, four scruples.
— Fennel,		

Make a powder.—Dose, from half a dram to one dram.—They are cephalic, stomachic, and carminative.

EXTRACT OF ROSEMARY.

Extractum Rorismarini. (HERBIP.)

R. Flowering Tops of Rosemary, } of each,
Alcohol, } one pound.
Water, } eight pounds.

Digest for three days at a moderate temperature ; then express, and set aside and decant the liquor : distil off the alcohol, and evaporate the remainder to the proper consistence.

INFUSION OF ROSEMARY. (FORM. HÔP. FR.)

R. Flowering Tops of Rosemary, } of each,
Liquorice, } two drams
Boiling Water, } and a half.
a sufficient quantity

to obtain two pounds of strained liquor.

Dose, one or two ounces.

HONEY OF ROSEMARY.

Mel Rorismarini seu anthosatum. (FULD. ; PALAT. ; SARD. ; WIRTEM. ; SPIELMANN)

R. Fresh Flowers of Rosemary, eight ounces.
Spring Water, four ounces.
Warm Despumated Honey, two pounds.

Digest for ten or twelve days in a warm place ; then boil in a water-bath, and express. (PALAT. ; WIRTEM.)

SARD. directs four pounds of honey and two pounds of flowers to be digested and then boiled in a water-bath.—SPIELMANN—eight ounces of dried herb to be digested for several days with twenty-four ounces of despumated honey.—FULD.—one part of infusion of rosemary to be mixed with four parts of despumated honey.

It is excitant, cephalic, anti-colical, carminative, and uterine.—It may be added to lavements.

AROMATIC BATH. (CADET DE GASSI-COURT)

R. Rosemary, }
Thyme, } of each, half a pound.
Sage, }
Origanum, }
Mint, }
Water, a sufficient quantity.

Add to the decoction,

Tincture of Soap, four ounces.
Muriate of Ammonia, two ounces.

Dissolve, and mix.

VINEGAR OF ROSEMARY.

Acetum Rorismarini. (GALL.)

R. Flowers of Rosemary, one part.
Vinegar, sixteen parts.

Macerate for fifteen days in a covered vessel, occasionally agitating; then strain and filter.

AROMATIC VINEGAR.

Acetum seu Acidum aceticum aromaticum seu odoratum. (EDINB.; DAN.; FERR.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; SUEC.; WIRTEM.; COXE; FEE; HUFELAND; SPIELMANN)

Acidum aceticum aromaticum.
(EDINB.; COXE)

R. Dried Tops of Rosemary, } of each,
— Leaves of Sage, } one
— Flowers of Lavender, } ounce.
Cloves, half an ounce.
Dilute Acetic Acid, two pounds.

Macerate for seven days, and filter the expressed liquor.

LUSIT.; SUEC.; FEE; and SWEDIAUR prescribe four ounces each, of rosemary and of sage, two ounces of lavender, two drams of cloves, and eight pounds of vinegar.

R. Herb of Wormwood, }
— Rosemary, } of each,
— Sage, } two ounces.
— Peppermint, }
Cinnamon, half an ounce.
Cloves, }
Nutmegs, } of each, two drams.
Vinegar, eight pounds.

Digest in a sand-bath, express and filter. (DAN.; OLDEN.)

R. Root of Zedoary, } of each,
Cloves, } half an ounce.

Herb of Wormwood, }
— Rosemary, } of each,
— Sage, } two ounces.
— Peppermint, }

Vinegar, eight pounds.

Macerate. Add to the expressed liquor
Spirit of Rosemary, four ounces.

Mix. (HANN.; POL.; BORUS.; HUFELAND)

R. Root of Round Cyperus, } of each,
— Lesser Galanga, } half an
— Zedoary, } ounce.

Herb of Rosemary, }
— Sage, } of each,
Flowers of Lavender, } three
— Lily of the Valley, } drams.
— Red Roses, }

Rhodium Wood, }
Cinnamon, } of each.
Cloves, } two drams.
Fresh Lemon Peel, }
Vinegar, two pounds.

Digest for three days, and filter. (WIRTEM.; SPIELMANN)

R. Root of Zedoary, } of each,
— Galanga, } half an ounce.
Herb of Rosemary, }
— Rue, } of each,
— Marum, } two drams.

Flowers of Lavender, }
Fresh Lemon Peel, } of each,
Cloves, } half an ounce.
Cubebs, }
Lesser Cardamoms, }
Vinegar, forty ounces.
— of Lily of the }
Valley, } of each,
— Clove-pink, } two ounces.

Macerate, and strain. (SPIELMANN)

R. Tops of Rosemary, } of each,
— Wormwood, } one ounce and
— Sage, } a half.
— Rue, }
Sweet Flag, } of each, two drams.
Cinnamon, }
Cloves, }
Vinegar, six pounds.

Infuse for eight days. Add to the expressed and strained liquor,

Camphor, one ounce.

Two drams of garlic may be added. (FERR.)

This last formula furnishes a preparation which much resembles the *Thieves' Vinegar*.

ESSENCE OF ROSEMARY.

Essentia Rorismarini. (BRUNS.)

R. Flowering Tops of Rosemary, one ounce and a half.
Spirit of Rosemary six ounces.

Digest, express, and filter.

It is excitant.

AROMATIC FOMENTATION.

Fomentum aromaticum. (AUGUSTIN)

- R. Herb of Rosemary, half an ounce.
 Red Wine, }
 Water, } of each, three ounces.
 Infuse, and express.

INFUSED OIL OF ROSEMARY.

Oleum Rorismarini. (HISP.)

- R. Flowers, one part.
 Olive Oil, three parts.

Digest for three or four days; then boil to the consumption of the humidity, and express.

ESSENTIAL OIL OF ROSEMARY.

Oleum Anthos athereum, Aetheroleum Anthos seu Rorismarini. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; HERBIP.; BRUGNATELLI; COXE; SWEDIAUR)

Oleum Rosmarini. (LOND.; DUBL.)*Ol. Volatile Rorismarini Officinalis.* (EDINB.)

- R. Tops of Rosemary, any quantity.
 Water, a sufficient quantity
 to prevent empyreuma. Distil.

AMER.; AMST.; BATAV.; BELG.; FERR.; GENEV.; ROSS.; SUEC.; BRUGNATELLI; and COXE prescribe the same.

ANT. prescribes three parts of flowers, and ten of water.—GALL.—five of flowers, and seven of water.—FULD.; and LIPP.—one of flowers, and three of water.—SWEDIAUR—one of flowers, and four of water.—DAN.—one of flowers, and six of water.—HANN.; OLDEN.; POL.; BORUS.; and SAX.—one of flowers, and eight of water.—BAVAR.—one of flowers, and nine of water.—AUSTR.—one of flowers, and twelve of water.

- R. Flowering Herb of Rosemary, twenty-five parts.
 Water, seventy-five parts.
 Muriate of Soda, three parts.
 Macerate for three days, and distil. (BRUNS.; PALAT.; SARD.)

- R. Herb of Rosemary, any quantity.
 Distilled Water of Rosemary, a sufficient quantity.

Distil. (HERBIP.)

Dose, from two to five drops.

NERVINE BALSAM.

Balsamum apoplecticum seu aromaticum seu cephalicum Saxonicum seu nervinum seu Scherzeri seu stomachicum, Linimentum aromaticum stimulan, Unguentum aromaticum seu nervinum. (BRUNS.; DAN.; GALL.; FENN.; FULD.; HANN.; HASS.; LIPP.; PALAT.; SAX.; SUEC.; WIRTEM.; HERBIP.; CADET DE GASSICOURT; NIEMANN; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

1. With Balsam of Peru.

- R. Expressed Oil of Nutmeg, one ounce.
 Oil of Rosemary, } of each, one dram.
 — Cloves, }
 — Amber, half a dram.
 Balsam of Peru, one dram.

Triturate together. (LIPP.)

PIDERIT prescribes one ounce of oil of nutmeg, one dram each, of oil of cloves and of oil of lavender, one scruple of oil of amber, and one dram and a half of balsam.—SAX.—one ounce of oil of nutmeg, one dram each, of oil of cloves and of oil of rosemary, one scruple of oil of amber, and one dram and a half of balsam.—HASS.—one ounce of oil of nutmeg, one dram each, of oil of cloves and of oil of peppermint, one scruple of oil of amber, and one dram and a half of balsam.—VAN MONS—three ounces of oil of nutmeg, half an ounce each, of oil of cloves, of oil of lavender, and of oil of amber, and two drams of balsam.—SWEDIAUR—three drams of oil of nutmeg, one scruple each, of oil of cloves and of oil of amber, and one dram and a half of balsam; or, one ounce of oil of nutmeg, one dram each, of oil of cloves and of oil of lavender, one scruple of oil of amber, and one dram and a half of balsam.—DAN.—one ounce of oil of nutmeg, half a dram each, of oil of cloves, of oil of lavender, and of oil of rosemary, half a scruple of oil of amber, and one dram of balsam.

- R. Oil of Nutmeg, three parts.
 Oil of Cloves, }
 — Benzoin, } of each one part.
 Balsam of Peru, }

Mix. (NIEMANN)

- R. Oil of Nutmeg, one ounce.
 — Wormwood, }
 — Orange, } of each,
 — Cloves, } half a dram.
 — Mace, }
 — Mint, }
 Balsam of Peru, one dram.

Mix. (PALAT.; WIRTEM.)

HANN. gives the same formula, but omits the oil of orange.

- R. Oil of Nutmeg, one ounce.
 — Rosemary, } of each, half a dram.
 — Lavender, }
 — Cloves, }
 — Amber, }
 Musk, triturated with } of each,
 Alcohol, } half a scruple.
 Balsam of Peru, one dram.

Mix. (BRUNS.)

- R. Oil of Nutmeg, three ounces and a half
 — Jessamine, } of each,
 — Cinnamon, } one dram and
 — Rhodium, } a half.
 — Lavender, } of each, one dram.
 — Marjoram, }
 — Cloves, }
 — Rue, } of each,
 — Amber, } one scruple.
 Musk, } of each,
 Ambergris, } sixteen grains.
 Civet, } half a scruple.
 Balsam of Peru, two scruples.
 Burnt Ivory, a sufficient quantity.

Mix. (WIRTEM.; SPIELMANN)

PALAT. prescribes one ounce of oil of nutmeg, one scruple each, of oil of rosemary, of oil of cloves, and of oil of cinnamon, half a scruple each, of oil of lavender, of oil of rhodium, of oil of marjoram, of oil of rue, and of oil of amber, one dram and a half of balsam, and a sufficient quantity of soot of amber.

- R. Oil of Palm, }
 — Nutmeg, } of each,
 Stag's Marrow, } four ounces.
 Beef Marrow, }
 Human Fat, } of each,
 Viper's Fat, } one ounce.
 Badger's Fat, }

Melt slowly, and add.

- Oil of Lavender, }
 — Mint, } of each, one dram.
 — Rosemary, }
 — Sage, }
 — Thyme, }
 — Cloves, }
 Camphor, two drams.
 Balsam of Peru, one ounce.

Mix. (SPIELMANN)

- R. Beef Marrow, } of each,
 Oil of Nutmeg, } four ounces.

Melt together. Pour the mass into a wide mouthed bottle, and add

- Oil of Rosemary, two drams.
 — Cloves, } of each, one dram.
 Camphor, }
 Balsam of Peru, dissolved in half
 an ounce of alcohol (36 de-
 grees), two drams.

Melt together in a water-bath. (GALL.; CADET DE GASSICOURT)

2. Without Balsam of Peru.

- R. Oil of Nutmeg, one ounce.
 Yellow Wax, two drams.
 Melt slowly, and add
 Oil of Cloves, } of each,
 — Lavender, } two drams.
 — Amber, } half a scruple.

Mix. (FENN.; SUEC.)

- R. Mutton Suet, two pounds.
 Oil of Chamomile, } of each,
 — Marjoram, } fourteen ounces.

Melt together, withdraw it from the fire, and add

- Oil of Rosemary, two ounces.

Mix. (HERBIP.)

- R. Oil of Nutmeg, one ounce.
 — Amber, } of each,
 — Cloves, } half a dram.
 — Lavender, } of each, one dram.
 — Juniper, }

Mix without heat. (FULD.)

- R. Oil of Nutmeg, one hundred and twenty parts.
 — Lavender, } of each, four parts.
 — Amber, }
 — Rosemary, } six parts.
 — Marjoram, } of each,
 — Peppermint, } three parts.
 — Mace, } two parts.

Triturate together. (NIEMANN)

CADET DE GASSICOURT prescribes thirty-three drams of oil of nutmeg, one dram and a half each, of oil of lavender and of oil of amber, one dram each of oil of origanum, of oil of sage, of oil of marjoram, and of oil of rosemary, and two scruples each, of oil of rue, of oil of mace, and of oil of mint. —SPIELMANN—one ounce and a half of oil of nutmeg, two drams each, of oil of marjoram, of oil of lavender, of oil of rosemary, of oil of amber, and of oil of wild thyme, and one scruple each, of oil of rue, of oil of penny-royal, of oil of balm, of oil of mint, and of oil of cardamoms.—BRUNS.—two ounces and a half of oil of nutmeg, half an ounce of oil of mint, one dram each, of oil of orange, of oil of cloves, and of oil of mace, half a dram each, of oil of rosemary, of oil of lavender, of oil of wormwood, of oil of cumin, and of oil of dill, a small quantity of rose oil, and half an ounce of white wax.—WIRTEM.—one scruple each, of oil of lavender and of oil of amber, fifteen grains each, of oil of penny-royal, of oil of marjoram, of oil of sage, of oil of rosemary and of oil of savory, half a scruple each, of oil of rue and of oil of mace, twelve grains each, of oil of mint and of oil of origanum, and half an ounce of oil of nutmeg.

- R. Oil of Nutmeg, one ounce.

Oil of Amber,	half a dram.
— Roses,	of each,
— Cinnamon,	one scruple.
— Lavender,	of each,
— Marjoram,	fifteen grains.
— Rue,	of each,
— Benzoin,	half a scruple.
— Cloves,	of each, four grains.
— Lemon,	of each, four grains.
Ambergris,	one dram and a half.
Civet,	four scruples.
Musk,	one dram.

Mix. (SWEDIAUR)

All these preparations are intrinsically the same; they scarcely differ from HOFFMANN'S Balsamum Vitæ except by the absence of alcohol. They are employed externally in frictions to the abdomen, in flatulent colic, obstinate diarrhœa, and the gripes of infants; to the temples, in cephalalgia; to the hands held before the eyes in weakness of sight and lippitude; and to the limbs of weak children. A few drops are also taken with sugar, in dyspepsia.

HOFFMANN'S BALSAMUM VITÆ.

Balsamum Vitæ Hoffmanni seu ad modum Hoffmann, Mixtura oleosa-balsamica seu Succinea, Liquor balsamico-aromaticus, Tinctura Succini aromatica. (BRUNS.; DAN.; OLDEN.; BORUS.; SAX.; SUEC.; WIRTEM.; AUGUSTIN.; BORIES.; CADET DE GASSICOURT.; NIEMANN.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

R. Oil of Rosemary,	half an ounce.
— Amber,	two drams.
Alcohol,	one pound.

Mix. (BORIES)

R. Dried Flowers of Lavender,	of each,
— Leaves of Marjoram,	one
— Rue,	ounce.
Fresh Lemon Peel,	half an ounce.
Lesser Cardamoms,	of each,
Cloves,	three drams.
Mace,	of each,
Proof Spirit,	thirty-two ounces.
Water,	sixty-four ounces.

Distil eighteen ounces of liquid, and dissolve in this product

Balsam of Peru,	one dram and a half.
Oil of Amber,	half a dram.

Preserve. (VAN MONS)

R. Oil of Lavender,	
— Marjoram,	
— Cloves,	of each,
— Mace,	one scruple.
— Cinnamon,	
— Lemon,	
— Rue,	ten drops.
Balsam of Peru,	half a dram.
Concentrated Alcohol,	ten ounces.

Digest without heat and filter through cotton. (OLDEN.; BORUS.; NIEMANN.)

SAX. prescribes one part each, of the oils of lavender, of marjoram, of cloves, of mace, of cinnamon, and of lemon, one half part each, of the oils of rue and of amber, one part and a half of balsam of Peru, and two hundred and forty parts of concentrated alcohol; to be infused for four days without heat.—DAN.—one scruple each, of the oils of lavender, of cloves, of mace, and of cinnamon, ten drops of oil of amber, half a dram of balsam of Peru, ten ounces of concentrated alcohol, and a little alkanet to colour the tincture.—BORIES.—ten drops each, of the oils of cinnamon, of lavender, of cloves, of lemon, of rosemary, and of amber, twenty drops of balsam of Peru, and two ounces of rectified spirit.

R. Balsam of Peru,	
Oil of Lavender,	
— Marjoram,	
— Cloves,	of each,
— Lemon,	one scruple.
— Mace,	
— Cinnamon,	
— Rue,	
— Amber,	
Ambergris,	

Mix together, and dissolve in

Concentrated Alcohol, ten ounces.

Preserve. (BRUNS.; WIRTEM.)

AUGUSTIN prescribes fifteen grains of balsam of Peru, ten drops each, of the oils of lavender, of marjoram, of cloves, of wild thyme, of rue, and of amber, and five ounces of alcohol.—SPIELMANN—one scruple each, of the oils of lavender, of marjoram, of cloves, of cubebs, of cardamoms, of cinnamon, and of lemon, two scruples of oil of mace, half a scruple each, of the oils of rue and of amber, of the balsam of Peru, and of ambergris, and twenty ounces of alcohol.—CADET DE GASSICOURT—one scruple each, of the oils of lavender, of marjoram, of cloves, of mace, of cinnamon, and of lemon, half a scruple each, of the oils of rue and of amber, and of ambergris, and five ounces of alcohol.—HUFELAND; and NIEMANN—half a dram each, of the oils of lavender, of rosemary, of bergamotte, of juniper, and of turpentine, and ten ounces of alcohol.—SWEDIAUR—ten drops each of the oils of lavender, of marjoram, of cloves, of mace, of cinnamon, of lemon, of rue, and of amber, half a dram of balsam of Peru, and ten ounces of alcohol; ten grains of musk and fifteen of ambergris may be added.—LIPP.—one scruple each, of the oils of rosemary, of marjoram, of cloves, of cinnamon, of rue, and of lemon, half a scruple of oil of amber, half a dram of balsam of Peru, ten grains of ambergris, fifteen grains of musk, and ten ounces of alcohol; to be digested without heat for four days.

- R. Oil of Amber, one part.
 Balsam of Peru, three parts.
 Lavender Water, } of each, one hundred
 Eau de Cologne, } and twenty parts.

Dissolve, digest for several days, and filter. (VAN MONS)

- R. Oil of Cloves, }
 — Cinnamon, } of each,
 — Lavender, } one scruple.
 — Nutmegs, }
 Tincture of Amber, four ounces.

Mix. (SUEC.)

It is excitant, nervine, cephalic, and stomatic.—Dose, from ten to twenty drops.—It is also used as an embrocation.

We have placed this preparation in the article of rosemary, although the oil of rosemary does not enter into all the formulae, that it might not be separated from the preceding ones, to which it has a great resemblance.

OPHTHALMIC BALSAM. (AUGUSTIN)

- R. HOFFMANN'S Balsam, one ounce.
 Solution of Ammonia, one dram.
 Oil of Chamomile, twenty drops.
 Concentrated Alcohol, half an ounce.

Mix.—It is recommended by LEBEL DE LEBENSTEIN, in lotions to the forehead, eyelids, and cheeks, in weakness of the sight.

COMPOUND OINTMENT OF ROSEMARY.

Unguentum Rorismarini compositum. (CAST. BORUS.)

- R. Yellow Wax, one ounce.
 Lard, three ounces.

Melt, and add to the mixture when half cold,

- Oil of Rosemary, }
 Subcarbonate of } of each,
 Ammonia, } half an ounce.

Mix. (CAST. BORUS.)

AROMATIC OINTMENT.

Unguentum aromaticum. (AUGUSTIN)

- R. Compound Ointment of Rosemary, two ounces.

- Oil of Lavender, }
 — Cajeput, } of each,
 — Mint, } one scruple.
 — Chamomile, }

Mix.—It is useful in asthenic, nervous, and paralytic affections.

DISTILLED WATER OF ROSEMARY.

Aqua Anthos. (AMST.; AUSTR.; BATAV.; BELG.; VAN MONS)

- R. Flowering Herb of Rosemary, one part.

Water, a sufficient quantity.

Infuse for twenty-four hours, and distil ten parts. (AMST.; BATAV.; BELG.)

WIRTEM. prescribes one part of herb and six parts of water.—AUSTR.—one of herb and twelve of water; four to be distilled.

SPIRIT OF ROSEMARY.

Spiritus Anthos seu Rorismarini. (LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; SPIELMANN; SWEDIAUR)

Spiritus Rosmarini. (LOND.)

- R. Fresh Tops of Rosemary,

two pounds.
 Proof Spirit, one gallon.
 Water, a sufficient quantity,

to prevent empyreuma. Macerate for twenty-four hours; then distil a gallon with a gentle heat.

Spiritus Rorismarini officinalis. (EDINB.)

- R. Fresh Tops of Rosemary,

two pounds.
 Alcohol (sp. gr. 835), eight pounds.

Distil seven pounds in a water-bath.

Spiritus Rorismarini. (DUBL.)

- R. Fresh Tops of Rosemary,

one pound and a half.
 Proof Spirit, one gallon.

Distil five pints with a moderate heat.

AMER.; and COXE give the same formula, as LOND.—HANN.; OLDEN.; PALAT.; BORUS.; and SAX. prescribe one part of flowers, four parts of alcohol, and a sufficient quantity of water.—BRUGNATELLI—one pound two ounces of flowers and eight pounds of proof spirit, three fourths to be distilled.—SUEC.—one part of flowers, four parts of alcohol, and two of water; two to be distilled.—FENN.—one of flowers, two of alcohol, and two of water; two to be distilled.—HISP.—one of flowers and two of proof spirit; one to be distilled.

- R. Flowers of Rosemary, twelve parts.
 Alcohol, six parts.
 Water, two parts.

Distil slowly six parts, and pour the product upon,

Fresh Flowers of Rosemary, twelve parts.
Water, two parts.

Distil six parts. (SWEDIAUR)

R. Flowering Herb of Rosemary, twelve parts.
Alcohol (22 degrees), thirty-six parts.
Distilled Water of Rosemary, twelve parts.

At the end of four days distil thirty parts. (GALL.)

ROSS. prescribes half a pound of rosemary, one pound of rosemary water, and four pounds of alcohol; three pounds to be distilled.

R. Flowering Herb of Rosemary, one part.
Alcohol (20 degrees), four parts.

Digest, and distil a spirit marking 28 degrees. (AMST.; ANT.)

FULD.; GENEV.; and LIPP. prescribe one part of flowers, and six parts of alcohol; four parts to be distilled.—SPIELMANN—one of flowers and three of alcohol.—BRUNS.; PALAT.; and WIRTEM.—one of flowers and four of alcohol.—LUSIT.—two and a half of flowers, and ten of alcohol; five to be distilled.—FERR.—one and a half of flowers and eight of alcohol; six to be distilled.

R. Dried Flowers of Rosemary, one part.
Alcohol, four parts.

Digest for twenty-four hours, and distil in a water-bath nearly to dryness; pour the product upon fresh flowers, and re-distil. (SARD.)

DAN. directs two thirds of a mixture of one pound of flowers and of six pounds of proof spirit to be distilled, and the product to be re-distilled with half a pound of fresh flowers.—HERBIP.—one pound and a half of flowers to be distilled with ten pounds of alcohol, and the product to be re-distilled with three ounces of flowers.

HUNGARY WATER; COMPOUND SPIRIT OF ROSEMARY.

Aqua Hungarica seu Regine Hungariæ seu Anthos composita, Alcohol *Anthos compositus*; *Eau de la reine de Hongrie*, *Eau de Hongrie*, *Esprit aromatique*. (BRUNS.; DAN.; FERR.; HANN.; WIRTEM.; SWEDIAUR; VAN MONS)

R. Flowers of Rosemary, one pound.
Lavender, four ounces.
Proof Spirit, six pounds.

Digest for twelve days, and distil three pounds. (DAN.)

R. Flowers of Rosemary, three parts.
Lavender, one part.
Proof Spirit, six parts.
Water, twelve parts.

Distil three fourths. (SWEDIAUR)

VAN MONS prescribes one part of lavender, three parts of rosemary, twelve of proof spirit, and forty of water; eighteen to be distilled.

R. Flowering Herb of Rosemary, } of each,
Wild } one
Thyme, } part.
Proof Spirit, twenty-four parts.
Water, forty-eight parts.

Distil twenty parts. (VAN MONS)

R. Flowering Herb of Rosemary, four pounds.
Sage, six ounces.
Ginger, two ounces.
Proof Spirit, twelve pounds.
Water, two pounds.

Distil eleven pounds. (BRUNS.; HANN.; WIRTEM.)

VAN MONS prescribes one part of ginger, three parts of sage, twenty-four of rosemary, ninety of proof spirit, and one hundred and eighty of water; sixty-six to be distilled.

R. Herb of Rosemary, two pounds.
Sage, three ounces.
Thyme, six ounces.
Alcohol (34 degrees), sixteen pounds.

Macerate for two days, distil in a water-bath, and rectify the product by re-distillation. (FERR.)

CARMINATIVE SPIRIT.

Spiritus carminativus; *Alcohol Helenii compositum*. (HISP.)

R. Root of Elecampane, } of each,
Carline Thistle, } one ounce.
Tops of Rosemary, } of each,
Marjoram, } four ounces.
Bay Berries }
Fennel, } of each, one ounce.
Caraway, }
Anise, }
Pimenta, } of each,
Cinnamon, } half an ounce.
Fresh Orange Peel, three drams.
Proof Spirit, three pounds.

Digest for two days in a covered vessel, and distil in a water-bath.

APOPLECTIC LIQUOR.

Aqua apoplectica. (PALAT.; WIRTEM.; SPIELMANN)

R. Flowering Herb of Sage, eight ounces.

Flowering Herb of Marjoram,	} of each, one ounce.
Wild	
Thyme,	
Flowers of Lily of the Valley,	
three ounces,	
Lavender,	} of each, two ounces.
Rosemary,	
Cinnamon,	one ounce and a half.
Cubebs,	half an ounce.
Cloves,	three drams.
Mace,	two drams.
White Wine,	twelve pounds.
Macerate for three days, and distil one half. (WIRTEM.)	
R. Root of Sweet Flag,	} of each, one ounce
Florentine Iris,	
Round Birthwort,	and a half.
Herb of Marjoram,	
Ground Pine,	} of each, one handful.
Rosemary,	
Flowers of Primrose,	
Seeds of Mountain	
Lasserwort,	} of each, one dram and a half.
Fennel,	
Cubebs,	} of each, three drams.
Nutmegs,	
Cloves,	
Cinnamon,	} of each, two drams.
Aloe Wood,	
Long Pepper,	
Sagapenum,	
Opoponax,	} of each, one dram and a half.
Myrrh,	
Sandarac,	
Cowslip Water,	eighteen ounces.

Lime-flower Water, } of each,
Sage Water, } eighteen ounces.
Infuse for eight days; and distil. (SPIELMANN)

R. Fresh Flowers of Lily of the Valley,	
Lavender,	} of each, one ounce.
Cowslip,	
Lime,	
Betony,	
Clove-pink,	
Herb of Rosemary,	
Marjoram,	} of each, one ounce
Sweet Basil,	
Sage,	
Wild Thyme,	} and a half.
Marum,	
Cinnamon,	two ounces.
Cloves,	six drams.
Nutmeg,	one ounce.
Lesser Cardamoms,	
Cubebs,	} of each, half an ounce.
Zedoary,	
Yellow Sanders,	
Alcohol,	} of each, five pounds.
Wine,	

Macerate the herbs and flowers in the alcohol, and aromatics in the wine; then mix the liquors and distil slowly: preserve the first product, which is clear and limpid, for internal use, and the succeeding, which is white and oily, for external use. (PALAT; WIRTEM.)

Dose, from two drams to one ounce, in flatulent colic. Externally, it is used in embrocations and suffumigations.

RUBIA TINCTORIA, LINN.—MADDER.

SYNONIMES : — Garance, FR.; Krapp, Krappwurz, Færberœthe, GERM.; Marena, BOHEM.; Krap, DAN.; and DUT.; Granza rubia, SPAN.; Robbia, ITAL.; Marzana, POL.; Ruida, PORT.; Krapp, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; COKE.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant cultivated in Europe. (*Tetrandria Monogynia*, LINN.; *Rubiaceæ*, JUSS.; *Fig. Flore Medic. IV. 177*)

The root (*Radix Rubiæ tinctoriæ seu sativæ seu majoris*) is employed. It is long, VOL. II.

branching, articulated, cylindrical, of the diameter of a goose-quill, and of a red colour. Its colour is more intense externally than internally; its odour is feeble, and its taste is bitterish and slightly styptic.

It contains much tannin, gallic acid, starch, oxalic acid, and a peculiar colouring principle which COLLIN and ROBQUET call *Alizarine*; *Izarinum*. DEBEREINER had previously ascertained the existence of another principle, which he designated *Erythrodanin*.

In the East Indies the *Rubia Manjista*, ROXB. (Fuh, ARAB.; Well madatta, CYNG.; Menjithe, HIND.; Puntvayr, MAL.; Runas, PERS.; Manjischtha, SANS.; Manjittie, TAM.; Mandestie, TEL), an indigenous plant of Nepaul and Thibet, is substituted for this species.

It is astringent, and is remarkable for the property of reddening the bones of the ani-
3 Y

imals who feed on it. It is recommended in rachitis.—Dose, from half a dram to one dram of the powder.

RESOLVENT SPECIES.

Species resolventes. (HUFELAND)

- | | |
|---------------------------------------|------------------------------|
| R. Root of Madder, | } of each,
an equal part. |
| —— Dog-grass, | |
| —— Dandelion, | |
| —— Soapwort, | |
| Herb of Fumitory,
Tops of Milfoil, | |
- Mix.

AQUEOUS EXTRACT OF MADDER.

Extractum Rubiæ tinctoriæ aquosum.
(FERR.)

- R. Root of Madder, one pound.
Water, three pounds.

Boil, express, clarify the liquor with white of egg, and evaporate it to the proper consistence.

ALCOHOLIC EXTRACT OF MADDER.

Extractum Rubiæ tinctoriæ alcoholicum.
(BRUNS.; HANN.)

- R. Root of Madder, two parts.
Alcohol, three parts.
Water, nine parts.

Digest in a covered vessel, and express; distil off the alcohol slowly, and evaporate the rest to the proper consistence. (HANN.)

BRUNS. directs the wood to be exhausted by alcohol, the mixed tinctures to be evaporated, the spirit distilled off, and the remainder evaporated to the proper consistence.

ANTI-RACHITIC POWDER. (SAINTE-MARIE)

- R. Madder, two drams.
Prepared Oyster Shells, four drams.

* Make eighteen* parcels.—Dose, two or *three daily, for a young child.

COMPOUND POWDER OF MADDER. (HERBIP.)

- R. Madder, half an ounce.
Myrrh, of each,
Sub-borate of Soda, } one dram.
Mix.

DECOCTION OF MADDER.

Decoctum Rubiæ tinctoriæ. (SWEDIAUR)

- R. Root of Madder, one ounce.
Mace, one dram.

Boil down to two thirds. Add when strained

- Aromatic Tincture, two drams.
Syrup of Citric Acid, two ounces.

Mix.—It is astringent, and is recommended in jaundice, chlorosis, amenorrhœa, and chronic catarrhal affections.—Dose, three ounces, three or four times a day.

ANTI-SCROFULOUS POTION. (CADET DE GASSICOURT; SAINTE-MARIE)

- R. Root of Madder, half an ounce.
Hips, one pinch.
Walnut Leaves, three.
Water, one pound and a half.

Boil down to one pound, and add to the strained liquor when cold

- Tincture of Tartarized Iron, one dram.

Mix.—Dose, two glassfuls, morning and evening.

TONIC AND ANTI-HERPETIC POTION. (PIERQUIN)

- R. Root of Madder, one ounce.
Flowers of Arnica, of each,
Stems of Bitter-sweet, } half an ounce.
Water, six glassfuls.

Boil, and add

- Leaves of Scurvy-grass, half an ounce.

Infuse, and strain.

ALTERATIVE BROTH. (BORIES; CADET DE GASSICOURT; PIERQUIN; SAINTE-MARIE)

- R. Neck of Mutton, four ounces.
Root of Soap-wort, half a dram.
—— Madder, two drams.
Leaves of Cichory, one handful.
Water, two pounds.

Boil down to two thirds.—To be taken at four doses, in the morning, fasting; in engorgements of the abdominal viscera. A scruple of acetate of potass is added to the first dose.

ANTI-ICTERIC POTION. (NIEMANN)

- R. Root of Madder, } of each,
—— Turmeric, one ounce.
—— Great Celan-
dine, of each,
Tops of Lesser Cen-
taury, one handful.
Spring Water, } of each,
Rhenish Wine, } two pounds.

Boil together, and add to the strained liquor

- Compound Syrup of Smallage, two ounces.

Mix.—It is extolled in jaundice, by SYDENHAM.

Dose, half a pound, morning and evening.

ANTI-RACHITIC DRINK. (NIEMANN;
SAINTE-MARIE)

R. Dried Root of Madder, half an ounce.
Tartrate of Potass, two drams.
Spring Water, sixty-four ounces.

Boil for an hour over a slow fire; then strain, and add

Despumated Honey, two ounces.

Mix.—It is extolled in rachitis. LEVRET gave eight ounces of it to children, daily. When the fresh root is employed twice the quantity should be used.

ALKALINE DECOCTION OF MADDER.

Decoctum Rubiæ compositum. (HERBIP.)

R. Root of Madder, } of each,
Rhubarb, } three drams.
Subcarbonate of Potass, two drams.
Spring Water, three pounds and a half.

Boil for one hour; then add to the strained liquor

Syrup of Ginger, two ounces and a half.

Mix.

TONIC APOZEM. (BORIES)

R. Root of Madder, } of each
Rest-harrow, } half an ounce.
Leaves of Horehound, }
Motherwort, } of each,
Mugwort, } half a
Rosemary, } handful.
Seeds of Parsley, one pinch.
Saffron, } of each, one scruple.
Cinnamon, }
Water, a sufficient quantity
to obtain eight ounces of strained liquor.
Add to this

Syrup of Horehound, one ounce.
Tartarized Iron, half a dram.

Mix.—It is extolled in dysmenorrhœa, chlorosis, leucorrhœa, and hæmorrhages occasioned by the suppression of some habitual discharge.

ASTRINGENT WINE.

Vinum adstringens. (AUGUSTIN)

R. Root of Madder, one ounce.
Tormentil, } of each,
Unripe Oranges, } half an ounce.
Cascarilla, six drams.
Extract of Logwood, three drams.
Wine, two pounds.

Digest, and strain.—Dose, from half an ounce to one ounce, every three hours.

RUBUS.

Six species of this genus of plants are used in medicine:

1. *Rubus Chamemorus*, LINN.—Mountain Bramble.

SYNONIMES:—Ronce de montagne, FR.; Multbeere, GERM.; Multebær, Myrebær, Laamanak, Latoch, DAN.; Aalbes blaadige bramboos, DUT.; Moruszká, POL.

DAN.; POL.; ROSS.; BERGIUS; MURRAY and GMELIN.

A plant of the north of Europe. (*Icosandria Polygynia*, LINN.; *Rosaceæ*, JUSS.; *Fig.* LINN. *Fl. Lap.* t. 5. t. 1)

The fruit (*Fructus* seu *Baccæ Chamemori*) is employed. It is formed of berries, inodorous, of an acidulous taste before maturity, and watery and not very agreeable afterwards.

2. *Rubus fruticosus*, LINN.—Common Bramble.

SYNONIMES:—Ronce noire, FR.; Strauchartige Himbeere, Gemeiner Brombeer-

strauch, GERM.; Brambær, DAN.; Zarca, SPAN.; Bramen, Brummeln, DUT.; Rovo, ITAL.; Jezyny, POL.; Sylva, PORT.; Brombær, SWED.

ANT.; BRUNS.; HISP.; GALL.; WIRTEM.; BRUGNATELLI; SPIELMANN.

A common shrub of Europe. (*Fig.* Lmk. *Enc. meth.* t. 441)

The fruit (*Baccæ Rubi nigri* seu *fruticosi* seu *Mori rubri*) is employed. It consists of inodorous berries of an agreeable taste.

3. *Rubus arcticus*.—Dwarf Bramble.

SYNONIMES:—Ronce septentrionale, FR.; Nordische Himbeere, GERM.; Akerbær, SWED.

BRUNS.; ROSS.; SUEC.; BERGIUS; MURRAY and GMELIN.

A shrub of the north of Europe. (*Fig.* LINN. *Fl. Lap.* t. 5. f. 2)

The fruit (*Baccæ Norlandicæ*) is employed. It consists of berries of an agreeable odour, and of a pleasant and vinous taste.

4. *Rubus trivialis*, MICH.—Dewberry.

SYNONIME :—Ronce commune, FR.

AMER. ; COXE.

A shrub of North America.—Its bark is astringent.

5. *Rubus villosus*, WILLD.—Blackberry.

SYNONIME :—Ronce velue, FR.

AMER. ; COXE.

A shrub of North America.—Its bark is astringent.

6. *Rubus Idæus*, LINN.—Raspberry.

SYNONIMES :—Framboisier, FR. ; Himbeere, GERM. ; Maliny, Ostruzini, BOHEM. ; Hindbær, Bryngebær, Himbær, DAN. ; Frambueso, Sanguesca, SPAN. ; Framboizen, DUT. ; Rovo ideo, ITAL. ; Maliny, POL. ; Amoreira frambeeza, PORT. ; Hallon, SWED.

AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIP. ; OLDEN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; WURCEB. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A common shrub of Europe. (*Fig. Flore Medic. III. 170*)

The fruit *Raspberries* (*Fructus seu Baccæ Rubi idæi*), is employed. It consists of berries, nearly round, red, grey, or white, composed of an assemblage of seeds enveloped each in a soft pulp, and supported upon a rather long conical receptacle.

DECOCTION OF BLACKBERRY.

Decoctio Rubi villosi. (AMER. ; NIEMANN)

R. Bark of the Root of *Rubus Villosus*,
one ounce.

Water, a sufficient quantity
to obtain eight ounces of strained decoction.

It is astringent, and is used in North America.—Dose, half a glassful several times a day.

SYRUP OF DWARF BRAMBLE.

Syrupus Rubi arctici. (SUEC.)

R. Juice, recently expressed and depurated, of the Berries of *Rubus arcticus*, one pound.
White Sugar, two pounds.

Make a syrup at a moderate heat.

JUICE OF RASPBERRIES.

Succus Rubi idæi. (BRUNS. ; HISP. ; FULD. ; HASS. ; PALAT. ; SARD. ; SAX. ; WIRTEM. ; PIDERIT)

R. Ripe Raspberries, any quantity.

Bruise them with the hands, set the mass in a cellar till a clear liquor appears at the surface, and strain with expression ; set apart the juice for several days, that it may become clarified, and preserve the clear part in long-necked bottles under a stratum of oil. (BRUNS. ; HISP. ; PALAT. ; SAX. ; WIRTEM.)

HASS. ; and PIDERIT direct, before the juice be put into bottles, that one fourth of sugar be added, and the mixture boiled for an instant.

ROB OF RASPBERRIES.

Rob Rubi idæi. (BRUNS. ; HISP. ; FULD. ; WIRTEM. ; HERBIP. ; PIDERIT ; SWEDIAUR)

R. Ripe Raspberries, any quantity.

Having bruised them, set aside the mass for three days ; then express the juice, and boil it over a slow fire, to the consistence of honey, adding one fourth of sugar. (BRUNS. ; FULD. ; WIRTEM. ; HERBIP. ; PIDERIT ; SWEDIAUR)

R. Sugar, boiled à la grande plume,
two pounds.
Juice of Raspberries, } of each,
Water, } one pound.

Boil to the consistence of jelly, and strain. (HISP.)

It is acidulous, and refreshing.

HONEY OF RASPBERRIES. (VAN MONS)

R. Recently expressed Juice of Raspberries, one part.
Honey, three parts.

Mix without heat, and decant after sufficient repose.

SYRUP OF RASPBERRIES.

Syrupus Rubi idæi seu de Frambæsiis.
(AMST. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; DAN. ; GALL. ; FENN. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; PALAT. ; POL. ; BORUS. ; SUEC. ; WIRTEM. ; HERBIP. ; COXE ; PIDERIT ; SWEDIAUR ; VAN MONS)

R. Raspberries, not perfectly ripe,
Sugar, } of each,
an equal part.

Mix with slight agitation, set over the fire, and boil a little. Strain the syrup through a close silk sieve. (GALL.)

R. Juice of Raspberries, recently expressed, two parts.
White Sugar, three parts and a half.
Dissolve in a warm water-bath. (VAN MONS)

R. Depurated Juice of Raspberries, twenty ounces.
White Sugar, three pounds.
Dissolve at a slow heat. (HANN.; OLDEN.; LUSIT.; POL.; BORUS.)

AMST.; BATAV.; and BELG. prescribe sixteen ounces of juice, and two pounds and a half of sugar.—DAN.; PALAT.; WIRTEM.; and HERBIP.—twenty ounces of juice, and thirty-two of sugar.—HASS.; and PIDERIT—ten ounces of juice, and thirteen and a half of sugar.—AUSTR.; FENN.; and SUEC.—one part of juice, and two parts of sugar.—SWEDIAUR—sixteen ounces of juice, and twenty-eight of sugar.—GENEV.—one pound and a half of juice, and two pounds and a half of sugar.—COXE—one pint of juice, and two pounds of sugar.—LIPP.—eighteen ounces of juice, and two pounds of sugar.—FULD.—an equal part of each.

RASPBERRY VINEGAR.

Acetum Rubi idæi. (GALL.; HANN.; OLDEN.; PALAT.; POL.; BORUS.; WIRTEM.; SARD.)

R. Fresh Raspberries, six pounds.
Vinegar, four pounds.

Macerate for four days, strain without expression, and filter. (GALL.)

HANN.; OLDEN.; POL.; BORUS.; and SARD. prescribe one part of raspberries, and two parts of vinegar; to be macerated for one month.—PALAT.; and WIRTEM.—one part of raspberries, and six parts of vinegar; to be macerated in the sun for several days.

R. Juice of Ripe Raspberries, two pounds.
Vinegar, half a pound.

Macerate for two days; then express strongly, and add to the liquor,

White Sugar, one pound and a half.

Digest over a slow fire, and decant when cold. (DAN.)

It is refreshing and cooling, and is added, in variable quantity, to drinks.

WATER OF RASPBERRIES.

Aqua Rubi idæi. (BAVAR.; BRUNS.; DAN.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; ROSS.; SAX.; PIDERIT; SWEDIAUR; VAN MONS)

R. Raspberries, bruised, one part.
Water, two parts.

Distil one part. (BRUNS.)

PALAT. prescribes one part of raspberries, and four parts of water; one part to be distilled.—DAN.—one of raspberries, and six of water; four to be distilled.—VAN MONS—one of raspberries, and twenty-four of water; six to be distilled.—ROSS.—four of raspberries, and ten of water; six to be distilled.—SWEDIAUR—ten of raspberries, and a sufficient quantity of water; twenty to be distilled.—LIPP.—three of raspberries, and sixteen of water; eight to be distilled.

R. Expressed Raspberries, one part.
Water, three parts.

Distil two parts. (HANN.)

BAVAR. prescribes one part of expressed fruit, and six parts of water; three parts to be distilled.—OLDEN.—two of expressed fruit, and a sufficient quantity of water; twenty to be distilled.—HASS.; and PIDERIT—three of expressed fruit, and sixteen of water; eight to be distilled.

R. Expressed Raspberries, one part.
Water, six parts.

Ferment for several days, and distil two parts. (SAX.)

R. Raspberries, any quantity.

Bruise, and beat them with an equal quantity of warm water; leave the mass to ferment, add twenty-four parts of cold water, and distil six parts. (VAN MONS)

R. Expressed Raspberries, ten pounds.
Water, thirty pounds.
Subcarbonate of Potass, two ounces.

Distil twenty pounds. (POL.)

FULD. prescribes eight pounds of raspberries twenty-four of water, and one pound of subcarbonate of potass; sixteen pounds to be distilled.

RUMEX.

Six species of this genus are mentioned in pharmacopœias:

1. *Rumex acetosa*, LINN.—Common Sorrel.

SYNONIMES:—Oseille des jardins, Surelle,

FR.; Sauerampfer, GERM.; Ssiowjk, BOHEM.; Suramper, DAN.; Azedera, SPAN.; Veldzuuring, DUT.; Acetosa, ITAL.; Sszsaw, POL.; Azedas, PORT.; Engsyra, SWED.

LOND.; EDINB.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HASS.; LUSIT.; BORUS.; ROSS.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

A plant cultivated in every part of Europe. (*Hexandria Trigynia*, LINN.; *Polygonæa*, JUSS.; *Fig. Zorn, Ic. pl. t. 70*)

The root, leaves, and seeds are employed.

The root (*Radix Acetosæ officinalis* seu *nostratis* seu *Oxalidis* seu *Acetosæ pratensis*) is fibrous and yellowish; several long slender fibres go off from a small head: under a reddish-yellow bark it presents a white substance. Its taste is slightly bitter and astringent.

The leaves are oblong, sagittate, obtuse, entire, smooth, inodorous, but of an acid taste from the presence of super-oxalate of potass and of lime.

The seeds are triangular, and of an acerb and bitterish taste.

2. *Rumex scutata*, LINN.—Small Sorrel.

SYNONIMES:—Oseille ronde, Petite Oseille, FR.

GALL.; HASS.; BERGIUS.; SPIELMANN.

A plant of Europe. (*Fig. Jacq. Ic. rar. I. t. 67*)

The herb (*Herba Acetosæ rotundifolia*) is employed. It consists of cordiform or oval, obtuse or pointed leaves having long foot-stalks, and furnished on each side of their base with divergent auricles (*oreillettes*). It has an acid and agreeable taste.

CONSERVE OF SORREL. (FERR.; GENEV.; ROSS.; SARD.)

R. Fresh Leaves of Sorrel, one part.
White Sugar, two parts.

Beat together till incorporated. (ROSS.)

SARD. prescribes an equal part of each.—FERR.; and GENEV.—one part of sorrel, and three parts of sugar.

JUICE OF SORREL.

Succus Acetosæ. (HASS.; SARD.; HERBIP.)

R. Fresh Leaves of Sorrel, any quantity.

Bruise them in a wooden mortar with a wooden pestle, express the juice, set it aside, and preserve the limpid part in bottles.

HERBIP. directs one dram of alcohol to be added to each pound of juice.

SYRUP OF SORREL.

Syrupus Acetosæ. (SARD.; WIRTEM.)

R. Depurated Juice of Sorrel, ten ounces.
White Sugar, sixteen ounces.

Boil together to the consistence of syrup. (WIRTEM.)

SARD. directs a sufficient quantity of the juice to be added to simple syrup, and then evaporated in a water-bath.

WATER OF SORREL. (SARD.)

R. Leaves of Sorrel, bruised, one part.
Water, two parts.

Distil two thirds.

SUPPURATIVE CATAPLASM. (GALL.; SAINTE-MARIE)

R. Leaves of Sorrel, bruised, one handful.
Lard, one ounce.
Linseed, bruised, four ounces.

Boil together, in beer, to the proper consistence. (SAINTE-MARIE)

R. Resolvent Meals, four ounces.
Emollient Decoction, a sufficient quantity.

Boil to the proper consistence, and add

Pulp of Lily, } of each,
Boiled Sorrel Leaves, } two ounces.
Resin Ointment, one ounce.

Mix. (GALL.)

3. *Rumex aquatica*, LINN.—Water Dock.

SYNONIMES:—Patience d'eau, Rhubarbe sauvage, FR.; Wasserrampfer, Wassermangold, Wassergrindwurz, Rostampferwurz, Butterweckenkraut, GERM.; Vandskreppe, DAN.; Waterpatich, DUT.; Labaca major, PORT.; Vatnsyra, SWED.

DUBL.; AMER.; AMST.; BATAV.; BELG.; BRUNS.; GALL.; FULD.; LIPP.; ROSS.; SAX.; WIRTEM.; BERGIUS.; COXE.; MURRAY and GMELIN.; SPIELMANN.

A common plant of Europe, found in marshy districts. (*Fig. Zorn, Ic. pl. t. 588*)

The root (*Radix Lapathi aquatici* seu *Herbæ Brittanica* seu *Hydrolapathi* seu *Rhapontici*) is employed. It is large, branching, annulated, of a blackish-brown colour without, and saffron coloured within. It is inodorous, but has an austere and bitterish taste.

It is tonic and diaphoretic, and is much employed in diseases of the skin.

4. *Rumex crispus*, LINN.—Curled Dock.

SYNONIMES:—Patience frisée, FR.; Krauser Ampfer, GERM.; Kruset skreppe,

DAN. ; Gekruide patich, DUT. ; Svinisyra, SWED.

GALL. ; COXE.

A plant of Europe. (*Fig. Curt. Fl. Lond. t. 204*)

The herb (*Herba Lapathi crispi*) is employed. It consists of a cylindrical, slightly branching stem, furnished with lanceolated leaves, very much waved, or rather curled on the margin. It has a very slightly styptic taste.

5. *Rumex sanguinea*, LINN.—Bloody Dock, Red Dock.

SYNONIMES :—Patience rouge, FR. ; Blut-ampfer, Blutkraut, GERM. ; Bluedkruid, DUT.

GALL.

A plant of North America and of the northern parts of Europe. (*Fig. Zorn, Ic. pl. t. 127*)

The root and herb are employed.

The root (*Radix Lapathi sanguinei*) has an acerb and astringent taste.

The herb consists of a branching purple coloured stem, and of cordiform, lanceolated, finely-serrated purple leaves.

EXTRACT OF DOCK.

Extractum Patientiæ. (GALL. ; VAN MONS)

R. Root of Dock, sliced, any quantity.

Cover it with water, and leave it to macerate for twelve hours ; then double the quantity of water, and continue the maceration for the same length of time. Strain and clarify the two liquors, and evaporate to the consistence of an extract. (VAN MONS)

R. Root of Dock, one pound.
Cold Water, four pounds.

Macerate for twenty-four hours, occasionally stirring, and strain ; macerate the residue with fresh water in the same manner, and strain again. Mix the liquors, and evaporate in a water-bath to the consistence of an extract. (GALL.)

Dose, from one to three scruples.

DECOCTION OF DOCK.

Decoctum Lapathi. (FORM. HÔP. FR. ; NIEMANN ; RATIER ; SPIELMANN)

R. Root of Dock, one ounce.
Water, two pounds.

Boil down to one half. (NIEMANN ; RATIER)

R. Root of Dock, six drams.
Water, a sufficient quantity

to obtain two pounds of strained decoction. Add towards the end

Root of Liquorice, two drams.

Infuse for a few minutes, and strain. (FORM. HÔP. FR. ; NIEMANN)

R. Root of Dock, two ounces.
—— Elecampane, } of each,
Juniper, } half an ounce.
Water, six pounds.

Boil to the consumption of half a pound, and add towards the end

Herb of Fumitory, one handful.

Strain. (SPIELMANN)

It is slightly excitant, and is prescribed as an alterative in cutaneous disorders.—Dose, several glassfuls in the course of the day.

6. *Rumex alpina*, LINN.—Bastard or Monk's Rhubarb.

SYNONIMES :—Rhubarbe des moines, Fausse rhubarbe, FR. ; Mönchsrhabarber, Alpenampfer, GERM. ; Skreppe, DAN. ; Romaza, SPAN. ; Spitzbladdige pattich, DUT. ; Romice, ITAL. ; Szczaw kobyli, POL. ; Babaca menor, PORT. ; Hæstysræ, SWED.

BELG. ; BRUNS. ; GALL. ; MURRAY and GMELIN ; SPIELMANN.

A plant of the Alps. (*Fig. Zorn, Ic. pl. t. 261*)

The root (*Radix Hippolapathi seu Rhubarbari monachorum*) is employed. It is thick, ligneous, fibrous, red without, and yellow within. It has a disagreeable odour, and a bitter, acerb, and nauseous taste.

RUSCUS.

Two species of this genus of plants are used in medicine :

1. *Ruscus aculeatus*, LINN.—Butchers-broom, Knee Holly.

SYNONIMES :—Fragon piquant, Petit houx,

Houx frelon, FR. ; Slachlicher Mäusedorn, Mäusedornbüsch, Myrtendorn, GERM. ; Lesnj myrtus, Gedlice wlaska, BOHEM.

ANT. ; BRUNS. ; HISP. ; GALL. ; FERR. ;

FORM. HÔP. FR.; GENEV.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the forests of Europe. (*Diæcia Monadelphica*, LINN.; *Asparagineæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 448*)

The root (*Radix Rusci* seu *Brusci* seu *Myrtacanthæ*) is employed. It is long, ligneous, white, and consists of numerous fibres of the thickness of quills. Its taste is at first sweetish, and then becomes bitter.

It is diuretic.—It forms one of the five major aperitive roots.

2. *Ruscus Hypoglossum*, LINN.

SYNONIMES:—Laurier Alexandrin, FR.; Alexandrinischer Loorbeer, GERM.; Cyjpkowa bylina, BOHEM.; Alexandrinse laurier, DUT.; Loreiro de Alexandria, PORT.

BRUNS.; GALL.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOUT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the south of Europe. (*Fig. Zorn, Ic. pl. t. 481*)

The leaves (*Herba Umlaria* seu *Hypoglossi* seu *Bislinguæ* seu *Bonifaciæ* seu *Lauri Alexandrini angustifolia*) are employed. They are oval and lanceolate, and bear towards the middle of their upper surface, a ligula in the angle of which the flowers rise.

They are destitute of odour but have a very astringent taste.

It is slightly astringent, but is not much used.

DIURETIC PTISAN. (PIERQUIN)

R. Root of Butchers-broom, six ounces.
Water, one pound.

Boil and strain.

R. Root of Butchers-broom, } of each,
—— Strawberry, } five drams.
—— Asparagus, }
Water, two pounds and a half.

Boil down to two pounds, then strain, and add

Syrup of Hyssop, } of each,
Compound Syrup of } one ounce.
Smallage, }

Mix.

RUTA GRAVEOLENS, LINN.—RUE.

SYNONIMES:—Rue des jardins, FR.; Raute, Gartenraute, GERM.; Sendib, ARAB.; Ruta, BOHEM.; ITAL.; POL.; and RUSS.; Aruda, CYNG.; Ruda, DAN.; Saturei, HIND.; Ruite, DUT.; Matskase-so, JAP.; Inghu, JAV.; Suddab, MAH.; Sadsa, MAL.; Arruda, PORT.; Brahmi, Somalata, SANS.; Winruta, SWED.; Aruda, TAM.; Saddapu aku, TEL.

LOND.; EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A small shrub of the south of Europe. (*Decandria, Monogynia*, LINN.; *Rutaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 163*)

The herb and seeds are employed.

The herb (*Herba Rutæ* seu *Rutæ hortensis*) consists of a cylindrical and branching stem, and of alternate, petiolated leaves of a glaucous green colour, almost doubly

winged, and composed of thickish, oval, obtuse leaflets, narrowed at the base. It has a strong, peculiar, and disagreeable odour, and a hot, acrid, and very bitter taste.

The seeds are small, blackish, reniform, wrinkled, somewhat angular, and of an aromatic taste.

It is excitant, stomachic, nervine, diaphoretic, carminative, anti-putrescent, anodyne, and anthelmintic.—Dose, from ten grains to two scruples, of the powder.

CONSERVE OF RUE.

Conserva Rutæ. (SARD.)

R. Leaves of Rue, } of each, one part.
White Sugar, }
Beat together.

CONFECTION OF RUE.

Confectio Rutæ. (LOND.; DUBL.; CADET DE GASSICOURT; VAN MONS)

R. Dried Leaves of } of each,
Rue, } one ounce and
Seeds of Cara- } a half.
way,

Bay Berries, one ounce and a half.
Sagapenum, half an ounce.
Black Pepper, two drams.
Despumated Honey,
sixteen ounces.

Pulverize and mix.

It is stimulant and emmenagogue.

RUBEFACIENT AND FEBRIFUGE PASTE.
(BORIES)

R. Fresh Tops of Rue, two ounces,
Powdered Mustard, two drams.

Beat together.—This paste is applied up-
on the wrist several hours before the pa-
roxysm.

AQUEOUS EXTRACT OF RUE.

Extractum Rutæ aquosum. (DUBL.; EDINB.;
BRUNS.; GENEV.; PALAT.; WIRTEM.)

Extractum Rutæ graveolentis.
(EDINB.)

Ext. Foliorum Rutæ. (DUBL.)

R. Leaves of Rue, one pound.
Pure Water, eight pounds.

Boil down to one half, and express.
Evaporate the liquor slowly to the
consistence of pilular mass.

GENEV. gives the same formula.

R. Herb of Rue, one pound.
Spring Water, six pounds.

Digest in a warm place for four days,
then boil a little, and express. Evaporate
the liquor to the proper consistence.
(BRUNS.; PALAT.; WIRTEM.)

Dose, from ten grains to half a dram.

ALCOHOLIC EXTRACT OF RUE.

Extractum Rutæ alcoholicum. (HERBIP.)

R. Herb of Rue, } of each, one pound.
Alcohol, }
Water, eight pounds.

Digest for three days in a moderate heat,
and express; then set aside, and decant the
liquor: distil off the alcohol, and evaporate
the residue to the proper consistence.

ANTI-LYSSIC POWDER.

*Pulvis anti-lyssus seu ad morsum canis ra-
bidi.* (PALAT.; WIRTEM.; SPIELMANN)

R. Leaves of Rue, }
Vervain, } of each,
Sage, } an
Plantain, } equal part.
Polypody,
Wormwood,
Mint,
Bastard Balm, }

VOL. II.

Leaves of Betony, } of each,
Tops of St. John's-Wort, } an
Lesser Centaury, } equal part.

Mix.—It is extolled by PAULMIER.—Dose,
from half a dram to two drams.

EMMENAGOGUE POTION. (PIERQUIN)

R. Juice of Rue, one ounce.
Oxymel of Squill, half an ounce.

Mix.—To be taken by spoonfuls, in hys-
terical affections.

DETERGENT INJECTION.

Injectio auricularis deterensiva. (SPIELMANN)

R. Herb of Rue, one handful.
Spring Water, six ounces.

Infuse for twelve hours, and strain. Add

Essence of Myrrh, half a dram.

Honey of Roses, six drams.

Mix.

VINEGAR OF RUE.

Acetum Rutæ seu rutaceum. (AMST.;
AUSTR.; BRUNS.; HASS.; OLDEN.; LU-
SIT.; PALAT.; BORUS.; SARD.; WIR-
TEM.; HERBIP.; PIDERIT; VAN MONS)

R. Fresh Leaves of Rue, one part.
Vinegar, ten parts.

Macerate without heat for several days,
express lightly, and filter. (VAN MONS)

SARD. prescribes one part of rue, four
parts of vinegar, and six days' digestion in
a moderate heat.—BRUNS.; HASS.; PA-
LAT.; WIRTEM.; and PIDERIT.—one of
rue, six of vinegar, and a few days' diges-
tion.—AMST.; and BORUS.—one of rue,
nine of vinegar, and three weeks' infusion
in the sun.—AUSTR.—one of rue, twelve of
vinegar, and three days' maceration.

R. Fresh Leaves of Rue, one pound.
Vinegar, six pounds.

Macerate for six days, boil a little, and
strain. Add to the liquor,

Alcohol, half a pound.

Mix. (LUSIT.)

AUSTR. prescribes three ounces of rue,
two pounds of distilled vinegar, several
days' digestion in a moderate heat, and the
addition of one eighth of alcohol.

Here, as in every vinegar, simple or com-
pound, the alcohol is intended to prevent
decomposition.—WIRTEM. thinks the addi-
tion of one-twentieth is sufficient.

CAMPHORATED VINEGAR OF RUE.

Acetum Rutæ camphoratum. (VAN MONS)

R. Flowering Tops of Rue, two ounces.
Vinegar, sixteen ounces.

Infuse in a moderate heat for twenty-four
3 Z

hours; then strain, and dissolve in the liquor.

Camphor, two drams.

Filter.—It is employed as a prophylactic.

INFUSED OIL OF RUE.

Oleum Rutæ infusum. (BRUNS.; HISP.; GALL.; SARD.; WIRTEM.; HERBIP.)

R. Fresh Herb of Rue, } of each,
Olive Oil, } an equal part.

Digest for three days, and express. (BRUNS.; WIRTEM.)

R. Fresh Herb of Rue, one part and a half.
Olive Oil, three parts.

Digest for three days in a water-bath and express when cold. Repeat the maceration with fresh herb. (SARD.)

GALL. directs two parts of oil to be macerated with one part of herb for twenty-four hours in warm ashes, the operation to be repeated a second time, the mixture to be then slightly boiled, and expressed.

R. Fresh Herb of Rue, one part.
Olive Oil, two parts.

Digest for three days at a moderate heat; then boil to the consumption of the humidity, and express. (HERBIP.)

HISP. prescribes one part of herb and three parts of oil.

It is excitant, anti-colical, anthelmintic, and nervine. It is employed in lavements and frictions.

OINTMENT OF RUE.

Unguentum seu Butyrum Rutæ. (HISP.)

R. Fresh Leaves of Rue, } of each
Wormwood, } two
Mint, } ounces.
Lard, sixteen ounces.

Boil to the consumption of the humidity, and strain with expression.

ESSENTIAL OIL OF RUE.

Oleum Rutæ æthereum, Ætheroleum Rutæ. (DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR)

Oleum Rutæ. (DUBL.)

R. Fresh Herb of Rue, any quantity.
Water, a sufficient quantity,
to prevent empyreuma. Distil.

AMST.; BATAV.; BELG.; FERR.; ROSS.; BRUGNATELLI; and COXE require the same.

BAVAR.; FULD.; and LIPP. prescribe one part of herb and three parts of water.—SWEDIAUR—one of herb and four of water.—GALL.—five of herb and seven of water.—ANT.—three of herb and ten of water.—HANN.; POL.; BORUS.; and SAX.; one of herb and eight of water.—AUSTR.—one of herb and twelve of water.

R. Fresh Herb of Rue, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Distil at the end of three days. (BRUNS.; HASS.; PALAT.; SARD.; PIDERIT)

R. Fresh Herb of Rue, any quantity.
Distilled Water of Rue, a sufficient quantity.

Distil. (HERBIP.)

BALSAM OF RUE.

Balsamum Rutæ. (WIRTEM.; SPIELMANN)

R. Oil of Nutmeg, two drams and a half.
Essential Oil of Rue, one dram.

Mix

DISTILLED WATER OF RUE.

Aqua Rutæ. (AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; HANN.; PALAT.; POL.; BORUS.; SARD.; SAX.; BRUGNATELLI; SWEDIAUR; VAN MONS)

R. Fresh Herb of Rue, one part.
Water, a sufficient quantity.

At the expiration of twelve hours, distil ten parts. (AMST.; BATAV.; BELG.; HANN.; POL.; BORUS.; SWEDIAUR)

PALAT. prescribes one part of herb and four parts of water; two parts to be distilled.—HERBIP.; and BRUGNATELLI—one of herb and six of water; four to be distilled.—BAVAR.—one of herb and sixteen of water; eight to be distilled.—SAX.—one of herb and fifteen of water; ten to be distilled.—AUSTR.—one of herb and twelve of water; four to be distilled.

R. Fresh Herb of Rue, one part.
Water, four parts.

Distil one half, and re-distil the product with fresh herb. (SARD.)

R. Fresh Herb of Rue, ten parts.
Proof Spirit, one half part.
Water, one hundred and sixty parts.

Distil forty parts. (VAN MONS)

ANTI-HYSTERIC JULEP. (SAINTE-MARIE)

R. Distilled Water of Rue, } of each,
Bryony, } two
Orange } ounces.
Flowers,
Syrup of French Lavender, } of each,
Valerian, } one
Peony, } ounce.

Mix.—To be taken at three or four doses.

SACCHARUM.—SUGAR.

SYNONIMES:—*Sacchar*; Σάκχαρ, Σάκχαρον.
—Sucre, FR.; Zucker, GERM.; Sukhir, ARAB.; Cukr, BOHEM.; Sukker, DAN.; Schukker, DUT.; and PERS.; Azucar, SPAN.; Cheni, HIND.; Suiker, DUT.; Zucchero, ITAL.; Saker, MAH.; Gula, MAL.; Cukiea, POL.; Assucarr, PORT.; Sarkara, SANS.; Socker, SWED.; Sak-karr, TEL.; Panchadara, TAM.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

A solid, white substance, easily crystallizable, inodorous, and of a very agreeable taste.

Sugar is extracted from the juice of the stem of the *Arundo Saccharifera*, LINN., a plant originally of eastern Asia. (*Triandra Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Rumph. Amb. V. 74*)

SIMPLE SYRUP.

Syrupus simplex seu communis seu Sacchari, Saccharum clarificatum. (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; SUEC.; WIRTEM.; HERBIP.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Syrupus Simplex.

R. Refined Sugar,
two pounds and a half.
Water, one pint.

Dissolve the sugar in the water by the heat of a water-bath; set the syrup aside for twenty-four hours, then take off the scum, and pour off the clear part from the fæces if there are any. (LOND.; DUBL.)

R. Refined Sugar, fifteen parts.
Water, eight parts.

Dissolve the sugar in the water by

the aid of a gentle heat, and then boil a little so as to form a syrup. (EDINB.)

LOND. directs syrups to be kept in a place, the temperature of which does not exceed 55 degrees.

HASS.; HERBIP.; and PIDERIT prescribe four parts of sugar and three of water.—AMER.; AMST.; and COXE—fifteen of sugar and eight of water.—ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; FENN.; FERR.; FULD.; SUEC.; WIRTEM.; and SWEDIAUR—two of sugar and one of water.—LIPP.—three pounds of sugar and twenty-seven ounces of water.—GENEV.—a sufficient quantity of each to form a thick liquid.

R. White Sugar, three parts.
Water, one part.

Dissolve at a gentle heat, and pour gradually into the syrup, the white of an egg mixed with one part of water; boil till nothing more than a little white scum appears on the surface of the liquor, and then strain it through flannel. (FORM. HÔP. FR.; GALL.)

The syrup may also be made without boiling, dissolving two parts of sugar in one part of water, by the heat of a stove. (GALL.)—FORM. HÔP. FR. requires the cold syrup to mark 35 degrees.

BLONDEAU has proposed the following process to obtain the syrup colourless.

R. Sugar, sixty pounds.
Washed Animal Carbon, sixty ounces.
Water, twenty-four pounds.

Dissolve over a clear fire, and add, at twice or thrice, three pounds of water mixed with the whites of six eggs: remove it from the fire, set aside, skim, and strain through flannel. (HENRI)

DUROZIER makes it according to the following process.

R. Sugar, forty pounds.
Cold Water, twenty-two pounds.

Agitate the solution with washed animal carbon; stir the mixture occasionally for twenty-four hours, and filter.

The following is by GRAMMAIRE:

R. Sugar, twenty pounds.
Water, ten pounds.
Whites of Eggs, three.

Put them together into an autoclave, and leave it on the fire for a quarter of an hour.

SYRUP OF CORAL.

Syrupus seu Tinctura Corallii. (GENEV.)

R. Red Coral, }
 Yellow Wax, } of each, two pounds.

Having melted the wax throw the coral into it, and boil till the colour of this shall have completely passed into the wax. Then decant, place the mixture over the fire again, and add

White Sugar, two pounds.

When the sugar is melted, remove it from the fire, and add

Boiling Water, one pound and a half, very gradually and constantly stirring to prevent the liquid from flowing over. Boil again to liquify the wax, and set it apart in

a cool place to consolidate; finally remove the crust, and preserve the syrup which it covers.

MANUS CHRISTI TROCHISCS.

Rotule Manus Christi, Saccharum perlatum.
 (BRUNS.; WIRTEM.; SPIELMANN)

R. White Sugar, any quantity.
 Rose Water, a sufficient quantity.

Make trochiscs. (SPIELMANN)

R. White Sugar, one pound.
 Rose Water, a sufficient quantity

to dissolve the sugar. Boil *à la plume*, and add

Mother of Pearl, in powder,
 half an ounce.

Make trochiscs. (BRUNS.; WIRTEM.)

SALEP, SALAB.—SALEP.

SYNONIMES:—Salep, FR.; Salibimisri, DUK.; and HIND.; Salamisrie, TAM.

AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; DAN.; GALL.; FENN.; HAMB.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE; BERGIUS; COXE; GUIBOURT; MURRAY and GME-LIN; PIDERIT; SPIELMANN.

An amylaceous substance in small pieces varying in size from the volume of a coffee grain to that of an almond; it is straw-coloured, semi-transparent, of an almost horny consistence, of a slightly hircine and spermatic odour, and of a sweet and mucilaginous taste.

Salep is prepared, in Persia, from the tubercles of unascertained orchideæ.

It is analeptic.

SALEP POWDER. (NIEMANN)

R. Salep, one ounce and a half.
 Sago, one ounce.
 Gum Tragacanth, one dram.

Mix.—A tea spoonful of this powder is given, morning and evening, mixed with water or milk, as a nutrient and demulcent.

CONSERVE OF ORCHIS.

Conditum Satyrii. (PALAT.; SARD.; WIRTEM.)

R. Fresh Root of Orchis, one pound.
 Water, eight pounds.

Macerate for several days, and then boil till the root becomes soft. Clarify in the decoction

White Sugar, one pound and a half.

Infuse the root in the syrup for three days, then boil, and preserve it in it or dry it in a stove. (SARD.)

PALAT.; and WIRTEM. direct the root to be softened in water by boiling, then having let it drip, sugar boiled *à la plume* to be poured over it.

SALEP JELLY.

Gelatina seu Mucilago Radicis Saleb. (ANT.; HANN.; LUSIT.; CAST. BORUS.; SWEDIAUR; VAN MONS)

R. Salep, in powder, one dram.
 Water, eight ounces.

Boil. (SAINTE-MARIE)

R. Salep Powder, two drams.
 Boiling Water, six ounces.

Triturate together. (ANT.)

CAST. BORUS. prescribes two drams of salep and twelve ounces of boiling water.

R. Salep, in powder, one dram.
 Water, eight ounces.

Boil, and add

White Sugar, a sufficient quantity to sweeten it agreeably.

VAN MONS prescribes one dram of salep, four ounces and a half of water, and one ounce and a half of simple syrup.—HANN.—two drams of salep, eight ounces of

water, and one ounce and a half of syrup of orange peel.

R. Salep, in powder, two drams.
Cold Water, a sufficient quantity
to make nine ounces of decoction. Add to it

Syrup of Poppy, one ounce and a half.
Water of Cherry Laurel, six drams.

Mix. (SWEDIAUR)

DECOCTION OF SALEP.

Decoctum de Radice Salep, Infusum Orchidis.
(LIPP.; HERBIP.; AUGUSTIN; SAUNDERS)

R. Salep, in powder, one dram.
Pure Water, sixteen ounces.

Boil till only one pound of strained liquor remain. (HERBIP.)

AUGUSTIN prescribes two drams of salep and twelve ounces of water.

R. Powdered Orchis, one dram.
Impregnate it with water till it has become quadrupled in volume; then add

Boiling Water, one pound and a half.

Boil for half an hour. (LIPP.)

R. Salep, in powder, two drams.
Spring Water, a sufficient quantity
to produce two pounds of strained decoction. Add

Syrup of Poppy, } of each,
Diacodium, } one ounce.

Mix. (SAUNDERS)

DEMULCENT POTION.

Mixtura demulcens. (HAMB.; SAINTE-MARIE)

R. Salep, in powder, two scruples.
Hot Water, ten ounces.

Add to the solution,
White Sugar, half an ounce.
Dissolve. (HAMB.)

R. Salep, one dram.
Boiling Water, a sufficient quantity.

Boil for a quarter of an hour, and to four ounces of the strained liquor, add,

SYDENHAM's Laudanum, fifteen drops.
Syrup of Diacodium, four drams.

Mix. (SAINTE-MARIE)

PECTORAL POTION. (SAINTE-MARIE)

R. Salep, in powder, } of each, two drams.
Sugar of Milk, }
Syrup of Diacodium, one ounce.

Syrup of Marshmallows, two ounces.
Mix.—To be taken by spoonfuls.

WHITE CHOCOLATE. (PIERQUIN)

R. Cocoa, four ounces.
Salep, six ounces.
Water, eight ounces.

Boil over a slow fire for half an hour, and add,

White Sugar, four ounces.
Flour, a sufficient quantity.

Make cakes of half an ounce each.

APHRODISIAC ELECTUARY.

Electuarium diaspyrion. (PALAT.; WIRTEM.; BORIS; SPIELMANN)

R. Cinnamon, two drams.
Vanilla, eight drams.
Root of Orchis, six drams.
Horehound, two scruples.
Preserved Eryngo Root, two ounces.
Conserve of Elecampane, a sufficient quantity.

Mix. (BORIS)

R. Fresh Root of Orchis, four ounces.
Sugar, boiled à la plume, twelve ounces.

Beat together, strain through a silk sieve, and incorporate with

Preserved Ginger, }
Pine Nuts, } of each, one ounce.
Pistachio Nuts, }

Then add,

Powdered Cloves, } of each,
Cinnamon, } half an
Scincus Marinus, } ounce.
Syrup of Ginger, a sufficient quantity.

Make an electuary. (WIRTEM.; SPIELMANN)

R. Fresh Root of Eryngo, }
Parsnip, } of each,
Orchis, } half
Preserved Walnuts, } an ounce.

Pine Nuts, }
Pistachio Nuts, }
Seeds of Anise, }
Rocket, } of each, five drams.
Ash, }
Nettle, }

Cloves, }
Cinnamon, }
Ginger, }
Scincus Marinus, two drams and a half.
Musk, seven grains.
Despumated Honey, three pounds.

Make an electuary. (PALAT.)

Dose, from four to six drams.

SALIX.—WILLOW.

Five species of this genus of plants are mentioned in pharmacopœias :

1. *Salix alba*, LINN.—Willow.

SYNONIMES :—Saufe blanc, Sauf commun, FR. ; Weide, GERM. ; Wřba, BOHEM. ; Pıl, DAN. ; Sauce, SPAN. ; Wilg, DUT. ; Salice, ITAL. ; Kora wierzbowa, POL. ; Salguerro, PORT. ; Pıl, SWED.

DUBL. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; GALL. ; FERR. ; FULD. ; ROSS. ; SAX. ; WIRTEM. ; HERBIP. ; WURCEB. ; BRUGNATELLI ; COXE ; MURRAY and GMELIN ; ZARDA.

A shrub or tree common in all Europe. (*Diœcia Diandria*, LINN. ; *Salicineæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 402*)

The bark (*Cortex Salicis albae*) of the branches three or four years old is employed. It is thin, flexible, and reddish. It has a peculiar odour, a very astringent, bitter, and balsamic taste.

2. *Salix fragilis*, LINN.

SYNONIMES :—Saufe fragile, FR. ; Bruchweide, GERM.

DUBL. ; BAVAR. ; BRUNS. ; DAN. ; FERR. ; HANN. ; LIPP. ; OLDEN. ; ROSS. ; SAX. ; COXE.

The shrub is met with in almost every part of Europe. (*Fig. LINN. Fl. Lipp. t. 8. f. 6*)

The bark (*Cortex Salicis fragilis*) of one or two years old is employed. It has a slightly balsamic odour and a very bitter and astringent taste.

3. *Salix capræa*, LINN.

SYNONIMES :—Saufe marceau, FR. ; Soolweide, Sohlweide, GERM.

LOND. ; EDINB. ; HISP. ; GALL. ; FERR. ; COXE.

A common shrub of Europe. (*Fig. Pall. Flor. Ross. t. 81. fig. 2*)

The bark (*Cortex Salicis Capræa*) is astringent.

4. *Salix vitellina*, LINN.

SYNONIMES :—Saufe osier jaune, FR. ; Dotterweide, GERM.

DAN. ; ROSS.

A shrub of the temperate climates of Europe. (*Fig. Hoffm. Fasc. 1. 57. t. 11, 12. f. 1—3. t. 24, f. 1*)

The bark (*Cortex Salicis vitellinae*) is astringent.

5. *Salix pentandra*, LINN.

SYNONIMES :—Saufe pentandre, FR. ; Baumwollenweide, GERM.

AUSTR. ; BATAV. ; BELG. ; DAN. ; HANN. ; OLDEN. ; POL. ; CAST. BORUS. ; ROSS. ; SAX. ; SUEC. ; BERGIUS ; SPIELMANN.

A rather common tree of Europe. (*Fig. Ed. Fl. Dan. t. 943*)

The bark (*Cortex Salicis pentandrae seu Laureæ*) of the branches of one or two years old is employed. It is thin, flexible, and of a brown colour without ; it has a balsamic odour, and an agreeable, balsamic, bitterish, and styptic taste.

These barks contain a peculiar alkaloid called *Salicine* by FONTANA who discovered it. They have been proposed, under the title of an astringent tonic, as a substitute for cinchona in intermittent fevers.—Dose, from four drams to one ounce, of the powder.

ANTISEPTIC CATAPLASM. (FULD.)

R. Bark of <i>Salix Alba</i> ,	} of each, two ounces.
in powder,	
Herb of Rue,	
Spirit of Camphor,	
Vinegar,	three ounces.
	a sufficient quantity.

Mix.

OINTMENT OF WILLOW.

Unguentum Salicis. (HUFELAND)

R. Juice of Willow Leaves, two ounces.
Lard, a sufficient quantity.

Mix with a gentle heat.

It is applied to sordid, scorbutic, and gangrenous ulcers.

AQUEOUS EXTRACT OF WILLOW.

Extractum Salicis aquosum. (AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; FULD. ; HANN. ; OLDEN. ; POL. ; BORUS. ; SAX. ; BRUGNATELLI)

R. Willow Bark, one pound.
Cold Water, ten pounds.

Macerate for two days, stirring from time to time, and strain ; set aside, and decant the liquor, and evaporate it to the consistence of an extract. (POL.)

- R. Willow Bark, one part.
Water, twelve parts.

Digest for twelve hours in a moderate heat, then boil for one hour, and strain. Evaporate the liquor to the consistence of an extract. (BAVAR.)

- R. Willow Bark, one part.
Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and strain with expression; boil the residue with

Water, four parts,
and express again. Mix the decoctions, decant after sufficient repose, and evaporate to the proper consistence. (ANT.; HANN.; OLDEN.; BORUS.)

SAX.; and AUSTR. direct the bark to be exhausted by several successive ebullitions, and the mixed decoctions to be evaporated in a water-bath, after decantation.—BRUGNATELLI—the bark to be boiled for two hours with six parts of water; the boiling to be repeated with more water, and the mixed liquors evaporated.—BATAV.—the bark to be digested for twenty-four hours, and then boiled for two, with ten parts of water, and the boiling to be repeated with eight parts of water for the same length of time.—AMST.—the bark to be subjected to two several boilings, one with twenty and the other with fifteen parts of water, each time to be reduced to one half.

Dose, from ten to twenty grains.

ALCOHOLIC EXTRACT OF WILLOW. (FULD.)

- R. Willow Bark, one pound.
Proof Spirit, six pounds.

Digest for several days, and express; boil the residue with

Water, three pounds,
and express again. Reduce the strained

liquors separately to the consistence of honey; then mix them, and evaporate.

BREMER'S INFUSO-DECOCTION OF WILLOW. (NIEMANN)

- R. Willow Bark, four ounces.
Spirit from grain, eight ounces.
Water, sixteen ounces.

Digest in a covered vessel, and strain. Boil the residue with

Water, thirty ounces,
till reduced to fifteen ounces. Mix the two strained liquors together.

It is febrifuge.—To be consumed between two paroxysms.

DECOCTION OF WILLOW BARK.

Decoction Corticis Salicis lauræ. (CAST. BORUS.; NIEMANN)

- R. Willow Bark, cut small, one ounce.
Spring Water, sixteen ounces.

Boil till reduced to eight ounces. (NIEMANN)

CAST. BORUS. prescribes one ounce and a half of bark and fifteen ounces of water; to be boiled down to nine.

It is tonic, and febrifuge.—To be taken by glassfuls.

ANTISEPTIC INJECTION. (FULD.; HAMB.)

- R. Willow Bark, two ounces.
Water, one pound.

Boil down to two thirds. Add to the strained decoction,

Camphor, dissolved in a sufficient quantity of alcohol, half a dram.
Despumated Honey, one ounce.

Mix.

SALSOLA.—SALTWORT.

Three species of this genus are employed :

1. *Salsola Soda*, LINN.—Common Saltwort.

SYNONIMES :—Soude commune, FR.; Salzkraut, GERM.; Sul Alikantska lauhowa, BOHEM.; Saltyder, DAN.; Soda, SPAN.; Loogkruid, DUT.; Soda, ITAL.; Solda, PORT.; Soudaert, SWED.

HISP.; GALL.; MURRAY and GMELIN; ZARDA.

A plant of the coasts of the Mediterranean. (*Pentandria Digynia*, LINN.; *Che-*

nopodea JUSS.; *Fig. Jacq. Hort. Vind. I. p. 28. t. 69*)

The herb (*Herba Vitri*) is employed. It consists of branching, reddish stems, furnished with sessile, long, narrow, unarmed leaves. It is inodorous, but has a disagreeable alkaline taste.

2. *Salsola Fragus*, LINN.

SYNONIME :—Soude épineuse, FR.

GALL.; MURRAY and GMELIN.

A plant of the south of Europe.

3. *Salvia Kali*, LINN.

SYNONIMES :—Soude Kali, FR.

GALL.; GUIBOURT; MURRAY and GME-
LIN.

A plant of the south of Europe.

This species and the preceding one may
be substituted for the first.

SALVIA.—SAGE.

Four species of this genus of plants are
indicated in pharmacopœias :1. *Salvia officinalis*, LINN.—Common Sage.SYNONIMES :—Sauge commune, FR.; Sal-
bey, GERM.; Ssalweg, BOHEM.; Salvie,
DAN.; Salvia, ITAL.; and SPAN.; Salie,
DUT.; Szalwia, POL.; Salva, PORT.;
Salwia, SWED.EDINB.; DUBL.; AMER.; AMST.; ANT.;
AUSTR.; BATAV.; BAVAR.; BELG.;
BRUNS.; DAN.; MIL. COPEN.; HISP.;
GALL.; FENN.; FERR.; FORM. HÔP. FR.;
FULD.; HAMB.; HANN.; HASS.; LIPP.;
OLDEN.; LUSIT.; POL.; CAST. BORUS.;
BORUS.; ROSS.; SAX.; SUEC.; WIR-
TEM.; HERBIP.; WURCEB.; BERGIUS;
BRUGNATELLI; COXE; GUIBOURT;
MURRAY and GMELIN; PIDERIT; SPIEL-
MANN; ZARDA.A plant of the south of Europe. (*Dian-
dria Monogynia*, LINN.; *Labiata* JUSS.;
Fig. Zorn, Ic. pl. t. 165)The herb (*Herba Salvia minoris*) consists
of a square simple stem, and of opposite,
lanceolate, oval, entire, crenelated, rugose
leaves of a greenish-grey colour. It has an
agreeable odour, and a styptic, aromatic, and
bitterish taste.The flowers are of a violet-blue colour,
forming a terminal spike. The plant is gathered
before their expansion.It is excitant, nervine, tonic, sub-astring-
ent, resolvent, and detersive.2. *Salvia Horminum*, LINN.—Purple-topped
Sage, Ormin.SYNONIMES :—Sauge Hormin, FR.; Edles
scharlachkraut, GERM.; Horminio,
SPAN.; Edele slarey, DUT.; Orminio,
ITAL.; Ormino, PORT.BRUNS.; HISP.; GENEV.; HERBIP.; BER-
GIUS; SPIELMANN.A plant of Europe. (*Fig. Zorn, Ic. pl.
t. 244*)

The herb and seeds are employed.

The herb (*Herba Hormini* seu *Gallitrichi*)
consists of obtuse, crenelated leaves, of a
disagreeable odour and bitterish taste.The seeds are glabrous, mucilaginous,
and bitterish.

It is excitant and tonic.

3. *Salvia pratensis*, LINN.SYNONIMES :—Sauge des près, FR.; Wie-
sensalbey, GERM.

GALL.

A plant common in Europe. (*Fig.
Blackw. Herb. t. 258*)The leaves (*Folia Hormini*) are petiolated,
prolonged, heart-shaped, crenelated, rugose,
glabrous, of an agreeable odour, and of a
bitter and astringent taste.4. *Salvia Sclarea*, LINN.—Common Clarey.SYNONIMES :—Sauge Sclarée, FR.; Zahmes
Scharlachkraut, GERM.; Graakuse, DAN.;
Maro cortuso, SPAN.; Tamne sclarey,
DUT.; Schiarea, ITAL.; Esclarea, PORT.HISP.; GALL.; FERR.; WIRTEM.; GUI-
BOURT; MURRAY and GMELIN.A plant of the south of Europe. (*Fig.
Zorn, Ic. pl. t. 484*)The herb (*Herba Sclarea* seu *Gallitrichi*)
consists of a thick, square, very branching,
stem, with opposite branches; its leaves
are long, cordiform, rugose, pointed, ser-
rated, and opposite. It has a strong, and
disagreeable odour, and a bitter, and aro-
matic taste.

CONSERVE OF SAGE.

Conserva Florum Salviae. (SARD.; WIR-
TEM.)R. Flowers of Sage, one part.
White Sugar, two parts.

Beat together. (WIRTEM.)

SARD. prescribes one part of flowers and
three parts of sugar boiled à la plume.

AROMATIC SPECIES.

Species seu Herbe aromaticæ. (AUSTR.; BAVAR.; GALL.; NIEMANN; VAN MONS)

R. Herb of Sage,
 ——— Balm,
 ——— Thyme,
 ——— Peppermint,
 ——— Hyssop,
 Flowers of Lavender,

} of each,
 an equal part.

Cut, and mix.

R. Leaves of Sage,
 ——— Balm,
 Flowering Tops of Wormwood,
 ——— Peppermint,
 ——— Wild Thyme,
 Chamomile Flowers,

} of each,
 an equal part.

Cut and mix. (BAVAR.)

R. Leaves of Sage,
 ——— Thyme,
 ——— Wild Thyme,
 ——— Hyssop,
 ——— Water-mint,
 ——— Wormwood,
 ——— Origanum,

} of each,
 an equal part.

Cut and mix. (GALL.)

R. Herb of Hyssop,
 ——— Horehound,
 ——— Origanum,
 ——— Rue,
 ——— Sage,
 ——— Savory,
 ——— Scordium,
 ——— Wild Thyme,

} of each,
 an equal part.

Cut, and mix. (AUSTR.)

R. Cubebs,
 ——— Nutmegs,
 ——— Cinnamon,
 ——— Ginger,
 ——— Cloves,
 ——— Orange Peel,
 ——— Mentha Crispa,
 ——— Rosemary,
 ——— Sage,

} of each, one part.
 } of each, two parts.
 four parts.
 } of each, eight parts.

Cut, and mix. (VAN MONS)

EXTRACT OF SAGE.

Extractum Salviæ. (AUSTR.)

R. Leaves of Sage, any quantity.
 Water, a sufficient quantity.

Exhaust the leaves by several successive boilings; then mix the decoctions, and after decantation, evaporate to the consistence of an extract.

INFUSION OF SAGE.

Infusum Salviæ. (MIL. COPEN.; FORM. HÔP. FR.; CAST. BORUS.; SUEC.; WURCEB.; SAUNDERS)

R. Herb of Sage, one dram.

VOL. II.

Boiling Water, twelve ounces.

Infuse for a quarter of an hour in a covered vessel, and strain with expression. (MIL. COPEN.)

CAST. BORUS.; and WURCEB. prescribe one ounce and a half of sage and eighteen ounces of water.—SUEC.—one ounce of sage and one pound of water.—HAMB.—one ounce of sage and a sufficient quantity of water to obtain eight ounces of strained liquor.

R. Herb of Sage,
 ——— Liquorice,
 ——— Boiling Water,

} of each, one ounce.
 a sufficient quantity

to obtain two pounds of strained infusion.

R. Leaves of Sage, two ounces.
 Petals of Red Roses, one ounce.
 Boiling Water, three ounces and a half.

Infuse without heat, and strain. (SAUNDERS)

It is excitant and aromatic; to be taken by glassfuls.

ACIDULATED INFUSION OF SAGE. (NIEMANN)

R. Herb of Sage, half an ounce.
 Boiling Water, a sufficient quantity

to obtain, after half an hour's infusion, six ounces of liquor. Add to this,

Diluted Sulphuric Acid, one dram.

Mix.—Dose, half a glassful every two hours.

STIMULANT GARGLE. (CADET DE GASSICOURT; SAUNDERS)

R. Infusion of Sage, two pounds.
 Walnut Electuary, two ounces.
 Alum, one dram.

Mix. (SAUNDERS)

R. Infusion of Sage, six ounces.
 Tincture of Cinchona, } of each,
 Syrup of Mulberries, } half an ounce.
 Spirit of Scurvy-grass, one dram.

Mix. (CADET DE GASSICOURT)

VINOUS INFUSION OF SAGE.

Infusum Salviæ vinosum. (NIEMANN)

R. Herb of Sage, one ounce.
 Boiling Water, one pound.

Digest for one hour; strain, and add to the cold liquor,

Red Wine, half a pound.

Mix.—Dose, half a glassful every two hours.

AROMATIC DECOCTION.

Decoctum incitans aromaticum. (BRERA)

R. Leaves of Sage, half an ounce.
 Water, a sufficient quantity,

4 A

to obtain two pounds of strained decoction.
Add to it

HALLER'S Acid Elixir, one dram.

Mix. — To be taken gradually in the course of the day.

AROMATIC WINE. (FORM. HÔP. FR. ;
GALL.)

R. Aromatic Species, four ounces.
Red Wine, twelve ounces.

Macerate for six days in a covered vessel;
then express, and filter. (GALL.)

R. Red Wine, ten parts.
Aromatic Tincture, one part.

Mix. (FORM. HÔP. FR.)

It is employed externally in lotions or fomentations.

WINE OF SAGE.

Vinum Salviæ. (HERBIP.)

R. Fresh Leaves of Sage, one ounce and a half.
Red Wine, one pound and a half.

Macerate for twenty-four hours, and strain.

EXCITANT FOMENTATION. (RATIER)

R. Wine of Sage, one pound.
Spirit of Camphor, eight ounces.

Mix.

RED LIQUOR; SPIRITUOUS VULNERARY
LIQUOR.

Aqua vulneraria spirituosâ. (GALL. ; VAN
MONS)

R. Fresh Leaves of Sage

_____ Rosemary,
_____ Savory,
_____ Origanum,
_____ Marjoram,
_____ Thyme
_____ Wild-

Thyme,

_____ Hyssop,
_____ Balm,
_____ Calamint,
_____ Sweet Basil,
_____ Water-

mint,

_____ Fennel,
_____ Angelica,
_____ Wormwood,
_____ Rue,

Flowering Tops of Lavender,
_____ St. John's

wort,

Alcohol (10 degrees), two pounds.

Macerate for eight days; then express and filter. (GALL.)

R. Fresh Leaves of Wormwood,
_____ Angelica,
_____ Sweet Basil,
_____ Southern-

wood,

_____ Fennel,
_____ Hyssop,
_____ Marjoram,
_____ Balm,
_____ Water-mint,
_____ Origanum,
_____ Rue,
_____ Rosemary,
_____ Sage,
_____ Wild Thyme,
_____ Thyme,

Flowers of Lavender,

Proof Spirit, thirty-two parts.

Digest without heat for eight days; then express and filter. (VAN MONS)

VINEGAR OF SAGE.

Acetum Salviæ. (GALL. ; VAN MONS)

R. Fresh Leaves of Sage, one part.
Vinegar, sixteen parts.

Macerate for fifteen days in a covered vessel, agitating from time to time; then strain and filter. (GALL.)

VAN MONS prescribes one part of sage and eight parts of vinegar, to be macerated for eight days.

OIL OF SAGE.

Oleum Salviæ æthereum, Ætheroleum Salviæ. (AUSTR. ; BRUNS. ; GALL. ; HANN. ;
PALAT. ; SARD. ; BRUGNATELLI)

R. Flowering Herb of Sage, any quantity,
Water, a sufficient quantity.

Distil. (BRUGNATELLI)

HANN. prescribes one part of herb and eight parts of water.—AUSTR.—one of herb and twelve of water.—GALL.—five of herb and seven of water.

R. Flowering Herb of Sage, twenty-five parts.

Water, seventy-five parts.
Muriate of Soda, three parts.

Distil at the expiration of three days: (BRUNS. ; PALAT. ; SARD.)

DISTILLED WATER OF SAGE

Aqua Salviæ. (AMST. ; AUSTR. ; BAVAR. ;
GALL. ; HANN. ; LIPP. ; OLDEN. ; PALAT. ; POL. ; BORUS. ; SARD. ; SAX. ;
HERBIP. ; SWEDIAUR. ; VAN MONS)

R. Flowering Herb of Sage, one part.
Water, a sufficient quantity.

At the expiration of twenty-four hours, distil ten parts. (AMST. ; HANN. ; OLDEN. ;
POL. ; BORUS. ; SWEDIAUR)

of
each,
one
part.

of each,
one
ounce.

GALL.; and PALAT. prescribe one part of herb and four parts of water; two parts to be distilled.—LIPP.; and HERBIP.—one of herb and six of water.—AUSTR.—one of herb and twelve of water; four to be distilled.—SAX.—one of herb and fifteen of water; ten to be distilled.—BAVAR.—one of herb and sixteen of water; four to be distilled.—SARD.—one of herb and four of water; two to be distilled, and the product to be re-distilled with fresh herb.

R. Flowering Herb of Sage, ten parts.
Water, one hundred and twenty parts.
Proof Spirit, one part.
Distil thirty parts. (VAN MONS)

COMPOUND WATER OF SAGE. (GALL.)

R. Vulnerary Species, two parts.
Water, twenty parts.
Distil eight parts.

SPIRIT OF SAGE.

Aqua Salviae spirituosus. (GALL.; FERR.)

R. Flowering Herb of Sage, one pound.
White Wine, four pounds.
Macerate for twenty-four hours, and distil one half. (GALL.)

R. Flowering Herb of Sage, one pound and a half.
Alcohol, eight pounds.
Distil six pounds. (FERR.)

VULNERARY SPIRIT; EAU D'ARQUE-BUSADE.

Aqua vulneraria seu traumatica seu scopetaria seu vulneraria spirituosus seu vinosa seu scopetaria aromatica seu Catapulta-rum, Alcohol Salviae vulnerarius. (AUSTR.; BELG.; BRUNS.; HISP.; GALL.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; WIRTEM.; BRUGNATELLI; CADET DE GASSICOURT; PIDERIT; NIEMANN; SPIELMANN; VAN MONS)

R. Herb of Hyssop, } of each,
Mentha Crispa, } two ounces.
Sage,
Rosemary,
Flowers of Rosemary } of each,
Lavender, } one ounce.
Proof Spirit, ten pounds.

Distil till the liquor ceases to pass over limpid. (LIPP.; WIRTEM.)

R. Herb of Wormwood, } of each,
Dill, } one ounce
Sage, } and a half.
Tansey,
Wild Thyme,
Peppermint,
Flowers of Lavender, }

Spring Water, twelve pounds.
Alcohol, eight pounds.

Infuse for several days, and distil eight pounds. (PIDERIT)

R. Herb of Wormwood, } of each,
Rue, } eighteen drams.
Sage,
Fennel, } of each,
Peppermint, } one ounce
Flowers of Lavender, } and a half.
Spring Water, twelve pounds.
Alcohol, eight pounds.

Infuse for two days, and distil eight pounds. (HASS.)

R. Fresh Herb of Agrimony, }
Ladies }
Mantle, }
Daisy, }
Chrysan- }
themum, }
Betony, } of each,
Bugloss, } two
Fennel, } handfuls.
St John's- }
wort, }
Plantain, }
Sanicle, }
Scrofularia }
Vervain, }
Birthwort, } of each,
Mugwort, } four
Golden Rod, } handfuls
Sage, }
Wine, forty pounds.

Digest till the mass begins to ferment, and distil slowly one half. (WIRTEM.)

R. Root of Comfrey, } of each,
Herb of Comfrey, } four handfuls.
Mugwort, }
Bugle, }
Betony, }
Ox-eye, }
Lesser Comfrey, } of each,
Scrofularia, } two
Plantain, } handfuls.
Agrimony, }
Vervain, }
Wormwood, }
Fennel, }
St. John's-wort, }
Birthwort, }
Small Houseleek, }
Veronica, } of each,
Lesser Centaury, } one
Milfoil, } handful.
Tobacco, }
Mint, }
Hyssop, }
White Wine, one hundred and forty-four ounces.

Infuse for three days, and distil. (SPIELMANN)

Root of Comfrey,	} of each, four ounces.	
Herb of Comfrey,		
— Bugle,		
— Sanicle,		
— Betony,		
— Scrofularia,		
— Plantain,		
— Agrimony,		
— St. John's-wort,		
— Periwinkle,		
— Ground Ivy,	} of each, eight ounces.	
— Mugwort,		
— Veronica,		
— Vervain,		
— Lesser Houseleek,		
— Milfoil,		
— Sage,		
— Angelica,		
— Tansey,		
— Wormwood,		
— Great Celandine,	} of each, four ounces.	
— Fennel,		
— Clematis,		
— Mint,		
— Hyssop,		
— Tobacco,		
— Rue,		
— Chamomile,		
— Scordium,		
— Marjoram,		
— Rosemary,	} of each, six ounces.	
Flowering Tops of Lavender,		
— Origanum,		
— Calamint,	} of each, fifty pounds.	
White Wine,		

Distil one half. (WIRTEM.)

PALAT. gives the same formula, adding eight ounces each, of Roman wormwood and of savory, and four ounces each, of sweet basil and of golden rod.

R. Leaves of Sage,	eight ounces.
— Rosemary,	} of each, one ounce.
— Wild Thyme,	
— Thyme,	
— Lesser Cardamoms,	} of each, six drams.
— Great Cardamoms,	
— Mace,	} of each, half an ounce.
— Nutmegs,	
— Cloves,	
— Zedoary,	
— Galanga,	
— Sweet Flag,	
White Wine,	eight pounds.
Alcohol,	four pounds.

After sufficient digestion distil three-fourths. (WIRTEM.)

R. Flowers of Lavender,	} of each, one ounce.	
— Fresh Herb of Hyssop,		
— Balm,		
— Mentha Crispa,		
— Origanum,		
— Rosemary,		
— Rue,		
— Sage,	} of each, two ounces.	
— Wild Thyme,		

Alcohol (0.850), ten ounces.
Water, twelve pounds.
After twenty-four hours, distil ten pounds. (AUSTRIAN.)

R. Fresh Herb of Sage,	} of each, four ounces.	
— Wormwood,		
— Peppermint,		
— Rue,		
— Rosemary,	} of each, six pounds.	
Flowers of Lavender,		
Alcohol,	} a sufficient quantity.	
Water,		

At the end of twenty-four hours, distil eighteen pounds. (HANN.; OLDEN.; POL.; BORUS.; NIEMANN)

R. Fresh Leaves of Sage,	} of each, four ounces.	
— Angelica,		
— Wormwood,		
— Savory,		
— Fennel,		
— Mint,		
— Hyssop,		
— Balm,		
— Sweet Basil,		
— Rue,		
— Thyme,		
— Marjoram,		
— Rosemary,		
— Origanum,	} of each, one hundred and twenty-eight ounces.	
— Calamint,		
— Wild Thyme,		
Flowers of Lavender,	} of each, one hundred and twenty-eight ounces.	
Alcohol,		

Distil. (SPIELMANN)

R. Flowering Herb of Wormwood,	} of each, two parts.	
— Sweet Basil,		
— Cat-mint,		
— Hyssop,		
— Marjoram,		
— Origanum,		
— Rosemary,		
— Rue,		
— Sage,		
— Thyme,		
Flowers of Lavender,	} of each, ninety-six parts.	
Proof Spirit,		
Water,	} four hundred and fifty-six parts.	

Distil one hundred and ninety-four parts. (VAN MONS)

BELG. prescribes one part each, of sage, of wormwood, of cat-mint, of hyssop, of balm, of sweet basil, of thyme, of rue, of marjoram, of rosemary, of origanum, and of lavender, forty-eight parts of proof spirit, and two hundred and seventy-six of water; ninety-two parts to be distilled.

R. Tops of Sage,	} of each, two ounces.	
— Tansey,		
— Fennel,		
— Thyme,		
— Lavender,		
— Wormwood,	} of each, two ounces.	

Tops of Water-mint,	}	of each, two ounces.
_____ Hyssop,		
_____ Chamomile,		
_____ Origanum,		
_____ Marjoram,		
_____ Calamint,		
Stem and Root of Angelica,	}	}
Alcohol (22 degrees),		
_____	twenty-four pounds.	

After sufficient maceration, distil twenty pounds in a water-bath. (GALL.; CADET DE GASSICOURT)

R. Flowers of Lavender,	}	of each, one ounce.
Fresh Herb of Wormwood,		
— Sage,		
— Marjoram,		
— Rosemary,		
— Hyssop,		
— Rue,		
— Balm,		
— Thyme,		
— Sweet Basil,		
— Origanum,		
— St. John's-		
wort,		
Alcohol (0.935),		four pounds.
Water,		one pound.

At the expiration of twenty-four hours, distil three pounds. (BRUGNATELLI)

R.	Leaves of Betony,	}	of each, fifteen ounces.
—	Ground Ivy,		
—	Mugwort,		
—	Milfoil,		
—	Sage,	}	of each, ten ounces.
—	Angelica,		
—	Tansey,		
—	Wormwood,		
—	Southernwood,	}	of each, eight ounces.
—	Fennel,		
—	Mint,		
—	Hyssop,		
—	Tobacco,	}	of each, four ounces.
—	Rue,		
—	Chamomile,		
—	Scordium,		
—	Marjoram,	}	
—	Rosemary,		
	Flowering Tops of Laven-	}	of each, six ounces.
	der,		
	— Origa-		
	num,		
	— Cala-	}	
	mint,		
	Wine,	sixty-seven pounds.	

Macerate for twenty-four hours, and distil one half. (BRUNS.)

R. Leaves of Betony,	}	of each, one pound.
— Agrimony,		
— Mugwort,	}	of each, eight ounces.
— Sage,		
— St. John's-wort,		
— Tansey,		
— Wormwood,	}	
— Mint,		

Leaves of Scordium,	}	of each, four ounces.
— Marjoram,		
— Rosemary,		
— Lavender,		
White Wine,	thirty pounds.	

Distil one half in a water-bath. (HISP.)

R. Herb of Sage,	}	of each, one part.
— Wormwood,		
— Peppermint,		
— Rue,		
— Rosemary,	}	
Flowers of Lavender,		
Water,		fifty parts.
Alcohol,	twenty-two parts.	

At the expiration of two days, distil thirty-six parts. (SAX.)

This liquor, which scarcely differs from the *Eau de Cologne* and the spirit of balm, is employed externally as resolvent, and sometimes internally in flatulent colic.

AROMATIC SPIRIT.

Elixir Vitæ mulierum, Aqua seu Balsamum Embryonum, Aqua aromatica. (AMST.; BRUNS.; PALAT.; BORUS.; WIRTEM.; NIEMANN.; SPIELMANN)

R.	Crumb of New Bread,	two pounds.
	Flowers of Clove-pink,	of each,
	——— Lime Tree,	one pound.
	Nutmegs,	three ounces and a half.
	Flowers of Lavender,	
	Herb of Sage,	of each,
	Seeds of Caraway,	three ounces.
	——— Fennel,	
	Cloves,	
	Cinnamon,	
	Ginger,	
	Cubebs,	
	Grains of Paradise,	of each,
	Zedoary,	two
	Herb of Mentha Crispa,	ounces.
	Root of Peony,	
	Seeds of Peony,	
	Misceltoe,	
	Mace,	half an ounce.
	Lesser Galanga,	three drams.
	Saffron,	two drams.
	Madeira,	ten pounds.
	Water of Borage,	of each,
	——— Bugloss,	four pounds.
	——— Lily of the Valley,	
		two pounds eight ounces.
	——— Strawberries,	of each,
	——— Roses,	sixteen
		ounces.

Digest for three days, distil one half of the liquid, over a slow fire. (WIRTEM.)

R. Galanga,	}	of each, half an ounce.
Zedoary,		
Herb of Mint,	}	of each, one ounce.
— Balm,		

Flowers of French Lavender, } of each,
 ——— Clove-pink, } five
 ——— Lime Tree, } drams.
 Seeds of Siler, } of each,
 ——— Fennel, } half an ounce.
 Nutmegs, } one ounce.
 Cinnamon, } six drams.

Fresh Orange Peel, } of each,
 ——— Lemon Peel, } one ounce and
 a half.

White Wine, } two pounds.

Digest for twenty-four hours, and distil one half. (PALAT.; WIRTEM.)

BRUNS. gives the same formula, but substitutes for the wine a mixture of five pounds of proof spirit and ten of water, and directs six pounds of the liquor to be distilled.

R. Root of Peony, }
 ——— Zedoary, } of each, two ounces.
 ——— Ginger, }
 ——— Galanga, } six drams.
 Herb of Sage, }
 ——— Balm, } of each,
 ——— Fennel, } three ounces.
 ——— Mint, }
 Flowers of Clove-pink, } of each,
 ——— Lime Tree, } six ounces.
 ——— Lavender, } of each,
 Nutmegs, } three ounces.
 Cloves, }
 Cinnamon, } of each,
 Cubebs, } two ounces.
 Grains of Paradise, }
 Mace, }
 Alcohol, } forty-four ounces.
 Spring Water, }
 two hundred and forty ounces.

Infuse for several days, and distil one

hundred and ninety-two ounces. (SPIELMANN)

R. Nutmegs, }
 Mace, }
 Cloves, } of each, half an ounce.
 Cinnamon, }
 Cubebs, }
 Ginger, }
 Zedoary, }
 Lesser Galanga, } two drams.
 Flowers of Lavender, } of each,
 Dried Herb of Mint, } half an
 ——— Sage, } ounce.
 Seeds of Coriander, } of each,
 ——— Fennel, } two drams.
 Proof Spirit, } one pound and a half.
 Water, } seven pounds and a half.

Macerate for three days, and distil three pounds and a half. (AMST.)

R. Herb of Sage, } eight ounces.
 ——— Rosemary, } of each
 ——— Peppermint, } four ounces.
 Flowers of Lavender, }
 Seeds of Fennel, } of each, two ounces.
 Cinnamon, }
 Alcohol, } four pounds.
 Water, } twenty pounds.

Macerate for twenty-four hours, and distil twelve pounds. (BORUS.; NIEMANN)

It is excitant, corroborant, cordial, and carminative.—Dose, from half an ounce to one ounce.—It is also applied externally. A crust of hot bread saturated with it used formerly to be applied, and it is said with success, to the abdomen in cases of threatened abortion. It is now better known how these stimulant applications should be appreciated.

SAMBUCUS.—ELDER.

Two species of this genus are employed in medicine:

1. *Sambucus Ebulus*, LINN.—Dwarf Elder.

SYNONIMES:—Hieble, Yèble, FR.; Zwerg-hollander, Attich, Attichhirschschwanz, GERM.; Chebdj, BOHEM.; Mmerhyld, DAN.; Yezgo, SPAN.; Lange vlier, DUT.; Ebbio, ITAL.; Chebd, POL.; Engos, PORT.; Sommarhyll, SWED.

AUSTR.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; LUSIT.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; MURRAY AND GMELIN.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

A shrub of Europe. (*Pentandria Trigy-*

nia, LINN.; *Caprifoliaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 40*)

The bark of the root, the leaves, fruit, and seeds are employed.

The inner bark of the root (*Radix Ebuli* seu *Sambuci humilis* seu *agrestis* seu *Chamaectis*) is white, and of a bitter and nauseous taste. The root itself is long, and of the thickness of the finger. It is esteemed emetic and hydragogue.—Two ounces of it are prescribed, infused in water: one ounce of the expressed juice may also be given.

The leaves are oval, lanceolate, acute, dentate, smooth and shining above, and of a deep green colour. They have a strong, disagreeable odour. They were formerly applied externally, in cases of œdema and erysipelas.

The fruit (*Baccæ seu Grana Actes*) consists of black, tri-spermal berries, of a sweet, bitter, and nauseous taste. They are reputed resolvent, sudorific, and diuretic.

The seeds, macerated in water and submitted to a press, furnish oil.

JUICE OF DWARF ELDER BERRIES.

Succus Ebuli. (GALL.; SARD.)

- R. Ripe Berries of Dwarf Elder,
any quantity.

Bruise them with the hands, leave the mass to ferment for three or four days, and express. Preserve the juice in bottles, under a stratum of oil.

ROB OR PULP OF DWARF ELDER.

Robb seu Pulpa Ebuli. (AUSTR.; BAVAR.; BRUNS.; FULD.; PALAT.; SARD.; WIRTEM.; VAN MONS)

- R. Fresh and Ripe Berries of
Dwarf Elder, one part.
Water, a sufficient quantity

to prevent them from burning. Having boiled them till they burst, express the juice, and evaporate it to the consistence of pulp, adding one sixteenth of sugar. (BAVAR.)

AUSTR. prescribes one part of sugar, and nine parts of juice.—VAN MONS—one of sugar, and ten of juice.—BRUNS.; FULD.; PALAT.; and WIRTEM.—one of sugar, and six of juice.

- R. Juice of Dwarf Elder Berries,
any quantity.

Strain it, and evaporate it in a water-bath to the consistence of an electuary. (SARD.)

It is diuretic and laxative.—Dose, 'two ounces.—It is seldom prescribed alone, but generally as an excipient.

DISTILLED WATER OF DWARF ELDER.

Aqua Ebuli. (WIRTEM.)

- R. Bruised Berries of Dwarf Elder,
any quantity.

Put them in a barrel, and leave them to ferment; when they have acquired a vinous odour, distil over a slow fire: rectify the product, if necessary.

2. *Sambucus nigra*, LINN.—Elder.

SYNONIMES:—Sureau noir, Sureau, FR.; Fliederhollerbaum, GERM.; Ukti, Khaman, ARAB.; Bez, BOHEM.; Hyld, DAN.; Saucos, SPAN.; Vlier, DUT.; Sambuco, ITAL.; Bzowy, POL.; Sabu-guiero, PORT.; Busina, RUSS.; Flæder, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GME-LIN.; PIDERIT.; SPIELMANN.

A shrub of Europe. (*Fig. Flore Medic. VI. 336*)

The bark, leaves, flowers, and fruit are employed.

The middle or inner bark of the branches (*Cortex seu Alburnum Sambuci*), from which the thin ash-coloured pellicle is usually separated, is of a green colour. It has a strong and nauseous odour, a taste at first sweetish, and afterwards acid and bitterish.

The leaves are unequally wing-cleft, of a deep green colour, glabrous, with leaflets oval, lanceolated, serrated, and acuminate. Their odour is disagreeable, and their taste acrid. The flowers are white, small, of a strong and disagreeable odour, and of a mucilaginous and bitterish taste.

The fruit consists of succulent, oval, black, umbilicated, tri-spermal berries, containing a reddish, acidulous, inodorous juice. They are called *Grana Actes* when dried. The American pharmacopœia replaces them by those of the *Sambucus Canadensis*, WILLD.

POWDER AGAINST ERYSIPELAS.

Pulvis ad Erysipelam. (CAST. BORUS.)

- R. Elder Flowers, } of each,
Wheat Bran, } an equal part.

Mix together.

SUDORIFIC SPECIES. (GALL.)

- R. Flowers of Elder, } of each,
—— Borage, } an equal part.
—— Red Poppy,
Sassafras Wood,

Mix.

JUICE OF ELDER BERRIES.

Succus Sambuci. (GALL.; SARD.)

- R. Ripe Elder Berries, any quantity

Bruise them with the hands, and leave the mass to ferment for three or four days; then express the juice, and preserve it in bottles, under a stratum of oil.

Dose, from half an ounce to two ounces.

ELDER POWDER.

Tragea Granorum Actes. (WIRTEM.; SPIELMANN)

- R. Juice of Elder Berries, any quantity.
Rye Meal, a sufficient quantity.

Reduce to a paste, and form it into small cakes; expose these before a fire till they become perfectly dry, and reduce them to powder: impregnate this with fresh juice, and dry again. Having repeated this operation three successive times, add to one ounce of the powdered mass,

Powdered Nutmeg, one dram.

Mix well.—It is recommended in dysentery.—Dose, one or two drams.

SYRUP OF ELDER.

Syrupus Sambuci. (AUSTR.)

- R. Recently expressed Juice of
Elder Berries, one pound.
White Sugar, two pounds.
Boil for a few moments.

ROB OF ELDER.

Roob Sambuci, Succus Sambuci inspissatus.
(EDINB.; AMER.; ANT.; AUSTR.; BAT-
TAV.; BAVAR.; BELG.; DAN.; GALL.;
FERR.; FULD.; GENEV.; HANN.; HASS.;
LIPP.; OLDEN.; PALAT.; POL.; BORUS.;
ROSS.; SARD.; WIRTEM.; HERBIP.;
BRUGNATELLI; COXE; SWEDIAUR;
VAN MONS)

Succus spissatus Sambuci nigrae.
(EDINB.)

- R. Ripe Elder Berries, five parts.
Refined Sugar, one part.
Boil slowly together to the consis-
tence of thick honey.

R. Ripe Elder Berries, any quantity.
Bruise them with the hands, express the juice, and strain it through flannel. Add half a pound of sugar to two pounds of this juice, and evaporate over a quick fire to the consistence of honey. (VAN MONS)

PALAT. prescribes one part of sugar, and two parts of juice.—AMER.; and COXE—one of sugar, and five of juice.—FULD.—one of sugar, and four of juice.—BRUNS.; HASS.; ROSS.; WIRTEM.; HERBIP.; and SWEDIAUR—one of sugar, and six of juice.—AUSTR.—one of sugar, and nine of juice.—OLDEN.; POL.; BORUS.; and SAX.—one of sugar, and sixteen of juice.

R. Ripe Elder Berries, any quantity.
Heat them in a covered tin vessel, constantly stirring, for half an hour, or till they shall have burst; then strain the liquor through a sieve, and express the residue

strongly. Mix the two liquors, add four pounds of sugar to fifteen of the juice, and evaporate to the consistence of honey, skinning well. (BELG.)

BATAV. prescribes four pounds of sugar, and ten of juice.—ANT.—one of sugar, and five of juice.—DAN.—one of sugar, and four of juice.—BAVAR.—one of sugar, and sixteen of juice.—FERR.—two of sugar, and four of juice.

- R. Expressed Juice of Elder Berries,
any quantity.

Set it aside to depurate by repose; then evaporate it to the proper consistence over a slow fire. (GALL.; GENEV.; LIPP.; SARD.; BRUGNATELLI)

It is diaphoretic.—Dose, from one ounce and a half to two ounces.

EXTRACT OF ELDER FLOWERS.

Extractum Florum Sambuci. (SARD.)

- R. Fresh Elder Flowers, eight pounds.

Macerate them for three days in an alembic; then distil in a water-bath about two ounces of liquor: pour upon the residue thrice its weight of water, and after six hours' digestion, boil down to one third. Express the flowers, clarify and strain the liquor, and evaporate to the consistence of an electuary, adding the distilled water to this when it is no more than warm.

ANTI-ASTHMATIC MIXTURE. (SAINTE-MARIE)

- R. Rob of Elder,
Pulp of Prunes,
Syrup of Red Poppy,
Nitrate of Potass, } of each,
two ounces.
one dram.

Mix.—Dose, a tea spoonful, three or four times a day.

HONEY OF ELDER FLOWERS.

Mel Sambuci. (WIRTEM.)

- R. Fresh Elder Flowers, one pound.
Despumated Honey, two pounds.
Elder-flower Water, six ounces.

Macerate for several days, then boil over a slow fire; express, and evaporate to the required consistence.

It used formerly to be added to emollient and resolvent gargles.

HONEY OF ELDER BERRIES.

Mel Fructuum Sambuci. (HISP.)

- R. Juice of Ripe Elder
Berries,
Honey,
Water, } of each,
two pounds.
a sufficient quantity.

Boil to the consistence of thick syrup, and strain.—Dose, from half an ounce to one ounce.

VINEGAR OF ELDER.

Acetum Sambuci seu sambucinum. (AMST.; BRUNS.; GALL.; LUSIT.; PALAT.; SARD.; SUEC.; WIRTEM.; HERBIP.; PIDERIT; SWEDIAUR)

R. Elder Flowers, one part.
Vinegar, twelve parts.

Macerate for four days, clarify with milk, and filter. (SUEC.)

GALL. prescribes one part of flowers, sixteen parts of vinegar, and fifteen days' maceration.—BRUNS.; PALAT.; WIRTEM.; and SWEDIAUR—one of flowers, and six of vinegar.—SARD.; and PIDERIT—one of flowers, four of vinegar, and six days' maceration.—HERBIP.—one of flowers, nine of vinegar, and three weeks' maceration.—AMST.—one of flowers, eight of vinegar, and four days' maceration.

R. Dried Elder Flowers, one pound.
Vinegar, six pounds.

Macerate for six days; then boil. Add to the strained liquor

Alcohol, half a pound.

Mix. (LUSIT.)

It is sudorific and detergent.—It is administered with hot water, sweetened, to produce diaphoresis. It is also employed in the form of gargle, in ulcerated sore throat, mixed with water and honey.

OXYMEL OF ELDER.

Oxymel Sambuci. (HERBIP.)

R. Vinegar of Elder, one part.
Honey, two parts.

Boil slowly to the consistence of syrup.

DECOCTION OF ELDER BARK.

Decoctum Corticis Sambuci. (NIEMANN)

R. Elder Bark, three handfuls.
Spring Water, } of each, one pound.
Milk, }

Boil down to one pound of strained liquor.

It is recommended by BOERHAAVE, in dropsies.—Dose, one half in the morning and the other in the evening.

INFUSION OF ELDER FLOWERS.

Infusum Florum Sambuci nigrae. (ANT.; MIL. COPEN.; GALL.; FORM. HÔP. FR.; HAMB.; DISP. CHA. PARIS; RATIER)

R. Elder Flowers, one dram.
Boiling Water, one pound.

VOL. II.

Digest for a quarter of an hour, and express. (MIL. COPEN.; DISP. CHA. PARIS; RATIER)

HAMB.; and FORM. HÔP. FR. prescribe one ounce of flowers and a sufficient quantity of boiling water to obtain eight ounces of strained liquor.—ANT.—one ounce of flowers and one pound of water; to be digested for an hour, gradually increasing the heat to the boiling point.

R. Elder Flowers, two drams.
Boiling Water, two pounds.

Infuse for a quarter of an hour; then strain, and add to the liquor,

Syrup of Maidenhair, one ounce.

Mix. (GALL.)

It is prescribed as diaphoretic in rheumatic and acute exanthematous affections.—It is administered hot and largely, sweetened with honey, oxymel, or tartaric syrup.

COMPOUND SULPHUREOUS WATER. (CADET DE GASSICOURT)

R. Infusion of Elder Flowers, } of each,
Barages Water, } one pound

Mix.—To be taken in twenty-four hours; in chronic catarrhs, tabes mesenterica, scrofula, and tetters.

DETERGENT GARGLE. (FULD.; SAUNDERS)

R. Elder Flowers, one ounce.
Boiling Water, one pound.

Infuse, strain, and add

Nitrate of Potass, half an ounce.

Tincture of Burnet Saxifrage,

three drams.

Oxymel, two ounces.

Mix. (FULD.)

R. Flowers of Elder, } of each,
Lesser } one handful.
Mullein, }

Boiling Water, a sufficient quantity,
to obtain half a pound of strained infusion.
Add to it

Nitrate of Potass, half a dram.

Sulphate of Potass, one dram.

Rob of Elder, } of each,

Syrup of Mulberries, } one ounce.

Mix. (SAUNDERS)

RESOLVENT FOMENTATION. (RATIER; SPIELMANN)

R. Infusion of Elder Flowers, one pound.
Spirit of Camphor, eight ounces.

Mix. (RATIER)

R. Elder Flowers, one handful.
Water, sixteen ounces.

4 B

Infuse for twelve hours. Add to the strained liquor,

Spirit of Camphor, one ounce.
Acetate of Lead, one dram.

Mix. (SPIELMANN)

R. Flowers of Melilot, } of each,
Elder, } two drams.
Leaves of Mint, } of each, one dram.
Sage, }
Boiling Water, two pounds.

Add to the strained liquor,
Proof Spirit, three ounces.

Mix. (RATIER)

ELDER OIL.

Oleum Sambuci. (HISP.; SARD.)

R. Elder Flowers, one part and a half.
Olive Oil, three parts.

Digest for two days in a water-bath; then set apart to cool, and express. Repeat the operation with fresh flowers. (SARD.)

HISP. prescribes one part of flowers and three parts of oil.

ELDER OINTMENT.

Unguentum Sambuci seu Sambucinum.
(LOND.; DUBL.; HISP.; COXE; SPIELMANN)

Unguentum Sambuci.

R. Elder Flowers, } of each,
Prepared Lard, } two pounds.

Boil the flowers in the lard, till they become crisp, and strain. (LOND.; COXE)

R. Fresh Elder Flowers, three pounds.
Prepared Lard, four pounds.
Mutton Suet, two pounds.

Boil the flowers with the lard; then express, and add the suet. (DUBL.)

R. Mutton Suet, three parts.
Olive Oil, one part.

Melt together, and digest in the mixture,
Elder Flowers, four parts.

Strain after some time. (SPIELMANN)

ELLIS directs four parts of flowers to be boiled with three of suet and one of olive oil.

R. Elder Bark, two pounds and a half.
Flowers, half a pound.
Lard, nine pounds.

Macerate for eight days, and boil to the consumption of the humidity. (HISP.)

ELDER-FLOWER WATER.

Aqua Florum Sambuci. (AMST.; AUSTR.;

BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT)

R. Elder Flowers, three parts.
Water, a sufficient quantity.

Distil six parts. (SUEC.)

AUST. prescribes six parts of flowers and a sufficient quantity of water; eight parts to be distilled.—GALL.; and SARD.—one of flowers and four of water, two to be distilled.—BAVAR.—one of flowers and sixteen of water; four to be distilled.—PALAT.—one of flowers and three of water; one and a half to be distilled.—AMST.; BATAV.; BELG.; DAN.; HANN.; OLDEN.; and POL.—one of flowers and a sufficient quantity of water; ten to be distilled.—HASS.; ROSS.; SAX.; and PIDERIT—one of flowers and eight of water; four to be distilled.—HERBIP. and BRUGNATELLI—one of flowers and six of water.—BORUS.—one of flowers and fifteen of water; ten to be distilled.—FERR.—three of flowers and six of water; six to be distilled.

R. Elder Flowers, three parts.
Water, twelve parts.

Enclose the flowers in a bag suspended in the middle of a cucurbit and not touching the water. Distil six parts. (FULD.; LIPP.)

R. Fresh Elder Flowers, sixteen parts.
Water, forty parts.

Macerate for two days, and when fermentation appears to have commenced, distil twenty parts. Add to the product,

Alcohol, one part.

Mix. (HISP.)

ELDER-BERRY WATER. (BRUNS.; WIRTEM.)

R. Bruised Elder Berries, any quantity.

Leave them to ferment in a barrel till they acquire a vinous odour; then distil slowly, and rectify the product if necessary.

ELDER JULEP.

Julapium Sambucinum. (WIRTEM.)

R. Elder Water, twenty ounces.
White Sugar, two pounds and a half.

Boil for an instant, and strain.

SPIRIT OF ELDER.

Spiritus Sambuci. (BRUNS.; PALAT.; WIRTEM.)

R. Elder Flowers, one part.
Proof Spirit, four parts.

Macerate for three days, and distil one half.

SANDARACA.—SANDARAC, GUM JUNIPER.

SYNONIMES :—*Gummi seu Resina Juniperi*, *Vernix sicca*; *σανδαράκη*.—Sandaraque, Gomme de genévrier, FR.; Sandarach, Wachholderharz, GERM.; Resina de enebro, Grasiilla, Sandaraca, SPAN.; Geneverharst, DUT.

AMST.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; HAMB.; HANN.; OLDEN.; BORUS.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A hard resin in small tears or transparent fragments, of a pale yellow or lemon co-

lour, becoming reddish with time. Its fracture is bright and smooth; its odour and taste resemble those of common resin.

BROUSONNET says that it flows from the *Thuya articulata*, DESF. (*Monæcia Monadelphica*, LINN.; *Conifera*, JUSS.; *Fig.* Vahl. *Symb.* 2. p. 96. t. 18), a tree of Barbary.

It was for a long time attributed to the *Juniperus communis*, LINN.; to the *J. Lycia*, LINN.; or, to the *J. oxycedrus*; the first of these sometimes presents between the wood and bark a very different resin of a greenish colour, called *German Sandarac*, *Sandaraca Germanica*.

It is excitant.

SANGUINARIA CANADENSIS, LINN.—BLOODROOT.

SYNONIMES :—Pacoon, Indian paint, Turmeric.—Sanguinaire du Canada, FR.

AMER.; COXE.

A plant of North America. (*Polyandria Monogynia*, LINN.; *Pavaveracea*, JUSS.; *Fig.* Big. *Med. bot.* t. 7)

The root and seeds are employed.

The root is thick and furnished in every part with a great many fibres. It yields a juice as red as blood. Its taste is acrid and bitterish. It is emetic.—Dose, from eight to twenty grains of the powder.—SMITH says that its virtues are analogous to those of the *Digitalis purpurea*, and that it also diminishes the frequency of the pulse.

The seeds, according to BARTON, have the same mode of action as those of *Datura stramonium*.

INFUSION OF BLOODROOT.

Infusum Sanguinariæ Canadensis. (NIEMANN)

R. Root of Bloodroot, one dram.
Boiling Water, four ounces.

Infuse, and strain.

It is recommended as emetic by TULLY.—Dose, from two drams to half an ounce, till the effect is produced.

TINCTURE OF BLOODROOT

Tinctura Sanguinariæ Canadensis. (AMER.; COXE; NIEMANN)

R. Root of Bloodroot, two ounces.
Dilute Alcohol, one pint.

Digest for ten days, and filter.

It is recommended in incipient phthisis, hydrothorax, at the commencement of croup, &c.—Dose, from ten to twenty drops or more, gradually augmented.

SANGUISORBA OFFICINALIS, LINN.—GREAT BURNET.

SYNONIMES :—Sanguisorbe, Pimprenelle des près, FR.; Blutkraut, Blutwurz, Wiesenknopf, GERM.; Pimpinelle, DAN.; Pimpinella, SPAN.; Blædkruid, DUT.;

Pimpinella maggiore, ITAL.; Sanguisorba, PORT.; Blodört, SWED.

BRUNS.; GALL.; WIRTEM.; MURRAY and GMELIN.

A plant of Europe. (*Tetrandria Monogynia*, LINN.; *Rosaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 184*)

The root (*Radix Sanguisorbæ* seu *Pimpinellæ Naticæ*) is employed. It is branch-

ing, of a reddish-brown-yellow externally, and of a yellowish-white internally, inodorous, and of a styptic taste.

It was formerly esteemed vulnerary.

SANICULA EUROPÆA, LINN.—SANICLE.

SYNONIMES :—Sanicle, FR.; Sanickel, GERM.; Sanikel, DAN.; DUT.; and SWED.; Sanicula, PORT.; and SPAN.; Sanicola, ITAL.; Zankiel, POL.

BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

A plant of Europe. (*Pentandria Digynia*, LINN.; *Umbellifera*, JUSS.; *Fig. Zorn, Ic. pl. t. 109*)

The herb (*Herba Saniculæ* seu *Diapensiæ*) is employed. It consists of simple radical leaves, round, five-lobed, serrated, shining, and having long foot-stalks, of slender stems, and of sessile, trifid caulinary leaves. It is destitute of odour, but has a bitter, sub-astringent, and saline taste.

It enters into several pectoral preparations.

SANTALINA CHAMÆCYPARISSIAS, LINN.—SWEET-SCENTED LAVENDER COTTON.

SYNONIMES :—Santoline à feuilles de cyprès, Aurone femelle, Garde-robe, FR.; Gartency pressen, Cypressenkraut, GERM.; Santolina, ITAL.; PORT.; and SPAN.; Cypreskruid, DUT.

GALL.; FERR.; WIRTEM.; BERGIUS.; BRUGNATELLI.; MURRAY and GMELIN.; SPIELMANN.

A plant of the mountains of the south of

Europe. (*Syngenesia Polygamia Æqualis*, LINN.; *Fig. Zorn, Ic. pl. t. 245*)

The herb (*Herba Santolinæ* seu *Abrotani feminei* seu *Abrotani montani*) is employed. It is an evergreen and is composed of den- tated, obtusely-edged, grey leaves disposed in four rows. It has a penetrating and agreeable odour, and an aromatic and bitter taste.

It is excitant, nervine, and anthelmintic.

SAPONARIA OFFICINALIS, LINN.—SOAPWORT.

SYNONIMES :—Saponaire, FR.; Seifenkraut, Speichelseifenkraut, Waschkraut, Speichelwurz, GERM.; Mydlj koren, BOHEM.; Sæbeurt, DAN.; Jabonera, SPAN.; Zeepkruid, DUT.; Saponaria, ITAL.; Mydelnik, POL.; Saboaira, PORT.; Sæpært, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; GUIBOURT.; MUR-

RAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A common plant of Europe. (*Hexandria Digynia*, LINN.; *Caryophyllææ*, JUSS.; *Fig. Flore Medic. VI. 311*)

The root (*Radix Saponariæ rubræ*) is slender, hard, very long, of the diameter of a goose-quill, nodose, of a reddish-brown externally and whitish internally, wrinkled, and nearly inodorous. Its taste is at first sweetish and mucilaginous, but soon becomes bitter, austere, and sub-acrid.

The herb consists of a glabrous, fistulous,

cylindrical, articulated stem, rather branching, and furnished with glabrous, very smooth, somewhat pointed leaves of a deep-green colour, nearly sessile, oval, lanceolated, very entire, presenting three nervures, and narrowed at the base. It is perfectly inodorous, but has a mucilaginous and bitterish taste.

Soapwort contains resin, gum, and a mucilaginous principle called *Saponine* by BUCHOLZ.

It is tonic and sudorific, and is principally employed in cutaneous and syphilitic diseases.

JUICE OF SOAPWORT.

Succus Saponariae. (HASS.)

R. Fresh Root or Herb of Soapwort, any quantity.
Water, four parts.

Bruise in a stone mortar, and express. Set apart the juice, and filter it.

Dose, from one to four ounces.

EXTRACT OF SOAPWORT.

Extractum Saponariae. (ANT.; AUSTR.; GENEV.; HANN.; SAX.; BRUGNATELLI; SWEDIAUR; VAN MONS)

R. Herb of Soapwort, ten pounds.
Bruise it in an iron mortar, adding

Water, six pounds.

Express the juice, boil it for an instant, and evaporate it slowly to the consistence of thick syrup. (SWEDIAUR)

R. Fresh Leaves of Soapwort, any quantity.

Bruise it with a little water, and strain the juice through a sieve; set aside, decant, coagulate it at the fire, and strain again: then having evaporated it to the consistence of pilular mass, remove it from the fire, incorporate the fecula, set apart, and evaporate finally to the proper consistence. (VAN MONS)

R. Fresh Root of Soapwort, any quantity.

Macerate for several hours in water; then add more water, boil for a few moments, clarify with white of egg, and evaporate to the consistence of extract. (VAN MONS)

R. Herb of Soapwort, one part.
Water, eight parts.

Infuse for twenty-four hours, then boil for a quarter of an hour, and strain with strong expression; boil the residue with

Water, four parts.

Mix the liquors, set apart for twenty-four hours, and evaporate to the required consistence. (SAX.)

R. Fresh Leaves of Soapwort, } of each,
Cold Water, } one part.

Macerate for twelve hours, and add,

Boiling Water, four parts.

Strain when cold, clarify with white of egg, set the liquor upon the fire, and when it begins to thicken, strain it through a sieve, and evaporate, constantly stirring. (VAN MONS)

R. Fresh Leaves of Soapwort, two pounds.
Water, ten pounds.

Boil for a quarter of an hour, and express; set aside and decant the liquor, and evaporate to the proper consistence. (ANT.; GENEV.)

LIPP. directs two pounds of root to be boiled in eight of water till reduced to four, and the decoction to be evaporated.

R. Root of Soapwort, one part.
Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and express; boil the residue with

Water, four parts.

Mix the strained liquors, decant after sufficient repose, and evaporate to the consistence of an extract. (HANN.)

AUST. directs the root to be exhausted by several successive ebullitions, and the mixed decoctions to be decanted and evaporated.—BRUGNATELLI directs it to be boiled for two hours with six parts of water, and a second time with fresh water; the mixed liquors to be then evaporated.

Dose, from half a dram to two drams.

DECOCTION OF SOAPWORT.

Decoctum Saponariae officinalis. (NIEMANN; SWEDIAUR; PH. SYPH.)

R. Root of Soapwort, two ounces.
Spring Water, four pounds.

Boil down to one half. (NIEMANN)

R. Herb of Soapwort, half a pound.
Water, eight pounds.

Boil down to one half. (SWEDIAUR; PH. SYPH.)

It is recommended in scrofula, syphilitic diseases, and cutaneous affections.—Dose, from two to four pounds daily.

RESOLVENT SPECIES. (FULD.; SARD.; FIDERIT)

R. Root of Dandelion, } of each,
—— Soapwort, } two ounces.
Herb of Lesser Centaury, } of each,
—— Horehound, } one ounce.
Flowers of Tansey, } of each,
—— Chamomile, } one ounce
and a half.

Cut, and mix. (FULD.; FIDERIT)

- R. Root of Dandelion, } of each,
 — Soapwort, } two parts.
 — Valerian, }
 Herb of Lesser Centaury, } of each,
 — White Horehound, } one part.
 Flowers of Chamomile, }
 two parts and a half.

Linseed, * four parts.

Cut, and bruise. (SARD.)

For lavements, in cases of obstructions of the abdominal viscera.

SATUREJA.—SAVORY.

Three species of this genus of plants are employed in medicine :

1. *Satureja Thymbra*, LINN.—Cretan Savory.

SYNONIME :—Sarriette de Crète, FR.

HISP. ; GALL.

A plant of Crete. (*Didymia Gymnospermia*, LINN. ; *Labiata*, JUSS.)

The herb has small, oval, pointed leaves.

2. *Satureja hortensis*, LINN.—Garden Savory.

SYNONIMES :—Sarriette des jardins, FR. ; Gartensaturei, Pfefferkraut, GERM. ; Satureye, BOHEM. ; Saer, DAN. ; Ajedrea, SPAN. ; Keulen, DUT. ; Santoreggia, ITAL. ; Ozabi, POL. ; Seguerelha, PORT. ; Kyndel, SWED.

AMST. ; AUSTR. ; BATAV. ; BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; HERBIP. ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A plant of the south of Europe. (Fig. Zorn, *Ic. pl. t. 216*)

The herb (*Herba Saturejæ seu Cunilæ sativæ*) consists of a very branching, square,

slender stem, and of opposite, sessile, simple, lanceolated, linear leaves. It has an agreeable and strongly aromatic odour, and a sub-acrid, warm, and aromatic taste.

3. *Satureja capitata*, LINN.—Cretan Thyme.

SYNONIME :—Thym de Crète, FR.

BRUNS. ; HISP. ; GALL. ; WIRTEM. ; BERGIUS ; SPIELMANN.

A plant of the south and east of Europe.

The flowering herb (*Herba et Spicæ Thymi Cretici*) is employed. It consists of a slender stem, of small, narrow, opposite, careened, pointed, ciliated leaves, and of white flowers disposed in a terminal spike. It has a strong and agreeable odour, and a warm and aromatic taste.

All these plants are excitant.

OIL OF SAVORY.

Oleum Saturejæ æthereum, Ætheroleum Saturejæ. (PALAT.)

- R. Flowering Herb of Savory, twenty-five parts.
 Water, seventy-five parts.
 Muriate of Soda, three parts.

Digest for three days, and distil.

SAXIFRAGA GRANULATA, LINN.—COMMON SAXIFRAGE.

SYNONIMES :—Saxifrage blanche, FR. ; Weisser Steinbrech, Hundsrebe, Keilant, GERM. ; Huid steenbræk, DAN. ; Witte Steenbreck, DUT. ; Saxifraga branca, PORT. ; Steenbræcke, SWED.

BRUNS. ; HISP. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

This plant is spread over every part of Europe. (*Hexandria Digynia*, LINN. ; *Saxifragæ*, JUSS. ; Fig. Zorn, *Ic. pl. t. 309*)

The root and herb are employed.

The root (*Radix Saxifragæ albæ*) is composed of small round tubercles of the size of coriander seeds, united together by fibrils, and covered with a red pellicle. They were formerly, improperly, designated *Semen Saxifragæ albæ*. It has a somewhat styptic and bitter taste, which disappears after desiccation.

The herb consists of a branching stem, and of reniform, lobed leaves. It has an acrid and sub-acid taste, which also is nearly quite dissipated by drying.

SCABIOSA.

Two species of this genus are employed in medicine :

1. *Scabiosa arvensis*, LINN.—Field Scabious.

SYNONIMES :—Scabieuse des champs, FR. ; Ackersscabiæse, GERM. ; Kawias, BOHEM. ; Scaburt, DAN. ; Escabiosa, SPAN. ; Gemeen scharfkruid, DUT. ; Scabiosa, ITAL. ; Dryakiew polne, POL. ; Escabiosa dos campos, PORT. ; Akervædd, SWED.

AMST. ; AUSTR. ; BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

A common plant of Europe. (*Tetrandria Monogynia*, LINN. ; *Dipsacæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 141*)

The root, herb, and flowers are employed.

The root is long, slender, white, of a scarcely sensible odour, and of a bitter and styptic taste.

The herb consists of a round, hairy, hollow stem, and of leaves, the inferior of which are oval, lanceolate and dentate, and the superior sub-pinnate.

The flowers are of a light purple colour.

WATER OF SCABIOUS.

Aqua Scabiosæ. (PALAT.)

R. Herb of Scabious, one part.

Water, three parts.

Distil one half.

2. *Scabiosa succisa*, LINN.—Devil's-bit.

SYNONIMES :—Mors du diable, FR. ; Teufels-abbiss, GERM. ; Czertkuss, BOHEM. ; Dievelsbid, DAN. ; Mordisco de diablo, SPAN. ; Duivelsbeet, DUT. ; Morso del diavolo, ITAL. ; Cisyna, POL. ; Morso del diabo, PORT. ; Engvædd, SWED.

ANT. ; BRUNS. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

A common plant of Europe. (*Fig. Zorn, Ic. pl. t. 150*)

The root and herb are employed.

The root (*Radix Succisæ seu Morsus diaboli*) is short, fibrous, yellowish without, whitish within, and as if gnawed off at the middle, on account of part of it dying at the end of every year.

The herb consists of a simple, round, hairy, branching stem, having opposite branches, furnished at each articulation with two small, opposite, lanceolate, entire leaves.

It has a scarcely-sensible odour, and a bitterish, sweet, and styptic taste.

It is slightly astringent, but is seldom used.

SCAMMONIUM.—SCAMMONY.

SYNONIMES :—*Diacrydium* ; σχαμμωνία.—Scammonée, FR. ; Scammonien, Purgierender windensafft, GERM. ; Skammonia, BOHEM. ; Sukmunia, ARAB. ; and DUK. ; Skammonium, DAN. ; and SWED. ; Escamonea, PORT. ; and SPAN. ; Mehmedeh, HIND. ; Scammoneum, DUT. ; Scamonea, ITAL. ; Skammonia, Sok suznico, POL.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; HANN. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; AINSLIE ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY

and GMELIN ; PARIS ; PIDERIT ; SAUNDERS ; SPIELMANN ; ZARDA.

There are three sorts of this gum-resin.

1. Aleppo Scammony ; *Scammonium de Aleppo seu Aleppense seu Halepense seu Syriacum*.—This is in irregular masses of an ash-grey colour, light, dry, and very friable. Its fracture is bright ; its powder of a brownish-white ; its odour somewhat nauseant, and its taste acrid and bitter.

It is the inspissated milky juice of the *Convolvulus Scammonia*, LINN., a plant of Syria. (*Pentandria Monogynia*, LINN. ; *Convolvulaceæ*, JUSS. ; *Fig. Zorn, Ic. pl. t. 214*)

2. Smyrna Scammony ; *Scammoneum*

Smyrnum.—This is more compact and heavy, less friable, of a deeper colour and more disagreeable odour than the preceding; it is also of less value.

It is obtained from the *Periploca Scammonaea*, LINN., a plant of Egypt. (*Pentandria Monogynia*, LINN.; *Apocynæ*, JUSS.; *Fig. Alp. de pl. Æg. t. 48*)

3. Montpellier Scammony.—This is entirely black, very hard and compact.

It is furnished by the *Cynanchum Mons-peliacum*, LINN., a shrub of the south of France and Spain. (*Pentandria Monogynia*, LINN.; *Apocynæ*, JUSS.; *Fig. Cav. Ic. rar. I. tab. 60*)

It is a drastic purgative.—Dose of the powder, from three to fifteen grains.

SULPHURETTED SCAMMONY.

Diacrydium sulphuratum. (PALAT.; SARD.; WIRTEM.)

R. Scammony, in powder, any quantity.

Spread it upon white paper, and keep this over a chafing-dish containing burning charcoal; throw sulphur upon the coals in small portions, and stir the scammony till it begins to melt and adhere to the paper.

GLYCYRRHIZED SCAMMONY.

Diacrydium glycyrrhizatum. (SARD.; WIRTEM.)

R. Scammony, half a pound.
Decoction of Liquorice, one pound.

Triturate together, set aside, evaporate the liquor to dryness, and pulverize the residuum.

The *Rosated Scammony*, *Diacrydium rosatum*, is obtained by substituting the infusion of roses for the decoction of liquorice, and the *Cydoniated Scammony*, *Diacrydium cydoniatum*, by using the depurated juice of quince.

RESIN OF SCAMMONY.

Resina Scammonii. (AMST.; ANT.; BRUNS.; GALL.; FULD.; PALAT.; ROSS.; SAX.; WIRTEM.; BRUGNATELLI)

R. Scammony, in powder, one part.
Alcohol (0.907), six parts.

Dissolve, filter, and distil off one part; precipitate the resin of the residue by adding water; wash it well and dry it with a very gentle heat. (AMST.; ANT.; BRUNS.; GALL.; ROSS.; SAX.; WIRTEM.; BRUGNATELLI)

PALAT. directs the tincture to be distilled nearly to dryness, and the residue to

be washed, dried, and pulverized.—Dose, from three to ten grains.

COMPOUND POWDER OF SCAMMONY.

Pulvis Scammonii compositus seu catholicus seu caryocostinus seu diacarthami. (LOND.; EDINB.; AMER.; ANT.; HISP.; FERR.; LUSIT.; SARD.; COXE; VAN MONS)

Pulvis Scammonæ compositus. (LOND.)

R. Scammony,	} of each,	two ounces.
Hard Extract of		
Jalap,		
Ginger,	half an ounce.	

Reduce them separately to powder, and mix.

Pulvis Scammonii compositus. (EDINB.)

R. Scammony,	} of each, an	equal part.
Supertartrate of		
Potass,		

Pulverize.

AMER.; LUSIT.; and COXE give the same formula as EDINB.—FERR. prescribes eight grains each, of scammony, of the supertartrate, and of sugar.

R. Scammony,	half an ounce.
Indian Jalap,	} of each, six drams.
Jalap,	
Supertartrate of Potass,	three drams.

Mix. (HISP.)

R. Scammony,	} of each, two ounces.
Hermodyctyls,	
Cloves,	} of each, one ounce.
Cinnamon,	
Ginger,	

Make a powder. (ANT.)

R. Cinnamon,	} of each, one part.
Arabian Costus,	
Ginger,	
Hermodyctyls,	
Scammony,	two parts.

Make a powder. (VAN MONS)

R. Seeds of Safflower,	six drams.
Gum Tragacanth,	} of each,
Hermodyctyls,	
Scammony,	half an ounce.
Cinnamon,	} of each, two drams.
Liquorice,	
Turbith,	
	one ounce.

Make a powder. (SARD.)

Dose, from ten to fifteen grains.

ANTI-ARTHRITIC POWDER.

Pulvis antiarthriticus purgans. (CADET DE GASSICOURT; SPIELMANN)

- | | |
|------------------------------|--|
| R. Seeds of Blessed Thistle, | } of each,
half
an ounce. |
| Safflower, | |
| Scammony, | |
| Sarsaparilla, | |
| China Root, | |
| Guaiac Wood, | } of each,
one ounce.
two drams. |
| Supertartrate of Potass, | |
| Senna, | |
| Cinnamon, | |

Make a powder. (SPIELMANN)

CADET DE GASSICOURT prescribes ten drams each, of the two seeds, half an ounce each, of the supertartrate and of senna, two drams each, of scammony, of sarsaparilla, of china root, and of guaiac, and one dram of cinnamon.

Dose, one dram.

RESOLVENT PILLS. (CADET DE GASSICOURT)

- | | |
|------------------------|----------------|
| R. Extract of Gentian, | half an ounce. |
| Ox-gall, | three drams. |
| Scammony, | two drams. |

Make one hundred and sixty-two pills.—Dose, from four to six daily, fasting or before dinner.

PURGATIVE PILLS.

Pilule Extracti catholici. (AMST.)

- | | |
|------------------------------|--------------|
| R. Aqueous Extract of Aloes, | one ounce. |
| Extract of Black Hellebore, | two drams. |
| Colocynth, | one dram. |
| Scammony, | three drams. |

Make a pilular mass.

PURGATIVE LOZENGES.

Tabellæ diacarthami. (GALL.; SARD.; PIERQUIN)

- | | |
|---|------------------------|
| R. Diacarthami Powder (form. of SARD.), | three ounces. |
| Sugar, | one pound. |
| Mucilage of Tragacanth, | a sufficient quantity. |

Make lozenges.

- | | |
|-----------------------------|------------------------|
| R. Supertartrate of Potass, | half an ounce. |
| Scammony, | three drams. |
| Sugar, | eight ounces. |
| Oil of Cinnamon, | four drops. |
| Rose Water, | a sufficient quantity. |

Make lozenges of one dram each. (PIERQUIN)

- | | |
|-----------------------|------------------------|
| R. Scammony, | three drams. |
| Senna, | four drams and a half. |
| Rhubarb, | one dram and a half. |
| Preserved Lemon Peel, | one ounce. |

VOL. II.

Sugar, six ounces six drams;
Mucilage of Tragacanth made with
Cinnamon Water,
a sufficient quantity.

Make tablets of six drams each. (GALL.)

Dose, from two to six drams, or more.

CARYOCOSTINE ELECTUARY. (VAN MONS)

R. Compound Powder of Scammony,

Honey,	one part. six parts.
--------	-------------------------

Mix.—Dose, as a purgative, from one to six drams.

HYDRAGOGUE ELECTUARY. (CADET DE GASSICOURT; PLANCHE)

- | | |
|---------------------|------------------------|
| R. Scammony, | } of each, two drams. |
| Root of Jalap, | |
| Squill, | one dram and a half. |
| Resin of Jalap, | half a dram. |
| Syrup of Buckthorn, | a sufficient quantity. |

Mix.—Dose, to twenty-four grains, in asthenic dropsy.

COMPOUND ELECTUARY OF SCAMMONY.

Electuarium Scammonii seu purgans, Confectio Scammoneæ. (LOND.; DUBL.; AMER.; COXE; RATIER; SPIELMANN; VAN MONS)

Electuarium Scammonææ. (LOND.)

Electuarium Scammonii. (DUBL.)

- | | |
|-----------------|------------------------|
| R. Scammony, | one ounce and a half. |
| Cloves, | } of each, six drams. |
| Ginger, | |
| Oil of Caraway, | half a dram. |
| Syrup of Roses, | a sufficient quantity. |

Triturate the dry ingredients to a fine powder, then add the syrup, and triturate again: finally incorporate the oil.

- | | |
|-----------------------|--------------------------|
| R. Scammony, | } of each,
one ounce. |
| Ginger, | |
| Oil of Cloves, | one scruple. |
| Syrup of Orange Peel, | a sufficient quantity. |

Mix. (AMER.; COXE)

- | | |
|-----------------|--------------------------|
| R. Coriander, | two drams. |
| Cloves, | } of each,
six drams. |
| Ginger, | |
| Scammony, | one ounce and a half. |
| Syrup of Roses, | a sufficient quantity. |

Mix. (VAN MONS)

- R. Scammony, half an ounce.
 Jalap, four ounces.
 Sweet Almonds, twenty.
 Senna, two ounces.
 Anise, } of each,
 Supertartrate of Potass, } one ounce.
 Pulp of Prunes, sixteen ounces.
 ——— Tamarinds, eight ounces.

Mix. (SPIELMANN)

Dose, one or two drams.

DIAPHGENIC ELECTUARY.

Electuarium Diaphgenicum. (GALL.;
 SARD.)

- R. Pulp of Dates, one pound.
 Turbith, four ounces.
 Liquorice, two ounces.
 Scammony, one ounce and a half.
 Purified Honey, two pounds and a half.

Boil the pulp with the honey to the consumption of the superfluous humidity, and incorporate the powders. (SARD.)

- R. Pulp of Dates, half a pound.
 Sweet Almonds, three ounces and a half.
 Sugar, half a pound.
 Despumated Honey, two pounds.
 Ginger, }
 Pepper, } of each,
 Mace, } two drams.
 Cinnamon, }
 Rue, }
 Seeds of Cretan Carrot, }
 ——— Fennel, }
 Root of Turbith, four ounces.
 Saffron, six grains.
 Scammony, one ounce and a half.

Mix intimately. (GALL.)

Dose, from half a dram to half an ounce.

TINCTURE OF SCAMMONY.

Tinctura Scammonii. (GALL.)

- R. Scammony, one part.
 Alcohol (22 degrees), four parts.

Digest for three days, and strain.—Dose, from twenty drops to one dram.

POLYCHRESTIC ELIXIR.

Elixir polychrestum Halense. (WIRTEM.;
 SPIELMANN)

- R. Compound Extract of Colocynth, one ounce.
 Alcohol, six ounces.
 Dissolve, and add
 Sulphuric Acid, three drams.
 Filter. (WIRTEM.; SPIELMANN)

- R. Compound Extract of Colocynth, one ounce.
 Simple Mixture, six ounces.
 Dissolve. (WIRTEM.)

It is purgative. The formula was for a long time kept secret; it is by TRILLER.—Dose, from thirty to sixty drops, or more.

PURGATIVE OIL.

Oleum purgans. (VAN MONS)

- R. Scammony, four scruples.
 Oil of Almonds, four ounces.
 Dissolve in a moderate heat.

It is mildly purgative.—Dose, from half an ounce to one ounce and a half, in emulsion, or with syrup of poppy.

DRASTIC SYRUP. (BORIES)

- R. Syrup of Lemon, five ounces.
 Scammony, in powder, four drams.
 Subcarbonate of Iron, } of each,
 Sulphuret of Antimony, } two drams.

Mix.

SYRUP OF SCAMMONY.

Syrupus de Scammonio; Sirop anti-arthritique, Sirop anti-hydropique. (GALL.;
 CADET DE GASSICOURT; VAN MONS)

- R. Scammony, in powder, half an ounce.
 White Sugar, four ounces.
 Alcohol (22 degrees), eight ounces.

Set the mixture over a fire in a silver pan, and when it has become warm, inflame it with a taper; stir incessantly with a spatula till the flame is extinguished; then, having removed the vessel from the fire, set the liquor aside to cool; strain it through flannel, and add

Syrup of Violets, four ounces.

Mix. (GALL.)

- R. Scammony, in powder, one part.
 Proof Spirit, sixteen parts.

Dissolve; inflame the liquor, and leave the flame to become extinguished of itself: then strain through flannel, and add

Syrup of Violets, eight parts.

Mix. (VAN MONS)

- R. Gum Arabic, one part.
 Scammony, two parts.

Triturate together, and with

Water, a sufficient quantity,
 make a thick emulsion. Incorporate this with

Syrup of Violets, twenty parts.

Mix. (VAN MONS)

PURGATIVE POTION.

Mixture cathartique Arabe. (CADET DE GASSICOURT)

- R. Orange-flower Water, } of each,
Syrup of Peach Flowers, } one ounce.
Spirit of Rosemary, } one dram.
Scammony, } twelve grains.

Mix.

- R. Syrup of Lemon, } one ounce.
Sulphuret of Antimony, } twelve grains.
Scammony, } one scruple.

Mix.—For one dose.—The second formula has been particularly extolled in dropsy.

PURGATIVE EMULSION. (GALL.)

- R. Scammony, } twelve grains.
White Sugar, } two drams.
Yolk of Egg, } one half.

Triturate together, and add gradually

- Simple Emulsion, } five ounces.
Orange-flower Water, } two drams.

Mix.

VERMIFUGE EMULSION. (SAUNDERS)

- R. Peach Kernels, } of each, two drams.
Bitter Almonds, }
Scammony, } half a scruple.
Water of Wormwood, } three ounces.

Mix.

SCILLA MARITIMA, LINN.—SQUILL, SEA ONION.

SYNONIMES:—Scille, Ognon marin, FR.; Meerzwiebel, Mauszwiebel, GERM.; Alaschil, ARAB.; Merska cybule, BOHEM.; Strandløg, DAN.; Escilla, Cebolla albarena, SPAN.; Zeaujuin, DUT.; Scilla, Cipolla marina, ITAL.; Korzen cebuli morskiey, POL.; Albarra, Cebolla albarra, PORT.; Sjelœk, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUST.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

A plant of the south of Europe, growing about the sea shore. (*Hexandria Monogynia*, LINN.; *Asphodeleæ*, JUSS.; *Fig. Flore Medic. VI. 318*)

The bulb (*Radix Scillæ marinæ* seu *Squillæ* seu *Scillæ rubræ* seu *Pancratii veri*) is employed. It is pyriform, large, and covered with several dry, thin, papyraceous tunics; under which are oval, fleshy, red or white scales. Its odour is pungent; and its taste, at first mucilaginous, is bitter, acrid, and nauseant. The red variety is only employed.

Squill contains, according to VOGEL, besides gum, woody fibre, &c. a peculiar bitter principle, *Scillitimum*, which is the source of its activity. According to TILLY, this principle is merely a mixture of uncrystallizable sugar, and of two matters, the

one very bitter, and the other excessively acrid.

In the East Indies, AINSLIE says that the bulb of the *Erythronium Indicum*, ROTTLER, is substituted for squill.

SYNONIMES:—Unsul, Iskil, ARAB.; Nurriala, CYNG.; Jungliepiaz, DUK.; Kanda, HIND.; Peyaz ideschti, PERS.; Nurri vungyum, TAM.; Addivitella guddalu, TEL.

Some authors have said that squill grows in Ceylon; but WHITE, of Bombay, thinks the bulb of the *Amaryllis Zeylanica*, has been mistaken for it.

It is excitant, diuretic, pectoral, and is prescribed in dropsies and chronic catarrhs. —Dose, from one to ten grains of the powder.

I. Preparations which contain Squill in Substance.

PULP OF SQUILL.

Pulpa Scillæ. (GALL.)

- R. Bulbs of Squill, any quantity.

Separate the roots and external envelopes and enclose the rest in paper. Roast them in warm ashes till the bulbs are easily bruised with the fingers; then remove the burnt parts, and make a pulp of the rest by pounding it in a mortar and passing it through a sieve.

The bulb may also be enclosed in a thin coat of paste, and then roasted in ashes or baked in an oven.

TROCHISCS OF SQUILL.

Trochisci de Scilla Damocratis seu scillitici.
(HISP.; PALAT.; SARD.; WIRTEM.; SPIELMANN)

R. Squill, twelve ounces.
Bruise it in a mortar, and add
Lupin Meal, eight ounces.

Divide the mass into trochiscs of two drams each. (PALAT.; WIRTEM.; SPIELMANN)

HISP.; and SARD. use the powdered root of *fraxinella* instead of the meal.

COMPOUND POWDER OF SQUILL.

Pulvis Scillæ compositus. (BRUNS.; MIL. COPEN.; FENN.; FULD.; GENEV.; PALAT.; CAST. BORUS.; SARD.; SUEC.; WIRTEM.; AUGUSTIN.; CADET DE GASSICOURT.; ELLIS.; NIEMANN.; PIDERIT.; SAUNDERS.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

R. Squill, six parts.
Root of Swallow-wort, one part.
Make a powder.

CADET DE GASSICOURT.; SWEDIAUR.; BRUNS.; PALAT.; SARD.; WIRTEM.; and SPIELMANN prescribe three parts of squill and one part of swallow-wort.—NIEMANN—two of squill and one of swallow-wort.

R. Squill, one part.
Root of Swallow-wort, six parts.
Nitrate of Potass, eight parts.

Make a powder. (PIDERIT)

R. Squill, one part.
Nitrate of Potass, four parts.

Mix. (SAINTE-MARIE)

ELLIS prescribes one part of squill and six parts of nitrate.—GENEV.—one of squill and eight of nitrate.

R. Squill, one dram.
Nitrate of Potass, two drams.
Cinnamon, one scruple.

Mix. (HERBIP.)

SAUNDERS prescribes ten grains of squill, fifteen of nitrate, and five of pimento.—SWEDIAUR—ten of squill, ten of pimento, and fifteen of nitrate.

R. Squill, one part.
Nitrate of Potass, } of each,
Supertartrate of Potass, } six parts.

Mix. (PIERQUIN)

In another place, PIERQUIN prescribes six parts of squill, twelve of nitrate, and twenty of supertartrate.

R. Squill, one part.
Nitrate of Potass, three parts.
Supertartrate of Potass, four parts.
Aromatic Powder, two parts.

Mix. (FENN.; SUEC.)

R. Squill, one part.
Supertartrate of Potass, twenty-five parts.
Oleosaccharum of Fennel, five parts.

Mix. (MIL. COPEN.)

R. Squill, } of each, twelve grains.
Ginger, }
Sulphate of Potass, two drams.

Mix. (FULD.)

R. Squill, one part.
Aromatic Powder, two parts.
Nitrate of Potass, three parts.
Sulphate of Potass, four parts.

Mix. (VAN MONS)

R. Acetate of Soda, } of each,
Oleosaccharum of } three drams.
Juniper, }
Squill, six grains.
Root of Sweet Flag, half a grain.
Opium, three grains.

Mix, and divide into six powders. (CAST. BORUS.)

R. Squill, two grains.
Leaf of Digitalis, one grain.
Supertartrate of Potass, half a dram.

Mix. (NIEMANN)

R. Squill, } of each, one grain.
Digitalis, }
Oil of Juniper, two drops.
Borated Tartar, } of each, one scruple.
Liquorice, }
Cinnamon, two grains.

Mix. (NIEMANN)

It is diuretic.—Dose, from ten to twenty grains.

R. Squill, two grains.
Juniper, eighteen grains.

Mix.—For one dose. (AUGUSTIN)

R. Squill, fifteen grains.
Florentine Iris, } of each, ten grains.
Liquorice, }
Sugar of Milk, two scruples.
Oil of Anise, ten drops.

Make three doses; one, night and morning. (AUGUSTIN)

R. Acetate of Soda, } of each,
Oleosaccharum of } three drams.
Juniper, }
Squill, six grains.
Sweet Flag, half a dram.
Opium, three grains.

Mix, and divide into six doses; one, three times a day in infusion of juniper. (AUGUSTIN)

BOLUSES OF SQUILL.

Bols contre l'œdème du poudon. (PIERQUIN.; RATIER)

R. Squill, six grains.

Sulphate of Potass, two grains.
Oxymel of Squill, a sufficient quantity.

Make one bolus. (RATIER)

- R. Root of Swallow-wort, half an ounce.
Squill, two drams.
Subcarbonate of Potass, one dram.
Syrup of Valerian, a sufficient quantity.
Make twenty-four boluses. (PIERQUIN)

PILLS OF SQUILL.

Pilule Scilliticæ. (LOND.; DUBL.; AMER.; AMST.; ANT.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; LUSIT.; CAST. BORUS.; SARD.; WIRTEM.; HERBIP.; AUGUSTIN.; BRUGNATELLI.; COXE.; CADET DE GASSICOURT.; HUFELAND.; NIEMANN.; PIDERIT.; PIERQUIN.; SPIELMANN.; SWEDIAUR.; VAN MONS.)

1. Without Soap.

- R. Squill, one ounce and a half.
Gum Ammoniac, half an ounce.
Oxymel of Squill, a sufficient quantity.
Make a pilular mass. (FORM. HÔP. FR.; GALL.)

FERR. prescribes an equal part of each, of squill and of gum, with a sufficient quantity of copaiba.

- R. Squill, half a dram.
Gum Ammoniac, three drams.
Nitrate of Potass, one dram.
Simple Syrup, a sufficient quantity.

Make pills of four grains each. (GENEV.)

- R. Gum Ammoniac, } of each, two parts.
Squill, }
Lesser Cardamoms, one part.
Oxymel of Squill, a sufficient quantity.

Make a pilular mass. (AMST.)

FULD.; and LIPP. prescribe an equal part each, of the three substances, with oxymel of squill.—LUSIT.; and CADET DE GASSICOURT—equal parts, with simple syrup.—HASS.; and PIDERIT—equal parts, without syrup or balsam.—HERBIP.; and PIERQUIN—equal parts, with copaiba.—AUGUSTIN.—two parts each, of squill and of gum, and one of cardamoms, without syrup, oxymel, or balsam.

- R. Squill, one part.
Gum Ammoniac, } of each,
Lesser Cardamoms, } three parts.
Extract of Liquorice, }
Simple Syrup, a sufficient quantity.

Make a pilular mass. (ANT.; BRUGNATELLI.; SWEDIAUR.)

- R. Squill, } of each,
Gum Ammoniac, } eight parts.
Wood-lice, }
Lesser Cardamoms, two parts.

Copaiba, one part.
Make a pilular mass. (BRUNS.)

2. With Soap.

Pilule Scillæ compositæ. (LOND.; DUBL.)

- R. Fresh Squill, dried, one dram.
Ginger, } of each,
Hard Soap, } three drams.
Gum Ammoniac, two drams.

Pulverize, mix, and add as much syrup as may be sufficient to form a pilular mass.

Pilule Scilliticæ. (EDINB.)

- R. Squill, dried, one scruple.
Gum Ammoniac, }
Cardamoms, } of each,
Extract of Li- } one dram.
quorice, }

Make a mass with syrup.

COXE prescribes one dram of squill, two drams of ginger, ten drops of oil of anise, and a sufficient quantity of jelly of soap.—BRUGNATELLI—one dram of squill, three each of ginger and of soap, two of gum ammoniac, and a sufficient quantity of syrup of ginger.—CADET DE GASSICOURT—three drams each of soap, and of ginger, two drams of gum ammoniac, one ounce of squill, and a sufficient quantity of simple syrup.—HANN.—one dram of squill, three drams each of ginger and of soap, two drams of gum ammoniac, and a sufficient quantity of mucilage of gum arabic.

- R. Squill, } of each,
Gum Ammoniac, } half an ounce.
Wood-lice, }
Soap, one ounce.
Copaiba, one dram.

Make pills of ten grains each. (SARD.; HERBIP.; HUFELAND.; NIEMANN)

- R. Venice Soap, two drams.
Gum Ammoniac, } of each, one dram.
Squill, }
Wood-lice, }
Oil of Anise, seven drops.
Honey of Roses, a sufficient quantity.

Make pills of one grain each. (NIEMANN)

BELG.; PIERQUIN.; and SPIELMANN give the same formula, but substitute the oxymel of Squill for the copaiba.

- R. White Soap, } of each, two drams.
Gum Ammoniac, }
Ginger, two scruples.
Squill, twelve grains.
Alcohol, a sufficient quantity.

Make pills of two grains each. (CAST. BORUS.; AUGUSTIN)

- R. Spanish Soap, eight parts.
 Gum Ammoniac, } of each,
 Root of Swallow-wort, } half
 Squill, } an ounce.
 Copaiba, one dram.

Make a pilular mass. (DAN.)

- R. Soap, two parts.
 Squill, } of each, one part.
 Nitre, }
 Copaiba, a sufficient quantity.

Make pills of four grains each. (CADET DE GASSICOURT)

- R. White Soap, two parts.
 Gum Ammoniac, } of each, one part.
 Squill, }
 Thick Emulsion of Copaiba, a sufficient quantity.

Make a pilular mass. (VAN MONS)

- R. Soap, seven parts.
 Gum Arabic, three parts.
 Squill, } of each, one part.
 Copaiba, }

Make a pilular mass. (HISP.)

- R. Squill, one dram.
 Soap, twenty-four grains.

Make forty pills. (AMER.; COXE)

The dose of these pills is, in general, from six to fifteen grains.

RESOLVENT PILLS.

Pilulæ deobstruentes seu diuretica resolventes. (BRERA; CADET DE GASSICOURT; SWEDIAUR)

- R. Gum Ammoniac, } of each,
 Soap, } one dram and a half.
 Squill, } of each,
 Aromatic Powder, } half a dram.
 Oil of Juniper, twenty drops.
 Syrup of Orange Peel, a sufficient quantity.

Make pills of five grains each.—Dose, four, every three days; in jaundice and chronic diseases of the liver.

- R. Soap, } of each,
 Extract of Dandelion, } one dram.
 Sulphate of Soda, four scruples.
 Squill, sixteen grains.
 Honey, a sufficient quantity.

Make pills of four grains each.—Dose, one every three hours. (BRERA)

PECTORAL PILLS.

Pilulæ pectorales. (VAN MONS)

- R. Gum Ammoniac, } of each,
 Sulphate of Potass, } six drams.
 Florentine Iris, half an ounce.
 Squill, two drams.

Soap, two ounces.
 Extract of Red Poppy, a sufficient quantity.

Make pills of five grains each.—Dose, five, thrice a day.

PHYSAGOGUE PILLS.

Pilulæ physagoga. (HERBIP.; NIEMANN; SAINTE-MARIE; SWEDIAUR)

- R. Squill, } of each,
 Extract of Elaterium, } half an ounce.
 Succinic Acid, one dram.
 Red Sulphuret of Antimony, half a dram.

Make pills of three grains each. (SWEDIAUR)

- R. Squill, half an ounce.
 Red Sulphuret of } of each,
 Antimony, } one dram.
 Succinic Acid, }
 Extract of Elaterium, half an ounce.
 Oil of Anise, a few drops.
 Extract of Liquorice, a sufficient quantity.

Make one hundred and fifty pills. (HERBIP.; NIEMANN; SAINTE-MARIE)

HYDRAGOGUE PILLS. (CADET DE GASSICOURT; PIDERIT; PIERQUIN)

- R. Extract of Colocynth, five drams.
 Squill, five drams and a half.

Make two hundred pills. (PIERQUIN)

- R. Squill, } of each,
 Proto-chloruret of } two drams and
 Mercury, } a half.
 Oxymel of Squill, a sufficient quantity.

Make one hundred pills. (PIERQUIN)

- R. Squill, } of each,
 Extract of } one dram and
 Buckbean, } a half.
 Digitalis, } of each, one dram.
 Assafoetida, }

Make one hundred pills. (CADET DE GASSICOURT)

- R. Extract of Dandelion, three drams and a half.
 Spanish Soap, one dram and a half.
 Galbanum, two drams.
 Squill, twenty-four grains.
 Proto-chloruret of Mercury, sixteen grains.
 Camphor, one scruple.
 Tincture of Orange, a sufficient quantity.

Make pills of two grains each. (PIDERIT)
 Dose, four, morning and evening.

- R. Assafoetida, two drams.
 Soap, one dram.
 Camphor, half a dram.
 Squill, twelve grains.

Oil of Juniper, twenty drops.

Make pills of two grains each.—Dose, ten, morning and evening. (AUGUSTIN)

R. Assafoetida, } of each,
 Extract of Valerian, } two drams.
 Squill, }
 Opium, } of each, ten grains.

Make pills of two grains each.—The same dose as for the preceding. (AUGUSTIN)

DIURETIC PILLS. (CADET DE GASSICOURT)

R. Supertartrate of Potass, four drams.
 Gum Ammoniac, two drams.
 Squill, one dram and a half.
 Saffron, two scruples.
 Oxymel of Squill, a sufficient quantity.

Make sixty pills. (SAINTE-MARIE)

R. Squill, twenty grains.
 Digitalis, eighteen grains.
 Rob of Juniper, a sufficient quantity.

Make ten pills.—Dose, one every two hours. (BRERA)

R. Wood-lice, in powder, two drams.
 Squill, thirty-six grains.
 Gum Ammoniac, three drams.
 Myrrh, one dram.
 Rob of Juniper, four drams.

Make pills of three grains each. (SAINTE-MARIE)

EXPECTORANT PILLS. (ELLIS; PARIS)

R. Myrrh, one dram and a half.
 Squill, half a dram.
 Extract of Henbane, two scruples.
 Water, a sufficient quantity.

Make thirty pills.—Dose, two, morning and evening.

INCISIVE PILLS. (CADET DE GASSICOURT; PIERQUIN)

R. Squill, one dram.
 Red Sulphuret of Antimony, eight grains.
 Aqueous Extract of Opium, six grains.
 Cocoa-nut Oil, a sufficient quantity.

Make thirty-six pills. (CADET DE GASSICOURT)

R. Squill, half an ounce.
 Ipecacuanha, two drams.
 Aqueous Extract of Opium, fifteen grains.
 Cocoa-nut Oil, one ounce.
 Syrup of Gum, a sufficient quantity.

Make pills of four grains each. (CADET DE GASSICOURT)

R. Squill, six grains.
 Amber, fifteen grains.
 Ipecacuanha, two grains.

Conserve of Elecampane, a sufficient quantity.

Make nine pills. (PIERQUIN)

Dose, two or three, thrice a day.

SQUILL LOZENGES. (PIERQUIN)

R. Squill, half a dram.
 Sugar, two ounces.
 Mucilage of Tragacanth, a sufficient quantity.

Make one hundred and twenty lozenges.

DEOBSTRUENT ELECTUARY.

Electuarium deobstruens. (SWEDIAUR)

R. Compound Powder of Squill } of each,
 (Form. of SWEDIAUR), } two
 Gum Ammoniac, } drams.
 Tartrate of Potass, }
 Muriate of Ammonia, one scruple.
 Syrup of Cinnamon, a sufficient quantity.

Mix.—It is prescribed in dropsy.—Dose, two drams, thrice a day.

ANTI-ASTHMATIC LOHOCH.

Lohoch ad Asthma seu de Scilla. (BRUNS; PALAT.; WIRTEM.; PIERQUIN; SPIELMANN)

R. Juice of Squill, } of each,
 Despumated Honey, } four ounces.
 Mix. (PIERQUIN)

R. Squill, in powder, eight grains.
 Gum Ammoniac, twenty grains.
 Syrup of Marshmallows, } of each,
 Oxymel of Squill, } one ounce.
 Borage Water, two ounces.

Mix. (PIERQUIN)

R. Squill, half an ounce.
 Florentine Iris, two drams.
 Horehound, } of each, one dram.
 Hyssop, }
 Myrrh, } of each, half a dram.
 Saffron, }
 Despumated Honey, six ounces.

Mix. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

OINTMENT OF SQUILL.

Unguentum Scillæ. (HUFELAND; NIEMANN)

R. Squill, three ounces.
 Caustic Ley, a sufficient quantity.
 Boil to a mucilage, and add,
 Lard, a sufficient quantity.

Make an ointment.

It is resolvent, and is employed in frictions to indolent tumours.

DIURETIC OINTMENT.

Unguentum diureticum resolvens. (BRERA)

R. Mercurial Ointment, one dram.
 Squill, half a dram.

Mix.—It is recommended to be rubbed upon the inner part of the thighs, upon the lumbar region, and upon the abdomen; in dropsical affections.

DIURETIC LINIMENT.

Linimentum diureticum gastro-scilliticum. (BRERA)

R. Squill, in powder, one dram.
 Calf's Gastric Juice, two ounces.

Mix.—To be rubbed upon the loins in dropsy, when the patient cannot take medicine by the mouth.

II. Preparations which only contain the active principle of Squill, in a more or less pure state.

A.

Extraction by Water.

EXTRACT OF SQUILL.

Extractum Scilla. (AUSTR.; BAVAR.; BELG.; FERR.; HERBIP.; BRUGNATELLI; VAN MONS)

R. Fresh Squill, any quantity.

Bruise then in a mortar, express the juice, and evaporate it, first over a slow fire, and then in a water-bath. (AUSTR.)

BELG. directs the squill to be pounded with a sufficient quantity of water to obtain a nearly liquid mass; this to be expressed; the residue to be triturated with a little water, and expressed again: the liquors to be mixed, set apart, decanted, and evaporated in a water-bath.—BRUGNATELLI—the squill to be pounded, mixed with water, and the juice expressed; the residue to be boiled with water; the two strained liquors to be then mixed and evaporated.

R. Squill, any quantity.

Boil for an hour in a sufficient quantity of water, and express. Having set apart and decanted the liquor, evaporate it to the consistence of thick honey. (FERR.; HERBIP.)

R. Fresh Squill, one part.
 Water, four parts.

Macerate for twenty-four hours, and strain without expression. Evaporate the liquor to the proper consistence. (BAVAR.; VAN MONS)

DIURETIC ELECTUARY.

Electuarium resolvens martiale diureticum. (BRERA)

R. Venice Soap, one dram.
 Extract of Hemlock, twelve grains.
 — Squill, one scruple.
 Deutoxide of Iron, one dram.
 Oxymel, a sufficient quantity.

Mix.—To be taken gradually.

DECOCTION OF SQUILL.

Decoctum Scilla. (AMER.; COXE; NIEMANN)

R. Squill, three drams.
 Juniper, four ounces.
 Seneka, three ounces.
 Water, four pounds.

Boil till reduced to one half. Strain, and add

Nitric Æther, four ounces.

Mix.—It is excitant, and is recommended in dropsies.—Dose, from half a spoonful to one spoonful.

HYDRAGOGUE DECOCTION. (SAUNDERS)

R. Dandelion, entire, two ounces.
 Root of Marshmallows, ten drams.
 Water, a sufficient quantity.

Boil for a quarter of an hour, and add,

Squill, one ounce and a half.
 Chamomile Flowers, two ounces.

Digest for an hour in a covered vessel. Add to fourteen ounces of the strained liquor,

Syrup of Liquorice, two ounces.

Mix.

PECTORAL AND INCISIVE APOZEM. (BORRIES)

R. Root of Dandelion, } of each,
 Flowers of Dandelion, } two ounces.
 Root of Marshmallows, one ounce.
 Flowers of Chamomile, three drams.
 Squill, one dram and a half.
 Extract of Liquorice, one ounce.
 Water, a sufficient quantity

to obtain three pounds of strained liquor. Add to this

Oxymel of Squill, } of each,
 Syrup of Marshmallows, } six drams.

Mix.—It is pectoral, and diuretic.—A glassful to be taken every hour.

DIURETIC DECOCTION.

Decoctum diureticum Hufelandi. (CADET DE GASSICOURT; VAN MONS)

R. Squill, half a dram.
 Water, six ounces.

Boil for an instant, remove it from the fire, and add

Root of Valerian, two drams.

Strain with slight expression. When the liquor is cold emulsionize it with

Gum Arabic, } of each, two drams.
Guaiac Resin, }

Strain again, and add

Alcoholic Solution of } of each,
Potass, }
Spirit of Nitric Æther, } two drams.

Mix. (VAN MONS)

CADET DE GASSICOURT directs half an ounce of squill to be boiled for twenty minutes in ten ounces of water; two drams of valerian root in powder to be added towards the end of the boiling: the liquor to be strained and mixed with one ounce of syrup of orange peel, three drams of mucilage of gum arabic, two drams each, of guaiac resin, of alcoholic solution of potass, and of alcoholized nitric acid, and twenty drops of tincture of opium.

Dose, a spoonful every two hours.

HONEY OF SQUILL.

Mel scilliticum. (GALL.; VAN MONS)

R. Squill, two ounces.
Water, three pounds.

Boil slightly for a few minutes; then digest for two days. Express, and add to the liquor,

Honey, one pound and a half.

Clarify, and boil to the consistence of syrup. (GALL.)

R. Dried Squill, one part.
Water, a sufficient quantity

to obtain twelve parts of infusion. Strain, and add

White Sugar, twelve parts.

Boil a little, strain through flannel, and, when cold, add

Honey, twelve parts.

Mix. (VAN MONS)

Dose, from half a dram to two drams.

AQUEOUS SYRUP OF SQUILL.

Syrupus de Scilla seu scilliticus seu Scilla.
(VAN MONS)

R. Dried Squill, one part.
Warm Water, twelve parts.

Infuse for twelve hours, and express. Add

White Sugar, sixteen parts.

Clarify, and boil to a syrup.

PECTORAL LOHOCH. (RATIER)

R. Simple Lohoch, four ounces.
VOL. II.

Honey of Squill, one ounce.

Mix.

DIURETIC POTION. (RATIER)

R. Water of Valerian, four ounces.
Peppermint, one ounce.
Nitric Æther, one dram.
Tincture of Opium, half a dram.
Honey of Squill, four drams.
Compound Syrup of Smalage, one ounce.

Mix.—This potion has been framed from HUFELAND's diuretic decoction.

COMPOUND HONEY OF SQUILL.

Mel Scilla compositum, Syrupus Scilla compositus. (AMER.; COXE; ELLIS; NIEMANN)

R. Dried Squill, bruised, } of each,
Root of Seneka, } four ounces.
Water, four pounds.

Boil slowly till reduced to one half, and strain. Add

Despumated Honey, two pounds.

Boil again till the liquid is reduced to three pounds. Dissolve in four ounces of the syrup,

Tartarized Antimony, one grain.

Mix.—It is excitant, resolvent, and diuretic, and is extolled in mucous asthma.—

Dose, from ten drops to a tea spoonful or more, every quarter of an hour, half-hour, or hour, according to the age of the patient or the indication.

COMPOUND SYRUP OF SQUILL. (FENN.; OLDEN.; SUEC.; SPIELMANN; VAN MONS)

R. Dried Squill, one ounce.
Ginger, half an ounce.
Dried Herb of Hyssop, two ounces.
Peppermint Water, two pounds.

Macerate for twenty-four hours in a covered vessel; then express, and to the liquor, depurated by repose, add

White Sugar, three pounds.

Boil a little. (FENN.; SUEC.; and SPIELMANN)

OLDEN. substitutes hyssop water for that of peppermint.

VAN MONS prescribes one part of ginger, two parts of squill, four of hyssop, twenty of water, infusion in a water-bath, and thirty-six parts of sugar.

B.

EXTRACTION BY WINE.

Wine of Squill.

Vinum Scilla seu Scilla maritimæ seu scilliticum. (AMST.; GALL.; FORM. HÔP. 4 D

FR. ; FULD. ; HAMB. ; LIPP. ; LUSIT. ;
SARD. ; SUEC. ; HERBIP. ; AUGUSTIN ;
CADET DE GASSICOURT ; NIEMANN ;
PIERQUIN ; SAUNDERS ; VAN MONS)

- R. Dried Squill, one ounce.
Malaga Wine, one pound.
Macerate for twelve hours, strain, and
filter. (GALL.)

LUSIT. prescribes one ounce of squill and
two pounds of wine.—SARD.—half an ounce
of squill and two pounds of wine.

- R. Squill, one dram.
Orange Peel, one ounce.
White Wine, two pounds.

Digest for twenty-four hours, and strain.
(HAMB.)

- R. Fresh Squill, six drams.
Cinnamon, three drams.
Wine, one pound and a half.

Infuse without heat for twelve hours, and
strain. (HERBIP.)

VAN MONS prescribes one part of squill,
one half part of cinnamon, forty-eight parts
of Madeira wine, and eight days' maceration.

- R. Fresh Squill, half an ounce.
Ginger, one dram.
Alcohol (0.931), two ounces.
White Wine, two pounds.

Strain at the expiration of three days.
(AMST.)

SUEC. prescribes one ounce of squill, one
dram of ginger, and two pounds of white
wine.—AUGUSTIN—one dram of squill, two
drams of sweet flag, and one pound of wine.

- R. Squill, one ounce.
Orange Peel, } of each,
Root of Sweet Flag, } two drams.
White Wine, one pound and a half.

Digest for three days, and add
Oxymel of Squill, two drams.

Mix. (CADET DE GASSICOURT ; PIER-
QUIN)

- R. Dried Squill, } of each, one dram.
Juniper, }
Root of Sweet Flag, } two drams.
White Wine, one pound.

Macerate for three days, and strain.
(FULD. ; NIEMANN)

LIPP. prescribes one ounce of white wine,
five grains of squill, two scruples of juni-
per, and ten grains of sweet flag.

- R. Squill, one dram.
Juniper, one ounce.
Lesser Cardamoms, half a dram.
White Wine, half a pound.

Digest for three days, in a moderate heat.
(NIEMANN)

- R. Squill, } of each,
Lesser Cardamoms, } one part.

Juniper, } of each, two parts.
Seeds of Carrot, }
Madeira Wine, thirty-two parts.

Macerate for several days without heat,
express lightly, and filter. (VAN MONS)

- R. Tincture of Squill, twenty grains.
White Wine, one ounce.

Mix. (FORM. HÔP. FR.)

It is excitant, sometimes nauseant, and
even emetic.—Dose, two or three spoonfuls
daily.

BITTER DIURETIC WINE.

*Vinum diureticum amarum seu amarum
scilliticum compositum, Infusum scilliti-
cum vinosum compositum.* (GALL. ; CA-
DET DE GASSICOURT ; RATIER ; SAUN-
DERS)

- R. Squill, six drams.
Cinnamon, } of each,
WINTER'S Bark, } four drams.
Root of Elecampane, one dram.
Wine, one pound.

Strain after sufficient extraction. (SAUN-
DERS)

- R. Squill, } of each,
Root of Swallow-wort, } half an ounce.
Cinchona, } of each,
WINTER'S Bark, } two ounces.
Lemon Peel, }
Stems of Bohemian Angelica,

half an ounce.
Leaves of Wormwood, } of each,
Balm, } one ounce.

Juniper, } of each, half an ounce.
Mace, }
White Wine, eight pounds.

Digest for four days at a moderate tem-
perature ; then strain, express, and filter.
(GALL.)

- R. Dried Squill, } of each,
Root of Swallow-wort, } one ounce.
Angelica, }
Cinchona, in powder, } of each,
WINTER'S Bark, } two ounces.
Lemon Peel, }
Juniper, } of each, two drams.
Mace, }
Dried Leaves of Worm- } of each,
wood, } half a dram.
Balm, }
Alcohol, two ounces.
White Wine, four pounds.

Digest in a sand-bath for twenty-four
hours, occasionally stirring ; then express
and filter. (CADET DE GASSICOURT)

- R. WINTER'S Bark, } of each, one ounce.
Cinchona, }
Cinnamon, }
Angelica Root, } of each,
Squill, } two ounces.
Juniper, }
Mace, }

Dried Leaves of Worm-wood, } of each,
Balm, } two handfuls.
White Wine, two pounds.
Macerate for twenty-four hours, and strain. (RATIER)

It is tonic, and is recommended in cases where the stomach requires to be stimulated, and in passive dropsical affections.—Dose, from one to four ounces daily.

ALKALINE WINE OF SQUILL.

Infusum Kali carbonici cum Scilla vinosum.
(AUGUSTIN)

R. Subcarbonate of Potass, half an ounce.
Squill, one scruple.
Cinnamon, two drams.
White Wine, eight ounces.

Infuse for twelve hours, and strain.—Dose, one or two spoonfuls every two or three hours.

C.

Extraction by Alcohol.

TINCTURE OF SQUILL.

Tinctura Scillæ seu Scillæ maritimæ.
(LOND.; DUBL.; EDINB.; ANT.; DAN.; GALL.; FORM. HÔP. FR.; HANN.; COXE; NIEMANN; VAN MONS)

Tinctura Scillæ.

R. Fresh Squill, dried, four ounces.
Proof Spirit, two pints.

Macerate for fourteen days, and filter. (LOND.)

DUBL. gives the same proportions, but requires only seven days' maceration; the liquor is then to be set apart that the feculencies may subside, and the clear tincture poured off.

R. Fresh Squill, dried, two ounces.
Proof Spirit, sixteen ounces.

Digest for seven days, and filter. (EDINB.)

FORM. HÔP. FR.; and GALL. prescribe one part of dried squill and four parts of alcohol (22 degrees); to be digested for six days.—NIEMANN gives the same formula as LOND.—VAN MONS prescribes one part of squill and six parts of proof spirit.—DAN.—six ounces of squill and two pounds of proof spirit.—HANN.—five ounces of squill and two pounds of proof spirit.—COXE—five ounces of squill and two pints of alcohol (0,930).

R. Fresh Squill, bruised, six ounces.
Alcohol (36 degrees), eight ounces.

Digest for four days; then express, and pour upon the residue

Alcohol (36 degrees), two ounces.

Digest again at a moderate heat, and strain.

Mix and filter the tinctures. (ANT.)

It is excitant and diuretic, and is principally prescribed in dropsical affections.—Dose, from ten to twenty drops, in emulsion or mucilaginous water.

ANTI-CATARRRHAL MIXTURE. (CADET DE GASSICOURT)

R. Paregoric Elixir, five ounces.
Tincture of Squill, one ounce.

Mix.—Dose, a tea spoonful, morning and evening.

DIURETIC MIXTURE. (ELLIS)

R. Tincture of Squill, two drams.
Nitric Æther, one ounce.

Mix.—Dose, a tea spoonful five or six times a day, in an infusion of juniper.

AUGUSTIN gives an analogous preparation under the very improper title of *Tinctura Scillæ*, of which the following is the formula:

R. Squill, one ounce and a half.
Muriatic Æther, } of each,
Spirit of Juniper, } half a pound.
Water, a sufficient quantity.

Distil six pounds.—Dose, from ten to thirty drops.

AROMATIC TINCTURE OF SQUILL. (SUEC.)

R. Dried Squill, one ounce.
Cinnamon, in powder, two drams.
Diluted Alcohol, half a pound.

Infuse for three days, and filter.

The cinnamon may be replaced by zest of orange peel. The tincture may also be prepared extemporaneously by adding tincture of orange peel to the simple tincture of squill. The aromatic tincture deserves to be preferred to the simple, as it is less disagreeable to the palate, and perhaps less irritating.

COMPOUND TINCTURE OF SQUILL.
(VAN MONS)

R. Vinegar of Squill, three parts.
Tincture of Orange Peel, one part and a half.

Extract of Buckbean, one part.

After several days' maceration, strain through flannel.

ALKALINE TINCTURE OF SQUILL.

Tinctura Scille kalina. (FENN.; FERR.; HASS.; LIPP.; WIRTEM.; HUFELAND; VAN MONS)

R. Squill, in powder, } of each,
Subcarbonate of Potass, } one ounce.

Triturate together in a mortar, and add

Alcohol, four ounces.

Digest for six hours, and strain with slight expression. (HASS.)

HUFELAND; and NIEMANN prescribe three ounces of squill, two drams of caustic potass, and one pound of alcohol; to be digested for three days.—LIPP.—one ounce each, of squill and of subcarbonate of potass, and eight ounces of alcohol.

R. Dried Squill, one ounce.
Cinnamon, two drams.
Solution of Potass, one dram.
Alcohol, half a pound.

Digest for three days, and filter. (FENN.)

VAN MONS prescribes one part of caustic potass, two parts of cinnamon, eight of squill, and forty-eight of proof spirit.

R. Subcarbonate of Potass, half an ounce.
Vinegar of Squill, twelve ounces.

Evaporate over a slow fire to the consistency of honey; then digest the residue with

Alcohol, six ounces,

Filter at the expiration of several days. (WIRTEM.)

VAN MONS prescribes eighteen ounces of vinegar and twelve of alcohol.

R. Squill, } of each,
Acetate of Potass, } one ounce.
Alcohol, six ounces.

Filter at the end of three days. (FERR.)

It is excitant, and one of the most active resolvents and diuretics. It is recommended in dropsies and asthma. The commencing dose should be thirty drops, and it should be augmented by five every two or three hours till the patient takes a hundred or more.

DIURETIC POTION. (ELLIS)

R. Subcarbonate of Potass, one dram.
Lemon Juice, a sufficient quantity
to dissolve the alkali. Add

Tincture of Squill, half an ounce.

Opium, thirty drops.

Cinnamon Water, four ounces.

Mix.—Dose, a spoonful every two or three hours.

WEDEL'S PECTORAL ELIXIR.

Elixir pectorale Wedelii. (BRUNS.; LIPP.; WIRTEM.; CADET DE GASSICOURT; HUFELAND; NIEMANN; SPIELMANN; VAN MONS)

R. Squill, } of each,
Root of Florentine Iris, } one ounce.
Elecampane, }
Liquorice, half an ounce.
Red Sanders, one ounce.
Herb of Rosa Solis, one ounce and a half.

Seeds of Anise, half an ounce.

Benzoin, six drams.

Myrrh, half an ounce.

Gum Ammoniac, two drams.

Saffron, three drams.

Proof Spirit, two pounds.

Digest the gums separately, and after sufficient extraction, mix the tincture with that of the other substances. (BRUNS.; WIRTEM.)

NIEMANN omits the rosa solis.—SPIELMANN adds one ounce of marshmallows root, and reduces the benzoin to four drams.

R. Squill, } of each,
Root of Iris, } one ounce.
Elecampane, }

Benzoin, }

Myrrh, }

Anise, }

Extract of Liquorice, half an ounce.

Gum Ammoniac, }

Saffron, three drams.

Alcohol, one pound.

Digest and strain. (HUFELAND)

The following formula is for the *Elixir pectorale Anglais* of CADET DE GASSICOURT.

R. Squill, } of each,
Root of Elecampane, } three ounces
Iris, } one dram.
Benzoin, two drams.

Liquorice, }

Anise, }

Myrrh, }

Gum Ammoniac, two scruples.

Saffron, seventeen grains.

Alcohol (12 degrees),

one pound six ounces.

Macerate for fifteen days, occasionally agitating, and filter.

R. Saffron, two scruples.
Proof Spirit, thirty-two ounces.

Infuse for two days, and add

Root of Liquorice, }

Seeds of Anise, } of each,
Root of Elecampane, } half an ounce.

Iris, }

Gum Ammoniac, one ounce.

Myrrh, two drams.

half an ounce.

Infuse again for two days, and add

Squill, one ounce.
Benzoin, three drams.

Digest, and filter. (VAN MONS)

R. Root of Elecampane, } of each,
Iris, } half an ounce.
Squill, }
Liquorice, } of each,
Seeds of Anise, } two drams.
Flowers of Red Poppy, }
Alcohol, fourteen ounces.

Digest in a moderate heat for several days. On the other hand, digest in three ounces of alcohol, for the same length of time,

Gum Ammoniac, one dram.
Myrrh, two drams.
Benzoin, three drams.

Strain with slight expression, and mix. (LIPP.)

It is excitant, and is recommended in asthma, catarrhs, cough, and hoarseness. —Dose, from forty to sixty drops.

D.

Extraction by Vinegar.

VINEGAR OF SQUILL.

Acetum Scillæ seu scilliticum, Acidum acetum scilliticum seu cum Scilla maritima. (LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI.; COXE.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

R. Squill, one part.
Vinegar, twelve parts.

Digest at a moderate heat for eight days; then express strongly, and filter. (DAN.; HANN.; OLDEN.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; BRUGNATELLI.; SPIELMANN)

BRUNS. prescribes one part and a half of squill, and twelve parts of vinegar; to be infused for forty-eight hours.—PALAT.—one of squill, and nine of vinegar; to be infused for forty days.—MIL. COPEN.; SARD.; and WIRTEM.—one of squill, and eight of vinegar; to be digested for fifteen days.—AUSTR.; HASS.; LIPP.; ROSS.; and PIDERIT.—one of squill, and six of vinegar; to be digested for three days.

R. Squill, one part.
Boiling Vinegar, eleven parts.
Acetic Acid, one part.

Macerate for eight days; then express, set aside, and decant. (FENN.)

FORM. HÔP. FR. prescribes eight parts of squill, ninety-two of vinegar, one of acetic acid (10 degrees), and fifteen days' infusion.

Acetum Scillæ.

R. Fresh Squill, dried, one pound.
Dilute Acetic Acid, six pints.
Proof Spirit, half a pint.

Macerate the squill in the acid at a gentle heat in a covered vessel for twenty-four hours; then express the liquor, set it apart that the feculencies may subside, and add the spirit to the decanted liquor. (LOND.)

R. Fresh Squill, dried, half a pound.
Vinegar from Wine, three pounds.
Rectified Spirit, four fluid ounces.

Digest the squill with the vinegar for four days, frequently agitating; then express, and when the feculencies have subsided, add the spirit. (DUBL.)

Acidum aceticum scilliticum. (EDINB.)

R. Dried Squill, one ounce.
Distilled Vinegar, fifteen ounces.
Alcohol, one ounce and a half.

Macerate the squill with the acid for seven days; then express the liquor, and add the alcohol to it. When the feculencies have subsided, pour off the clear part.

R. Squill, sliced, two ounces.
Vinegar, twelve ounces.

Digest for eight days at a moderate heat in a covered vessel; then express, and add to the liquor

Alcohol, one ounce.

Set aside, decant, and preserve. (BATAV.; BAVAR.; BELG.)

AMST. prescribes half a pound of squill, four pounds of vinegar, twenty-four hours' maceration in a gentle heat, and the addition of one twelfth of alcohol (0.907) to the strained liquor.—HERBIP.—half a pound of

squill, four pounds of vinegar, four days' digestion, and three ounces of alcohol.—**LUSIT.**—one pound of squill, six pounds of vinegar, six days' maceration, and then a slight boiling, and the addition of half a pound of alcohol.—**HISP.**—one pound of squill, six pounds of vinegar, twenty-four hours' digestion at a moderate heat, and three ounces of alcohol.—**GALL.**—half a pound of dried squill, six pounds of vinegar, one ounce of alcohol (22 degrees), and fifteen days' maceration.—**POL.**—one ounce of squill, one pound of vinegar, and one ounce and a half of alcohol.—**SWEDIAUR.**—one pound of squill, eight pounds of vinegar, two or three days' digestion at a moderate heat, and eight ounces of proof spirit.—**VAN MONS.**—one part of squill, four parts of vinegar, and two of proof spirit; or, one of squill, twelve of vinegar, and one of proof spirit; to be macerated without heat for eight days.—**AMER.**; and **COXE.**—half a pound of squill, three pints of vinegar, and four ounces of alcohol (0.840).—**GENEV.**—four ounces of squill, two pounds of vinegar, and three ounces of proof spirit.—**FULD.**—four ounces of squill, four pounds of vinegar, and two ounces of proof spirit.—**FERR.**—one pound of squill, six pounds of vinegar, and three ounces of alcohol.

R. Fresh Squill, one pound.
Vinegar, concentrated by freezing,
one pound and a half.

Digest for a week in a covered vessel, frequently agitating; then express, and pour upon the residue,

Vinegar, concentrated by freezing,
half a pound.

Express, set aside the mixed liquors, filter, and add

Alcohol (38 degrees), two ounces.

Mix. (ANT.)

It is excitant, and is principally prescribed in chronic affections of the mucous membrane of the lungs, and in hydropical affections. It is also administered in the form of gargle in inflammation of the tonsils, and in certain cases of deafness.—Dose, from twenty drops to one dram, thrice a day, in some aromatic water, or in emulsion of gum ammoniac.

ACETOSE SYRUP OF SQUILL.

Oxysaccharum scilliticum. (EDINB.; AMER.; BRUNS.; HANN.; HERBIP.; WURCEB.; BRUGNATELLI; COXE; NIEMANN)

Syrupus Scillæ maritimæ. (EDINB.; AMER.; COXE)

R. Vinegar of Squill, four parts.
Refined Sugar, seven parts.

Dissolve the sugar with a gentle heat so as to form a syrup.

NIEMANN prescribes two pounds of vinegar of squill, and three pounds of sugar.—**HERBIP.**—one and a half of vinegar, and three and a half of sugar.—**BRUGNATELLI.**—two of vinegar, and three and a half of sugar.

R. Vinegar of Squill, one pound and a half.
Cinnamon, }
Ginger, } of each, one ounce.

Digest for three days; then strain, and add

White Sugar, thirty ounces.

Boil for a few moments, and strain. (BRUNS.; HANN.; WIRTEM.)

R. Vinegar of Squill, half a pound.
Syrup of Cinnamon, }
Ginger, } of each, one pound.

Mix. (SWEDIAUR)

EXPECTORANT POTION. (ELLIS)

R. Syrup of Squill, half an ounce.
Compound Tincture of Camphor, two drams.
Antimonial Wine, one dram.
Gum Arabic, half a dram.
Water, two ounces.

Mix.—Dose, a tea spoonful every hour or two hours. This potion is intended for children.

OXYMEL OF SQUILL.

Mel Scillæ acetatum, Oxymel scilliticum, Mel Aceti Scillæ seu Scillæ maritimæ. (LOND.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; COXE; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Oxymel Scillæ. (LOND.; DUBL.)

R. Despumated Honey, three pounds.
Vinegar of Squill, two pints.

Boil in a glass vessel, over a slow fire, to a proper consistence.

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; PIDERIT; SPIELMANN; and SWEDIAUR pre-

scribe one part of vinegar of squill, and two parts of honey.—PALAT.; and POL.—two of the vinegar, and three of honey.—AMER.; and COXE—the same as LOND.—MIL. COPE.—an equal part of each.—DAN.—thirty ounces of honey, reduced to six or eight, and twelve ounces of the vinegar.—GENEV.—one part of the vinegar, and four parts of honey.

The following formula is taken from the *Journal de Pharmacie*:

R. Dried Squill, one pound two ounces.
Vinegar, six pounds.
Alcohol, two ounces.

Macerate for a month, and strain and filter. Incorporate, in a water-bath, four pounds of honey with one pound one ounce of this scillitic vinegar.

It is prescribed by spoonfuls, but more frequently in potions, in the dose of from one to four drams.

PECTORAL LOHOC

Linctus expectorans. (LIPP.; PIERQUIN; SWEDIAUR)

R. Oxy mel of Squill,
from half an ounce to one ounce.
Syrup of Marshmallows, one ounce.

Mix. (SPIELMANN)

R. Oxy mel of Squill, } of each,
Syrup of Marshmallows, } an equal
Mucilage of Gum Arabic, } part.

Mix. (LIPP.)

R. Oxy mel of Squill, half an ounce.
Syrup of Tolu, one ounce.
Oil of Almonds, one ounce and a half.

Mix. (SWEDIAUR)

Dose, a tea spoonful.

SCILLITIC GARGLE.

Gargarisma e Scilla. (SWEDIAUR)

R. Oxy mel of Squill, one ounce.
Cinnamon Water, ten ounces.

ANTI-CATARRHAL POTION. (SAINTE-MARIE)

R. Root of Zedoary, one dram.
Camphorosma, one pinch.
Water, eight ounces.

Boil down to five ounces, and add to the cold liquor

Muriate of Ammonia, six grains.
Spirit of Nitric Ether, six drops.
Oxy mel of Squill, one ounce.

Mix.—A fourth part to be taken every two hours.

DIURETIC POTION. (FORM. HÔP. FR.; BRERA; CADET DE GASSICOURT; RATIER; SAUNDERS)

R. Oxy mel of Squill, half an ounce.
Mint Water, one ounce.
Water of the three Nuts, four ounces.

Mix. (CADET DE GASSICOURT)

R. Oxy mel of Squill, half an ounce.
Infusion of Hyssop, one ounce.
Alcoholized Nitric Acid, four ounces.

Mix. (FORM. HÔP. FR.)

R. Oxy mel of Squill, one dram and a half.
Cinnamon Water, six drams.
Compound Spirit of Lavender, one dram.

Subcarbonate of Potass, ten grains.

Mix. (SAUNDERS)

R. Compound Decoction of Cichory, three ounces.

Orange-flower Water, half an ounce.
Oxy mel of Squill, two drams.
Simple Syrup, one ounce.

Tincture of Opium } of each,
Castor, } six drops.
Musk, }

Spirit of Nitric Ether, } of each,
Sulphuric Ether, } ten drops.

Mix. (RATIER)

R. Juniper, half an ounce.
Boiling Water, a sufficient quantity

to obtain eight ounces of infusion. Add when strained

Sulphuric Ether, twenty drops.
Oxy mel of Squill, half an ounce.

Mix. (BRERA)

R. Juniper, one ounce.
Water, a sufficient quantity

to obtain six ounces of infusion. Add to the strained liquor

Acetate of Potass, two drams.
Spirit of Soot, thirty drops.
Succinated Spirit of Ammonia, two drams.

Oxy mel of Squill, one dram.

Mix. (BRERA)

R. Emulsion of Bitter Almonds, one pound.

Nitrate of Potass, two drams.
Oxy mel of Squill, one ounce.

Mix. (BRERA)

R. Decoction of Dog-grass, two pounds.
HOFFMANN'S LIQUOR, twenty drops.
Oxy mel of Squill, one ounce.

Mix. (BRERA)

To be taken by spoonfuls.

DIURETIC BROTH. (PIERQUIN)

- R. Root of Dandelion, one ounce.
 ——— Liquorice, half an ounce.
 Woodlice, thirty.
 Leaves of Dandelion, half a handful.
 ——— Seneca, } of each,
 Camphorosma, } one pinch.
 Oxymel of Squill, half an ounce.
 Red Sulphuret of Antimony, one grain.

PECTORAL MIXTURE. (BRERA; CADET DE GASSICOURT; SAUNDERS)

- R. Oxymel of Squill, } of each,
 Spirit of Pimenta, } one ounce.
 Mint Water, two ounces.

Mix. (SAUNDERS)

- R. Decoction of Barley, half a pound.
 Hyssop Water, four ounces.
 Oxymel of Squill, three ounces.
 Vinegar of Squill, six drams.
 Sulphate of Soda, one dram.

Mix. (CADET DE GASSICOURT)

- R. Gum Ammoniac, one dram.
 Vinegar of Squill, six ounces.
 Oxymel of Squill, one ounce.

Mix. (CADET DE GASSICOURT)

- R. Root of Seneca, three drams.
 Water, a sufficient quantity
 to obtain six ounces of decoction. Strain,
 and add

- Ammoniacal Emulsion, one dram.
 Camphor, half a dram.
 Oxymel of Squill, one ounce.

Mix. (BRERA)

Dose, one ounce every half hour.

SCILLITIC JULEP. (RATIER)

- R. Pectoral Species, } of each,
 Gum Arabic, } twenty grains.
 Simple Syrup, } of each,
 Oxymel of Squill, } one ounce.
 Water, four ounces.

Mix.

INCISIVE POTION. (FORM. HÔP. FR.; ELLIS; RATIER)

- R. Gum Potion, four ounces.
 Oxymel of Squill, half an ounce.

Mix. (RATIER)

- R. Gum Ammoniac, one half part.
 Pulverize, and add gradually
 Vinegar of Squill, thirty parts.
 Infusion of Hyssop, one hundred parts.

Mix. (FORM. HÔP. FR.)

- R. Gum Ammoniac, one dram.
 Water, four ounces.
 Oxymel of Squill, two ounces.
 Compound Tincture of Camphor, half an ounce.

Mix.—Dose, one spoonful every two hours. (ELLIS)

PECTORAL SYRUP. (CADET DE GASSICOURT)

- R. Oxymel of Squill, one ounce and a half.
 Syrup of Ipecacuanha, } of each,
 ——— Poppy, } two ounces.
 ——— Orange Flowers, half an ounce.

Mix.—Dose, two spoonfuls, every hour, in a cupful of pectoral infusion.—It is recommended in hooping cough.

E.

Extraction by Oil.

COMPOUND OIL OF BRYONY.

Oleum Bryoniae compositum. (HISP.)

- R. Fresh Root of Bryony, } of each,
 ——— Elaterium, } one pound.
 ——— Squill, } of each,
 ——— Iris, } half a
 ——— Dwarf Elder, } pound.
 Olive Oil, four pounds.

Boil slowly to the consumption of the humidity, and strain.

COMPOUND OINTMENT OF BRYONY.

Unguentum Bryoniae seu Agrippæ. (HISP.; SARD.; WIRTEM.)

- R. Juice of Bryony, three pounds.
 ——— Elaterium, two pounds.
 Squill, } of each,
 Root of Iris, } six ounces.
 ——— Male Fern, }

Macerate for twelve hours; then boil, express, evaporate to the consistence of soft extract, and add

- White Wax, fifteen ounces.
 Oil of Mucilages, four pounds.

Mix. (SARD.)

- R. Root of Bryony, two pounds.
 ——— Elaterium, one pound.
 ——— Dwarf Elder, } of each,
 ——— Male Fern, } three ounces.
 ——— Iris, }
 Squill, half a pound.
 Water-chesnut, two ounces.
 Olive Oil, six pounds.

Macerate for eight days; then boil, express, and add

- Yellow Wax, one pound.
 Melt in a water-bath. (WIRTEM.)

- R. Compound Oil of Bryony, four pounds.
 White Wax, one pound.

Melt together slowly. (HISP.)

It is feebly stimulant, resolvent, and antihydropic.

SCLERODERMA CERVINUM, PERS.—HART'S TRUFFLE.

SYNONIMES :—Truffe de cerf, FR.; Hirschtrüffel, GERM.; Hertsbull, DUT.

BRUNS.; WIRTEM.; SPIELMANN.

A mushroom (*Boletus cervinus*, *Tubera cervina*), which grows in almost all parts of Europe. (*Fig. Mic. Gen.* 220. t. 99. f. 4)

It is globular, smooth, rootless, dry, membranous, coriaceous, and of a blackish-grey colour. Its substance is spongy, and becomes converted into a blackish-red powder. It has a strong odour and taste which are removed by desiccation.

The tincture of this mushroom has been esteemed aphrodisiac.

SCOLOPENDRIUM OFFICINALE, SM.—HART'S-TONGUE.

SYNONIMES. — Scolopendre, Langue de cerf, Langue de bœuf, FR.; Hirschzunge, GERM.; Gelenj gazyk, Psany trank, Ceterak, BOHEM.; Hiortetongue, DAN.; Lengua de ciervo, SPAN.; Hertstong, DUT.; Lingua di cervo, ITAL.; Jalení Szczaw, POL.; Lingua cervina, PORT.; Hiorttunga, SWED.

BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; WIRTEM.; HERBIP.;

BERGIUS; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the south of Europe. (*Cryptogamia*, LINN.; *Filices*, JUSS.; *Fig. Zorn*, *Ic. pl.* t. 47)

The herb (*Herba Scolopendrii* seu *Lingua cervina* seu *Phyllitidis* seu *Lonchitidis*) is employed. It consists of long leaves, heart-shaped at their inferior part, with hairy footstalks. It has an astringent taste.

SCORZONERA.

Two species of this genus of plants are mentioned in pharmacopœias :

1. *Scorzonera humilis*, LINN.

SYNONIMES :—Scorzonère d'Allemagne ou de Bohême, FR.; Waldskorzonere, Nattermilch, GERM.

AMST.; BRUNS.; GALL.; FERR.; FULD.; HANN.; LIPP.; OLDEN.; BORUS.; WIRTEM.; HERBIP.; MURRAY and GMELIN.

A plant of the north of Europe. (*Syngenesia Polygamia Æqualis*, LINN.; *Synantherea*, CASS.; *Fig. Jacq. Austr.* VI. t. 36)

The root (*Radix Scorzonera latifolia sinuata* seu *serpentaria*) is employed. It is anulated, thick, ligneous, black without, white within, and when fresh, full of a milky juice. It has a bitterish and austere taste.

2. *Scorzonera Hispanica*, LINN.—Garden Scorzonera.

SYNONIMES :—Scorzoncre d'Espagne, SALVOL. II.

sifis noir, FR.; Garten scorzonere, Schlangengras, GERM.; Hadj mordowy koren, BOHEM.; Skorzonera, DAN.; Escorzonera, SPAN.; Spaansche scorzonera, DUT.; Scorzonera, ITAL.; Escorzonreira, PORT.; Scorzonera, SWED.

HISP.; GALL.; FERR.; HANN.; OLDEN.; BORUS.; SAX.; BERGIUS; BRUGNATELLI; SPIELMANN; ZARDA.

A plant of the temperate and warm climates of Europe, and cultivated in kitchen gardens. (*Fig. Zorn Ic. pl.* t. 405)

The root (*Radix Scorzonera*) is employed.

It is long, of the thickness of the finger, black externally and white internally, wrinkled annularly, and full of a milky juice which flows when it is cut. It is sold, after being rasped and dried, in long pieces, which are of a brownish-red-yellow colour. It is inodorous. Its taste is sweetish, bitter, and farinaceous when fresh, but is only mucilaginous when dried.

It is excitant and diaphoretic.

EXTRACT OF SCORZONERA.

Extractum Scorzonerae. (BRUNS.)

- R. Root of Scorzonera, one pound.
 Spring Water, six pounds.

Macerate for four days in a warm place; then boil a little, and express: evaporate the liquor to the proper consistence.

BEZOARDIC DECOCTION OF SCORZONERA.

Decoctum Scorzonerae bezoardicum, Bezoardicum Curvi incompletum. (HISP.)

- R. Root of Scorzonera, one ounce.
 Bruised Lemon Pips, two drams.
 Water, six pounds.

Boil down to four pounds; then strain, and add

Bezoardic Powder of Cinchona,
 half an ounce.

Honey of Elder, three ounces.

Mix.—Dose, four ounces.

APERIENT BEZOARDIC DECOCTION OF SCORZONERA.

Decoctum Scorzonerae bezoardicum solutivum, Bezoardicum completum seu purgans Curvi. (HISP.)

- R. Bezoardic Decoction of Scorzonera, four pounds.
 Senna, half an ounce.

Boil slowly, strain, and add to the liquor

Syrup of Pale Roses, six ounces.
 Bezoardic Powder of Cinchona, half an ounce.

Mix. — It is purgative. — Dose, four ounces.

SCROPHULARIA.—SCROFULARIA, FIGWORT.

Two species of this genus are used in medicine:

1. *Scrophularia aquatica*, LINN.—Water Figwort.

SYNONIMES:—Scrofulaire aquatique, FR.; Wasserbraunwurz, Wasserkropfwurz, GERM.; Escrofularia aquatica, SPAN.; Water speenkruid, DUT.; Scrofularia aquatica, ITAL.; Escrofularia dos rios, PORT.

HISP.; GALL.; GENEV.; WIRTEM.; MURRAY and GMELIN.

A plant growing on the borders of streams all over Europe. (*Didynamia Angiospermia*, LINN.; *Scrofularia*, JUSS.; *Fig. Flore Medic. VI. 321*)

The herb (*Herba Scrophulariae aquaticae seu Betonicae aquaticae seu Ficariae seu Ferrariae seu Castranulae*) is employed. It consists of a glabrous, tetragonal, branching stem, and of opposite, petiolated, oval, nearly cordiform, crenelated, green, glabrous leaves. It has a disagreeable, fetid, repugnant odour, and a bitter, sub-acrid, nauseous taste.

2. *Scrophularia nodosa*, LINN.—Knot-rooted Figwort.

SYNONIMES:—Scrofulaire noueuse, FR.; Knotige braunwurz, Feigwurzengewurz, Knollenwurz, GERM.; Swinsky neb kyticnjkowy koren, Roschodnjak aneb prut-znik, BOHEM.; Fredewnik, POL.

DUBL.; BRUNS.; HISP.; FERR.; GENEV.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; MURRAY and GMELIN; SPIELMANN; ZARDA.

A common plant of Europe. (*Didynamia Gymnospermia*, LINN.; *Scrofularia*, JUSS.; *Fig. Zorn, Ic. pl. t. 28*)

The root and herb are employed.

The root (*Radix Scrophulariae seu Scrophulariae fetidae seu vulgaris*) is thick, knotty, white, of a disagreeable odour, and of a bitter, sub-acrid taste, which it partly loses by desiccation.

The herb consists of an angular stem, and of entire, oval, tri-nerved leaves. It is inodorous, but has a bitter and nauseous taste.

OIL OF FIGWORT. (VAN MONS)

- R. Fresh Herb and Root of Figwort, three parts.
 Olive Oil, one part.

Infuse, and express.

SECALE CEREALE, LINN.—RYE.

SYNONIMES:—Seigle commun, FR.; Roggen, GERM.; Rug, DAN.; Centeno, SPAN.; Rog, DUT.; Segale, ITAL.; Rez, POL.; Sontejo, PORT.; Ræg, SWED.

BAVAR.; HISP.; GALL.; FERR.; GENEV.; CAST. BORUS.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant said to be originally of Crete and the borders of the Caspian Sea. (*Triandria Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Blackw. Herb. t. 424*)

The seeds (*Semen Secalis* seu *Frustrum*) are employed. They are oblong, cylindrical, rather pointed, and of a reddish colour.

RESOLVENT MEALS. (SPIELMANN)

R. Flour of Rye,	} of each, an equal part.
_____ Wheat,	
_____ Barley,	
_____ Beans,	

Mix.

ABSORBENT POWDER.

Pulvis absorbens pro usu externo. (BRERA)

R. Flour of Rye,	four ounces.
_____ Lupins,	eight ounces.

Mix.—The parts affected with erysipelas are to be covered with it, night and morning.

SECALE CORNUTUM.—SPURRED OR HORNED RYE.

SYNONIME:—Ergot, FR.

AMER.; COXE.

A sort of horn or spur which is developed on the spikes of some of the gramineæ, but principally on those of rye. It is usually curved and long, seldom globular, and most frequently distinguished by three blunt angles and by longitudinal lines. Its colour is violet with various shades. Its substance is firm and of a dull white colour: it breaks with sound, and presents a clean fracture. Its smell is disagreeable when fresh, and when found together in a large quantity. Its taste is sub-acrid and nauseous. It attains the length of from an inch to an inch and a half, and is two or three lines thick.

DECANDOLLE says it is a species of fungus, *Sclerotium clarum*, but the more probable opinion is, that it is a morbid degenerescence of the ovarium of the gramineæ on which it is found, occasioned, as MARTIN FIELD thinks, by the puncture of an insect of the genus *Musca*, which does not deposit its eggs in the wound, but leaves in it a blackish liquid.

It contains, according to VAUQUELIN, a yellow colouring matter, a violet colouring matter, a limpid oil, a free acid, which appears to be the phosphoric, free ammonia, and an azotic matter.

It is exceedingly irritant, the use of it occasioning tetanus, gangrene of the extremities, and death. In small doses it seems to excite the contraction of the ute-

rus; and it has from this property been recommended in protracted labour when arising from inertia of the uterus, and to arrest hæmorrhage resulting from a deficiency in its tone.—Dose, from ten to thirty grains in six ounces of vehicle.

INFUSION OF SPURRED RYE.

Infusum Secalis Cornuti. (NIEMANN)

R. Spurred Rye,	half a dram.
Boiling Water,	four ounces.

Infuse, and strain.

One third to be given, and another third twenty-four hours after, if the case seems to require it.

DECOCTION OF SPURRED RYE.

Decoction Secalis Cornuti. (AUGUSTIN; NIEMANN)

R. Spurred Rye,	one ounce.
Spring Water,	one pound and a half.

Boil down to one pound.—Many practitioners prefer this to the infusion.—Dose, a spoonful, every quarter of an hour.

OCYTIC POTION. (PIERQUIN)

R. Spurred Rye,	one dram.
Simple Syrup,	three ounces.
Tincture of Opium,	twenty drops.
Essence of Bergamotte,	a sufficient quantity.

Mix.

SEDUM.

Three species of this genus of plants are used in medicine :

1. *Sedum Rhodiola*, CAND.—Rosewort.

SYNONIMES :—Orpin, Orpin rose, FR. ; Rosenwurz, Rhodiserwurz, GERM. ; Rosenrod, DAN. ; Roosenwortel, DUT. ; Rodiola, ITAL. ; Rosenrot, SWED.

BRUNS. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; SPIELMANN.

A plant of the high mountains of Europe. (*Diecia Octandria*, LINN. ; *Crassulea*, CAND. ; *Fig. Zorn, Ic. pl. t. 180*)

The root (*Radix Rhodiæ* seu *Rosariæ*) is employed. It is branched, white, fleshy, and tuberculous, particularly at its upper part. It has a very agreeable rosaceous odour ; its taste is sub-acrid and styptic.

It is slightly astringent.

2. *Sedum acre*, LINN.—Wall-pepper.

SYNONIMES :—Vermiculaire, Vermiculaire brulante, Poivre de muraille, FR. ; Katzenrœnlein, Mauerpfeffer, Kleines Hauslauch, GERM. ; Netresk, Tucny muzjk neymensjj, BOHEM. ; Helleknop, DAN. ; Siempreviva menor, Uvas de gato, SPAN. ; Sherp huislock, DUT. ; Sedo acre, Erba pignuola, ITAL. ; Vermicularia, PORT. ; Matodile, RUSS. ; Fetknoppar, Hællegræs, SWED.

AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; HISP. ; GALL. ; FERR. ; GENEV. ; SAX. ; SUEC. ; WIRTEM. ; BERGIUS ; GUIBOUT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A common plant of Europe. (*Decandria Pentagynia*, LINN. ; *Crassulea*, JUSS. ; *Fig. Flore Medic. IV. 209*)

The herb (*Herba Sedi minoris* seu *Illecebræ vermicularis*) consists of a glabrous

stem furnished with loose, oval, rather triangular, short, succulent leaves of a light green colour, and growing very near to each other.

It is destitute of odour but has an acrid, pungent, pepperish taste, which is removed by desiccation.

It is excitant, emetic, and purgative internally ; rubefacient externally.

ANTI-SCORBUTIC BEER. (NIEMANN)

R. Wall Pepper, eight handfuls.
Beer, eight pounds.

Boil down to one half.

Dose, three or four ounces in the morning, fasting.

3. *Sedum Telephium*, LINN.—Orpine.

SYNONIMES :—Joubarbe des vignes, Herbe aux charpentiers, FR. ; Fitte Henne, Wundkraut, GERM. ; Kræfurt, DAN. ; Letefolio, SPAN. ; Hemels leutel, DUT. ; Sopra vivolo, ITAL. ; Wronie masla, POL. ; Telethio bastardo, PORT. ; Kuringkal, SWED.

ANT. ; BRUNS. ; HISP. ; GALL. ; FERR. ; WIRTEM. ; BERGIUS ; GUIBOUT ; MURRAY and GMELIN ; SPIELMANN.

A plant of Europe. (*Hexandria Pentagynia*, LINN. ; *Crassulea* JUSS. ; *Fig. Zorn, Ic. pl. t. 486*)

The root and leaves are employed.

The root (*Radix Crassulæ* seu *Crassulæ majoris* seu *Faburæ* seu *Telephii vulgaris* seu *Illecebræ majoris*) is tuberculous, thick, fleshy, and whitish.

The leaves are sessile, oval, of a light-green colour, sometimes reddish, fleshy, and succulent, dentated on the borders, and of a saline, acidulous, and mucous taste.

It is esteemed vulnerary.

SELINUM.—SPIGNELE.

Three species of this genus are mentioned in pharmacopœias :

1. *Selinum Cervaria*, RO.—Broad-leaved Spiguel.

SYNONIMES :—Carotte de montagne, Persil de montagne, FR. ; Hirschwurz,

Hirschheil, Hirschketersilje, Bergpetersilje, GERM. ; Gekruiste beerwortel, DUT.

WIRTEM. ; SPIELMANN.

A plant of the mountains of Europe. (*Pentandria Digynia*, LINN. ; *Umbellifera*, JUSS. ; *Fig. Jacq. Austr. I. t. 69*)

The root (*Radix Cervarea, nigra* seu *Gen-tiana nigra*) is employed. It is long, thick, annulated, blackish externally, and white internally. It has a strong and agreeable odour, and an aromatic and sharp taste.

2. *Selinum palustre*, LINN.—Marsh Spiguel.

SYNONIMES :—Persil des marais, FR. ; Sumpfselenich, Milcpeterling, Sumpfpetersilje, GERM. ; Wandmerke, DAN. ; Apio lechal, SPAN. ; Wilde eppe, DUT. ; Finskingfära, SWED.

BRUNS. ; WIRTEM. ; SPIELMANN.

A common plant of Europe, growing in marshy places. (*Fig. Jacq. Austr. II. t. 152*)

The root (*Radix Tyllselini* seu *Olsnitii*) is employed. It is fusiform, branching, yellowish or blackish without, and white within. It has a strong, aromatic, and disagreeable odour, and a strong, bitterish, and burning taste.

3. *Selinum Oreoselinum*, Ro.—Divaricated Spiguel.

SYNONIMES :—Selin de montagne, Persil de montagne, FR. ; Bergpetersilje, Grundheil, GERM. ; Vilde quale rodde, DAN. ; Apio de montana, SPAN. ; Bergselie, DUT. ; Apio de montagna, ITAL. ; Opich gorny, Pietruszka gorna, POL. ; Apio di montanna, PORT. ; Vild morat, SWED.

BRUNS. ; GALL. ; HANN. ; POL. ; BORUS. ;

SAX. ; WIRTEM. ; MURRAY and GME-LIN ; SPIELMANN.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 400*)

The root, herb, and seeds are employed.

The root (*Radix Oreoselini* seu *Apii montani*) is of the thickness of the finger, very long ; fibrous, covered with a blackish epidermis, underneath which it is white, and in the middle yellowish. It has an agreeable and aromatic odour, and an aromatic and bitterish taste.

The herb consists of a round, slightly striated, smooth stem of a reddish-brown colour, and of trebly-winged leaves of a light-green colour, smooth, and having lanceolated leaflets. Its odour is agreeable ; its taste is bitter and aromatic.

The seeds are oval, flattish, smooth on one side, grooved on the other, red, and surrounded with a white border. They have a strong and a bitter taste.

It is excitant, diaphoretic, and nerveine.

TINCTURE OF SPIGUEL.

Tinctura Oreoselini. (VAN MONS)

R. Fresh Leaves of Spiguel, two ounces.
Seeds of Spiguel, one ounce.
Proof Spirit, fourteen ounces.

Infuse for several days, and strain.

It is excitant and carminative, and is recommended as a stimulant for the stomach.
—Dose, one ounce, twice a day, after meals.

SEMEN CINÆ, SEU SANTONICI.—SEMEN CONTRA.

SYNONIMES :—*Artemisia Santonica Semi-na*, (DUBL.)—*Semen Cynæ* seu *Sinæ* seu *contra vermes* seu *lumbricorum* seu *sanc-tum* seu *Zedoaria*, *Sementina*.—*Semen-contra*, Barbotine, Sementine, Graine de Zédoaire, FR. ; Zittwersaame, Wurmsaame, GERM. ; Seme santo, ITAL. ; Cyt-warowe nasiene, POL.

DUBL. ; EDINB. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; FORM. HÔP. FR. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HER-BIP. ; WURCEB. ; BERGIUS ; COXE ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

There are three sorts of semen contra in commerce, the Aleppo, the East Indian, and the Barbary. It consists of one third of small linear, striated seeds, obtuse at both

ends, and of a greenish-yellow colour ; of one third of small, stunted, shagreened tops, obtuse at the larger end, composed of small tubercles of the same colour as the seeds ; and of one third of extraneous substances, fragments of branches of the plant or of other vegetables, &c. It has an aromatic, very strong, and somewhat anisated odour, and a bitter and acrid taste.

It is generally supposed to be obtained from the *Artemisia Judaica*, LINN., a plant (*Syngenesia Polygamia Superflua*, LINN. ; *Synanthereæ*, CASS.) of Arabia and Judea (*Fig. Zorn, Ic. pl. t. 367*), and from the *Artemisia contra*, LINN, a plant of Persia. (*Fig. Hand. ber. pharm. bot. LXXIV. f. 355*)

SAUNDERS attributes it to a species of *Chenopodium*.

It is anthelmintic.—Dose, from twenty grains to two drams.

INFUSION OF SEMEN CONTRA.

Infusum Seminis Santonici. (AUGUSTIN;
BRERA; NIEMANN)

R. Semen Contra, two drams.
Water or Wine, three ounces.

Infuse, and strain.—Dose, a spoonful,
three or four times a day. (NIEMANN)

R. Root of Valerian, } of each,
Semen Contra, } half an ounce.
Boiling Water, nine ounces.

Infuse, and add to the strained liquor.

Peppermint Water, two ounces.
Extract of Walnut
Rind, } of each,
Ammoniated Tincture of Valerian, } three drams.

Mix.—Dose, a spoonful every hour. (AUGUSTIN)

R. Semen Contra, half an ounce.
Fennel Water, } of each,
Peppermint Water, } four ounces.
Camphor, dissolved in Sulphuric
Æther, one dram.

Mix.—Dose, a spoonful, every hour.
(AUGUSTIN)

R. Semen Contra, two drams.
Rhubarb, one dram.
Boiling Water, a sufficient quantity.

Infuse for a quarter of an hour and strain.
(BRERA)

ANTHELMINTIC POTION.

Potio anthelmintica parum laxans. (BRERA.)

R. Rhubarb, one dram.
Semen Contra, two drams.
Boiling Water, a sufficient quantity,
to obtain six ounces of infusion. Add to
the strained liquor,
Compound Syrup of Cichory,
one ounce.

Mix.—To be taken at twice.

DECOCTION OF SEMEN CONTRA. (RATIER)

R. Semen Contra, half an ounce.
Boiling Water, two pounds.
Boil for a short time.

VERMIFUGE DECOCTION.

Decoctum amarum anthelminticum.
(BRERA)

R. Semen Contra, } of each,
Root of Gentian, } two drams.
Tops of Wormwood, one handful.
Water, a sufficient quantity.
to obtain one pound of strained liquor.
Boil for half an hour.

VERMIFUGE POWDER.

*Pulvis vermifugus seu anthelminticus seu
contra vermes.* (BRUNS.; FERR.; GE-
NEV.; LIPP.; WURCEB.; AUGUSTIN;
SAUNDERS)

R. Semen Contra, } of each, one
Leaves of Plantain, } ounce and a half.
Soap of Jalap, half an ounce.
White Sugar, one ounce.

Make a powder. (LIPP.)

R. Semen Contra, half an ounce.
Senna, one dram.
Coriander, } of each,
Calcined Hartshorn, } half a dram.

Make a powder. (BRUNS.)

R. Semen Contra, half an ounce.
Tartrate of Potass, three drams.
White Sugar, two drams.

Mix.—(SAUNDERS).

R. Semen Contra, } of each,
Root of Jalap, } half a scruple.
Sulphuret of Mercury and
Antimony, six grains.

Make a powder. (WURCEB.)

R. Semen Contra, one scruple.
Muriate of Ammonia, half a scruple.
Sulphate of Iron, two grains.

Make a powder. (WURCEB.)

R. Semen Contra, } of each,
Flowers of Tansey, } three drams.
Sulphate of Iron, one dram.

Make a powder. (GENEV.)

R. Semen Contra, } of each,
Root of Valerian, } half a scruple.
Jalap, } half a scruple.

Make a powder. (FERR.)

R. Semen Contra, from ten to fifteen grains.
Oxide of Zinc, from one to three grains.

Make a powder. (AUGUSTIN)

R. Semen Contra, } of each,
Extract of Tansey, } six grains.
Iron Filings, four grains.
Oil of Valerian, one drop.

Make a powder. (AUGUSTIN)

R. Semen Contra, half an ounce.
Extract of Cinchona, } of each,
Sulphate of Iron, } one scruple.

Make a powder. (AUGUSTIN)

ANTHELMINTIC BOLUS. (FORM. HÔP. FR.)

R. Semen Contra, ten grains.
Proto-chloruret of Mercury, two grains.
Despumated Honey,
a sufficient quantity.

Make a bolus.

ANTHELMINTIC PILLS. (AUGUSTIN; NIE-MANN)

- R. Extract of Tansey, } of each,
 Semen Contra, } six grains.
 Sulphate of Iron, } four grains.
 Resin of Jalap, triturated with
 sweet almonds, } six grains.
 Oil of Tansey, } one drop.

Make eleven pills.—For one dose, to an adult. Three or five may be given to a child.

VERMIFUGE CONFECTION.

Confectio Seminis Cinæ. (LIPP.; PALAT.; SAX.; WIRTEM.; HERBIP.; PIDERIT)

- R. Semen Contra, any quantity.
 Set it over the fire in a copper pan, and when it is very hot, add, by spoonfuls,
 Syrup, boiled to a proper consistence,
 a sufficient quantity,
 to cover the seeds with a thick crust.

ANTHELMINTIC TROCHISCS.

Tabellæ contra vermes seu de Semine Santonico, Morsuli contra vermes seu anthelmintici, Trochisci anthelmintici, Tragemata de Semine Santonico. (HASS.; PALAT.; SARD.; WIRTEM.; HERBIP.; PIDERIT; SPIELMANN)

- R. Semen Contra, } of each, six drams.
 Serpentaria, }
 Jalap, } of each, one ounce.
 Rhubarb, }
 Thick Syrup, } three pounds.
 Pulverize, and make trochiscs. (SARD.)
- R. Semen Contra, one ounce.
 Myrrh, half an ounce.
 White Sugar, one pound.
 Mucilage of Tragacanth,
 a sufficient quantity.

Make trochiscs. (SARD.)

- R. Semen Contra, } of each,
 Corsican Moss, } half an ounce.
 Black Sulphuret of Mercury, }
 three drams.
 Cinnamon, } two drams.
 Resin of Jalap, } two scruples.
 Sugar, boiled à la plume, seven ounces.
 Make trochiscs. (PALAT.; SARD.; WIRTEM.; SPIELMANN)

- R. Semen Contra, one ounce and a half.
 Sulphate of Iron, half an ounce.
 White Sugar, eight ounces.
 Pulverize, and with ten drams of this powder, and mucilage of tragacanth, make thirty trochiscs. (HASS.; PIDERIT)

ANTHELMINTIC ELECTUARY. (LIPP.; AUGUSTIN)

- R. Semen Contra, } of each,
 Valerian, } one dram and a half.

Jalap, one dram.
 Oxymel of Squill, } of each, one ounce.
 Simple Syrup, }

Mix.—Dose, from one to three drams. (LIPP.)

- R. Semen Contra, }
 Jalap, } of each,
 Valerian, } one dram.
 Sulphate of Soda, }
 Oxymel of Squill, } four ounces.

Mix.—Dose, a tea spoonful, every two or three hours. (AUGUSTIN)

ANTHELMINTIC LOHOCH. (HASS.)

- R. Semen Contra, }
 Root of Valerian, } of each, one dram.
 Jalap, }
 Honey of Squill, a sufficient quantity,
 to make with the powders a thick mass.

VERMIFUGE POTION. (PIERQUIN; SWEDIAUR)

- R. Sulphate of Potass, two drams.
 Rhubarb, one dram and a half.
 Semen Contra, one dram.
 Oxymel of Squill, half an ounce.
 Water, two ounces and a half.

Mix. (SWEDIAUR)

- R. Strong Decoction of Semen Contra, two ounces.
 Mint Water, } of each,
 Orange-flower Water, } one ounce.
 Tincture of Cinchona, } two ounces.
 Syrup of Vinegar, a sufficient quantity.

Mix. (PIERQUIN)

- R. Purslain Water, } of each,
 Cichory Water, } three ounces.
 Syrup of Lemon, } eleven scruples.
 Confection of Hyacinth, } of each,
 Semen Contra, } half a dram.
 Orange-flower Water, } half an ounce.

Mix. (PIERQUIN)

- R. Syrup of Wormwood, } of each,
 Bugloss, } one ounce.
 Lemon, } half an ounce.
 Semen Contra, } of each,
 Calcined Hartshorn, } one scruple.
 Alkermes Confection, }
 Dog-grass Water, } of each, a
 Wormwood Water, } sufficient quantity

Mix. (PIERQUIN)

- R. Lettuce Water, } of each,
 Decoction of Dog-grass, } three ounces.
 Confection of Hyacinth, } of each,
 Semen Contra, } half a dram.
 Syrup of Lemon, } six drams.
 Oil of Almonds, } one ounce.

Mix. (PIERQUIN)

Each of these potions are to be taken at one dose.

SEMPERVIVUM TECTORUM, LINN.—GREAT HOUSELEEK.

SYNONIMES :—Joubarbé des toits, Grande joubarbe, FR.; Grosses Hauslauch, Hauswurzkraut, GERM.; Netresk Tucymuzjik wettsj, BOHEM.; Hauslæg, DAN.; Siempreviva de tejados, SPAN.; Donderboard, Husslook, DUT.; Siemprevivo, ITAL.; Hazi zœld, MAG.; Rozchadnik-wielka, POL.; Sempreviva, PORT.; Tschesnok dikoi, RUSS.; Husslæk, SWED.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HAMB.; LIPP.; OLDEN.; ROSS.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A common plant of Europe. (*Dodecandria Dodecagynia*, LINN.; *Crassuleæ*, JUSS.; *Fig. Flore Medic. IV. 208*)

The herb (*Herba Sempervivi seu Sedi majoris*) is employed. It consists of a rosette of fleshy, oval, acute, green leaves, glabrous on both surfaces, and ciliated at the borders. It has scarcely any sensible odour, and a watery, fresh, acerb, styptic, and somewhat saline, taste.

Dose of the juice, two ounces.

SYRUP OF HOUSELEEK.

Syrupus Sedi majoris seu Sempervivi.
(WIRTEM.)

R. Juice of Houseleek, reduced to one half by evaporation, ten ounces.
White Sugar, sixteen ounces.

Boil for a few moments, and strain.

It is refreshing, and cooling.

DEMULCENT LINIMENT. (CADET DE GASSICOURT; SAINTE-MARIE)

R. Juice of Houseleek, } of each,
Olive Oil, } an equal part.

Shake together. (SAINTE-MARIE)

CADET DE GASSICOURT prescribes three ounces of juice, three of lard, and four of oil of almonds.

It is excellent for subduing irritations of the skin, and is said to appease the pain of ulcerated cancer.

SENECIO.

Two species of this genus of plants are employed :

1. *Senecio vulgaris*, LINN.—Groundsel.

SYNONIMES :—Seneçon, FR.; Gemeine Kreutzkraut, GERM.; Kaasurt, DAN.; Hierba cuna, SPAN.; Gemeen kruiskruid, DUT.; Senecione, ITAL.; Przymietowie ziele, POL.; Tarneirinha, PORT.; Kras-townik, RUSS.; Stonært, SWED.

BRUNS.; HISP.; GALL.; WIRTEM.; MURRAY and GMELIN.; SPIELMANN.

A very common plant of Europe. (*Syngenesia Polygamia Superflua*, LINN.; *Synanthèreæ*, CASS.; *Fig. Flore Medic. VI. 324*)

The flowering herb (*Herba Senecionis seu Erigeri*) is employed. It consists of a fistulous, nearly glabrous, branching stem, bearing sessile, alternate, amplexicaul, ciliated, sinuated, denticulated, irregular, gla-

brous leaves, and furnished with flowers which form a very loose terminal corymb. It is destitute of odour; its taste is herbaceous, slightly acid, and somewhat saline.

FINAZZI says that a table spoonful of the juice, administered before an expected fit of hysterical convulsion, will prevent its coming on.

2. *Senecio Jacobææ*, LINN.—Ragwort.

SYNONIMES :—Jacobée, FR.; Jacobskraut, Jacobskreutzkraut, GERM.; Jacobs-kruid, DUT.

GALL.; SPIELMANN.

A common plant of Europe. (*Syngenesia Polygamia Equalis*, LINN.; *Synanthèreæ*, CASS.; *Fig. (Ed. Fl. Dan. t. 944)*)

The herb (*Herba Jacobææ*) is employed. It consists of lyrate, bipinnatifid, dentated, glabrous leaves. It has a disagreeably, bitter and acrid taste.

It is tonic.

SERICUM CRUDUM.—RAW SILK.

SYNONIMES :—*Serici Folliculi*.—Soie, Con-
con de Soie, FR.; Seidenbälglein,
GERM.; Capulio de jeda, SPAN.

HISP.; GALL.; WIRTEM.; GUIBOURT;
SPIELMANN.

An oval pouch of very fine silk of a close

tissue, most frequently of a fine yellow co-
lour, but sometimes white, spun by the
caterpillar of the *Bombix Mori*, in which to
become changed to a chrysalis; it is a le-
pidopterous insect (*Nocturnes Bombycites*,
LAT.) originally of Asia. (*Fig. Roes. Ins.*
III. VII. IX.)

SERUM LACTIS.—WHEY.

SYNONIME :—Petit-lait, FR.

The serous portion of milk.

WHEY IN POWDER. (BRUNS.; NIEMANN)

R. Sugar of Milk, two drams.
White Sugar, one ounce.
Gum Arabic, half a dram.

Mix.—Dissolve it in water.

HOFFMANN'S WHEY

Serum Lactis dulce Hoffmanni. (BRUNS.)

R. Cow's Milk, any quantity.

Evaporate it to dryness, in a tinned cop-
per pan, over a slow fire, constantly stir-
ring; remove the oily portion which usually
floats over the powder, add the same quan-
tity of boiling water as there was milk, boil
a little over a slow fire, and filter.

SWEET WHEY.

Serum Lactis dulce seu sincerum. (HISP.;
GALL.; FERR.; FULD.; HASS.; LIPP.;
LUSIT.; SUEC.; NIEMANN)

R. Boiling Milk, one pound.
Rennet, mixed with water, one dram.

Boil, strain, and add the beaten whites of
three eggs. Boil again for an instant, and
strain through a linen cloth. (FERR.;
FULD.; HASS.; NIEMANN)

R. Cow's Milk, two thousand parts.
Rennet, mixed with a little water,
two parts and a half.

Stand the mixture in hot ashes, and when
coagula begins to be formed, augment the
heat gradually, but not to ebullition, till
the caseous portions adhere to each other;
then pour off the whey, and having let the
curd drip on a small osier hurdle, collect
the liquor, and mix it with the former.
(GALL.; LIPP.; LUSIT.; SUEC.)

VOL. II.

HISP. employs artichoke flowers instead
of rennet.

CHALYBEATE WHEY.

Serum Lactis chalybeatum. (BRUNS.)

R. Whey, any quantity.

Extinguish a red hot iron in it several
times, and then filter.

VINOUS WHEY.

Serum Lactis vinosum. (BRUNS.; FULD.;
HASS.; LIPP.; LUSIT.; AUGUSTIN;
BRUGNATELLI; ELLIS; SAUNDERS;
SAINTE-MARIE; SWEDIAUR)

R. Cow's Milk, four parts.
Acidulous White Wine, one part.

Boil together; strain when coagulated,
and clarify. (BRUNS.; FULD.; HASS.;
SAINTE-MARIE)

LIPP. prescribes three parts of milk and
one part of wine.—ELLIS—half a pint of
milk and one or two ounces of wine.

R. Cow's Milk, } of each, two pounds.
Water, }
Rhenish Wine, one ounce and a half.

Boil, strain, and clarify. (SWEDIAUR)

R. Whey, four ounces.
White Wine, eight ounces.

Boil, strain, and clarify with white of
egg. (SAUNDERS)

LUSIT.; and SAINTE-MARIE prescribe
eight ounces of milk and two pounds of
wine.—AUGUSTIN—three pounds of whey
and one pound of wine.

ACIDULOUS WHEY.

Serum Lactis acidum seu tartarisatum.
(BRUNS.; HISP.; GALL.; FORM. HOP.
FR.; FULD.; HANN.; HASS.; LIPP.;
LUSIT.; SUEC.; NIEMANN; RATIER;
SPIELMANN; VAN MONS)

4 F

R. Cow's Milk, two pounds.
Supertartrate of Potass, four scruples.

Boil the milk, and immediately sprinkle the supertartrate into it; then remove it from the fire, and stir slowly and thoroughly with a spatula: set aside, and when cold strain the liquor through flannel. Shake a beaten white of egg with the whey, set it again on the fire, and at the first boil, pour it upon the strained residue which was set aside. (BRUNS.; FORM. HÔP. FR.; HASS.; NIEMANN; SPIELMANN; VAN MONS)

LIPP. prescribes eighteen parts of milk and one part of supertartrate of potass.—FULD.—thirty-six of milk and one of supertartrate.

Instead of the supertartrate, a little vinegar or lemon juice may be used. (HANN.; HASS.; RATIER)

R. Whey, four pounds.
Whites of Eggs, three.
Supertartrate of Potass, twenty-four grains.

Beat the whites of egg with a little of the whey, then add the rest of this, and boil.

When ebullition has commenced, add the supertartrate gradually, constantly stirring the liquor; then strain, and filter. (HISP.; GALL.; LUSIT.; SUEC.)

ACETOSE WHEY.

Serum Lactis acetosum. (SAUNDERS; SWEDIAUR)

R. Cow's Milk, } of each,
Spring Water, } two pounds.

Boil, and add

Vinegar, one ounce and a half.

When the curd is formed, strain.

COMPOUND WHEY.

Serum Lactis compositum. (BRUNS.)

R. Acidulous Whey, two pounds and a half.
Lemon Juice, one ounce.
Vitriolated Conserve of Roses, six drams.

Mix.

SESAMUM ORIENTALE, LINN.—SESAME.

AMER.; HISP.; GALL.; WIRTEM.; AIN-SLIE; COXE; MURRAY and GMELIN; SPIELMANN.

A plant of Syria and Egypt. (*Didynamia Angiosperma*, JUSS.; *Fig. Burm. Zeyl.* t. 38, f. 1)

The seeds, *Sesame Seeds* (*Sesamum*; *σησαμὸν*.—Sesame, Sesame d'Orient, FR.; Sesam, GERM. and DUT.; Sumsum, ARAB.; Ellu, CAN.; Tal, GUZ.; Till, HIND.; Kunjid, PERS.; Talla, SANS.; Yellu, TAM.; Nuwula, TEL.), are em-

ployed. They are oval, yellow, oleaginous; and of a sweet and agreeable taste.

OIL OF SESAME.

Oleum Sesami orientalis. (HISP.; PALAT.)

R. Sesame Seeds, any quantity.

Bruise them in a mortar, and express. (HISP.)

PALAT. directs them to be torrifed a little before expression.

SESELI TORTUOSUM, LINN.

SYNONIMES:—*Séséli* de Marseille, FR.; Büschelblättriger seseli, GERM.; Comino de Marsella, PORT.

BRUNS.; HISP.; GALL.; WIRTEM.: GUIBOURT; MURRAY and GMELIN.

A plant of the south of Europe. (*Pen-*

tandria Digynia, LINN.; *Umbelliferae*, JUSS.; *Fig. Regn. Bot.* t. 313)

The seeds (*Semen Seseleos Massiliensis*) are employed. They are long, striated, greenish, of an aromatic odour, and of an acrid and aromatic taste.

SILENE.

Three species of this genus of plants are used in medicine :

1. *Silene Behen*, CAND.

SYNONIMES :—Behen commun, FR. ; Wildes Seifenkraut, Junggesell enknopf, GERM.

GALL. ; GENEV.

A common plant of Europe. (*Decandria Trigynia*, LINN. ; *Caryophylleæ*, JUSS. ; *Fig. Céd. Fl. Dan. t. 857*)

The root (*Radix Behen nostratis*) is employed. It is long, articulated, and white.

It is slightly astringent, but is not much used.

2. *Silene Otites*, CAND.—Field Campion.

SYNONIMES :—Silène Otites, FR. ; Feldtiebender Taubenknopf, GERM.

BRUNS. ; WIRTEM.

A plant of Europe. (*Fig. Céd. Fl. Dan. t. 518*)

The herb (*Herba Viscaginis*) is employed. It consists of a simple, viscous stem, and of sparse leaves, the lower ones of which are spatulate, and the upper, lanceolated or linear. It has an astringent and bitterish taste.

3. *Silene Virginica*, LINN.—Catch-fly.

SYNONIME :—Silène de Virginie, FR.

COXE.

A plant of Virginia.

The root is employed. It is considered anthelmintic.

SIMAROUBA AMARA, AUBL.

SYNONIMES :—Simarouba, FR. ; Simarubarinde, Ruhrrinde, GERM. ; Simaruba, ITAL. ; PORT. ; and SPAN.

LOND. ; EDINB. ; DUBL. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOUT ; MURRAY and GMELIN ; PARIS ; PIDERIT ; SPIELMANN ; ZARDA.

A tree of South America. (*Decandria Monogynia*, LINN. ; *Simaroubeæ*, CAND. ; *Fig. Flore Medic. VI. 327*)

The bark (*Cortex Simarubæ*) is employed. It is in long, quilled pieces, thin, flexible, tenaceous, fibrous, and somewhat verrucose externally. It is of a pale yellow without and yellowish-grey within. It is inodorous, but of a bitter taste.

It is very seldom taken from the trunk, but nearly always from the root.

It contains, according to MORIN, quasine, a resinous matter, a volatile oil, alumine, malic acid, a little gallic acid, and various salts.

It is tonic, and is prescribed in intermittent fevers, and mucous discharges of all kinds.—Dose of the powder, from a scruple to half a dram.

DECOCTION OF SIMAROUBA.

Decoctum Simarubæ seu Corticis Simarubæ. (FORM. HÔP. FR. ; FULD. ; GENEV. ; NIEMANN ; SAUNDERS ; SWEDIAUR)

R. Simarouba Bark, half an ounce.
Water, a sufficient quantity
to obtain two pounds of strained liquor.
Boil for a quarter of an hour. (FORM. HÔP. FR.)

NIEMANN prescribes from two drams to half an ounce of bark and two pounds of water ; to be boiled down to six ounces.—SAUNDERS—two drams of bark and two pounds of water ; to be boiled down to one half.—GENEV.—two drams of root and one pound of water ; to be boiled down to twelve ounces.—FULD.—half an ounce of bark and one pound of water, to be boiled down to eight ounces.

Dose, one ounce, every two or three hours, or four ounces, thrice a day.

INFUSION OF SIMAROUBA.

Infusum Simarubæ. (LOND. ; DUBL. ; COXE ; NIEMANN)

R. Simarouba Bark, bruised, half a dram.

Boiling Water, half a pint.

Macerate for two hours in a lightly-covered vessel, and strain.

ASTRINGENT POTION.

Haustus adstringens. (BRERA)

R. Simarouba Bark, half an ounce.
 Water, a sufficient quantity
 to obtain eight ounces of decoction. Strain,
 and add,

Almond Emulsion, two ounces.
 Tincture of Opium, twenty drops.

Mix.—To be taken gradually in atonic
 diarrhœa.

R. Simarouba Bark, half an ounce.
 Water, a sufficient quantity
 to obtain eight ounces of strained decoction.
 Add

Tincture of Opium, one scruple.
 Alcohol, one ounce.
 Cinnamon Water, half an ounce.

Mix.—To be taken as the preceding.

SINAPIS.—MUSTARD.

Two species of this genus of plants are
 used in medicine:

1. *Sinapis alba*, LINN.—White Mustard.

EDINB.; DUBL.; DAN.; GALL.; HANN.;
 LIPP.; POL.; BORUS.; WIRTEM.; EL-
 LIS.; GUIBOUT.; SPIELMANN.

A common plant of Europe. (*Tetradyna-
 miam Siliquosa*, LINN.; *Crucifera*, JUSS.;
Fig. Zorn, Ic. pl. t. 92)

The seeds (*Semen Sinapios alba* seu *Eru-
 cæ*) are employed. They are small, round,
 yellowish, of a fugaceous odour, and of an
 acrid and bitter taste.

2. *Sinapis nigra*, LINN.—Black Mustard.

SYNONIMES:—Moutarde, FR.; Senf,
 GERM.; Sennep, DAN.; Mostaro, SPAN.;
 Mosterd, DUT.; Senapa, Senevra, ITAL.;
 Gorczyca czarna, POL.; Mostarda, PORT.;
 Senap, SWED.

LOND.; DUBL.; AMER.; AMST.; ANT.;
 AUSTR.; BATAV.; BAVAR.; BELG.;
 BRUNS.; DAN.; HISP.; GALL.; FERR.;
 FENN.; FORM. HÔP. FR.; FULD.; GE-
 NEV.; HAMB.; HANN.; HASS.; LIPP.;
 OLDEN.; LUSIT.; POL.; CAST. BORUS.;
 BORUS.; SAX.; SUEC.; WIRTEM.; HER-
 BIP.; WURCEB.; BERGIUS.; BRUGNA-
 TELLI.; COXE.; GUIBOUT.; MURRAY
 and GMELIN.; PARIS.; PIDERIT.; SAUN-
 DERS.; SPIELMANN.

A common plant of Europe. (*Fig. Zorn,
 Ic. pl. t. 152*)

The seeds (*Semen Sinapios nigra* seu *Eru-
 cæ nigra*) are employed. They are small,
 round, black, or blackish-brown, of a fu-
 gacious odour, and of an acrid and bitter
 taste.

These two seeds are strongly stimulant.

They are classed among the anti-scorbu-
 tics.—HENRY and GARROT have discovered
 in it *Sulpho-sinapic Acid*, which approaches

the sulphuric, but differs from it in its ele-
 mentary composition.

DISCUTIENT OINTMENT.

Unguentum Sinapis. (AUGUSTIN)

R. Seeds of Mustard, three ounces.
 Oil of Almonds, half an ounce.
 Lemon Juice, a sufficient quantity.

Mix.—It is extolled by FRANK, in ecchy-
 moses.

STIMULANT BOLUSES.

Boli ad Paralysis. (CADET DE GASSI-
 COURT.; SAUNDERS.; SWEDIAUR)

R. Seeds of Mustard, half a dram.
 — Caraway, } of each,
 Cinnamon, } four grains.
 Ginger, } two grains.
 Simple Syrup, a sufficient quantity.

Make a bolus. (SAUNDERS)

CADET DE GASSICOURT.; and SWEDIAUR
 omit the ginger, but prescribe syrup of gin-
 ger instead of simple syrup.

To be taken twice a day in paralysis.

ANTI-SCORBUTIC ELECTUARY. (VAN
MONS)

R. Powdered Cinnamon, } of each,
 — Mustard, } one part.
 Orange Peel, } of each,
 Extract of Buckbean, } two parts.
 Conserve of Beccabunga, } of each,
 — Horse-radish, } three
 — Scurvy-grass, } parts.
 — Cress, }

Mix.

SINAPISM, MUSTARD CATAPLASM.

Cataplasma epispasticum seu *Sinapis* seu *Si-
 napeos*, *Linimentum rubefaciens*, *Sinapis-
 mus*. (LOND.; DUBL.; BRUNS.; MIL.
 COPEN.; HISP.; GALL.; FERR.; FORM.
 HÔP. FR.; HANN.; HASS.; LUSIT.;

SAX.; WURCEB.; AUGUSTIN; BRUGNATELLI; CADET DE GASSICOURT; COXE; ELLIS; DISP. CHA. PARIS; RATIER; SAUNDERS; SWEDIAUR; VAN MONS)

Cataplasma Sinapis. (LOND.)

- R. Seeds of Mustard, } of each,
 Linseed, } half a pound.
 Hot Vinegar, } a sufficient quantity.

Pulverize, and mix, so as to form a cataplasim.

Cataplasma Sinapeos. (DUBL.; COXE)

- R. Mustard Seeds, in } of each,
 powder, } half
 Crumb of Bread, } a pound.
 Vinegar, a sufficient quantity.

Mix. The cataplasim may be rendered still more acrid by the addition of two ounces of finely-scraped horse-radish.

RATIER prescribes four ounces of linseed poultice and half an ounce of flour of mustard.—WURCEB.—four parts of rye flour, two parts of mustard flour and a sufficient quantity of vinegar.—SWEDIAUR—two parts each, of rye and of mustard flour, and a sufficient quantity of water.—SAUNDERS—two parts each, of wheat and of mustard flour, and a sufficient quantity of vinegar.

R. Flour of Mustard, any quantity.
 Vinegar, a sufficient quantity
 to make a firm paste. (MIL. COPEN.; GALL.; FORM. HÔP. FR.; HANN.; CADET DE GASSICOURT; ELLIS; DISP. CHA. PARIS; RATIER)

The quantity of the mustard prescribed varies from one ounce and a half to four ounces.

R. Flour of Mustard, } of each,
 Crumb of Bread, } an equal part.
 Vinegar, a sufficient quantity
 to make a paste of a softish consistence. (LUSIT.; BRUGNATELLI)

SWEDIAUR; and VAN MONS prescribe one part of mustard, two parts of yeast, and a sufficient quantity of hot vinegar.

R. Yeast, } of each,
 Flower of Mustard, } an equal part.
 Vinegar or Vinegar of Squill, } a sufficient quantity.

Mix. (SAX.)

RATIER prescribes four parts of yeast and one part of mustard.

- R. Oatmeal, four ounces.
 Vinegar, six ounces.

Boil to the proper consistence, adding towards the end,

Powdered Elder Flowers, three ounces.
 Mustard, three drams.

Mix. (AUGUSTIN)

- R. Yeast, three ounces.
 Root of Horse-radish, } of each,
 rasped, } one ounce
 Mustard, bruised, } and a half.
 Vinegar, a sufficient quantity.

Mix. (HANN.)

HENRI; and SWEDIAUR prescribe four parts each, of wheat flour and of flour of mustard, one part of horse-radish, and a sufficient quantity of vinegar.

- R. Yeast, } of each, three ounces.
 Mustard, }
 Black Pepper, bruised, one ounce.
 Vinegar, a sufficient quantity.

Make a paste. (HASS.)

HANN. prescribes one ounce and a half of mustard, one dram of pepper, one ounce each, of horse-radish and of yeast, and a sufficient quantity of vinegar.—CADET DE GASSICOURT—half a pound of mustard, one dram each, of pepper, and of ginger, and a sufficient quantity of oxymel.

- R. Yeast, two ounces.
 Muriate of Soda, } of each,
 Flour of Mustard, } half an ounce.
 Vinegar, a sufficient quantity

to make a paste. (FERR.)

- R. Yeast, one ounce and a half.
 Flour of Mustard, four ounces.
 Muriate of Ammonia, one dram.
 Herb of Rue, bruised, half an ounce.
 Vinegar, a sufficient quantity.

Make a soft paste. (HISP.)

- R. Yeast, } of each, two drams.
 Soft Soap, }
 Mustard, } of each,
 Herb of Rue, bruised, } half an ounce.
 Onions, bruised, one ounce.
 Muriate of Soda, half an ounce.
 Vinegar, a sufficient quantity.

Make a soft paste. (BRUNS.)

WATER OF MUSTARD. (HERBIP.)

- R. Mustard Seeds, one part.
 Water, eight parts.

Macerate for twenty-four hours, and distil.

INFUSED OIL OF MUSTARD.

Oleum Seminum Sinapi infusum. (NIE-MANN)

- R. Mustard freed from its Oil by
 expression, one ounce.

Essential Oil of Rosemary, eight ounces.

Digest for three days, and filter.—To be used as a liniment in paralysis.

WINE OF MUSTARD.

Vinum Sinapios, Collutorium Sinapi. (HERBIP.; AUGUSLIN)

R. Mustard Seeds, bruised, half an ounce.
Wine, one pound.

Macerate for six hours, and decant. (HERBIP.)

R. Mustard Seeds, one ounce.
White Wine, one pound.

Infuse for six hours. Add to the strained liquor

Tincture of Cinnamon, two ounces.

Mix. (AUGUSTIN)

It is sialagogue, and is useful externally, in dropsies.

MUSTARD WHEY.

Serum Lactis sinapinum seu cum Sinapi. (FULD.; HASS.; LIPP.; LUSIT.; NIEMANN.; SAUNDERS.; SWEDIAUR)

R. Cow's Milk, two pounds.
Mustard Seeds, bruised, two ounces.

Boil together till the curd falls to the bottom of the vessel, and filter. (HASS.; LIPP.; NIEMANN.; SAUNDERS)

LUSIT. prescribes one ounce of mustard, and two pounds of milk.—FULD.—one ounce and a half of mustard, and two pounds of milk.

R. Cow's Milk, one pound.
Mustard Seeds, one ounce.

Triturate together, and add

Rhenish Wine, a sufficient quantity.

Coagulate by boiling, and strain. (SWEDIAUR)

It is excitant, and is recommended in arthrodynia, gout, and paralysis.—Dose, one or two pounds, to be taken in the course of the night.

DIURETIC BEER.

R. Ale, sixty pounds.
Mustard Seeds, } of each,
Juniper, } half a pound.
Carrot Seeds, six ounces.

Infuse for several days, and strain. (CADET DE GASSICOURT)

SAUNDERS prescribes one ounce each, of mustard and of juniper, three drams of carrot seeds, two drams of wormwood, and one gallon of beer.—VAN MONS—one part of carrot seeds, two parts each, of juniper and of mustard, and sixty-four of strong beer.

R. Root of Horse-radish, } of each,
Mustard Seeds, } one ounce.
Juniper, } of each,
Subcarbonate of Potass, } six drams.
Strong Beer, six pounds.

Macerate without heat for four days, and strain.

It is given in catarrhal affections of the bladder, and in dropsies.

ALTERATIVE BEER. (SAUNDERS.; SWEDIAUR)

R. Black Mustard, one ounce.
Long Birthwort, six drams.
Lesser Centaury, two drams.
Savine, one dram.
Weak Beer, twelve pounds.

Macerate for four days, and strain.

STIMULANT BEER. (SWEDIAUR)

R. Root of Valerian, one ounce.
Black Mustard, six drams.
Serpentaria, three drams.
Leaves of Rosemary, half an ounce.
Weak Beer, eight pounds.

Macerate without heat for two days, and strain.

SIALAGOGUE GARGLE. (SPIELMANN)

R. Water of Angelica, } of each,
Hyssop, } six ounces.
Spirit of Scurvy Grass, half an ounce.
Ammonia, four scruples.
Flour of Mustard, one dram.
Honey of Roses, four ounces.

Mix.

ASTRINGENT GARGLE. (PIERQUIN)

R. Flour of Mustard, one dram.
Vinegar of Roses, } of each,
White Sugar, } one ounce.
Water, three ounces.

Mix.

STIMULANT GARGLE. (NIEMANN)

R. Flour of Mustard, one dram and a half.
Vinegar, one ounce.
Spring Water, four ounces.

Mix.

SISON AMOMUM, LINN.—FIELD STONE-WORT.

SYNONIMES:—Amome, Amome faux, FR.; Deutsches Amomum, GERM.; Wsecko korenj, BOHEM.

GALL.; WIRTEM.; HERBIP.; SPIELMANN; ZARDA.

A plant of Europe. (*Pentandria Digynia*,

LINN.; *Umbelliferae*, JUSS.; *Fig.* Blackw. *Herb.* t. 442)

The seeds (*Semen Sisonis Amomi*) are employed. They are small, grey, shagreened, sweet, and of an aromatic taste and odour.

It is excitant, and is considered diuretic.

SISYMBRIUM SOPHIA, LINN.

SYNONIMES:—Sagesse des Chirurgiens, FR.; Sophienkraut, GERM.; Barbierförs-stand, DAN.; Talictro, SPAN.; Fiekruid, DUT.; Talitro, ITAL.; Herva sophia, PORT.; Stillfræ, SWED.

GALL.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe. (*Tetradynamia Siliquosa*, LINN.; *Cruciferae*, JUSS.; *Fig.* Zorn, *Ic. pl.* t. 333)

The herb and seeds are employed.

The herb (*Herba Sophiæ chirurgorum*) is composed of a round, branching, somewhat hairy stem, and of alternate, decomposite, glabrous leaves.

The seeds are very small, oval, smooth, and of a reddish-brown colour. It has, like the herb, an acrid and burning taste.

It is excitant, and anti-scorbutic.

SIUM.

Six species of this genus of plants are used in medicine:

1. *Sium angustifolium*, LINN.—Narrow-leaved Water-parnsip.

SYNONIMES:—Berle à feuilles étroites, FR.; Schmalblättriger Mez, GERM.; Berrera de hoja angusta, SPAN.; Smalbladiger watereppe, DUT.

GALL.

A plant of Europe. (*Pentandria Monogynia*, LINN.; *Umbelliferae*, JUSS.; *Fig.* Jacq. *Flor. Austr.* t. 67)

The herb is employed; it consists of a hollow stem, furnished with incised leaflets nearly auriculated at the base. It has a bituminous odour, and a bitter and sub-acrid taste.

It is excitant and diuretic.

2. *Sium latifolium*, LINN.

SYNONIMES:—Berle à large feuilles, FR.; Breitblättriger Merk, GERM.

GALL.

A plant of Europe. (*Fig.* Moris. *Hist.* II. 9. t. 5. f. 1)

The herb is employed; it consists of a hollow stem, and of winged leaves furnished with lanceolated, serrated leaflets.

This plant is nearly always substituted for the preceding.

3. *Sium nodiflorum*, LINN.—Creeping Water-parnsip.

SYNONIMES:—Berle nodiflore, FR.; Knotenblümiger Merk, Kleiner Eppich, GERM.

DUBL.; COKE.

A plant of Europe. (*Fig.* Moris. *Hist.* II. 9. t. 5. f. 3)

The herb is employed; it consists of a fistulous stem, and of simply winged leaves, with lanceolated, dentated leaflets.

It is diuretic.—The expressed juice is, according to MORRISON, useful in various cutaneous disorders. WITHERING used to prescribe three or four ounces of it to adults, taken in the morning, and three spoonfuls to children, twice a day.

4. *Sium Sisarum*.—LINN.

SYNONIMES:—Berle Chervi, FR.; Zucker-

wurzel, Merk, Zuckerwurzel, GERM.; Sisaro, Chirivia de quaresma, SPAN.

HISP.

A plant said to be originally of China. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Moris. Hist.* 3. II. 9. t. 4. f. 8)

The root is composed of several oblong, rugose tuberosities.

The seeds are ovoid, glabrous, and striated.

This plant is said to be a specific against the effects occasioned by the abuse of mercury.

5. *Sium Ammi*, LINN.—Small Stonewort.

SYNONIMES:—Ammi des boutiques, FR.; Cretischer ammey, Mohrenkümmel, GERM.; Amus, ARAB.; Ammi, BOHEM.; Aymadavum, CAN.; Assamodum, CYNG.; Ajuwan, DUK.; Nankhah, PERS.; Ajamodum, Brahnadarbha, SANS.; Womun, TAM.; and TEL.

BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; AINSLIE.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of the south of Europe. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Zorn, Ic. pl.* t. 256)

The seeds (*Semen Ammios veri* seu *Cretici* seu *Feniculi Lusitanici*) are employed. They are small, oval, deeply striated, fawn-coloured, of an agreeable odour, and of an aromatic and bitter taste.

It is excitant and carminative.

6. *Sium Ninsi*, LINN.

SYNONIMES:—Ninsi, FR.; Ninsing, Ninsing Merk, GERM.; Ninsiwortel, DUT.

BRUNS.; GALL.; WIRTEM.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.

A plant of the mountains of China. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Burm. Flore Ind.* t. 29. f. 1)

The root (*Radix Ninsi* seu *Ninsing* seu *Ninsini*) is employed. It is fusiform, annulated, whitish, horny and semi-transparent or yellowish-brown and opaque, as it may or may not have been blanched with boiling water, and deprived of its epidermis. It varies in size from that of a quill to that of the little finger. Its odour is strong and disagreeable, and its taste is at first sweetish, then bitter.

The Chinese attribute almost miraculous virtues to this root, though, like that of ginseng, these are really but very few and limited. In Europe it is rather culinary than medicinal.

SMILAX.

Two species of this genus of plants are employed in medicine:

1. *Smilax China*, LINN.—China.

SYNONIMES:—Squine, Esquine, FR.; Chinawurzel, GERM.; Khush sinie, ARAB.; Koren chyny, BOHEM.; China alla, CYNG.; Chinarbd, DAN.; Chob chinie, DUK.; and HIND.; Esquina, PORT.; and SPAN.; China wortel, DUT.; Icina, ITAL.; Chub chiny, PERS.; Kinarot, SWED.; Paringay puttay, TAM.

AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; GENEV.; OLDEN.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of China and Japan. (*Diæcia Hexandria*, LINN.; *Asparagineæ*, JUSS.; *Fig. Flore Medic.* VI. 329)

The root, *China Root* (*Radix Chinae verae* seu *Chinae Orientalis* seu *Chinae ponderosæ*)

is employed. It is very large, ligneous, nodose, tuberculous, heavy, of a reddish-brown externally, of a reddish-white internally, and inodorous. Its taste is viscous and rather acerb.

It contains much starch, and but few active principles.

It was for a long time reputed a powerful sudorific, but it is now thought to be nearly inert.

DECOCTION OF CHINA ROOT.

Decoctum Chinae Radicis. (RATIER; SWEDIAUR)

R. China Root, one ounce.
Water, two pounds.

Boil, and strain. (RATIER)

R. China Root, of each,
Corinth Raisins, one ounce.
Spring Water, six pounds.

Boil down to one fourth, and strain. (SWEDIAUR)

It is reputed sudorific, but is emollient and demulcent.

COMPOUND SYRUP OF CHINA ROOT.

Syrupus Chinæ compositus. (HISP.)

R	Root of China,	} of each,
	Sarsaparilla,	
	Guaiac Wood, rasped,	} three ounces.
	Red Sanders Wood,	
	rasped,	} of each,
	Water,	
		one ounce.
		twelve pounds.

Digest for twenty-four hours, and then boil down to one half. Add to the strained liquor

White Sugar, four pounds.

Clarify, and make a syrup.

2. *Smilax Sarsaparilla*, LINN. — *Sarsaparilla*.

SYNONIMES: — *Salsaparilla*, *Sarsaparilla*, *Sarsa*, *Zarsa*, *Zarsaparilla*. — *Salseparille*, FR.; *Sasseparille*, *Sarsaparille*, GERM.; *Sarsaparylle*, BOHEM.; *Sarsapiril*, DAN.; *Zarzaparilla*, SPAN.; *Sarzarilla*, DUT.; *Salsaparglia*, ITAL.; *Sarsaparyle*, POL.; *Solsaparilha*, PORT.; *Sassaparill*, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY AND GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

A plant of Peru and Brazil. (*Fig. Flore Medic. VI. 308*)

The root is employed. Four principal varieties of it are met with in commerce:

1. *Brasilian or Portuguese Sarsaparilla*.—This is in bundles of the length of six or eight feet, and about two in circumference, formed of interlaced roots of about the thickness of a goose quill, pliable, grooved longitudinally, having a thick bark, and usually deprived of the stock and fibres. It is reddish or blackish-brown externally, and white and farinaceous internally. It is inodorous, but has a bitterish taste. This sort is the most esteemed.

It comes from Peru and Maranham.

2. *Honduras Sarsaparilla*.—This has a dull brown bark sometimes whitish. It is very long, and more fibrous than the preceding, and the stock is left attached to it. The bark is pinkish internally, and covers a white ligneous axis, the taste of which is rapid and amylaceous, while that of the bark is mucilaginous and often distinctly bitter.

It comes from the Bay of Honduras.

VOL. II.

3. *Red or Jamaica Sarsaparilla*.—This is from six to seven feet long, round and compressed by desiccation, covered with an epidermis of an orange-red, reddish-grey, or whitish colour; the bark is sometimes moist, and its soft part white; the stock is left attached. A small quantity of it is found in the bundles coming from Honduras. Although it is but little esteemed, it deserves to be preferred to the two others, as it has a bitterer, stronger, and more aromatic taste, as well as a more decided odour, which is developed by rubbing it.

The medicinal properties of this root are referable, according to PALOTTA, to a peculiar alkaloid, *Parigline*. FOLCHI has also extracted from the medullary part of *sarsaparilla* another alkaloid called *Smilacine*.

It is generally esteemed diuretic, and, particularly, diaphoretic; but although its action in both these respects is sufficiently feeble, it is evidently excitant, which property the *parigline* possesses in a very marked degree. It is administered chiefly in syphilitic affections, chronic rheumatism, and various cutaneous diseases.

It is given, in powder, in the dose of from half a dram to one dram.

It is incompatible with the infusion of galls, lime water, the acetate of lead, and the nitrate of mercury.

4. *East Indian Sarsaparilla*.—This term is applied to the root of the *Periploca Indica*, LINN., a shrub of the East Indies (*Pentandria Monogynia*, LINN.; *Apocynæa*, JUSS.) This root (Oshba, ARAB.; Erramassumul, IRIMUSU, CYNG.; Mugrabu, HIND.; Shativa, SANS.; Nunnarivayr, TAM.; Sugundapala, TEL.) is prescribed in decoction, in the dose of three or four ounces, in scrofula, syphilitic affections, and cutaneous diseases.

SUDORIFIC AND TONIC POWDER. (PIERQUIN)

R.	Sarsaparilla,	} of each,
	Sassafras,	
	Root of Butcher's-broom,	} an equal part.

Pulverize, and mix.—Dose, one dram, every morning, in white wine.

ARABIAN ELECTUARY. (PIERQUIN)

R.	Sarsaparilla,	} of each,
	China Root,	
	Torrified Filberts,	} four ounces.
	Cloves,	} five ounces.
	Honey,	
		a sufficient quantity.

Mix.—This electuary is much employed in Provence, in syphilitic ulcers of the throat, and in what is called chronic or degenerated syphilis.

4 G

EXTRACT OF SARSAPARILLA.

Extractum Sarsaparillæ. (LOND.; DUBL.; AMST.; BATAV.; HISP.; SARD.; WIRTEM.; ELLIS)

R. Sarsaparilla, sliced, one pound.
Boiling Water, one gallon.

Macerate for twenty-four hours, and boil down to four pints. Strain the decoction while hot, and evaporate it to the proper consistence. (LOND.; DUBL.; COXE)

R. Sarsaparilla, one part.
Boiling Water, ten parts.

Digest for twenty-four hours, then boil for two, and strain; boil the residue, for the same length of time, with

Water, four parts.

Mix the liquors, and evaporate to the consistence of an extract. (BATAV.; HISP.)

AMST. directs the root to be boiled first with twenty and then with fifteen parts of water, each time, till reduced to one half.

R. Sarsaparilla, one pound.
Boiling Water, six pounds.

Macerate for three days, and boil a little. Express, and evaporate to the consistence of an extract. (WIRTEM.)

SARD. directs the decoction to be boiled down to two-thirds before straining and evaporating.

FLUID EXTRACT OF SARSAPARILLA.

Extractum Sarsaparillæ fluidum.
(DUBL.)

R. Sarsaparilla, sliced, one pound.
Water, twelve pints.

Boil for an hour, and pour off the liquor; boil the root again with

Water, twelve pints,
and strain with strong expression.

Set apart the mixed decoctions till the feculencies have subsided; then after having continued the boiling till the liquor is reduced to thirty ounces, add

Rectified Spirit,
two fluid ounces.

Mix.

DECOCTION OF SARSAPARILLA.

Decoctum Sarsaparillæ seu Sarsæ seu Smilacis Sarsaparillæ. (LOND.; DUBL.;

EDINB.; AMER.; LUSIT.; SUEC.; BRUGNATELLI; COXE; NIEMANN; RATIER; SAUNDERS; SWEDIAUR; PH. SYPH.)

Decoctum Sarsaparillæ. (LOND.; DUBL.)

R. Sarsaparilla, sliced, four ounces.
Boiling Water, four pints.

Macerate for four hours in a vessel placed near the fire and lightly covered; then having taken out the root and bruised it, put it again into the liquor, and continue the maceration for two hours longer: finally boil it down to two pints, and strain the decoction.

Decoctum Smilacis Sarsaparillæ.
(EDINB.)

R. Sarsaparilla, sliced, six ounces.
Water, eight pounds.

Digest for two hours in a temperature of about 195 degrees; then having taken out and bruised the root, boil it in the liquor till this is reduced to four pounds; then express, and strain the decoction.

AMER.; and COXE direct one ounce and a half of sarsaparilla to be digested for two hours in two pints of boiling water, the root to be taken out and bruised, then to be returned, and the liquor boiled to the consumption of one half.—PH. SYPH.—three ounces of root, three pounds of boiling water, and four hours' infusion; the infusion to be continued with the bruised root for seven hours, and the liquor to be then boiled till reduced to one half: or, three ounces of root, three pounds of boiling water, and twelve hours' digestion; then ebullition till the liquor is reduced to two pounds.—BRUGNATELLI—six ounces of root, sixteen pounds of water, and after digestion, ebullition till the liquor is reduced to one half.—LUSIT.—the same process, but with ten pounds of water.—NIEMANN; and RATIER—one ounce of sarsaparilla and two pounds of water; to be reduced to one third.—NIEMANN—the same as LOND.—SUEC.—four ounces of sarsaparilla, sixty-four ounces of boiling water, four hours' maceration, then two hours' digestion after having bruised the root, and the reduction of the liquor by boiling to thirty-two ounces.

It is prescribed in syphilitic and rheumatic affections.—Dose, half a pound, three or four times a day.

SAINTE-MARIE'S PTISAN.

- R. Sarsaparilla, sliced, four ounces.
 Water, six pounds.
 Boil down to four pounds. Then having removed it from the fire, add
 Liquorice, four drams.

Infuse, set aside, and decant.

It is recommended in syphilitic affections. —The quantity prescribed should be taken warm, in the morning fasting, in six doses, at intervals of a quarter of an hour.

ANTI-SYPHILITIC LIQUOR OF ROUEN.
 (NIEMANN)

- R. Decoction of Sarsaparilla, thirty-two ounces.
 Nitrate of Potass, three drams.
 Hydrochloric Æther, half an ounce.
 Spirit of Camphor, two drams.
 Oil of Anise, five drops.
 Mix.—Dose, a spoonful, twice or thrice a day.

SYRUP OF SARSAPARILLA.

Syrupus Sarsaparillæ. (LOND.; DUBL.; FEE)

- R. Sarsaparilla, one pound.
 Boiling Water, one gallon.
 Refined Sugar, one pound.

Macerate the root in the water for twenty-four hours; then boil down to four pints, and strain while hot. Add the sugar, and boil till of a proper consistence. (LOND.; DUBL.)

- R. Sarsaparilla, sliced, three pounds.
 Boiling Water, thirty pounds.

Infuse for twenty-four hours; then boil till reduced to twelve pounds, and strain with expression: boil the residue with

Water, thirty pounds.

Reduce to twelve, and again express.

Having mixed the two decoctions, evaporate till reduced to ten pounds, set aside, filter, and add

Simple Syrup, } of each,
 Honey, } three pounds.

Boil rapidly to the proper consistence.

ARABIAN PTISAN. (PIERQUIN)

- R. Sarsaparilla, two ounces.
 China Root, one ounce.
 Water, eight pounds.

Boil down to four pounds.

This is much used in Provence, where it is administered with the electuary of the same name.

COMPOUND DECOCTION OF SARSAPARILLA.

Decoctum Sarsaparillæ compositum seu sudoriferum seu Ligni Guajaci compositum. (HISP.; GALL.; SARD.; CADET DE GASSICOURT; RATIER; SPIELMANN; SWEDIAUR; VAN MONS)

- R. Sarsaparilla, one ounce and a half.
 Water, five pounds.
 Boil for half an hour, and add

Guaiac Shavings, one ounce and a half.

Remove it from the fire at the end of a quarter of an hour, and add

Liquorice, grated, half an ounce.
 Sassafras Wood, two drams.

Cover the vessel, strain the liquor when cold, set it aside, and pour off three pounds. (VAN MONS)

- R. Sarsaparilla, } of each,
 Guaiac, rasped, } one ounce and a half.
 Warm Water, four pounds.

Infuse for twelve hours in a covered vessel, and then boil down to three pounds, and add towards the end

Sassafras Wood, rasped, two drams.
 Liquorice, grated, half an ounce.

Infuse for half an hour, and strain. (GALL.)

HISP. prescribes two ounces each, of sarsaparilla and of guaiac, eight pounds of water, to be reduced to four; one ounce of sassafras, and one dram of liquorice.

- R. Sarsaparilla, two ounces.
 Guaiac Shavings, one ounce.
 China Root, half an ounce.
 Water, two pounds.

Boil down to one half, and infuse at the end

Sassafras Wood, two drams.

Strain with expression. (CADET DE GASSICOURT)

- R. Sarsaparilla, thirty ounces.
 Boiling Water, twenty-four pounds.

Infuse for twenty-four hours, boil down to eight pounds; repeat this operation twice again with the residue. Take twenty pounds of the mixed and strained decoctions, and add

Flowers of Borage, } of each,
 Petals of Pale Roses, } two ounces.
 Senna,
 Seeds of Anise, }

Boil down to twelve pounds, and strain. (SWEDIAUR)

- R. Sarsaparilla, } of each, two ounces.
 China Root, }
 Red Sanders, six drams.
 Hartshorn, one ounce and a half.

Digest for twenty-four hours, boil down to one half, and add

Yellow Sanders, six drams.
White Sugar, six ounces.

Strain. (HISF.)

R. Sarsaparilla, one ounce.
Water, three pounds.
Sulphuret of Antimony, in a bag, two drams.

Isinglass, one dram.

Boil down to two thirds, and strain. (RATIER)

R. Sarsaparilla, } of each,
China Root, } half an ounce.
Guaiac Shavings, } of each,
Sulphuret of Antimony, } two ounces.
Boiling Water, eight pounds.

Macerate for twenty-four hours; then boil for an hour in a covered vessel, and add

Sassafras Shavings, one ounce.

Root of Avens, half an ounce.

Juniper, bruised, two drams.

Boil for a quarter of an hour, and strain. (SPIELMANN)

This decoction may be rendered laxative by infusing in it one ounce of senna.

R. Sarsaparilla, two ounces.
China Root, } of each,
Guaiac Shavings, } one ounce.
Sulphuret of Antimony, one ounce and a half.
Water, six pounds.

Digest for twelve hours in warm ashes; then boil slowly till reduced to two pounds, and add towards the end

Grated Liquorice, six drams.

Strain the liquor, and pour it boiling upon

Sassafras Shavings, three drams.

Cover the vessel till perfectly cold, and strain. (SARD.)

It is recommended in syphilitic, catarrhal, and exanthematic affections.—Dose, several glassfuls, daily.

SUDORIFIC PTISAN. (PIERQUIN; RATIER; SAINTE-MARIE)

R. Sarsaparilla, two ounces.
China Root, } of each, one ounce.
Bitter-sweet, }
Water, three pounds.

Boil down to two pounds, and strain. (PIERQUIN)

Dose, a glassful, morning and evening.

It may be mixed with one third of milk and sweetened with sugar.

R. Sarsaparilla, } of each, one ounce.
Bitter-sweet, }
Root of Soapwort, two drams.
Water, six glassfuls.

Boil down to two thirds, and add

Cow's Milk, two glassfuls.

Mix. (SAINTE-MARIE)

To be taken in the course of the day, at six doses.

R. Sarsaparilla, } of each, half an ounce.
China Root, }
Guaiac, }
Sassafras, two drams.
Water, two pounds.

Boil together. (RATIER)

To be taken as the preceding.

STIMULANT SUDORIFIC DECOCTION.

Decoctum diaphoreticum incitans. (BRERA)

R. Sarsaparilla, } of each, three drams.
Guaiac Wood, }
Bitter-sweet, } of each, two drams.
Fennel, }
Water, a sufficient quantity,
to obtain one pound of decoction. To be taken at twice or more.

NUTRITIVE ANTI-SYPHILITIC DECOCTION.

Decoctum anti-syphiliticum nutriens. (BRERA)

R. Sarsaparilla, one ounce.
Iceland Moss, washed, one ounce and a half.
Water, a sufficient quantity,
to obtain fourteen ounces of decoction.—Add to it when strained.

Simple Syrup, one ounce.

Mix.—To be taken at twice, morning and evening.

COMPOUND DECOCTION OF ISINGLASS. (NIEMANN)

R. Sarsaparilla, three ounces.
Sassafras Bark, } of each, one ounce.
——— Wood, }
Boiling Water, nine pounds.

Digest for twelve hours; after which boil, and add

Isinglass, three ounces,
previously dissolved in a sufficient quantity of water to make eight pounds of liquor.

It is recommended by EIMBKE, in syphilitic diseases.—Dose, from twenty to fifty ounces daily.

FELTZ' PTISAN. (BORIES; CADET DE GAS-SICOURT; PIERQUIN; RATIER)

R. Sulphuret of Antimony, in a bag, four ounces.
Water, twelve pounds.
Sarsaparilla, two ounces.
China Root, one ounce.
Isinglass, } of each,
Bark of Box, } one ounce, and a half.
——— Ivy, }

Boil down to one half, and strain through a sieve; set aside, decant, and add

Deutochloruret of Mercury,
three grains.

Dissolve. (BORIES; CADET DE GASSICOURT; PIERQUIN)

R. Sulphuret of Antimony, powdered
and enclosed in a bag, four ounces.

Suspend it in a glazed earthen vessel, and pour into this

Water, six pounds.

Boil for twelve hours to reduce it to one half. Add

Sarsaparilla, three ounces.
Isinglass, half an ounce.

Leave the vessel in warm ashes for twenty-four hours; then strain the liquor through a fine linen cloth. (RATIER).

The last is the only formula which is followed at the present day. The sulphuret of antimony is useless; by omitting that, the preparation is similar to the preceding one.

If isinglass which has been whitened with sulphuric acid is employed, that substance always retaining a certain quantity of the acid, and this acting upon the sulphuret, forms a portion of sulphate of antimony. This is the cause of the pain and vomiting which this decoction sometimes occasions. CHEVALLIER and IDT however are of opinion that these accidents are sometimes owing to a portion of arsenic mixed with the sulphuret, and which is dissolved by the water. The sulphuret should therefore be abandoned, which would be the best, or, if retained, the isinglass should be replaced by gum arabic, in the proportion of one ounce to one pound and a half of reduced decoction.

VIGAROUS' ALTERATIVE PTISAN. (BORIES; PIERQUIN)

R. Sarsaparilla, six ounces.
Senna, three ounces.

Guaiac Shavings,	} of each, one ounce and a half.
Sassafras Shavings,	
China Root,	
Florentine Iris,	
Sulphuret of Antimony,	
Anise,	
Supertartrate of Potass,	
Long Birthwort,	
Round Birthwort	
Jalap, bruised	
Polypody of the Oak,	} twelve.
Fresh Walnuts, with their rinds,	
bruised,	
White Wine,	three pounds.

Infuse for twenty-four hours in hot ashes, then add

Spring Water, six pounds.

Evaporate slowly to two thirds, and strain through flannel.

The residue serves to make a weaker ptisan with two or three pounds of white wine and six or seven of water, to be also reduced to two-thirds.

Three glassfuls of the first are to be taken in the course of the day, and the second serves for common drink.

SAINTE-MARIE prepares with the decoction an *Anti-syphilitic Rob*, of which the following is the formula:

R. VIGAROUS' Ptisan, one pound.
Sugar, one or two pounds.

Boil to the consistence of a rob.

This rob is reputed anti-syphilitic. — Dose, a spoonful four times a day. After each dose, a large glassful of concentrated decoction of sarsaparilla should be taken, and the dose of the rob increased till the patient takes nine or ten spoonfuls in the course of the day.

CATHOLIC DECOCTION. (SARD.)

R. Root of Dog-grass, one ounce and a half.	
Sarsaparilla,	one ounce.
Leaves of Cichory,	} of each, half a hand- ful.
——— Agrimony,	
——— Betony,	
——— Maidenhair,	
——— Burnet Saxifrage,	
Senna,	} of each, half an ounce.
Dodder,	
Root of Liquorice,	two drams.
Seeds of Anise,	one dram.
Water,	four pounds.

Boil the sarsaparilla and dog-grass in the water till this is reduced to one half; then add the liquorice and the herbs. Macerate the senna, dodder, and anise in six ounces of boiling water for several hours, and strain. Mix the decoction with this infusion.

PURGATIVE ANTI-SYPHILITIC PTISAN.

Decoctum *anti-venereum laxans*. (GALL.; CADET DE GASSICOURT; SPIELMANN)

R. Sarsaparilla,	} of each, one ounce.
Guaiac Shavings,	
Subcarbonate of Potass,	
	twenty-five grains,
Water,	four pounds.

Macerate for twelve hours, occasionally stirring, and then boil down to three pounds. Infuse in the decoction for half an hour,

Senna,	} of each, two drams.
Sassafras Shavings,	
Liquorice,	
Rhubarb,	
Coriander,	} of each, one dram.

Strain with slight expression, and decant when cold. (GALL.; CADET DE GASSICOURT)

- R. Sarsaparilla, } of each,
 China Root, } two ounces.
 Sassafras Shavings, }
 Sulphuret of Antimony, one ounce.
 Spring Water, two pounds and a half.

Boil down to two pounds and add,

- Root of Polypodium, half an ounce.
 ——— Rhubarb, half a dram.
 ——— Liquorice, } of each, one dram.
 Senna, }
 Sassafras Wood, }
 Subcarbonate of Potass, four grains.
 Coriander, half a dram.

Infuse for twelve hours, then boil for a few moments, and strain. (SPIELMANN)

To be taken in the course of the day.

VINACHE'S PTISAN. (CADET DE GASSICOURT)

- R. Sulphuret of Antimony, in a bag, two ounces.

- Sarsaparilla, } of each,
 China Root, } one ounce and a half.
 Guaiac Wood, }
 Water, three pounds.

Boil down to one pound, and add

- Sassafras, } of each, half an ounce.
 Senna, }

Infuse, and strain, set aside, and decant.

It is employed in syphilitic and cutaneous diseases.—Dose, several glassfuls, daily.

JAUSSEURAND'S PTISAN. (BORIES)

- R. Sarsaparilla, } of each,
 China Root, } two ounces.
 Sulphuret of Antimony, }
 Subcarbonate of Potass, one dram.
 Senna, } of each, one ounce.
 Sassafras Wood, }
 Rhubarb, } of each,
 Cinchona, bruised, } half an ounce.
 Water, a sufficient quantity.

Boil for eight hours in a covered vessel, over a slow fire, till only six pounds of liquor remain. Infuse for the same length of time, and decant.

It is principally recommended in scrofula.—Dose, three glassfuls daily.

ASTRUC'S SUDORIFIC PTISAN. (PIERQUIN)

- R. Sarsaparilla, } of each, two ounces.
 Sassafras, }
 Guaiac, } of each,
 China Root, } one ounce and a half.
 Sulphuret of Antimony, two drams.
 Water, twelve pounds.

Infuse with heat for twelve hours, then boil slowly, and add towards the end,

- Senna, half an ounce.
 Liquorice, one ounce and a half.

Strain.

ANTI-ARTHRITIC DECOCTION. (CADET DE GASSICOURT)

- R. Sarsaparilla, four ounces.
 Sulphuret of Antimony, six ounces.
 Water, two pounds.

Boil down to one half. Infuse in the remainder,

- Root of Liquorice, one ounce.
 Seeds of Anise, two drams.

It is recommended by QUARIN.—To be taken by glassfuls.

LISBON PTISAN, LISBON DIET-DRINK.

Decoctum Lusitanicum seu anti-venereum Ulyssiponense seu Sarsaparillæ compositum seu cum Mezereo. (LOND.; DUBL.; AMER.; LUSIT.; SUEC.; COXE; FEE; NIEMANN; SPIELMANN; SWEDIAUR; PH. SYPH.)

Decoctum Sarsaparillæ compositum. (LOND.; DUBL.)

- R. Decoction of Sarsaparilla, boiling, four pints.
 Root of Sassafras, } of
 sliced, } each,
 Guaiac Wood, rasped, } one
 Liquorice Root, bruised, } ounce.
 Bark of Mezereon Root, three drams.

Boil for a quarter of an hour, and strain.

- R. Sarsaparilla, six ounces.
 Sassafras, } of each, one ounce.
 Guaiac, }
 Water, ten pounds.

Macerate for six hours in a gentle heat, and boil down to one half. Add towards the end,

- Liquorice, grated, one ounce.
 Mezereon Bark, three drams.

Strain. (SWEDIAUR)

- R. Sarsaparilla, three ounces.
 Mezereon Bark, two drams.
 Spring Water, three pounds.

Boil down to two pounds, and add towards the end,

- Liquorice, grated, one ounce and a half.

Strain two pounds of liquor. (PH. SYPH.; NIEMANN)

- R. Red Sanders, } of each, one ounce.
 White Sanders, }
 Guaiac Wood, }
 Rhodium Wood, }
 Sassafras Wood, }
 Sarsaparilla, three ounces.
 Sulphuret of Antimony, two ounces.

Mezereon Bark, half an ounce.
Boiling Water, ten pounds.
Infuse, boil down to one half, and add towards the end,

Grated Liquorice, half an ounce.
Strain. (FEE; NIEMANN; SPIELMANN)

R. Sarsaparilla, } of each,
Sassafras, } three ounces.
Red Sanders, }
Guaiac, }
Mezereon Bark, one ounce.
Coriander, from six drams to one ounce.
Spring Water, twenty pounds.

Boil down to one half. (PH. SYPH.)

It is recommended in syphilitic diseases, and affections caused by the abuse of mercury.—Dose, from fifteen to fifty ounces daily, augmenting it gradually.—The third formula, by GIRTANNER, is preferable to the rest.

ANTI-SYPHILITIC PTISAN. SAINTE-MARIE)

R. Sarsaparilla, two ounces.
Mezereon Bark, half a dram.
Spring Water, twenty pounds.

Boil down to two thirds, and add to the strained liquor, a spoonful of alcoholic solution of deuto-chloruret of mercury.

To be taken in the course of the day.—PLENK extolled it highly.

CALLAC'S DECOCTION.

Ptisana Callaci, Decoctum Sarsaparillæ excellentæ. (HISP.; NIEMANN)

R. Sarsaparilla, two ounces.
Proto-chloruret of Mercury, }
in a bag, two drams.
Water, fifteen pounds.

Boil to the consumption of one third, and add

Senna, one ounce.
Seeds of Coriander, six drams.
Alum, half a dram.

Boil again for a quarter of an hour, and strain the decoction before it is cold. (HISP.)

NIEMANN prescribes but two drams of coriander.

It used formerly to be prescribed in syphilis.—Dose, six ounces, twice or thrice a day, for a month or six weeks.

PURGATIVE AND SUDORIFIC PTISAN. (SAINTE-MARIE)

R. Sarsaparilla, four ounces.
Water, three pounds.

Boil slowly for three hours over a small fire, and add towards the end,

Senna, } of each,
Sulphate of Magnesia, } one ounce and a half.
Liquorice, bruised, four drams.

Strain after twenty-four hours infusion.—Dose, six ounces, thrice a day, for three or four weeks. It is said to be very useful in chronic rheumatism.

ZITTMANN'S DECOCTION. (NIEMANN)

R. Sarsaparilla, twelve ounces.
Spring Water, ninety pounds.

Digest for twenty-four hours; then throw into the decoction a bag containing

Styptic Powder (form. of EDINB.), one ounce and a half.

Proto-chloruret of Mercury, half an ounce

Red Sulphuret of Antimony, one dram.

Boil down to thirty pounds, and add towards the end,

Seeds of Anise, } of each,
Fennel, } half an ounce

Senna, three ounces.
Liquorice, one ounce and a half.

Set apart the liquor, under the name of *Strong decoction*. Add to the residue,

Sarsaparilla, six ounces.
Water, ninety pounds.

Boil, and add towards the end,

Lemon Peel, } of each,
Cinnamon, } three drams.
Lesser Cardamoms, }
Liquorice, }

Strain, and set apart the liquor under the name of *Weak decoction*.

This preparation is still more absurd than some of the preceding. THEDEN extolled it in syphilitic diseases. It has been lately brought forward again in France as a novelty.

ANTI-SYPHILITIC ROB. (VAN MONS)

R. Sarsaparilla, fifteen parts.
Water, one hundred and twenty parts.

Boil at three different times, reducing the liquor each time to one half. Mix the three strained decoctions, clarify, reduce the whole to eighteen parts, and add

Senna, } of each,
Flowers of Borage, } one part.
Musk-rose, }
Seeds of Cumin, }

Strain, set aside, decant, and add,

White Sugar, fifteen parts.

Dissolve in a gentle heat, and at the moment of its becoming cold, add

Honey, fifteen parts.

Mix.

SYRUP OF SARSAPARILLA AND OF GUAIAIC.
(AMER. ; COXE)

R. Sarsaparilla, sliced, } of each,
Guaiac, rasped, } one pound.
Water, } ten pounds.

Boil for an hour, and express ; boil the residue with

Water, } ten pounds,
for two hours, adding towards the end,

Petals of Roses, }
Gum Arabic, } of each, one ounce.
Senna, }
Ginger, } half an ounce.

Strain, and add,

Despumated Honey, } of each,
Sugar, } three pounds.

Make a syrup.

This syrup differs from the following only in name :

STEVENS' SYRUP. (NIEMANN)

R. Sarsaparilla, } of each, one pound.
Guaiac Wood, }
Petals of Red Roses, } one ounce.
Water, } a sufficient quantity.

Boil for an hour, and decant. Boil the residue with a similar quantity of water for two hours, adding towards the end,

Gum Arabic, } of each, one ounce.
Senna, }
Ginger, } half an ounce.

Mix the two decoctions, and add

Honey, } of each, three pounds.
Sugar, }

Make a syrup.

At the *Hôpital des Vénériens* at Paris the following formula is followed :

R. Sarsaparilla, } of each, one pound.
Guaiac, }
Water, } twelve pounds.
Honey, } of each,
Sugar, } one pound and a half.

Make a syrup. (RATIER)

CUISINIER'S SYRUP.

Syrupus de Sarsaparilla compositus seu de Sarsaparilla et Senna compositus seu sudorificus seu Cuisinieri seu Smilacis Sarsaparilla. (AMER. ; ANT. ; GALL. ; FORM. HÔP. FR. ; CADET DE GASSICOURT ; COXE ; NIEMANN ; SWEDIAUR)

R. Sarsaparilla, } two pounds.
Warm Water, } twelve pounds.

Infuse for twenty-four hours ; then boil for a quarter of an hour, and express ; boil the residue with

Water, } ten pounds,
till reduced to six pounds. Repeat this ope-

ration two or three times. Mix the decoctions, and boil the whole slowly with

Flowers of Borage, }
White Roses, } of each,
Senna, } two ounces.
Seeds of Anise, }

Boil down to one half, and add to the strained liquor,

Honey, } of each, two pounds.
White Sugar, }

Boil to the consistence of thick syrup. (GALL. ; CADET DE GASSICOURT ; NIEMANN)

HENRI thinks that a prolonged maceration of the sarsaparilla followed by a short boiling, would be sufficient, and that the other substances require only to be infused either separately or in the product of the former operation.

FORM. HÔP. FR. indicates the same process as GALL., but substitutes leaves of borage for the flowers of borage and the roses.—AMER. ; and COXE also suppress the flowers of borage, replacing them by an equal quantity of liquorice, and retaining the roses.—SWEDIAUR directs two pounds of sarsaparilla to be boiled twice in nine pounds of water till reduced to three, one ounce and a half each, of senna, of flowers of borage, of musk roses, and of cumin, to be infused in the two mixed decoctions, the liquor to be strained, and a syrup made with it and two pounds each, of honey and of sugar.—ANT.—two pounds and a half of sarsaparilla to be infused in twenty-four pounds of hot water for twenty-four hours, and then to be boiled till the liquor is reduced to eight pounds ; this operation to be repeated twice again with the residue : twenty-four pounds of the decoction thus obtained to be boiled down to one half, two ounces each, of flowers of borage, of red roses, of senna and of anise, being added towards the end : the syrup to be finally made by evaporating the liquor with two pounds each, of honey and of sugar.

PELLETAN recommends two pounds of sarsaparilla to be boiled several times successively in a sufficient quantity of water ; the decoctions to be reduced together to two pounds : two ounces of senna to be boiled separately in two pounds of water for a quarter of an hour, and two ounces each, of flowers of bugloss, of flowers of borage, of pale roses, and of anise, to be infused in the mixed liquors. A syrup to be finally made with the strained liquor and two pounds each, of honey and of white sugar.

GESNOUIN has recommended the following process :

R. Sarsaparilla, sliced, }
twenty-four pounds.

Boiling Water,
one hundred and seventy pounds.

Infuse for twelve or fifteen hours, then
boil for two, adding, a quarter of an hour
before withdrawing it from the fire,

Senna, one pound eight ounces.

Strain, and pour upon the residue,

Boiling Water, fifteen pounds.

Express, and add

Honey, eighteen pounds.

White Sugar, thirty pounds.

Clarify with white of egg; pour it then
upon

Cumin, bruised, one pound eight ounces.

Boil to the consistence of syrup, and add
to it while boiling,

Extract of Borage, four pounds.

Mix.

KERAUDREN has reduced this formula,
in the following manner:

R. Syrup of Sarsaparilla or of Borage,
twelve pounds.
——— Senna, eight ounces.

Heat in a water-bath and add,

Extract of Borage, one pound.

Rob of Elder, three ounces.

Mix, and strain.

LAGNEAU has reduced it still more:

R. Syrup of Sarsaparilla, four pounds.
——— Senna, three ounces.

Heat, and add,

Extract of Borage, five ounces.
Rob of Elder, one ounce.

Mix.

VINOUS SYRUP OF SARSAPARILLA. (BOR-
RIES; PIERQUIN)

R. Sarsaparilla, one pound and a half.
Root of Long Birthwort, }
——— Round Birthwort, }
——— Jalap, }
——— Polypodium, } of each,
China Root, } six
Guaiac Wood, } ounces.
Sassafras Wood, }
Supertartrate of Potass, }
Sulphuret of Antimony, }
Anise, }
Senna, } twelve ounces.
White Wine, } twelve pounds.

Macerate for six hours in a covered ves-
sel; then add

Water, fifteen pounds.

Boil for an hour over a slow fire; then
decant, and pour upon the residue,

White Wine, six pounds.

Macerate for three hours, and add further,

Water, eight pounds.

Boil for half an hour, and decant. Mix
the two liquors, set apart, decant, and add

Molasses, } of each,
Brown Sugar, } nine pounds.

Boil to the consistence of syrup, and
clarify.

SODIUM.

This metal is not used in medicine, but
the following compounds of it are em-
ployed:

PROTOXIDE OF SODIUM.

Soda caustica seu pura.—Caustic Soda.

I. In the liquid state.

Lixivium causticum seu Sodæ.—Soap-boil-
ers' Ley, Soda Ley, Solution of Soda.

FERR.; FORM. HÔP. FR.; LUSIT.; SAX.;
SWEDIAUR; HENRI; VAN MONS.

R. Subcarbonate of Soda, two parts.
Quick Lime, one part.

Boil with a sufficient quantity of water,
strain the ley, and evaporate till eleven
drams of it fill a vessel which is capable of
containing eight drams of water, or till,
when cold, it marks 36 degrees of the are-
ometer.

VOL. II.

II. In the solid state.

ANT.; FERR.; BRUGNATELLI; VAN
MONS.

R. Soda Ley, any quantity.

Evaporate till it flows like oil, and till a
drop placed on a cold body becomes solid;
then pour it upon a marble slab, or into a
mould.

Caustic soda may serve the same pur-
pose as potass.

WAX SOAP. (VAN MONS)

R. Yellow Wax, three parts.

Melt, and add

Soda Ley, two parts.

Keep it over the fire, constantly stirring,
till the mass becomes susceptible of solu-
tion in water without being decomposed.

4 H

PROTOXI-SULPHURET OF SODIUM.

Sodæ Sulphuretum, Sulphuretum Sodii Protoxidi, Natrum Sulphuratum.

AMER.; FERR.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS.

R. Sublimed Sulphur, } of each,
Subcarbonate of Soda, } an equal part.

Fuse together with a gradually augmented heat, pour the mass upon a marble slab, and break it into small pieces when cold. (AMER.; FERR.; BRUGNATELLI; COXE)

SWEDIAUR prescribes one part of sulphur, and two parts of subcarbonate.

R. Roll Sulphur three parts.
Caustic Soda, seven parts and a half.

Mix together, and fuse in a matrass placed in a sand-bath. (VAN MONS)

This is what VAN MONS calls simple sulphuret of soda; the following formula he gives for the super-sulphuret of soda:

R. Roll Sulphur, three parts and a fourth.
Fused Caustic Soda, four parts.

Fuse together in a glass matrass placed in a gradually heated sand-bath.

The properties of the protoxi-sulphuret of sodium are the same as those of the protoxi-sulphuret of potassium. But the preparation of soda should be preferred, as, according to CHEVALLIER and IDT, "the carbonate of soda varying but little in its degree of purity, the sulphuret may always be depended on for uniformity of composition, whereas that of potass varies from twelve to twenty-five per cent. when prepared with the potass of commerce." However, the sulphuret of soda is very rarely employed.

ANTI-PSORIC LIQUOR. (VAN MONS)

R. Sulphuret of Soda, one dram.
Muriate of Ammonia, two scruples and a half.

Dissolve each salt in six ounces of water, then filter, and mix the solutions.

LOTION AGAINST TINEA. (BORIES; CADET DE GASSICOURT)

R. Sulphuret of Soda, three ounces.
Medicinal Soap, one dram and a half.
Lime Water, seven ounces and a half.
Alcohol, two drams.

Dissolve, and mix.—The head is to be covered with a linen cloth wetted with this lotion, every two days.

SULPHURETTED SOAP OF SODA. (VAN MONS)

R. Marrow Soap, one part.

Sulphuret of Soda, two parts.
Alcohol, six parts.
Dissolve, and strain.

ANTI-PSORIC OINTMENT. (SWEDIAUR)

R. Sulphuret of Soda, three drams.
Lard, one ounce and a half.
Mix by trituration.

SULPHURO-SAPONACEOUS LINIMENT. (VAN MONS)

R. Oil of Almonds, } of each,
Sulphur, } one part and a half.
Sulphuret of Soda, four parts and a half.
Lard, six parts and a half.

Triturate the powders with the lard, and add the oil gradually.

It is employed in itch.

PROTOXI-CHLORURET OF SODIUM.

Eau de javelle, Chlorure de Protoxide de Sodium, Chlorure de Soude. (FEE; HENRI; MAGENDIE)

R. Subcarbonate of Soda, five pounds.
Distilled Water, twenty pounds.

Pour the solution, marking 12 degrees, R., into a flask which it fills only to three fourths, and pass into it a current of chlorine disengaged from a mixture of one pound of hydrochlorate of soda, one pound and a half of sulphuric acid, and twelve ounces of water.

The chloruret is sufficiently strong when eighteen parts of sulphate of indigo are rendered colourless by it.

It is employed, diluted with ten or fifteen parts of water, as an application to hospital gangrene, degenerated syphilitic ulcers, gangrenous wounds, in which it rapidly accelerates cicatrization, and to ulcerated cancer, the fetor of which it destroys at the same time that it calms pain. ALIBERT employs it with advantage as a lotion to spreading herpetic ulceration. It has also been used with success in a case of tinea favosa. It has been given internally in the dose of twenty-five or thirty drops, in one pint of vehicle.

CHLORURET OF SODIUM, MURIATE OF SODA.

Chloruretum Sodii, Hydrochlorinas natricus, Muria, Murias Sodæ, Natrum muriaticum, Sal communis seu marinus seu culinaris, Sal gemme.—Quadri-chloruret of Sodium, Hydrochlorate of Protoxide of Sodium, Hydrochlorate of Soda, Salt.

SYNONIMES:—Quadrichlorure de sodium B., Muriate de Soude, Hydro-chlorate de protoxide de sodium, Proto-hydro-

chlorate de sodium, Hydrochlorate de Soude, Sel de cuisine ou de gabelle, Sel gemme ou marin, Soude muriatée H., FR.; Küchensalz, GERM.; Salt, DAN.; and SWED.; Melh, ARAB.; Uyah, BAL.; Yen, CHIN.; Sunu, CYNG.; Nimmuk, DUK.; Sal, POL.; PORT.; and SPAN.; Lun, HIND.; Zout, DUT.; Sale, ITAL.; Mit, Mah, Garam, MAL.; Hemerk, PERS.; Lavana, SANS.; Uppu, TAM.; Lavanum, TEL.

LOND.; EDINE.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; HERBIP.; WURCEB.; AINSLIE.; GUIBOURT.; PARIS.; PIDERIT.; SPIELMANN.

It is unnecessary to speak of the numerous domestic uses of this compound. It is not extensively used in medicine. It is frequently added to pediluvia to render them more stimulant. It has also, on account of its evidently stimulant properties, been administered internally, in diseases attributed to debility, and externally, in various cutaneous affections, particularly tinea and itch.

RESOLVENT APPLICATION.

Sachet resolutif. (SAINTE-MARIE)

R. Muriate of Soda, } of each,
Sal Gemme, } one ounce.
Muriate of Ammonia, four drams.
Mix.

VERMIFUGE POWDER. (ELLIS)

R. Muriate of Soda, three drams.
Cochineal, fifteen grains.

Make six doses, one to be taken every morning.—This powder is recommended by RUSH.

VERMIFUGE LAVEMENT.

Clyisma ad Ascarides. (SWEDIAUR)

R. Red Wine, six ounces.
Muriate of Soda, a sufficient quantity,
or as much as the wine will dissolve.

PURGATIVE LAVEMENT.

Clyisma purgans. (BRERA; ELLIS; RATIER)

R. Muriate of Soda, one ounce.
Emollient Decoction, two pounds.
Dissolve. (RATIER)
R. Decoction of Barley, four ounces.
Muriate of Soda, half an ounce.

Olive Oil, four ounces.

Mix. (BRERA)

R. Muriate of Soda, half an ounce.
Castor Oil, } of each, two ounces.
Simple Syrup, }
Decoction of Linseed,
one pound and a half.

Mix. (ELLIS)

ANTI-SPASMODIC LAVEMENT.

Clyisma anti-spasmodico-resolvens. (BRERA)

R. Muriate of Soda, four drams.
Linseed Oil, four ounces.
Infusion of Chamomile, three ounces.

Mix.

EXCITANT MIXTURE.

Solutio seu Mixtura citro-muriatica. (BORRIES; CADET DE GASSICOURT; PIERQUIN; SWEDIAUR)

R. Lemon Juice, any quantity.
Muriate of Soda, a sufficient quantity,
or as much as the juice will dissolve.

BROUSSONNET recommends it in adynamic fevers.—Dose, a spoonful every hour.

OINTMENT OF MURIATE OF SODA. (SAINTE-MARIE)

R. Muriate of Soda, one ounce and a half.
GALEN'S Cerate, three ounces.
Rose Ointment, one ounce.

Mix.—To be rubbed, night and morning, on parts affected with tinea.

ARTIFICIAL SEA WATER.

Aqua marina. (GENEV.; AUGUSTIN; BRUGNATELLI; NIEMANN; and SWEDIAUR)

R. Muriate of Soda, half an ounce.
Sulphate of Magnesia, } of each,
Syrup of Buckthorn, } one dram.
Water, two pounds.

Mix. (GENEV.)

R. Muriate of Soda, ten ounces.
Magnesia, ten drams.
Lime, two ounces.
Sulphate of Soda, } of each,
Magnesia, } six drams.
Water, fifty pounds.

Dissolve. (NIEMANN; SWEDIAUR)

BRUGNATELLI gives the same formula but omits the muriate of lime.—AUGUSTIN prescribes two pints of water, twenty-four grains of muriate of soda, four of sulphate of magnesia, fifteen each, of sulphate of lime, of carbonate of magnesia, and of carbonate

of lime, and a current of carbonic acid gas to be passed through the liquor.

This water has been extolled as an excellent resolvent.

SELTZER WATER. (SUEC.)

- R. Muriate of Soda, five grains.
 Subcarbonate of Soda, twenty-four grains.
 ————— Magnesia, twenty-nine grains.
 ————— Lime, seventeen grains.
 Water, eighty-two ounces.
 Saturate with carbonic acid.

PLYMOUTH WATER. (SUEC.)

- R. Muriate of Soda, half a scruple.
 Sulphate of Magnesia, thirty-five grains.
 ————— Lime, thirty-six grains.
 Subcarbonate of Lime, fifteen grains.
 Iron Filings, one ounce.
 Water, eighty-two ounces.
 Saturate with carbonic acid.

SELTZ WATER. (GALL.; VAN MONS)

- R. Muriate of Soda, twenty-two grains.
 Subcarbonate of Soda, four grains.
 ————— Magnesia, two grains.
 Water, charged with five times its volume of carbonic acid gas, twenty ounces and a half.
 Dissolve. (GALL.)
 R. Water, three pounds.
 Pour it into a bottle; introduce to the bottom of this by means of a tube,
 Hydrochloric Acid, seventy grains.
 Then add successively,
 Subcarbonate of Lime, six grains.
 ————— Magnesia, ten grains.
 Neutral Carbonate of Soda, fifty-four grains.
 Stop the bottle well. (VAN MONS)

BALARUC WATER. (GALL.)

- R. Muriate of Soda, one dram and a half.
 ————— Lime, eighteen grains.
 ————— Magnesia, fifty-six grains.
 Subcarbonate of Magnesia, one grain.
 Water, containing twice its volume of carbonic acid, twenty ounces and a half.
 Dissolve.

BOURBONNE-LES-BAINS WATER. (GALL.)

- R. Muriate of Soda, one dram.
 ————— Lime, ten grains.
 Water, charged with twice its volume of carbonic acid, twenty ounces and a half.
 Dissolve.

SUB-PROTO-CARBONATE OF SODIUM.

Soda, Natron, Natrum, Alkali minerale, Carbonas Sodæ, Subcarbonas Sodæ, Hypercarbonas Sodæ, Natrum carbonicum, Subcarbonas natrium, Carbonas Sodæ alkalinus seu incompletus, Hypercarbonas Sodæ. — Subcarbonate of Protoxide of Sodium, Subcarbonate of Soda, Fixed mineral alkali.

1. *That which is found in commerce.*

Soda, Barilla, Carbonas natrium seu Soda impurum. — Soda, Barilla.

LOND.; DUBL.; EDINB.; AMER.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENÈV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; COXE.; GUIBOUT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.

Four sorts of soda are distinguished in commerce.

A. *Native Soda*, in fine needles, or in pulverulent, snowy efflorescences. It is tolerably abundant in Egypt, in the desert of Thiaïat, to the west of Delta, and in the plains of Upper Hungary.

B. That which is obtained by the incineration of a great number of marine plants, particularly the *Atripliceæ* and *Fucacææ*, as *Salsola*, *Chenopodium*, *Atriplex* and *Fucus*. It is in solid, compact, semifused masses, of the hardness of stone, of a bluish-grey colour, and of a saline and lixivial taste. It contains subcarbonate of soda, in very variable quantity, sulphate of soda, muriate of soda, subcarbonate of lime, alumine, silex, and iron, sometimes also sulphate of potass, and chloruret and even ioduret of potassium.

C. That which is obtained by the decomposition of muriate of soda by litharge.

D. That which results from the decomposition of sulphate of soda with charcoal and chalk. It is composed of caustic soda, of subcarbonate of lime, of sulphuret of lime with excess of base, and of charcoal.

2. *Purification of the Soda of Commerce.*

Sodæ Subcarbonas. (LOND.)

- R. Impure Soda, in powder, one pound.
 Boiling Distilled Water, four pints.

Boil together for half an hour, and filter the solution; evaporate to one half, and set it aside that crystals may form. Reject the residuary liquor.

Subcarbonas Sodæ. (EDINB.)

- R. Impure Subcarbonate of Soda, any quantity.

Bruise it, and then boil it in water till all the saline matter is dissolved. Filter the solution and evaporate it in an iron vessel, so that crystals may form on cooling.

Sodæ Carbonas. (DUBL.)

- R. Barilla, in powder, one part.
Water, two parts.

Boil together in a covered vessel for two hours, occasionally stirring, and filter; then bruise the barilla that remains with an equal quantity of water, and again boil it: this may be repeated a third time. Having mixed and filtered the liquors, evaporate to dryness, in a shallow iron vessel, taking care that the saline mass which will remain be not liquified by too great a heat: stir it with an iron spatula till it becomes white; then dissolve it in boiling water, and after due evaporation, set the solution aside, that crystals may form as it cools.

These will be the purer if the barilla before each boiling be exposed for some time to the air. The crystallization should be effected when the air is at the freezing temperature, and with a solution the specific gravity of which is, to that of water, as 1220 is to 1000. If the salt is not very pure, repeat the solution and crystallization.

- R. Soda of Commerce, any quantity.

Having pulverized it, dissolve it in water, evaporate the solution till it marks 28 or 30 degrees by the areometer, and set it apart to crystallize.

This process is the same for all the other pharmacopœias.

DRIED SUBCARBONATE OF SODA.

Sodæ Subcarbonas exsiccata. (LOND.)

- R. Subcarbonate of Soda, one pound.

Expose it to a boiling heat in a clean iron vessel till it becomes per-

fectly dry, stirring it assiduously with an iron spatula. Finally, reduce it to powder.

Sodæ Carbonas siccatum. (DUBL.)

- R. Crystals of Carbonate of Soda, any quantity.

Liquify them in a silver crucible over a fire; then, having augmented the heat, stir the fused salt till, by the evaporation of the water of crystallization, the salt has become dry. Reduce it to fine powder, and preserve it in stopped bottles.

The subcarbonate of soda is prescribed in scrofula, passive dropsies, engorgements of the abdominal viscera, and in cases of urinary calculi and gravel the base of which is formed by uric acid.—Dose, from ten to twenty grains.

SOLUTION OF SUBCARBONATE OF SODA.

Sodæ Carbonatis Aqua. (DUBL.)

- R. Carbonate of Soda, any quantity.

Dissolve it in distilled water, and let the specific gravity of the solution be to distilled water as 1024 is to 1000.

A solution of this specific gravity may be made by dissolving one ounce of the carbonate in one pint of distilled water

STOMACHIC BOLUSES.

Boli stomachici. (VAN MONS)

- R. Subcarbonate of Soda, half a scruple,
Aromatic Powder, four scruples.
Syrup of Cinchona, a sufficient quantity.

Make three boluses.

LITHONTRIPTIC PILLS.

Pilulæ Subcarbonatis Sodæ. (EDINB.; AMER.; AUGUSTIN; COXE; ELLIS; NIEMANN)

- R. Dried Subcarbonate of Soda, four parts,
Hard Soap, three parts.
Simple Syrup, a sufficient quantity.

Make a pilular mass.

Dose, a scruple every two hours.

AMER.; COXE; and ELLIS prescribe one dram of subcarbonate and half a dram of soap; to be made into forty pills.

ABSORBENT PILLS. (ELLIS)

- R. Subcarbonate of Soda, } of each,
 Rhubarb, } one scruple.
 Extract of Gentian, }
 Proto-chloruret of Mercury, } three grains.

Make twenty pills.—Dose, two; in dyspepsia.

ALKALINE OINTMENT. (RATIER)

- R. Subcarbonate of Soda, two drams.
 SYDENHAM'S Laudanum, one dram.
 Lard, one ounce.

Mix.—CULLERIER employed it in various cutaneous affections, when there was much pain.

- R. Subcarbonate of Soda, } of each,
 Sulphuret of Potass, } three drams.
 Lard, } three ounces.

Mix.—ALIBERT directed it to be rubbed upon the head in tinea every day, after having removed the crusts or scales with emollient cataplasms.

ANTI-NEPHRITIC LIQUOR.

Aqua anti-nephritica. (FULD.; HAMB.; LIPP.; NIEMANN)

- R. Subcarbonate of Soda, two drams.
 Lime Water, two pounds.
 Dissolve.—Dose, two or three spoonfuls.

ABSORBENT MIXTURE.

Mixtura antacida. (ELLIS; SWEDIAUR)

- R. Subcarbonate of Soda, } of each,
 White Sugar, } two drams.
 Yolks of Eggs, } two.
 Honey, } two drams.

Mix. (SWEDIAUR)

- R. Subcarbonate of Soda, four scruples.
 Infusion of Gentian, } of each,
 Cinnamon Water, } three ounces.
 Tincture of Cardamoms, } one ounce and a half.

Mix.—Dose, a spoonful every two or three hours. (ELLIS)

- R. Subcarbonate of Soda, one dram.
 White Sugar, half an ounce.
 Mucilage of Gum Arabic, two ounces.
 Water, thirty ounces.
 Spirit of Pimenta, two ounces.

Mix.—Dose, two ounces, three or four times a day. (SWEDIAUR)

- R. Subcarbonate of Soda, two scruples.
 Rhubarb, two drams.
 Simple Syrup, one ounce.
 Mint Water, eight ounces.

Dissolve, and mix.—Dose, two spoonfuls, thrice a day. (SWEDIAUR)

DIGESTIVE POTION.

Infusum amarum cum Natro. (AUGUSTIN)

- R. Root of Gentian, two drams.
 Unripe Orange, one dram and a half.
 Lesser Cardamoms, half a dram.
 Subcarbonate of Soda, one dram.
 Boiling Water, six ounces.

Infuse for six hours, and strain.—Dose, a spoonful, thrice a day.

ANTI-SCROFULOUS POTION. (RATIER)

- R. Subcarbonate of Soda, half a dram.
 Tincture of Gentian, half an ounce.

Dissolve.

PESCHIER has employed in goitre for many years past, a more or less concentrated solution of subcarbonate of soda; from two to four drams of the salt in eight ounces of water, of which he gives twice a day in half a glassful of wine or sweetened aromatic water. HUFELAND prescribes four spoonfuls daily of a solution of two drams of subcarbonate in six ounces of balm water and half an ounce of cinnamon water.

POTION AGAINST URINARY GRAVEL. (SAINTE-MARIE)

- R. Linseed, bruised, } of each,
 Montpellier Maidenhair, } one pinch.
 Iceplant Herb, }
 Water, } one pound.

Boil for a quarter of an hour, and add to the strained liquor

Subcarbonate of Soda, one scruple.

Dissolve.—To be taken in the course of the day.

ALKALINE BALSAMIC SOLUTION.

Solutio balsamica alkalisata. (SWEDIAUR)

- R. Subcarbonate of Soda, } of each,
 Sugar, } one ounce
 Myrrh, } and a half.
 Peruvian Balsam, } half an ounce.
 Canadian Balsam, } ten drams.
 Water, } one pound and a half.

Mix.—One ounce of this solution is to be mixed with a pound and a half of water, and taken in the course of the day.

LITHONTRIPTIC INJECTION. (BORIES)

- R. Subcarbonate of Soda, one dram.
 Soap, two ounces.
 Water, twelve ounces.

Mix.—It is recommended as a solvent for urinary calculi formed of uric acid.

LITHONTRIPTIC POTION. (ELLIS)

- R. Subcarbonate of Soda, one dram.
 Infusion of Quassia, four ounces.
 Tincture of Calumba, one ounce.

Dissolve.—Dose, a spoonful four times a day.

EXPECTORANT POTION. (ELLIS)

- R. Subcarbonate of Soda, twelve grains.
 Wine of Ipecacuanha, fifteen drops.
 Tincture of Opium, four drops.
 Water, one ounce.

Dissolve.—Dose, a tea-spoonful, every two or three hours; in hooping cough.

STOMACHIC LIQUOR.***Infusum Quassiae cum Natro.* (AUGUSTIN)**

- R. Quassia Wood, one dram and a half.
 Subcarbonate of Soda, two drams and a half.
 Boiling Water, one pound.

Infuse for an hour, and strain.

Dose, from two to four spoonfuls, daily, in dyspepsia.

PROTO-CARBONATE OF SODIUM.

Carbonas Sodæ seu natrium, Bicarbonas Sodæ.—Carbonate, or Bicarbonate, or Neutral Carbonate, of Soda, Carbonate of Protoxide of Sodium.

1. In the solid state.

LOND.; DUBL.; EDINB.; BATAV.; BAVAR.; BELG.; FENN.; SUEC.

***Sodæ Carbonas.* (LOND.)**

- R. Subcarbonate of Soda, one pound.
 Distilled Water, three pints.

Dissolve; and transmit carbonic acid through the solution by means of a proper apparatus till it is saturated; then set it apart to crystallize. Dry the crystals, folded and compressed in blotting paper. Evaporate the remainder of the solution with a heat not exceeding 120 degrees, that more crystals may be obtained. Dry these in the same manner as the former.

***Carbonas Sodæ.* (EDINB.)**

- R. Subcarbonate of Soda, two parts.
 Water, three parts.

Dissolve, and submit the solution to a current of carbonic acid gas till this ceases to be absorbed. Filter the liquor, and evaporate it with a heat not exceeding 180 degrees, that it may crystallize.

The carbonic acid may be obtained from equal weights of pulverized carbonate of lime and of sulphuric acid largely diluted with water.

***Sodæ Bicarbonas.* (DUBL.)**

- R. Carbonate of Soda, two parts.
 Water, three parts.

Dissolve; and by means of a proper apparatus, pass through the solution a current of carbonic acid gas, extricated from white marble by means of diluted hydrochloric acid, till crystals form: then, with a heat not exceeding 120 degrees, evaporate the liquor to obtain more crystals. Dry the crystals and preserve them in well-stopped bottles.

- R. Pure Subcarbonate of Potass, one part.
 Hot Water, two parts.

Pass a current of carbonic acid gas into the filtered and cold solution, and preserve the crystals which form spontaneously. (AMER.; BATAV.; BAVAR.; BELG.; FENN.; SUEC.; COXE)

- R. Subcarbonate of Soda, } of each,
 Water, } one pound.

Add to the solution

Subcarbonate of Ammonia, three ounces.

Apply to the mixture for three hours, a heat of 80 degrees C., till all the ammonia is driven off; then set aside the liquor to crystallize. (AMER.; COXE)

2. In the liquid state.

Aqua Supercarbonatis Sodæ seu Carbonatis Sodæ acidula. (EDINB.; FENN.; HANN.; POL.; ROSS.; SUEC.; NIEMANN; SWEDIAUR; VAN MONS)

***Aqua Supercarbonatis Sodæ.* (EDINB.)**

- R. Water, ten pounds.
 Subcarbonate of Soda, two ounces.

Disssolve; and submit the solution to a stream of carbonic acid gas pro-

cured from three ounces of carbonate of lime and the same quantity of sulphuric acid carefully diluted with three pounds of water. It may be well prepared in NOOTH'S apparatus, but if a large quantity is required, an apparatus capable of affording greater pressure will be necessary.

Preserve the liquor in well corked bottles.

Aqua Carbonatis Sodæ acidula.
(DUBL.)

R. Subcarbonate of Soda,
any quantity.

Dissolve it in water in such proportion that each pint may contain one dram of the subcarbonate: then in a proper apparatus pass through the solution a stream of carbonic acid gas, extracted from white marble by means of hydrochloric acid diluted with six times its weight of water, till the carbonic acid is in excess in the liquor.

The other pharmacopœias give the same process as EDINB.

This preparation, which is in very extensive use, does not sensibly differ from the preceding therapeutically.

GASEOUS ALKALINE WATER; SODA WATER.

Aqua Sodæ. (AMER.; COXE; ELLIS)

R. Subcarbonate of Soda, three drams.
Water, eight pounds.

Charge the solution with ten times its volume of carbonic acid gas. (AMER.; COXE)

R. Subcarbonate of Soda, eighteen grains.
Hydrochloric Acid, half a dram.
Water, twelve ounces.

Preserve in a well stopped bottle. (RATIER)

FEBRIFUGE POTION. (BORIES)

R. Distilled Water, eight ounces.
Carbonic Acid, a sufficient quantity
to make a volume double that of the water.
Add to the liquor

Subcarbonate of Soda, half a dram.
Camphor, twenty grains.
Syrup of Poppy, six drams.

Mix.—One half to be taken four hours before the paroxysm, and the remainder at the moment of its coming on. Its use should be continued for seven or eight days.

DIGESTIVE TROCHISCS. (FEE; HENRI;
MAGENDIE)

R. Bicarbonate of Soda, two hundred and fifty parts.
White Sugar, four thousand
seven hundred and fifty parts.
Essential Oil of Mint, eight parts.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs of eighteen grains each.
—Dose, two or three, before each meal, in
calculous and gouty cases.

SUB-PROTO-BORATE OF SODIUM.

Boras Sodæ alkalescens, Borax, Borax Venetus, Hypoboras Sodæ, Sub-boras natricum seu Sodæ.—Sub-borate of Soda, Borate of Soda, Borax, Tinkal, Sedative Salt.

SYNONIMES:—Souborate de Protoxide de Sodium, Sous-borate de Soude, Borate sur-saturé de Soude, Borate de Soude avec excès de base, Soude boratée, H., Borate de Soude, B., Borax, Borax brut, Tinkal, Tinkar, Fr.; Naturliches Bedativsalz, GERM.; Buruk, ARAB.; Piger, BAL.; JAV.; and MAL.; Sansipuscara, CYNG.; Sohaga, DUK.; and HIND.; Borace, ITAL.; Chautarya, NEP.; Tunkar, PERS.; Borax, POL.; and SWED.; Tincal, PORT.; Tunkana, SANS.; Velligarum, Vengarum, TAM.

LOND.; EDINB.; DUBL.; AMER.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENÈV.; HAMB. HANN.; HASS.; LIPP.; OLDEN.; CAST. BORUS.; BORUS.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; WURCEB.; AINSLIE; GUIBOUT; PARIS; SPIELMANN.

In hexahedral prisms, more or less depressed, and terminated by dihedral pyramids; it is sometimes colourless, sometimes yellowish or greenish, always covered with an earthy coating: it is unctuous to the touch, of a saponaceous odour and a weak alkaline taste; its fracture is vitreous, and it is efflorescent.

SPECIES DIABORACIS. (WIRTEM.)

R. Venice Borax, half an ounce.
Cassia Lignea, } of each,
Turmeric, } one dram.
Savine Wood, rasped, } of each,
Prepared Amber, } half a dram.
Bone of Stag's Heart, } of each,
Miseltoe, } one scruple.
Leaves of Wallflower, }

Make a powder.—MYSNIGHT recom-

mended it to accelerate parturition.—Dose, from a scruple to half a dram.

PULVIS AD PARTUM PROMOVENDUM.
(PALAT.; WIRTEM.)

- R. Sub-borate of Soda, half an ounce.
Castor, } of each,
Saffron, } one dram and a half.
Oil of Cinnamon, eight drops.
— Amber, six drops.

Mix, and make a powder. (WIRTEM.)

- R. Cassia Lignea, } of each,
Cinnamon, } two drams.
Sub-borate of Soda, } of each,
Prepared Amber, } one dram.
Saffron, half a dram.
Oil of Cinnamon, three drops.

Make a powder. (PALAT.)

Dose, from one scruple to half a dram.

MIXTURA AD PARTUM. (SWEDIAUR)

- R. Sub-borate of Soda, two drams.
Water of Chamomile, four ounces.
Syrup of Orange Peel, half an ounce.
Solution of Succinate of Ammonia, two drams.
SYDENHAM'S Laudanum, half a dram.

Dissolve, and mix.—Dose, two spoonfuls every half hour.

ASTRINGENT POTION.

Infusum Salep cum Adstringentibus. (AUGUSTIN)

- R. Salep, one dram.
Boiling Water, ten ounces.

Add to the strained liquor,

- Extract of Cinchona, } of each,
— Logwood, } one dram.
Sub-borate of Soda, half a dram.
Syrup of Marshmallows, one ounce.

Dissolve.—To be taken by spoonfuls in chronic diarrhoea, and particularly in asthenic and malignant aphthae.

- R. Salep, two drams.
Linseed, half a dram.
Boiling Water, ten ounces.

Infuse, and strain. Add

- Sub-borate of Soda, half a dram.
Extract of Logwood, } of each,
— Cinchona, } one dram.
Syrup of Marshmallows, one ounce.

Mix.—To be taken by spoonfuls, in the same cases as the preceding.

BORATED LINCTUS.

Collutorium adstringens, Linimentum de Borate seu ad Aphthas, Mel Boracis seu boraxatum seu Sub-boratis Sodæ, Linctus de Borace seu ad Aphthas. (LOND.; DUBL.;

MIL. COPEN.; FORM. HÔP. FR.; HAMB.; SUEC.; AUGUSTIN; COXE; PIERQUIN; SAUNDERS; SWEDIAUR; PH. SYPH.; VAN MONS)

Mel Boracis. (LOND.; DUBL.)

- R. Sub-borate of Soda, in powder, one dram.

Despumated Honey, one ounce.

Mix.

HAMB.; and COXE give the same formula.—FORM. HÔP. FR. prescribes one part of borax, and ten parts of honey.—PIERQUIN; and VAN MONS—one of borax, and twelve of honey.—AUGUSTIN—an equal part of each.

- R. Sub-borate of Soda, one dram.
Oil of Almonds, one ounce.
Yolk of Egg, one.

Triturate together. (SAUNDERS; SWEDIAUR)

- R. Sub-borate of Soda, one or two drams.
Despumated Honey, one ounce.
Extract of Walnut Rind, one or two drams.

Triturate together. (SWEDIAUR)

- R. Sub-borate of Soda, one dram.
Syrup of Mulberries, from one to three ounces.

Mix. (AUGUSTIN)

- R. Sub-borate of Soda, one or two drams.
Mucilage of Quince, } of each,
Syrup of Mulberries, } one ounce.

Mix. (PH. SYPH.; SWEDIAUR)

- R. Sub-borate of Soda, one dram and a half.
Spring Water, } of each, one ounce.
Honey, }

Triturate together. (MIL. COPEN.; SUEC.)

BORATED GARGLE.

Gargarisma e Borace seu ad Aphthas seu boraxatum, Mixtura boraxata, Liquor ad Aphthas: (MIL. COPEN.; GENEV.; HAMB.; AUGUSTIN; BREERA; CADET DE GASSICOURT; ELLIS; PIERQUIN; RATIER; SAUNDERS; SWEDIAUR; PH. SYPH.)

- R. Sub-borate of Soda, two drams.
Mucilage of Quince, half a pound.
Honey of Roses, one ounce.

Mix. (GENEV.; SAUNDERS)

SWEDIAUR directs one ounce of quince seeds to be boiled in sixteen ounces of water till reduced to twelve, and three drams of borax and one ounce and a half of honey of roses to be added to the strained liquor.

- R. Sub-borate of Soda, six drams.
 Infusion of Sage, ten ounces.
 Despumated Honey, two ounces.

Mix. (HAMB.)

MIL. COPEN. prescribes two drams of borax, eight ounces of infusion of sage, and one ounce of honey.—BRERA—one scruple of borax, one pound of decoction of sage, and one ounce of proof spirit.

- R. Sub-borate of Soda, two drams.
 Water, } of each,
 Tincture of Myrrh, } one ounce.
 Honey of Roses, two ounces.

Mix. (CADET DE GASSICOURT ; SWEDIAUR)

AUGUSTIN ; and PH. SYPH. prescribe one ounce of borax, one pound of water, and two ounces each, of tincture of myrrh and of honey.—ELLIS—one dram of borax, four ounces of water, half an ounce of tincture of myrrh, and one ounce of honey.

- R. Sub-borate of Soda, two drams.
 Decoction of Barley, one pound and a half.
 Syrup of Gum, one ounce.

Mix. (RATIER)

- R. Sub-borate of Soda, two drams.
 Infusion of Roses, ten ounces.
 Syrup of Mulberries, half an ounce.

Mix. (BRERA)

- R. Sub-borate of Soda, three drams.
 Decoction of Cinchona, twelve ounces.
 Honey, one ounce and a half.

Mix. (RATIER)

- R. Sub-borate of Soda, two or three drams.
 Honey, one ounce.
 Extract of Walnut, half an ounce.
 Alcohol, one or two ounces.
 Boiling Water, one pound.

Mix. (SWEDIAUR)

- R. Sub-borate of Soda, } of each,
 Sulphate of Copper, } eight grains.
 Tincture of Cochineal, one dram.
 Water, one ounce.
 Spirit of Scurvy-grass, two ounces.
 Syrup of Raspberries, a sufficient quantity.

Mix. (PIERQUIN)

BORATED COLLYRIUM.

Collyrium boracicum. (AUGUSTIN)

- R. Sub-borate of Soda, half a dram.
 Sugar, one dram.
 Rose Water, two ounces.

Mix.

COSMETIC LIQUOR. (AUGUSTIN)

- R. Almonds, two ounces.

- Rose Water, } of each,
 Orange-flower Water, } eight ounces.
 Make an emulsion, and add

Tincture of Benzoin, two drams.
 Sub-borate of Soda, one dram.

Mix.

SUB-PROTOPHOSPHATE OF SODIUM.

Phosphas Sodæ seu Natricum, Soda phosphorata, Natrum phosphoricum seu oxyphosphorodes, Alkali minerale phosphoratum.—Phosphate of Soda, Sub-phosphate of Soda, Sub-phosphate of Protoxide of Sodium. (DUBL. ; EDINB. ; AMER. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; DAN. ; GALL. ; FERR. ; FENN. ; HANN. ; OLDEN. ; POL. ; BORUS. ; SAX. ; SUEC. ; BRUGNATELLI ; COXE ; SWEDIAUR ; VAN MONS)

Phosphas Sodæ.

- R. Bones burnt to whiteness and reduced to powder, ten pounds.
 Sulphuric Acid, six pounds.
 Subcarbonate of Soda, a sufficient quantity.

Mix together in an earthen vessel, and add nine pounds of water. Keep the mixture in a vapour-bath for three days, after which add nine pounds more of boiling water, and strain through a strong linen cloth, pouring boiling water gradually upon it till all the phosphoric acid is washed out. Set the strained liquor apart that the feculencies may subside ; then having poured it off, evaporate it to nine pounds. To this liquor, heated in an earthen vessel, add a warm solution of subcarbonate of soda till the effervescence ceases ; then strain, and set the liquor aside that crystals may form. Having removed these, add to the liquor, if necessary, a little subcarbonate of soda, that the phosphoric acid may be entirely saturated, and dispose it, by repeated evaporation, to yield crystals, till they are no longer produced. Finally, preserve the crystals in a well-stopped vessel. (EDINB.)

- R. Burnt Bones, in powder, five pounds.

Sulphuric Acid of commerce,
three pounds and a half.

Mix in an earthen vessel, add gradually five pints of water, and stir the mixture. Digest for three days, occasionally stirring, and adding water to prevent the ingredients from becoming dry; then pour over them five parts of boiling water, and strain through a linen cloth, pouring on boiling water, at intervals, till all the acid is washed out. Set apart the liquor that the impurities may subside, then pour it off and evaporate it to one half; then add

Carbonate of Soda,
three pounds ten ounces,

dissolved in a sufficient quantity of hot water. Filter, and obtain crystals by repeated evaporation and refrigeration. Preserve the crystals in a well-stopped vessel. If the salt is not perfectly pure, the solution and crystallization may be repeated. (DUBL.)

R. Super-sulphate of Lime, any quantity.
Subcarbonate of Soda,

a sufficient quantity

to saturate perfectly the acid. Pour off the clear liquor, evaporate it to 25 degrees of the areometer, and set it apart to crystallize. (AMER.; BATAV.; BAVAR.; BELG.; GALL.; PENN.; FERR.; SUEC.; COXE; VAN MONS)

R. Phosphoric Acid, one part.
Water, eight parts.

Pour gradually into the mixture,
Solution of Subcarbonate of Soda,
a sufficient quantity;

or, till the alkali remains in slight excess. Filter the liquor, evaporate, and crystallize. Preserve the crystals in a very dry place. (ANT.; BATAV.; DAN.; HANN.; OLDEN.; POL.; BORUS.; SAX.; BRUGNATELLI; SWEDIAUR)

It is mildly purgative.—Dose, from half an ounce to an ounce and a half, in broth.

LAXATIVE POTION.

Mixtura Nitri phosphorici. (AUGUSTIN)

R. Phosphate of Soda, } of each,
Syrup of Marshmallows, } half an ounce.
Water, four ounces.

Mix.—To be taken in divided doses, at intervals of an hour.

PHOSPHATE OF SODA AND OF AMMONIA.
(BAVAR.)

R. Sub-phosphate of Soda, six parts.
Hot Distilled Water, two parts,
holding in solution,

Hydrochlorate of Ammonia, one part.

Dissolve over a slow fire. Set apart the solution in a cool place for twenty-four hours; then remove and dry the crystals which are formed.

SUPER-PROTO-SULPHATE OF SODIUM.

Sulphas Sodæ acidulus. (VAN MONS)

R. Sulphuric Acid, } of each, four ounces.
Water,

Add to the mixture,

Subcarbonate of Soda, eight ounces.

Filter, and crystallize.

It is purgative.—Dose, from two to six drams.

PROTO-SULPHATE OF SODIUM.

Alkali minerale sulphuricum seu vitriolatum, Natrum sulphuricum, Sal catharticum seu mirabile Glauberi, Soda vitriolata, Sulphas natricus seu Sodæ.—Sulphate of Soda, Glauber's Salt.

1. *That which is met with in commerce.*

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; MIL. COPEN.; HISP.; GALL.; FENN.; FULD.; HAMB.; LIPP.; OLDEN.; CAST. BORUS.; SAX.; SUEC.; WURCEB.; COXE; GUIBOUT; PARIS.

It is in long six-sided prisms terminated by a dihedral summit, transparent, colourless, efflorescent, at first of a refreshing taste then bitter.

2. *Prepared according to various formulæ.*

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; GALL.; FERR.; GENEV.; HANN.; HASS.; OLDEN.; LUSIT.; BORUS.; ROSS.; SARD.; WIRTEM.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS.

Sodæ Sulphas.

R. Residuary Salt after the distillation of Muriatic Acid,
two pounds.

Boiling Water,

two pints and a half.

Dissolve, and gradually add,

Subcarbonate of Soda,

a sufficient quantity for the saturation of the acid. Boil the liquor till a pellicle appears on its surface; then having filtered it set aside to crystallize. Pour off the water, and dry the crystals in bibulous paper. (LOND.)

Dissolve in water the acidulous salt remaining after the distillation of muriatic acid, and saturate the superfluous acid by the admixture of carbonate of lime in powder; then set apart that the impurities may subside. Pour off the clear liquor, filter it, and evaporate that crystals may form. (EDINB.)

Dissolve the residuary salt after the distillation of muriatic acid in a sufficient quantity of boiling water. Evaporate the solution to a proper degree of concentration, and set it apart, that crystals may form as it slowly cools. (DUBL.)

AMST. ; AUSTR. ; BATAV. ; BRUNS. ; DAN. ; FERR. ; GENEV. ; HANN. ; HASS. ; OLDEN. ; LUSIT. ; BORUS. ; ROSS. ; SARD. ; WIRTEM. ; BRUGNATELLI ; PIDERIT ; SPIELMANN ; SWEDIAUR ; and VAN MONS give the same formula as LOND.

R. Subcarbonate of Soda, any quantity.

Dissolve it in water, and pour into the liquor,

Sulphuric Acid, a sufficient quantity to saturate the alkali. Evaporate the solution, and set it aside to crystallize. (AMST. ; ANT. ; BATAV. ; BELG. ; GALL. ; FERR.)

It forms a mild purgative in the dose of from an ounce to an ounce and a half. In that of from three to five scruples, it is diuretic. It is also prescribed as a deobstruent.

PURGATIVE POTION. (FORM. HÔP. FR. ; HAMB. ; CAST. BORUS. ; AUGUSTIN ; NIEMANN ; RATIER ; SAUNDERS)

R. Sulphate of Soda, sixteen parts.
Water, five hundred parts.

Dissolve. (FORM. HÔP. FR.)

R. Sulphate of Soda, one ounce.
Water, eight ounces.
Honey, half an ounce.

Dissolve. (CAST. BORUS.)

R. Sulphate of Soda, half an ounce.
Nitrate of Potass, one dram.
Lemon Juice, one ounce and a half.
Syrup of Raspberries, two ounces.

Water of Raspberries, ten ounces.

Mix. (SAUNDERS)

R. Sulphate of Soda, one ounce.
Sugar, half an ounce.
Lemon Juice, one ounce and a half.
Gruel, one pound.

Dissolve.—A glassful to be taken every hour. (AUGUSTIN)

R. Sulphate of Soda, one ounce.
Peppermint Water, } of each,
Infusion of Rhubarb, } three ounces.
Oxymel of Squill, one ounce.

Dissolve. (HAMB.)

R. Senna, three drams.
Sulphate of Soda, two drams.
Manna, one dram.
Boiling Water, four ounces.

Infuse, and strain. (RATIER)

R. Pulp of Tamarinds, one ounce and a half.

Water, a sufficient quantity to obtain four ounces of decoction. Add, when strained,

Sulphate of Soda, one ounce and a half.
Syrup of Pale Roses, six drams.

Mix. (SAUNDERS)

R. Sulphate of Soda, one ounce and a half.
Manna, one ounce.
Water, six ounces.

Dissolve.—To be taken by spoonfuls. (AUGUSTIN)

R. Sulphate of Soda, two drams.
Magnesia, one ounce.
Infusion of Orange Peel, eight ounces.
Red Wine, a small quantity.

Dissolve. (NIEMANN)

PURGATIVE LAVEMENT. (LIPP. ; PH. SYPH. ; RATIER ; SAUNDERS)

R. Sulphate of Soda, one ounce.
Linseed Oil, two ounces.
Decoction of Barley, six ounces.

Mix. (PH. SYPH.)

R. Emollient Species, two ounces.
Senna, half an ounce.
Sulphate of Soda, two drams.
Boiling Water, one pound.

Infuse and strain. (RATIER)

RATIER also prescribes two drams of senna, half an ounce of sulphate, and two pounds of emollient decoction; or, half an ounce each, of senna and of sulphate, and one pound of water.—LIPP.—eight ounces of emollient decoction, three ounces of oxymel, and one ounce of sulphate.

R. Herb of Rue, } of each,
Flowers of Chamomile, } half
Elder, } an ounce.
Water, one pound.

Boil down to two thirds; then strain, and add,

Sulphate of Soda,	three drams.
Oxymel,	of each,
Oil of Chamomile,	one ounce.

Mix. (SAUNDERS)

ALTERATIVE AND TONIC APOZEM. (PIERQUIN)

R. Root of Dock,	one ounce.
——— Strawberry,	half an ounce.
——— Elecampane,	one dram.
Leaves of Cichory,	of each,
——— Chervil,	one handful.
——— Cress,	
Sulphate of Soda,	one dram.
Compound Syrup of Cichory,	one ounce.

Boil and mix.—To be taken every morning.

ALTERATIVE BROTH. (PIERQUIN)

R. Chicken Broth, one pound and a half.
Root of Dock, two ounces.

Boil for eight minutes. Add towards the end,

Leaves of Cress,	of each,
——— Balm,	one handful.
——— Cowslip,	

Strain, and add,

Sulphate of Soda, two ounces.

Dissolve.

APERIENT APOZEM. (GALL.; CADET DE GASSICOURT)

R. Fresh Leaves of Cichory,	of each,
——— Bugloss,	one ounce.
——— Borage,	
Boiling Water,	two pounds.

Infuse for a quarter of an hour; then strain, and add

Sulphate of Soda,	one dram.
Syrup of Violets,	one ounce.

Mix.

STIMULANT LOTION.

Lotio ad Tineam. (SWEDIAUR)

R. Sulphate of Soda,	six ounces.
White Soap,	three drams.
Lime Water,	eighteen ounces.
Proof Spirit,	six drams.

Mix.

SALINE GELATINOUS SOLUTION. (CADET DE GASSICOURT)

R. Water,	one pound.
Subcarbonate of Soda,	of each,
Flemish Glue,	one ounce.
Sulphate of Soda,	of each,
Muriate of Soda,	half an ounce.

Petroleum, five drops.
Dissolve, and mix.

CATHARTIC DIURETIC POWDER. (VAN MONS)

R. Tartarized Antimony,	of each,
Opium,	one grain.
Sulphate of Soda,	two ounces.
Gum Arabic,	of each, two drams.
Liquorice,	

Make a powder.—To be taken three or four times a day in water.

GUINDRE'S SALT. (CADET DE GASSICOURT, NIEMANN)

R. Sulphate of Soda,	one ounce.
Nitrate of Potass,	twelve grains.
Tartarized Antimony,	half a grain.

Mix.—It is purgative. To be taken in the morning dissolved in broth or any other convenient vehicle.

RESOLVENT LIQUOR. (CADET DE GASSICOURT)

R. Sulphate of Soda,	one or two ounces.
Nitrate of Potass,	ten grains.
Tartarized Antimony,	half a grain.
Water,	two pounds.

Dissolve.

PURGATIVE AND EMETIC LIQUOR.

Eau minérale. (FORM. HÔP. FR.; RATIER)

R. Sulphate of Soda,	three drams.
Tartarized Antimony,	three grains.
Warm Water,	ten ounces.

Dissolve.

CATHARTIC POTION. (CADET DE GASSICOURT)

R. Sulphate of Soda,	of each,
Manna,	half an ounce.
Tartarized Antimony,	two grains.
Water,	four ounces.
Oxymel of Squill,	one ounce.

Dissolve.—Dose, a spoonful, every two hours.

SULPHATE OF SODA AND OF MAGNESIA.

Sulphas Sodæ et Magnesiae, Sal Astracanen-sis artificialis. (VAN MONS)

R. Supersulphate of Soda, any quantity,
Warm Water, a sufficient quantity
to dissolve the salt exactly. Saturate it with

Calcined Magnesia,

a sufficient quantity,

Filter the liquor, and crystallize.

It is laxative or diuretic, according to the

dose.—From half a dram to two drams are prescribed several times a day, or from one to two ounces at once.

PROTO-SULPHITE OF SODIUM. (VAN MONS)

- R. Subcarbonate of Soda, one part.
Water, four parts.

Dissolve; saturate the solution with sulphurous acid gas, heat in a warm water-bath, and set apart to crystallize.

PROTO-HYPOSULPHATE OF SODIUM. (GALL.; VAN MONS)

- R. Subcarbonate of Soda, eight parts.
Water, sixteen parts.

Add to the solution,

Sublimed Sulphur, one part.

Pass a current of sulphurous acid gas into the solution; filter after saturation, evaporate, and crystallize. (GALL.)

VAN MONS directs a current of hydro-sulphuric acid gas into a solution of sulphite of soda, or a current of sulphurous acid gas into a solution of hydro-sulphate of soda.

PROTO-NITRATE OF SODIUM.

Nitras Sodæ seu natriicum seu cubicum.—Nitrate of Soda, Nitrate of Protoxide of Sodium, Cubic Nitre, Rhomboidal Nitre.

NIEMANN.

- R. Subcarbonate of Soda, any quantity.
Diluted Nitric Acid,

a sufficient quantity

to saturate the alkali. Crystallize the solution.

This salt has been recommended in dysentery.

SOLUTION OF NITRATE OF SODA.

Solutio Nitratis Sodæ. (NIEMANN)

- R. Nitrate of Soda, half an ounce.
Gum Arabic, ten grains.
Water, eight ounces.
Dissolve.—Dose, a spoonful every hour.

HYDROSULPHATE OF SODA. (VAN MONS)

- R. Calcined Caustic Soda, one part.
Water, six parts.

Dissolve; pass a current of hydrosulphuric acid gas into the solution, and preserve it when it has become clear by repose.

SULPHUREOUS BATH. (CADET DE GASSI-COURT)

- R. Hydrosulphate of Soda, ten ounces.
Saline Gelatinous Solution, four ounces.

Mix, and add it to the water of the bath, at the moment of the patient entering it.

PROTO-ACETATE OF SODIUM.

Acetas Sodæ seu natricum, Alkali minerale acetatum, Natrum aceticum, Soda acetata seu acerosa, Terra foliata mineralis seu Tartari crystallisabilis.—Acetate of Soda, Acetate of Protoxide of Sodium, Acetate of Soda. (DUBL.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; GALL.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SWEDIAUR.; VAN MONS)

Sodæ Acetas. (DUBL.)

- R. Subcarbonate of Soda, any quantity

Distilled Vinegar,

a sufficient quantity

to saturate the alkali. Evaporate the filtered solution till it attains the specific gravity of 12.76. Carefully dry the crystals, which form as the liquor cools, and preserve them in well-stopped bottles.

The same process is given by the other pharmacopœias, with the exception of their directing the evaporation to be continued till a pellicle forms.

It is diuretic, or cathartic, according to the dose, which is from two to four drams.

ANTI-LACTEAL PILLS. (BORIES)

- R. Acetate of Soda, five drams.
Camphor, } of each, two drams.
Nitrate of Potass, }
Rob of Elder, a sufficient quantity.

Make one hundred and twenty pills.—

Dose, two, morning and evening.

DIURETIC POWDER. (AUGUSTIN)

- R. Acetate of Soda, } of each,
Oleosaccharum of Juniper, } three drams.
Squill, } six grains.
Sweet Flag, } half a dram.
Opium, } three grains.

Make six doses.—One, thrice a day, in infusion of juniper.

LAXATIVE PILLS.

Pilulæ e Rheo cum Natro acetico. (AUGUSTIN)

- R. Rhubarb, } of each,
Acetate of Soda, } one dram.
Inspissated Ox-gall, }
Gum Arabic, } half a dram.

Make a pilular mass.—Dose, from twenty to thirty grains, twice a day.

PROTO-MECONATE OF SODIUM.

Meconate of Soda. (FERR.)

R. Acetate of Soda, one ounce.
Alcoholic Tincture of Opium, four ounces.

Digest for several days; then filter to separate the crystals which are formed, and

purify them by means of boiling alcohol, which does not dissolve the meconate of soda.

This salt has been proposed as a certain remedy against tænia; but it should be employed with great circumspection, as it is violently poisonous in the dose of half a grain.

PROTO-OLEO-MARGARATE OF SODIUM.

See Proto-oleo-margarate of Potassium.

SOLANUM.

Five species of this genus are used in medicine:

1. *Solanum Dulcamara*, LINN. — Bitter-sweet, Woody Nightshade.

SYNONIMES:—Douce-amere, Morelle grim-pante, FR.; Bittersüsstengel, Hirsch-kraut, Rother Nachtschatten, Hindisch-krautstengel, GERM.; Sladka horka, BOHEM.; Hundebaer, Trolldær, DAN.; Dulciamargo, SPAN.; Bitterzoet, DUT.; Dulcamara, ITAL.; Glistrik Psinki wodne, POL.; Docamarga, PORT.; Qweswod, SWED.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENÈV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; COKE; GUIBOURT; MURRAY; and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A very common plant of Europe. (*Pentandria Monogynia*, LINN.; *Solanæ*, JUSS.; *Fig. Flore Medic.* III. 153)

The young branches are employed, gathered before they have become quite ligneous (*Stipites Dulcamaræ* seu *Amaræ dulcis* seu *Solani scandentis*). They are several feet long, of the thickness of the stock of a goose-quill or more, of a yellowish colour, flexible, branched, glabrous, and covered with a rugose bark. They have a virous odour, particularly when fresh. Their taste is at first bitter and then sweet.

DEFOSSES has discovered in them *Solanine*, an alkaloid, which constitutes their active principle.

This plant is excitant, irritant, and sudorific. It is employed in syphilitic diseases, rheumatism, and itch. It frequently occasions nausea and giddiness.

PILLS OF BITTER-SWEET. (SWEDIAUR)

R. Extract of Bitter-sweet, one ounce.
Sulphuret of Antimony, half an ounce.
Bitter-sweet, a sufficient quantity.

Make pills of four grains each.—Dose, three or four daily.

EXTRACT OF BITTER-SWEET.

Extractum Dulcamaræ. (AMST.)

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HANN.; LIPP.; OLDEN.; PALAT.; BORUS.; SAX.; WIRTEM.; HERBIP.; VAN MONS.

R. Stalks of Bitter-sweet, any quantity.

Bruise them in a mortar, and express strongly; set apart the juice for two hours, and evaporate it to the consistence of an extract. (LIPP.)

R. Stalks of Bitter-sweet, any quantity.

Boil for an hour in a sufficient quantity of water, and express; then set aside, decant, and evaporate the decoction to the consistence of thick honey. (BAVAR.; HERBIP.)

R. Dried Stalks of Bitter-sweet, any quantity.

Having cut and bruised them, macerate them in cold water, and boil slightly with the addition of more water; strain at the end of half an hour, and having clarified the decoction, evaporate it, and strain the liquid again through a sieve, when it begins to thicken. (VAN MONS)

R. Stalks of Bitter-sweet, one part.
Water, eight parts.

Macerate for twenty four hours, then boil for a quarter of an hour, and express strongly; boil the residue with

Water, four parts.

Mix the decoctions, set aside for twenty-four hours, and evaporate. (ANT.; SAX.)

BATAV.; and BELG. prescribe twenty-four hours' digestion and two hours' ebullition with ten parts of water, the ebullition of the residue for two hours with eight parts of water, and the evaporation of the mixed liquors.—AMST.—the ebullition of the stalks with twenty pounds of water, till reduced to one half, then with fifteen till reduced to one half, and the evaporation of the mixed liquors.

R. Stalks of Bitter-sweet, one part.
Boiling Water, eight parts.

Boil slowly for a quarter of an hour, and express; boil the residue with

Water, four parts.
and express again. Set aside the mixed decoctions, decant after sufficient repose, and evaporate slowly to the consistence of an extract. (HANN.; OLDEN.; BORUS.)

AUSTR.; and POL. direct the stalks to be exhausted by several successive ebullitions, the liquors to be mixed, decanted, and evaporated in a water-bath.—BRUGNATELLI—the stalks to be boiled for two hours with six parts of water, then a second time with fresh water, and the mixed decoctions to be evaporated.

R. Stalks of Bitter-sweet, half a pound.
Spring Water, two pounds.

Digest in a moderate heat for two days, and decant the liquor; repeat the digestion till the water comes off tasteless; then mix the liquors, and evaporate to the consistence of an extract. (PALAT.; WIRTEM.)

Dose, from five to ten grains.

ANTI-PHTHISICAL MIXTURE. (PIERQUIN)

R. Extract of Bitter-sweet, one scruple.
Oxymel of Squill, } of each,
Syrup of Poppy, } six drams.
Red Poppy, }
Water, eight ounces.
Orange-flower Water, one ounce.

Mix.—Dose, half a spoonful or one spoonful, four times a day.

RESOLVENT MIXTURE. (CADET DE GAS-SICOURT)

R. Extract of Bitter-sweet, three drams.
Seneka, two drams.
Hemlock, one dram.
Antimonial Wine, } of each,
Cinnamon Water, } one ounce.

Mix.—Dose, from forty to eighty drops, four times a day, in white swellings of the joints.

INFUSION OF BITTER-SWEET.

Infusum Dulcamaræ. (AUGUSTIN; NIEMANN)

R. Stalks of Bitter-sweet,
from two to four drams.

Boiling Water, one pound.

Infuse for half an hour; then boil for a quarter of an hour, and strain.

It is diaphoretic.—Dose, a spoonful every two hours.

DECOCTION OF BITTER-SWEET.

Decoctum Dulcamaræ. (LOND.; DUBL.; AMER.; FORM. HÔP. FR.; GENEV.; CAST. BORUS.; COXE; ELLIS; RATIER; SAUNDERS; SWEDIAUR; PH. SYPH.)

R. Stalks of Bitter-sweet, sliced,
one ounce.

Water, one pint and a half.

Boil down to one pint, and strain.
(LOND.; DUBL.)

AMER.; COXE; and ELLIS give the same formula.—SWEDIAUR—from half an ounce to one ounce of stalks and three pounds of water, to be boiled down to two thirds.—RATIER—one ounce of stalks and two pounds of water.—PH. SYPH.—half a dram of stalks and one pound of water, to be boiled down to one half.—GENEV.—one ounce of stalks and four pounds of water, to be boiled down to two.—AUGUSTIN—two drams of stalks and two pounds of water, to be boiled down to one.

R. Stalks of Bitter-sweet, one ounce.
Water, a sufficient quantity,

to obtain two pounds of strained liquor. Boil for a quarter of an hour, then infuse for two hours, and express. (FORM. HÔP. FR.)

R. Stalks of Bitter-sweet, one ounce.
Water, twenty-four ounces.

Boil till reduced to sixteen ounces. Strain, and add

Simple Syrup, one ounce.

Mix. (CAST. BORUS.)

R. Stalks of Bitter-sweet, two ounces.
Water, a sufficient quantity,

to obtain two pounds of strained liquor. Add

Syrup of Red Poppy, } of each,
Oxymel, } two ounces.

Mix.—(SAUNDERS)

It is stimulant, and is recommended in diseases of the skin, syphilitic affections, arthrodynia, dysmenorrhœa, amenorrhœa, and jaundice.—Dose, one pound, daily, either alone or with milk.

COMPOUND DECOCTION OF BITTER-SWEET. (AUGUSTIN)

R. Bitter-sweet, half an ounce.
Root of Liquorice, }
Burdock, } of each,
Wood of Sassafras, } two drams.
Guaiaac, }

Spring Water, two pounds.

Boil down to sixteen ounces of strained liquor.

It is recommended in acute rheumatism, and syphilitic diseases.

PECTORAL POTION.

Haustus pectoralis resolvens. (BRERA)

R. Bitter-sweet, two drams.
Water, a sufficient quantity

to obtain five ounces of strained decoction. Strain, and add

Extract of Soap-wort, one dram.
Oxymel, one ounce.

Mix.—To be taken gradually, to promote expectoration.

ALTERATIVE PTISAN. (CADET DE GASSICOURT)

R. Stalks of Bitter-sweet,
Herb of Fumitory,
Bark of Elm,
Root of Burdock,
Red Dock, } of each,
half an ounce.

Water, two pounds and a half.

Boil down to half a pound; strain, and add

Syrup of Sarsaparilla, two ounces.

Mix.

CARDIAC OINTMENT. (SPIELMANN)

R. Stalks of Bitter-sweet,
Fresh Herb of Mother-wort,
Germander,
Mallow,
Groundsel,
Scabious,
Houseleek,
Red Pimpernel,
Knapweed, } of each,
two handfuls.

Butter, one hundred and twenty-eight ounces.

Boil to the consumption of the humidity, and strain.

To be rubbed upon the abdomen; for children.

2. *Solanum Lycopersicum*, LINN.—Tomata, Love-apple.

SYNONIMES:—Tomate, Pomme d'amour, FR.; Liebesapfel, Goldapfel, GERM.; Elskovsæble, DAN.; Tomates, SPAN.; Appeltje der liefde, DUT.; Pomo d'amore, ITAL.; Tomato, PORT.; Karleksæple, SWED.

GALL.

The fruit (*Mala Lycopersica*) is employed. It consists of round berries, variously lobed, fleshy, succulent, smooth,
VOL. II.

shining, of a yellow or red colour, containing a great many seeds in the middle of an acidulous pulp.

3. *Solanum Melongena*, LINN.—Egg-plant.

SYNONIMES:—Aubergine, Mélongène, FR.; Eierfrucht, GERM.; Berinjela, SPAN.; Eijerdragende nagtschade, DUT.; Melanzana, ITAL.; Beringela, PORT.

HISP.; BERGIUS.

A plant cultivated in the south of Europe.

The fruit (*Pomum Melongenæ seu Mala insana*) is employed. It consists of a berry of very variable form and colour, sometimes white, and with the exact form of an egg, and sometimes prolonged and of a violet colour.

It is alimentary and demulcent, and is applied to painful piles

4. *Solanum nigrum*, LINN.—Garden Nightshade.

SYNONIMES:—Morelle noire, Morelle, FR.; Schwarzer Nachtschatten, GERM.; Psy vjno, BOHEM.; Svineurt, Soebær, DAN.; Hierbamorra, SPAN.; Zwarte nachtschade, DUT.; Solano nero, ITAL.; Psinki ziele, POL.; Herba moira, PORT.; Hansletsgras, SWED.

ANT.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENÈV.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

A very common plant of Europe. (*Fig. Zorn, Ic. pl. t. 44*)

The herb (*Herba Solani nigri*) consists of a round, unarmed, smooth, branching stem, and of oval, pointed, serrated, leaves. It has a disagreeable and stupefying odour, and a nauseous taste.

Its berries, according to DEFOSSÉS, contain *Solanine*.

It is feebly sudorific, emollient, and sedative, externally.

NARCOTIC DECOCTION. (RATIER)

R. Dried Leaves of Garden Nightshade, two ounces.
Poppy Heads, four.
Water, two pounds.

Boil, and strain.

It is employed in fomentations, local baths, injections, and lavements.

OIL OF GARDEN NIGHTSHADE.

Oleum Solani nigri. (HISP.; GALL.)

R. Herb of Garden Nightshade, with its seeds, one part.

4 K

Olive Oil, three parts.

Digest in warm ashes, and boil to the consumption of the humidity. (HISP.)

GALL. prescribes one part of leaves and two parts of oil.

OINTMENT OF GARDEN NIGHTSHADE.

Unguentum Solani. (SPIELMANN)

R. Expressed Juice of the leaves and berries of Garden Nightshade, thirty-six ounces.

Olive Oil, fifty-four ounces.

Subcarbonate of Lead, eight ounces.

Semi-vitreous Oxide of Lead, thirty-six ounces.

Boil, and add

White Wax, six ounces.

Powdered Olibanum, three ounces.

Mix.

WATER OF GARDEN NIGHTSHADE. (BRUNS.; GALL.; PALAT.)

R. Herb of Black Nightshade, one part.
Water, three parts.

Distil one half. (BRUNS.; PALAT.)

GALL. directs twelve parts and a half of water to be poured upon five of herb, and two parts distilled; the product to be mixed with five of herb and ten of water, and ten to be again distilled: the operation may be repeated a third time if thought requisite.

5. *Solanum tuberosum*, LINN.—Potato.

SYNONIMES:—Pomme de terre, FR.; Kartoffel, GERM.; Rata inada, CYNG.; Jord-eable, DAN.; Patata de la Mancha, SPAN.; Aardappel, DUT.; Ziemne jablko,

POL.; Batata da terra, PORT.; Jord-pæron, SWED.; Wal arai kilangu, TAM.; Uralay gudda, TEL.

GALL.; FERR.; AINSLIE; GUIBOURT; MURRAY and GMELIN.

A plant originally of America, from whence it was transported to Europe by SIR WALTER RALEIGH in 1584. (*Fig. Bauh. Prodr.* t. 39)

The root and herb are employed.

The culinary uses of the root are generally known.

The fecula which it contains in very large quantity is used in medicine:

The herb is reputed anti-spasmodic.—LATHAM recommends the extract of it.

FECULA OF POTATO; POTATO STARCH.

Fæcula Solani Tuberosi. (BELG.; GALL.; FERR.; NIEMANN; VAN MONS)

R. Potatoes, washed, any quantity.

Rasp them, and mix the pulp with a sufficient quantity of water to enable it to be passed through a sieve. Receive the strained turbid liquor in an earthenware vessel, and set it aside; purify the fecula with repeated ablutions, and dry it at a gentle heat in a dark place.

JELLY OF POTATO STARCH.

Gelatina ex Amylo Solani tuberosi. (NIEMANN)

R. Potato Starch, one ounce.
Water, one pound.
Simple Syrup, one ounce.

Boil.

SOLIDAGO VIRGA AUREA, LINN.—GOLDEN ROD, SARACEN'S WOUNDWORT.

SYNONIMES:—Verge d'or, FR.; Goldruthe, Heidenwundkraut, GERM.; Swalnjh pros-trednj, Wysoky trank, BOHEM.; Hei-densche wondkruid, DUT.

AMER.; AMST.; BRUNS.; HISP.; GALL.; FULD.; GENEV.; LIPP.; WIRTEM.; HERBIP.; BERGIUS; COXE; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A plant common in Europe. (*Syngenesia Polygamia Superflua*, LINN.; *Synantherea*, CASS.; *Fig. Zorn, Ic. pl.* t. 383)

The herb (*Herbæ Virgæ aureæ seu Consolidæ Saracenicæ seu Solidaginis Sarase-*

nica) consists of a pubescent, branching stem, and of leaves, the lower ones of which are elliptical and the upper lanceolated. It has a bitter, styptic, and somewhat aromatic taste.

It is diuretic.

EXTRACT OF GOLDEN ROD.

Extractum Virgæ aureæ. (HERBIP.)

R. Herb of Golden Rod, any quantity.

Boil it for an hour in a sufficient quantity of water, and express; set aside and decant the liquor, and evaporate it to the consistence of thick honey.

SONCHUS OLERACEUS, LINN.—SOW THISTLE.

SYNONIMES:—Laiteron, FR.; Kohlgänsedistel, GERM.; Svinetistel, DAN.; Corraja, SPAN.; Milkdistel, DUT.; Sonco, ITAL.; Mleczne, POL.; Serrutha, PORT.; Mjoelk tistel, SWED.

BRUNS.; GALL.; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe. (*Syngenesia*

Polygamia Æqualis, LINN.; *Synantherææ*, CASS.; Fig. Zorn, *Ic. pl. t. 148*)

The herb (*Herba Sonchi lavis et asperi*) is employed. It consists of a branching, cylindrical, glabrous, hollow stem, and of alternate, embracing, smooth, usually run-cinated leaves, having acute, dentated lobes, of which the terminal is triangular and very large. It has a bitterish taste.

SORBUS.

Two species of this genus are employed.

1. *Sorbus domestica*, LINN.—Sorb, or Service Tree.

SYNONIMES:—Cormier, FR.; Spierlingsbaum, Eschræleinbaum, GERM.

GALL.; WIRTEM.; BERGIUS; BRUGNATELLI.

A tree of Europe. (*Icosandria Monogynia*, LINN.; *Rosacææ*, JUSS.; Fig. Jacq. *Austr. V. t. 447*)

The fruit, *Sorb Apple* (*Fructus Sorbi*), is employed. It is a small pyriform apple, containing three oblong, cartilaginous seeds. It is at first very acerb, but becomes sweet when over ripe, though it always preserves a little astringency.

2. *Sorbus aucuparia*, LINN.—Mountain Ash.

SYNONIMES:—Sorbier, Sorbier des oiseleurs, FR.; Vogelbeerbaum, GERM.; Fuglebær, DAN.; Serbar de cazador, SPAN.; Sorbenboom, DUT.; Sorbo selvatico, ITAL.; Jakszebina, POL.; Sereveira brava, PORT.; Rænn, SWED.

BRUNS.; FERR.; FULD.; LIPP.; SUEC.; BERGIUS; SPIELMANN.

A tree met with in all parts of Europe. (*Icosandria Trigynia*, LINN.; *Rosacææ*, JUSS.; Fig. Zorn, *Ic. pl. t. 440*)

The fruit (*Baccæ seu Poma Sorbi aucupariæ*) is employed. It consists of small round berries of a vivid red colour, soft, umbilicated at the summit, and containing three cartilaginous seeds.

Their taste is astringent, acid, and disagreeable.

The *Sorbic Acid* which is contained in them, according to LABILLARDIERE and BRACONNOT, is merely malic acid.

They are astringent, diuretic, hydragogue, and anti-scorbutic.

ROB OF THE FRUIT OF MOUNTAIN ASH.

Roob Sorbi aucupariæ. (FULD.; LIPP.)

R. Berries of Mountain Ash, eight pounds.

Pour a sufficient quantity of water over them to cover them by a finger's breadth, and set the vessel over a slow fire for half an hour; then to the strongly expressed liquor add

Sugar, one pound, and evaporate slowly to the proper consistence.

Dose, from half an ounce to one ounce.

SPIGELIA.

Two species of this genus are used in medicine:

1. *Spigelia anthelmia*, LINN.—Anthelmintic Spigelia.

SYNONIMES:—Spigélie anthelminétique, FR.; Arapubaca, BRAZ.; Espigelia, PORT.

AMST.; BATAV.; BRUNS.; DAN.; GALL.; FERR.; ROSS.; SAX.; WIRTEM.; HERBIP.; GUIBOURT; MURRAY and GMELIN.

A plant of North America. (*Pentandria Monogynia*, LINN.; *Gentianææ*, JUSS.; *Fig. Desc. Flore Ant.* 1. 61)

The root and herb are employed.

The root (*Radix Spigeliæ anthelmia*) is small and fibrous.

The herb consists of a round stem, at the summit of which are four large verticillated leaves.

2. *Spigelia Marylandica*, LINN.—Maryland Spigelia, Snake Root, Indian Pink, Carolina Pink.

LOND.; DUBL.; EDINB.; AMER.; AMST.; BATAV.; BELG.; DAN.; HISP.; GALL.; FERR.; FULD.; LIPP.; LUSIT.; ROSS.; SAX.; SUEC.; WIRTEM.; BERGIUS; BRUGNATELLI; COXE; MURRAY and GMELIN.

A plant of North America. (*Fig. Cat. Carol.* 2. p. 78. t. 78)

The root and herb are employed.

The root (*Radix Spigeliæ Marylandicæ*) is simple and fibrous.

The herb consists of a tetragonal stem, and of opposite, sessile, oval, long, entire, pointed, smooth leaves. It has a fetid and narcotic odour, and a bitter and nauseous taste.

It is excitant, anthelmintic, and laxative. —Dose of the powder, twelve grains to children of twelve years old, and one dram for adults.

VERMIFUGE POWDER. (ELLIS)

R. Maryland Spigelia, sixteen grains.
Proto-chloruret of Mercury, eight grains.

Mix.—For one dose.

DECOCTION OF SPIGELIA.

Decoctum Spigeliæ. (FULD.; HERBIP.)

R. Herb of Anthelmintic Spigelia, two handfuls.
Spring Water, two pounds.
Boil, and add to the strained decoction,
Fresh Lemon Juice, six drams.
Syrup of Peach Flowers, two ounces.
Water, two pounds.

Mix.—Dose, two or three ounces, every six hours; for an adult.

INFUSION OF ANTHELMINTIC SPIGELIA.

Infusum anthelminticum. (AMER.; FULD.; COXE; ELLIS; NIEMANN; SAUNDERS; SWEDIAUR)

R. Herb of Anthelmintic Spigelia, half an ounce.
Boiling Water, six ounces.

Macerate for two hours, and strain. (ELLIS; SAUNDERS; SWEDIAUR)

AMER.; and COXE prescribe two drams of spigelia, half a pound of water, and four hours' digestion.

R. Herb of Spigelia, three drams.
Boiling Water, four ounces.

Add to the strained infusion,
Oil of Rue, two drops.

Mix. (FULD.)

Dose, from one ounce and a half to two ounces, in the evening.—Two drams of seneka may be added.

VERMIFUGE INFUSION. (ELLIS)

R. Maryland Spigelia, half an ounce.
Senna, two drams.
Manna, one ounce.
Fennel, three drams.
Boiling Water, one pound.

Infuse.—Dose, a glassful, three or four times a day.

SYRUP OF SPIGELIA. (CADET DE GASSI-COURT)

R. Leaves of Anthelmintic Spigelia, two ounces.
Seeds of Anise, bruised, half a dram.
Water, six ounces.

Infuse in a water-bath for six or seven hours; then express strongly, and add

Aqueous Extract of Jalap, half an ounce.

Mix, filter, and add

White Sugar, twice the quantity.

Dissolve. —Dose, two tea-spoonfuls, daily, to children; and a table-spoonful, morning and evening, to adults, in any proper vehicle.

SPILANTHUS Acmella, LINN.—COMMON SPILANTHUS.

SYNONIMES:—Acmelle, Cresson des Indes, FR.; Akmella, Indianische Harnkraut, GERM.

BRUNS.; GALL.; WIRTEM.; MURRAY and GMELIN; SPIELMANN.

A plant of Ceylon. (*Syngenesia Po-*

lygamia Æqualis, LINN.; *Synantherææ*, CASS.; *Fig. Rumph. Amb. VI. t. 65*)

The herb (*Herba Acnellæ* seu *Attmellæ* seu *Admellæ* seu *Accmellæ* seu *Achmellæ*) is employed. It consists of a straight stem, and of oval, dentated leaves of a bitter and aromatic taste, becoming burning after being chewed for a long time, and inducing an abundant flow of saliva.

It is brought to us, the leaves green and

in fragments, and mixed with a great proportion of rather long and thick stems. It is then nearly insipid. From its deariness, the herbs of the *Sigesbechia orientalis*, LINN.; of the *Coreopsis bidens*, LINN.; and of the *Bidens tripartitum*, LINN.; are frequently mixed with it.

It is bitter, aromatic, excitant, and diuretic.

SPINACIA OLERACEA, LINN.—SPINAGE.

BRUNS.; GALL.; BERGIUS.; MURRAY and GMELIN.

A plant cultivated in the kitchen gardens of Europe. (*Diœcia Pentandria*, LINN.; *Atriplicææ*, JUSS.; *Fig. Enc. meth. t. 814*)

Its native country is not known.

The herb (*Herba Spinacææ* seu *Spinachia*) is employed. It consists of a straight, branching stem, and of petiolated, arrow-shaped leaves, and of a deep green colour. Its taste is vapid.

It is emollient.

SPIRÆA.

Two species of this genus are used :

1. *Spiræa Filipendula*, LINN.—Dropwort.

SYNONIMES:—*Filipendule*, FR.; *Spierstaude*, *Filipendelwurz*, *Rothe Steinbrech-wurz*, GERM.; *Rød Steenbrek*, DAN.; *Roode Steenbreck*, DUT.; *Filipendula*, ITAL.; *Kropidelco mnicyse*, POL.; *Brudbrød*, SWED.

BRUNS.; HISP.; GALL.; LIPP.; WIRTEM.; HERBIP.; BERGIUS.; GUIBOUT.; MURRAY and GMELIN.; SPIELMANN.

A plant of Europe. (*Icosandria Pentagynia*, LINN.; *Rosacææ*, JUSS.; *Fig. Zorn, Ic. pl. t. 394*)

The root (*Radix Saxifragæ rubræ* seu *Filipendulæ*) is employed. It consists of a great number of small fibres furnished with fleshy, round, or oblong tubercles. It is of a deep-brown or blackish colour without, and of a slightly-reddish white within. It is destitute of odour except in autumn when it diffuses one of orange; its taste is sweet, but after desiccation it becomes astringent and bitterish.

The Cossacks employ an infusion of the leaves against tænia.

2. *Spiræa ulmaria*, LINN.—Queen of the Meadows.

SYNONIMES:—*Reine des près*, FR.; *Geisbartwurz*, *Wiesenköniginn*, *Sumpfspierstaude*, GERM.; *Mædurn*, DAN.; *Reyna del prado*, SPAN.; *Reynette*, DUT.; *Regina degli prati*, ITAL.; *Rozia brodka*, POL.; *Rainha dos prados*, PORT.; *Micæært*, SWED.

BRUNS.; HISP.; GALL.; GENEV.; WIRTEM.; BERGIUS.; MURRAY and GMELIN.; SPIELMANN.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 141*)

The root, leaves, and flowers are employed.

The root (*Radix Barbæ caprinæ* seu *Barbulæ caprinæ* seu *Ulmariæ* seu *Regina pratensis*) is employed. It is cylindrical, unequal, furnished with a great many small fibres, of a blackish-brown colour externally, and yellowish or reddish internally, and of a bitterish and styptic taste.

The leaves are pinnate, the leaflets unequally three lobed, and the leaflets are alternately large and small.

The flowers are white. They have a feeble but agreeable odour of bitter almonds which becomes dissipated by drying. Their taste is sweet and sub-astringent.

SPONGIA MARINA.—SPONGE.

SYNONIMES :—Eponge, Eponge de mer, FR. ; Badeschwamm, Meerschwamm, GERM. ; Isfenj, ARAB. ; Esponjas, SPAN. ; Muabadul, HIND. ; Spons, DUT. ; Spugna, ITAL. ; Univatta, JAP. ; Abermurdeh, PERS. ; Esponja, PORT. ; Badswamp, SWED.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; CAST. BORUS. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; WURCEB. ; AINSLIE ; BRUGNATELLI ; COXE ; GUIBOURT ; PARIS ; PIDERIT ; SPIELMANN.

The fibrous, reticulated, or rather felted skeleton of various organized marine bodies, occupying the extreme limits of the animal kingdom.

The *Spongia lacinulosa*, LMK. (*Fig. Esp. die Pflanzenth.* II. t. 1517) of the mediterranean, Red Sea, and Indian Ocean, is most generally employed. It is round, flattish, sublobate, soft, tomentose, porous, and covered with numerous narrow and irregular notches on its surface.

FYFE has discovered iodine in it.

PREPARED SPONGE.

Spongia preparata. (AMST. ; ANT. ; AUSTR. ; BAVAR. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; BORUS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BRUGNATELLI ; PIDERIT ; SPIELMANN ; SWEDIAUR ; VAN MONS)

1. *Without Wax.*

Wash and cleanse a fine sponge, and bind it closely and firmly with a string while it is still moist; having fastened the string with a slip-knot, dry the sponge carefully, and keep it in a dry place. (FERR. ; FORM. HÔP. FR. ; GALL. ; SWEDIAUR)

HISP. directs the sponge to be cut into long strips, and dipped in white of egg, before it is bound up.—VAN MONS directs it to be saturated with mucilage of gum arabic.

2. *With Wax; Spongia cerata.*

Wash, cleanse, and dry a fine sponge; then dip it into melted yellow wax, and press it between two tin plates heated in a water-bath. Remove it when cold. (AMST. ; ANT. ; AUSTR. ; BAVAR. ; BRUNS. ; DAN. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; BORUS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BRUGNATELLI ; PIDERIT ; SPIELMANN ; SWEDIAUR ; VAN MONS)

It is used for the dilatation of fistulous and other openings.

BURNT SPONGE.

Spongia usta. (LOND.)

Cut sponge into small pieces, and bruise it in order to free it from any adherent extraneous matters; then burn it in a covered iron vessel till it is black and friable, and finally reduce it to a very fine powder.

STACHYS RECTA, LINN.—WHITE STACHYS.

SYNONIMES :—Crapaudine, FR. ; Berufskraut, GERM. ; Estaquis recta, Yerba de feridura, SPAN.

HISP. ; WIRTEM. ; SPIELMANN.

A plant of the south of Europe. (*Didynamia Gymnospermia*, LINN. ; *Labiata*, JUSS. ; *Fig. Jacq. Austr.* IV. t. 359)

The herb (*Herba Sideritis hirsuta*) presents a square, rather branching, and rather hairy stem, and elliptical or lanceolated, heart-shaped, crenelated, rough leaves. It has an agreeable odour, and an aromatic taste.

It is stimulant.

STANNUM.—TIN.

SYNONIMES:—*Jupiter*; *Καττιτερος*, *σλαγγος*.

—ETAIN, FR.; ZINN, GERM.; TIN, DAN.; and DUT.; RESAS, ARAB.; YANG-SEIH, CHIN.; RUNGA, DUK.; ESTANNO, SPAN.; KULAI, HIND.; STAGNO, ITAL.; TIMA, MAL.; URZIZ, PERS.; CYNA, POL.; ESTANTHO, PORT.; OLOWO, RUSS.; TRAPU, RANGA, SANS.; TENN, SWED.; TAJARUM, TAM.; GALAI, TEL.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE; COXE; GUIBOURT; SAUNDERS; SPIELMANN.

A solid metal, nearly as white as silver, soft, laminable, scarcely ductile, fusible at 210 degrees, not volatile, scarcely alterable by damp air. Its specific gravity is 7.291. It makes a peculiar cracking sound when bent.

POWDERED TIN.

Stannum pulveratum seu præparatum seu granulatum seu raspatum, Limatura Stanni, Oxydulum Stanni, Pulvis Stanni philosophice præparatus. (DUBL.; AMER.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; GALL.; FERR.; FULD.; HASS.; LIPP.; OLDEN.; PALAT.; ROSS.; SARD.; SUEC.; HERBIP.; COXE; SAUNDERS; SWEDIAUR)

Pulvis Stanni. (DUBL.)

R. Tin, any quantity.

Melt it in an iron mortar over a fire, then stir it as it cools, till it becomes a powder, and when cold pass it through a sieve.

R. English Tin, any quantity.

File it with a very fine file, and preserve the sifted powder for use. (AMER.; BAVAR.; BRUNS.; FERR.; PALAT.; COXE; SWEDIAUR)

LIPP. directs the filings to be boiled in vinegar till the liquor ceases to be precipitated by subcarbonate of potass.

R. Pure Tin, any quantity.

Melt it, and pour it into a rough wooden box rubbed with chalk; then close the box and shake it rapidly. Pass the powder through a sieve. (AUSTR.; BATAV.; GALL.; OLDEN.; PALAT.; SARD.; SUEC.; HERBIP.)

R. Tin Filings, two parts.
Chalk, in powder, one part.
Triturate together in a mortar. (ROSS.)

SAUNDERS directs three drams of tin and one ounce of coral to be triturated together.

R. Leaf Tin, } of each,
White Sugar, } an equal part.

Triturate together; then having sifted the powder, boil it in a large quantity of water, and set aside to crystallize. (ANT.)

BELG. directs muriate of soda to be triturated rapidly in a heated mortar with melted tin, till the whole is reduced to very fine powder; this to be well washed, dried, and sifted.

R. Tin Filings, two parts.
Crab's Eyes, one part.

Triturate together. (FULD.)

VERMIFUGE POWDER. (CADET DE GASSICOURT)

R. Powdered Tin, one ounce.
Sulphuret of Mercury, two scruples.

Mix.—A sixth part to be taken twice a day in syrup, treacle, or honey.

FERRUGINOUS POWDER OF TIN.

Pulvis Stanni martiatus. (AUGUSTIN; SAUNDERS; SWEDIAUR)

R. Tin Filings, two drams.
Sulphate of Iron, five grains.

Triturate together, and make five parcels.
—Dose, one every two hours.

VERMIFUGE BOLUS.

Bolus e Stanno. (SAUNDERS; SWEDIAUR)

R. Powdered Tin, half a scruple.
Conserve of Orange Peel, one scruple.
Simple Syrup, a sufficient quantity.

Mix.

VERMIFUGE ELECTUARY. (CADET DE GASSICOURT; ELLIS; PIERQUIN; RATIER)

R. Powdered Tin, one ounce.
Despumated Honey, four ounces.

Mix.—Dose two or three spoonfuls in the morning. (ELLIS)

R. Powdered Tin, one ounce.
Extract of Mugwort, } of each,
Powdered Jalap, } one dram.
Compound Syrup of Cichory, a sufficient quantity.

Mix.—To be taken at twelve or fifteen doses, at intervals of half an hour. (RATIER)

- R. Powdered Tin, one ounce.
 Root of Male Fern, six drams.
 Semen Contra, half an ounce.
 Jalap, } of each,
 Sulphate of Potass, } one dram.
 Honey, a sufficient quantity.

Mix.—Dose, half a tea spoonful in the morning, repeated every half hour. (AUGUSTIN; CADET DE GASSICOURT; PIERQUIN)

AMALGAM OF TIN. (AMER.; HISP.; HERBIP.; COXE; SAUNDERS)

- R. Tin, two ounces.
 Mercury, three ounces.

Heat the mercury in a crucible till it begins to be volatilized; then add the tin fused separately, and agitate the mixture in a marble mortar, adding a little muriate of soda: lastly wash it well, and dry it in a linen cloth. (HISP.)

SAUNDERS directs three drams of mercury and two ounces of tin to be united, and the amalgam triturated to powder in a marble mortar.—AMER.; and COXE—five parts of tin and two of mercury to be united and the amalgam triturated with one part of prepared chalk till it has entirely lost its metallic lustre.—HERBIP.—one ounce of tin to be amalgamated with one ounce of mercury, and the mixture when cold to be powdered with one ounce of prepared oyster shells.

VERMIFUGE POWDER. (SAUNDERS)

- R. Amalgam of Tin, half a dram.
 Rhubarb, } of each,
 Deutoxide of Antimony, } ten grains.

Mix.—To be taken twice a day.

ANTHELMINTIC PILLS. (VAN MONS)

- R. Amalgam of Tin, any quantity.
 Set it over hot ashes that the metals may become oxidized. Add

Extract of Wormwood,
 a sufficient quantity.

Make pills of six grains each.—Dose, four, six, or eight, in the morning.

VERMIFUGE ELECTUARY.

Electuarium joviale ad taniam. (BRUNS.; SARD.; HERBIP.; SPIELMANN; VAN MONS)

- R. Pure Tin, }
 Mercury, } of each, one ounce.

Make an amalgam; pulverize it with Prepared Oyster Shells, one ounce.
 Add gradually,
 Conserve of Wormwood, two ounces.

Syrup of Mint, a sufficient quantity, to make an electuary. (BRUNS.; SARD.; HERBIP.; SPIELMANN)

VAN MONS substitutes calcined magnesia for the subcarbonate of lime, and simple syrup for that of mint.

Dose, two drams, twice a day.

ANTI-HÆMORRHOIDAL OINTMENT.

Unguentum empiricum Flaki in tumoribus hæmorrhoidalibus. (BRERA)

- R. Tin Filings, }
 Mercury, } of each, two drams.

Make an amalgam, and add

Rose Ointment, one ounce.
 Nitric Oxide of Mercury, two drams.
 Oil of Peppermint, twenty drops.

Triturate together.

PROTOXIDE OF TIN.

Stannum sub-oxdatum seu pulveratum, Pulvis Stanni. (PALAT.; NIEMANN; SWEDIAUR; VAN MONS.)

- R. Pure Tin, any quantity.

Keep it in fusion in an open vessel, till it is entirely converted into a grey powder; then triturate and sift it. (PALAT.; SWEDIAUR; VAN MONS)

- R. Chloruret of Tin, dissolved in water, any quantity.

Solution of Potass, a sufficient quantity, to precipitate and re-dissolve the oxide. Filter the liquor, and precipitate the oxide with hydrochloric acid. (NIEMANN)

It is anthelmintic and purgative, and is principally prescribed against tinea, though it is also extolled in phthisis pulmonalis.—Dose, five or six grains.

POTIER'S ANTI-HECTIC.

Anti-hecticum Poterii, Bezoardicum joviale. (SARD.; WIRTEM.; SPIELMANN; VAN MONS)

- R. Regulus of Antimony, four parts.
 Fine Tin, five parts and a half.

Fuse together, and pour the mass upon a cold metal slab; pulverize, and mix it with

Nitrate of Potass, fifteen parts.

Project the mixture into a crucible at a white heat, and still keep it at the same heat for some time after it is fused; then having let it cool, wash it with cold water, and dry it slowly. (VAN MONS)

WIRTEM.; and SPIELMANN prescribe six ounces of regulus, two of tin, and twenty-four of nitre.—SARD.—six of regulus, six of tin, and thirty-six of nitre. This prepa-

ration was formerly employed to arrest sanguineous and seminal discharges, and to diminish the colliquative sweats of consumptive patients.—Dose, half a dram, morning and evening.

PERSULPHURET OF TIN, MOSAIC GOLD.

Sulphuretum Stanni, Aurum mosaicum.
(ANT.; FERR.; BRUGNATELLI; CADET DE GASSICOURT; NIEMANN; VAN MONS)

R. Powdered Tin, three ounces.
Sublimed Sulphur, one ounce.

Put them into a crucible placed in burning charcoal, and apply heat till a vivid flame is produced; then remove the crucible, cover it, set it apart to cool, and finally levigate and sift the mass. (ANT.; FERR.; BRUGNATELLI; CADET DE GASSICOURT)

VAN MONS directs seven parts of oxide of tin to be calcined with four and a half of sulphur.

R. Chloruret of Tin, any quantity.

Pour into it,

Solution of Hydrosulphate of Potass,
a sufficient quantity;

or, till it ceases to produce any precipitate. Dry this, and heat it to redness. (VAN MONS)

R. Tin, twelve parts.
Mercury, three parts.
Sublimed Sulphur, seven parts.

Hydrochlorate of Ammonia,

three parts.

Triturate together, and apply a gentle heat till sulphurous acid gas ceases to be disengaged; then calcine slowly till a scaly, yellow, brilliant mass remains. (NIEMANN)

VERMIFUGE POWDER. (VAN MONS)

R. Persulphuret of Tin, three parts.
Mercury, one part.

Triturate to the extinction of the mercury.

It is principally recommended against tænia.—Dose, from two drams to half an ounce, in the morning, in conserve of wormwood.

CHLORURET OF TIN.

Murias Stanni seu Oxyduli Stanni, Liquor fumans Libanii, Stanni Butyrum, Stanni Deuto-hydrochloras. (BAVAR.; NIEMANN; VAN MONS)

R. Tin, one part.
Concentrated Hydrochloric Acid, three parts.

Dissolve with heat, and crystallize.

It is violently purgative.—Dose, two grains.—BAVAR. only indicates it as a chemical test dissolved in nine parts of water, to ascertain the presence of gold.

STATICE LIMONICUM, LINN.—SEA LAVENDER.

SYNONIMES :—Marsh Rosemary, Lavender Thrift.—Behen rouge, FR.; Rother Behen, Limonienkraut, Meerneekengras, GERM.

BATAV.; BRÜNS.; GALL.; GENEV.; PALAT.; WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A plant which grows in the north of both continents, on the sea shore. (*Pentandria*

Pentagynia, LINN.; *Plombagineæ*, JUSS.; *Fig. Céd. Fl. Dan. t. 315*)

The root (*Radix Behen rubri seu Limonii*) is employed. It is met with in the shops, in round, wrinkled, arched, solid pieces, of a brown colour externally and reddish internally. Its odour is feeble, and its taste astringent.

A congeneric species, the *Statice Caroliniana* (*Fig. Walter, Flor. Carol. 118*), is indicated by AMER.; and COXE.

STIZOLOBIUM PRURIENS, P.—COWHAGE.

SYNONIMES :—*Dolichos pruriens*, (LOND.; EDINB.; DUBL.)—Pois à gratter, FR.; Kuhkretze, Juckende faseln, GERM.; Kosambiliwail, Dewipaghura, CYNG.; Klænde bæne, DAN.; Kaunchkurikebinge, DUK.; Kiwack, HIND.; Jeuk-boontjes, DUT.; *Dolico pizzicante*, ITAL.;

RAWÉ, JAP.; Kapikachlin, SANS.; Punaykalie, TAM.; Piliadugukaila, TEL.

LOND.; DUBL.; EDINB.; DAN.; FERR.; SAX.; SUBC.; WIRTEM.; AINSLIE; BERGIUS; COXE; MURRAY and GMELIN.

A shrub of South America. (*Diadelphia Decandria*, LINN.; *Leguminosæ*, JUSS.; *Fig. Desc. Flor. Ant.* t. 49)

The fruit (*Siliqua hirsuta*) is employed. It is a pod of from two to four inches in length, and of about the thickness of the finger, coriaceous, arched, and covered with sharp, brittle hairs, easily detached, shining, and of a reddish-brown colour (*Dolichi*

Pubes seu Setæ, *Stizolobium*, *Setæ seu Lanugo Siliquæ hirsutæ*).

It is esteemed anthelmintic.—Dose, from three to six grains in syrup.

VERMIFUGE ELECTUARY.

R. Hairs of Cowhage, one dram.
Honey, a sufficient quantity.

Mix.—Dose, a tea spoonful, to a child of two or three years old.

STRYCHNINA, STRYCHNINUM, STRYCHNIA.—STRYCHNINE.

GALL.; FERR.; CADET DE GASSICOURT;
MAGENDIE; NIEMANN.

R. Alcoholic Extract of Nux Vomica,
any quantity.

Having mixed it with water, add solution of acetate of lead to the liquor till it ceases to throw down any precipitate; separate the lead by passing a current of hydrosulphuric acid gas into the liquor; then filter, and boil it with magnesia. Wash the precipitate with cold water, dissolve it in alcohol, and evaporate the solution to dryness.

It is violently stimulant and poisonous, serious effects being frequently produced by the fourth of a grain. It is prescribed in palsies, commencing with the sixteenth of a grain, and increasing the dose very gradually.

PILLS OF STRYCHNINE.

Pilule Strychnini. (FERR.; BRERA; BORIES; CADET DE GASSICOURT; ELLIS; FEE; HENRI; MAGENDIE; PIERQUIN; NIEMANN; RATIER)

R. Strychnine, two grains.
Conserve of Roses, half a dram.

Make twenty-four pills, and silver them.
(FERR.; BORIES; CADET DE GASSICOURT; ELLIS; FEE; HENRI; MAGENDIE; NIEMANN; PIERQUIN; RATIER)

BRERA prescribes from half a grain to six grains of strychnine and a sufficient quantity of crumb of bread, of despumated honey, and of liquorice powder, to make four pills; one of which is to be taken every two hours in the morning.—It is very likely that there is an error in this formula.

TINCTURE OF STRYCHNINE.

Tinctura Strychnini. (FERR.; BORIES; FEE; MAGENDIE; NIEMANN; PIERQUIN; RATIER)

R. Strychnine, three grains.
Alcohol (36 degrees), one ounce.

Mix.—Dose, from six to sixty drops.

STIMULANT POTION. (FEE; MAGENDIE; PIERQUIN; RATIER)

R. Strychnine, one grain.
Distilled Water, two ounces.
White Sugar, two drams.
Acetic Acid, two drops.

Mix.—A spoonful to be taken morning and evening.

POWDER OF STRYCHNINE. (BRERA)

R. Strychnine, one sixth of a grain.
Deutoxide of Iron, six grains.
Sugar, half a dram.

Mix.—To be taken morning and evening.

STRYCHNOS COLUBRINA, LINN.—SNAKEWOOD.

SYNONIMES:—Bois de couleuvre, FR.; Schlangenhholz, GERM.; Kuchila-luta, BENG.; Mahapenala, CYNG.; Wideropait; JAV.; Modira caniram, MALAB.

BRUNS.; HISP.; GALL.; PALAT.; WIRTEM.; AINSIE; BERGIUS; GUIBOUT; MURRAY; and GMELIN; SPIELMANN.

A tree of the Moluccas. (*Pentandria Monogynia*, LINN.; *Apocynææ*, JUSS.; *Fig. Rumph. Amb.* 2. c. 46. t. 37)

The root (*Lignum colubrinum*) is employed. It is in heavy pieces, of a yellowish-white colour, covered with a brownish, ru-

gose, tuberculous bark. It is destitute of odour, and is exceedingly bitter when old.

PELLETIER and CAVENTOU have found strychnine in it.

This wood is a reputed specific against the bites of venomous animals. It should be employed with circumspection.—BOERHAAVE extolled it in intermittent fevers.

STYRAX LIQUIDAMBAR.—LIQUID STORAX.

SYNONIMES: — *Ambra liquida*. — Styra, Styra liquide, Liquidambar, Fr.; Flüssiger Storax, GERM.; Tekaucy Styra, BOHEM.; Estoraque liquido, SPAN.; Vloëjende storax, DUT.; Storax rzadki, POL.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

There are two sorts of this kind of storax.

1. *Copalm Liquid Storax, Balsam or Oil of Copalm, Louisiana Copalm*; of the consistence of honey, transparent, of a reddish-yellow colour becoming of a deep red or blackish with time, of a very agreeable odour, and of an acid and aromatic taste. It solidifies into a blackish-brown, brittle resin.

2. *Common Liquid Storax*; of the consistence of thick ointment, of a reddish, greyish, or brownish colour, scarcely translucent, of a strong and agreeable odour and of an acrid and bitter taste.

This balsam flows from the *Liquidambar styraciflua*, a tree of Mexico. (*Monœcia Polyandria*, LINN.; *Amentaceæ*, JUSS.; *Fig. Flore Medic. VI. 331*)

BONASTRE has discovered in it a peculiar resinous and crystallizable matter, which he calls *Styacine*.

It is stimulant, and is reserved almost exclusively for external use.

OINTMENT OF STORAX.

Unguentum Styracis resinosum seu e Styraçe compositum. (ANT.; BELG.; BRUNS.; GALL.; FORM. HÔP. FR.; GENEV.; HANN.; SARD.; WIRTEM.; HERBIP.; CADET DE GASSICOURT.; NIEMANN.; SPIELMANN.; VAN MONS.)

R. Walnut Oil,
three hundred and fifty parts.

Liquid Storax, eight parts.

Melt at a gentle heat, occasionally stirring; then set aside, strain, and add

Resin, four hundred and eighty parts.
Elemi, } of each, one hundred
Yellow Wax, } and ninety-two parts.

Strain through a linen cloth. (GALL.; FORM. HÔP. FR.)

GENEV.; SARD.; and SPIELMANN prescribe one pound of walnut oil, half a pound of storax, one pound of resin, and half a pound each of elemi and of wax.

R. Olive Oil, two ounces and a half.
Liquid Storax, } of each seven
Elemi, } ounces and a half.
Yellow Wax, }
Resin, two ounces.

Melt together. (BRUNS.; HANN.; WIRTEM.; NIEMANN)

HERBIP.; and VAN MONS prescribe one part each of oil, of liquid storax, of elemi, and of wax, and two parts of resin.—SUEC.—one and a half of oil, and seven each, of the four other ingredients.

R. Turpentine, fifteen ounces.
Walnut Oil, eleven ounces.
Elemi, } of each,
Yellow Wax, } seven ounces.

Melt slowly together; remove from the fire, and add

Liquid Storax, seven ounces.

Evaporate the humidity at a gentle heat. (ANT.; BELG.)

R. Resin, } of each, one pound.
Lard, }
Walnut Oil, fifteen ounces.
Elemi, } of each, six ounces.
Yellow Wax, }

Add to the melted mass,

Liquid Storax, seven ounces.

Mix. (CADET DE GASSICOERT)

STORAX PLASTER.

Emplastrum Styracis Stomachicum.
(HISP.)

R. Yellow Wax, } of each, two pounds.
Olibanum, }
Turpentine, } three pounds.

Add to the melted mixture,

Liquid Storax, } of each, three
Mastic, } ounces and a half.
Labdanum, }

Benzoin, } two ounces two scruples.
 Pimenta, } four ounces and a half.
 Oil of Mint, } one ounce of each,
 ——— Wormwood, } two drams.

Mix.

R. Yellow Wax, two pounds and a half.
 Turpentine, three pounds.
 Tacamahac, two pounds and a half.
 Liquid Storax, } of each, four
 Mastic, } ounces and a half.
 Labdanum, }
 Benzoin, }
 Cloves, } of each, eighteen drams.
 Nutmeg, }
 Oil of Mint, } of each,
 ——— Wormwood, } two drams.

Soften the storax, and beat it with the labdanum, adding a little wax; then melt the wax with the turpentine, and strain; finally add the powders and the oils.

DIGESTIVE OINTMENT. (RATIER)

R. Ointment of Storax, one pound.
 Olive Oil, half a pound.

Mix.

ANTI-SCORBUTIC OINTMENT. (SAINTE-MARIE)

R. Storax Ointment, five ounces.
 FIORAVENTI'S Balsam, one ounce.

Mix.

BALSAMUM AD LORICANDUM CADAVER CONDITUM. (SPIELMANN)

R. Balsam of Peru, } of each,
 ——— Copaiba, } twenty-four
 Liquid Storax, } ounces.
 Oil of Nutmeg, } eight ounces.
 ——— Lavender, } two ounces.
 ——— Thyme, } half an ounce.

Melt slowly together.

STYRAX SOLIDUS.—STORAX.

Storax, Styrae Resina, Styrae Balsamum; Στίραξ. — Störax, Styrae solide, Fr.; Storax, Judenweyrauch, GERM.; Storax, DAN.; and SWED.; Usteruk, ARAB.; Estoraque, PORT.; and SPAN.; Styrae, DUT.; Storace, ITAL.

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HASS.; OLDEN.; LUSIT.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY.; and GME-LIN.; PIDERIT.; SPIELMANN.; ZARDA.

There are distinguished:

1. *Storax in tears; Styrae granulata seu in granis.* This sort is in small, shining, transparent grains, rather unctuous, softening between the teeth, of an agreeable odour and resinous taste.

2. *Storax in cakes; Styrae in massis.* This sort is in reddish, odoriferous masses, sometimes of a soft consistence, and separating a honey-like liquor, of a pleasant odour, and of a bitter and resinous taste.

The first variety, which is now very rare, was the only sort known to the ancients, who called it *Styrae calamita seu cannulata*, which name has been since applied to the second. This is called *Styrae amygd-*

lata when the masses are of the size of an almond.

This balsam is said to be produced by the *Styrae officinale*, LINN.; a tree of the south of Europe and of the East. (*Decandria, Monogynia*, LINN.; *Ebenacea*, JUSS.; *Fig. Enc. Meth. t. 369*) Perhaps it comes from the *liquid ambar orientale*, LMK.

It is stimulant.—Dose, from ten grains to half a dram.

PURIFIED STORAX.

Styrae purificata. (DUBL.)

Dissolve balsam of storax in rectified spirit, and strain the solution; then distil off the spirit with a gentle heat till the balsam becomes of a proper consistence.

PILLS OF STORAX.

Pilulae e Styrae. (DUBL.)

R. Purified Storax, three drams.

Soft Purified Opium, } of each,
 Saffron, } one
 } dram.

Mix well.

SUCCINUM.—AMBER.

SYNONIMES:—*Ambra flava*; *Ηλεκτρον*.—

Succin, Ambre jaune, Karabé, FR.; Agtstein, Bernstein, GERM.; Kernulbehr, ARAB.; Hambar, BAL.; Ambar, CYNG.; and MAL.; Beensteen, DAN.; Karuba, DUK.; and PERS.; Sucino, SPAN.; Kepur, HIND.; Barnsteen, DUT.; Ambra gialla, Succino, ITAL.; Barsztyn, POL.; Alumbre, PORT.; Jantar, RUSS.; Bernsten, SWED.; Umbir, TAM.; and TEL.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; COXE; GUIBOURT; PIDERIT; SPIELMANN.

A solid substance of various colours; some of a more or less deep yellow, some white, some red, and some of a reddish-brown approaching to black: it is diaphanous, translucent, or opaque; sometimes clouded. Its fracture is conchoid and brilliant; it is susceptible of polish, though not very hard, and it is fragile. It diffuses an agreeable odour, and becomes electric when rubbed; it burns with a yellow and greenish flame, diffusing a thick smoke and a strong and penetrating but not disagreeable odour.

FUMIGATORY POWDER.

Pulvis fumalis seu ad suffiandum seu pro Suffimigio. (BRUNS.; LIPP.; CAST. BORUS.)

R. Olibanum, } of each, one part.
Amber, }

Pulverize. (CAST. BORUS.)

R. Juniper, } of each, four ounces.
Mastic, }
Amber, }
Cascarilla, } one ounce.

Make a coarse powder. (LIPP.)

R. Amber, } twenty ounces.
Olibanum, } ten ounces.
Mastic, } four ounces.
Labdanum, }
Benzoin, } of each,
Storax, } two ounces.
Rhodium Wood, }
Yellow Sanders, }
Petals of Red Roses, } four ounces.
Flowers of Lavender, } three ounces.
Juniper, } twelve ounces.

Make a coarse powder. (BRUNS.)

ÆTHEREAL TINCTURE OF AMBER.

Tinctura Succini ætherea seu balsamica. (SAX.; SUEC.; WIRTEM.; HERBIP.; NIEMANN; SWEDIAUR; VAN MONS)

R. Amber, two ounces.
Sulphuric Æther, two pounds.

Digest for four days in a sand-bath, at a moderate heat, and strain. (HERBIP.)

R. Amber, powdered and slightly torrified, two ounces.

Alcohol, seven ounces and a half.

After several days' digestion in a covered vessel, filter, and add

Sulphuric Æther, two parts and a half.

Preserve. (NIEMANN)

R. Amber, powdered and slightly torrified, one ounce.

Spirit of Sulphuric Æther, five ounces.

Digest for four days in a covered vessel, and strain. (SWEDIAUR)

R. Amber, in coarse powder, one part.
Spirit of Sulphuric Æther, three parts.

Macerate without heat for eight days, with occasional agitation, and filter. (SAX.; VAN MONS)

SUEC. prescribes one ounce of amber, and four ounces of spirit.—WIRTEM.—one ounce of amber, and three and a half of spirit.—FULD.—one part of amber, and six parts of spirit.

It is excitant, and is reputed anti-spasmodic, nerve, anti-catarrhal, and uterine.—Dose, from fifteen to thirty drops, on sugar or with water.

ALCOHOLIC TINCTURE OF AMBER.

Tinctura seu Essentia Succini alcoholica, Alcohol cum Succino. (AMST.; ANT.; BELG.; BRUNS.; GALL.; GENEV.; HANN.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; WIRTEM.; VAN MONS)

R. White Amber, in powder, one ounce.
Alcohol (38 degrees), six ounces.

Digest for three days at a moderate heat, then decant, and pour upon the residue

Alcohol (38 degrees), two ounces.

Digest again for two days, and strain. Mix the tinctures, and filter. (ANT.)

R. White Sugar, one part.
Alcohol (32 degrees), eight parts.

Digest without heat for ten days; then strain and filter. (BELG.)

AMST. prescribes one part of amber, eight parts of alcohol (0.835), and eight days' digestion at a moderate heat.—WIRTEM.—one ounce and a half of amber, six ounces of concentrated alcohol, and several days' digestion without heat.—PALAT.—one ounce of amber, eight ounces of concentrated alcohol, and several days' digestion at a moderate heat.—LIPP.; and BRUNS.—one ounce of amber, one pound of alcohol, and eight days' digestion at a gentle heat.—GALL.—one part of amber, sixteen parts of alcohol (26 degrees), and six days' digestion, followed by a repose of six days.—GENEV.—one ounce of amber, ten ounces of alcohol, and eight days' digestion.

R. Amber, powdered and slightly
torrified in an iron pan, one part.
Concentrated Alcohol, four parts.

Digest for several days at a gentle heat in a covered vessel, occasionally agitating, and filter. (BRUNS.; HANN.; OLDEN.; POL.; BORUS.; SAX.; VAN MONS)

It is slightly stimulant, and is reputed anti-catarthal, balsamic, nervine, and vulnerary.—Dose, from forty to sixty drops.

SUCCINATED JULEP.

Julapium succinatum. (GENEV.)

R. Tincture of Amber, two drams.
Sugar, one ounce.
Water, six ounces.

ALKALINE TINCTURE OF AMBER.

Tinctura seu Essentia Succini alkalisata.
(BRUNS.; SAUNDERS; WIRTEM.; SPIELMANN)

R. Amber, in powder, two ounces.
Solution of Subcarbonate of Potass, half an ounce.

Make a paste; then, having dried it, pour upon it

Alcohol, eight ounces,
or, a sufficient quantity to cover it by three or four fingers' breadth. Digest for several days in a sand-bath, and filter. (BRUNS.; SARD.)

R. Amber, four ounces.
Subcarbonate of Potass, six ounces.

Triturate together. Add

Proof Spirit, sixteen ounces.

Macerate for several days, and distil about eight ounces of the liquor; then, having evaporated the rest to dryness, pour upon it the spirit first obtained, and after sufficient digestion, filter. (WIRTEM.)

R. Powdered Amber, } of each,
Subcarbonate of Potass, } twelve ounces.
Concentrated Alcohol, twenty-four ounces.

Distil the spirit, and infuse in it for eight days, in a water-bath, a powder composed of

Amber, eight ounces.
Deliquesced Subcarbonate of Potass, a sufficient quantity

to reduce it to a paste, to be then dried. Filter. (SPIELMANN)

It is excitant, and is reputed nervine, anti-catarthal, and resolvent.—Dose, from twenty to forty drops.

RESIN OF AMBER.

Resina Succini. (LIPP.)

R. Amber, two pounds.
Alcohol, four pounds.

Digest for three days in a sand-bath, and decant; treat the residuum in the same manner with more alcohol till this at last comes off colourless. Having mixed and distilled the tinctures, wash the resin, and dry it gently.

OIL OF AMBER.

Oleum Succini aethereum, Aetheroleum seu Pyrooleum Succini. (LOND.; DUBL.; EDINB.; AMER.; AMST.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SARD.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Oleum Succini.

Put amber into an alembic, and distil in a sand-bath, with a gradually raised heat, an acid liquor, an oil, and a salt impregnated with the oil. Re-distil the oil a second and third time. (LOND.)

R. Amber, in powder, } of each, an
Pure Sand, } equal part.

Mix them together in a glass retort which the mixture only fills to one half; then having adapted to it a large receiver, distil in a sand-bath with a gradually augmented heat. A watery liquor tinged with a little yellow oil will first come over; then a yellow oil with an acid salt, and lastly, a black and reddish oil. Pour the fluid from the receiver, and separate the oil from the watery liquor. (EDINB.)

R. Amber, bruised, any quantity.

Put it into a luted stoneware retort communicating with a large tubulated receiver. Distil with a gradually augmented heat till nothing more passes over, and separate the oil by means of a funnel.

This oil is only employed after it has been rectified. There are several processes for that purpose.

Oleum Succini purissimum. (EDINB.)

Distil oil of amber, mixed with six times its quantity of water, from a glass retort, till two thirds of the water have passed over into the receiver; then separate the purified volatile oil from the water, and keep it in well-stopped phials.

Oleum Succini Rectificatum. (DUBL.)

R. The Oil which comes over in the preparation of succinic acid, one pound.
Water, six pints.

Distil till two-thirds of the water have passed over into the receiver; then separate the oil.

R. Emphyreumatic Oil of Amber, any quantity.

Fill a retort with it to one half, and distil, by slight ebullition, till the oil begins to pass over more coloured. (BAVAR.; POL.; BRUGNATELLI; HENRI; VAN MONS)

POL. requires three successive rectifications.—GENEV.; and HERBIP.—four.

R. Oil of Amber, one part.
Wood Ashes, two parts.

Distil slowly. (BRUNS.; FERR.; HASS.; LIPP.; PALAT.; WIRTEM.; PIDERIT)

R. Oil of Amber, } of each, an equal part.
Charcoal, }

Distil in a glass retort. (FENN.; FULD.; SUEC.)

R. Oil of Amber, } of each, one part.
Charcoal, }
Water, six parts.

Distil slowly. (OLDEN.)

SARD. prescribes either pure chalk or burnt hartshorn.

R. Oil of Amber, one part.
Water, six parts.

Distil four parts. Separate the oil from the water. (AMER.; BATAV.; BELG.; DAN.; HANN.; BORUS.; SAX.; COXE; SWEDIAUR)

R. Oil of Amber, one pound.
Water, six pounds.

Distil slowly four ounces; then change the receiver, and distil five ounces more: re-distil this with water, and draw off three ounces more. Mix this second product with the first, and re-distil the whole with water. (AMST.)

HISP. requires the distillation of the oil with twelve parts of water, till it becomes perfectly limpid.

It is stimulant, anti-spasmodic, nervine, resolvent, diaphoretic, diuretic, and emmenagogue.—Dose, from five to fifteen drops.—It is not much used.

OLEOSACCHARUM OF AMBER.

Oleosaccharum Succini. (PALAT.; WIRTEM.)

R. White Sugar, one ounce.
Oil of Amber, sixteen drops.
Triturate together.

BALSAM OF AMBER.

Balsamum Succini. (WIRTEM.)

R. Oil of Amber, one dram.
— Nutmeg, two drams and a half.

Mix.—It is employed externally, in hysteria and cephalalgia.

ARTIFICIAL MUSK.

Moschus artificialis, Resina Succini balsamica, Oleum Succini oxydatum, Succinum oxygenatum. (AMER.; LIPP.; OLDEN.; SUEC.; BRUGNATELLI; COXE; NIEMANN; VAN MONS)

R. Oil of Amber, two parts.
Nitric Acid, six parts.

Digest for twenty-four hours, and wash the product with cold water. (LIPP.; NIEMANN)

AMER.; and COXE prescribe one part of oil and two parts of acid.—OLDEN.—two of oil and five of acid.—VAN MONS—one of oil and three of acid.—SUEC.; and BRUGNATELLI—one of oil and four of acid.

It is stimulant, and is recommended as a substitute for the true musk.

TINCTURE OF ARTIFICIAL MUSK.

Tinctura Moschi artificialis seu Resinae Succini seu Succini balsamica. (LIPP.; SUEC.; NIEMANN; VAN MONS)

R. Artificial Musk, one part.
Concentrated Alcohol, eight parts.

Dissolve at a moderate heat. (LIPP.; NIEMANN)

SUEC. prescribes one part of musk and

seven parts of alcohol.—VAN MONS—one of musk and sixteen of alcohol.

It is excitant, and is recommended in epilepsy.—Dose, forty drops.

FETID TINCTURE OF AMBER. (FERR.)

R. Rectified Oil of Amber, two drams.
Subcarbonate of Potass, } of each,
Medicinal Soap, } one dram.
Alcohol, } four ounces.

Filter at the expiration of a few days.

By adding a few drops of this tincture to solution of ammonia, a kind of *Eau de Luce* is immediately produced.

EXPECTORANT POTION. (ELLIS)

R. Oil of Amber, eighty drops.

Gum Arabic, } of each, half an ounce.
Sugar, }

Tincture of Balsam of Tolu,

Water, } two drams.
four ounces.

Dissolve, and mix.—Dose, one spoonful, every two or three hours, in whooping cough.

SUCCINATED ANODYNE LIQUOR.

Liquor succinicus anodymus. (SWEDIAUR)

R. Rectified Oil of Amber, one pound and a half.
Concentrated Sulphuric Acid, half a pound.

Distil one pound in a glass retort.—It is extolled, in Germany, against epilepsy.—Dose, twenty or thirty drops, in syrup of wormwood or of mint.

SULPHUR; *Θείον*.—SULPHUR.

SYNONIMES: — Soufre, FR.; Schwefel, GERM.; Kabrit, ARAB.; Lew, CHIN.; Gundaka, CYNG.; Svolv, DAN.; Gunduck, DUK.; Azufre, SPAN.; Gundhuk, HIND.; Schwavel, DUT.; Zolfo, ITAL.; Gunduk, MAH.; Blerong, MAL.; Gowgird, PERS.; Siarka, POL.; Enxofre, PORT.; Sura, RUSS.; Ganhak, SANS.; Skafvel, SWED.; Gendagum, TAM.; Ghendagum, TEL.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE.; COXE.; GUIBOURT.; PARIS.; PIDERIT.; SAUNDERS.; SPIELMANN.

A solid substance of a lemon-yellow colour, very friable at the least concussion, and presenting a brilliant fracture. It is crystallizable in needles or in octohædra, fusible, volatilizable, and inflammable; insipid and inodorous, but exhaling a slight odour when rubbed. It is met with in commerce, under two forms:

1. *Roll Sulphur*; in cylinders of a more or less dull and greyish yellow, almost always containing needle-shaped crystals in their centre.

2. *Flowers of Sulphur, Sublimed Sulphur: Flores Sulphuris*; in fine powder, of a beautiful yellow colour.

The pharmacopœias of ANT.; BRUNS.;

HISP.; FERR.; FULD.; HASS.; PALAT.; SARD.; WIRTEM.; HERBIP.; PIDERIT.; SWEDIAUR; and VAN MONS are the only ones which require the apothecary himself to prepare it by subliming the sulphur of commerce, either alone or previously mixed with washed sand. They direct it to be well washed afterwards in order to free it from any sulphureous acid which may be formed.

This washing is required by the other pharmacopœias who direct the flowers of sulphur of commerce to be used. It is then called *Sulphur depuratum* sen. *lotum*, *Flores Sulphuris loti*.

WASHED SULPHUR.

Sulphur lotum. (LOND.)

R. Sublimed Sulphur, one pound.

Pour boiling water upon it, that whatever acid may be present with it, may be entirely washed away; then dry it.

Sulphur Sublimatum lotum.

R. Sublimed Sulphur, one part.
Water, four parts.

Boil the sulphur in the water for a short time; then pour off this water, and by repeated effusions of cold water, wash away all the acid; finally dry the sulphur. (EDINB.)

Pour warm water upon sublimed sulphur repeatedly till the water employed ceases to acquire acidity, which may be ascertained by testing with litmus.

Dry the sulphur on bibulous paper. (DUBL.)

There is also another process for the obtaining of pure sulphur, the product of which process is called *Precipitated Sulphur*, *Magistery of Sulphur*, *Milk of Sulphur*, *Hydrate of Sulphur*; *Sulphur præcipitatum*, *Lac seu Magisterium sulphuris*. (LOND.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; HANN.; LIPP.; OLDEN.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI.; COXE.; PIDERIT.; SPIELMANN.; VAN MONS). This process consists in decomposing the aqueous solution either of sulphuret of potass, or of sulphuret of soda (ANT.), or of sulphuret of lime. (LOND.; COXE), with hydrochloric acid, carefully washing and drying the precipitate.

PRECIPITATED SULPHUR.

Sulphur præcipitatum. (LOND.; COXE)

R. Sublimed Sulphur, one pound.
Fresh-burnt Lime, two pounds.
Water, four gallons.

Boil the sulphur and lime together in the water; then filter the liquor through paper, and drop into it Muriatic Acid,

a sufficient quantity

to precipitate the sulphur. Wash this repeatedly with water till it becomes tasteless.

Sulphur is an excitant, which, according to the dose and particularly according to the subject, acts as a stimulant, expectorant, diaphoretic, diuretic, or purgative. It is one of the most powerful and most valuable agents in the materia medica, and is employed under a great number of different forms, and to fulfil numerous indications.

Dose, from twelve to twenty grains twice or thrice a day, as a stimulant, and one or two drams and more, as a purgative.

CARBURET OF SULPHUR.

Alcohol Sulphuris. (VAN MONS)

R. Sulphuret of Iron, in powder,
five parts and a half.
Charcoal, one part and a half.

VOL. II.

Mix by continued trituration, and fill a retort with the mixture to one fourth; distil by heating to incandescence, and set apart the liquid which collects at the bottom of the water contained in the receiver.

Some experiments which have been made with this substance do not appear to have succeeded.

DENTIFRICE POWDER. (PIERQUIN)

R. Sublimed Sulphur, one ounce.
Lac, in powder, a sufficient quantity.
Make a very fine powder.

ANTI-PSORIC POWDER.

Pulvis contra Scabiem. (HAMB.; HERBIP.; AUGUSTIN.; CADET DE GASSICOURT.; VAN MONS)

R. Sulphur, } of each,
Root of Liquorice, } six drams.
Camphor, half a dram.

Mix.—Dose, half a dram twice a day. (AUGUSTIN)

R. Sublimed Sulphur, } of each,
Charcoal, } an equal part.
Brick, }

Make a powder. (CADET DE GASSICOURT)

R. Sublimed Sulphur, } of each, half
Sulphuret of Antimony, } an ounce.
Magnesia, two drams.

Make a powder. (HAMB.)

R. Sublimed Sulphur, } of
Sulphuret of Mercury and of } each,
Antimony, } one
Mercury, } ounce.

Make a very fine powder. (HERBIP.)

R. Nitrate of Potass, } of each,
Sugar, } two drams.
Supertartrate of Potass, six drams.
Sublimed Sulphur, one ounce and a half.

Make a powder. (VAN MONS)

The second of these powders is only applied externally incorporated with a little oil. The two succeeding may be used both externally and internally. The last is for internal use.

DIAPHORETIC POWDER. (HUFELAND; SAINTE-MARIE)

R. Sublimed Sulphur, four drams.
Camphor, one scruple.

Make twelve parcels.—Dose, one or two daily, in a cup of tea. (SAINTE-MARIE)

R. Sulphur, two drams.
Red Sulphuret of Antimony, } of each,
Camphor, } eight grains.
Sugar, two scruples.

4 M

Mix.—Dose, half a dram, three or four times a day. (HUFELAND)

ANTI-PLEURITIC POWDER.

Pulvis pleuriticus Mynsichti. (WIRTEM.)

R. Sublimed Sulphur, half an ounce.
Wild Boar's Teeth, } of each,
Pike's Jaws, } one ounce.
Flowers of White Poppy, half an ounce.

Make a powder.—Dose, from a scruple to half a dram.—It was to the osseous particles that superstitious credulity attributed the reputed efficacy of this powder, which however is not absolutely inert as those are into which only the phosphate of lime enters. The sulphur renders it excitant.

LAXATIVE POWDER. (ELLIS ; RATIER)

R. Sublimed Sulphur, } of each,
Supertartrate of Potass, } one ounce.
White Sugar, a sufficient quantity.

Mix. (RATIER)

ELLIS prescribes half an ounce of sulphur and one ounce of supertartrate.

This powder is mixed with honey.

Delicate stomachs better support a mixture at equal parts of precipitated sulphur and of calcined magnesia, a spoonful of which mixture may be taken four or five times a day in milk. (ELLIS)

INCISIVE POWDER.

Poudre anti-asthmatique. (GALL. ; FORM. HÔP. FR.)

R. White Sugar, three parts.
Sublimed Sulphur, two parts.
Dried Squill, one part.

Mix.

ANTI-DYSENTERIC POWDER. (VAN MONS)

R. Sublimed Sulphur, eight parts.
Fennel, one part.
Gum Arabic, } of each, sixteen parts.
White Sugar, }

Mix.

ABSORBENT PURGATIVE POWDER.

Pulvis catharticus absorbens. (SWEDIAUR)

R. Subcarbonate of Lime, } of each,
Sublimed Sulphur, } one ounce.

Mix.—Dose, from ten to twenty grains, to children, and from half a dram to one dram, to adults.

ANTI-CATARRHAL POWDER.

Pulvis sulphureus seu sulphureus seu catarrhalis. (FULD. ; HAMB. ; CAST. BO-

RUS. ; CADET DE GASSICOURT ; HUFELAND)

R. Sublimed Sulphur, } of each,
Liquorice, } six drams.
Camphor, half a dram.

Make a powder. (CAST. BORUS.)

R. Sublimed Sulphur, } of each,
Nitrate of Potass, } two drams.
Liquorice, }
Fennel, }

Mix. (FULD.)

HAMB. gives the same formula, but substitutes hydrochlorate of ammonia for the nitre.

R. Sublimed Sulphur, two drams.
Supertartrate of Potass, six drams.
Red Sulphuret of Antimony, fifteen grains.

Make a powder. (HAMB.)

R. Sugar, three ounces.
Florentine Iris, nine scruples.
Sublimed Sulphur, four scruples.
Saffron, one dram.
Gum Arabic, } of each, one scruple.
Starch, }
Liquorice, }

Make a powder. (CADET DE GASSICOURT)

R. Sublimed Sulphur, one ounce and a half.
Fennel, } of each, one ounce.
Florentine Iris, }
Liquorice, }
Senna, six drams.

Make a powder. (HUFELAND)

Dose, one or two drams.

RESOLVENT PECTORAL POWDER.

Pulvis pectoralis resolvens. (BRUNS. ; PALAT. ; WIRTEM.)

R. Simple Diaireos Species, } of each,
Cold Diatragacanth } half an ounce.
Species, }
Liquorice, } of each,
Marshmallows, } three drams.
Armenian Bole, } of each,
Sublimed Sulphur, } two drams.
Terra Sigillata, one dram and a half.
White Sugar Candy, six ounces.

Make a powder. (PALAT.)

R. Root of Arum, } of each,
Florentine Iris, } half an ounce.
Deutoxide of Antimony, }
Sublimed Sulphur, }
Native Cinnabar, }
Crabs' Eyes, }

Make a powder. (BRUNS. ; WIRTEM.)

Dose, from a scruple to half a dram.

BALSAMIC PECTORAL POWDER.

Pulvis pectoralis seu pectoralis balsamicus.
(BRUNS.; WIRTEM.; BORIES; CADET
DE GASSICOURT; NIEMANN; SPIEL-
MANN; SWEDIAUR)

- R. Root of Liquorice, one ounce.
 — Florentine Iris, two drams.
 Sublimed Sulphur, half an ounce.
 Benzoic Acid, one scruple.
 White Sugar, two ounces.
 Oil of Anise, } of each, ten drops.
 — Fennel, }

Make a powder. (BRUNS.; WIRTEM.; NIEMANN)

SPIELMANN prescribes six drams each, of liquorice and of iris, one ounce of sulphur, one dram of benzoic acid, two ounces of sugar, and ten drops each, of the two oils.—CADET DE GASSICOURT; and SWEDIAUR—two drams each, of liquorice and of iris, half an ounce of sublimed sulphur, one scruple of benzoic acid, two ounces of sugar, and ten drops each, of the two oils.

- R. Benzoic Acid, six grains.
 Sublimed Sulphur, ten grains.
 Cascarilla, twelve grains.
 Sugar Candy, twenty grains.

Mix. (PIERQUIN)

This last is for one dose.

BOLUS OF SULPHUR. (FORM. HÔP. FR.)

- R. Washed Sulphur, eight grains.
 Honey, a sufficient quantity.
 Make a bolus.

COMPOUND BOLUS OF SULPHUR. (FORM. HÔP. FR.)

- R. Washed Sulphur, } of each,
 Anise, in powder, } eight grains.
 Senna, in powder, } four grains.
 Honey, a sufficient quantity.

Make a bolus.

DIAPHORETIC BOLUSES. (CADET DE GASSICOURT)

- R. Sublimed Sulphur, } of each,
 Supertartrate of Potass, } one scruple.
 Guaiac Resin, ten grains.
 Simple Syrup, a sufficient quantity.

Make two boluses.—One to be taken night and morning, in diseases of the skin and chronic rheumatism.

BOLUSES AGAINST CYSTITIS.

Boli in Cystitide lenta. (BRERA)

- R. Sublimed Sulphur, one dram.
 Venice Turpentine, two scruples.

Make six boluses.—Dose, one every two

or three hours.—They are said to be particularly serviceable when some cutaneous disease exists as a cause or complication of the complaint.

ANTI-ASTHMATIC BOLUSES. (CADET DE GASSICOURT)

- R. Conserve of Smallage, } of each,
 Extract of Elecampane, } one dram.
 Sublimed Sulphur, } of each,
 Gum Ammoniac, } half a dram.
 Oxymel of Squill, a sufficient quantity.

Make boluses of ten grains each.—Dose, one, two, or three, five or six times a day.

ANTI-HÆMORRHOIDAL BOLUSES. (CADET DE GASSICOURT; SAUNDERS; SAINTE-MARIE)

- R. Sublimed Sulphur, four scruples.
 Compound Electuary of Rhubarb, eight scruples.

Make four boluses. (CADET DE GASSICOURT; SAINTE-MARIE)

- R. Sublimed Sulphur, two scruples.
 Confection of Senna, two drams.
 Syrup of Buckthorn, a sufficient quantity.

Make two boluses. (SAUNDERS)

Dose, one every morning.

TROCHISCS OF SULPHUR.

Rotule pectorales, Trochisci e Sulphure.
(AMST.; BRUNS.; GALL.; AUGUSTIN;
CADET DE GASSICOURT; PIERQUIN;
SAUNDERS; SWEDIAUR; VAN MONS)

- R. Sublimed Sulphur, one part.
 White Sugar, eight parts.
 Mucilage of Tragacanth, a sufficient quantity.

Make trochiscs. (GALL.)

PIERQUIN prescribes one part of sublimed sulphur, and six parts of sugar.—PIERQUIN, in another place; AMST.; and SAINTE-MARIE—one of sulphur, and two of sugar.

- R. Sublimed Sulphur, one part.
 White Sugar, twelve parts.

Triturate the sulphur with one part of sugar, and incorporate with the rest of the sugar boiled *à la plume*. (VAN MONS)

- R. Powdered Florentine Iris, half an ounce.
 — Liquorice, eight scruples.
 — Angelica, two scruples.
 Sublimed Sulphur, one dram.
 Extract of Elecampane, } of each,
 — Horehound, } two scruples.
 — Saffron, one scruple.

White Sugar, sixteen ounces.
Essential Oil of Fennel, fifteen drops.
Make trochiscs. (BRUNS.)

- R. Powdered Liquorice, } of each,
Florentine Iris, } one dram.
Sublimed Sulphur, }
Saffron, } ten grains.
White Sugar, half an ounce.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs of half a scruple each.
(AUGUSTIN)

- R. Sublimed Sulphur, two drams.
Benzoic Acid, twelve grains.
Florentine Iris, half a dram.
Oil of Anise, twelve drops.
White Sugar, five ounces and a half.
Mucilage of Tragacanth,
a sufficient quantity.

Make trochiscs. (GALL.; CADET DE
GASSICOURT)

- R. Pulp of Marshmallows } of each,
Root, } one ounce.
Great Comfrey }
Root,
Sublimed Sulphur, two drams.
Benzoic Acid, half a dram.
Sugar dissolved in White Lily Water,
eight ounces.

Make trochiscs. (PIERQUIN)

- R. Precipitated Sulphur, two ounces.
Benzoic Acid, one scruple.
White Sugar, four ounces.
Mucilage of Gum Arabic,
a sufficient quantity.

Make trochiscs. (SWEDIAUR)

- R. Oxalic Acid, one part.
Sublimed Sulphur, sixteen parts.

Incorporate in

Sugar, melted at a moderate heat,
one hundred and twenty-eight parts.

Make trochiscs. (VAN MONS)

ANTI-PSORIC TROCHISCS.

Morsuli ad Scabiem. (SPIELMANN)

- R. Sublimed Sulphur, one ounce.
Sulphuret of Antimony, } of each,
Species Imperatoris, } two drams.
Sweet Almonds, cut, one ounce.
White Sugar, boiled *à la plume*,
eight ounces.
Essence of Lemon, ten drops.

Make lozenges.

ELECTUARY OF SULPHUR. (HENRI;
RATIER)

- R. Sublimed Sulphur,
one hundred and sixty-six parts,

Honey, three hundred and fifty parts.

Mix. (HENRI)

RATIER prescribes one part of honey and
three parts of sulphur.

PURGATIVE ELECTUARY. (ELLIS)

- R. Sublimed Sulphur, half an ounce.
Confection of Senna,
one ounce and a half.
Simple Syrup, a sufficient quantity.
Mix. — Dose, a tea spoonful night and
morning.

PECTORAL MARMALADE.

Conditum pectorale. (SAINTE-MARIE;
VAN MONS)

- R. Sublimed Sulphur, } of each,
Fennel, } half an ounce.
Liquorice, } one ounce.
Syrup of Red Poppy,
a sufficient quantity.

Mix. (VAN MONS)

- R. Honey, eight ounces.
Sublimed Sulphur, } of each,
Root of Elecampane, } two drams.
Extract of Liquorice, }
Rose Water, }

Mix. (SAINTE-MARIE)

- R. Sublimed Sulphur, } of each,
Saffron, } two drams.
Rob of Elder, }
Extract of Liquorice, }
Honey, four ounces.

Mix. (SAINTE-MARIE)

Dose, the volume of a nutmeg, three or
four times a day.

ANTI-RHEUMATISMAL ELECTUARY. (NIE-
MANN)

- R. Resin of Guaiac, } of each,
Rhubarb, } two drams.
Supertartrate of Potass, one ounce.
Sublimed Sulphur, two ounces.
Nutmeg, one.
Despumated Honey, one pound.

Mix.

ANTI-PHTHISICAL ELECTUARY. (NIE-
MANN; PIERQUIN)

- R. Spermaceti, } of each,
Sublimed Sulphur, } two drams.
Crab's Eyes, }
Conserve of Roses, half an ounce.
Honey, a sufficient quantity.

Mix. — Dose, one or two drams in the
course of the day. — DUFRESNOY has pro-
posed to add three drams of edible mush-
room (*Galarhaus piperatus*, FRIES.) to ren-
der it more efficacious.

EXPECTORANT LOHOCH. (HANN.; BORIES;
PIERQUIN; SAINTE-MARIE)

R. Sublimed Sulphur, one dram.
Sugar, two drams.
Linseed Oil, one ounce.
Balsamic Syrup, one ounce and a half.

Mix. (HANN.)

R. Syrup of Violets, one ounce and a half.
Oxymel, one ounce.
Sublimed Sulphur, half a dram.
Benzoic Acid, fifteen grains.

Mix. (BORIES; PIERQUIN)

R. Honey, } of each,
Pulp of Raisins, } two ounces.
Sublimed Sulphur, } of each, one dram.
Benzoic Acid. }
Syrup of Wild Mustard,
a sufficient quantity.

Mix. (PIERQUIN)

SAINTE-MARIE gives the same formula, but substitutes the pulp of rennet apples for that of raisins.

SULPHUROUS POTION.

Mixtura Sulphuris præcipitati. (AUGUSTIN)

R. Precipitated Sulphur, two drams.
Fennel Water, three ounces.
Cinnamon Water, one ounce.
Syrup of Opium, half an ounce.

Mix.—Dose, one spoonful every two hours; in diabetes and in affections from the abuse of mercury.

TINCTURE OF SULPHUR. (WIRTEM.)

R. Sulphur, one ounce and a half.

Fuse it slowly in an earthen dish, and add gradually to it,

Subcarbonate of Potass, three ounces.

Increase the heat till the mass flows; then pour it into a heated mortar: reduce it to powder, and add,

Concentrated Alcohol, ten ounces.

Digest and strain.

It is stimulant, and is recommended in affections of the chest, itch, and scorbutic ulcers.—Dose, fifty or sixty drops in an appropriate vehicle.

SIMPLE BALSAM OF SULPHUR.

Balsamum Sulphuris simplex, Oleum sulphuratum, Linimentum e Sulphure. (LOND.; EDINB.; AMST.; ANT.; BELG.; BRUNS.; DAN.; HISP.; FULD.; HANN.; LIPP.; OLDEN.; PALAT.; BORUS.; SARD.; SAX.; WIRTEM.; BRUGNATELLI; CADET DE GASSICOURT; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Oleum Sulphuretum.

R. Washed Sulphur, two ounces.
Olive Oil, one pint.

Add the sulphur gradually to the oil heated in a very large iron pot, stirring the mixture after each addition till they perfectly unite. (LOND.)

R. Olive Oil, eight parts.
Sublimed Sulphur, one part.

Boil with a gentle heat in a large iron vessel, stirring constantly, till they unite. (EDINB.)

R. Sublimed Sulphur, one part.
Oil of Almonds, six parts.

Digest in a sand-bath for several days, and decant. (SARD.; WIRTEM.)

SPIELMANN prescribes the same proportions, but requires the oil of walnuts instead of that of almonds.—ANT.; BELG.; DAN.; FERR.; HANN.; OLDEN.; BORUS.; SAX.; and VAN MONS—one part of sulphur and four parts of linseed oil.—BRUNS.—one of sulphur and four of oil of almonds.—FULD.; and COXE—one of sulphur and four of olive oil.—HISP.; and PIDERIT—one of sulphur and four of walnut oil.—LIPP.—one of sulphur and four of linseed oil.—CADET DE GASSICOURT—one of sulphur and three of walnut oil.—BRUNS.—one of sulphur and two of linseed oil.—AMST.; BRUGNATELLI; and SWEDIAUR—one of sulphur and eight of olive oil.—SWEDIAUR—an equal part each, of sulphur and of olive oil.

R. Sublimed Sulphur, two ounces.
Oil of Clove-pink, one pound and a half.
Wine, half a pound.

Digest for eight days; then boil to the consumption of the humidity. (PALAT.)

ANISATED BALSAM OF SULPHUR.

Balsamum Sulphuris anisatum. (ANT.; BRUNS.; DAN.; HISP.; GALL.; LIPP.; PALAT.; SARD.; WIRTEM.; CADET DE GASSICOURT; NIEMANN; PIERQUIN; VAN MONS)

R. Sublimed Sulphur, one part.
Oil of Anise, four parts.

Digest in a sand-bath. (ANT.; BRUNS.; HISP.; GALL.; CADET DE GASSICOURT; PIERQUIN; VAN MONS)

PALAT.; SARD.; and WIRTEM. prescribe one part of sulphur and six parts of oil.

R. Simple Balsam of Sulphur, one part.
Oil of Anise, five parts.

Mix. (DAN.; LIPP.; NIEMANN)

EMULSION OF ANISATED BALSAM OF SULPHUR. (VAN MONS)

- R. Gum Arabic, two drams.
 Anisated Balsam of Sulphur, one dram.
 Honey, one ounce.
 Hyssop Water, eight ounces.

Mix.

SUCCINATED BALSAM OF SULPHUR.

Balsamum Sulphuris succinatum. (HISP.; SARD.; VAN MONS)

- R. Rectified Oil of Amber, six parts.
 Sublimed Sulphur, one part.

Digest in a sand-bath. (SARD.; VAN MONS)

HISP. prescribes one part of the oil and four parts of sulphur.

TEREBINTHINATED BALSAM OF SULPHUR.

Balsamus Vitæ Rulandi, Balsamum sulphuris terebinthinatum, Oleum Terebinthine sulphuratum. (ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; FERR.; HANN.; LIPP.; OLDEN.; PALAT.; BORUS.; SARD.; SUEC.; WIRTEM.; HERBIP.; CADET DE GASSICOURT; PIDERIT; VAN MONS)

- R. Sublimed Sulphur, one part.
 Spirit of Turpentine, four parts.

Dissolve in a sand-bath. (HISP.; FERR.; VAN MONS)

PALAT.; SARD.; WIRTEM.; CADET DE GASSICOURT; BRUNS.; and HERBIP. prescribe one part of sulphur and five parts of spirit.

- R. Sublimed Sulphur, two parts.
 Linseed Oil, six parts.

Dissolve, and add to three parts of the solution.

- Spirit of Turpentine, eight parts.
 Mix. (SUEC.)

ANT.; BATAV.; BELG.; HANN.; OLDEN.; and BORUS. prescribe one part of simple balsam of sulphur and three parts of spirit.—PIDERIT—one of balsam and four of spirit.—DAN.—one of balsam and eight of spirit.—LIPP.—one of balsam and five of spirit.—FERR.—three of balsam and thirty-two of spirit.

- R. Sulphur, two ounces.
 Spirit of Turpentine, three ounces.
 Linseed Oil, two pounds.

Dissolve the sulphur in the oil; then add the spirit, and leave the mixture to digest in a very gentle heat. (BRUNS.; WIRTEM.)

It is excitant, and was formerly reputed diuretic.—Dose, from six to twelve drops.

ÆTHEREAL BALSAM.

Balsamum æthereum. (AUGUSTIN)

- R. Terebinthinated Balsam } of each,
 of Sulphur, } half a dram.
 Sulphuric Æther, }
 DIPPEL's Animal Oil, six drops.

Mix.—It is extolled by BUCKING, in what he calls pituitous colic, attributed to atony of the stomach.—Dose from two to ten drops, thrice a day, on sugar or in a little water.

CERATE OF SULPHUR.

Ceratum e Sulphure. (RATIER; SWEDIAUR)

- R. Sublimed Sulphur, four ounces.
 Cerate, one pound.
 Mix. (RATIER)

- R. Sublimed Sulphur, } of each,
 Bay Berries, } one ounce.
 Fresh Butter, a sufficient quantity.

Triturate together.

SIMPLE OINTMENT OF SULPHUR.

Unguentum Sulphuris seu e Sulphure seu sulphuratum simplex seu ad Scabiem. (LOND.; DUBL.; EDINB.; AMER.; BAVAR.; HAMB.; HANN.; LUSIT.; CAST. BORUS.; BORUS.; SAX.; AUGUSTIN; CADET DE GASSICOURT; COXE; RATIER; SWEDIAUR)

Unguentum Sulphuris.

- R. Sublimed Sulphur, three ounces.
 Prepared Lard, half a pound.
 Mix. (LOND.)

EDINB.; and DUBL. prescribe one part of sulphur and four parts of lard.

HANN.; CAST. BORUS.; BORUS.; SAX.; RATIER; and SWEDIAUR prescribe one part of sulphur and two parts of lard.—BAVAR.—one of sulphur and three of lard.—AMER.; and COXE—one of sulphur and four of lard.—HAMB.—one of sulphur and eight of lard.—AUGUSTIN—one of sulphur and a sufficient quantity of lard.

- R. Sulphur, one part.
 Rose Ointment, two parts.
 Triturate together. (LUSIT.)

- R. Walnut Oil, half a pound.
 Sublimed Sulphur, two ounces.

Dissolve in a water-bath, and add

- Lard, half a pound.
 Yellow Wax, one ounce.
 Alkanet, a sufficient quantity,

to colour the mass. Strain through a linen cloth. (CADET DE GASSICOURT)

SULPHURO-SAPONACEOUS OINTMENT.

Linimentum psoricum. (MIL. COPEN; AUGUSTIN; HENRI; NIEMANN; RATIER)

- R. Sublimed Sulphur, } of each, one part.
 White Soap, }
 Water, three parts.

Dissolve the soap in the water; then strain, and add the sulphur, assiduously stirring. (HENRI; RATIER)

- R. Subcarbonate of Potass, two ounces.
 Boiling Water, one ounce.
 Olive Oil, four ounces.

Make a soap, and add gradually to it, constantly stirring,

Sublimed Sulphur, five ounces.

Mix. (RATIER)

- R. Soft Soap, two parts.
 Sublimed Sulphur, one part.

Triturate together. (MIL. COPEN.)

NIEMANN prescribes an equal part each, of sulphur and of soap, and a sufficient quantity of water.

- R. Sublimed Sulphur, } of each,
 Soft Soap, } half a pound.
 Root of White Hellebore, two ounces.
 Nitrate of Potass, one dram.
 Lard, one pound and a half.

Mix. (AUGUSTIN)

SULPHURO-SAPONACEOUS LOTION. (RATIER)

- R. White Soap, three ounces.
 Water, fifteen pounds.

Dissolve, and add

Sulphur, three ounces.

Mix.

COMPOUND OINTMENT OF SULPHUR.

Unguentum sulphuratum seu Sulphuris seu Sulphuris compositum seu anti-psoricum seu contra Scabiem seu ad Scabiem. (LOND.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; DAN.; HISP.; GALL.; FORM. HÔP. FR.; FULD.; HANN.; LIPP.; POL.; BORUS.; ROSS.; SAX.; SARD.; WURCEB.; AUGUSTIN; BORIES; BRUGNATELLI; CADET DE GASSICOURT; COXE; PIDERIT; PIERQUIN; RATIER; SAUNDERS; SAINTE-MARIE; SPIELMANN; SWEDIAUR; VAN MONS)

Unguentum Sulphuris compositum. (LOND.; HANN.)

- R. Sublimed Sulphur, half a pound.
 White Hellebore Root, in powder, two ounces.

Nitrate of Potass, one dram.
 Soft Soap, half a pound.
 Prepared Lard, one pound and a half.

Mix.

SWEDIAUR prescribes one pound of sulphur, three ounces of white hellebore, one ounce and a half of nitre, one pound of soap, and three pounds of lard.

- R. Sublimed Sulphur, one ounce.
 White Precipitate, } of each,
 Benzoic Acid, } one dram.
 Oil of Lemon, one ounce.
 Sulphuric Acid, sixteen grains.
 Nitrate of Potass, two drams.
 Lard, half a pound.

Melt the lard and incorporate the other ingredients, constantly stirring till cold. (AMER.; COXE)

- R. Lard, eighteen ounces.
 Rose Oil, six ounces.
 Sublimed Sulphur, nine ounces.
 Powdered Cloves, } of each, one
 Grains of } ounce and a half.
 Paradise,
 Ginger,

Mix. (SPIELMANN)

- R. Lard, one pound.
 Extract of Ox-gall, two ounces.
 Sublimed Sulphur, } of each,
 Hepatic Aloes, } half an ounce.
 Colocynth, }
 Oil of Wormwood, three drams.

Melt the lard, and incorporate. (BRUGNATELLI)

- R. Juice of Elaterium, one part.
 Coagulate it over a slow fire; then remove the fecula, let it drip, and mix it with
 Olive Oil, one half part.
 Butter, one fourth part.

Evaporate to the consumption of the humidity; then triturate, and add

- Wax, one tenth part.
 Extract of Bile, } of each,
 Galbanum, } one ninetieth part.

Add to the mass when half cold,

- Scammony, } of each,
 Aloes, } one fiftieth part.
 Colocynth, }
 Mezereon Bark, }
 Euphorbium, } of each,
 Myrrh, } one hundredth part.
 Long Pepper, }
 Ginger,

Mix. (BRUGNATELLI)

- R. Sublimed Sulphur, } of each,
 Soap, } half a pound.
 Root of Black Hellebore, two ounces.
 Lard, two pounds and a half.

Mix. (SAX.)

- R. Sublimed Sulphur, one ounce.
Powdered White Hellebore, two drams.
Lard, two ounces.
Mix. (ANT.)

- R. Sublimed Sulphur, sixteen parts.
Muriate of Ammonia, one part.
Lard, thirty-two parts.

Mix. (PIDERIT)

AMST.; and NIEMANN prescribe eight parts of sulphur, one part of muriate and sixteen parts of lard.—RATIER—two ounces of sulphur, one dram of muriate, four ounces of lard, and one scruple of essence of lemon.—PIERQUIN—two ounces of sulphur, two drams of muriate, two ounces of lard, and a sufficient quantity of oil of orange flowers.—AUGUSTIN—three ounces of sulphur, two drams of muriate, and a sufficient quantity of lard to form an ointment.

- R. Sublimed Sulphur, two parts.
Decrepitated Muriate of Soda, one part.
Lard, eight parts.

Melt the lard, and incorporate. (FORM. HÔP. FR.)

- R. Root of Dock, eight ounces.
Leaves of Fumitory, } of each,
Bay, } half a handful.
Tobacco, }
Water, a sufficient quantity.

Boil; then bruise, and pass the pulp through a sieve. Add to six ounces of it

- Sublimed Sulphur, half an ounce.
Decrepitated Muriate of Soda, six drams.
Lard, three ounces.

Triturate together. (BORIES)

- R. Sublimed Sulphur, two ounces.
Subcarbonate of Lead, half an ounce.
White Precipitate, two drams.
Lard, washed with rose water, one pound.

Mix. (SARD.)

- R. Sublimed Sulphur, } of each, one part.
Sulphate of Zinc, }
Bay Berries, }
Rose Ointment, four parts.

Mix well. (HANN.)

SAINTE-MARIE prescribes one part each, of sulphur, of the sulphate, and of the berries, and a sufficient quantity of linseed oil.

- R. Sublimed Sulphur, } of each, one part.
Sulphate of Zinc, }
Oil of Bay, }
Lard, six parts.

Triturate together. (SAX.)

HISP. prescribes two parts each, of sulphur and of sulphate, and four each, of oil of bay and of lard.—BATAV.; BELG.; and

LIPP.—two of sulphur, one of sulphate, three of oil, and six of lard.—SWEDIAUR—an equal part each, of sulphur, of sulphate, and of oil, without lard.—FULD.—two of sulphur, one of sulphate, and six of lard, without oil.—POL.; and BORUS.—two each of sulphur and of sulphate, and eight of lard, without oil.—AUSTR.—one ounce and a half each, of sulphur and of sulphate, and one pound of simple ointment.

- R. Sublimed Sulphur, two parts.
Oxide of Zinc, one part and a half.
Oil of Bay, three parts.
Lard, six parts.

Mix. (VAN MONS)

- R. Sublimed Sulphur, } of each,
Sulphate of Zinc, } six parts.
Lard, sixteen parts.

Triturate together. (WURCEB)

DAN.; and ROSS. prescribe two parts of sulphur, one of sulphate, and twelve of lard.

- R. Sublimed Sulphur, two parts.
Subcarbonate of Potass, one part.
Lard, eight parts.

Triturate together. (GALL.; CADET DE GASSICOURT; SWEDIAUR)

SAUNDERS prescribes two ounces of sulphur, half an ounce of subcarbonate, and half a scruple of essence of lemon.—DAN.—one ounce of sulphur, one ounce of bay berries, two ounces of root of elecampane, half an ounce of subcarbonate, and ten ounces of lard.

ANTI-HERPATIC OINTMENT. (PIERQUIN)

- R. Sublimed Sulphur, one dram.
Pulp of Dock Root, one ounce.
Lard, two ounces.
Lemon Juice, two spoonfuls.

Mix.

- R. Sublimed Sulphur, } of each,
Benzoin, } two ounces.
Fresh Butter, a sufficient quantity.

Triturate together.

OINTMENT AGAINST TINEA. (BORIES)

- R. Sublimed Sulphur, } of each
Powdered Charcoal, } two ounces.
Soot, } of each,
Cinchona, } one ounce.
Cerate, a sufficient quantity.

Triturate together.

SULPHUR PLASTER.

Emplastrum sulphuratum resinosum seu Guillermi servitoris. (HISP.)

- R. Resin, four pounds.
Yellow Wax, one pound.
Turpentine, eight ounces.

Add to the melted mixture,

Powdered Ginger, one ounce and a half.	} of each, one ounce.
———— Bay Berries,	
———— Sulphur,	
———— Anise,	
———— Pennyroyal,	

Powdered Olibanum,	} of each, one ounce. two ounces.
———— Saffron,	
———— Mastic,	
———— Pimenta,	

Mix.

SWIETENIA.

Two species of this genus are mentioned in pharmacopœias :

1. *Swietenia Mahagoni*, LINN.—Mahogany.

SYNONIMES :—Acajou à meubles, Mahogon d'Amerique, FR.

DAN. ; BRUGNATELLI ; GUIBOUT ; MURRAY and GMELIN.

A large tree of South America. (*Decandria Monogynia*, LINN. ; *Meliaceæ*, JUSS. ; *Fig. Cat. Carol. II. tab. 20*)

The bark of the trunk and of the branches (*Cortex Ligni Mahagoni*) is employed. It is in nearly flat pieces, of about a foot in length, rough, and of a brown-grey colour : that of the branches is quilled, of the length of the hand or less, of a pale whitish-grey without, and of a deep brown-red within ; it is inodorous, but of a very astringent and bitterish taste.

2. *Swietenia febrifuga*, ROXB.

SYNONIMES :—Mahogon fébrifuge, FR. ; Rohun, BENG. ; Swamy, CAN. ; Patranga, SANS. ; Schemmarum, TAM. ; Sormido, TEL.

DUBL. ; DAN. ; COXE ; GUIBOUT.

A tree of the Coromandel Coast. (*Fig. Roxb. Pl. Corom. I. tab. 17*)

The bark (*Cortex Soymidæ*) is employed. It is in pieces of different sizes, generally one or two feet long, from one to eight inches broad, and of a moderate thickness. It is covered with a thin, rough, ash-grey, and a little spotted, epidermis, and is itself of an iron-red colour, brittle, compact, and susceptible of a polish. It has an agreeable and feebly aromatic odour, and a very bitter, astringent, and rather balsamic taste.

It is febrifuge.—Dose, of the powder, one scruple.

SYMPHYTUM.—COMFREY.

Two species of this genus of plants are used in medicine :

1. *Symphytum Officinale*, LINN.—Great Comfrey.

SYNONIMES ,—Grande Consonde, FR. ; Schwarzwurz, Beinwell, Wallwurz beinwell, Schmeerwurz, GERM. ; Kostiwal, Swalnj, Wysaky trunk, BOHEM. ; Kul sukkerod, DAN. ; Consuelda major, SPAN. ; Smeerwortel, DUT. ; Consolida maggiore, ITAL. ; Zywokost, POL. ; Consolda mayor, PORT. ; Vallært, SWED.

AMST. ; ANT. ; AUSTR. ; BATAV. ; BELG. ; BRUNS. ; HISP. ; GALL. ; FERR. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; LIPP. ; PALAT. ; ROSS. ; SAX. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; GUIBOUT ; MURRAY and GMELIN ; PIDE-RIT ; SPIELMANN ; ZARDA.

A common plant of Europe. (*Pentan-*
VOL. II.

dria Monogynia, LINN. ; *Borragineæ*, JUSS. ; *Fig. Flore Medic. III. 130*)

The root (*Radix consolida majoris seu Symphyti*) is employed. It is thick, not very branching, fibrous, long, of a blackish-brown without, and white within. It is inodorous, and has a viscous taste. It contains much mucilage, with gallic acid.

2. *Symphytum tuberosum*, LINN.

HISP.

A plant of the south of Europe. (*Fig. Jacq. Austr. t. 225*)

The root is employed ; it is bulged, and tuberculous at distances.

These two plants are slightly astringent. They are used, especially the first, in hæmoptysis and diarrhœa.

CONSERVE OF GREAT COMFREY.

Conserva Symphyti. (SARD.)

R. Root of Great Comfrey, any quantity.

Having sliced it, boil it in water till it may be bruised with the fingers; then pound it in a marble mortar, and strain the pulp through a silk sieve. Add to this two parts of sugar, and boil over a slow fire, constantly stirring.

ANTI-HÆMORRHAGIC POWDER. (NIE-MANN)

R. Powdered Brick, one ounce.
 Gum Traga- } of each,
 canth, } half an
 Arabic, } ounce.
 Root of Great Comfrey, six drams.
 Leaves of Betony, two drams and a half.

Mix.—It is recommended by PURMANN in hæmorrhages which cannot be restrained by ligature.

MUCILAGE OF GREAT COMFREY.

Mucilago Consolidæ majoris. (SARD.)

R. Root of Great Comfrey, one pound.
 Boiling Water, four pounds.

Macerate till the water is rendered very thick, and strain with slight expression.

DECOCTION OF GREAT COMFREY.

Decoctum Radicis Consolidæ. (FORM. HÔP. FR.; NIEMANN; RATIER)

R. Root of Great Comfrey, half an ounce.
 Spring Water, sixteen ounces.

Boil down to one half. (NIEMANN)

RATIER prescribes one ounce of root, and two pounds of water.

R. Root of Great Comfrey, half an ounce.
 Water, a sufficient quantity
 to obtain two pounds of strained liquor.

Boil for a quarter of an hour, adding towards the end,

Grated Liquorice, two drams.

Infuse for a few minutes, and strain. (FORM. HÔP. FR.)

It is astringent, and is recommended in internal hæmorrhages.—Dose, two spoonfuls every hour, of the first decoction, and a cupful of the two others.

SYRUP OF GREAT COMFREY.

Syrupus Symphyti seu e Symphyto. (HISP.; GALL.; SARD.; VAN MONS)

R. Root of Great Comfrey, one part.
 Warm Water, twenty parts.

Infuse for five or six hours, and strain

with expression. Dissolve in the strained liquor,

Sugar, twelve parts.

Evaporate to the consistence of syrup. (VAN MONS)

R. Sliced Root of Great Comfrey, six ounces.

Water, four pounds.

Boil a little. Add to the strained decoction,

Sugar, six pounds.

Boil to a syrup. (GALL.)

R. Sliced Root of Great Comfrey, two ounces.

Water, six pounds.

Boil till reduced to four pounds, and add towards the end,

Grated Liquorice, half an ounce.

Dissolve in the liquor,

White Sugar, two pounds.

Clarify, and make a syrup. (HISP.)

R. Depurated Juice of Comfrey, two pounds and a half.

Root of Great Comfrey, sliced, three ounces.

Macerate for an hour, and boil. Add to the strained decoction,

White Sugar, two pounds.

Clarify, boil to a syrup, and add

Syrup of Kermes, one pound.

Mix. (SARD.)

ASTRINGENT BROTH. (CADET DE GASSI-COURT; PIERQUIN; SAINTE-MARIE)

R. Root of Great Comfrey, } of each,

 Tormentil, } half an

 Bistort, } ounce.

 Chicken Broth, two pounds.

Boil down to one half.—To be taken cold in the course of the day, in uterine hæmorrhages. It may be sweetened with syrup of quince or of comfrey.

ASTRINGENT DRINK. (CADET DE GASSI-COURT; SAINTE-MARIE)

R. Root of Great Comfrey, } of each,

 Catechu, } two drams.

 Water, one pound.

Boil down to ten ounces. Add to the strained decoction,

Syrup of Quince, two ounces.

Hordeated Cinnamon Water, one ounce.

Mix.—To be taken by spoonfuls, every hour, in atonic diarrhoea.

ASTRINGENT PTISAN. (SAINTE-MARIE)

R. Root of Great Comfrey, one ounce.

Red Roses, one pinch.

Hips, twelve.
Catechu, one scruple.
Water, one pound and a half.

Boil for a quarter of an hour and strain.
—To be taken cold, by glassfuls, sweetened with syrup of quince or of gooseberries.

EMOLLIENT GARGLE. (FULD.)

R. Root of Great Comfrey, four ounces.
Dried Figs, one ounce.
Water, three pounds.

Boil down to two thirds.

BOYLE'S SYRUP.

Syrupus Boyleanus. (SPIELMANN)

R. Root of Great Comfrey, six ounces.
Herb of Great Plantain, three ounces.

Bruise together, and express the juice.
After having depurated it, add

White Sugar, an equal weight.

Boil to a syrup.

It is principally recommended in hæmoptysis.

SYRUP AGAINST HOOPING COUGH.

Syrupus ad Pertussim. (SAINTE-MARIE)

R. Fresh Leaves of Lungwort, four handfuls.
Root of Great Comfrey, } of each,
Corinth Raisins, } two ounces.
Liquorice, bruised, } three ounces.
Water, } two pounds.

Boil down to one pound and a half.
Add to the strained decoction,

White Sugar, twenty-four ounces.

Boil for a few moments.

COMPOUND SYRUP OF COMFREY.

Syrupus e Symphyto Fernelii. (BRUNS.; PALAT.; WIRTEM.)

R. Red Roses, } of each,
Fresh Roots and Leaves } three ounces.
of Great Comfrey, }
Betony, }
Plantain, } of each,
Burnet Saxifrage, } one ounce.
Scabious, }
Knot Grass, }
Coltsfoot, }

Express the juice of these fresh plants, and after depuration add to every nine ounces of it,

White Sugar, sixteen ounces.

Make a syrup. (BRUNS.; WIRTEM.)

PALAT. directs the juice to be boiled, and two pounds and a half of sugar to be dissolved in three pounds of it.

ANTI-HERNIAL PLASTER.

Emplastrum ad Hernias seu Picis nigre resigum. (HISP.; PALAT.; WIRTEM.; SPIELMANN)

R. Venice Turpentine, four ounces.
Mastic, one ounce and a half.

Melt together, and add

Yellow Wax, six ounces.
Powdered Myrrh, } of each,
Olibanum, } six
Dragon's Blood, } drams.
Root of Great
Comfrey, } of each,
Seeds of Thorough-wax, } one ounce.
Blood stone, } of each,
Calcined } half an
Vitriol, } ounce.

Mix. (WIRTEM.)

R. Yellow Wax, four ounces.
Turpentine, six ounces.

Melt together, and add,

Powdered Acacia Juice, }
Hypocist, } of each,
Mastic, } one ounce.
Myrrh, }
Olibanum, }
Root of Great
Comfrey, }
Alum, } of each,
Dragon's Blood, } six drams.
Cypress Nuts, }
Blood stone, } of each,
Seeds of Thorough-wax, } half
Sulphate of } an ounce.
Copper, }

Mix. (SPIELMANN)

R. Black Resin, one pound and a half.
Yellow Wax, } of each,
Turpentine, } eight ounces.
Labdanum, } two ounces.

Melt together, and add,

Juice of Hypocist, dissolved in decoction of Cypress nuts and inspissated, one ounce.

Having removed it from the fire, add further,

Powdered Mastic, } of each,
Root of Great } two drams
Comfrey, } and a half.
Cypress Nuts, } one ounce.
Armenian } of each,
Bole, } half an
Blood stone, } ounce.

Mix. (HISP.)

R. Yellow Wax, two ounces.
Turpentine, } of each,
Black Peruvian } half an ounce.
Balsam, }

Resin, one ounce.
Melt together, and add
Powdered Mastic, }
Dragon's } of each,
Blood, } half an ounce.
Olibanum, } of each,
Myrrh, } two drams.
Oil of Eggs, a sufficient quantity.

Mix. (PALAT.)

It was formerly recommended to be applied to the herniæ of children. The use of styptics in such cases is now properly estimated.

PLASTER AGAINST FRACTURES.

Emplastrum catagmaticum seu ad fracturas seu ad fracturas et luxationes. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

R. Yellow Wax, half a pound.

Resin, one pound.
Turpentine, two ounces.
Melt, and add,
Powdered Root of Great
Comfrey, }
Queen } of each,
of the Meadows, } two
Armenian Bole, } ounces.

Mix. (WIRTEM.)

PALAT. prescribes three pounds of resin, one pound of wax, half a pound of turpentine, five ounces of root of comfrey, three ounces of root of ulmaria, six ounces of bole, and one ounce of hæmatitis.—BRUNS.—thirty-two ounces of resin, eight of turpentine, four of comfrey, and four of ulmaria.—SPIELMANN—twenty-four ounces of resin, six of turpentine, and three each, of bistort, of comfrey, and of tormentil.

It is astringent and corroborant.

TACAMAHACA.—TACAMAHAC.

SYNONIMES:—Tacamahaca, FR.; Hack und tack, Takamahak harz, GERM.; Takamaka, BOHEM.; Hak og tak, DAN.; Tacamahaca, SPAN.

AMST.; ANT.; BATAV.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; OLDEN.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A resin in irregular masses, appearing to be formed of agglutinated pieces, of a yellowish or greenish colour, semi-transparent internally, undulated with whitish zones, farinuous and of a greyish colour externally, slightly friable, of an aromatic and terebinthine odour, and of an indistinct taste becoming acrid after some time.

Besides this kind, called *Tacamahaca communis* seu *in sortis*, there is also another, *Tacamahaca sublimis* seu *in testa* which is in tears or small grains, of a whitish-grey externally and reddish or yellowish internally.

This resin flows from incisions made in the bark of the *Fagara octandra*, LINN. (*Ocandria Monogynia*, LINN.; *Terebinthaceæ*, JUSS.), a tree of South America, and also, it is said, of Madagascar (*Fig. Jacq. Amer. t. 71. f. 1.*—3). It has been erroneously attributed to the *Populus balsamifera*, LINN.; a tree (*Diæcia Octandria*, LINN.; *Amentaceæ*, JUSS.) of the north of

the two continents (*Fig. Pall. Fl. Ross. I. t. 41*), which yields an odorous resin called *North American Tacamahac* (*Baume Tocot, Tacamahaca de l'Amerique septentrionale*, FR.); but some authors think this latter is produced by the *Calophyllum inophyllum*, LINN.; a tree (*Polyandria Monogynia*, LINN.; *Guttiferæ*, JUSS.) of the East Indies. (*Fig. Rheed. Hort. Malab. 4. t. 38*)

It is stimulant and resolvent.

OIL OF TACAMAHAC.

Oleum Tacamahacæ. (WIRTEM.)

R. Tacamahac, one part.
White Sand, washed, one part and a half.

Distil in a glass retort placed in a sand-bath, separate the oil from the acid liquor which passes over with it, and rectify it.

STOMACHIC PLASTER.

Emplastrum stomachicum seu de Tacamahaca seu tacamahacinum seu de Crusta Panis. (BRUNS.; FULD.; PALAT.; SARD.; SAX.; WIRTEM.; AUGUSTIN; SPIELMANN)

R. Tacamahac, three ounces.
Caranna, six drams.
Balsam of Peru, } of each,
Turpentine, } one ounce and a half.

Melt together over a slow fire. (SARD.)

R. Yellow Wax, eight ounces.
Turpentine, one ounce.

Melt, and add to the mixture when half cold,

Powdered Tacamahac,	four ounces.
—— Balsam of Peru,	of each,
Oil of Mint,	two drams.
—— Cloves,	one ounce.

Mix. (SAX.; WIRTEM.; SPIELMANN)

FULD. prescribes one pound and a half of wax, half a pound of turpentine, three ounces each, of tacamahac and of nutmegs, half an ounce of balsam of Peru, and two drams of oil of mint.—VAN MONS—sixty-four parts of wax, eight of turpentine, thirty-two of tacamahac, one of oil of cloves, and two each, of oil of nutmeg and of balsam of Peru.

R. Galbanum,	} of each, an equal part.
Tacamahac,	
Yellow Wax,	
Assafœtida,	
Cumin,	
Turpentine,	

Make a plaster. (AUGUSTIN)

R. Tacamahac, one ounce.

Mastic,	} of each, half an ounce.
Storax,	
Mace,	} of each, one dram.
Nutmeg,	
Cloves,	
Balsam of Peru,	two drams.
Oil of Caraway,	of each,
—— Mint,	twenty drops.
Yellow Wax,	four ounces.
Turpentine,	two ounces.

Make a plaster. (BRUNS.)

R. Crust of Bread, grated, two ounces.
 Tacamahac, one ounce.
 Mastic, }
 Cloves, } of each,
 Nutmegs, } half an ounce.
 Oil of Mint, }
 Yellow Wax, six ounces.
 Turpentine, }
 Oil of Mastic, } of each, one ounce.
 —— Quince, }

Make a plaster. (PALAT.; WIRTEM.; SPIELMANN)

This plaster is applied to the epigastrium to arrest vomiting and diarrhœa, and also as tonic and carminative.

TAMARINDUS INDICA, LINN.—TAMARIND TREE.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FULD.; HAMB.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOUT.; MURRAY and GME-LIN.; PARIS.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

A fine tree of the East Indies. (*Triandra Monogynia*, LINN.; *Leguminosæ*, JUSS.; *Fig. Flore Medic. VI. 338*)

The fruit, *Tamarinds*; *Tamarindi*, *Fructus Tamarindorum*, are employed.

SYNONIMES:—Tamarins, FR.; Tamarinden, Sauerdatteln, GERM.; Umblic, ARAB.; DUK.; and HIND.; Tamaryndy, Indyanske daktyle, BOHEM.; Tamalaki, BAND.; Cay-me, COCHIN.; Mahasi ambala, CYNG.; Tamarin, DAN.; and SUEC.; Tamarindo, ITAL.; SPAN.; and PORT.; Tamarinde, DUT.; Kamal, JAV.; Heghka, MAL.; Tumiri hindi, PERS.; Amlika, Tintili, SANS.; Pullie, TAM.; Chinta pundo, TEL.

It consists of rather compressed pods, from three to five inches in length, indehiscent, of a brown-grey colour, filled with a

thick pulp, containing several rather large, shining, angular, compressed seeds.

The pulp is of a glutinous consistence, of a dark-brown colour, of a vinous odour, and of an agreeable and acidulous taste, becoming nauseous in time.

It contains, according to VAUQUELIN, starch, sugar, jelly (pectic acid), gum, citric acid, super-proto-tartrate of potassium, tartaric acid, and malic acid.

PULP OF TAMARINDS.

Pulpa Tamarindorum. (AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FULD.; HANN.; HASS.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS)

R. Tamarinds, any quantity.

Triturate them with water to form a homogeneous mass; rub this on the under side of a hair-sieve with a wooden spatula: remove the pulp which passes through with a silver spatula, and preserve it in a pot covered with paper. (ANT.; BATAV.; BELG.; VAN MONS)

AMST. and GALL. direct the tamarinds to be triturated with boiling water.—LU-

sit. directs them to be softened by previous ebullition in water.

R. Tamarinds, any quantity.

Triturate with a little boiling water; then strain the pulp through a silk sieve, and evaporate it to the consistence of honey; then add one-sixth of sugar, and evaporate again to the required consistence. (HANN.; OLDEN.; POL.; BORUS.; SAX.; SWEDIAUR)

BRUNS.; HASS.; PALAT.; WIRTEM.; and PIDERIT prescribe two parts of pulp and one part of sugar.—DAN.; ROSS.; and HERBIP.—three of pulp and one of sugar.—SPIELMANN—four of pulp and one of sugar.—FENN.—twenty-four of pulp and one of sugar.

R. Tamarinds, bruised, any quantity.
Water, a sufficient quantity.

Boil for some time, and strain through flannel; add half an ounce of iron filings to each pound of the strained liquor, and set aside for twenty-four hours; then pour off the clear part, plunge a polished iron plate in it, and if no particle of copper is deposited upon the plate, evaporate the liquor to the consistence of honey. Add three ounces of sugar to each pound of the decoction, and evaporate to the required consistence. (FULD.; LIPP.)

It is laxative, refreshing, anti-septic, and anti-alkaline.—Dose, from two drams to two ounces.

CONSERVE OF TAMARINDS.

R. Pulp of Tamarinds, one part.
White Sugar, two parts.
Mix intimately.

INFUSION OF TAMARINDS.

Infusum Tamarindorum. (CADET DE GASSICOURT)

R. Tamarinds, one or two ounces.
Boiling Water, two pounds.

Boil for ten minutes, and strain without expression.

LAXATIVE DRINK. (CADET DE GASSICOURT)

R. Tamarinds, two ounces.
Boiling Water, two pounds.

Mix, and add to the strained liquor

Honey, one ounce.

Mix.

COMPOUND INFUSION OF TAMARINDS.

Infusum Tamarindorum compositum.
(BRUNS.; LUSIT.)

R. Pulp of Tamarinds, } of each, one
Manna, } ounce and a half.
Orange Peel, two drams.
Boiling Water, eight ounces.

Digest till the manna and pulp become dissolved, and strain. (LUSIT.)

R. Tamarinds, freed from their stones, half a pound.
Oleosaccharum of Lemon, one ounce.
Lemon Juice, half an ounce.
Supertartrate of Potass, two drams.
Boiling Water, eight pounds.

Infuse, and strain. (BRUNS.)

It is laxative.

DECOCTION OF TAMARINDS.

Decoctum Tamarindorum. (AMER.; GALL.; FULD.; NIEMANN; VAN MONS)

R. Tamarinds, two ounces.
Water, three pounds.

Boil for a quarter of an hour, and strain. (RATIER)

R. Tamarinds, one or two ounces.
Water, two pounds.

Boil for eight minutes, and strain without expression. Add

Syrup of Maidenhair, one ounce.

Mix. (GALL.)

NIEMANN prescribes two ounces of tamarinds, four pounds of water reduced to three, and one ounce and a half of syrup of raspberries.

R. Tamarinds, one ounce.
Corinth Raisins, two ounces.
Water, a sufficient quantity

to obtain sixteen ounces of strained liquor.

Boil for a few moments, and strain without expression.

ANT. prescribes two ounces of tamarinds, four of raisins, and two pounds and a half of water, to obtain two pounds of strained liquor.

R. Pulp of Tamarinds, two ounces.
Water, four pounds.

Boil to the consumption of one fourth. Add

Nitrate of Potass, three drams.
Rob of Gooseberries, one ounce and a half.

Mix. (FULD.)

To be taken by glassfuls.

COMPOUND DECOCTION OF TAMARINDS.

Decoctum Tamarindi compositum. (HERBIP.; VAN MONS)

R. Tamarinds, } of each,
Corinth Raisins, } three ounces.
Boiling Water, a sufficient quantity
to obtain three pounds of strained decoction.
Add

Tartarized Antimony, half a grain.
Syrup of Mulberries, one ounce and a half.

Mix. (VAN MONS)

R. Tamarinds, six ounces.
Supertartrate of Potass, three drams.
Spring Water, a sufficient quantity
to obtain one pound of strained decoction.
Add while hot,

Senna, } of each,
Rhubarb, } one dram and a half.

Digest for four hours; then dissolve in the liquor

Manna, ten drams.

Strain, and add

Sulphuric Æther, six drops.

Mix. (HERBIP.)

Dose, from two to four ounces every hour.

TAMARIND WHEY.

Serum Lactis tamarindatum seu temperans.
(HANN.; AUGUSTIN; BRERA; SWEDIAUR)

R. Cow's Milk, two pounds.
Tamarinds, two ounces.

Boil, and strain after coagulation.
(HANN.; SWEDIAUR)

LIPP.; and AUGUSTIN direct four parts of whey to be boiled with one part of tamarinds, and strained.—BRERA—one ounce and a half of pulp, and a sufficient quantity of milk to obtain two pounds of strained whey.

It forms an agreeable, acidulous, refreshing, and slightly laxative drink.

ANTI-SCORBUTIC WHEY. (PALAT.; BORRIES)

R. Sorrel, } of each, one handful.
Betony, }
Clove-pinks, half a handful.
Tamarinds, one ounce and a half.
Clarified Whey, three pounds.

Infuse for an hour at a moderate heat.
Add to the strained liquor

Syrup of Lemon, } of each, one
Violets, } ounce and a half.

Mix. (BORRIES)

R. Tamarinds, half an ounce.
Sorrel, one handful and a half.
Betony, } of each, half a handful.
Chervil, }
Whey, three pounds.

Infuse for an hour, and express. Add

Syrup of Lemon Juice, } of each,
Raspberries, } one ounce.
Violets, }

Mix. (PALAT.)

Dose, one ounce every half hour.

LAXATIVE POTION.

Potio seu Mixtura laxans. (FULD.; HAMB.; BRERA; PIERQUIN; SAINTE-MARIE)

R. Tamarinds, three ounces.
Water, twelve ounces.

Boil down to ten ounces; then add to the strained liquor,

Manna, two ounces.

Strain again. (HAMB.)

R. Pulp of Tamarinds, one ounce.
Water, a sufficient quantity

to obtain eight ounces of decoction. Add
Supertartrate of Potass, one ounce.

Dissolve.—To be taken at twice. (BRERA)

R. Pulp of Tamarinds, one ounce and a half.

Sulphate of Soda, one ounce.

Boiling Water, nine ounces.

Syrup of Manna, one ounce.

Dissolve, and mix. (FULD.)

R. Pulp of Tamarinds, } of each,
Manna, } two ounces,
Nitrate of Potass, } two drams.
Water, two pounds.

Dissolve, and strain. (FULD.)

R. Tamarinds, one ounce.
Yellow Sanders, one dram.
Manna, two ounces and a half.
Water, nine ounces.
Infusion of Ipecacuanha, one ounce.

Mix. (PIERQUIN)

R. Supertartrate of Potass, six drams.
Nitrate of Potass, two drams.
Tamarinds, four ounces.
Water, one pound and a half.

Boil for half an hour. Add to the strained decoction,

Syrup of Violets, two ounces.

Mix. (SAINTE-MARIE)

R. Tartrate of Potass, two drams.
Senna, three drams.
Tamarinds, one ounce.
Liquorice, two drams.
Lemon, sliced, one half.
Chervil, one handful.
Montpellier Scammony, seven grains.
Boiling Water, one pound and a half.

Infuse for twelve hours, and express.
(PIERQUIN)
To be taken by glassfuls.

LAXATIVE ELECTUARY.

Electuarium laxativum seu Tamarindorum.
(PALAT.; WIRTEM.; PH. SYPH.; SWEDIAUR.; VAN MONS)

R. Pulp of Tamarinds, four ounces.
—— Raisins, } of each, one
—— Prunes, } ounce and a half.
Senna,
Supertartrate of Potass, six drams.
Cinnamon, } of each, two drams.
Fennel, }
Oil of Nutmeg, ten drops.
Syrup of Buckthorn,
a sufficient quantity.

Make an electuary. (WIRTEM.)

R. Senna, six ounces.
Fennel, two drams.
White Wine, } of each, two pounds.
Water, }

Boil down to one half. Add to the expressed liquor,

Pulp of Prunes, } of each,
—— Raisins, } half a pound.
—— Tamarinds, }
Supertartrate of Potass, one ounce.
Senna, three ounces.
Oil of Anise, } of each, half a dram.
—— Nutmeg, }

Mix. (PALAT.)

R. Supertartrate of Potass, one dram.
Senna, three drams.
Manna, three ounces and a half.
Pulp of Cassia, } of each, one
—— Tamarinds, } ounce and a half.
Syrup of Violets, a sufficient quantity.

Mix. (VAN MONS)

R. Pulp of Tamarinds, three ounces.
Leaves of Senna, one dram and a half.
Nitrate of Potass, } of each,
Sulphur, } three drams.
Syrup of Orange Peel,
a sufficient quantity.

Mix. (SWEDIAUR)

PH. SYPH. gives the same formula, but omits the senna.

Dose, from half an ounce to one ounce.

TAMARIX.—TAMARISK.

Two species of this genus of plants are used in medicine.

1. *Tamarix Germanica*, LINN.—German Tamarisk.

SYNONIMES:—*Tamaris d'Allemagne*, FR.; *Deutscher Tamarisken*, GERM.

AMST.; FERR.; WIRTEM.; HERBIP.; SPIELMANN.

A tree of Germany. (*Pentandria Trigynia*, LINN.; *Portulacæ*, JUSS.; *Fig. Pall. Fl. Ross.* II. t. 80)

The bark and leaves are employed.

The bark (*Cortex Tamarisci* seu *Tamariscis*) of the trunk and particularly of the root, is grey or brown-grey without and yellow or reddish within, unequal, and cracked. It has a pleasant and balsamic odour, and a very astringent and bitterish taste.

The leaves are sessile, linear, lanceolated, and imbricated.

2. *Tamarix Gallica*, LINN.—French Tamarisk.

SYNONIMES:—*Tamaris de France*, FR.; *Französischer Tamarisk*, GERM.; *Fransche Tamarish*, DUT.; *Tamarisco*, ITAL.; *Tamarqueira*, PORT.

ANT.; BATAV.; BRUNS.; HISP.; GALL.; FERR.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY; and GMELIN.

A shrub of the south of Europe. (*Fig. Zorn, Ic. pl.* t. 312)

The bark and leaves of this species are also employed.

The bark (*Cortex Tamarisci* seu *Tamarisci Gallici*) is thin, quilled, brown externally and pale yellow internally. Its taste is bitter and somewhat astringent. The leaves are lanceolated, very small, pointed, amplexicaul, and imbricated.

It is tonic, but is not much used. It was formerly classed among the alteratives.

TAMNUS COMMUNIS, LINN.—COMMON TAMNUS, BLACK BRYONY.

SYNONIMES :—Sceau de notre dame, Vigne noire, Bryone noire, Fel ; Jungferwurz, Schmeerwurz, Schmeerswurz schwarzebe, GERM.

GALL. ; GUIBOURT.

A plant of the south of Europe. (*Diaccia*

Hexandria, LINN. ; *Alismaceæ*, JUSS. ; *Fig.* Blackw. *Herb.* t. 457)

The root (*Radix Tamni* seu *Bryonia nigra*) is employed. It is round, tuberculous, black without and white within. It has an acrid and rather disagreeable taste.

It is considered diuretic.

TANACETUM VULGARE, LINN.—TANSEY.

SYNONIMES :—Tanaïsie, FR. ; Rheinfarn, GERM. ; Wratje, BOHEM. ; Rheinfan, DAN. ; Tanaceto, ITAL. ; and SPAN. ; Seinevaren, DUT. ; Wrotyez, POL. ; Tanasia, PORT. ; Renfana, SWED.

EDINB. ; DUBL. ; AMER. ; AMST. ; ANT. ; AUSTR. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FERR. ; FENN. ; FORM. HÔP. FR. ; FULD. ; GENÈV. ; HAMB. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; WURCEB. ; BERGIUS ; BRUGNATELLI ; COXE ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A plant of Europe. (*Syngenesia Polygamia Superflua*, LINN. ; *Synanthèreæ*, CASS. ; *Fig. Flore Medic.* VI. 339)

The herb, flowers, and seeds are employed.

The herb (*Herba Tanacetii* seu *Athanasia*) consists of a striated, cylindrical, branching stem, and of ample, alternate, petiolate, even, glabrous leaves, the inferior of which are twice winged ; the leaflets present three notches, are nearly sessile, narrow, linear, acute, and slightly decurrent. The taste is aromatic, warm, and bitter.

The flowers are radiated calathides, of a brilliant yellow colour, and of a balsamic and bitter taste.

The seeds are small, long, slightly curved, deeply grooved, and furnished with a small border at the superior extremity, which is broader than the inferior. Their taste is bitter and camphorous.

The odour is strong and balsamic. That of the flowers is disagreeable, camphorous, and aromatic.

It is excitant, diaphoretic, and vermifuge. —Dose, of the powder, from twenty to forty grains.

VOL. II.

CONSERVE OF TANSEY.

Conserva Tanacetii, Confectio Florum Tanacetii. (FULD. ; SARD. ; VAN MONS)

R. Leaves of Tansey, bruised, one part.
White Sugar, two parts.

Triturate together, and heat in a water-bath. (VAN MONS)

SARD. prescribes an equal part of each.

R. Flowers of Tansey, without calices, one part.

Put them into a copper pan placed over a gentle fire, and pour over them gradually,

Syrup, boiled à la plume, two parts.

Stir till the mass is perfectly dry, and repeat this till the flowers become entirely covered with sugar. (FULD.)

COMPOUND POWDER OF TANSEY.

Pulvis Tanacetii compositus. (SUEC.)

R. Tops of Tansey, }
Semen Contra, } of each, one dram,
Gamboge, }
Nux Vomica, }
Sulphate of Iron, }

Pulverize separately, and mix.

NERVINE SPECIES.

Species nervinæ pro cucuphis. (HASS. ; PIDERIT)

R. Herb of Peppermint, }
— Rue, }
— Wild Thyme, } of each,
— Marjoram, } two ounces.
— Thyme, }
— Sage, }
— Savine, }
— Balm, }
Juniper, }
Flowers of Red Roses, } of each,
— Lavender, } three ounces.
— Tansey, } four ounces.

Bay Berries,	two ounces.
Lesser Cardamoms,	one ounce.
Anise,	of each,
Caraway,	one ounce and a half.
Mace,	half an ounce.

Cut, and bruise.

AQUEOUS EXTRACT OF TANSEY.

Extractum Tanacetii aquosum. (BRUNS.; FULD.; LIPP.; SARD.; SAX.; WIRTEM.; VAN MONS)

R. Herb of Tansey,	one pound.
Spring Water,	six pounds.

Digest for four days in a warm place, and boil a little. Evaporate the expressed liquor to the consistence of an extract. (BRUNS.; FULD.; WIRTEM.)

LIPP. directs one pound of herb, to be boiled with eight pounds of water, and expressed; a little cold water to be poured upon the residue, and expressed again: the liquors to be mixed and then evaporated.

R. Herb of Tansey,	one part.
Water,	eight parts.

Macerate for twenty-four hours, then boil for a quarter of an hour, and express; boil the residue with

Water,	four parts.
--------	-------------

Mix the decoctions, set apart for twenty-four hours, and evaporate the decanted liquor to the proper consistence. (SAX.)

R. Herb of Tansey, bruised,	eight pounds.
-----------------------------	---------------

Macerate for three days in an alembic; then having distilled off about two ounces of liquor, pour upon the herb,

Water,	twenty-four pounds,
--------	---------------------

and boil till reduced to one third. Clarify and strain the liquor, and evaporate to the consistence of an electuary, adding the distilled water to this when nearly cold. (SARD.)

FERR. directs one pound of herb to be boiled in three pounds of water; the strained decoction to be clarified with white of egg, and evaporated.

R. Leaves of Tansey,	any quantity.
----------------------	---------------

Bruise them, and triturate them with a little water for a considerable time; then express, coagulate, and set apart the fecula: then infuse the residue at a moderate heat in the limpid juice, express again, clarify, and evaporate to the consistence of an extract; finally, having added the fecula, continue the evaporation till the mass has acquired a proper consistence. (VAN MONS)

Dose, from six to twenty grains.

ALCOHOLIC EXTRACT OF TANSEY.

Extractum Tanacetii alcoholicum. (HERBIP.)

R. Tops of Tansey,	} of each, one pound.
Alcohol,	
Water,	

Digest for three days at a gentle heat, and express; then, having set aside and decanted the liquor, distil off the alcohol, and evaporate to the proper consistence.

VERMIFUGE LINIMENT. (VAN MONS)

R. Fresh Green Fecula of Tansey,	one part.
Lard,	one part and a half.

Melt gradually, wash the mixture with hot water, and decant; then by successive meltings and decantations, free it as much as possible from water.

INFUSION OF TANSEY.

Infusum Tanacetii. (NIEMANN)

R. Herb of Tansey,	half an ounce.
Boiling Water,	ten ounces.

Infuse, and strain.

It is diaphoretic.—Dose, a cupful every two hours.—The infusion made with milk, and administered as a lavement, is useful to expel ascarides.

VERMIFUGE LAVEMENT.

Enema ad Ascarides. (SWEDIAUR)

R. Herb of Tansey,	half an ounce.
Water,	eight ounces.

Boil down to six ounces, and strain.

TINCTURE OF TANSEY.

Tinctura Tanacetii. (FULD.)

R. Tops of Tansey,	one part.
Alcohol,	four parts.

Digest for three days, at a moderate heat, and express.

Dose, from thirty to sixty drops.

VERMIFUGE ESSENCE.

Essentia contra Vermes. (BRUNS.; VAN MONS)

R. Herb of Tansey,	} of each,
Wormwood,	
Lesser Centaury,	
Root of Gentian,	} an ounce.
Flowers of Tansey,	
Semen Contra,	} of each,
Myrrh,	
Spirit of Mint,	one pound.

Digest, express, and filter. (BRUNS.)

- R. Fresh Leaves of Tansey, one part.
 — Root of Elecampane, } of each,
 Rhubarb, } three
 Santolina, } parts.
 Madeira Wine, two parts.
 Proof Spirit, twenty parts.

Infuse without heat for eight days; then express and filter. (VAN MONS)

ESSENTIAL OIL OF TANSEY.

Oleum Tanacetii æthereum, Ætheroleum Tanacetii. (AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; ROSS.; SAX.; SUEC.; HERBIP.; BRUGNATELLI; PIDERIT; SWEDIAUR)

- R. Flowering Herb of Tansey, any quantity.
 Water, a sufficient quantity.
 Distil. (AMST.; BATAV.; BELG.; FENN.; FERR.; ROSS.; SUEC.; BRUGNATELLI)

FULD.; and LIPP. prescribe one part of herb and three parts of water.—SWEDIAUR—
 one of herb and four of water.—DAN.—
 one of herb and six of water.—HANN.;

OLDEN.; and SAX.—one of herb and eight of water.—AUSTR.—one of herb and twelve of water.—GALL.—five of herb and seven of water.—ANT.—three of herb and ten of water.

- R. Herb of Tansey, twenty-five parts.
 Water, seventy-five parts.
 Muriate of Soda, three parts.

Macerate for three days, and distil. (BRUNS.; HASS.; PIDERIT)

- R. Herb of Tansey, any quantity.
 Distilled Water of Tansey, a sufficient quantity.
 Distil. (HERBIP.)

DISTILLED WATER OF TANSEY.

Aqua Tanacetii. (AUSTR.; HERBIP.; PIDERIT)

- R. Flowering Herb of Tansey, one part.
 Water, six parts.
 Distil, and separate the oil from the water. (HERBIP.)

AUSTR. prescribes one part of herb and twelve parts of water; four parts to be distilled.—PIDERIT—one of herb and eight of water; four to be distilled.

TARAXACUM DENS LEONIS, DESF.—DANDELION.

SYNONIMES:—Pissenlit, FR.; Löwenzahn, Pfaffenröhrlein, Pappenkraut, GERM.; Plesska, BOHEM.; Lœvetand, DAN.; Diente de leon, SPAN.; Paapenkruid, DUT.; Dente di leone, ITAL.; Wolowe oczy, Swini mlecz, POL.; Dente de leao, PORT.; Lejontand, Maskrosor, SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; GUIBOUT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN; ZARDA.

A plant common to all Europe. (*Syngenesia Polygamia equalis*, LINN.; *Synantherææ*, CASS.; *Fig. Zorn, Ic. pl. t. 69*)

The root and herb are employed.

The root (*Radix Taraxaci seu Dentis Leonis*) is cylindrical, long, of the thickness of the finger, sometimes branched, furnished with fibrils. It is of a dark-brown colour externally and white and milky internally;

it is destitute of odour, and has a bitter, then sweet, and rather saline taste: it loses its bitterness by desiccation.

The herb consists of a rosette of long, runcinate, denticulated, smooth leaves of a light-green colour, inodorous, and of a bitter taste.

JUICE OF DANDELION.

Succus Taraxaci. (HASS.; SARD.; HERBIP.)

- R. Fresh Root and Herb of Dandelion, any quantity.

Bruise in a stone mortar; then express, set aside, and filter the juice. (HASS.)

HERBIP. directs one dram of alcohol to be added to each pound of juice.—SARD. requires it to be clarified.

Dose, from two to four ounces.

DECOCTION OF DANDELION.

Decoctum Taraxaci seu Leontodontis Traxaci. (ANT.; NIEMANN)

- R. Root of Dandelion, two ounces.
 Water, two pounds.

Boil down to about one half, and set aside and decant the liquor. (ANT.)

NIEMANN prescribes one ounce and a half of root, and three pounds of water; to be reduced to one third.

It is diuretic and deobstruent.—Dose, one pound daily.

COMPOUND DECOCTION OF DANDELION.

Decoctum Taraxaci compositum seu resolvens. (MIL. COPEN.; NIEMANN; SWEDIAUR)

R. Root of Dandelion, four ounces.
Herb of Buckbean, half an ounce.
Water, forty-eight ounces.

Boil down to one half, and express strongly. Add to the liquor when half cold

Muriate of Ammonia, two drams,
or, Sulphate of Magnesia, one ounce.
Sulphuric Æther, two drams.

Mix. (MIL. COPEN.)

SWEDIAUR prescribes one pound and a half each, of herb and root of dandelion and of dog-grass, six pounds of water, reduced to three, and the addition to the strained liquor of half a pound of oxymel, six drams of sulphate of potass, and twenty-five drops of spirit of sulphuric æther.—NIEMANN—two ounces and a half each, of dog-grass, of root of dandelion, and of root of cichory, and a sufficient quantity of water to obtain one pound of strained decoction, and the addition to this of from one to three ounces of compound syrup of smallage, two drams of sulphate of potass, and twenty-five drops of sulphuric æther.

It is recommended by STOLL, in obstinate quartan fevers, enlargements of the abdominal viscera, jaundice, and in cases of biliary calculi.—Dose, half a pound, four or six times a day.

DEOBSSTRUENT DECOCTION.

Decoctum resolvens. (FULD.; HAMB.; SAINTE-MARIE; VAN MONS)

R. Root of Dandelion, } of each,
—— Dog-grass, } two ounces.
Water, three pounds.

Boil down to two pounds; then add to the strained decoction,

Sulphate of Potass, half an ounce.
Oxymel, two ounces.

Mix. (SAINTE-MARIE; VAN MONS)

R. Root of Dandelion, four ounces.
Spring Water, two pounds.

Boil down to one half, and add to the strained liquor

Supertartrate of Potass, one ounce.

Sub-borate of Soda, three drams.
Glycyrrhized Honey, two ounces.

Dissolve. (HAMB.)

R. Root of Dandelion, two ounces.
Spring Water, one pound.

Boil down to one half, and add when strained,

Extract of Great Celandine, two drams.
Tartarized Antimony, two grains.

Dissolve. (HAMB.)

R. Root of Dandelion, four ounces.
Supertartrate of Potass, } of each,
Sub-borate of Soda, } half an ounce.
Water, three pounds.

Boil down to one half, and add

Despumated Honey, two ounces.

Mix. (FULD.)

EXTRACT OF DANDELION.

Extractum seu Mellago Taraxaci seu Leontodontis Taraxaci. (LOND.; DUBL.; AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; FERR.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; HERBIP.; BRUGNATELLI; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Extractum Taraxaci. (LOND.)

Ext. Herbx et Radicis Taraxaci. (DUBL.)

R. Fresh Root of Dandelion, one pound.
Boiling Water, one gallon.

Macerate for twenty-four hours, and then boil down to one half. Strain the decoction while hot, and evaporate to the proper consistence. (LOND.; COXE)

DUBL. directs the herb and root to be employed.

R. Fresh Root and Herb of Dandelion, one pound.
Water, ten pounds.

Boil for two hours, and express; set apart the decoction, strain it through flannel, and evaporate till the extract no longer adheres to the fingers. (AMST.)

ROSS. directs one part of herb and root to be boiled in two or three of water; the decoction to be filtered and evaporated.—HASS.; and PIDERIT—one of herb and root to be boiled in three of water; the decoction to be expressed and evaporated over a slow fire.—AUSTR.—equal parts of root and of herb to be boiled in a sufficient quantity of

water, and the decanted decoction evaporated to the consistence of liquid extract.—OLDEN.; POL.; BORUS.; and SAX.—twelve pounds of root to be bruised, gradually adding six pounds of water; the juice to be expressed, boiled for an instant, strained through flannel, and evaporated to the consistence of honey.—HISP.—the root to be boiled in a sufficient quantity of water, and the decanted decoction evaporated slowly, constantly stirring.—HERBIP.—the boiling to be continued for an hour, and then evaporated to the consistence of honey.—ANT.—two pounds of root and herb to be boiled for a quarter of an hour in ten of water, and the decanted decoction evaporated.—BAVAR.—one part of root and herb to be boiled in six of water for a quarter of an hour, and the strained liquor evaporated.—FERR.—one pound of root to be boiled in three pounds of water; the decoction to be clarified with white of egg and evaporated.

R. Fresh Herb and Root of Dandelion,
two pounds.
Boiling Water, ten pounds.

Digest for twenty-four hours, and boil for one hour; then decant when cold, and express; when the mixed liquors are depurated by repose, evaporate to the proper consistence. (BATAV.; BELG.)

R. Dried Root of Dandelion, any quantity.
Macerate in cold water, and digest at two different times in hot water, clarify the mixed infusions with white of egg, and evaporate to the consistence of extract. (VAN MONS)

R. Fresh Herb of Dandelion, any quantity.
Bruise it in a stone mortar, sprinkling it with a little water; then express the juice, and evaporate it in a water-bath, constantly stirring towards the end with a spatula. (BRUNS.; HISP.; GENEV.; LUSIT.)

DAN.; and SWEDIAUR direct two pounds of root and herb to be bruised, adding six pounds of water; the juice to be expressed, after having boiled it for a few moments, and evaporated to the consistence of thick syrup.

R. Root of Dandelion, } of each,
Herb of Dandelion, } eight pounds.

Bruise in a stone mortar, and express the juice; boil the residue with

Water, two parts,
and express. Mix the liquors, and after twelve hours repose, evaporate to the consistence of syrup. (LIPP.)

R. Herb and Root of Dandelion, one part.
Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and express; boil the residue with

Water, four parts,
and strain again. Mix the liquors, decant after sufficient repose, and evaporate slowly to the consistence of an extract. (HANN.)

SUEC. directs one part of root to be boiled a little with two parts of water; then to be boiled again with fresh water; the mixed decoctions to be evaporated in a water-bath.—FENN.; and FULD.—the same process to be followed, but six parts of water to be used the first time, and the ebullition to be continued for a quarter of an hour.—BRUGNATELLI—the first ebullition to be continued for two hours.

It is slightly stimulant or tonic, and has been much extolled in abdominal obstructions and dropsy. The liquid extract may be given, in the dose of a spoonful, several times a day.

RESOLVENT PILLS. (SAINTE-MARIE)

R. Extract of Dandelion, } of each,
White Soap, } one ounce.
Solution of Acetate of Potass,
a sufficient quantity.

Make pills of four grains each.—Dose, four or six, daily.

DEOBSTRUENT ELIXIR.

Elixir resolvens. (FULD.; HAMB.)

R. Extract of Dandelion, three drams.
Great Celandine, two drams.
Mint Water, eight ounces.

Mix. (HAMB.)

FULD. prescribes six drams of extract of dandelion, two of extract of great celandine, six ounces of distilled water, and two of peppermint water.

To be taken by spoonfuls.

TEREBINTHINA.—TURPENTINE.

Although it has lately been attempted to extend the appellation of turpentine to every liquid resin, yet custom has prevailed: the four following resinous substances are still only recognized under that denomination:

1. *Terebinthina Cypria* seu *Chia* seu *pistacina*.—Chio or Cyprian Turpentine.

SYNONIMES:—Térébenthine de Chio, de Scio, de Chypre ou de térébinthe, FR.; Cyprischer Terpenthin, GERM.

LOND.; BAVAR.; GALL.; FERR.; FORM. HÔP. FR.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; SAUNDERS; SPIELMANN.

A very thick, glutinous, transparent substance, of a greenish-citrine colour, of an agreeable odour, and of a pleasant aromatic taste without bitterness or acidity.

It flows from the *Pistacia Terebinthus*, LINN.; a small tree of the Archipelago. (*Diæcia Pentandria*, LINN.; *Terebinthaceæ*, JUSS.; Fig. Blackw. Herb. t. 478)

2. *Terebinthina communis* seu *Pini sylvestris* seu *pineæ*, *Balsamum nativum Pini sylvestris*.—Turpentine, Common Turpentine.

SYNONIMES:—Térébinthine commune, de Bourdeaux ou de pin, FR.; Gemeiner Terpenthin, GERM.; Ratinge rumie, ARAB.; Almindelig terperntin, DAN.; Trementina comun, SPAN.; Gemeen terperntin, DUT.; Trementina commune, ITAL.; Kota, NEP.; Zungbarie, PERS.; Tjok terperntin, SWED.

LOND.; DUBL.; EDINB.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; MIL. COPEN.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; HAMB.; HANN.; LIPP.; OLDEN.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; AINSLIE; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; PARIS.

A syrupy, whitish, turbid, consistent liquid, of a disagreeable odour, and of an acrid, bitter, and nauseous taste.

It flows from the *Pinus sylvestris*, LINN. (*Monæcia Monadelphia*, LINN.; *Conifera*, JUSS.; Fig. Flore Medic. V. 272), and from the *Pinus maritima*, LMK. (Fig. Zorn, Ic. pl. t. 526)

3. *Terebinthina laricea* seu *laigna* seu *laricina* seu *Veneta*, *Resina liquida Pini Laricis*, *Pini Laricis Terebinthina* seu *Balsamum*.—Venice Turpentine.

SYNONIMES:—Térébinthine de mélèze, ou de Venise, FR.; Venetianischer Terperntin, GERM.; Venedisk terperntin, DAN.; and SWED.; Termentina de Venecia, SPAN.; Terpentyna modrzewowa, POL.; Terebinthina fina, PORT.

EDINB.; DUBL.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SUEC.; WIRTEM.; WURCEB.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN.

This resin is rather liquid, transparent, of

a greenish hue, of a strong and not unpleasant odour, and of a warm, acrid, and bitter taste.

It is obtained by piercing the trunk of the *Larix Europæa*, CAND., a tree common on the Alps. (*Monæcia Monadelphia*, LINN.; *Conifera*, JUSS.; Fig. Blackw. Herb. t. 477)

4. *Terebinthina abiegna* seu *abietina* seu *Argentoratensis*.—Strasburg Turpentine.

SYNONIMES:—Térébinthine de sapin ou de Strasbourg, FR.; Strasbourgher Terperntin, GERM.; Trementina de abeto, SPAN.

ANT.; HISP.; GALL.; FORM. HÔP. FR.; LIPP.; BERGIUS; COXE; GUIBOURT; PARIS; SPIELMANN.

A rather fluid resin, transparent or slightly opaque, of a strong odour, and of an acrid and very bitter taste.

It is obtained from the *Abies pectinata*, CAND., a tree of Switzerland. (*Monæcia Monadelphia*, LINN.; *Conifera*, JUSS.; Fig. Zorn, Ic. pl. t. 528)

Like all the resins, turpentine is powerfully excitant: its action, when introduced into the stomach, at a certain dose variable according to the individual, is not confined to the digestive apparatus, but extends to other organs, either by sympathy or by absorption. Among these its secondary effects, those which manifest themselves upon the urinary apparatus were for a long time only observed, but those also which are perceptible in the respiratory organs and in the sensorium have been latterly attended to; so that turpentine, to which formerly only a specific action upon the urinary organs was assigned, is now also reputed to possess one not less specific, upon the thoracic viscera and upon the nerves. It is prescribed in gonorrhœa, particularly in the chronic stage, leucorrhœa, chronic catarrhs of the bladder, chronic catarrhs of the lungs, and in neuralgiæ.

Turpentine principally consists of two substances, a resin and a volatile oil. The practitioner should therefore consider it under three different forms, as it presents these two substances in natural combination, and as it presents each of them separately. This has not yet been sufficiently attended to, for the three substances do not act with the same energy upon the living tissues; if they were classed according to their energy, the first place should be assigned to the volatile oil, then would follow the unaltered turpentine, and lastly the pure residuum. We do not intend to follow this order, as it would, without sufficient occasion, interfere with the course followed in the other articles.

I. Preparations which contain the entire Turpentine.

A

Preparations which contain it dissolved in an alcoholic liquid.

TINCTURE OF TURPENTINE.

Tinctura seu Essentia Terebinthinæ; Baume vulnérable commun. (GALL.; VAN MONS)

R. Clear Turpentine, one part.
Alcohol (26 degrees), four parts.

Digest for six days, occasionally agitating, and strain.

B

Preparations which contain Turpentine suspended in an aqueous liquid.

EMULSION OF TURPENTINE.

Lac terebinthinaceum, Emulsio terebinthinacea. (NIEMANN; PIERQUIN; SAUNDERS; SWEDIAUR; VAN MONS)

R. Chio Turpentine, one ounce and a half.

Wash it well in proof spirit, and then mix it, in a marble mortar, with

Yolk of Egg, one,
adding gradually,

Water of Wall-Pellitory, twelve ounces.

Make an emulsion. (PIERQUIN)

R. Gum Arabic, three drams.
Water, a sufficient quantity,

to make a thick mucilage. Triturate this with Turpentine, two drams.

Adding gradually,
Water, nine ounces.

Simple Syrup, one ounce and a half.

Make an emulsion. (SWEDIAUR)

VAN MONS prescribes one dram of gum, two drams of turpentine, eight ounces of water, and one ounce of syrup.

R. Chio Turpentine, two drams.

Triturate it with a sufficient quantity of yolk of egg, and add

Common Julep, eight ounces.

Mix. (SAUNDERS)

NIEMANN prescribes half an ounce of turpentine, two yolks of eggs, two ounces of syrup of marshmallows, and eight ounces of water or of white wine.

R. Turpentine, two drams.
Mucilage of Gum Arabic, half an ounce.

Triturate, adding gradually,
Honey, one ounce and a half.

Juniper Water, nine ounces.
Spirit of Nitric Æther, one dram.

Mix. (VAN MONS)

It is excitant, and has been extolled in chronic diseases of the bladder and urethra.

TEREBINTHINE LOHOCH.

Lohoch leniens balsamicum. (SPIELMANN)

R. Gum Tragacanth, in powder, sixteen grains.
Common Ptisan, half an ounce.

Make a mucilage, and incorporate gradually,

Syrup of Marshmallows, } of each,
Oil of Almonds, } one dram.

Turpentine, triturated with Yolk of Egg, one scruple.

Pectoral Ptisan, three ounces and a half.

Mix.

POTION OF TURPENTINE.

Potio terebinthinata. (SWEDIAUR)

R. Venice Turpentine, four pounds.
Myrrh, one ounce.

Triturate, adding gradually,
Water, twenty-four pounds.

Set aside, and put the decanted liquor into a dozen bottles.

It is recommended in diseases of the urinary and respiratory passages, cutaneous affections, dysmenorrhœa, leucorrhœa, hysteria, in the suppurative stage of variola, &c.—It is administered in the same as a tar water.—Sometimes an ounce of sub-carbonate of potass may be added to it with advantage.

DIURETIC MIXTURE. (BORIES; PIERQUIN)

R. Turpentine, one dram.
Yolk of Egg, one.

Triturate, and add

Parsley Water, } of each,
Strawberry Water, } three ounces.
Honey, half an ounce.
Syrup of Marshmallows, one ounce.

A spoonful to be taken four times a day.
—BARTHEZ employed it extensively.

BALSAMIC MIXTURE. (BORIES)

R. Turpentine, two drams.
Yolk of Egg, one.
Extract of Rhubarb, } of each,
Nitrate of Potass, } one dram.
Infusion of Liquorice, ten ounces.
Syrup of Marshmallows, one ounce.

Mix.—BOERHAAVE prescribed it in gonorrhœa.—Dose, a spoonful every half hour.

TEREBINTHINATE INJECTION. (NIEMANN; SAUNDERS)

R. Strasburg Turpentine, three drams.

Triturate it with a sufficient quantity of yolk of egg, and add gradually,

Thick Infusion of Linseed, one pound.

Make an emulsion. (SAUNDERS)

R. Venice Turpentine, half an ounce,

Distilled Water, eight ounces.

Tincture of Benzoin, two drams.

Mix intimately. (NIEMANN)

DETHARDING recommended the latter in deafness and dulness of hearing. The former is used in leucorrhœa.

TEREBINTHINATE LAVEMENT. (FORM. HÔP. FR.; NIEMANN; SARD.; SPIELMANN; SWEDIAUR)

R. Venice Turpentine, from half an ounce to one ounce.

Triturate it with yolk of egg or mucilage, and add gradually,

Spring Water, six or eight ounces.

Mix. (FORM. HÔP. FR.; NIEMANN)

R. Venice Turpentine, one ounce.

Triturate with yolk of egg, and add

Decoction of Barley, six ounces.

Mix. (SWEDIAUR)

R. Venice Turpentine, any quantity.

Yolk of Egg, a sufficient quantity.

Decoction of Barley, six ounces.

Honey, or Confection of Senna, six drams.

Mix. (SAUNDERS)

R. Turpentine, triturated with yolk of egg, half an ounce.

Emollient Decoction, ten ounces.

Linseed Oil, one ounce.

Triturate together. (SPIELMANN)

R. Venice Turpentine, triturated

with yolk of egg,

from one dram to half an ounce.

Emollient Decoction, one pound.

Linseed Oil, } of each,

Syrup of Buckthorn, } one ounce.

Mix. (SWEDIAUR)

It is recommended in constipation, nephritis, calculous ischuria, and colic.

SEDATIVE LAVEMENT. (AUGUSTIN; SAINTE-MARIE)

R. Turpentine, one dram.

Fresh Yolk of Egg, one.

Triturate, and add

Theriac, four drams.

Warm Milk, four ounces.

Mix.—VAN SWIETEN employed it to appease the tenesmus and intestinal disturbances of consumptive patients. It has also been successfully employed in what has been called intestinal consumptions, that is, in those which succeed acute or chronic enteritis. (SAINTE-MARIE)

AUGUSTIN prescribes eight ounces of milk, half an ounce of turpentine triturated with yolk of egg, half an ounce of theriac, and one ounce of honey of roses.

ASTRINGENT LAVEMENT. (SAINTE-MARIE)

R. Turpentine, four drams.

Linseed Oil, two ounces.

Yolk of Egg, two drams.

Triturate in a mortar, adding gradually,

Lime Water, six ounces.

Mix.—It is prescribed by WINTRINGHAM in the same cases as the preceding.

ASTRINGENT BALSAMIC LAVEMENT. (BORIES; PIERQUIN)

R. Chamomile Flowers, two handfuls.

Milk, a sufficient quantity,

to obtain eight ounces of strained decoction. Add

Turpentine, triturated with yolk of egg, three drams.

Honey of Roses, one ounce.

Mix. (BORIES)

R. Saffron, } of each,

St. Johns-wort, } one pinch.

Milk, a sufficient quantity.

Boil and add to the strained liquor,

Turpentine, triturated with two

yolks of eggs, half an ounce.

Theriac, two drams.

Mix. (BORIES; PIERQUIN)

This lavement may be useful in the same cases as the preceding.

TEREBINTHINATE INJECTION.

Injectio terebinthinacea. (AUGUSTIN)

R. Turpentine, two ounces.

Yolk of Egg, one ounce.

Honey, two ounces.

Alcohol, half an ounce.

Water, nine ounces.

Mix.

R. Medicinal Soap, one ounce.

Turpentine, half an ounce.

Water, eight ounces.

Tincture of Benzoin, three drams.

Mix.

C.

Preparations which contain Turpentine mixed with fat bodies.

ASTRINGENT MIXTURE. (NIEMANN)

- R. Venice Turpentine, }
Copaiba, } of each,
Oil of Amber, } an equal part.

Mix.—It is recommended in blenorrhœa, and seminal emissions.—Dose, from thirty to sixty drops, thrice a day, with a spoonful of sugar, taking afterwards a glass of wine.

ANTI-NEPHRITIC MARMALADE.

Conditum nephriticum. (VAN MONS)

- R. Turpentine, four scruples.
Oil of Almonds, two ounces.
Extract of Juniper, four ounces.
Prepared Crab's Eyes, half an ounce.

Mix.—It is useful in chronic nephritis.—Dose, two drams, several times a day.

OINTMENT OF TURPENTINE.

Unguentum Terebinthinatum. (AUSTR.)

- R. Simple Ointment, nine ounces.
Turpentine, two pounds.

Melt together over a slow fire, and stir till cold.

COMMON SPARADRAP. (AUSTR.; FORM. HÔP FR.; PALAT.; VAN MONS)

- R. Turpentine, one part.
Olive Oil, four parts.
Yellow Wax, eight parts.

Melt together in a water-bath, and spread it over strips of linen.

VAN MONS prescribes one part of turpentine, one and a half of mutton suet, and four parts of white wax.—AUSTR.—six ounces of yellow wax, two of mutton suet, and one and a half each, of lard and of turpentine.—PALAT.—half a pound of white wax, four ounces of stag's suet, one ounce and a half of turpentine, and a sufficient quantity of rose oil.

EMOLLIENT OINTMENT.

Unguentum lenitivum. (GENEV.; SWEDIAUR)

- R. Mucilage, one pound.
Fresh Butter, two pounds.
Yellow Wax, half a pound.

Boil over a slow fire to the consumption of the humidity, and add

Venice Turpentine, one ounce.

Mix.—There are sometimes added to each pound of this ointment, three ounces of

spirit of turpentine and one dram of opium.—It is recommended in milky enlargements of the breasts, nephritic pains, ischury, and rigidity of the joints. (SWEDIAUR)

GENEV. prescribes three pounds and a half of olive oil, one pound of yellow wax, half a pound of resin, and three ounces of turpentine.

CAUSTIC PAPER.

Papier à cautère. (CADET DE GASSI-COURT)

- R. Pure Turpentine, } of each, one
White Wax, } ounce and a half.
Spermaceti, } one dram.

Melt in a water-bath, and spread the mass upon smooth paper.

CAUSTIC PEAS.

Pisa pro fomiculis. (NIEMANN)

- R. Venice Turpentine, any quantity.
Yellow Wax, one ounce and a half.
Powdered Turmeric, one ounce.
Iris, half an ounce.

Mix, and form masses of the size of peas.

DEPILLATORY OINTMENT. (SWEDIAUR)

- R. Venice Turpentine, ten drams.
Resin, one ounce.

Melt together and preserve it in cold water.—A little is to be taken between the fingers wetted, the hairs seized, and torn up by a quick movement.

EMPLASTRUM GRATIA DEI. (VAN MONS)

- R. Yellow Wax, } of each, nine parts.
Resin, }

Melt together, and add

Turpentine, five parts,
then, when the mass is becoming solidified,
Powdered Olibanum, three parts.

Mix.

FRAHM'S BALSAM.

Unguentum terebinthinatum seu Balsami et Olei Laricis, Balsamum Frahmii. (BAVAR.; NIEMANN; SWEDIAUR; VAN MONS)

- R. Yellow Wax, one part.

Melt it over a slow fire; then add a mixture, melted apart, of

Venice Turpentine, } of each,
Spirit of Turpentine, } one part.

Mix. (BAVAR.; NIEMANN; VAN MONS)

SWEDIAUR prescribes one part of the spirit, and four parts each, of wax and of turpentine.

It is much extolled as an application to chronic ulcers of the legs.

ANTI-HÆMORRHOIDAL LINIMENT. (CADET DE GASSICOURT)

R. Venice Turpentine, }
Honey, } of each, one ounce.
Olive Oil, }
Triturate together.

DIGESTIVE LIQUOR.

Digestivum liquefactum. (SPIELMANN)

R. Turpentine, two ounces.
Yolk of Egg, one ounce.
Triturate, adding gradually,
Honey, two ounces.
Spring Water, four ounces.
Alcohol, six ounces.

It is employed by PLENK, as an injection, in fistulæ.

DEMULCENT AND BALSAMIC OINTMENT.
(SAINTE-MARIE)

R. Proof Spirit, }
Wheat Flour, }
White Sugar, } of each, one ounce.
— of Egg, }
Turpentine, }
Olive Oil, }
Honey, }

Incorporate the honey with the turpentine; then add the other ingredients gradually.

It is a useful application to ulcers, excoriations, and wounds attended with severe pain.

ARCEUS' BALSAM; COMPOUND OINTMENT
OF ELEM.

Balsamum Arcai, Unguentum Amyridis elemifera seu Arcai seu balsamicum seu basilicum seu digestivum flavum seu Elemi seu Elemi compositum seu Elemi et Terebinthina seu Resina flava seu Resina Elemi seu Sebi Arcai. (LOND.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; WIRTEM.; BRUGNATELLI; CADET DE GASSICOURT; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Unguentum Elemi compositum.
(LOND.)

R. Elemi, one pound.
Turpentine, ten ounces.

Prepared Suet, two pounds.
Olive Oil, two fluid ounces.

Melt the elemi with the suet; then remove the mixture from the fire, add the turpentine and oil, and strain the whole through a linen cloth.

R. Turpentine, sixteen ounces.
Elemi, two pounds.
Resin, two pounds and a half.
Olive Oil, one pound and a half.
Yellow Wax, four ounces.

Melt together. (FERR.)

R. Venice Turpentine, }
Elemi, } of each,
Stag's Suet, } two ounces.
Oil of St. John's-wort, one ounce.
Red Sanders, one dram.

Melt the elemi with the turpentine over a slow fire, then add the oil and suet melted together, and finally incorporate the powdered sanders wood. (SARD.; WIRTEM.)

R. Turpentine, }
Elemi, } of each, six parts.

Melt together over a slow fire, constantly stirring; then remove from the fire, and incorporate gradually

Melted Stag's Suet, six parts.
Oil of St. John's-wort, two parts.

Strain through a flannel, and triturate till perfectly cold. (LIPP.)

AMST.; ANT.; AUSTR.; BAVAR.; BELG.; OLDEN.; POL.; BORUS.; and SAX. prescribe an equal part each, of turpentine, of elemi, of lard, and of mutton suet.—LUSIT.; and COXE—the same as LOND.—GALL.; and FORM. HÔP. FR.—three parts each, of turpentine and of elemi, six of suet, and five of lard.—HISP.—five each, of turpentine, of elemi, and of lard, and six of suet.—ROSS.—six each, of turpentine and of elemi, eight of lard, and four of suet.—FULD.; HASS.; and PIDERIT—three each, of turpentine, of elemi, and of stag's suet, and two of oil of St. John's-wort.—SPIELMANN—six each, of turpentine and of elemi, nine of stag's suet, and two of St. John's-wort.—BRUNS.; and PALAT.—six each, of turpentine and of elemi, two of goat's suet, and one of oil of St. John's-wort.

R. Turpentine, }
Elemi, } of each, six parts.
Mutton Suet, ten parts.

Melt together. (BRUGNATELLI)

ANT. prescribes one part each, of turpentine and of elemi, and two parts of mutton suet.—GENEV.—fourteen of turpentine, twelve of elemi, and twenty-four of mutton suet.

This stimulant ointment is rather exten-

sively used in surgical practice : it is one of the best of the kind.—The third formula is that of the inventor.

VULNERARY BALSAM. (WIRTEM. ; NIE-MANN)

R. Venice Turpentine, one ounce and a half.
Elemi, half an ounce.

Add to the melted mixture

Oil of St. John's-wort, one ounce and a half.
— Wax, two scruples.

Mix. (WIRTEM.)

R. Venice Turpentine, } of each, one
Elemi, } ounce and a half.
Goat's Suet, two ounces.
Balsam of Tolu, one ounce.

Make an ointment.

It is for the same purposes as the preceding, from which it scarcely differs.

CAMPHORATED OINTMENT OF TURPENTINE. (FORM. HÔP. FR.)

R. ARCÆUS' Balsam, one hundred and fifty parts.

Camphor, one part.

Triturate together.

PLENK'S DIGESTIVE.

Baume d'Arcaus liquide. (CADET DE GASSICOURT ; SPIELMANN ; VAN MONS)

R. ARCÆUS' Balsam, } of each, one ounce.
Yolk of Egg, }

Triturate, and mix with

Proof Spirit, eight ounces.

Preserve. (SPIELMANN)

R. Clear Turpentine, } of each,
Honey, } two ounces.
Yolk of Egg, one ounce.
Alcohol, six ounces.
Water, four ounces.

Triturate together. (CADET DE GASSICOURT)

R. ARCÆUS' Balsam, two parts.
Rectified Petroleum, one part.

Triturate together. (VAN MONS)

It is suppurative,—It may be employed in injections.

SIMPLE DIGESTIVE OINTMENT.

Unguentum digestivum seu digestivum simplex seu lacteum, Unguentum Terebinthinae seu de Terebinthina, Balsamum terebinthinatum. (DAN. ; GALL. ; FULD. ; HASS. ; LIPP. ; CAST. BORUS. ; ROSS. ; WURCEB. ; BRUGNATELLI ; PIDERIT ; RATIER)

R. Venice Turpentine, one pound.

Yolks of Eggs, eighty.

Mix. (ROSS.)

R. Venice Turpentine, one pound.
Yolks of Eggs, twelve.

Mix, and add

Honey of Roses, half a pound.

Preserve. (WURCEB.)

R. Turpentine, half a pound.
Yolks of Eggs, four.

Triturate together, and add

Oil of St. John's-wort, one ounce and a half.

Mix. (HASS. ; PIDERIT)

GALL. prescribes two parts of turpentine, one part of yolk of egg, and a sufficient quantity of the oil to obtain a diffuent ointment.—LIPP.—one pound of turpentine, eight yolks of eggs, and three ounces of olive oil.

R. Venice Turpentine, } of each,
Olive Oil, } one pound.
White Wax, half a pound.

Melt together, and stir till perfectly cold. (BRUGNATELLI)

CAST. BORUS. prescribes two parts of wax, two of turpentine, and six of lard.

R. Venice Turpentine, two drams.
Yellow Wax, one dram and a half.

Melt slowly, and add

Spirit of Turpentine, two drams.

Mix. (DAN.)

R. Venice Turpentine, half an ounce.
Honey of Roses, one ounce.
Oil of St. John's-wort, three drams.
Yolk of Egg, one.

Mix. (FULD.)

R. ARCÆUS' Balsam, one ounce.
Cerate, two ounces.
Yolk of Egg, one.

Mix. (RATIER)

It is applied as a stimulant to wounds and ulcers.

COMPOUND DIGESTIVE OINTMENT.

Unguentum digestivum aloeticum seu myrrhatum. (BRUNS. ; FULD. ; HANN. ; LIPP. ; OLDEN. ; PALAT. ; POL. ; BORUS. ; SAX. ; SUEC. ; WIRTEM. ; PIDERIT ; RATIER ; SPIELMANN ; SWEDIAUR ; VAN MONS)

1. With Aloes.

R. Aloes, one part.
Honey of Roses, four parts.

Triturate together, and add

Clear Turpentine, twelve parts.

- Oil of St. John's-wort, three parts.
 Mix. (HANN.; BORUS.; VAN MONS)
- R. Venice Turpentine, half a pound.
 Honey, four ounces.
 Tincture of Aloes, three ounces.
 Yolks of Eggs, four.
 Mix. (POL.)
- R. Turpentine, two ounces.
 Oil of St. John's-wort, half an ounce.
 Mix, and add
 Yolk of Egg, one.
 Tincture of Aloes, } of each, one
 Myrrh, } dram and a half.
 Mix well. (VAN MONS)

2. Without Aloes.

- R. Venice Turpentine, half a pound.
 Yolks of Eggs, four.
 Triturate together, and add
 Powdered Myrrh, six drams.
 Olive Oil, one ounce and a half.
 Mix. (SUEC.)
- R. Venice Turpentine, nine ounces.
 Oil of Myrrh, one ounce and a half.
 Add
 Yolks of Eggs, two.
 Mix well, (SWEDIAUR)
- R. Strasburg Turpentine, eight ounces.
 Yolks of Eggs, four.
 Triturate together, and add,
 Oil of St. John's-wort, one ounce and a half.
 Powdered Olibanum, } of each,
 Myrrh, } three drams.
 Mix. (BRUNS.; WIRTEM.; PIDERIT; SPIELMANN)
- PALAT. prescribes one pound of turpentine, three yolks of eggs, three ounces of oil of St. John's-wort, and half an ounce each, of the two powders.
- R. Turpentine, half a pound.
 Yolks of Eggs, four.
 Triturate together, and add
 Oil of St. John's-wort, one ounce and a half.
 Powdered Myrrh, } of each,
 Olibanum, } three drams.
 Saffron, macerated in alcohol, a sufficient quantity
 to colour the mass. (OLDEN.)
- R. ARCAÛS' Balsam, two ounces.
 Yolks of Eggs, two.
 Olive Oil, two ounces.
 Mix. (RATIER)
- R. ARCAÛS' Balsam, } of each,
 Ointment of Storax, } two drams.
 Yolks of Eggs, two.

- Spirit of Turpentine, one ounce.
 Mix. (RATIER)

FULD. directs two drams each, of myrrh and of olibanum, to be added to two ounces of the simple digestive ointment.—LIPP.—adds four grains each, of myrrh and of olibanum to each dram.—The detergent property of this ointment may also be increased by the addition of *Unguentum Egyptiacum* (GALL.), or of verdigris (SWEDIAUR).

OPIATE DIGESTIVE OINTMENT, (RATIER)

- R. Simple Digestive Ointment, four ounces.
 SYDENHAM's Laudanum, one ounce.
 Mix.

Two drams of powdered opium may be substituted for the laudanum.

LOCATELLI'S BALSAM.

Balsamum Locatelli seu Italicum seu Italicum rubrum. (AMST.; BATAV.; BRUNS.; FULD.; GENEV.; LIPP.; PALAT.; SAX.; SARD.; WIRTEM.; HENRI; SPIELMANN; VAN MONS)

- R. Olive Oil, one pound.
 Turpentine, } of each, half a pound.
 Yellow Wax, }
 Red Sanders, one ounce.

Melt the wax with a portion of the oil; then add the remainder and the turpentine, and finally the wood in powder. Stir till perfectly cold. (GENEV.)

- R. Turpentine, four parts.
 Red Sanders, one part.

Digest for an hour in warm ashes in a covered vessel; then melt separately,

- Yellow Wax, six parts.

Add
 Olive Oil, twelve parts,
 and finally the wood and turpentine.

Leave the mixture over the fire for a quarter of an hour; then express strongly, and add at the moment of its cooling,

- Balsam of Peru, one part.

Stir till cold. (VAN MONS)

- R. Yellow Wax, four ounces.
 Olive Oil, six ounces.

Melt over a slow fire, and add

- Venice Turpentine, six ounces.
 Red Sanders, half an ounce.

Strain, and add further,

- Balsam of Peru, two drams.

Preserve. (BATAV.)

AMST. prescribes four ounces of wax, six of oil, six of turpentine, one ounce of wood, and one of balsam.—PALAT.—six ounces

of wax, nine of oil, five of turpentine, a sufficient quantity of wood, and six drams of balsam.

- R. Olive Oil, six ounces.
 Yellow Wax, four ounces.
 Madeira Wine, five ounces.

Boil slowly to the consumption of the humidity, and add

- Venice Turpentine, six ounces.
 Balsam of Peru, two drams.
 Red Sanders, half an ounce.

Preserve. (WIRTEM.; SPIELMANN)

HENRI prescribes six ounces of wax, nine of oil, two of Spanish wine, nine of turpentine, one of red sanders, and one and a half of balsam of Peru.

- R. Olive Oil, two pounds.
 Madeira Wine, one pound.
 Venice Turpentine, eight ounces.

Boil slowly to the consumption of the humidity, and add

- Balsam of Peru, } of each, one ounce.
 Dragon's Blood, }
 Storax, }

Stir till perfectly cold. (SARD.)

- R. Yellow Wax, one part.
 Olive Oil, one part and a half.

Add to the melted mixture,

- Venice Turpentine, one part and a half.

Add further, when taken from the fire,

- Black Balsam of Peru,

- Kino, one sixteenth part.
 one twelfth part.

Preserve. (BRUNS.; SAX.)

- R. Olive Oil, four ounces.
 Yellow Wax, one ounce and a half.

Melt slowly, and add

- Turpentine, three ounces.

Remove from the fire, and add further, constantly stirring,

- Copaiba, half an ounce.
 Kino, three drams.

Preserve. (FULD.; LIPP.)

It is employed externally, as a detergent and vulnerary.—It is seldom used now.

CAMPHORATED OINTMENT OF TURPENTINE. (GALL.; CADET DE GASSICOURT; VAN MONS)

- R. Red Wine, } of each, three pounds.
 Olive Oil, }
 Yellow Wax, } of each, half a pound.
 Rose Water, }
 Red Sanders, two ounces.

Boil for half an hour, stirring with a wooden spatula. Add

- Venice Turpentine, one pound.

Incorporate; and when the mass is nearly cold, add

- Camphor, in powder, two ounces.

Strain; and after some hours' repose, express. (CADET DE GASSICOURT)

- R. Yellow Wax, half a pound.
 Olive Oil, one pound.

Melt; then keep the mixture over the fire, for an hour, with

- Red Sanders, two ounces.

Then add

- Turpentine, one pound.

Express strongly; and when the mass is nearly cold, add to it,

- Camphor, in powder, two drams.

Stir till perfectly cold. (VAN MONS)

- R. Olive Oil, three hundred and eighty-four parts.
 Yellow Wax, sixty-four parts.
 Red Sanders, fourteen parts.
 Turpentine, one hundred and twenty-eight parts.

Digest in a gentle heat; and just before it becomes cold, add

- Camphor, two parts.

Mix. (GALL.)

It used formerly to be extolled as an application to bruises, ulcerations, and even gangrene.

BALSAMUM CHIRON. (HENRI)

- R. Olive Oil, six ounces.
 Turpentine, two ounces.
 Yellow Wax, one ounce.
 Root of Alkanet, a sufficient quantity.

Having melted these together, add successively

- Black Peruvian Balsam, one dram and a half.

- Camphor, in powder, twelve grains.

Mix.

This is scarcely different from the preceding.

PECTORAL BALSAM. (SPIELMANN)

- R. Turpentine, } of each,
 Oil of Almonds, } eighteen drams.
 — St. John's-wort, two ounces.
 Spermaceti, half an ounce.
 Dragon's Blood, } of each,
 Aqueous Extract of Opium, } two
 Balsam of Peru, } drams.
 Butter, one ounce and a half.

Mix.

MEIBOM directed from ten to fifteen drops to be taken daily, in the chimerical hope of thus preventing pulmonary consumption.

SPARADRAP OF PLASTERS. (GALL.)

- R. Simple Diachylon Plaster, } of each,
 Gummy Diachylon Plaster, } fifteen
 Plaster of Yellow Wax, } parts.
 — Turpentine, twenty-five parts.

Melt together in a water-bath, and spread the mass upon strips of linen.

ANTI-APOPLECTIC PLASTER. (WIRTEM.; SPIELMANN)

- R. Venice Turpentine, three drams.
 Galbanum, } of each, two drams.
 Opoponax, }
 Root of Pyrethrum, } of each,
 Long Pepper, } one dram.
 Castor, }
 Oil of Amber, } one scruple.

Make a plaster. (WIRTEM.)

- R. Walnut Oil, twelve ounces.
 Turpentine, four ounces.
 Resin, twenty ounces.
 Yellow Wax, sixty-four ounces.

Melt together, and add,

- | | |
|----------------------------|--|
| Powdered Amber, | } of each,
one ounce
and a half. |
| — Tacamahac, | |
| — Caranna, | |
| — Sandarac, | |
| — Olibanum, | } of each,
one ounce. |
| — Mastic, | |
| — Storax, | |
| — Benzoin, | |
| — Root of Round Birthwort, | } of each,
half an ounce. |
| — Pyrethrum, | |
| — Angelica, | |
| — Iris, | |

Mix; and when the mass is half cold, add to it,

- | | |
|-----------------|--------------------------------|
| Fat of Castor, | } of each, two drams. |
| Camphor, | |
| Oil of Mace | } of each,
two
scruples. |
| — Marjoram, | |
| — Cloves, | |
| — Rhodium Wood, | |
| — Lavender, | } of each,
twelve drops. |
| — Sage, | |
| — Rosemary, | |
| — Chamomile, | |
| — Amber, | |

Mix. (SPIELMANN)

It used formerly to be applied, in apoplexy and paralysis, to the shaven head.

BALSAMUM VIRIDE. (PALAT.)

- R. Herb of Lavender, }
 — Sage, } of each,
 — Hyssop, } half
 — Marjoram, } an ounce.
 — Thyme,
 — Caraway,
 — Water-mint,

- | | |
|------------------------|---------------------------------|
| Herb of Mentha Crispa, | } of each,
half
an ounce. |
| — Plantain, | |
| — Rosemary, | |
| — Ladies-Mantle, | |
| — Golden Rod, | } |
| Flowers of Chamomile, | |
| — Red Roses, | |

- | | |
|---------------|------------------------|
| Olive Oil, | } of each, two pounds, |
| Oil of Roses, | |
| Linseed Oil, | one pound. |

Infuse; then add

- | | |
|-------------|-----------------------|
| Turpentine, | } of each, one pound. |
| Sagapenum, | |
| Olibanum, | half a pound. |

Digest in a water-bath; then express strongly. Add to the mass when half cold,

- | | |
|-----------------------|----------------------------|
| Oil of Lavender, | } of each,
four ounces. |
| Spirit of Turpentine, | |
| Petroleum, | |

Decant after a few days.

It used to be formerly employed internally and externally, as stimulant, alterative, and agglutinant.

D.

Preparations which contain Turpentine mixed with dry substances.

PILLS OF TURPENTINE. (MIL. COPEN.; GENEV.; CADET DE GASSICOURT; ELLIS; PIERQUIN; SAINTE-MARIE; SWEDIAUR)

- R. Guaiac Resin, one dram.
 Turpentine, a sufficient quantity
 to make fifteen pills.—Dose, one, thrice a day. (ELLIS)

- R. Turpentine, half a dram.
 Balsam of Tolu, two scruples.
 Crabs' Eyes, a sufficient quantity.

Make pills of three grains each. (GENEV.)

- R. Turpentine, } of each, two drams.
 Rhubarb, }

Make pills of two grains each. (MIL. COPEN.)

Dose, five.

- R. Gum Arabic, half an ounce.
 Water, a sufficient quantity

to make a mucilage.

- Turpentine, six drams.
 Aperitive Powder, a sufficient quantity.

Make pills of five grains each. (VAN MONS)

Dose, three, four times a day.—The turpentine may be substituted by copaiba.

- R. Turpentine, two drams.
 Powdered Liquorice, a sufficient quantity.

Make pills of four grains each. (CADET DE GASSICOURT; SAINTE-MARIE)

These various pills are principally prescribed as diuretic. The third formula is by DE HAEN.

ANTI-GONORRHOIC BOLUSES. (PIERQUIN ;
SAINTE-MARIE) .

R. Venice Turpentine, one or two scruples.

Triturate with yolk of egg, and add

Liquorice Powder, a sufficient quantity
to make four boluses.

Dose, one every six hours.—This formula is by ASTRUC.

ASTRINGENT PILLS. (SAINTE-MARIE)

R. Venice Turpentine, two ounces.
Extract of Rhubarb, three drams.
Camphor, two drams.

Make pills of four grains each.

They are prescribed in leucorrhœa and
blenorrhœa.—Dose, three, thrice a day.

TRAUMATIC PILLS. (FULD.)

R. Venice Turpentine, } of each,
Extract of Liquorice, } one dram.
Milfoil, }
Powdered Mastic, } of each,
Olibanum, } half a dram.

Make a pillular mass.

Dose, from ten to twenty grains.

BALSAMIC PILLS. (CADET DE GASSI-
COURT)R. Chio Turpentine, } of each,
Spermaceti, } half an ounce.
Myrrh, } two drams.
Olibanum, } a sufficient quantity.Make pills of three grains each.—Dose,
one, every three hours, in pulmonary con-
sumption.RESOLVENT PILLS: (PIERQUIN ; SAINTE-
MARIE)R. Turpentine, twenty drops.
Extract of Liquorice, two drams.
White Soap, four drams.
Compound Syrup of Smallage,
a sufficient quantity.Make pills of four grains each. (PIER-
QUIN ; SAINTE-MARIE)Dose, three, thrice a day ; in cases of
biliary calculi.R. Turpentine, two scruples.
Extract of Rhubarb, one dram.
Dandelion, four drams.
White Soap, two drams.
Compound Syrup of Smallage,
a sufficient quantity.Make pills of four grains each. (SAINTE-
MARIE)To be taken in the same dose as the pre-
ceding, in abdominal obstructions.

These two formulæ are by TRONCHIN.

PILLS AGAINST TÆNIÆ. (AUGUSTIN)

R. Venice Turpentine, one dram.
Jalap Soap, half a dram.
Extract of Henbane, four grains.
Proto-chloruret of Mercury,
eight grains.Make pills of twelve [*two ?* EDIT.] grains
each.HUFELAND directs four to be taken every
three hours, for two or three days succes-
sively.

CYPHEOS TROCHISCS.

Trochisci Cyphæos, Cyphoides Galli. (PA-
LAT ; SPIELMANN)R. Chio Turpentine, six drams.
Pulp of Raisins, two ounces and a half.
Bdellium, half an ounce.
Myrrh, }
Aloe Wood, } of each, one dram.
Saffron, }
Cinnamon, }
Spikenard, }
Cassia Lignea, }
China Root, } three drams.
Sweet Flag, } two drams.
Honey, } five ounces.
Greek Wine, a sufficient quantity.

Mix carefully. (SPIELMANN)

R. Raisins, } of each,
Yellow Resin, } three ounces.
China Root, } of each,
Myrrh, } one ounce and a half.
Sweet Flag, } nine drams.
Cinnamon, } four drams.
Bdellium, }
Spikenard, } of each, three drams.
Cassia Lignea, }
Round Cyperus, }
Juniper, }
Asphaltum, } two drams and a half.
Saffron, } one dram.
Honey, } of each,
Cretan Wine, } a sufficient quantity.Dissolve the bdellium, myrrh, and resin
in the wine ; then having evaporated the
solution to the consistence of honey, mix it
with the honey, the raisins triturated to-
gether for a long time, and add the other
substances reduced to fine powder. Dry
the trochiscs in the shade. (PALAT.)These trochiscs enter into the mithri-
date.II. *Preparations which contain the oil or
spirit of Turpentine exclusively or princi-
pally.*

OIL OR SPIRIT OF TURPENTINE.

*Ætheroleum seu Spiritus seu Alcohol seu
Oleum Terebinthina, Oleum volatile seu*

æthereum Pini seu Laricis, Spiritus Olei Terebinthina æthereus.

1. *That which is met with in commerce.* (LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; FENN.; FORM. HÔP. FR.; LIPP.; ROSS.; SUEC.; COXÆ; GUIBOUT; MURRAY and GMELIN; PARIS; SPIELMANN)

A colourless liquid, of a disagreeable and very strong odour, and of an acrid and pungent taste. Its specific gravity is 0.870.

From the *Pinus Mugho* is obtained a similar product, but of a greenish colour and pleasant odour, bearing the name of *Oleum templinum*.

Several pharmacopœias recommend the spirit of commerce to be rectified by distilling it with water.

Oleum Terebinthinæ rectificatum.

- R. Oil of Turpentine, one pint.
Water, four pints.

Distil the oil. (LOND.)

- R. Oil of Turpentine, two pints.
Water, four pints.

Distil one pint and a half of the oil. (DUBL.)

Oleum Volatile Pini purissimum.
(EDINB.; COXE)

- R. Oil of Turpentine, one part.
Water, four parts.

Distil as long as any oil passes over.

ANT.; BAVAR.; FENN.; and BORUS. prescribe one part of oil and two parts of water.

VAN MONS observes that at every such rectification, part of the oil becomes decomposed, that acetic acid is formed, that more or less residuum is left, and that the oil becomes less and less soluble in alcohol.

2. *Prepared by the apothecary.* (DUBL.; ANT.; AUSTR.; BRUNS.; HISP.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; PIDERIT; SWEDIAUR; VAN MONS)

Oleum Terebinthinæ. (DUBL.)

- R. Common Turpentine, five pounds.
Water, four pints.

Distil the oil from a copper alembic. Yellow resin remains in the retort after the distillation.

- R. Venice Turpentine, } of each,
Water, } an equal weight.

Put them into a glass cucurbit placed in a sand-bath; then lute the capital, and distil. Separate the distilled oil from the water. (VAN MONS)

AUSTR.; HANN.; OLDEN.; POL.; BORUS.; ROSS.; and SWEDIAUR give no relative proportions of the water and turpentine.—AMER.; and COXE give the same formula as DUBL.—HASS.; SUEC.; HERBIP.; and PIDERIT—one part of turpentine and three parts of water.—FULD.; LIPP.; PALAT.; SAX.; and WIRTEM.—one of turpentine and six of water.—LUSIT.—five of turpentine and six of water.—HISP.—one of turpentine and forty of water.

The oil or spirit of turpentine is more stimulant than the turpentine itself; it has been recommended and employed in the same cases.

Preparations which contain spirit of turpentine dissolved in æther.

TEREBINTHINATED ÆTHER. (CADET DE GASSICOURT; VAN MONS)

- R. Sulphuric Acid, } of each,
Rectified Alcohol, } an equal part.

Mix gradually together, and distil the æther. Pour into this,

Spirit of Turpentine,
a sufficient quantity

to saturate it. Shake the mixture well, set it apart for a day, and distil. (VAN MONS)

- R. Alcohol, two pounds.
Spirit of Turpentine, half a pound.

Mix, and add gradually,
Concentrated Nitric Acid, two pounds.

Distil one half of the mixture, at a gentle heat. (CADET DE GASSICOURT; VAN MONS)

It is employed both externally and internally, in cases of biliary calculi, jaundice, engorgements of the liver, and rheumatism.

Dose, from twenty to forty drops, in honey or yolk of egg.

ÆTHEREAL LINIMENT OF TURPENTINE.
(VAN MONS)

- R. Spirit of Turpentine, any quantity.
Spirit of Nitric Æther,
a sufficient quantity

to dissolve the oil.

It is for the same purposes as the preceding.

RESOLVENT MIXTURE. (MIL. COPEN.; AUGUSTIN.; CADET DE GASSICOURT.; NIEMANN.; PIERQUIN)

R. Spirit of Turpentine, one part.
Sulphuric Æther, three parts.

Dissolve. (AUGUSTIN.; NIEMANN)

MIL. COPEN.; and CADET DE GASSICOURT prescribe one part of spirit and two of æther.—PIERQUIN—two of spirit and three of æther.—CADET DE GASSICOURT, in another place,—one of spirit and fourteen of æther.

It is recommended by DURANDE in cases of biliary calculi.—Dose, from two scruples to a dram, daily, in whey or broth.

DIURETIC MIXTURE. (CADET DE GASSICOURT)

R. Spirit of Turpentine, one ounce.
Scurvy-grass, half an ounce.
Hydrochloric Acid, of each,
Alcoholized Nitric Acid, three drams.
Succinic Acid, of each,
Compound Tincture of Aloes, two drams.

Mix.—Dose, from four to fifteen drops, in white wine, for several days.

B.

Preparations which contain spirit of turpentine dissolved in alcoholic liquors, or mixed with water.

DIURETIC WINE. (PIERQUIN)

R. Spirit of Turpentine, two drams.
Lemon Juice, one ounce.
White Wine, four ounces.

Mix.—To be taken at one dose.

ANTI-ICTERIC SPIRIT.

Spiritus anti-ictericus. (GENEV.; AUGUSTIN.; NIEMANN)

R. Spirit of Turpentine, one ounce and a half.
Alcohol, half a pound.

Distil at a moderate heat, remove the oil which floats upon the product, and preserve the latter.

Dose, five drops, on sugar.

TEREBINTHINATED LOTION. (NIEMANN)

R. Spirit of Turpentine, of each,
Alcohol, four ounces.
Camphor, }

Dissolve.—HEBERDEN and NIEMANN
VOL. II.

extol it in the morbus pedicularis of old persons.

VULNERARY BALSAM.

Liquor vulnerarius; Essentia vulneraria Stahlii. (FULD.; WIRTEM.; NIEMANN.; SPIELMANN)

R. Spirit of Turpentine, one part.
Tincture of Myrrh, two parts.
Amber, four parts.

Digest in a warm place.

It used to be employed in injections to fistulous ulcers, and as an application to wounds and ulcers of non-vascular parts.—It might properly be substituted for the compound tincture of benzoin.

FIORAVENTI'S BALSAM.

Balsamum Fioraventi. (BATAV.; GALL.; FORM. HÔP. FR.; BRUGNATELLI; SPIELMANN.; VAN MONS)

R. Cloves,	} of each, one part.
Nutmegs,	
Cinnamon,	
Ginger,	
Myrrh,	
Galbanum,	} of each, two parts.
Liquid Storax,	
Bay Berries,	three parts.
Venice Turpentine,	four parts.
Alcohol (10 degrees),	two hundred
Water,	and sixteen
	parts.

Distil one hundred and eight parts, and preserve the product. (BATAV.)

R. Fresh Bay Berries, four ounces.
Elemi, of each, one ounce.
Tacamahac, }
Galbanum, }
Olibanum, } of each,
Myrrh, } three ounces.
Gum Hedera, }
Liquid Storax, two ounces.
Aloe Wood, three ounces.
Lesser Galanga, }
Cloves, } of each,
Cinnamon, } one ounce.
Nutmegs, }
Zedoary, }
Ginger, }
Dittany of Crete, }
Socotrine Aloes, }
Amber, }
Alcohol, ninety-six ounces.

Digest for several days, and add

Turpentine, sixteen ounces.

Distil.—The spirit which passes over is called *Balsamum Fioraventi spirituosum*. The residue, distilled slowly in a retort, yields a yellow oil, called *Balsamum Fioraventi oleosum*. By increasing the heat,

a black and empyreumatic mass is obtained, called *Balsamum Fioraventi nigrum*. (SPIELMANN)

R. Turpentine, five hundred and sixteen parts.

Elemi,	}	of each,	ninety-six parts.
Tacamahac,			
Amber,			
Galbanum,			
Myrrh,	}	of each,	thirty-two parts.
Liquid Storax,			
Leaves of Dittany,			
Aloes,			
Fresh Bay Berries,	}	of each,	forty-eight parts.
one hundred and twenty-eight parts.			
Galanga,			
Zedoary,			
Ginger,	}	of each,	thirty thousand parts.
Cinnamon,			
Cloves,			
Nutmegs,			
Alcohol (22 degrees),	}	of each,	thirty thousand parts.

Macerate for six days; then distil two thousand five hundred parts in a water-bath. (GALL.; FORM. HÔP. FR.)

R. Turpentine, eight parts.
 Bay Berries, } of each, three parts.
 Myrrh, }
 Liquid Storax, two parts.
 Nutmegs, }
 Cloves, } of each, one part.
 Ginger, }
 Cinnamon, }
 Proof Spirit, }
 two hundred and sixteen parts.

Distil about one fourth. (BRUGNATELLI; NIEMANN; VAN MONS)

This was formerly a very celebrated stimulant, and was classed among the analeptics, nervines, anti-nephritics, diuretics, alexipharmics, anti-hysterics, vulneraries, and carminatives.

EAU D'ANHALT.

Aqua Anhaltina, Spiritus Anhaltinus.
 (BRUNS; HANN.; PALAT.; SARD.; WIRTEM.; CADET DE GASSICOURT;
 NIEMANN; SPIELMANN; VAN MONS)

R. Turpentine, half a pound.
 Male Olibanum, one ounce.
 Aloe Wood, three ounces.
 Mastic, }
 Cloves, } of each, six drams.
 Nutmegs, }
 Cubebs, }
 Cinnamon, }
 Saffron, two drams and a half.
 Seeds of Fennel, } of each,
 Bay Berries, } half an ounce.
 Alcohol, five pounds.

Digest for three days, and distil in a water-bath. (PALAT.)

SPIELMANN directs the supernatant oil to be separated from the product, and allows a bag containing fifteen grains of musk to be suspended in it.

R. Leaves of Rosemary, one ounce.
 Herb of Rue, } of each,
 Seeds of Fennel, } half an ounce.
 Bay Berries, }
 Cloves, }
 Cubebs, } of each, six drams.
 Nutmegs, }
 Cinnamon, }
 Mastic, two ounces.
 Venice Turpentine, half a pound.
 Rectified Spirit, five pounds.

Digest for six days, and add

Spring Water, two pounds.

Distil five pounds in a water-bath. (BRUNS.)

R. Root of Lesser Galanga, } of each,
 Flowers of Rosemary, } six drams.
 Seeds of Fennel, } of each,
 Bay Berries, } one ounce.
 Cloves, }
 Cubebs, } of each, six drams.
 Nutmegs, }
 Cinnamon, }
 Mastic, }
 Olibanum, } one ounce.
 Venice Turpentine, half a pound.
 Rectified Spirit, six pounds.

Digest for about six days, and add

Water, one pound.

Distil five pounds in a water-bath. (HANN.; SARD.; WIRTEM.)

R. Bay Berries, } of each,
 Seeds of Fennel, } half an ounce.
 Cinnamon, }
 Cubebs, } of each, six drams.
 Lesser Galanga, }
 Cloves, }
 Mastic, }
 Nutmegs, }
 Rosemary, }
 Olibanum, } four ounces.
 Proof Spirit, ninety-six ounces.
 Water, }
 two hundred and eighty-eight ounces.

Distil eighty ounces of spirit, and redistil this product with

Turpentine, four ounces.
 Water, one hundred and sixty ounces;
 to obtain sixty-six ounces of spirit.

We omit the formula of CADET DE GASSICOURT, as it is erroneous.

This liquor, which scarcely differs from FIORAVENTI'S balsam in properties as well as in composition, is employed in the same cases as that is, and is also applied exter-

nally to the abdomen to arrest vomiting and diarrhoea, and to the limbs in paralysis.—
Dose, sixty drops or more.

SCHAUER'S BALSAM.

Balsamus Schauerianus. (BRUNS.; NIEMANN; SPIELMANN)

- R. Venice Turpentine, one ounce and a half.
 Galbanum, nine drams.
 Root of Comfrey, of each,
 ——— Fraxinella, six drams.
 ——— Master-wort, of each,
 Mastic, half an ounce.
 Root of Sweet Flag, }
 ——— Galanga, }
 Storax, of each,
 Olibanum, three drams.
 Ginger, }
 Cinnamon, }
 Benzoin, }
 Nutmegs, }
 Cubebs, }
 Cloves, of each,
 White Sanders, two drams.
 Lesser Cardamoms, }
 Camphor, }
 Aloe Wood, }
 Root of Avens, of each,
 ——— Elecampane, one dram.
 Hepatic Aloes, }
 Myrrh, half a dram.
 Root of Angelica, fifteen grains.
 Concentrated Alcohol, eight pounds.

Digest, and distil by the heat of a water-bath, to dryness.

This also, like FIORENTI'S balsam, is stimulant, resolvent, and vulnerary. It used formerly to be extolled as an application to contusions and ecchymoses.

BALSAMUM VITÆ ALBUM; SPIRITUS BAL-SAMICUS. (WIRTEM.)

- R. Olibanum, }
 Myrrh, of each, three ounces.
 Galbanum, }
 Aloe Wood, }
 Cloves, }
 Cinnamon, }
 Nutmegs, of each, one ounce.
 Galanga, }
 Zedoary, }
 Ginger, }
 Bay Berries, four ounces.
 Venice Turpentine, twelve ounces.
 Alcohol, twelve pounds.

Digest, and distil in a water-bath.

This balsam, also resembling that of FIORENTI, is also reputed stomachic, diuretic, carminative, and nervine, either externally, or internally, when the dose is from forty to sixty drops.

ANTI-GONORRHOIC LIQUOR. (WIRTEM.)

- R. Venice Turpentine, ten ounces.
 Root of Florentine Iris, two ounces.
 Dried Herb of Mint, three ounces.
 ——— Dittany of Crete, one ounce.
 ——— Rue, of each,
 ——— Seeds of Agnus Castus, two ounces
 ——— White Wine, and a half.
ten pounds.

Distil one half in a water-bath.

HOFFMANN prescribed this in gonorrhœa and leucorrhœa, in the dose of four spoonfuls daily, at twice.

ASTRINGENT AROMATIC LIQUOR. (BORIES)

- R. Flowers of White Nettle, one pound.
 ——— Pale Roses, two ounces.
 Seeds of Nettle, one ounce.
 Tops of Rosemary, eight ounces.
 Leaves of Plantain, two ounces.
 Acorns, twenty-four.
 Root of Bistort, two ounces.
 White Wine, eight pounds.

Infuse for twenty-four hours, and add

Venice Turpentine, four ounces.

Distil in a water-bath to dryness; then add to the product,

Subcarbonate of Potass, one ounce.

Rob of Elder, four ounces.

White Sugar, one pound.

Mix.—It is excitant, and is extolled in leucorrhœa.—Dose, two or three ounces, in the morning, fasting.

C.

Preparations which contain spirit of Turpentine suspended in watery liquids.

TEREBINTHINATED LAVEMENT. (ELLIS; NIEMANN; RATIER)

- R. Spirit of Turpentine, one ounce.
 Triturate it with yolk of egg, and add gradually,
 Spring Water, eight ounces.
 Mix. (NIEMANN)
 It is extolled by CROSS against ascarides.
 R. Spirit of Turpentine, one ounce.
 Yolk of Egg, one.
 Decoction of Poppy Heads, eight ounces.

Mix.—It is recommended by RECAMIER in lumbar neuralgia. (RATIER)

- R. Spirit of Turpentine, one ounce and a half.
 Yolks of Eggs, two.
 Solution of Gum Arabic, one pound.

Mix. (ELLIS)

DIURETIC LINIMENT. (HANN.; CADET DE GASSICOURT; NIEMANN)

- R. Spirit of Turpentine, half an ounce.
Yolk of Egg, one.
Peppermint Water, six ounces.

Mix. (HANN.; NIEMANN)

CADET DE GASSICOURT prescribes one ounce of spirit and three ounces of peppermint water.

To be rubbed upon the inguinal region in spasmodic ischury.

DIURETIC JULEP. (BORIES; PIERQUIN)

- R. Spirit of Turpentine, fifteen drops.
Woodlice, in powder, half a dram.
Juice of Asparagus, one ounce.
Decoction of Asparagus, } of each,
Water of Rupture-wort, } three ounces
Syrup of Maidenhair, six drams.

Mix.—To be taken at twice, morning and evening; in ischury, rheumatism, and gout.

TEREBINTHINATED LOBOCH. (RATIER)

- R. Spirit of Turpentine, three drams.
Yolk of Egg, one.
Syrup of Mint, two ounces.
Orange Flowers, } of each,
Ether, } one ounce.
Tincture of Cinnamon, half a dram.

Mix.—RECAMIER employs it in neuralgia. —Dose, three spoonfuls daily.—A dram of laudanum may be added to it if it excites vomiting. (RATIER)

POTION AGAINST TÆNIÆ. (CADET DE GASSICOURT)

- R. Spirit of Turpentine, three ounces.
Honey, six drams.
Mint Water, three drams.

Mix.—To be taken at thrice in the course of the day.

TEREBINTHINATED EMULSION. (NIEMANN; VAN MONS)

- R. Spirit of Turpentine, one dram.
Triturate it with yolk of egg or with two drams of mucilage of gum arabic, and add gradually,

Water, six or eight ounces.

Mix. (NIEMANN)

- R. Spirit of Turpentine, half an ounce.
Gum Arabic, two drams.
Water, a sufficient quantity.

Make an emulsion, and add

Honey, two ounces.

Water, a sufficient quantity,
to make the whole six ounces. (VAN MONS)

Dose, a spoonful, several times a day.

COSMETIC LIQUOR. (PALAT.)

- R. Whites of boiled Eggs, two pounds.
Venice Turpentine, half a pound.
Honey, eight ounces.
Storax, one ounce.
Benzoin, two drams.
Rose Water, ten ounces.

Distil in a water-bath, to dryness.

DIURETIC POTION. (ELLIS)

- R. Spirit of Turpentine, one hundred drops.
Gum Arabic, } of each, one dram.
White Sugar, }
Mint Water, four ounces.

Mix.—Dose, four or five spoonfuls, daily.

STIMULANT POTION. (ELLIS)

- R. Spirit of Turpentine, one hundred and twenty drops.
Gum Arabic, } of each, two drams.
Sugar, }
Tincture of Opium, sixty drops.
Compound Tincture of Lavender, two drams.
Mint, five ounces.

Mix.—Dose, a spoonful every two hours.

D.

Preparations which contain spirit of Turpentine mixed with saccharine substances.

OLEO-SACCHARUM OF TURPENTINE.

Pulvis terebinthinatus. (AUGUSTIN; NIEMANN)

- R. Spirit of Turpentine, one scruple.
Pyrotartaric Acid, ten grains.
White Sugar, a sufficient quantity,
to absorb the whole. Triturate, and make a powder.

It is extolled by GÖZE against tæniæ.—A powder to be taken thrice a day.—Catharsis is to be then effected by gamboge.

TEREBINTHINATE ELECTUARY.

Electuarium terebinthinatum. (GENEV.; RATIER)

- R. Spirit of Turpentine, two drams.
Gum Arabic, in powder, one ounce and a half.
Sugar, half an ounce.
Syrup of Orange Flowers, one ounce.

Mix.—It is employed by RECAMIER in neuralgia.—Dose, one third, daily, at thrice, between two wafers. (RATIER)

- R. Spirit of Turpentine, half an ounce.
Honey, one ounce.
Liquorice, a sufficient quantity.

Mix, (GENEV.)

FEBRIFUGE MARMALADE.

Conditum Febrifugum. (VAN MONS)

- R. Spirit of Turpentine, two drams.
 Pure Turpentine, one ounce and a half.
 Triturate together with yolk of egg, and add

Manna, two ounces.

Mix.—It is recommended against tæniæ.
 —Dose, a teaspoonful, morning and evening.

ANTI-ARTHRITIC BOLUS. (SAUNDERS)

- R. Spirit of Turpentine, twenty-four drops.
 Honey, thirty-six grains.
 Mix.

TEREBINTHINATED HONEY.

Linctus Olei Terebinthinæ. (ELLIS; NIEMANN; RATIER; SWEDIAUR)

- R. Spirit of Turpentine, half an ounce.
 Triturate with yolk of egg, and add
 Honey, three ounces.
 Mix. (SWEDIAUR)
 R. Spirit of Turpentine, two drams.
 Honey, one ounce.
 Mix. (ELLIS; NIEMANN)

HOME has proposed it in lumbago, and RECAMIER has since recommended it in all neuralgiæ: he substitutes honey of roses for the simple honey, varying the quantity of it from one ounce to four ounces.—Dose, three spoonfuls in the course of the day.

E.

Preparations which contain spirit of turpentine mixed with fat bodies.

VERMIFUGE POTION. (BRERA)

- R. Spirit of Turpentine, two drams.
 Walnut Oil, one ounce and a half.
 Mix.—For one dose.

- R. Spirit of Turpentine, five drams.
 BRERA elsewhere prescribes five drams of spirit and three ounces of oil; for one dose.

ANTHELMINTIC OIL. (NIEMANN)

- R. Spirit of Turpentine, four parts.
 Oil of Hartshorn, one part.
 Distil two thirds of the mixture.

NEUROHR says he has given with success from ten to a hundred drops of the mixture against tæniæ.

- R. Spirit of Turpentine, three parts.
 Oil of Hartshorn, one part.
 Distil three fourths.

CHABERT administered the latter product in the dose of from one to two tea spoonfuls morning and evening, or two tea spoonfuls of it in mucilaginous decoction for a lavement.

BREMSEY prescribes two tea spoonfuls morning and evening in a little water; this is certainly too strong a dose, particularly as a commencing one. ALDERS has found it a very efficacious remedy.—CHABERT only joined the spirit of turpentine to the oil of hartshorn in order to conceal the taste of the latter; however, it is the principal ingredient of the remedy, and GRUNDLE has committed a great mistake in saying that it was useless.

LINIMENT OF TURPENTINE. (LOND.; DUBL.; AMER.; COXE; RATIER)

Linimentum Terebinthinæ. (LOND.; DUBL.)

- R. Spirit of Turpentine, half a pint.
 Melt and add to it,
 Resin Cerate, one pound.
 Mix.

AMER.; and COXE give the same formula. It forms a useful liniment in rheumatism.

- R. Spirit of Turpentine, one ounce.
 Oil of Chamomile, two ounces.
 SYDENHAM'S Laudanum, one dram.

Mix.—It is employed by RECAMIER in neuralgiæ. (RATIER)

OLEUM DE TRIBUS. (VAN MONS)

- R. Spirit of Turpentine, } of each,
 Oil of Lavender, }
 — Brick, } an equal part.

Mix.

F.

Preparations which contain spirit of turpentine combined or mixed with acids.

NARCOTIC BALSAM. (VAN MONS)

- R. Spirit of Turpentine, } of each,
 Sulphuric Acid, } an equal part.
 Mix together gradually. When cold, heat the mass with hot water, and wash it till all the free acid is removed from it.

ACHARD ascertained that the spirit of turpentine would combine with sulphuric acid. He calls the substance which results acid soap (*savon acide*); it is completely soluble in alcohol and in water. This soap forms the basis of the following preparation:

ASTRINGENT BALSAM.

Balsamum adstringens seu ischæmaticum.
(BRUNS.; FULD.; AUGUSTIN.; CADET DE
GASSICOURT.; PIERQUIN.; SPIELMANN.;
VAN MONS.)

R. Sulphuric Acid, ten drams.

Instil into it slowly,

Spirit of Turpentine, half an ounce.

After the cessation of the effervescence,
and when the liquor has become cold, add
very carefully,

Alcohol, four ounces.

Preserve. (BRUNS.)

PIERQUIN.; and SPIELMANN prescribe
half an ounce each, of acid and of spirit,
and three ounces of alcohol.—CADET DE
GASSICOURT—half an ounce of acid, four
ounces of spirit, and three of alcohol.

R. Clear Turpentine, two parts.

Concentrated Sulphuric Acid,
two parts and a half.

Mix gradually in a glass mortar placed
in a warm water-bath. When cold, add at
several times,

Alcohol, eight parts.

Set apart for several days, and preserve
the decanted liquor. (VAN MONS.)

FULD.; and AUGUSTIN prescribe three
parts of turpentine, five of sulphuric acid,
and twenty-four of rectified alcohol.

It is excitant and astringent, and has been
recommended in hæmoptysis.—Dose, one
or two scruples, in an appropriate liquid.

ARTIFICIAL CAMPHOR. (VAN MONS)

R. Muriate of Soda, in powder,
five parts and a half.

Put it into a matrass, and pour upon it,

Concentrated Sulphuric Acid,
five parts.

Pass, through a tube, the gas disengaged
from the mixture, into a flask containing,

Spirit of Turpentine, ten parts.

Leave it to digest without heat for several
hours; then heat slightly, applying linen
cloths wetted continually with cold water
round the flask. When no more gas passes
over, let the apparatus cool; then pour off
the liquor, let the concrete part drip, press
it in bibulous paper, and preserve it in a
closely stopped bottle.

SOLUTION OF ARTIFICIAL CAMPHOR.
(VAN MONS)

R. Artificial Camphor, washed, one part.
Rectified Alcohol, six parts.

Dissolve without heat.

POTT'S ARTHRITIC LIQUOR.

Sapo seu Liquor seu Spiritus arthriticus
Pottii, Linimentum resolvens, Spiritus
Salis terebinthinatus. (LIPP.; SAX.;
HERBIP.; CADET DE GASSICOURT.; NIE-
MANN.; SWEDIAUR)

R. Muriate of Soda, in } of each,
powder, } two ounces.
Spirit of Turpentine, }
Add gradually and carefully,
Concentrated Sulphuric Acid,
one ounce.

Distil the mixture in a sand-bath. (LIPP.;
SAX.; HERBIP.; NIEMANN)

CADET DE GASSICOURT.; and SWEDIAUR
prescribe one part of hydrochloric acid and
two parts of spirit.

The hydrochloric acid acts upon the spirit
of turpentine in the same manner as the
sulphuric. It forms two different com-
pounds with it: one, containing a smaller
proportion of acid, is the artificial camphor,
indicated above; the other, containing a
greater proportion of acid, is liquid, accord-
ing to the observations of HOUTON—LA-
BILLARDIERE: it is the latter which forms
the basis of the arthritic liquor.

ANTI-RHEUMATIC LINIMENT.

Lotus ad Rheumatismum. (SPIELMANN)

R. Spirit of Turpentine, } of each,
Hydrochloric Acid, } one ounce.
Spirit of Camphor, } six ounces.

Mix.

RICHARD'S RESOLVENT. (VAN MONS)

R. Spirit of Turpentine, } of each,
Concentrated Hydro- } one part.
chloric Acid, }

Put the mixture into a retort, and agi-
tate it frequently. At the end of one or two
days, add

Alcohol, eight parts.

Distil eight parts, and dissolve in the
product,

Camphor, one half part.

BALSAM AGAINST CHILBLAINS. (BORIES;
RATIER)

R. FIORAVENTI's Balsam, four ounces.
Hydrochloric Acid, thirty-two drops.

Mix.

RATIER prescribes two ounces of olive
oil, one dram of balsam of Peru, two drams
each, of spermaceti, of white wax, and of
hydrochloric acid, and six drams of pure
water.

R. Olive Oil, five parts.

Spirit of Turpentine, two parts.
Sulphuric Acid, one half part.

Mix. (CADET DE GASSICOURT)

HENRI prescribes one hundred and sixty parts of oil, sixty-four of spirit, and sixteen of acid.

To be rubbed upon the parts threatened with chilblains, morning and evening.

G.

Preparations which contain spirit of turpentine combined with alkalis.

STARKEY'S SOAP.

Sapo Starkeyanus seu tartareus seu chemicus pro casu externo seu Potassæ terebinthinaceus seu tartareus Starkeyi, Corrector Starkeyi seu Opii. (BRUNS.; HISP.; GALL.; FERR.; FULD.; HASS.; LIPP.; SARD.; WIRTEM.; HERBIP.; AUGUSTIN.; PIERQUIN.; SPIELMANN.; SWEDIAUR.)

R. Subcarbonate of Potass, any quantity.

Calcine it strongly, and throw the mass while hot into a heated mortar, reduce it to powder, and pour upon it a little Venice turpentine, triturating well: continue to add turpentine by portions of about a dram, till it ceases to be absorbed; then dissolve the mass in concentrated alcohol, set aside and filter the solution, and finally evaporate it slowly. (FULD.; LIPP.)

BRUNS.; SARD.; and WIRTEM. direct the subcarbonate to be calcined for an hour or two, and then to be pulverized; a sufficient quantity of spirit of turpentine, to be then added drop by drop, as will preserve it from the contact of air: the vessel to be then put into a cellar, and after some time a little warm spirit of turpentine added: this is to be continued till the alkali has absorbed about thrice its weight of the spirit.

R. Dried Subcarbonate of Potass, } of each,
Spirit of Turpentine, } an equal part.
Venice Turpentine,

Triturate the subcarbonate in a marble mortar with a glass pestle, first adding the spirit and then the turpentine. When the mixture has acquired the proper consistence let it be levigated. (GALL.)

R. Common Turpentine, one ounce.

Liquify in an earthen vessel, and add

Solution of Potass, half an ounce.

Stir from time to time till the mixture has acquired the consistence of soap. (HISP.)

PIERQUIN prescribes three ounces of dried potass, two of water, and four of spirit of turpentine.

R. Caustic Potass, one ounce.

Fuse it over a very slow fire, and add

Spirit of Turpentine, three ounces.

Stir till perfectly incorporated. (HASS.; HERBIP.; SPIELMANN.)

FERR.; and SWEDIAUR prescribe ten parts of potass, and eight of spirit.

R. Turpentine, one ounce.
Spirit of Turpentine, half an ounce.
Oil of Almonds, } of each,
Solution of Potass, } one ounce.

Mix. (AUGUSTIN.)

This preparation derives its name from its supposed property of correcting the noxious qualities of opium, of hellebore, and of other substances of the same kind. It is balsamic, stimulant, deobstruent, diuretic, and anti-spasmodic. It has been principally recommended in gonorrhœa and dropsies.—Dose, from eight to twelve grains, in a watery vehicle or in pills.

H.

Preparations which contain spirit of turpentine combined with soaps.

SAPONACEOUS BALSAM OF TURPENTINE.

Balsamum saponaceum terebinthinatum seu Vitæ externum, Linimentum stimulans, Sapo terebinthinatus, Spiritus Saponis compositus.

1. Without Alcohol.

R. Hard Venice Soap, sixteen parts.
Spirit of Turpentine, thirty-two parts.

Add to the Solution,

Solution of Subcarbonate of Potass, three parts.

Mix. (FULD.; HASS.; LIPP.; CAST. BORUS.; HERBIP.)

BRUNS. prescribes twenty-four parts of soap, twenty-four of spirit, and one part of subcarbonate.—DAN.; HANN.; BORUS.; SAX.; and NIEMANN—six each, of soap and of spirit, and one of subcarbonate.—NIEMANN, after THEDEN,—eighteen of soap, fifteen of spirit, and one of subcarbonate.—SPIELMANN—nine of soap, eighteen of spirit, and one of subcarbonate.—HERBIP.; and SWEDIAUR—four of soap, eight of spirit, and one of subcarbonate.

R. Soap, scraped, two parts.
Spirit of Turpentine, three parts.

Dissolve with trituration, and add

Calcined Caustic Potass, one part and half,

dissolved in

Water, one half part.

Mix. (VAN MONS)

- R. White Soap, four parts.
 Camphor, one part.
 Triturate together, adding gradually,
 Spirit of Turpentine, sixteen parts.
 Preserve the solution.

2. With Alcohol.

- R. Marrow Soap, four parts.
 Turpentine, one part.
 Tincture of Soda, four parts.
 Dissolve in a close vessel placed in a water-bath, and strain. (VAN MONS)

- R. White Soap, } of each,
 Spirit of Turpentine, } one part.

Triturate together, and add

Alcohol, eight parts.

Digest in a warm water-bath. Set the solution aside for several days, and filter. (HASS.; LIPP.; WIRTEM.; PIDERIT; VAN MONS)

The solution of soap in spirit of turpentine, prepared separately, is designated in HASS.; and PIDERIT, under the name of *Corpus pro Balsamo Saponis*.

- R. Camphor, in powder, one part.
 Rectified Alcohol, sixteen parts.

Add to the solution

White Soap, scraped, four parts.

Digest over a slow fire, and add

Spirit of Turpentine, one part.

Mix. (MIL. COPEN.)

- R. Subcarbonate of Potass, one ounce and a half.
 Spring Water, twenty-four ounces.

Dissolve, and add

Spirit of Turpentine, three ounces.

Venice Soap, of each,

Spirit of Juniper, } two ounces.

Oil of Culilawan, three ounces.

Mix. (SPIELMANN)

- R. Soap, } of each,
 Spirit of Turpentine, } half a pound.

Dissolve, and add

Spirit of Wild Thyme, four pounds.

Solution of Ammonia, from two to eight ounces.

Mix. (SWEDIAUR)

CADET DE GASSICOURT prescribes one ounce of soap, half a pound of spirit of turpentine, five ounces of spirit of wild thyme, and from two drams to one ounce of solution of ammonia.

- R. Marrow Soap, one part.
 Spirit of Ammonia, four parts.
 ————Turpentine, eight parts.
 ————Wild Thyme, sixty-four parts.

Filter the solution. (VAN MONS)

Several of these formulæ resemble those for opodeldoch. The saponaceous balsam of turpentine possesses nearly the same properties, and is applied to the same purposes as that is.—It is excitant, resolvent, and discutient.—It is applied externally, in luxations and to disperse certain tumours.

RESOLVENT FOMENTATION. (FULD.; HAMB.; LIPP.; SPIELMANN)

- R. Saponaceous Balsam of Turpentine, one part.
 Water, twenty-four parts.
 Dissolve.

BASVILLE'S BALSAM.

Balsamum Basvillianum. (AUGUSTIN; NIEMANN)

- R. Spirit of Turpentine, three ounces.
 Subcarbonate of Potass, one ounce and a half.
 Alicant Soap, scraped, two ounces.
 Distilled Water, one ounce and a half.
 Spirit of Juniper, two ounces.

Mix.—This preparation scarcely differs from the saponaceous balsam of turpentine. It seems to be framed from SPIELMANN'S formula. It differs still less from the following:

PEREBOOM'S NERVINE LIQUOR. (HANN.; NIEMANN; VAN MONS)

- R. Subcarbonate of Potass, three parts.
 Alicant Soap, four parts.
 Triturate together, moistening with
 Warm Water, two parts.
 Incorporate gradually
 Spirit of Turpentine, six parts.
 Cajeput Oil, one part.

Then add, at several times,

Warm Water, forty-six parts.

Filter at the end of several days. (VAN MONS)

HANN.; and NIEMANN prescribe three parts of subcarbonate, four of soap, forty of water, six of spirit, one of oil of cajeput, and four of spirit of juniper.

It is excitant, and is employed externally in paralysis.

ANTI-ARTHRITIC LINIMENT. (CADET DE GASSICOURT)

- R. Camphor, one scruple.
 Spirit of Turpentine, two drams.
 Add to the solution,
 Soft Soap, one ounce.
 Nervine Ointment, half an ounce.
 Cumin, in powder, two drams.

Succinated Salt of Hartshorn,
fifteen grains.

Mix.—It is recommended by HOME in long continued rheumatic affections, particularly sciatica.

III. Preparations which contain the resinous part of turpentine exclusively or principally.

The resinous part of turpentine is presented under several different forms, depending generally on its greater or less degree of purity. We shall mention those kinds only which enter into the materia medica of the various pharmacopœias.

1. *Resina alba, Pix Burgundica*.—Burgundy Pitch.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN.

It is obtained by melting over a slow fire the *Barras* or *Gallipot*, that is, the common turpentine solidified at the surface of the incisions, and filtering the melted mass through straw.

2. *Resina communis*.—Common Resin.

It is prepared by melting the preceding, or by working melted colophony with one fourth of galipot in water, in order to destroy its transparency and give it a dull yellow colour. It scarcely differs from the preceding with which it is confounded in most pharmacopœias.

3. *Resina flava, Colophonium, Pix Græca*.—Yellow Resin, Colophony.

AMST.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

This is the residuum of the common turpentine after the oil or spirit has been distilled from it; it is then re-melted and set aside to solidify.

4. *Resina flava, Terebinthina cocta*.—Yellow Resin.

DUBL.; AUSTR.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; GENEV.; HANN.; VOL. II.

HASS.; OLDEN.; PALAT.; POL.; BORUS.; SAX.; WIRTEM.; HERBIP.; BRUGNATELLI; VAN MONS.

This only differs from colophony in its containing water, and from common resin in its containing still more of it than that. It is obtained either by taking what remains in the alembic after the distillation of turpentine with water in obtaining spirit of turpentine, or by boiling one part of turpentine in three parts of water till the mass when poured into cold water assumes a ductile consistence.

GALL. directs it to be formed into pills of six grains each, and preserved in water.

5. *Pix solida seu atra seu navalis seu vegetabilis, Palampissa*.—Pitch.

LOND.; DUBL.; EDINB.; AMST.; BATAV.; BAVAR.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; HANN.; HASS.; LIPP.; LUSIT.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SPIELMANN.

It is obtained by the descending combustion of the straw which has served to filter common turpentine.

It is resin partly altered by heat, and mixed with carbon. It is somewhat similar to tar, but much less impure, and does not appear to contain any acetic acid.

The *terebinthina cocta* is administered internally as balsamic and diuretic. It is much prescribed in obstinate gonorrhœa, gout, and chronic rheumatism, in the dose of half a dram, several times a day. The other varieties are more particularly reserved for external use.

A.

Preparations into which the terebinthina cocta enters.

ASTRINGENT MARMALADE.

Conditum corroborans. (VAN MONS)

R. Terebinthina cocta, one ounce.
Copaiba, half an ounce.

Incorporate in a heated mortar; then add

Mucilage of Gum Arabic, one ounce.
Conserve of Red Roses, four ounces.

Mix.—Dose, two or three drams, thrice a day, in inveterate gonorrhœa.

TEREBINTHINATE PILLS.

Pilulæ e Terebinthina seu diuretice corroborantes seu stimulantés ad gonorrhœam.

4 R

(CAST. BORUS.; SARD.; WIRTEM.; PH. SYPH.; AUGUSTIN; NIEMANN; SPIELMANN; SWEDIAUR)

R. Terebinthina cocta, two drams.
Powdered Rhubarb, one dram.

Make thirty pills. (PH. SYPH.)

Dose, four or six, morning and evening.

R. Terebinthina cocta, two ounces.
Powdered Rhubarb, one ounce.
Syrup of Tolu, a sufficient quantity.

Make pills of four grains each. (SWE-
DIAUR)

To be given in the same dose as the pre-
ceding.

R. Terebinthina cocta, } of each,
Soap, } one dram.
Powdered Rhubarb, half an ounce.

Make pills of two grains each. (CAST.
BORUS.; NIEMANN)

Dose, from ten to fifteen, morning and
evening.

AUGUSTIN prescribes half a dram of tere-
binthina cocta, half a dram of soap, and one
dram of rhubarb.

R. Terebinthina cocta, one ounce.
Rhubarb, two drams.
Amber, } of each,
Mastic, } one dram.
Liquorice, }
Cuttle Fish Bone,
Venice Turpentine, a sufficient quantity.

Make a pilular mass. (SARD.)

R. Terebinthina cocta, one ounce and a half.

Amber, }
Mastic, } of each,
Olibanum, } two drams.
Extract of Liquorice,
Rhubarb,
Cuttle Fish Bone,
Armenian Bole, }

Make a pilular mass. (WIRTEM.)

R. Terebinthina cocta, one ounce and a half.
Armenian Bole, } of each,
Cuttle Fish Bone, } half an ounce.
Amber, } of each, two drams.
Mastic, }
Rhubarb, } one dram.
Syrup of Tolu, a sufficient quantity.

Make a pilular mass. (SPIELMANN)

The three last formulæ are properly
abandoned.

Dose, one scruple, thrice a day, in go-
norrhœa.

ASTRINGENT PILLS. (SAINTE-MARIE)

R. Terebinthina cocta, } of each,
Dragon's Blood, } two drams
Clear Turpentine, a sufficient quantity

Make pills of four grains each.

Dose, three, daily, in gonorrhœa.

R. Terebinthina cocta, four drams.
Amber, two drams.
Cuttle Fish Bone, one dram.
Guaiac Resin, four scruples.
Copaiba, a sufficient quantity.

Make one hundred and fifty pills.

Dose, twelve daily at thrice, in leucor-
rhœa and gonorrhœa.

BALSAMIC PILLS.

Pilulæ balsamicæ Junkeri. (AUGUSTIN;
NIEMANN; SPIELMANN)

R. Mastic, one part.
Myrrh, } of each,
Terebinthina cocta, } one part and
Amber, } a half.
Jalap, three parts.
Extract of Fumitory, } of each,
Blessed Thistle, } two
Wormwood, } parts.
Scurvy-grass, }

Make a pilular mass. (NIEMANN)

These pills, invented by JUNKER, were
formerly considered emmenagogue.

R. Terebinthina cocta, half an ounce.
Soap, one ounce.
Resin of Jalap, } of each,
Aqueous Extract of Aloes, } three
Extract of Lesser Centaury, } drams.
Gentian, }

Make a pilular mass. (SPIELMANN)

Dose, to eighty grains.

OINTMENT AGAINST CHILBLAINS. (SWE-
DIAUR)

R. Terebinthina cocta, } of each,
Cocoa-nut Oil, } three ounces.

Melt together over a slow fire, and add
Balsam of Peru, half a dram.

Mix.

OLEO-RESINOUS CERATE. (VAN MONS)

R. White Wax, } of each,
Terebinthina cocta, } one part.
Olive Oil, two parts.

Melt over a slow fire; then strain, and
stir till perfectly cold.

It is a useful application to slight exco-
riations.

ADHESIVE SPARADRAP. (VAN MONS)

- R. Terebinthina cocta, } of each,
 White Wax, } an equal part.

Melt together, and spread the mass upon linen or paper.

- R. Terebinthina cocta, } one part.
 Simple Plaster, } three parts.

Melt, and spread in the same manner.

FOURCROY'S BALSAM. (CADET DE GASSICOURT)

- R. Olive-Oil, four pounds.
 Powdered Root of Angelica, } of each,
 Scorzo- } two
 nera, } ounces.
 St. John's-wort, }
 Ivy Berries, }

Boil over a very slow fire; then macerate for one night; set it over a slow fire on the following morning, and when the fluid is about to boil, add

- Theriac, } of each,
 Saffron, } two drams.
 Extract of Juniper, }
 Aloes, } one dram.

Boil, and strain; then, having set it aside for a few hours, put it again over the fire, and add

- Turpentine, ten ounces.

Heat till a terebinthinaceous odour is no longer exhaled; then remove it from the fire, and add

- Powdered Olibanum, } of each,
 Storax, } one dram and
 Benzoin, } a half.

Mix till perfectly cold; then set aside for two or three days, strain through a napkin, and preserve the liquid in a bottle.

It is recommended in chaps of the skin, ulcers, chilblains, rheumatism, and sprains, rubbed or simply applied.

B.

Preparations into which resin properly so called enters.

RESINOUS FUMIGATION. (SAINTE-MARIE)

- R. Common Resin, } of each,
 Yellow Wax, } an equal part.

Put them into an earthen vessel placed in a chafing dish, or over a spirit lamp.

This fumigation has been recommended in pulmonary consumption. It has seemed useful in certain chronic catarrhs accompanied by abundant expectoration.

OINTMENT AGAINST TINEA. (CADET DE GASSICOURT; HENRI)

- R. Wheat Flour, five parts.
 Vinegar, forty parts.

Make a paste; add to it, a melted and strained mixture of

- Burgundy Pitch, } of each, four parts.
 Pitch, }

Mix. (HENRI)

CADET DE GASSICOURT prescribes four parts each, of flour, and of each of the two pitches, and thirty parts of vinegar.

It is to be spread upon a piece of leather and applied to the part affected, and removed at the end of twelve hours.

CERATE OR OINTMENT OF RESIN.

Cera arborea, Ceratum citrinum seu Resinæ Pini seu resinosum, Emplastrum attrahens seu cereum seu Cera seu citrinum seu Cera terebinthinatum seu Picis, seu piceum seu adhasivum seu Picis navalis seu simplex seu sticticum seu roborans, Unguentum flavum seu Resinæ albæ seu resinosum. (LOND.; EDINB.; DUBL.; AMER.; AMST.; AUSTR.; BATAV.; BRUNS.; DAN.; MIL. COPEN.; FENN.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; HUFFELAND; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

Emplastrum Cerae. (LOND.)

- R. Yellow Wax, } of each,
 Prepared Suet, } three pounds.
 Yellow Resin, } one pound.

Melt together, and strain.

Emplastrum simplex. (EDINB.; COXE)

- R. Yellow Wax, three parts.
 Mutton Suet, } of each,
 White Resin, } two parts.

Melt slowly, and stir assiduously till the mixture cools.

VAN MONS prescribes two parts of resin, two of suet, and three of wax (*Emplastrum simplex*).—HERBIP.; COXE; and SWEDIAUR—one part of yellow resin, three parts of suet, and three of yellow wax (*Emplastrum commune seu Cera*).—SPIELMANN—two each, of resin and of wax, and one of suet (*Emplastrum attrahens*).—SWEDIAUR—sixteen of resin, one of suet, and six of wax; or, sixteen of resin, one of suet, and ten of wax (*Emplastrum adhasivum seu roborans*).

Unguentum Resinæ albæ. (DUBL.)

- R. Yellow Wax, one pound.
 White Resin, two pounds.

Prepared Lard, four pounds.
Make an ointment, and strain it
through a sieve while hot.

Unguentum Resinosum. (EDINB.;
AMER.; COXE)

R. Lard, eight parts.
Resin, five parts.
Yellow Wax, two parts.

Melt together slowly, and stir till
the mixture solidifies.

R. Lard, eight pounds.
Turmeric, in powder, two ounces.
Water, four ounces.

Boil slowly to the consumption of the
humidity. Add

Yellow Wax, } of each,
Resin, } half a pound.

Melt, and strain. (BORUS.; SAX.)—
This is the *Unguentum flavum*.

R. Resin, eighteen parts.
Burgundy Pitch, twenty-eight parts.
Yellow Wax, twenty parts.

Melt slowly together. (BATAV.)

VAN MONS prescribes four parts and
a half of black pitch, five of resin, and six of
yellow wax.—AMST.—twelve of resin, four-
teen of Burgundy pitch, and ten of wax.—
SAX.—nine of resin, fourteen of Burgundy
pitch, and ten of wax.—This is the *Emplas-
trum Picis seu piceum*.

Unguentum Picis Nigri (LOND.)

R. Pitch, } of each,
Yellow Resin, } nine ounces.
Olive Oil, one pint.

Melt together, and strain through
a linen cloth.

R. Resin, four parts.
Yellow Wax, one half part.
Turpentine, two parts.

Melt together. (VAN MONS). This is
the *Unguentum simplex seu Ceratum Resinæ*.

SWEDIAUR prescribes thirty-two parts of
Burgundy pitch, and one part each, of wax
and of turpentine (*Emplastrum piceum*).—
VAN MONS, in another place,—one and a
half of resin, three of yellow wax, and one
of turpentine (*Emplastrum citrinum*).

R. Resin, sixteen ounces.
Yellow Wax, ten ounces.
Turpentine, three ounces.

Add to the melted mass,

Powdered Turmeric, three drams.

Mix. (SPIELMANN).—It is the *Emplas-
trum citrinum*.

R. Resin, } of each, twelve parts.
Yellow Wax, }
Pitch, }
Goat's Suet, one part and a half.

Melt together, and strain. (DAN.; MIL.
COPEN.; HERBIP.)—This is the *Emplas-
trum piceum seu sticticum*.

LIPP. prescribes six parts of resin, six of
Burgundy pitch, four of yellow wax, and
two of stag's suet (*Emplastrum piceum seu
adhasivum*).

R. Resin, twenty-four parts.
Pitch, sixteen parts.
Yellow Wax, eight parts.
Turpentine, four parts.

Melt together. (FENN.; SUEC.)—This
is the *Emplastrum Picis*.

VAN MONS prescribes six parts of resin,
twelve of pitch, three of wax, and one of
turpentine (*Ceratum nigrum*).

R. Resin, eight parts.
Yellow Wax, four parts.
Suet, two parts.
Turpentine, one part.

Melt together, and strain. (FULD.;
HASS.; LIPP.; PIDERIT)—This is the *Em-
plastrum citrinum seu cereum seu attrahens,
Ceratum citrinum seu Resinæ Pini*.

AUSTR.; and BORUS. prescribe two parts
of resin, four of yellow wax, and one each,
of turpentine and of suet (*Ceratum seu Em-
plastrum citrinum*).—HANN.; and SAX.—
two of resin, four of wax, and one each, of
suet and of turpentine (*Ceratum seu Em-
plastrum citrinum, Cera arborea*).—VAN
MONS—four of resin, three of wax, and one
each of suet and of turpentine (*Ceratum
adhasivum*)—HUFELAND—six of resin, four
of wax, two of suet, and one of turpentine
(*Emplastrum citrinum*).—FERR.—sixteen
of wax, fourteen of resin, twelve of suet,
and three of turpentine (*Ceratum citrinum*).

R. Resin, twenty-four parts.
Yellow Wax, sixteen parts.
Mutton Suet, eight parts.
Turpentine, four parts.

Melt together, and add

Powdered Turmeric, one part.

Mix. (BRUNS.; OLDEN.; PALAT.; BO-
RUS.; ROSS.; WIRTEM.; HERBIP.)—This
is the *Emplastrum seu Ceratum citrinum*.

R. Pitch, } of each, two parts.
Resin, }
Mutton Suet, }
Turpentine, }
Yellow Wax, three parts.

Melt together. (NIEMANN)—This is the
*Emplastrum Picis navalis seu sticticum
Hamburgense*.

R. Yellow Wax, one pound.
Resin, half a pound.
Mutton Suet, four ounces.

Olive Oil, } of each, two ounces.
 Turpentine, }
 Powdered Turmeric, two drams.

Melt, and strain. (FENN.)—This is the *Emplastrum Cerae terebinthinatum seu citrinum*.

Some of these formulæ are very similar to those for the basilicon ointment, both with regard to the nature and number of the ingredients, and the respective proportions of each.

COMPOUND PLASTER OF PITCH.

Emplastrum Picis compositum. (LOND.; CADET DE GASSICOURT; COXE; VAN MONS)

R. Burgundy Pitch, two pounds.
 Resin of Spruce Fir, one pound.
 Yellow Resin, } of each,
 ——— Wax, } four ounces.
 Expressed Oil of Nutmeg, one ounce.

Olive Oil, } of each, two ounces.
 Water, }

To the pitch, resin, and wax, melted together, add the resin of spruce fir, then the oil of nutmeg, the olive oil, and the water. Mix the whole and evaporate to a proper consistence. (LOND.; COXE.)

VAN MONS prescribes four parts of Burgundy pitch, twelve of resin, twelve of black pitch, four of yellow wax, and one part of oil of nutmeg.—CADET DE GASSICOURT—one pound each, of lard and of Burgundy pitch, ten ounces of yellow wax, five ounces each, of resin and of pitch, and ten drams of olibanum.

GREEN DEFENSIVE PLASTER. (BRUNS.; WIRTEM.)

R. Yellow Wax, one pound.
 Resin, two ounces.
 Goat's Suet, six ounces.
 Turpentine, one ounce.
 Powdered Herb of Ladies-mantle, six ounces.

Mix.

To give a deeper colour to the plaster, a scruple of indigo may be added to each pound of it.

BRUNS. requires rue instead of ladies-mantle.

ANTI-RACHITIC PLASTER. (NIEMANN)

R. Resin, one ounce.
 Yellow Wax, half an ounce.

Empyreumatic Oil of Tartar, one dram.

Melt together.

This plaster is said to have been applied with advantage to the vertebral column of rickety children.

RESINOUS SPARADRAP. (VAN MONS)

R. Resin, } of each, one part.
 Turpentine, }
 White Wax, four parts.

Melt together, and spread the plaster upon linen or paper.

ANDRE' DE LA CROIX' PLASTER.

Emplastrum Andreæ a Cruce seu Resinarum glutinans. (HISP.; FORM. HÔP. FR.; CADET DE GASSICOURT; VAN MONS)

R. White Resin, eight parts.
 Elemi, two parts.
 Turpentine, } of each, one part.
 Oil of Bay, }

Melt and strain. (FORM. HÔP. FR.; CADET DE GASSICOURT; VAN MONS)

HISP. prescribes sixteen parts of white resin, five of elemi, two each, of turpentine and of olive oil, and one of white wax.

SUPPURATIVE PLASTER.

Emplastrum maturans. (SWEDIAUR)

R. Resin, half a pound.
 Gum Plaster, one pound.

Melt together.

BASILICON OINTMENT.

Unguentum basilicum seu tetrapharmacum seu Picis seu Picis empyreumaticum seu piceum compositum seu regium seu Terebinthinæ resinosum.

1. Yellow Basilicon; *Unguentum basilicum flavum seu pallidum, Unguentum Resinæ flavæ, Ceratum Resinæ, Unguentum emolliens seu aureum seu citrinum*. (LOND.; AMST.; DAN.; MIL. COPEN.; HISP.; FENN.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SARD.; SAX.; SUEC.; WIRTEM.; SWEDIAUR; VAN MONS)

Ceratum Resinæ. (LOND.)

R. Yellow Resin, } of each,
 ——— Wax, 142 } one pound.
 degrees,
 Olive Oil, one pint

Melt the resin and wax together over a slow fire, then add the oil,

and strain the mixture while hot through a linen cloth.

AMST. prescribes one part each, of resin and of wax, and three parts of oil.—LUSIT. prescribes an equal part of each ingredient.

R. Palm Oil, } of each, two pounds.
 Linseed Oil, }
 Yellow Wax, half a pound.
 ——— Resin, one ounce and a half.

Melt slowly together. (AMST.)—This is the *Unguentum emolliens*.

R. Resin, } of each, one part.
 Yellow Wax, }
 Lard, } of each, four parts.
 Olive Oil, }

Melt over a slow fire. (FULD.)

SWEDIAUR prescribes one part each, of resin and of wax, and six each, of lard and of oil.

R. Resin, four parts.
 Yellow Wax, } of each, two parts.
 Goat's Suet, }
 Olive Oil, three parts.

Melt together, and strain. (HISP.)

SWEDIAUR; and VAN MONS prescribe one part each, of the four first substances and two parts of the oil.

R. Resin, } of each, one part.
 Yellow Wax, }
 Turpentine, }
 Oil of Rape, two parts.

Melt slowly and strain. (MIL. COPEN.)

R. Resin, } of each, one part.
 Yellow Wax, }
 Mutton Suet, }
 Lard, three parts.

Melt together, and add when removed from the fire,

Turpentine, one half part.

Mix. (DAN.; HANN.; HASS.; OLDEN.; POL.; BORUS.; SAX.)

ROSS. prescribes one part each, of resin, of wax, of suet, and of turpentine, and four of lard.—FENN.; and SUEC.—three each, of resin, of wax, of suet, and of turpentine, and twelve of oil.—LIPP.—one each, of resin, of wax, of suet, and of turpentine, and five of oil.—GENEV.—twelve each, of resin, of Burgundy pitch, and of yellow wax, three of turpentine, and fourteen of olive oil.

R. Olive Oil, two pounds.
 Turpentine, one pound.
 Yellow Wax, } of each, seven ounces.
 Resin, }
 Mastic, } of each, two ounces.
 Olibanum, }
 Saffron, two drams.

Melt the wax, resin, mastic, and turpentine in the oil, and add the olibanum and

saffron, in powder, to the mass when half cold. (SARD.)

WIRTEM. prescribes two pounds and a half of oil, half a pound of yellow wax, two ounces of turpentine, one ounce each, of resin, of mastic, and of olibanum, and one dram of saffron.

2. *Black Basilicon; Unguentum Basilicum nigrum seu Resina nigra.* (LOND.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; PALAT.; SARD.; WIRTEM.; BRUGNATELLI; SPIELMANN; SWEDIAUR; VAN MONS)

Unguentum Resinae nigrae. (LOND.)

R. Black Resin, } of each,
 Yellow Resin, } nine ounces.
 ——— Wax, }
 Olive Oil, one pint.

Melt together, and express through a linen cloth.

R. Pitch, } of each, one part.
 Yellow Resin, }
 ——— Wax, }
 Olive Oil, four parts.

Melt the two first together; then add the wax and oil, and strain through a linen cloth. (AMST.; BATAV.; BAVAR.; BELG.; GALL.; FORM. HÔP. FR.; SARD.; SPIELMANN)

R. Pitch, } of each, one part.
 Resin, }
 Yellow Wax, }
 Lard, }
 Olive Oil, two parts.

Melt together. (ANT.; VAN MONS)

HISP. prescribes two parts each, of the four first substances, and three of the oil.

R. Pitch, } of each, one part.
 Yellow Wax, }
 Resin, }
 Turpentine, }
 Olive Oil, eight parts.

Melt together. (BRUGNATELLI)

BRUGNATELLI also prescribes an equal part each, of pitch, of resin, of turpentine, and of oil.

R. Yellow Wax, } of each three ounces.
 Mutton Suet, }
 Resin, }
 Pitch, }
 Turpentine, }
 Olive Oil, one pound.

Melt together, and add
 Powdered Olibanum,
 one ounce and a half.

Mix. (BRUNS.; WIRTEM.)

SWEDIAUR prescribes half a pound each, of pitch, of resin, and of yellow wax, one pound and a half of oil, three ounces of turpentine, and one ounce of olibanum.

R. Yellow Wax, two ounces.
Saffron, two scruples.
Mastic, }
Myrrh, }
Pitch, } of each, two drams.
Suet, }
Olibanum, }
Resin, } of each, half an ounce.
Turpentine, }
Olive Oil, three ounces.

Make an ointment. (PALAT.)

BROWN RESINOUS OINTMENT.

Unguentum fuscum. (BRUNS.; WIRTEM.)

R. Olive Oil, one pound and a half.
Yellow Wax, four ounces.
Resin, }
Pitch, } of each, two ounces.
Mastic, }
Turpentine, } of each, one ounce.

Melt, strain, and add when nearly cold,

Powdered Galbanum, } of each,
Sagapenum, } one ounce.
Olibanum, }
Round Birthwort, half an ounce.

Mix.

RESINOUS PLASTER; ADHESIVE PLASTER.

Emplastrum adhæsivum seu Lithargyricum Resinæ seu resinosum seu commune cum Resina seu Resinæ seu polychrestum seu commune adhæsivum seu lithargyricum cum Resina Pini. (LOND.; EDINB.; DUBL.; AMER.; AMST.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HANN.; HASS.; LUSIT.; POL.; BORUS.; SAX.; SUEC.; WURCEB.; AUGUSTIN.; BRUGNATELLI.; COXE.; PIDERIT.; SWEDIAUR.; VAN MONS.)

Emplastrum Resinæ. (LOND.)

R. Yellow Resin, half a pound.
Lead Plaster, three pounds.

Melt the plaster with a gentle heat, then add the resin in powder, and mix.

Emplastrum Resinosum. (EDINB.)

R. Plaster of Semivitreous
Oxide of Lead, five parts.
Resin, one part.

Melt slowly; then stir continually till cold.

Emplastrum Lithargyri cum Resina. (DUBL.)

R. Litharge Plaster, three pounds and a half.
Yellow Resin, half a pound.

Melt the plaster with a moderate heat; then add the resin in very fine powder, that it may melt quickly, and form a plaster.

AMST.; BATAV.; BELG.; GALL.; GENEV.; LUSIT.; and VAN MONS give the same formula as LOND.—AMER.; and COXE—the same as EDINB.—BRUGNATELLI prescribes eight parts of plaster and one part of resin.—ROSS.; WIRTEM.; and SPIELMANN—two of plaster and one of resin.—VAN MONS—five of plaster and one of terebinthina cocta. FERR.—six of simple plaster, one of plaster, of subcarbonate of lead, and one of resin.—AUGUSTIN—an equal part each, of resinous ointment, and of resin.

R. Olive Oil, two pounds.
Semivitreous Oxide of Lead, one pound.
Subcarbonate of Lead, four ounces.

Boil slowly, constantly stirring and occasionally adding a little water; to the consistence of a plaster, then add

Yellow Wax, eight ounces.
Turpentine, two ounces.

Mix. (FENN.; SUEC.)

BAVAR. directs eight parts of the oxide to be boiled in twelve of lard, and six of resin, and three of pitch to be added.

R. Simple Plaster, twenty-four parts.
Burgundy Pitch, six parts.
Turpentine, one part.

Melt together. (HASS.)

HANN.; POL.; BORUS.; and VAN MONS prescribe twenty-four parts of plaster, twelve of Burgundy pitch, and one of turpentine.—SWEDIAUR—sixteen of plaster, four of pitch, and one of turpentine.—SAX.—twelve of plaster, six of pitch, and one of turpentine.—FULD.—four of plaster, sixteen of resin, and four of turpentine.—PIDERIT—six of plaster, twelve of resin, and four of turpentine.

R. Simple Plaster, two pounds.
Resin, one pound.
Yellow Wax, one ounce and a half.
Melt together. (WURCEB.)

R. Simple Plaster, thirty-eight parts.
Yellow Wax, sixteen parts.

Melt, and add
Turpentine, three parts.
Lard, one part.

Mix. (VAN MONS)

- R. Simple Plaster, one pound and a half.
 Marshmallows Ointment, } of each,
 Olive Oil, } six ounces.
 Turpentine, three ounces.
 Yellow Wax, two ounces.

Mix. (BRUNS.)

- R. Simple Plaster, six pounds.
 Isinglass, dissolved in two ounces
 of Proof Spirit, half an ounce.
 Yolk of Wool, } of each, four ounces.
 Resin, }
 Yellow Wax, }
 Turpentine, half a pound.

Melt together the resin, wax and turpentine; then add the melted plaster, then the yolk triturated with the isinglass, and apply heat till the humidity is dissipated. (HISP.)

FERRUGINOUS ADHESIVE PLASTER.

Unguentum Lithargyri resinosum martiale.
 (SAX.)

- R. Resin, six parts.
 Turpentine, four parts.

Melt together, and add
 Powdered Mastic, two parts.
 Subcarbonate of Iron, one part.

Then incorporate
 Simple Plaster, twelve parts.

CORROBORANT PLASTER.

Emplastrum roborans seu stypticum.
 (FULD.)

- R. Adhesive Plaster (Form. of FULD.),
 two ounces.

Melt, and add
 Powdered Camphor, } of each,
 Oil of Caraway, } one dram.

Mix.

TETRANTHERA PICHURIM.—PICHURIM.

ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; HANN.; LIPP.; LUSIT.; BORUS.; ROSS.; SAX.; SÆC.; WIRTEM.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A shrub of Brazil. (*Enneandria Monogynia*, LINN.; *Laurineæ*, JUSS.; *Fig. Desc. Fl. Ant.* III. t. 213)

The seed, *Pichurim Bean* (*Faba Pichurim*.—Fève Pichurim, FR.; Brasilianische Bohnen, Pichurimbohne, GERM.; Kryddbøna, SWED.), is employed. It is an oval,

oblong nut, an inch or more in length, of an olive-red colour, nearly glabrous, heavy, convex on one side, and concave on the other. Its odour and taste resemble those of sassafras and of nutmeg.

It is employed in diarrhœa, dysentery, and flatulent colic.—Dose, two scruples.

ANTI-LEUCORRHOIC POWDER.

Pulvis contra Fluorem album. (HAMB.)

- R. Pichurim Bean, } of each,
 White Sugar, } half an ounce.

Make twelve doses.

TEUCRIUM.

Nine species of this genus of plants are used in medicine:

1. *Teucrium Chamædrys*, LINN.—Common Germander.

SYNONIMES:—Germandrée commune, Petit Chêne, FR.; Gamander, Gamenderlein, GERM.; Kamanderle, Ozankamenss, BOHEM.; Camedrio, PORT.; and SPAN.; Gamanderlyn, DUT.; Camedrio, Querciola, Calamandrina, ITAL.; Ozanka, POL.; Ekegræs, SWED.

DURL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; LIPP.; LUSIT.; CAST. BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the temperate and southern parts of Europe. (*Didynamia Gymnospermia*, LINN.; *Labiata*, JUSS.; *Fig. Flore Medic.* III. 183)

The flowering herb (*Herba Chamædryos*

seu *Trissaginis* seu *Quercula minoris*) is employed. It consists of a square, slender, hairy stem, and of opposite, petiolated, oval, smooth leaves of a greyish-green colour above, paler beneath, deeply crenelated, and of purpurine flowers, fixed, to the number of two or three, by short peduncles, in the axillæ of the superior leaves. It has a very feeble aromatic odour, which is lost by desiccation, and a moderately styptic and bitter taste.

It is excitant.

2. *Teucrium flavum*, LINN. — Yellow-Flowered Germander.

SYNONIMES : — Germandrée jaune, FR. ; Gelber Garnander, GERM.

HISP.

A plant of Spain and Italy. (Fig. Lob. Ic. 490. 1)

The flowering herb (*Herba Teucrii flavi*) is employed. It consists of a pubescent stem, furnished with oval leaves, crenelated inferiorly, entire at the summit, and of yellow flowers in lateral racemes. It has a bitter taste.

It is excitant.

EXTRACT OF GERMANDER. (FORM. HÔP. FR. ; GALL. ; SARD.)

R. Tops of Germander, any quantity.
Boiling Water, a sufficient quantity.

Infuse ; then strain, and evaporate over a slow fire to the consistence of an extract. (FORM. HÔP. FR. ; GALL.)

R. Fresh and bruised Herb of Germander, eight pounds.

Macerate it for three days in an alembic ; then distil off about two ounces of liquor ; pour upon the residue,

Water, twenty-four pounds ; and, after six hours' digestion, boil down to one third ; then, having clarified and strained the liquor, evaporate it to the consistence of an electuary, adding the distilled water to this when it is only warm. (SARD.)

DECOCTION OF GERMANDER.

Decoctum Chamædrys amarum. (HISP.)

R. Tops of Germander, half an ounce.
—— Wormwood, } of each,
—— Lesser Centaury, } two drams.
Water, two pounds.

Boil down to one half ; and add towards the end,

Chamomile Flowers, two drams.
Senna, one ounce and a half.

The senna is omitted, when the laxative effect is not desired.—Dose, from four to six ounces.

VOL. II.

INFUSION OF GERMANDER. (FORM. HÔP. FR. ; GALL. ; SPIELMANN)

R. Herb of Germander, one ounce.
Boiling Water, one pound and a half.

Infuse. (FORM. HOP. FR. ; SPIELMANN)

R. Green Leaves of Germander, one ounce.
Boiling Water, two pounds.

Infuse, and add when strained,

Syrup of Maidenhair or Honey, one ounce.

Mix. (GALL.)

3. *Teucrium Chamæpitys*, LINN.—Ground Pine.

SYNONIMES : — Ivette commune, FR. ; Schlagkraut, Feldcypresse, Erdweihrauch, GERM. ; Ywa, Polnj cypris, BOHEM. ; Camepiteos, Pinillo oloroso, SPAN. ; Veldey-press, DUT. ; Camepizio, ITAL. ; Iwinka, Piznowa, POL. ; Chamepite, PORT.

AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; HISP. ; GALL. ; FERR. ; FULD. ; GENEV. ; LIPP. ; ROSS. ; SAX. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; SPIELMANN ; ZARDA.

A common plant of Europe. (Fig. Zorn, Ic. pl. t. 120)

The following herb (*Herba Chamæpityos* seu *Iva arthritica* seu *Chamæmori*) is employed. It consists of a branching, hairy stem, of leaves divided almost to the mid-rib, and by three linear notches, and of yellow, sessile, solitary flowers, in the axillæ of the superior leaves. It has a resinous and feebly balsamic odour, which becomes dissipated by desiccation. Its taste is acrid, bitter, and styptic.

It is tonic and stomachic.

EXTRACT OF GROUND PINE. (SARD.)

R. Fresh Herb of Ground Pine, bruised, eight pounds,

Digest for three days in an alembic ; then distil off about two ounces of liquor in a water-bath ; pour upon the residue,

Water, twenty-four pounds.
and boil down to one third ; then clarify and strain the decoction, and evaporate it to the consistence of an electuary, adding the distilled water to this when it is merely warm.

4. *Teucrium Iva*, LINN.

SYNONIMES : — Ivette musquée, FR. ; Bisamginsel, GERM.

GALL.

4 S

A plant of the south of Europe. (*Fig. Cav. II. t. 120*)

The herb (*Herba Ivæ seu Chamæpytos Monspeiaceæ*) is employed. It consists of a branching stem bearing entire leaves furnished with but two teeth at the summit. It has a strong, resinous, and slightly musky odour, particularly during hot weather, and a bitter taste.

It is excitant and stomachic.

5. *Teucrium Marum*, LINN.—Marum.

SYNONIMES:—Marum, FR.; Amberkraut, Ratzenkraut, GERM.; Lathymo, CHIN.; Kocij korenj, BOHEM.; Maro cortuso, SPAN.; Erba gatta, ITAL.; Ambrowe ziele, POL.; Katgræs, SWED.

DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A plant of the south of Europe and of the Levant. (*Fig. Zorn, Ic. pl. t. 60*)

The flowering herb (*Herba Mari veri seu Syriaci*) is employed. It consists of slender cottony stems, furnished with small, oval, lanceolated, petiolated leaves of greyish-green colour above and white and cottony beneath, and of purpurine flowers, solitary in the axillæ of the superior leaves, but so approximated as to form a long lateral, secund raceme. It has a very aromatic and penetrating odour, and a pungent, acrid, aromatic, and bitterish taste.

It is stimulant, diaphoretic, diuretic, emmenagogue, pectoral, and sternutatory.

ETHEREOUS TINCTURE OF MARUM.

Tinctura Mari veri æthereæ. (OLDEN.; NIEMANN)

R. Herb of Marum,	} of each,
HOFFMANN'S Anodyne	
Liquor,	
Concentrated Alcohol,	three ounces.

Digest for four days, at a slow temperature, in a covered vessel, occasionally shaking the vessel; then filter.

It is extolled in paralysis. It may be given to those with whom the infusion of the plant occasions nausea.—Dose, forty drops.

COMPOUND AROMATIC LIQUOR.

Aqua vulneraria seu styptica Clementina. (FERR.)

R. Juice of Marjoram,	} of each,
Sanicle,	
	three ounces.

Juice of Vervain,	} of each,
Houseleek,	
Burnet Saxifrage,	
Milfoil,	
Bryony,	} three ounces.
Marum,	
Muriate of Soda,	} of each,
Subcarbonate of Potass,	
Supertartrate of Potass,	
Aloes,	four ounces.
Vinegar,	one ounce and a half.
	one pound.

Digest for twenty-four hours, and distil to dryness. Pulverize the residue, add it to the product, and re-distil. Repeat this operation three times successively.—Dose, to half an ounce.

6. *Teucrium Creticum*, LINN.

SYNONIME:—Polium de Crête, FR.

GALL.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of Crete and Egypt. (*Fig. Alp. Exot. p. 103, t. 102*)

The flowering herb (*Herba et Summitates Teucrii seu Polii Cretici seu Rorismarini Stæchadis facie*) is employed. It consists of a square stem furnished with sessile, lanceolated, entire leaves, and of violet-coloured, petiolated flowers, disposed by threes in axillary racemes. It has a strong, agreeable, and aromatic odour, and a very bitter taste.

7. *Teucrium montanum*, LINN.

SYNONIME:—Polium de montagne, FR.

GALL.

A plant of the south of Europe. (*Fig. Zorn, Ic. pl. t. 362*)

The tops (*Summitates Polii montani*) are employed. They consist of cottony, whitish branches, furnished with lanceolated, entire leaves, curved at the borders, cottony beneath, and of whitish or yellowish flowers, nearly sessile, forming a kind of round, terminal umbel.

It has a pleasant odour, and a bitter taste.

8. *Teucrium Scordium*, LINN.—Scordium, Water Germander.

SYNONIMES:—Scordium, Germandrée d'eau, FR.; Knoblauchsgamander, Skordienkraut, Lachenknoblauch, GERM.; Plany neb wodnj cesnek, BOHEM.; Lægsurt, DAN.; Scordio, ITAL.; and SPAN.; Waterbock, DUT.; Czosnkowe ziele, POL.; Escordio, PORT.; Gamander, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant growing in every part of Europe. (*Fig. Zorn, Ic. pl. t. 36*)

The herb (*Herba Scordii* seu *Chamædryis aquatica*) is employed. It consists of a hairy, branching stem, and of oval, oblong, sessile, crenelated, or dentated leaves. It has, in the fresh state, an alliaceous and balsamic odour, which diminishes by desiccation. Its taste is bitter, acrid and rather styptic.

9. *Teucrium Scorodonia*, LINN.

SYNONIME :—*Scordium faux*, FR.

GALL.; WIRTEM.; GUIBOURT.

A plant of Europe. (*Fig. Bulh. Herb. t. 301*)

The herb (*Herba Scorodonia* seu *Salvia sylvestris* seu *agrestis*) is employed. It consists of petiolated, cordiform, dentated, and wrinkled leaves.

It is stimulant, tonic, and diaphoretic.

CONSERVE OF SCORDIUM.

Conserva Scordii. (HISP.)

R. Fresh Leaves of Scordium, one part.

Bruise them in a marble mortar with a wooden pestle, adding gradually,

White Sugar, two parts.

Incorporate.

EXTRACT OF SCORDIUM.

Extractum Scordii. (BRUNS.; PALAT.; SARD.; WIRTEM.)

R. Herb of Scordium, one pound.
Spring Water, six pounds.

Digest for four days in a warm place; then boil a little, and express. Evaporate the liquor to the proper consistence. (BRUNS.; PALAT.; WIRTEM.)

R. Herb of Scordium, eight pounds.

Digest for three days in an alembic; then distil off about two ounces of liquor, and pour upon the residue,

Water, twenty-four pounds.

Boil down to one third. Then having clarified and strained the liquor, evaporate it to the consistence of an electuary, and

add the distilled water to this when it is only warm. (SARD.)

Dose, from half a dram to one dram.

DISTILLED WATER OF SCORDIUM.

Aqua Scordii. (BRUNS.; GALL.; PALAT.; SARD.)

R. Flowering Herb of Scordium, one part.
Water, four parts.

Distil one half, and re-distil with fresh herb. (SARD.)

GALL. prescribes five parts of herb and twenty of water; ten parts to be distilled. —BRUNS.—one of herb and three of water; one half to be distilled. —PALAT.—one of herb and four of water; one half to be distilled.

Dose, from one ounce to three ounces.

SYRUP OF SCORDIUM.

Syrupus Scordii seu *de Scordio.* (GALL.; WIRTEM.)

R. Herb of Scordium, four ounces.
Boiling Water, one pound.

Infuse during one night; then express, clarify, and dissolve in nine ounces of the strained liquor,

White Sugar, sixteen ounces.

Make a syrup. (WIRTEM.)

R. Dried Herb of Scordium, one ounce.
Distilled Water of Scordium, two pounds.

Digest in a covered vessel in a water-bath for two hours; then strain, filter, and add to the liquor,

White Sugar, two parts.

Dissolve in a water-bath, and strain through flannel. (GALL.)

Dose, from half an ounce to one ounce.

VERMIFUGE INFUSION.

Infusio anthelmintico-incitans. (BRERA)

R. Leaves of Scordium, half an ounce.
Water, a sufficient quantity
to obtain eight ounces of strained infusion.
Add

Sulphuric Æther, half a dram.

To be taken by spoonfuls.

VERMIFUGE POTION.

Mixtura anthelmintico-incitans. (BRERA)

R. Corsican Moss, two drams.
Leaves of Scordium, half an ounce.
Water, a sufficient quantity
to obtain ten ounces of strained infusion.
Add

Sulphuric Æther, half a dram.
Syrup of Cinnamon, half an ounce.
Mix.—To be taken gradually.

ANTISEPTIC DECOCTION. (CADET DE GASSICOURT)

R. Leaves of Scordium, }
——— Alliaria, } of each,
——— White Hore- } two ounces.
hound, }
Water, } four ounces.

Boil, and add

Oxymel of Squill, half a pound.
Theriacal Vinegar, one ounce.
Nitrate of Potass, three drams.

Mix.—It is recommended in putrid diseases with difficulty of expectoration.

AROMATIC FOMENTATION.

Fomentum aromaticum. (AUGUSTIN)

R. Herb of Scordium, } of each,
Flowers of Chamomile, } one ounce.
Boiling Water, } three drams.

Boil a little, and add to the strained liquor,

Spirit of Camphor, four ounces.

Mix.

TINCTURE OF SCORDIUM.

Tinctura seu Essentia Scordii. (BRUNS.; PALAT.; WIRTEM.)

R. Herb of Scordium, three ounces.
Alcohol, sixteen ounces.

After three days' digestion, pour the strained liquor upon

Herb of Scordium, two ounces.

Macerate for some time; then express, and filter. (PALAT.; WIRTEM.)

BRUNS. directs one ounce of scordium to be digested in four ounces of concentrated alcohol.

It is stimulant, and is reputed alexipharmic and diaphoretic.—Dose, sixty drops.

STAHL'S ALEXIPHARMIC ESSENCE.

Essentia alexipharmaca Stahl. (BRUNS.; LIPP.; PALAT.; WIRTEM.; CADET DE GASSICOURT; SPIELMANN)

R. Root of Burnet Saxifrage, } of each,
——— Swallow-wort, } one ounce
——— Carline Thistle, } and a half.
——— Angelica, }
——— Master-wort, } of each,
——— Elecampane, } one ounce.

Herb of Scordium, three ounces.

Alcohol, two pounds and a half.

Digest, express, and filter.

BRUNS.; WIRTEM.; PALAT.; and SPIELMANN prescribe one ounce and a half each, of burnet saxifrage and of swallow-wort, half an ounce of carline thistle, and and one ounce each of angelica, of master-wort, and of elecampane; but SPIELMANN prescribes six ounces of scordium and two pounds of alcohol, while PALAT. omits the scordium, and requires only one pound of alcohol. The formula of CADET DE GASSICOURT is too mutilated for us to report it here.

R. Root of Angelica, } of each,
——— Master-wort, } half an ounce.
——— Elecampane, }
——— Carline Thistle, } two drams.
——— Swallow-wort, }
——— Burnet Saxi- } of each,
frage, } three drams.
Herb of Scordium, two ounces.
Alcohol, one pound.

Digest for two days at a moderate heat; then express.

It is excitant, tonic, stomachic, carminative, diuretic, and sudorific.—Dose, from forty to fifty drops.

STAHL'S TEMPERATE ALEXIPHARMIC ESSENCE.

Essentia alexipharmaca temperata Stahl. (WIRTEM.; SPIELMANN)

R. Root of Burnet Saxifrage, } of each,
——— Swallow-wort, } three ounces.
Herb of Scordium, two ounces.
Alcohol, three pounds.

After sufficient digestion, filter, and distil off one third of the spirit in a water-bath. Preserve the remainder. (WIRTEM.)

R. Tincture of Burnet Saxifrage, } of each,
——— Swallow- } one ounce.
wort, }
——— Scordium, two ounces.

Mix. (SPIELMANN)

It is excitant, but less so than the preceding.—Dose, from forty to sixty drops.

COMPOUND ESSENCE OF SCORDIUM.

Essentia de Scordio composita, Diascordium liquidum, Tinctura Scordii composita. (SAX.; WIRTEM.; VAN MONS)

R. Herb of Scordium, two ounces.
Lemon Peel, one ounce.
Myrrh, } of each,
Root of Gentian, } six drams.
——— Angelica, }
——— Swallow-wort, } of each,
——— Lovage, } half an ounce.

Root of Zedoary, }
 ——— Contrayerva, } of each,
 ——— Tormentil, } half an ounce.
 ——— Valerian,

Amber, }
 Aloe Wood, } of each, three drams.
 Saffron, } two drams and a half.
 Elder Flowers, } two drams.
 Opium, } one dram.

Make a powder of the whole, and pour upon it,

Spirit of Juniper, three pounds.

Digest at a moderate heat in a covered vessel, and filter. (WIRTEM.)

R. Dried Herb of Scordium, one part.

————— Blessed Thistle, }
 ————— one half part.

Root of Angelica, one fourth part.
 Alcohol, eight parts.

Digest for eight days in a covered vessel; then express and filter. (SAX.)

R. Leaves of Germander, four parts.

————— Blessed Thistle, two parts.

Fresh Root of Angelica, one part.

Proof Spirit, thirty-two parts.

Infuse without heat for several days, then express, and filter. (VAN MONS)

It is reputed alexipharmic, anodyne, and stomachic.—One ounce contains about two grains of opium.

HOFFMANN'S SPIRITUOUS BALSAMIC ELIXIR.

Elixir balsamicum spirituosum Hoffmanni.
 (BRUNS.; WIRTEM.; SPIELMANN)

R. Herb of Scordium, one ounce.

————— Lesser Centaury, } of each.

Orange Peel, } one ounce

Myrrh, } and a half.

Root of Zedoary, } of each,

Nutmegs, } three drams.

Aloe Wood one dram and a half.

Saffron, one dram.

Amber, three drams.

Alcohol, one pound and a half.

Digest, express and filter. Add

Oil of Cardamoms, } of each, ten drops,

————— Cloves, }

dissolved in

Spirit of Ammonia, one ounce.

Mix.—It is excitant, and was formerly extolled as alexipharmic, anti-catarrhal, stomachic, and principally recommended by HOFFMANN in chronic diarrhoea.—Dose, sixty drops, twice a day.

THALICTRUM FLAVUM, LINN.—MEADOW RUE.

SYNONIMES:—Pigamon, Pigamon jaune, Rue des prés, Fausse rhubarbe, Rhubarbe des pauvres, FR.; Wiesenrauteheilblatt, Wiesenrautewurz, GERM.; Kneppanden, DAN.; Waterruid, DUT.; Wzodowiec, POL.; Torrvaerkroes, SWED.

BRUNS.; GALL.; GENEV.; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN.

A plant common in the meadows of Eu-

rope. (*Polyandria, Polygynia*, LINN.; *Ranunculaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 406*)

The root (*Radix Thalictri flavi seu magni seu Rhubarbari pauperum*) is employed. It is of a light brown colour in the fresh state, and of a deep brown when dried, yellow within, and full of a sweetish juice, leaving an impression of bitterness and acidity. It tinges with yellow the saliva, the fæces, and urine. It is purgative.—Dose, from one ounce to two ounces.

THAPSIA ASCLEPIUM, LINN.—NARROW LEAVED CARROT.

SYNONIMES:—Thapsie, Thapsie Asclepion, FR.; Tapsia, Comino rostico, SPAN.

HISP.

A plant of the south of Europe. (*Pentandria Digynia*, LINN.; *Umbelliferae*, JUSS.; *Fig. Mor. Hist. t. 18*)

The root and seeds are employed. They are stimulant and are reputed resolvent.

ANTI-RHEUMATIC OIL.

Oleum Thapsiæ anti-rheumaticum. (HISP.)

R. Root of Narrow-leaved

Carrot, } of each,

Leaves of Rosemary, } one pound.

Red Wine, }

Olive Oil, } four pounds.

Digest for four days; then boil to the consumption of the humidity; and strain.

THEA VIRIDIS, LINN.—GREEN TEA.

SYNONIMES:—Thé, FR.; Thee, GERM.; BOHEM.; DUT.; and SWED.; Cha, ARAB.; CHIN.; DUK.; and PERS.; The DAN.; Te, ITAL.; PORT; and SPAN.; Tjao, JAV.; Tjcha, RUSS.

BRUNS.; HISP.; GALL.; FERR.; WIRTEM.; AINSLIE; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A shrub of China. (*Polyandria Monogynia*, LINN.; *Aurantiaceæ*, JUSS.; *Fig. Flore Medic. VI. 339*)

The leaves are employed; they are moderately petiolated, green, somewhat shining, hard, glabrous, oval, lanceolated, or

elliptical, serrated at the superior portion, entire at the base, dried and rolled, either only lengthwise, or transversely besides; sometimes, when cut in pieces, they present small, oblong, or round, greenish, wrinkled masses, of an aromatic and pleasant odour, and of a bitter and rather styp-tic taste.

In commerce there are a great many kinds of tea, the principal of which bear the names of *Heyswen*, *Schulang*, *Saot-chaon*, or *Bony*; and *Pekao*. The two last, the least esteemed, belong to the *Thea Bohea*, LINN., which is considered to be merely a variety of the *T. viridis*.

It is excitant and diaphoretic.—Dose, one or two drams in a pint of boiling water.

THEOBROMA CACAO, LINN.—COCOA TREE.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A tree of South America. (*Polyadelphia Pentandria*, LINN.; *Malvaceæ*, JUSS.; *Fig. Flore Medic. II. 83*)

The seed, called *Cocoa Nut*; *Fructus seu Nuces seu Nuclei Cacao*, is employed. It is ovoid, the size of a filbert, fleshy, of a violaceous odour, and of an oleaginous and bitterish taste.

RESTORATIVE POWDER. (HAMB.)

R. Powdered Torrified Cocoa Nut, eight ounces.
 ———— Rice, } of each,
 ———— White Sugar, } four ounces.
 ———— Cinnamon, one dram.

Mix.

COCOA PASTE.

Pasta cacaotina. (HASS.; LIPP.; SAX.; NIEMANN; PIDERIT)

R. Cocoa Nuts, any quantity.

Detach the husks by a slight torrifaction, or by immersion in boiling water; then dry the white part, reduce to coarse powder,

and form a paste by pounding in a heated mortar.

This paste comes to us from America, where it is prepared with nuts of an inferior quality.

CHOCOLATE OF HEALTH.

Chocolata medica seu simplex, *Cacao tabulata seu preparata*, *Pasta cacaotina saccharata*; *Chocolat de santé*. (GALL.; FULD.; SAX.; NIEMANN)

R. Cocoa Paste, } of each, an equal part.
 White Sugar, }
 Triturate together, and form tablets.
 (SAX.; NIEMANN)

Caraccas Cocoa, torrified and prepared, one thousand seven hundred and fifty-eight parts.
 West India Cocoa Nuts, prepared in the same manner, three thousand parts.
 White Sugar, five thousand parts.
 Cinnamon, in powder, forty parts.

Pound the nuts in a heated iron mortar with one fourth of the sugar, then bruise it by portions with an iron roller on a hot polished slab; add the remainder of the sugar and the cinnamon, and having submitted it again to the action of the roller for seven or eight minutes longer, dry it in iron moulds. (GALL.)

FULD. prescribes one pound of cocoa nut and eight ounces of sugar, without cinnamon.

This chocolate scarcely deserves the title applied to it; it is more difficult of digestion than the following.

VANILLA CHOCOLATE.

Chocolat à la vanille. (GALL.; WIRTEM.; NIEMANN; SPIELMANN)

R. Caraccas Cocoa, slightly torrifed, sixteen ounces.

Pulverize it, and reduce it to paste in a heated iron mortar, adding

Sugar, twelve ounces.

Mix intimately, and add

Vanilla, triturated with three drams of sugar, one dram.

Put it into moulds. (SPIELMANN)

The *Chocolat à une vanille* is that which contains one scruple of vanilla in one pound of chocolate; the *Chocolat à deux vanilles* contains two scruples to the pound.

WIRTEM. prescribes one pound each, of torrifed cocoa nut and of sugar, and four of powdered vanilla.—GALL.—forty parts of vanilla, in the proportions indicated in the preceding paragraph.

According to WIRTEM., two drams each, of cinnamon and of cloves are sometimes added; sometimes also, a few drops of balsam of Peru.—NIEMANN indicates, after WIEGLEB, under the name of *Chocolata aromatica*: cocoa nut, sixteen ounces; sugar, sixteen ounces; cinnamon, half an ounce; cloves, two drams; lesser cardamoms, one dram; vanilla, powdered with sugar, one dram.

In general, one ounce of chocolate is sufficient for four or six ounces of water, which may be partly substituted by milk.

COCOA-NUT OIL.

Butyrum seu Oleum Cacao, Oleum Cacao spissatum seu Theobromæ Cacao expressum seu Fructus Cacao. (AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; HANN.; HASS.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SARD.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI; PIDERIT; SPIELMANN; SWEDIAUR; VAN MONS)

R. Cocoa Nuts, any quantity.

Break and remove the pellicle; winnow and pulverize, pass through a hair sieve, and pound in a heated mortar; enclose the paste in a ticking bag, and press it between two heated tin plates; melt the oil in hot water, set it aside, strain, and pour it into paper moulds. (VAN MONS)

R. Cocoa Nuts, slightly torrifed, any quantity.

Remove the bark, bruise the nuts, heat the pulp slightly, enclose it in a bag, and submit it to a tin press heated by immersion in hot water; digest the oil with two parts of water, in a sand-bath, and preserve it when cold. (ANT.; DAN.; HISP.; OLDEN.; SUEC.; HERBIP.; BRUGNATELLI; SPIELMANN; SWEDIAUR; VAN MONS)

R. Cocoa Nuts, any quantity.

Torrify it slightly, remove the bark, and bruise the nuts; put the paste into a linen bag, and expose it to the vapour of boiling water; when it is well impregnated with it, submit it to the action of a heated press. (BATAV.; BELG.; FENN.; FERR.; HANN.; HASS.; LIPP.; POL.; BORUS.; SAX.; BRUGNATELLI; PIDERIT; VAN MONS)

R. Cocoa Nuts, any quantity.

Torrify slightly; then having bruised them in a mortar, triturate on a marble slab; then moisten the paste with a sufficient quantity of water, enclose it in a bag, and express between two heated tin plates; melt the oil in a water-bath, filter it through paper, and preserve it when cold. (GALL.)

R. Cocoa Nuts, any quantity.

Torrify slightly in an iron stove; then, having bruised them in a mortar, triturate in a hot stove: boil the paste in water, till the oil rises to the surface; then, having set this apart to congeal, liquify it in a water-bath, filter it while hot through paper, and preserve it when cold. (AMST.; AUSTR.; BAVAR.; BRUNS.; FULD.; LUSIT.; PALAT.; SARD.; WIRTEM.; BRUGNATELLI; PIDERIT; SWEDIAUR; VAN MONS)

It is demulcent and alimentary.—LENTIN prescribed, thrice a day, one dram with yolk of egg or fat broth, in cardialgia attributed to excoriation of the stomach.

PILLS OF COCOA-NUT OIL. (HENRI)

R. Cocoa-nut Oil, two drams.
Powdered Marshmallows Root,
a sufficient quantity

to make a pilular mass.

PECTORAL CREAM. (CADET DE GASSICOURT; PIERQUIN)

R. Cocoa-nut Oil, two ounces.
Syrup of Tolu, } of each,
Maidenhair, } one ounce.
White Sugar, half an ounce.

Mix.—To be taken by spoonfuls. (CADET DE GASSICOURT; PIERQUIN)

This formula is by TRONCHIN.

R. Cocoa-nut Oil, }
Syrup of Red Poppy, } of each,
Orange-flower Water, } one ounce.
Oil of Almonds, }

Mix.—To be taken by spoonfuls. (CADET DE GASSICOURT)

EMOLLIENT SUPPOSITORY. (CADET DE GASSICOURT)

R. Cocoa-nut Oil, } of each,
Hard Cerate, } an equal part.

Make a conical bougie.

COSMETIC CERATE. (VAN MONS)

R. White Wax, one part.
Melt, and add

Oil of Almonds, two parts.
Cœcoo-nut Oil, one part.

Stir till perfectly cold.

LABIAL CERATE. (VAN MONS)

R. White Wax, one part.
Oil of Almonds, three parts.
Cocoa-nut Oil, one part and a half.
Alkanet, one half part.

Macerate for a quarter of an hour, and express; when the mass begins to solidify, add

Essence of Bergamotte, one half part.
Mix.

THLASPI.

Four species of this genus are employed :

1. *Thlaspi campestre*, LINN.

GALL.

A common plant of Europe. (*Tetradynamia Siliculosa*, LINN.; *Cruciferae*, JUSS.; *Fig. Blackw. Herb. t. 409*)

The seeds (*Semen Thlaspeos*) are employed. They are round, flattish, of a deep-brown colour, and of an acrid and pungent taste.

2. *Thlaspi arvense*, LINN.—Penny Cress.

SYNONIMES :—*Thlaspi des champs*, FR.; *Bawernsenf*, GERM.; *Penjzek*, aneb *sedlska horcice*, BOHEM.; *Pengeurt*, DAN.; *Wilde mosterd*, DUT.; *Penningegræs*, SWED.

BRUNS.; GALL.; WIRTEM.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; SPIELMANN; ZARDA.

The seeds (*Semen Thlaspeos*) are employed.

They are very small, round, and rather flattened.

Their odour is strong and alliaceous, and their taste subacrid and bitter.

3. *Thlaspi sativum*, CAND.—Common Cress.

SYNONIMES :—*Cresson Alénois*, *Cresson des jardins*, FR.; *Gartenkresse*, GERM.; *Hansekerse*, DAN.; *Nastuerco*, SPAN.; *Tuinkers*, DUT.; *Crescione*, ITAL.; *Nasturoga*, PORT.; *Trægardskress*, SWED.

ANT.; BATAV.; BRUNS.; HISP.; GALL.; FORM. HÔP. FR.; GENEV.; BERGIUS;

GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 16*)

The herb and seeds are employed.

The herb (*Herba Nasturtii hortensis*) consists of a branching stem, and of alternate, petiolated, long, smooth, and multifid leaves.

The seeds are oblong, smooth, and of a brown colour.

The plant exhales, by rubbing, a feeble but agreeable odour; its taste is bitterish and acrid.

It is excitant and anti-scorbutic.

4. *Thlaspi Bursa Pastoris*, LINN.—Shepherd's Purse.

SYNONIMES :—*Bourse*, *Bourse à pasteur*, FR.; *Täschelkraut*, *Taschenkraut*, *Hirtenasche*, *Hirtenäschelkraut*, GERM.; *Bolsa de pastor*, SPAN.; *Bearsjeskruid*, DUT.

BRUNS.; HISP.; GALL.; GENEV.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe. (*Fig. Zorn, Ic. pl. t. 158*)

The herb (*Herba Bursæ pastoris seu Cancræ seu Sanginariæ*) is employed. It presents an assemblage of radical pinnatifid leaves. It has a vapid odour, resembling that of cabbage, and an acrid and astringent taste.

It is very feebly astringent, and is scarcely ever used.

THUYA OCCIDENTALIS, LINN.—AMERICAN ARBOR VITÆ.

SYNONIMES:—Thuya du Canada, Cèdre blanc, FR.; Lebensbaum, GERM.; Livestræ, DAN.; Arbol de la vida, SPAN.; Levensboom, DUT.; Albero di vita, ITAL.; Aarvore da vida, PORT.; Lifvets-træd, SWED.

BRUNS.; FERR.; WIRTEM.; MURRAY and GMELIN; SPIELMANN.

A tree of North America. (*Monœcia Monadelphica*, LINN.; *Conifera*, JUSS.; *Fig. Blackw. Herb. t. 210*)

The wood and leaves are employed.

The wood (*Lignum Arboris vitæ*) is esteemed diaphoretic. It has been recommended in the treatment of syphilitic diseases.

The leaves are short, opposite, scaly, greenish, of a strong odour when rubbed, and of a bitterish taste. The decoction was formerly reputed resolvent.

THYMUS.—THYME.

Of the species of this genus of plants which are mentioned in pharmacopœias, we quote the following four:

1. *Thymus Alpinus*, LINN.—Alpine Thyme.

SYNONIMES:—Thym des Alpes, FR.; Alpenthymiam, Alpenquendel, GERM.

BRUNS.

A plant of the Alps. (*Didymia Gynospermia*, LINN.; *Labiata*, JUSS.; *Fig. Jacq. Fl. Austr. I. t. 97*)

The flowering herb (*Herba Clinopodii montani*) is employed. It consists of round, branching stems, and of petiolated, round, obtuse, somewhat serrated, concave leaves. It has a very agreeable aromatic taste and smell.

2. *Thymus mastichina*, LINN.—Mastic Thyme.

SYNONIMES:—Thym ciliée, Marjolaine sauvage, FR.; Mastix Thymian, GERM.; Meiorana silvestre, SPAN.

HISP.

A plant of Spain and Italy. (*Fig. Blackw. Herb. t. 134*)

The flowering herb (*Herba Mastichinae Gallorum seu Mari vulgaris*) is employed. It consists of a cylindrical stem, and of petiolated, oval, slightly serrated leaves, smooth on both sides. The flowers are white, and arranged in verticilli. It has an agreeable odour and taste.

3. *Thymus vulgaris*, LINN.—Garden Thyme.

SYNONIMES:—Thym commun, FR.; Gemeiner Thymian, GERM.; Wlaska nekzjmska materj dausska, BOHEM.; Thy-

VOL. II.

mian, DAN.; Tomillo, SPAN.; Gemeen Thym, DUT.; Timo, ITAL.; Tymian, Szmer wloski, POL.; Tomilho, PORT.; Timjan, SWED.

AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A plant of the south of Europe. (*Fig. Zorn. Ic. pl. t. 458*)

The flowering herb (*Herba et Flores Thymi vulgaris seu hortensis*) is employed. It consists of a cylindrical, slightly branching, rough stem, furnished with small, opposite, sessile leaves with slightly rolled edges; narrow, nearly obtuse, slightly pubescent beneath, of an ash-grey colour above, and bearing at the summit of the branches, verticilli of purpurine flowers forming a head or a spike, accompanied by bractæ. It has an agreeable, strong, and aromatic odour, and a warm and aromatic taste.

All these plants are stimulant.

INFUSION OF THYME. (FORM. HÔP. FR.)

R. Tops of Thyme, } of each,
Grated Liquorice, } two drams.
Boiling Water, a sufficient quantity
to obtain two pounds of strained liquor.

ESSENTIAL OIL OF THYME.

Oleum Thymi æthereum, Ætheroleum Thymi.
(BAVAR.; BRUNS.; DAN.; GALL.; HANN.; PALAT.; SARD.; SAX.)

R. Flowering Herb of Thyme, five parts.
4 T

Water, seven parts.
Distil. (GALL.)

DAN. prescribes one part of flowers and six parts of water.—HANN.; and SAX.—one of flowers and eight of water.—BAVAR.—one of flowers and nine of water.

R. Flowering Herb of Thyme, twenty-five parts.

Water, seventy-five parts.
Muriate of Soda, three parts.

Distil at the end of three days.

Dose, from two to four drops.

DISTILLED WATER OF THYME.

Aqua Thymi. (GALL.; HERBIP.)

R. Flowering Herb of Thyme, five parts.
Water, twenty parts.

Distil ten parts. (GALL.)

HERBIP. prescribes one part of herb and six parts of water.

Dose, from two to four ounces.

SPIRIT OF THYME.

Spiritus seu Alcohol Thymi. (BRUGNATELLI)

R. Flowering Herb of Thyme, one pound two ounces.
Proof Spirit, eight pounds.

Distil three fourths.

Dose, from half a dram to one dram.

AROMATIC SPIRIT.

Spiritus aromaticus. (LIPP.)

R. Herb of Thyme, three ounces.
—— Marum, one ounce.
Root of Zedoary, } of each,
Caraway, } two ounces.
Alcohol, eight ounces.

Macerate for twenty-four hours, and add,
Water, eight pounds.

Distil slowly eight pounds.

4. *Thymus Serpyllum*, LINN.—Wild Thyme,
Mother of Thyme.

SYNONIMES:—Serpolet, FR.; Quendel, Feldkümmel, GERM.; Materj dausska, BOHEM.; Wild Thymian, DAN.; Serpol, SPAN.; Wild Tym, DUT.; Sermollino, Serpillo, ITAL.; Macierzanka, POL.; Serpao, PORT.; Schadownik, RUSS.; Backtimian, SWED.

AMST.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENÈV.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; WURCEB.; BER-

GIUS.; BRUGNATELLI.; GUIBOUT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A plant common in every part of Europe. (Fig. Flore Medic. VI. 326)

The flowering herb (*Herba et Flores Serpylli*) is employed. It presents a branching, rough stem, furnished with small, nearly sessile leaves, ciliated at the borders, oval or lanceolated, and bearing small groups of very short, axillary, pedicellated flowers, forming very short spikes, and frequently a kind of terminal head. Its odour is agreeable, and its taste bitter and balsamic.

OIL OF WILD THYME.

Oleum Serpylli æthereum, Ætheroleum Serpylli. (AUSTR.; BAVAR.; BRUNS.; HANN.; PALAT.; SAX.)

R. Flowering Herb of Wild Thyme, one part.
Water, eight parts.

Distil. (HANN.; SAX.)

BAVAR. prescribes one part of wild thyme and nine parts of water.—AUST.—one of wild thyme and twelve of water.

R. Flowering Herb of Wild Thyme, twenty-five parts.
Water, seventy-five parts.
Muriate of Soda, three parts.

Macerate for three days, and distil. (BRUNS.; PALAT.)

DISTILLED WATER OF WILD THYME.

Aqua Serpylli. (SARD.)

R. Flowering Herb of Wild Thyme, one part.
Water, four parts.
Distil one half, and re-distil with fresh herb.

SPIRIT OF WILD THYME.

Spiritus Serpylli. (AUSTR.; BRUNS.; DAN.; FULD.; HANN.; HASS.; LIPP.; PALAT.; POL.; BORUS.; SARD.; SAX.; WIRTEM.; HERBIP.; PIDERIT.; SWEDIAUR)

R. Flowering Herb of Wild Thyme, one part.
Alcohol, four parts.

At the end of three days, distil one half. (BRUNS.; HANN.; HASS.; PALAT.; POL.; BORUS.; SAX.; WIRTEM.; PIDERIT)

LIPP. prescribes one part of herb and six parts of alcohol; one half to be distilled.—DAN.; and FULD.—one of herb and six of alcohol; two thirds to be distilled, and re-distilled with one half part of fresh herb.—HERBIP.—one pound and a half of herb and

ten pounds of alcohol; the product to be re-distilled with three bunces of herb.—
SARD.—one part of herb and four parts of alcohol; the product to be re-distilled with fresh herb.

R. Fresh Herb of Wild Thyme, one part.
Alcohol (0.910), four parts and a half.
Water, six parts.

Macerate for twenty-four hours, and distil four parts (0.913). (AUSTR.)

VAN MONS prescribes one part of herb, four parts of proof spirit, and twelve of water; three and a half to be distilled.—
SWEDIAUR—twelve of herb, six of alcohol, and two of water; six to be distilled; the product to be poured upon twelve of fresh herb with two of water, and six re-distilled.

COMPOUND SPIRIT OF WILD THYME.

Aqua benedicta Serpylli seu Serpylli composita. (SARD.; WIRTEM.; SPIELMANN)

R. Flowering Herb of Wild Thyme, four handfuls.
Flowers of Chamomile, } of each,
Tops of Savory, } one handful.
Thyme, }
Wine, six pounds.

Macerate for three days, and distil one half. Infuse in the product,

Sweet Flag, } of each,
Cassia Lignea, } half an ounce.
Cinnamon, }
Mace, }

Digest for twenty-four hours, and distil to dryness. (SARD.)

R. Flowering Herb of Wild Thyme, two pounds.
Savory, } of each,
Thyme, } two
Flowers of Chamomile, } ounces.
Cinnamon, three ounces.
Mace, half an ounce.
White Wine, eighteen pounds.

Digest for two days, and distil slowly eight pounds. (WIRTEM.)

R. Root of Zedoary, } of each, one dram.
Galanga, }
Peony, one ounce and a half.
Herb of Wild Thyme, six handfuls.
Fennel, four handfuls.
Vervain, } of each,
Balm, } two
Flowers of Lime, } handfuls.
Lily of the Valley, }
Lavender, }
Rosemary, }

Cubebs, one dram.
Cinnamon, half a dram.
Mace, } of each, two drams.
Nutmegs, }
Misteloe, one ounce and a half.
Wine, one hundred and ninety-two ounces.

Distil. (SPIELMANN)

It is cephalic, cardiac, stomachic, carminative, and emmenagogue. — Dose, from half an ounce to one ounce.

TILIA EUROPÆA, LINN.—LIME TREE, LINDEN TREE.

SYNONIMES:—Tilleul, FR.; Linde, DUT.; and GERM.; Ljpa, BOHEM.; Lind, DAN.; and SWED.; Tilo, SPAN.; Tiglio, ITAL.; Lipo, POL.

AMST.; ANT.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; OLDEN.; POL.; SAX.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A common tree of Europe. (*Polyandria Monogynia*, LINN.; *Tiliaceæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 281*)

The flowers (*Flores Tiliæ*) are employed. They are of a yellowish colour. Their odour is agreeable, but is removed in a great measure by desiccation. Their taste is sweet and mucilaginous.

They are anti-spasmodic, and slightly

diaphoretic, particularly when the aqueous vehicle is abundant and very warm.

CONSERVE OF LIME FLOWERS.

Conserva Florum Tiliæ. (SARD.)

R. Lime Flowers, one part.
Sugar, boiled *à la plume*, three parts.
Preserve.

INFUSION OF LIME FLOWERS. (GALL.; FORM. HÔP. FR.; DISP. CHA. PARIS; RATIER)

R. Lime Flowers, two drams.
Boiling Water, two pounds.
Infuse, and strain. (FORM. HÔP. FR.; RATIER)

R. Infusion of Lime Flowers, two pounds.
Orange-flower Water, two ounces.
Mix. (RATIER)

R. Lime Flowers, two drams.
Boiling Water, two pounds.
Infuse for a quarter of an hour; then strain, and add
Syrup of Marshmallows, one ounce.
Mix. (GALL.)

DISP. CHA. PARIS prescribes two drams each, of lime flowers and of liquorice root, and one pound and a half of boiling water.

It is slightly anti-spasmodic.—The first infusion is generally acidulated with oxymel or with tartaric syrup, and sweetened with syrup of gum, and sometimes with syrup of poppy. The second is recommended by CHAUSSIER in nervous affections with hysteric spasms. Two drams of HOFFMANN's liquor is frequently added. Both should be taken hot.

LIME-FLOWER WATER.

Aqua Tilie. (GALL.; FULD.; GENEV.; OLDEN.; PALAT.; SAX.; SARD.; HERBIP.; SWEDIAUR)

R. Lime Flowers, two parts.
Water, a sufficient quantity.

Distil twenty parts. (SWEDIAUR)

PALAT. prescribes one part of flowers and three parts of water; one part and a half to be distilled.—GALL.; and SARD.—one of flowers and four of water; one half to be distilled.—HERBIP.—one of flowers and six of water.—SAX.—one of flowers and twenty of water; two thirds to be distilled.—OLDEN.—one of flowers and thirty of water; twenty to be distilled.—GENEV.—four of flowers and a sufficient quantity of water; six to be distilled.

R. Lime Flowers, three parts.
Water, twelve parts.

Enclose the flowers in a bag suspended in the centre of a cucurbit in such a manner that it may not touch the water, and distil six parts without ebullition.

Dose, from two to four ounces.

TORDYLIUM OFFICINALE, LINN.

SYNONIME :—Séséli de Crète, Fr.

BRUNS.; HISP.; WIRTEM.; BERGIUS;
MURRAY and GMELIN; ZARDA.

A plant of the south of Europe. (*Pentandria Digynia*, LINN.; *Umbellifera*, JUSS.; *Fig. Zorn, Ic. pl. t. 278*)

The root and seeds are employed.

The root (*Radix Tordylii seu Seseli Cre-tici seu montani*) is long, slender, fibrous, and white.

The seeds are long, nearly flat, rough to the touch, marked by three projecting lines, and furnished with a white border. It has an aromatic, and a similar though feeble taste.

It is excitant.

TORMENTILLA ERECTA, LINN.—TORMENTIL.

SYNONIMES :—Tormentille, Fr.; Tormen-tillwurz, Blutkraut, Feigwarzakraut, Sie-benfingerkraut, GERM.; Natrzniak, Trzene korenj, BOHEM.; Tormentilla, ITAL.; PORT.; and SPAN.; Korze ziele pepawy, POL.

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; SPIELMANN; ZARDA.

A plant of Europe. (*Icosandria Poly-*

gynia, LINN.; *Rosacea*, JUSS.; *Fig. Zorn, Ic. pl. t. 358*)

The root (*Radix Tormentillæ sylvestris seu Heptaphylli seu Septifolii*) is employed. It is nodose, of the thickness of the finger, oblong, and furnished with numerous fibres. It is of a brown-red colour externally, and pink or yellowish-brown, sometimes spotted internally; it is inodorous, but of a very astringent and bitterish taste.

It is astringent.—Dose, from half a dram to one dram, of the powder.

ASTRINGENT POWDER.

Pulvis tormentillæ compositus. (GENEV.)

R. Terra sigillata, six ounces.

Cinnamon,	four ounces.
Tormentil,	} of each, three ounces.
Gum Arabic,	
Long Pepper,	
	half an ounce.

Mix.—A forty-eighth of opium is sometimes added.

EXTRACT OF TORMENTIL.

Extractum Tormentillæ. (AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; HISP.; LIPP.; LUSIT.; SAX.; SARD.; WIRTEM.; HERBIP.)

R. Root of Tormentil,	one pound.
Spring Water,	six pounds.

Digest for four days in a warm place, and then boil a little. Evaporate the expressed liquor to the proper consistence. (BRUNS.; PALAT.; WIRTEM.)

SARD. directs the liquor to be boiled down to two thirds before straining and evaporating.

R. Root of Tormentil,	one part.
Water,	eight parts.

Macerate for twenty-four hours, then boil for a quarter of an hour, and express strongly; boil the residue with

Water,	four parts.
--------	-------------

Mix, and after twenty-four hours' repose, evaporate to the proper consistence. (ANT.; LIPP.)

BATAV. requires twenty-four hours' digestion, and two hours' ebullition with ten parts of water, and a second ebullition with eight parts of water.—AMST.—an ebullition first with twenty and then with fifteen parts of water, to be reduced each time to one half.—HISP.—twenty-four hours' digestion, and two hours' ebullition, first with ten parts of water, and then with a fresh quantity, to be reduced each time to one half.

R. Root of Tormentil,	any quantity.
Water,	a sufficient quantity.

Exhaust the root by successive ebullitions; then mix the decoctions, and after decantation, evaporate to the consistence of an extract. (AUSTR.)

HERBIP. requires only one ebullition, but prolonged for two hours.—BAVAR.—one part of root to be first boiled in eight and then in four parts of water, after twenty-four hours' previous maceration.

Dose, from one scruple to half a dram.

ASTRINGENT PILLS. (CADET DE GASSI-COURT)

R. Extract of Tormentil,	one dram.
Mastic,	two drams.
Gum Arabic,	half an ounce.

Terebinthina cocta,
a sufficient quantity.

Make pills of four grains each. They are recommended by QUARIN towards the end of gonorrhœa.—Dose, five or six, morning and evening.

DISTILLED WATER OF TORMENTIL.

Aqua Tormentillæ. (PALAT.)

R. Root of Tormentil,	one part.
Water,	two parts.

Distil one part and a half.

TRAUMATIC ESSENCE.

Essentia traumatica. (WIRTEM.; SPIEL-MANN)

R. Root of Tormentil,	} of each, half an ounce.
Herb of Sanicle,	
——— Sundew,	} of each, two drams.
Flowers of St. John's-wort,	
——— Red Roses,	
Dragon's Blood,	} of each, three drams.
Mummy,	
Catechu,	
Myrrh,	
Red Sanders,	two drams.
Nutmegs,	one dram.
Alcohol,	sixteen ounces.

Digest for three days at a moderate heat; then express and filter. (WIRTEM.)

R. Root of Tormentil,	half an ounce.
Flowers of Red Roses,	} of each, two drams.
——— St. John's-wort,	
Red Sanders,	} of each, one ounce.
Mummy,	
Dragon's Blood,	} of each, half an ounce.
Catechu,	
Myrrh,	} one dram.
Nutmegs,	
Alcohol,	twenty ounces.

Digest, and filter. (SPIELMANN)

It is astringent, and is recommended to arrest internal hæmorrhages and diarrhœa.—Dose, from fifty to sixty drops.

TINCTURE OF TORMENTIL.

Tinctura Tormentillæ. (BELG.)

R. Root of Tormentil,	one part.
Alcohol (15 degrees),	eight parts.

Macerate for six days without heat; then express and filter.

DECOCTION OF TORMENTIL.

Decoctum Radicis Tormentillæ. (NIE-MANN)

R. Root of Tormentil,	half an ounce,
Spring Water,	one pound and a half.

Boil, and strain.—Dose, two spoonfuls every two hours.

ASTRINGENT DECOCTION.

Decoctum adstringens. (SAUNDERS; SWEDIAUR; VAN MONS)

R. Root of Tormentil,	}	of each,
Oak Bark,		two drams.
Pomegranate Bark,		
Water,	}	of each,
Milk,		one pound.

Boil for a quarter of an hour, adding towards the end,

Cinnamon, two drams.

Strain when cold. (SWEDIAUR; VAN MONS)

R. Root of Tormentil,	}	of each,
Dried Flowers of Pomegranate,		half
Pomegranate Bark,		an ounce.
Cinchona,		one ounce.
Spring Water,		a sufficient quantity

to obtain two pounds of strained decoction. (SAUNDERS)

It is recommended in diarrhoea and blennorrhœa.—Dose, two ounces, three or four times a day. By leaving out the cinnamon, this decoction may also be employed in the form of lavement or injection.

ASTRINGENT PTISAN. (CADET DE GASSICOURT; PIERQUIN; SPIELMANN)

R. Root of Tormentil,	six drams.
Spring Water,	twelve pounds.

Boil for half an hour, adding towards the end,

Root of Great Comfrey,	six drams.
——— Liquorice,	half an ounce.

Strain. (SPIELMANN)

PIERQUIN prescribes three ounces of tormentil, thirty-four pounds of water, three ounces of comfrey, and two ounces of liquorice.

R. Root of Tormentil,	}	of each,
——— Dog-grass,		half an ounce.
Pomegranate Bark,	}	of each,
Barberry,		two drams.
Sumach,		
Hartshorn Shavings,		three drams.
Water,		four pounds.

Boil down to three pounds, and add to the strained liquor,

Honey, two ounces.

Strain. (PIERQUIN)

R. Boiling Water,	two pounds.	
Dried Petals of Red Roses,	half an ounce.	
Root of Tormentil,	}	of each,
——— Bistort,		one ounce.

Boil the roots and infuse the petals; then add to the strained liquor,

Syrup of Pomegranate, one ounce and a half.

——— Gum, one ounce.

Dilute Sulphuric Acid, twenty drops.

Mix. (CADET DE GASSICOURT)

ASTRINGENT GARGLE. (SAUNDERS)

R. Root of Tormentil,	two ounces.
Water,	a sufficient quantity

to obtain two pounds of strained liquor, after half an hour's boiling. Add

Honey of Roses, one ounce and a half.

Sulphuric Acid, ten drops.

Mix.

TORNASOLIS CÆRULEA.—LITMUS, TURNSOL.

SYNONIMES : — *Lacmus*, *Lacca cærulea*, *Lacca musica*.—Tournesol, FR.; Blaues Lack, Lackmus, GERM.; Lackmuss, DUT.

A colouring substance, of which there are two kinds met with in commerce :

1. *Cake or Stone Litmus*.

DUBL.; AUSTR.; BELG.; BRUNS.; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

This is in small straight parallelopipedons of a blue colour, prepared with several species of lichens reduced to powder

mixed with subcarbonate of potass and chalk, and fermented with human urine.

2. *Litmus cloths*; *Becetta*, *Nezetta*.

BRUNS.; WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GMELIN.

These are old rags saturated with the juice of the tops of the *Croton tinctorium*, LINN.; (*Monœcia Monadelphia*, LINN.; *Euphorbiaceæ*, JUSS.), a plant of the south of Europe (*Fig. Lob. Ic. 26*), then dried and exposed to the vapour disengaged from a mixture of urine and quick lime till they acquire a violet-blue colour.

TRAGOPOGON PRATENSE, LINN.—SALSEFI.

SYNONIMES: — Salsifis, Barbe de bouc, FR.; Boksbart, Wiesenbocksbart, GERM.; Salsifi, SPAN.; Salsifica, ITAL.; Sersifim, PORT.

BRUNS.; GALL.; FERR.; WIRTEM.

A plant common to the meadows of Europe. (*Syngenesia Polygamia Æqualis*, LINN.; *Synanthereæ*, CASS.; *Fig. Zorn, Ic. pl. t. 179*)

The root (*Radix Tragopogi* seu *Barba Hirci*) is employed. It is long, cylindrical, and whitish. It has a sweet taste.

The *Tragopogon porrifolium*, LINN.; (*Fig. Œd. Fl. Dan. t. 797*) is frequently substituted for it.

It sometimes enters into pectoral decoctions.

TRAPA NATANS, LINN.—WATER CHESNUT, WATER CALTROPS.

SYNONIMES: — Macre, Macre flottante, Châtaigine d'eau, Noix d'eau, FR.; Was- sernuss, GERM.; Svømmende hornøed, DAN.; Castana de aqua, SPAN.; Water- nooten, DUT.; Tribulo aquatico, ITAL.; Orzech wodoje, POL.; Arolho dos char- cos, PORT.; Vatnøet, SWED.

WIRTEM.; BERGIUS; MURRAY and GME- LIN; SPIELMANN.

A plant of Europe, common in stagnant waters.

The fruit used formerly to be employed as astringent, and the leaves as resolvent. Neither are used at the present day.

TREMELLA AURICULA JUDÆ, PERS.—JEW'S EAR.

SYNONIMES: — Oreile de Judas, FR.; Hol- derschwämmchen, Juda's ohr, GERM.; Bezowa hauba, BOHEM.; Judasoor, DUT.

BRUNS.; GALL.; LUSIT.; WIRTEM.; HER- BIP.; BERGIUS; MURRAY and GMELIN; SPIELMANN; ZARDA.

A mushroom (*Auricula Judæ* seu *Fungus Sambuci*) growing on decayed trees. (*Fig. Zorn, Ic. pl. t. 500*)

It is in the form of an ear and of a green- ish grey colour.

It used formerly to be employed in an- gina, boiled in milk or vinegar.

TRIFOLIUM PRATENSE, LINN.—PURPLE TREFOIL.

SYNONIMES: — Trèfle des près, FR.; Ro- ther Wiesenkleee, GERM.; Røedklover, DAN.; Røedkøeping, SWED.

GALL.; WIRTEM.

A common plant of Europe. (*Diadel- phia Decandria*, LINN.; *Leguminosæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 93*)

The flowering herb (*Herba cum Floribus Trifolii pratensis* seu *purpurei*) is em- ployed. It consists of trifoliate leaves, with oval, entire, rough leaflets, frequently marked with a white or black spot, and of red or purple flowers, disposed in rounded, obtuse spikes, surrounded by membranous bractæ.

It is said to be slightly laxative.

TRIGONELLA FÆNUM GRÆCUM, LINN.—FENUGREEK.

SYNONIMES:—Fenu-grec, FR.; Fœnugræk, Kuhkornklee, Bockshorn, GERM.; Bukkehorn, Fœnugraec, DAN.; Alholva, SPAN.; Fene-grick, DUT.; Pieno Greco, ITAL.; Fengræk, POL.; Fenogregio, PORT.; Fenugrek, SWED.

LOND.; AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOUT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.

A plant of Europe. (*Diadelphia Decandria*, LINN.; *Leguminosæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 116*)

The seeds (*Semen Fenugræci*) are employed. They are nearly quadrangular, flattish, obtuse at both extremities, and marked with an oblique groove. They are of a deep yellow or brown colour; their odour is strong, sweetish, and disagreeable

when bruised; their taste is bitter, farinaceous, and mucilaginous.

MUCILAGE OF FENUGREEK. (SARD.)

R. Seeds of Fenugreek, one part.
Water, ten parts.

Macerate for twelve hours; then boil, and express.

MUCILAGINOUS OIL. (FEE)

R. Seeds of Fenugreek, bruised, eight ounces.
Linseed Oil, two pounds.

Infuse for ten days, and strain.

RESOLVENT MEALS. (GALL.)

R. Fenugreek Seeds, }
Beans, } of each,
Tares, } an equal part.
Lupins, }
Pulverize, and mix.

TRIOSTEUM PERFOLIATUM, LINN.—FEVER-ROOT, BASTARD IPECACUANHA, TINKER-WEED.

AMER.; BRUNS.; COXE.

A plant of North America. (*Pentandria Monogynia*, LINN.; *Caprifoliaceæ*, JUSS.; *Fig. Dill. Eth. t. 293, f. 378*)

The bark of the root is employed. It is purgative and sometimes emetic.

TRIPOLITIANA TERRA.—TRIPOLI.

SYNONIMES:—*Tripela*.—Tripoli, FR.; Tripel, Trippelerde, GERM.

BRUNS.; WIRTEM.; GUIBOUT.; SPIELMANN.

A hard substance, rough and dry to the

touch, the colour of which varies from white to yellow, to red, to green, and to brown. That tripoli is most esteemed which is called Venetian, and which comes from Corfu; it is schistose, porous, and of a yellowish-red colour.

TRITICUM REPENS, LINN.—DOG-GRASS, WHEAT-GRASS, QUICK-GRASS, COUCH-GRASS.

SYNONIMES:—Chludent, FR.; Quecke, Queckengras, Graswurzel, Hundsquecken, GERM.; Peyr, Trawa, BOHEM.; Hun-

degræs, Kroppegræs, Qvikketæan, DAN.; Grama, SPAN.; Kueekgras, Hondsgas, DUT.; Gramigna, Caprinella, Dente cæ

nino, ITAL.; Psia partza, POL.; Grama canina, PORT.; Pyrei, RUSS.; Qwickrot, SWED.

AMER.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOUT.; PIDERIT.; SPIELMANN.; ZARDA.

A very common plant of Europe. (*Triandria Digynia*, LINN.; *Gramineæ*, JUSS.; *Fig. Hist. Gram.* 2, p. 17, t. 21)

The creeping shoots, improperly called roots (*Radix Graminis* seu *Graminis canini* seu *albi*), are employed; they are long, slender, nodose, articulated, smooth; whitish or yellowish-white, inodorous, of a sweet and slightly saccharine taste when fresh; and insipid or very feebly astringent when dried.

In the fresh state, dog-grass furnishes a mucoso-saccharine matter, which renders it demulcent, though in a very inconsiderable degree; when dried, this matter, which is contained in the medulla, disappears, and then the envelope only remains, which communicates to water a slightly astringent quality, particularly when the ebullition is prolonged.

The decoctions of dog-grass act principally from the water contained in them, and on this account the popular employment of them is of use.

SPECIES FOR PTISAN. (FULD.)

R. Root of Dog-grass, four ounces.
—— Scorzonera, two ounces.

Cut, and mix.

JUICE OF DOG-GRASS.

Succus Tritici. (GALL.; HASS.)

R. Fresh Herb of Dog-grass, sixteen parts.

Bruise it in a stone mortar, adding gradually,

Water, one part.

Express the juice, set it aside, and filter it without heat through bibulous paper.

HASS. prescribes the root instead of the herb.

EXTRACT OF DOG-GRASS.

Extractum seu Mellago Graminis seu Tritici repentis. (AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; DAN.; FERR.; FULD.; HANN.; HASS.; LIPP.; OL-

DEN.; POL.; BORUS.; SAX.; HERBIP.; PIDERIT.; SWEDIAUR.; VAN MONS)

R. Fresh Root of Dog-grass, six pounds.

Having cut it finely, enclose it in a linen bag, and express strongly; boil the residue for an hour with

Water, eight pounds,

and express again. Mix the liquors, set aside for twelve hours, and having strained through flannel, evaporate to the consistence of thick syrup. (LIPP.)

R. Root of Dog-grass, bruised, } of each,
Water, } one part.

After twelve hours' infusion, add

Hot Water, two parts,

then stir till cold, and express. Heat the residue for several hours with twice its weight of water, at a temperature of 45 or 50 degrees of R., and express again. Mix the liquors; then clarify, and evaporate to the consistence of an extract. (VAN MONS)

R. Fresh Root of Dog-grass, twelve pounds.

Bruise it in a mortar, adding

Water, six pounds.

Express the juice, boil it for a few moments, and evaporate it, when strained, to the consistence of honey. (DAN.; OLDEN.; POL.; BORUS.; SAX.; SWEDIAUR)

R. Fresh Root of Dog-grass, one pound.
Spring Water, eight pounds.

Boil till reduced to one half. Strain the decoction, and evaporate it to the consistence of an extract. (HASS.; PIDERIT)

AUSTR.; and HERBIP. direct the root to be boiled in a sufficient quantity of water; the decoction to be set aside, decanted and evaporated to the consistence of thick honey.

R. Root of Dog-grass, one part.
Boiling Water, eight parts.

Boil slowly for a quarter of an hour, and express; boil the residue with

Water, four parts,

and express again. Mix the decoctions, and having decanted after sufficient repose, evaporate to the consistence of an extract. (ANT.; HANN.)

BATAV. prescribes twenty-four hours' maceration and two hours' boiling with ten parts of water; then a second ebullition of two hours with other eight parts of water. —AMST. directs the root to be boiled first with twenty then with fifteen parts of water till reduced each time to one half. —BAVAR. —twenty-four hours' maceration of one part of root in four parts of water; the addition of four other parts of water, and an ebullition of twenty-four hours: then a second ebullition of the residue with four

other parts of water, and the mixed liquors evaporated to the consistence of honey, or of solid extract.—FULD.—six days' maceration of one pound of root in six pounds of water, then an hours' ebullition, and expression; the ebullition of the residue with three pounds of water: the repetition of this till the water comes off colourless and insipid, and the final evaporation of the mixed liquors to the consistence of honey.

This extract is esteemed resolvent and diuretic.—Dose, a spoonful.

DECOCTION OF DOG-GRASS.

Decoctum Graminis. (ANT.; GALL.; CAST. BORUS.; CADET DE GASSICOURT.; RATIER)

R. Root of Dog-grass, sliced, eight ounces.
Water, four pounds.

Boil down to three fourths, and strain.—ANT.—prescribes two ounces of dog-grass and two pounds of water; to be boiled down to one half.

R. Root of Dog-grass, one ounce.

Boil it for a short time in a sufficient quantity of water, and reject this; then bruise the root, and boil it with

Water, two pounds eight ounces, till only two pounds remain. Add towards the end,

Grated Liquorice, two drams.

Strain when cold. (GALL.; CADET DE GASSICOURT.; DISP. CHA. PARIS; RATIER)

This last decoction is only emollient; the first is slightly tonic.

COMPOUND DECOCTION OF DOG-GRASS.

Decoctum Graminis compositum. (HISP.; SAUNDERS)

R. Root of Dog-grass, } of each,
Madder, } half an ounce.
Water, three pounds.

Boil down to two thirds; and add towards the end,

Grated Liquorice. one dram.

Express, and add

Nitrate of Potass, one dram.

Dissolve. (HISP.)

R. Root of Dog-grass, } of each,
Dandelion, } two ounces.
Water, a sufficient quantity,

to obtain one pound and a half of strained decoction. Boil for half an hour, and add

Sulphate of Potass, half an ounce.

Syrup of Fumitory, two ounces.

Spirit of Nitric Æther, one dram.

Mix. (SAUNDERS)

It is excitant, and deobstruent.

ALKALINE DIURETIC DECOCTION.

Decoctum diureticum alkalinum. (VAN MONS)

R. Saturated Decoction of Dog-grass
and of Dandelion, three pounds.
Subcarbonate of Soda, three drams.
Nitrate of Potass,

one dram and a half.

Honey, one ounce and a half.

Dissolve.—It is excitant, and is recommended in lacteal metastasis and in anasarca.—Dose, a cupful, every two hours.

TROPEOLUM MAJUS, LINN.—INDIAN CRESS, NASTURTIIUM.

SYNONIMES:—Capucine, Cresson des Indes, Cresson du Mexique, FR.; Indianische kresse, Capucinerkresse, GERM.; Indiansk karse, DAN.; Capcichinos, SPAN.; Spaensche kers, DUT.; Capucino, ITAL.; Mastraceo da Paro, PORT.; Indianisk kresse, SWED.

BRUNS.; GALL.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A plant of South America. (*Octandria Monogynia*, LINN.; *Tropæoleæ*, Cand.; Fig. Flore Medic. II. 96)

The herb and flowers are used.

The herb (*Herba Nasturtii Indici* seu *Cardami majoris*) consists of a cylindrical, glabrous stem, and of alternate, plane, rounded leaves; they present five obscure lobes, are peltate, and supported upon long foot stalks.

The flowers consist of five petals of an orange-yellow or scarlet colour.

The taste and odour are strong and similar to those of cress.

It is excitant, and reputed anti-scorbutic.

TUSSILAGO.

Two species of this genus of plants are employed in medicine:

1. *Tussilago Farfara*, LINN.—Coltsfoot.

SYNONIMES:—Pas d'âne, FR.; Huflattich, GERM.; Konske kopyto, BOHEM.; Hovvurt, Hovblad, DAN.; Tusilago, SPAN.; Hoefblad, DUT.; Tosilagine, ITAL.; Podbial, POL.; Tosilagem, PORT.; Hæsthof, SWED.

LOND.; EDINB.; DUBL.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HAMB.; HANN.; HASS.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; PARIS.; PIDERIT.; SPIELMANN.; ZARDA.

A plant common in all parts of Europe. (*Syngenesia Polygamia superflua*, LINN.; *Synanthereæ*, CASS.; Fig. Zorn, *lc. pl. t. 64*)

The root, leaves, and flowers are employed.

The root (*Radix Farfaræ* seu *Tussilaginis* seu *Bechii* seu *Ungulæ Caballinæ*) is long, thin, fibrous, whitish, inodorous, and insipid.

The leaves are rounded, somewhat heart-shaped at the base, angular, dentated, of a deep green colour and smooth above, and tomentose beneath; they have long foot-stalks.

They are inodorous, and have a mucilaginous and slightly bitter, and acerb taste.

The flowers are yellow and have a pleasant taste.

It is slightly excitant and pectoral.

PECTORAL SPECIES. (CAST. BORUS.; WURCEB.)

R. Leaves of Coltsfoot,	two ounces.
Root of Liquorice,	one ounce.
Flowers of Red Poppy,	of each,
Anise,	two drams.

Cut and bruise.

CONSERVE OF COLTSFOOT.

Conserva Farfaræ. (PALAT.; WIRTEM.; VAN MONS)

R. Flowers of Coltsfoot,	one part.
White Sugar,	two parts.

Beat together.

It is recommended in catarrhal cough.—Dose, two drams, four times a day.

LOHOCH OF COLTSFOOT.

Lohoch de Farfara. (WIRTEM.; PIDERIT.; SPIELMANN)

R. Fresh Root of Coltsfoot, any quantity.

Boil it in water; then bruise it, and pass it through a hair sieve. Boil the pulp with twice its weight of honey.

SYRUP OF COLTSFOOT.

Syrupus Tussilaginis. (BRUNS.; GALL.; SARD.; WIRTEM.)

R. Fresh Flowers of Coltsfoot,	four pounds.
Boiling Water,	eight pounds.

Infuse for twelve hours, and express slightly: to the decanted liquor, add

White Sugar,	twice the quantity.
--------------	---------------------

Dissolve in a water-bath. (GALL.)

BRUNS.; and WIRTEM. prescribe one pound and a half of flowers, two pounds of boiling water, twenty-four hours' maceration, and sixteen ounces of sugar to nine of the strained liquor.

SARD. allows either an equal part each, of depurated juice and of simple syrup to be mixed together, or two parts of infusion with one part of white sugar.

WATER OF COLTSFOOT. (PALAT.)

R. Flowers of Coltsfoot,	one part.
Water,	three parts.

Distil one part and a half.

RESUMPTIVE ELECTUARY.

Electuarium resumptivum. (WIRTEM.; SPIELMANN)

R. Root of Liquorice,	} of each,
Herb of Ground Ivy,	
Veronica,	} half an ounce.
Boiling Water,	
Macerate for twelve hours. Boil in the strained liquor,	sixty ounces.
Fresh Root of Colts-	} of each,
foot,	
Corinth Raisins,	
	sixteen ounces.

Pass the softened root and raisins through a silk-sieve; then emulsionize the decoction with

Sweet Almonds, blanched,	seven ounces.
--------------------------	---------------

Mix the pulp with the emulsion, and boil the whole with

White Sugar, thirty-two ounces.
Then add
Lemon Peels, three.
Powdered Cinnamon, half an ounce.
Cloves, three drams.

Mix.—Dose, from half an ounce to an ounce, two or three times a day in obstinate cough.

2. *Tussilago Petasites*.—Butterbur.

SYNONIMES:—Pétasite, FR.; Pestilenzwurz, Pestwurzmönnlein, Neunkraftstrosshuf, GERM.; Demesyl, Wetssjkonkske kopyto, BOHEM.; Storbladet hestihoc, DAN.; Petasites, Lombrerera, SPAN.;

Poddeblad, DUT.; Petasite, ITAL.; Lipieznik, POL.; Pestilenzrot, SWED.

BAVAR.; BRUNS.; HISP.; GALL.; LIPP.; WIRTEM.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A plant common to most places in the north of Europe. (*Fig. Zorn, Ic. pl. t. 58*)

The root (*Radix Petasitidis* seu *Petasitis majoris* seu *vulgaris* seu *rubentis*) is employed. It is branching, fibrous, and of the thickness of the finger, brown externally, and yellow or red internally. It has a strong and disagreeable odour; and a bitter and repugnant taste.

ULMUS.—ELM.

Two species of this genus are employed in medicine:

1. *Ulmus fulva*, MICH.—Rough-leaved Elm Tree, Red Elm, Slippery Elm.

SYNONIME:—Orme d'Amerique, FR.

AMER.; COXE.

A tree of North America. (*Pentandria Digynia*, LINN.; *Amentaceæ*, JUSS.)

The inner bark is employed. It is emollient and mucilaginous.

2. *Ulmus campestris*, LINN.—Elm Tree.

SYNONIMES:—Orme pyramidal, FR.; Ulme, Rüster, GERM.; Alm, DAN.; and SWED.; Olmo, ITAL.; PORT.; and SPAN.; Olm, DUT.; Kora wiazowa, POL.

LOND.; DUBL.; EDINB.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FENN.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; BRUGNATELLI; COXE; SPIELMANN.

A large tree of Europe.

The inner bark of the branches (*Cortex ulmi interior*) is employed. It is smooth, thin, of a reddish-brown colour, and inodorous. It has a bitter, astringent, sub-acrid, and very mucilaginous taste, after it has been chewed for some time. It has been extolled in ascites, cutaneous diseases, scrofula, scurvy, and intermittent fevers.

EXTRACT OF ELM. (AUSTR.)

R. Inner Bark of Elm, any quantity.
Water, a sufficient quantity.

Exhaust the bark by several successive ebullitions; then mix the liquors, and having decanted them, evaporate in a water-bath.

INFUSION ON ELM.

Infusum Ulmi. (AMER.; COXE; NIEMANN)

R. Elm Bark, sliced, one ounce.
Boiling Water, one pint.

Macerate for twelve hours in a covered vessel, occasionally agitating, and strain.

DECOCTION OF ELM BARK.

Decoctum Ulmi campestris. (LOND.; DUBL.; EDINB.; GENEV.; COXE; NIEMANN; SAUNDERS; SWEDIAUR; PH. SYPH.)

Decoctum Ulmi. (LOND.; DUBL.)

R. Fresh Elm Bark, bruised, four ounces.
Water, four pints.
Boil to two pints and strain.

Decoctum Ulmi campestris. (EDINB.)

R. Fresh Elm Bark, bruised, four ounces.
Water, five pounds.
Boil down to two pounds and a half, and strain.

The other pharmacopœias give the same formula as LOND.

It is recommended in leprosy and other cutaneous disorders.—Dose, from four to eight ounces twice or thrice a day.

DIAPHORETIC DECOCTION.

Decoctum diaphoreticum. (BRERA)

R. Elm Bark, two ounces.

Dock, one ounce.
 Water, a sufficient quantity
 to obtain three pounds of strained liquor.
 It is prescribed in chronic exanthemata.

URTICA.—NETTLE.

Three species of this genus are used in medicine:

1. *Urtica urens*, LINN.—Stinging Nettle.

ANT.; BAVAR.; BRUNS.; GALL.; FERR.; LIPP.; BRUGNATELLI.; MURRAY and GMELIN.

A very common plant of Europe. (*Monæcia Tetrandria*, LINN.; *Urtica*, JUSS.; *Fig. CEd. Fl. Dan. t. 739*)

The herb, flowers, and seeds are employed.

The herb (*Herba Urticæ minoris*) presents a branching stem and oval, deeply-toothed leaves bristled with extremely acute hairs. It has a bitterish taste.

The flowers are green and assembled in short axillary racemes, nearly sessile.

The seeds are oval, shining, and of a light brown colour.

2. *Urtica dioica*, LINN.—Common Nettle.

SYNONIMES:—Ortie dioïque, FR.; Grosse Brennessel, GERM.; Kopriwa paliwa nefwetssj, BOHEM.; Stor brændenælde, DAN.; Ortiga mayor, SPAN.; Groote brandenatel, DUT.; Ortica, ITAL.; Pokrzywa, POL.; Ortiga major, PORT.; Brænn ætsta, SWED.

AMST.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; GENEV.; HASS.; WIRTEM.; HERBIP.; BERGIUS.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of Europe. (*Fig. Zorn, Ic. pl. t. 465*)

The root and herb are employed.

The root (*Radix Urticæ majoris*) is long, ligneous, and branching; of a brownish colour without, and whitish within.

The herb consists of a quadrangular stem furnished with petiolated, cordiform, pointed, serrated leaves, covered with prickly hairs.

3. *Urtica Pilulifera*, LINN.

BRUNS.; WIRTEM.; MURRAY and GMELIN.; SPIELMANN.

A plant of the south of Europe. (*Fig. Blackw. Herb. tab. 321*)

The seeds (*Semen Urticæ Romanæ seu Hispanica seu piluliferæ*) are employed. They are rounded, flattish, of a blackish-brown colour, somewhat resembling linseed but smaller.

EXTRACT OF NETTLE.

Inspissamentum Urticarum. (HISP.)

R. Leaves and Stalks of Nettle,
 any quantity.

Bruise them in a stone mortar; then express the juice, and evaporate it when defecated, to the consistence of thick honey, in a water-bath.

SYRUP OF NETTLE.

Syrupus de Urtica seu Urticarum. (HISP.; GALL.)

R. Juice of Nettle, depurated } of each,
 by boiling and strained, } an
 White Sugar, } equal part.

Boil slowly to the consistence of syrup. (GALL.)

HISP. directs equal parts of nettle juice and of white sugar, to be clarified and boiled to a syrup with a sufficient quantity of water.

This syrup is considered diuretic.

VACCINIUM.

Three species of this genus are employed in medicine:

1. *Vaccinium Myrtillus*, LINN.—Common Blueberry.

SYNONIMES: Airelle, Myrtille, Raisin des bois, FR.; Heidelbeere, GERM.; Myrtus, Gahody cerne, Wranj oka, BOHEM.; Blæbar, DAN.; Mirtillo, Arandano,

SPAN.; Blaaw bessen, DUT.; Mirtillo, ITAL.; Borowkie zorna, POL.; Blabær, SWED.

AMST.; BRUNS.; HISP.; GALL.; FENN.; FULD.; LIPP.; ROSS.; SUEC.; WIRTEM.; BERGIUS; BRUGNATELLI; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A small shrub of the north of Europe. (*Octandria Monogynia*, LINN.; *Ericineæ*, JUSS.; *Fig. Zorn, Ic. pl. t. 81*)

The fruit *Baccæ Myrtilli* is employed. It consists of berries of the size of a pea, filled with a deep red juice. They have a sweet, acid, and sub-astringent taste.

These berries are slightly styptic, refrigerant, and anti-scorbutic.

JUICE OF BLUEBERRIES.

Succus e Baccis Myrtillosum. (FULD.; PIDERIT)

R. Ripe Blueberries, any quantity.

Bruise and put them in a cellar for several days; then express the juice, and having left it in the cellar for several days, strain it, and pour it into long necked bottles; pour a little oil upon the surface, and preserve in a cool place.

It is added, as acidulous, to mixtures. It is substituted for vinegar in the north of Europe.

ROB OF BLUEBERRIES.

Rob Myrtillosum seu Vaccinii Myrtilli. (FULD.; LIPP.; SWEDIAUR)

R. Blueberries, any quantity.

Bruise them in a mortar, and having set them aside for several days, express the juice and boil it rapidly to the consistence of honey with one fourth of sugar.

It is used instead of the rob of mulberries in the north of Europe.

SYRUP OF BLUEBERRIES.

Syrupus Myrtilli seu Myrtillosum. (BRUNS.; HISP.; FULD.; PALAT.; SUEC.; WIRTEM.; PIDERIT)

R. Expressed juice of Blueberries, one pound.

White Sugar, two pounds.

Dissolve at a gentle heat. (SUEC.)

HISP. prescribes four pounds of sugar and two pounds two ounces of juice.—BRUNS.; PALAT.; WIRTEM.; HERBIP.; and PIDERIT—sixteen ounces of sugar and ten of juice.—FULD.—an equal part of each.

JOUBERT'S ASTRINGENT SYRUP. (PIERQUIN)

R. Blueberries, thirty drams.

Barberries	} of each,	twenty drams.
Leaves of Sumac,		
Pomegranate Flowers,	} of each, three ounces.	
Medlars,		
Unripe Sorbs,	} of each,	two pounds.
Juice of Pomegranate,		
Quince,		

Boil together. Add when strained,

Sugar, five pounds.

Flavour with

Yellow Sanders,	half an ounce.
or Cinnamon,	two drams.

2. *Vaccinium Oxycoccus*, LINN.—Cranberry.

SYNONIMES.—Canneberge des marais, FR.; Moosbeere, GERM.; Tranebær, DAN.; *Vacernia laganusa*, SPAN.; Veen bessen, DUT.; Ossicocco, ITAL.; Canaberga, PORT.

DAN.; GALL.; FENN.; ROSS.; SUEC.; WIRTEM.; BERGIUS; MURRAY and GMELIN; SPIELMANN.

A small shrub of the north of Europe. (*Fig. CEd. Fl. Dan. t. 80*)

The fruit (*Baccæ Oxycocci*) is employed. It consists of ovoid berries of a red colour sprinkled with purple points; they are filled with a red juice of a very acid taste.

It is refrigerant and anti-scorbutic.

JUICE OF CRANBERRIES. (SUEC.; NIEMANN)

R. Cranberries, any quantity.

Express the juice; then boil it for a few moments, and having strained it, enclose it in well-stopped bottles.

SYRUP OF CRANBERRIES. (SUEC.; NIEMANN)

R. Juice of Cranberries, one pound.

White Sugar, two pounds.

Make a syrup at a gentle heat.

ROB OF CRANBERRIES.

Rob Oxycoccus seu Vaccinii Oxycocci. (ROSS.; SWEDIAUR)

R. Cranberries, any quantity.

Bruise them in a mortar, and set aside for three days; then express the juice and boil it slowly to the consistence of honey, with one fourth of sugar. (SWEDIAUR)

Ross. prescribes one pound of sugar and six pounds of juice.

It is acidulous and refrigerant.

3. *Vaccinium Vitis Idæa*, LINN.—Red Bilberry.

SYNONIMES.—Canneberge ponctuée, FR. Prisselbeere, Mehlbeere, Rothe bernitz

beere, GERM.; Tyttebær, DAN.; Arandona de fruto enearnado, SPAN.; Vösesbessen, DUT.; Vite del Monte Ida, ITAL.; Lingon, SWED.

DAN.; GALL.; FULD.; LIPP.; ROSS.; SUEC.; BERGIUS; GUIBOURT.

A small shrub of the north of Europe and of the Alps. (*Fig. Zorn, Ic. pl. t. 87*)

The fruit (*Baccæ Vitis Idææ*) is employed. It consists of round berries, of a beautiful red colour, and of an acid and bitterish taste.

The leaves were for a long time reputed lithontriptic, and are even now classed among the diuretics. These possess a slight astringency.

ROB OF RED BILBERRIES.

Rob Vitis Idææ seu Baccarum Vitis Idææ.
(FULD.; LIPP.; SAX.; SWEDIAUR)

R. Red Bilberries, any quantity.

Bruise them in a mortar, and set aside for three days; then express the juice, set it apart, and having decanted it, evaporate it the consistence of honey, over a slow fire, with one fourth of sugar. (FULD.; LIPP.; SWEDIAUR)

SAX. prescribes one part of sugar and sixteen parts of juice.

It is acidulous and refreshing.

VALERIANA.

Three species of this genus of plants are used:

1. *Valeriana Celtica*, LINN.—Celtic Nard.

SYNONIMES:—Nard Celtique, FR.; Nardenbaldrian, Alpenbaldrian, GERM.; Espicanardo Celtico, SPAN.

AMST.; ANT.; AUSTR.; BRUNS.; HISP.; GALL.; FERR.; PALAT.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A plant of the Alps. (*Triandria Monogynia*, LINN.; *Valerianææ*, JUSS.; *Fig. Jacq. Collect. t. I. p. 24.*)

The root (*Radix Valerianæ Celticæ* seu *Spicæ Celticæ* seu *Alpinæ*, *Nardus Celtica*) is composed of long fibrils covered with small yellowish scales. It has a strong and agreeable odour, and an acrid, bitter, and aromatic taste.

It is tonic, stomachic, and diuretic.

COMPOUND POWDER OF CELTIC NARD.
(HERBIP.)

R. Celtic Nard, half an ounce.
Sulphate of Zinc, one scruple.
Make a powder.

2. *Valeriana Phu*, LINN.—Garden Valerian.

SYNONIMES:—Grande Valeriane, FR.; Grosser baldrian, GERM.; Koslik velssj, BOHEM.; Stor baldrian, DAN.

BRUNS.; DAN.; HISP.; GALL.; SUEC.;

WIRTEM.; BERGIUS; GUIBOURT; MURRAY and GMELIN.

A plant of the temperate parts of Europe. (*Triandria Digynia*, LINN.; *Valerianææ*, JUSS.; *Fig. Herb. t. 250*)

The root (*Radix Valerianæ majoris*) is employed. It is thick, long, wrinkled, fibrous, of a light-brown colour without, and white within.

It has a nauseous odour, and a disagreeable, bitter, and acrid taste.

3. *Valeriana officinalis*, LINN.—Valerian, Wild Valerian.

SYNONIMES:—Valériane sauvage, FR.; Gemeiner baldrian, GERM.; Koslik obecky, BOHEM.; Baldrian, Cienrod velerian, Vild nardus, Valandsrod, Venderad, Vendmosrod, DAN.; Valeriana, ITAL.; PORT.; and SPAN.; Wilde valerian, DUT.; Kozlki, koztkowy, POL.; Balderjan, RUSS.; Wandelrot, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FERR.; FENN.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; CAST. BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; PIDERIT; SAUNDERS; SPIELMANN; ZARDA.

A plant which grows in all parts of Europe. (*Fig. Flore Medic. VI. 343*)

The root (*Radix Valerianæ minoris seu sylvestris*) is employed. It is yellowish externally and whitish internally, and consists of numerous long fibrils. It is nearly inodorous in the fresh state, but has a peculiar, fetid, and penetrating odour when dried. Its taste is acrid and bitter.

It contains, according to TROMMSDORF, a greenish, volatile oil of a camphorous odour, gum, starch, resin, and a peculiar principle.

It is a very powerful excitant, and is supposed to act in a special manner on the nervous system. It is principally prescribed in nervous affections, and has been employed as a febrifuge.—Dose, of the powder, from half a dram to one ounce, or even more, twice or thrice a day.

I. Preparations which contain Valerian in substance.

ANTI-SPASMODIC SPECIES. (HERBIP.; HUFELAND; PIDERIT)

R. Root of Valerian, three ounces.
Leaves of Orange, two ounces.
Herb of Milfoil, one ounce.

Cut and mix. (HERBIP.; PIDERIT)

R. Root of Valerian, } of each,
—— Avens, }
Herb of Peppermint, } an equal part.
Leaves of Orange, }

Cut and mix. (HUFELAND)

Dose, two tea-spoonfuls; for infusion.

ANTI-SPASMODIC POWDER.

Pulvis nervinus seu anti-spasticus seu epilepticus seu roborans anti-spasmodicus seu Valerianæ aromaticus. (MIL. COPEN.; FORM. HÔP. FR.; FULD.; HAMB.; CAST. BORUS.; AUGUSTIN; BRERA; HUFELAND; NIEMANN; SAINTE-MARIE)

R. Root of Valerian, } of each,
Leaves of Orange, } two drams.

Make a powder. (FULD.)

R. Root of Valerian, twenty parts.
Cinnamon, three parts.

Make a powder. (FORM. HÔP. FR.)

AUGUSTIN prescribes two drams of valerian, one scruple of cinnamon, and one dram and a half of oleosaccharum of valerian.

R. Root of Valerian, one ounce.
Quassia, half an ounce.

Make a powder. (HAMB.)

R. Root of Valerian, one scruple.
Camphor, one grain.

Pulverize. (MIL. COPEN.)

R. Root of Valerian, half a dram.

Cinnamon, five grains.
Oleosaccharum of Valerian, one scruple.

Make a powder. (CAST. BORUS.)

R. Root of Valerian, half an ounce.
Orange Peel, } of each,
White Sugar, } two drams.
Oil of Valerian, six drops.

Make a powder. (HAMB.)

R. Crab's Eyes, } of each,
Hartshorn Shavings, } an equal part.
Root of Valerian, }
Miseltoe, }

Make a powder. (HUFELAND; NIEMANN)

R. Root of Valerian, half an ounce.
—— Peony, two drams.
Hartshorn Shavings, three drams.
Oil of Nutmeg, three grains.
Sugar, three ounces.

Make a powder. (SAINTE-MARIE)

R. Root of Valerian, half an ounce.
Orange Peel, } of each,
Tartrate of Potass, } two drams.

Make a powder. (FULD.)

R. Root of Valerian, one dram.
Leaves of Orange, one scruple.
Muriate of Ammonia, two grains.
Cajeput Oil, four drops.

Make a powder. (NIEMANN)

R. Valerian, half a dram.
Subcarbonate of Magnesia, } of each,
Muriate of Ammonia, } three grains.
Cajeput Oil, two drops.

Make a powder. (AUGUSTIN; NIEMANN)

R. Valerian, one dram.
Florentine Iris, one dram and a half.
Liquorice, two drams.
Anise, half a dram.
Saffron, eight grains.
Subcarbonate of Magnesia, one dram.

Make a powder. (AUGUSTIN)

R. Valerian, one ounce.
Oxide of Zinc, one scruple.
Musk, ten grains.

Make a powder. (BRERA)

The dose of all these powders is, in general, about twenty grains.

ANTHELMINTIC POWDER. (AUGUSTIN)

R. Root of Valerian, two drams.
—— Jalap, } of each,
Black Sulphuret of Mercury, } one dram.

Mix.—Dose, a tea-spoonful, from time to time.

ANTI-EPILEPTIC BOLUSES.

Boli ad epilepticos seu e Valerian. (FORM. HÔP. FR.; RATIER; SAUNDERS; SWEDIAUR)

R. Root of Valerian, ten parts.
Cinnamon, three parts.
Simple Syrup, a sufficient quantity.

Make a mass. (FORM. HÔP. FR.)

R. Root of Valerian, half a dram.
Simple Syrup, one dram and a half.

Mix. (SAUNDERS)

SWEDIAUR prescribes syrup of orange-flowers. Ten grains of castor, or seven of black oxide of iron, may be added.

R. Root of Valerian, twenty grains.
Sulphate of Potass, two grains.
Syrup of Orange Peel, a sufficient quantity.

Mix. (RATIER)

ANTI-EPILEPTIC PILLS. (CADET DE GASSICOURT; RATIER; SAINTE-MARIE)

R. Valerian, six drams.
Galbanum, } of each,
Sagapenum, } one dram and a half.
Assafetida, one dram.

Make pills of three grains each. (CADET DE GASSICOURT)

R. Valerian, thirty grains.
Castor, four grains.
Oxide of Zinc, twenty grains.
Simple Syrup, a sufficient quantity.

Make three pills. (RATIER)

They are prescribed by DUPUYTREN.

R. Cocoa-nut Oil, four drams.
Acetate of Potass, } of each,
Valerian, two drams.
Castor, one dram.
Cinnabar, one scruple.
Syrup of Orange Flowers, a sufficient quantity.

Make pills of three grains each. (SAINTE-MARIE)

R. Valerian, one ounce.
Iron Filings, } of each, two drams.
Catechu, } of each, two drams.
Cascarilla, } of each, two drams.
Syrup of Cinchona, a sufficient quantity.

Make pills of three grains each. (SAINTE-MARIE)

ANTI-SPASMODIC ELECTUARY.

Conditum epilepticum. (SAINTE-MARIE; VAN MONS)

R. Root of Valerian, two ounces.

VOL. II.

Subcarbonate of Magnesia, two drams.
Syrup of Peony, a sufficient quantity.

Mix. (SAINTE-MARIE)

R. Root of Valerian, } of each,
Leaves of Orange, } six drams.
Syrup of Wormwood, a sufficient quantity.

Mix. (VAN MONS)

Dose, a tea-spoonful twice or thrice a day.

ANTHELMINTIC ELECTUARY.

Electuarium anthelminticum. (HAMB.; HANN.; ROSS.; WURCEB.; NIEMANN)

R. Sulphate of Potass, } of each,
Root of Jalap, } one dram.
—— Valerian, } of each, two drams.
Oxymel of Squill, four ounces.

Mix. (ROSS.; NIEMANN)

R. Root of Valerian, two drams.
—— Jalap, two scruples.
Semen Contra, two drams.
Glycyzied Honey, three ounces.

Mix. (HAMB.)

R. Semen Contra, one ounce.
Valerian, } of each, two drams.
Jalap, } of each, two drams.
Sulphate of Potass, half an ounce.
Oxymel of Squill, five ounces.

Mix. (HANN.)

WURCEB. prescribes one dram and a half of valerian, one dram of jalap, half an ounce of semen contra, two drams of sulphate of potass, six drams of oxymel of squill, and a sufficient quantity of despumated honey.

Dose, one or two drams to children, and half an ounce to adults, four times a day.

VERMIFUGE LINCTUS.

Linctus vermifugus. (SAUNDERS)

R. Powdered Valerian, one dram.
Olive Oil, one ounce.

Mix.

ANTI-EPILEPTIC POTION. (BORIES)

R. Black Cherry Water, } of each,
Lime-flower Water, } two ounces.
Orange-flower Water, } of each, two drams.
Powdered Valerian, half a dram.
Alkermes Syrup, three drams.

Mix.—To be taken at twice.

EXCITANT AND ANTI-SPASMODIC POTION. (PIERQUIN)

R. Root of Valerian, twenty-four grains.
HOFFMANN'S Anodyne Liqueur, thirty drops.

Mint Water,	}	of each,
Hordeated Cinnamon		
Water,		
Mix.		two ounces.

II. Preparations which contain only the active principle of Valerian.

A.

Extraction by Water.

INFUSION OF VALERIAN.

Infusum Valerianæ. (DUBL.; AMER.; HAMB.; FORM. HÔP. FR.; CAST. BORUS.; WURCEB.; COXE.; DISP. CHA. PARIS.; HUFELAND.; NIEMANN.; RATIER.; SAUNDERS.; SWEDIAUR.)

R. Root of Valerian, in coarse powder, two drams.
Boiling Water, seven ounces.

Digest for one hour, and strain when cold. (DUBL.)

AMER.; and COXE give the same formula as LOND.—RATIER prescribes one ounce of root, and two pounds of water.—MIL. COPEN.; and NIEMANN—one ounce of root, and one pound of water.—HAMB.—one ounce of root, and a sufficient quantity of water to obtain eight ounces of strained infusion.—CAST. BORUS.; and WURCEB.—one ounce and a half of root, and one pound and a half of water.—DISP. CHA. PARIS.; —two drams each, of valerian and of liquorice, and one pound and a half of water.—HUFELAND—half an ounce of valerian, and a sufficient quantity of water to obtain eight ounces of infusion.

It is excitant, nervine, and diaphoretic.—Dose, one or two spoonfuls every hour.

R. Root of Valerian, one ounce.
Boiling Water, twelve ounces.
Macerate for two hours, and strain. Add Compound Spirit of Lavender, six drams.

Or,

Ammoniated Tincture of Valerian, three drams.

Mix. (SAUNDERS.; SWEDIAUR.)

SYRUP OF VALERIAN.

Syrupus Valerianæ Radicis. (HANN.)

R. Root of Valerian, eight ounces.
Boiling Water, three pounds.
Infuse for several hours. Add to twenty ounces of the strained liquor,

White Sugar, four pounds.
Dissolve.

COMPOUND INFUSION OF VALERIAN.

Infusum Valerianæ compositum seu tonico-nervosum. (MIL. COPEN.; AUGUSTIN)

R. Root of Valerian, } of each,
—— Sweet Flag, } one ounce.
Flowers of Leopard's-bane, two drams.
Boiling Water, twelve ounces.

Digest till cold. Add to the expressed liquor

Tincture of Buck-bean, three drams.
Sulphuric Ether, two drams.

Mix.—Dose, half an ounce. (MIL. COPEN.)

R. Root of Valerian, from two to six drams.
Boiling Water, eight ounces.
Infuse, and add to the liquor when cold,
Cinnamon Water, two ounces.
HOFFMANN'S Liquor, two drams.
Syrup, one ounce.

Mix. (AUGUSTIN)

Dose, a spoonful every hour or two.

R. Root of Valerian, two drams.
Boiling Water, one ounce.
Infuse. Add to the strained liquor,
Cinnamon Water, one ounce.
Living Wood-lice, fifty.

Digest for two hours in a sand-bath; then express, and add

Syrup of Orange Peel, one ounce.
Tincture of Opium, six drops.

Mix, and at the moment of administration, add a cubic inch of pure hydrogen gas. This potion has been recommended in amaurosis. (AUGUSTIN)

NERVINE INFUSION. (SAUNDERS)

R. Root of Valerian, one ounce and a half.
Decoction of Cinchona, two pounds.
Infuse, and add to the strained liquor,
Syrup of Chamomile, one ounce and a half.

Camphor, triturated with Mucilage, twenty grains.

Mix.

ANTHELMINTIC LAVEMENT.

R. Herb of Wormwood, } of each,
Root of Valerian, } one ounce.
Tansey, } of each,
Orange Peel, } half an ounce.

Pulverize, and mix.—Two spoonfuls of this powder is to be infused for one night in a pound of boiling water, and expressed.—For two lavements, to each of which a spoonful of oil of hartshorn is to be added.—It is praised by BREMSER.

ANTI-SPASMODIC DRINK.

- | | | |
|-----------------------|---|------------------------|
| R. Miseltoe, | } | of each,
two drams. |
| Root of Valerian, | | |
| Peony, | | |
| Orange Leaves, | | |
| Water, | | one pinch. |
| one pound and a half. | | |

Boil down to one pound.

EXCITANT POTION. (FORM. HÔP. FR.; AUGUSTIN; BRERA)

- R. Valerian, two drams and a half.
 Water, a sufficient quantity
 to obtain four ounces of strained decoction.
 Add

Simple Syrup, one ounce.
 Tincture of Cinnamon, ten grains.

Mix. (FORM. HÔP. FR.)

- R. Valerian, one dram.
 Decoction of Cinchona,
 a sufficient quantity

to obtain eight ounces of infusion. Add

Sulphuric Æther, half a dram.
 Tincture of Opium, thirty drops.

Mix. (BREERA)

- R. Camphor, one scruple.
 Sulphuric Æther, two drams.

Dissolve, and add

Extract of Valerian, six drams.
 Peppermint Water, six ounces.
 SYDENHAM'S Laudanum, one dram.

Mix.—Dose, half a spoonful every two or three hours. (AUGUSTIN)

AQUEOUS EXTRACT OF VALERIAN.

Extractum Valerianæ aquosum. (DUBL.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; HISP.; GALL.; FULD.; HANN.; LIPP.; BORUS.; SAX.; SUEC.; PIDERIT)

Extractum Valerianæ. (DUBL.)

- R. Root of Valerian, in
 coarse powder, six ounces.
 Boiling Water, three pints.

Digest at a medium heat for twenty-four hours in a covered vessel; then express the liquor, and evaporate it to the proper consistence.

- R. Root of Valerian, any quantity.
 Enclose it in a woollen bag, and pour boiling water upon it till it passes through tasteless. Evaporate the liquor slowly to the proper consistence. (FULD.)

- R. Root of Valerian, one part.
 Water, six parts.

Macerate for twenty-four hours, and express; set aside for two hours, then decant,

and evaporate the liquor to the proper consistence. (BAVAR.)

- R. Root of Valerian, one pound.
 Cold Water, four pounds.

Macerate for twenty-four hours, occasionally stirring, and strain; macerate the residue with fresh water, and strain again. Mix the two liquors, and evaporate to the consistence of an extract. (GALL.)

HANN.; BORUS.; and SAX. require two days' maceration in twelve parts of water, evaporation to one-sixth, filtration, and final evaporation in a water-bath.

- R. Root of Valerian, one pound.
 Spring Water, six pounds.

Digest for four days in a warm place; then, having boiled a little, express, and evaporate the liquor to the consistence of an extract. (BRUNS.)

PIDERIT directs one pound of root to be boiled in eight pounds of water to the reduction of one half.

- R. Root of Valerian, one part.
 Boiling Water, eight parts.

Boil slightly for a quarter of an hour, and express; boil the residue with four parts of water, and strain again. Mix the two decoctions; decant after sufficient repose, and evaporate to the proper consistence. (ANT.; FERR.; COXE)

AUSTR. directs the root to be exhausted by several successive boilings, the liquors to be mixed, decanted after sufficient repose, and evaporated in a water-bath.—SUEC.—one part of root to be boiled with two parts of water; the boiling to be repeated with fresh water, and the mixed decoctions evaporated in a water-bath.—BATAV.—one part of root to be digested for twenty-four hours, and boiled for two, with ten parts of water, and the residue to be boiled for two hours with ten other parts of water.—AMST.—the root to be first boiled with twenty and then with fifteen parts of water, to be each time reduced to one half.—HISP.—the root to be digested for twenty-four hours and boiled for two, with ten parts of water, and the residue boiled with more water, reduced to half.

Dose, from one scruple to one dram.

B.

Extraction by Beer.

ANTI-SPASMODIC BEER. (VAN MONS)

- R. Root of Valerian, five parts.
 Leaves of Rosemary, } of each,
 Sage, } one part and
 a half.
 Serpentaria, one part.

Strong Beer,
one hundred and twenty-eight parts.
Macerate for eight or ten days, and strain.

CEPHALIC BEER. (CADET DE GASSICOURT; SAUNDERS)

R. Root of Valerian, one ounce.
Mustard, six drams.
Rosemary, four drams.
Serpentaria, three drams.
Beer, twelve pounds.

Macerate for two days, and strain. (SAUNDERS)

CADET DE GASSICOURT prescribes ten ounces of valerian, six of mustard, three each, of sage and of rosemary, two of serpentaria, and sixty pounds of beer.

C.

Extraction by Wine.

WINE OF VALERIAN.

Infusum Valerianæ vinosum. (NIEMANN)

R. Root of Valerian, one ounce.
White Wine, one pound.

Digest for twenty-four hours, and strain.—Dose, a spoonful every hour or two.

D.

Extraction by Alcohol.

TINCTURE OF VALERIAN.

Tinctura seu Essentia Valerianæ sylvestris seu officinalis. (LOND.; DUBL.; AMER.; AMST.; ANT.; AUSTR.; BELG.; BRUNS.; MIL. COPEN.; FULD.; HANN.; POL.; BORUS.; SAX.; BRUGNATELLI; HUFELAND; SWEDIAUR; VAN MONS)

Tinctura Valerianæ.

R. Root of Valerian, in powder, four ounces.

Proof Spirit, two pints.

Macerate for fourteen days, and filter. (LOND.)

DUBL. requires but seven days' digestion.

AMER.; and COXE give the same formula as DUBL.—AMST.; AUSTR.; and HUFELAND prescribe two ounces of root, and one pound of alcohol (0.910).—ANT.—one part of root, and six parts of alcohol (20 degrees).—BATAV.; and BELG.—one of root, and six of alcohol (15 degrees).—VAN MONS—one of root, and six of proof spirit.—BRUNS.—one of root, and five of alcohol.—HANN.; POL.; and BORUS.—five ounces of root, and two pounds of alcohol.—SAX.—one part of root, and four parts of

alcohol.—MIL. COPEN.; and SWEDIAUR—one of root, and six of alcohol.

R. Root of Valerian, two ounces.
Cinnamon, six ounces.
Alcohol, two ounces.

Infuse, and express: (FULD.)

It is excitant, anti-spasmodic, nervine, vermifuge, and diaphoretic.—Dose, from thirty to sixty drops.

ALCOHOLIC EXTRACT OF VALERIAN.

Extractum Valerianæ alcoholicum. (BATAV.; DAN.; LIPP.; POL.; HERBIP.)

R. Root of Valerian, } of each,
Alcohol, } two pounds.
Water, } nine pounds.

Digest for twelve hours, and express; then distil off the alcohol, and evaporate the residue to the proper consistence. (POL.)

BATAV. prescribes two parts of root, three of alcohol, and nine of water.—DAN.—one of root, two of alcohol, and four of water.

R. Root of Valerian, any quantity.
Alcohol, a sufficient quantity

to cover it by four fingers' breadth. Digest in a warm place, and decant; boil the residue for one hour, in water, and express; then evaporate the tincture to the consistence of honey, adding the decoction towards the end, and stirring constantly. (LIPP.; HERBIP.)

CAMPHORATED TINCTURE OF VALERIAN. (HAMB.)

R. Root of Valerian, two ounces.
Camphor, one dram.
Alcohol, eight ounces.

Digest to perfect extraction, and strain.

COMPOUND TINCTURE OF VALERIAN.

Tinctura seu Essentia Valerianæ composita. (BRUNS.; WIRTEM.; PIDERIT; VAN MONS)

R. Root of Valerian, } of each,
———Serpentaria, } one ounce.
Camphor, } one dram.
Alcohol, } eight ounces.

Digest, express, and filter. (BRUNS.; WIRTEM.; PIDERIT)

R. Root of Valerian, one ounce.
———Serpentaria, half an ounce.
Proof Spirit, eight ounces.

Infuse, express, and filter. Add

Camphor, one dram.

Dissolve. (VAN MONS)

It is excitant, and is reputed alexiterious, uterine, and anti-epileptic.—Dose, from forty to sixty drops.

ANTI-SPASMODIC AND TONIC TINCTURE.
(PIERQUIN)

- R. Root of Valerian, } of each,
Powdered Cinchona, } two drams.
Ipecacuanha, } thirty grains.
Boiling Decoction of
two Poppy Heads, sixteen ounces.
Infuse for twenty-four hours in a covered
vessel, and then add
Syrup of Ground Ivy, one ounce.
Mix.—Dose, a spoonful thrice a day.

NERVINE AND TONIC TINCTURE. (CADET
DE GASSICOURT; VAN MONS)

- R. Root of Valerian, } of each,
Herb of Peppermint, } one ounce and
a half.
Castor, one ounce.
Saffron, half a dram.
Essence of Mint, one dram.
Alcohol, two pounds.
Digest, express, and filter. (CADET DE
GASSICOURT)
R. Castor, two parts.
Saffron, one part.
Proof Spirit, } of each,
Peppermint Water, } twenty parts.
Infuse without heat for several days; then
add

Root of Valerian, two parts.

At the end of several days express and
filter. (VAN MONS)

It is excitant, and is recommended in af-
fections supposed to result from debility
and spasm of the stomach.—Dose, twenty
or thirty drops.

AMMONIATED TINCTURE OF VALERIAN.

Tinctura seu Essentia Valerianæ ammoniata
seu volatilis. (LOND.; EDINB.; DUBL.;
AMER.; ANT.; DAN.; GENEV.; HANN.;
OLDEN.; LUSIT.; POL.; BORUS.; SAX.;
SUEC.; HERBIP.; COXE; SWEDIAUR;
VAN MONS)

Tinctura Valerianæ ammoniata.

- R. Root of Valerian, bruised,
four ounces.
Aromatic Spirit of Ammonia,
two pints.
Macerate for fourteen days, and
strain. (LOND.; AMER.; COXE)
R. Root of Valerian, bruised,
four ounces.
Spirit of Ammonia,
two pounds and a half.

Digest for seven days, and filter.
(EDINB.)

- R. Root of Valerian, in powder,
two ounces.
Spirit of Ammonia, one pint.

Digest for seven days, and strain.
(DUBL.)

- R. Root of Valerian, one part.
Spirit of Ammonia, six parts.

Digest without heat for several days in a
covered vessel, occasionally shaking, and
strain. (DAN.; GENEV.; HANN.; OL-
DEN.; POL.; BORUS.; SAX.; HERBIP.;
NIEMANN; SWEDIAUR)

VAN MONS prescribes one part of root,
and eight parts of spirit of ammonia.—LU-
SIT.—four ounces of root, and two pounds
and a half of aromatic spirit; to be mace-
rated for eight days.

- R. Root of Valerian, in powder,
two ounces.
Alcoholic Solution of Subcar-
bonate of Ammonia, one pound.

Macerate for eight days, frequently agi-
tating, and strain. (SUEC.)

- R. Valerian, in coarse powder,
one ounce and a half.
Alcohol (30 degrees), four ounces.
Solution of Ammonia (19 degrees),
two ounces.

Macerate for six days in a covered vessel,
occasionally agitating, and filter. (ANT.)

It is excitant, and has been recommended
in several spasmodic diseases, particularly
in epilepsy and hysteria.—Dose, from one
dram to three drams, in milk or gruel.—
NIEMANN proposes, as a milder prepara-
tion, a mixture of one part of solution of
subcarbonate of ammonia, and two parts of
simple alcoholic tincture of valerian.

COMPOUND AMMONIATED TINCTURE OF
VALERIAN. (PIERQUIN; VAN MONS)

- R. Root of Valerian, two ounces.
Oil of Lemon, one dram.
Cloves, half a dram.
Spirit of Ammonia half an ounce.
Alcohol, sixteen ounces.

Digest in a sand-bath for two or three
days, and filter. (PIERQUIN)

- R. Root of Valerian, one ounce.
Cloves, } of each, one dram.
Mace, }
Fresh Lemon Peel,
one dram and a half.
Aromatic Spirit of Ammonia,
twenty-four ounces.

Digest without heat for several days; then
express and filter.

It is recommended in epilepsy.—Dose, twenty or thirty drops, in infusion of valerian.

ACOUSTIC LIQUOR.

Aqua acoustica Ludovici. (PALAT.; SPIELMANN)

R. Root of Valerian	} of each,	half an ounce.
—— Lovage,		
Herb of Rosemary,		
Flowers of Lavender,		
Bay Berries,	} of each, two drams.	one pound.
Castor,		
Camphor,		
Spirit of Juniper,		

Macerate for four days, and add

Solution of Ammonia,	half an ounce.
Oil of Juniper,	half a dram.

Mix.—A few drops, on cotton, are to be introduced into the meatus auditorius.

E.

Extraction by Æther.

ÆTHEREAL TINCTURE OF VALERIAN.

Tinctura Valerianæ ætherea. (FENN.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; WURCEB.; AUGUSTIN.; BRUGNATELLI.; CADET DE GASSICOURT.; PIERQUIN.; NIEMANN.; SWEDIAUR.; VAN MONS)

R. Root of Valerian, in coarse powder,	one ounce.
Spirit of Sulphuric Æther,	eight ounces.

Digest without heat for three days in a covered vessel, and filter. (FENN.; HANN.; POL.; BORUS.; SAX.; NIEMANN.; SWEDIAUR.)

HASS. prescribes two ounces of root, and one pound of the spirit.—PIERQUIN—eight ounces of root, and one pound and a half of the spirit.—VAN MONS—one part of root, and ten parts of the spirit.—BRUGNATELLI—three ounces of root, and one pound of the spirit: this is the only one which is required to be prepared with heat.—HAMB.; LIPP.; OLDEN.; and CADET DE GASSICOURT—one part of root, and four parts of the spirit.

R. Root of Valerian, in powder,	one ounce.
Concentrated Alcohol,	six ounces.

Digest at a moderate heat in a covered vessel. Add to the filtered tincture,

Sulphuric Acid,	two drams.
-----------------	------------

Mix well. (SUEC.; WURCEB.)

AUGUSTIN prescribes sixteen ounces of valerian, twenty of alcohol, and twelve of sulphuric acid.

It is excitant, and is recommended in nervous disorders, particularly in spasms, convulsions, and hysteria.—Dose, from one to four scruples, three or four times a day, in infusion of balm or of peppermint.

CAMPHORATED ÆTHEREAL TINCTURE OF VALERIAN.

Tinctura Valerianæ camphorata. (AUGUSTIN)

R. Camphor,	half a dram.
Æthereal Tincture of Valerian,	three drams.

Dissolve.—Dose, twenty-five or thirty drops, in gout.

ANODYNE TINCTURE. (NIEMANN)

R. Root of Valerian,	one ounce.
Bitter Orange Peel,	two drams.
Sulphuric Æther,	four ounces.

Digest for eight days at a very gentle heat, and then boil for an instant.

It is excitant and anti-spasmodic.—Dose, from forty to fifty drops.

ANTI-SPASMODIC TINCTURE. (NIEMANN)

R. Root of Valerian,	} of each,	one ounce.
Herb of Marum,		
Spirit of Sulphuric Æther,		

Digest for six days, and filter.

R. Root of Valerian,	} of each,	half an ounce.	
Herb of Marum,			
Alcohol,			four ounces.
Sulphuric Acid, diluted with water,			one dram.

Digest for four days at a moderate heat.

It is excitant, anti-spasmodic, and anti-epileptic.—Dose, a tea-spoonful, in wine.

ANTI-SPASMODIC ELIXIR. (FULD.; VAN MONS)

R. Tincture of Valerian,	two ounces.
Spirit of Sulphuric Æther,	two drams.

Mix, and add

Camphor,	one dram.
----------	-----------

Dissolve. (VAN MONS)

R. Root of Valerian,	} of each,	one ounce.
—— Serpentina,		
Alcohol,		

Digest, express, and filter. Add

Sulphuric Æther,	two ounces.
Camphor,	one dram.

Dissolve. (VAN MONS)

R. Camphor, one dram.
Sulphuric Æther, two drams.
Add to the solution,
Extract of Valerian, one ounce.
Chamomile Water, six ounces.

Mix. (FULD.)

Dose, one or two scruples, several times a day.

F.

Extraction by Distillation.

ESSENTIAL OIL OF VALERIAN.

Oleum Valerianæ æthereum, Ætheroleum Valerianæ. (AUSTR.; BATAV.; BELG.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SWEDIAUR)

R. Root of Valerian, any quantity.
Water, a sufficient quantity.

Distil. (BELG.; ROSS.)

SWEDIAUR prescribes one part of root and four parts of water.—HANN.; OLDEN.; POL.; BORUS.; and SAX.—one of root and eight of water.—BAVAR.—one of root and nine of water.—AUSTR.—one of root and four of water.

R. Root of Valerian, four parts.
Water, sixteen parts.

Macerate for four days, and add

Muriate of Soda, one part.

Distil. (HASS.)

R. Root of Valerian, six pounds.
Alcohol, one pound.

At the end of two days, add

Water, eighteen pounds.

Distil. (LIPP.)

OLEOSACCHARUM OF VALERIAN.

Oleosaccharum Valerianæ. (AUSTR.; HANN.; BORUS.)

R. White Sugar, one dram.

Oil of Valerian, three grains.
Triturate together. (AUSTR.)

HANN.; and BORUS. prescribe one ounce of sugar and twenty pounds of oil.

DISTILLED WATER OF VALERIAN.

Aqua Valerianæ. (AUSTR.; BAVAR.; BRUNS.; GALL.; HANN.; HASS.; LIPP.; POL.; SAX.; VAN MONS)

R. Root of Valerian, one part.
Water, a sufficient quantity.

At the end of twenty-four hours, distil six (HASS.), or seven (HANN.), parts.

BRUNS. prescribes one part of root and three parts of water, one half to be distilled.—GALL.—one of root and five of water; two to be distilled.—VAN MONS—one of root and ten of water; two to be distilled.—AUSTR.—one of root and twelve of water; four to be distilled.—POL.; and SAX.—one of root and twelve of water; nine to be distilled.—BAVAR.; and NIEMANN—one of root and eighteen of water; ten to be distilled.

R. Root of Valerian, two pounds.
Water, twelve pounds.

Enclose the root in a bag suspended in the centre of a cucurbit, so that it may not touch the water, and distil six pounds. (LIPP.)

COMPOUND SPIRIT OF VALERIAN.

Aqua Valerianæ composita. (PIDERIT)

R. Root of Valerian, nine ounces.
—— Lovage, three ounces.
Herb of Pennyroyal, two ounces.
—— Savine, one ounce.
Spring Water, } of each, eight pounds.
Alcohol, }

After two days' digestion, distil eight pounds.

VALERIANELLA OLITORIA, ME.

SYNONIME.—Mache, Doucette, Fr.

GALL.; BERGIUS.

A common plant of Europe. (*Triandria Monogynia*, LINN.; *Valerianæa*, JUSS.)

The herb is employed; it consists of a branching stem, furnished with oblong, sessile, entire or slightly dentated leaves, and rising from a rosette of similar leaves. It has a vapid taste.

It is emollient.

VANILLA AROMATICA, Sw.—VANILLA.

SYNONIMES :—Vanille, FR. ; and GERM. ; Wanylie, BOHEM. ; Vaynilla, SPAN. ; Baniļe, DUT. ; Vaniglia, ITAL. ; Tlix-ochitl, MEX. ; Wanilia, POL. ; Vanilj, SWED.

ANT. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; HANN. ; POL. ; BORUS. ; ROSS. ; SAX. ; SUEC. ; WIRTEM. ; HERBIP. ; BERGIUS ; BRUGNATELLI ; GUIBOURT ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN ; ZARDA.

A parasitical and sarmentose plant of South America. (*Gynandria Diandria*, LINN. ; *Orchideæ* JUSS. ; *Fig. Flore Medic.* VI. 345)

The fruit (*Siliquæ Vanilla seu Vanillæ seu Vaniglia seu Banilla seu Araci aromatici*) are employed. It consists of pods of from four to eight inches long, and two or three lines thick, straight, curved at their origin, terminated by a kind of pediculated nipple, cylindrical, grooved, shining, compressed, bivalved, with thick, though rather soft, and shining valves, enclosing a soft brown pulp surrounding numerous round, blackish, shining seeds. They are of a blackish-brown colour ; their odour is strong and agreeable ; and their taste warm and pungent.

They are stimulant.—Dose of the powder, from fifteen grains to a dram.

POWDER OF VANILLA. (GALL.)

R. Vanilla, cut small, one part.
White Sugar, two parts.

Reduce to a very fine powder in an iron mortar, and pass it through a silk sieve.

FEE ; and HENRI prescribe four parts of sugar.

INDIAN WAKAKA. (CADET DE GASSI-COURT)

Sugar, four ounces.
Chocolate, one ounce and a half.
Sugar of Vanilla, ten drams.
Cinnamon, } of each, one dram.
Terra Orleana, }

Pulverize, and mix.—Dose, a spoonful, in milk, chocolate, or broth.

SYRUP OF VANILLA. (PLANCHE ; VAN MONS)

R. Vanilla, in powder, four parts.
Gum Arabic, one part.
Water, six parts.

Triturate together ; then express strongly, and add to the liquor, a cold syrup made with

White Sugar, twenty-two parts.
Water, twelve parts.

Then dissolve slowly in the mixture,

White Sugar, twenty-two parts.

R. Vanilla, two ounces.
White Sugar, seventeen ounces.
River Water, ten ounces.
Alcohol (21 degrees), six drams.

Cut the vanilla into small pieces, and levigate it, adding a little sugar and alcohol, till reduced to a soft and homogeneous paste. Put this into a matrass with the remainder of the sugar and the water ; add the white of an egg, keep it in a water-bath for eighteen or twenty hours, shaking the mixture occasionally ; then having set it aside for one day, strain it through a sieve.

TINCTURE OF VANILLA.

Tinctura Vanigliæ. (SAX. ; NIEMANN)

R. Vanilla, one part.
Alcohol, ten parts.

Digest for eight days at a moderate heat. (SAX.)

NIEMANN prescribes, after PFAFF, one part of vanilla and six parts of alcohol.

It is excitant, reputed nervine, and anaesthetic.—Dose, forty drops, in wine.

LIQUID ALKERMES. (VAN MONS.)

R. Ambergris, one scruple.
Cloves, half a dram.
Cinnamon, two drams.
Vanilla, } of each, half an ounce.
Cochineal, }
Proof Spirit, three drams.

Infuse for two days in a water-bath ; then filter, and dissolve in the liquor, in a water-bath at the temperature of 60 degrees, a mixture of

R. White Sugar, three pounds.
Honey, six pounds.

Decant the syrup, when depurated, and strain it through a woollen sieve.

DISTILLED WATER OF VANILLA.

Aqua Vanilla. (NIEMANN)

R. Vanilla, one pound.
Subcarbonate of Potass, three ounces.
Water, twelve pounds.

Digest for twenty-four hours, and distil one half.

SPIRIT OF VANILLA.

Spiritus Vanilla. (NIEMANN)

- R. Vanilla, six ounces.
Subcarbonate of Potass, one ounce and a half.

Alcohol, six pounds.
Macerate for twenty-four hours, and add
Water, six pounds.
Distil slowly six pounds.

VERATRINUM.—VERATRINE.

Veratrina, Veratria.

MAGENDIE.

- R. Seeds of Sabadilla, any quantity.

Exhaust it with boiling alcohol, and filter the tincture; set it aside to cool, filter again, and evaporate to the consistence of an extract: dissolve this in water, filter the liquor, and evaporate it slowly. Separate the precipitate which is formed, pour solution of acetate of lead into the strained liquor, and separate the precipitate again by filtration. Pass a current of hydrocyanic acid through the liquor, and filter again, then concentrate the liquor by evaporation, treat it with magnesia, and finally exhaust the precipitate which is formed, by boiling alcohol, and evaporate the mixed tinctures to dryness.

It is violently stimulant and poisonous. It purges at the dose of a quarter of a grain.

SOLUTION OF VERATRINE. (MAGENDIE)

- R. Sulphate of Veratrine, one grain.
Distilled Water, two ounces.
Dissolve.

TINCTURE OF VERATRINE. (MAGENDIE)

- R. Veratrine, four ounces.
Alcohol, one ounce.

Dissolve.—Dose, from ten to twenty-five drops in a potion.—It is used externally, in embrocations, in gout, dropsy, and white swelling.

PILLS OF VERATRINE. (MAGENDIE)

- R. Veratrine, half a grain.
Gum Arabic, } of each,
Syrup of Gum, } a sufficient quantity.

Make pills of one grain each.

Dose, from one to three, daily.

OINTMENT OF VERATRINE. (MAGENDIE)

- R. Veratrine, four grains.
Lard, one ounce.

Mix.—To be rubbed upon the parts affected in gout, anasarca, and chronic rheumatism.

VERATRUM.

Two species of this genus of plants are employed in medicine:

1. *Veratrum Sabadilla*, RE.—Sabadilla.

SYNONIMES:—Cevadille, FR.; Sabadille, Mexikanischer Læusesaamen, Læusemøerder Læusekørner, Mönchensaat, GERM.; Sabadillkorn, DAN.; Nasikio sabatyli, POL.; Sabadill, Husarfrøe, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; CAST. BORUS.; LUSIT.; ROSS.; SAX.; SUEC.; VOL. II.

WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

A plant of Mexico. (*Polygamia Monœcia*, LINN.; *Colchicaceæ*, JUSS.)

The fruit and seeds are employed.

The fruit is an oval capsule, pointed at one end, obtuse at the other, about half an inch long, of a light brown colour, inodorous, and of an acrid, burning, and durable taste.

The seeds (*Semen Sabadilli* seu *Sabadille*) are blackish, rugose, and slightly convex.

It contains, according to MEISSNER, PELLETIER, and CAVENTOU, *Cevadic Acid* and *Veratrine* or *Sabadilline*.

It is excitant and irritant, occasioning, when taken internally, burning heat of the stomach and vomiting; externally, it inflames the skin, particularly in young subjects. It is used to destroy lice and also intestinal worms. SEELIGER has employed it successfully against tænia. It should be prescribed with great circumspection.

CAPUCINS' POWDER. (NIEMANN)

R. Seeds of Sabadilla,	} of each, an equal part.
—— Stavesacre,	
—— Parsley,	
Leaves of Tobacco,	

Make a fine powder.

To destroy lice on the head.—This powder frequently produces vertigo and other unpleasant symptoms.

OINTMENT AGAINST LICE. (FULD.)

R. Powdered Sabadilla,	four ounces.
—— Mustard,	} of each, two ounces.
—— Pyrethrum,	
Prepared Lard,	thirty-two ounces.

Mix.

ANTHELMINTIC PILLS. (CADET DE GASSICOURT; SWEDIAUR)

R. Sabadilla,	} of each, half an ounce.
Honey,	

Make pills of five grains each. (SWEDIAUR).—CADET DE GASSICOURT adds twenty drops of oil of fennel.

LAVEMENT AGAINST ASCARIDES.

Enema Sabadilla. (AUGUSTIN; NIEMANN; SWEDIAUR)

R. Sabadilla,	two drams.
Spring Water,	ten ounces.

Boil, and to seven ounces of the strained liquor, add

Cow's Milk,	seven ounces.
-------------	---------------

Mix.

Bedsteads have been recommended to be washed with the following liquor to destroy bugs:

R. Seeds of Sabadilla,	two ounces.
Vinegar,	one ounce.

Digest for two days, and strain.

2. *Veratrum album*, LINN.—White Hellebore.

SYNONIMES:—Ellébore blanc, FR.; Weissnieswurz, Lausekraut, Krätzwurz, GERM.; Cemerice bj la, Keychawk bj la,

BOHEM.; Wedegambre blanco, SPAN.; Wrtblæmige nieswortel, DUT.; Elleboro bianco, ITAL.; Biala ciemierzycza, POL.; Helleboro, PORT.; Hwit prustrot, SWED.

LOND.; DUBL.; EDINB.; AMER.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; HANN.; LIPP.; PALAT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS; BRUGNATELLI; COXE; GUIBOURT; MURRAY and GMELIN; PARIS; SAUNDERS; SPIELMANN; ZARDA.

A plant which grows on the high mountains of Europe. (*Fig. Jacq. Austr. IV. t. 335*)

The root (*Radix Hellebori albi* seu *Veratri*) is employed. It is cylindrical obtuse, about two inches long, sometimes more than an inch thick, furnished with numerous fibrils. It is ash-grey or reddish coloured without, and whitish within; its odour is very disagreeable but is removed by desiccation; its taste is bitterish, burning and nauseous.

It contains, according to PELLETIER and CAVENTOU, *Veratrine*, which is the source of its activity.

It is a violent drastic, and was formerly much employed in mania, hydrophobia, and melancholy, though not much used at present.—Dose, from one grain to twenty-four grains, gradually augmented.

ERRHINE POWDER.

Pulvis sternutatorius. (HANN.; PALAT.; WIRTEM.; SPIELMANN)

R. Powdered White Hellebore,	two drams.
—— Root of Iris,	one dram.
—— Rice,	one ounce and a half.

Oleo-balsamic Mixture, twenty drops.

Mix. (HANN.)

R. Herb of Marjoram,	one dram and a half.
—— Betony,	} of each, half a dram.
—— Pyrethrum,	
Root of White Hellebore,	} of each, one dram.
—— Black Hellebore,	
Flowers of Pennyroyal,	one scruple.

Make a powder.

POLYCHREST PILLS.

Pilula polychrestæ Starkeyi. (BRUNS.; WIRTEM.; SPIELMANN)

R. Extract of Opium,	} of each, one ounce.
Powdered Liquorice,	
—— White Hellebore,	
—— Black Hellebore,	

STARKEY'S Soap, three ounces.

Make a pilular mass, (BRUNS.; WIRTEM.)—SPIELMANN prescribes one ounce of extract of opium, two ounces of white hellebore, and four of STARKEY'S soap.

Dose, three grains or more, to be gradually increased.

DECOCTION OF WHITE HELLEBORE.

Decoctum Veratri. (LOND.; DUBL.; AMER.; AUGUSTIN; COXE; NIEMANN; SWEDIAUR)

R. Root of White Hellebore,
bruised, one ounce.
Water, two pints.
Rectified Spirit,
two fluid ounces.

Boil the root with the water down to one pint, and strain; then add the decoction to the spirit when cold.

AMER.; COXE; NIEMANN; and SWEDIAUR give the same formula.

It is principally used externally in tinea, tetters, itch, and other obstinate diseases of the skin.

R. Root of White Hellebore,
from two to four drams.
Water, a sufficient quantity,
to obtain eight ounces of strained decoction.
Add

Subcarbonate of Potass, one ounce.
Despumated Honey, two ounces.

Mix.—To be taken by spoonfuls, in melancholy. (AUGUSTIN)

WINE OF WHITE HELLEBORE.

Vinum Veratri seu Veratri albi. (LOND.; AMER.; COXE; NIEMANN)

R. Root of White Hellebore,
bruised, eight ounces.
Proof Spirit, one pint.
Distilled Water,
one pint and a half.

Macerate for fourteen days, and filter.

R. Root of White Hellebore, eight ounces.
White Wine, two pints and a half.
Macerate for fifteen days, and strain. (NIEMANN)

AMER.; and COXE prescribe four ounces of root and one pint of wine; to be digested fourteen days.

It is emetic and purgative, and is recommended to repress irritability, and is prescribed with this view, in mania and melancholy.—Dose, ten drops and more, to be gradually augmented.

TINCTURE OF WHITE HELLEBORE.

Tinctura Ellebori albi seu Veratri. (EDINB.; HANN.; NIEMANN; SWEDIAUR)

Tinctura Veratri albi. (EDINB.)

R. Root of White Hellebore,
bruised, eight ounces.
Proof Spirit,
one pound and a half.

Digest for seven days, and filter.

HANN. prescribes five ounces of root and two pounds of rectified spirit.—SWEDIAUR—eight ounces of root and two pounds and a half of alcohol (0,935).—NIEMANN.—eight ounces of root and two pounds of alcohol (0,930).

It is stimulant and emetic.—Dose, from five to fifteen drops, in a proper vehicle.

LOTION OF HELLEBORE.

Lotio e Veratro. (SWEDIAUR)

R. Root of White Hellebore,
half an ounce.
Boiling Water,
from twenty to thirty ounces.
Strain when cold; then add to the liquor,
Tincture of White Hellebore,
four ounces.

Mix.—It is recommended in prurigo, tinea, and psoriasis, morning and evening.

VINEGAR OF WHITE HELLEBORE.

Acetum Veratri. (NIEMANN)

R. Root of White Hellebore, one dram.
Vinegar, twenty ounces.

Boil and strain.

It is recommended by WENDT, in mania.—Dose, a spoonful, every two or three hours.

OINTMENT OF WHITE HELLEBORE. (LOND.; DUBL.; SAX.; COXE; SWEDIAUR)

Unguentum Veratri.

R. Root of White Hellebore,
in powder, two ounces.
Prepared Lard, eight ounces.
Oil of Lemon, twenty minims.
Mix. (LOND.)

- R. Prepared Lard, one pound.
 Root of White Hellebore,
 in powder, three ounces.
 Make an ointment. (DUBL.)

SWEDIAUR prescribes one part of hellebore, and three parts of lard.—SAX.; and COXE give the same formula as LOND.
 It is used in cutaneous disorders.

VERBASCUM.—MULLEIN.

Two species of this genus of plants are employed:

1. *Verbascum Thapsus*, LINN.—Great Mullein.

SYNONIMES:—Bouillon blanc, FR.; Wollkraut, Königskerzkraut, GERM.; Diwizna, BOHEM.; Kongelys, DAN.; Gordolobo, SPAN.; Wollekruid, DUT.; Verbascos, ITAL.; Dziewanna ziele, POL.; Verbasco branco, PORT.; Kungslis, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; FULD.; GENÈV.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GMELIN.; PIDERIT.; SPIELMANN.; ZARDA.

A common plant of Europe. (*Pentandria Monogynia*, LINN.; *Solanææ*, JUSS.; Fig. Zorn, *Ic. pl. t. 197*)

The leaves and flowers are employed.

The leaves (*Folia seu Herba Verbasci albi*) are lanceolated, very cottony on both sides, dentated, and soft. They have a pleasant odour, and a mucilaginous, bitterish, and slightly styptic taste.

The flowers are yellow, rather large, of an agreeable odour, and of a sweet and mucilaginous taste.

It is emollient.

The *Verbascum phlomoïdes*, LINN. (GALL.), (Fig. Lob. *Ic. 560, 561*), a plant of Europe, may be substituted for it. The leaves of this are oblong-ovate, pointed, the inferior narrowed into a winged foot-stalk, and the superior heart-shaped at the base. The flowers are yellow or white.

The flowers have been analysed by MOXON; they yield a yellowish volatile oil, a fatty matter similar to oleic acid, phosphoric acid, malic acid, uncrystallizable sugar, gum, chlorophylle, a yellow resin, and salts.

2. *Verbascum nigrum*, LINN.—Black Mullein.

SYNONIMES:—Bouillon noir, FR.; Schwarze königskerze, GERM.

GALL.; MURRAY and GMELIN.

A common plant of Europe. (Fig. Zorn, *Ic. pl. t. 25*)

The root and flowers are employed.

The root (*Radix Verbasci nigri*) is branching, hairy, of a light brown colour externally, and of a yellowish brown internally.

The flowers are yellow, rather small, inodorous, and of a mucilaginous taste.

It is emollient.

EXTRACT OF GREAT MULLEIN.

Extractum Verbasci. (HERBIP.)

- R. Dried Flowers of Great Mullein, any quantity.
 Alcohol, a sufficient quantity.

Digest at a moderate heat in a covered vessel, and pour off the liquor; repeat this operation till the alcohol comes off colourless. Mix the tinctures, and distil two-thirds in a water-bath, and evaporate the remainder to the consistence of honey.

COMPOUND DECOCTION OF GREAT MULLEIN.

Decoctum Verbasci anodynum, Fotus anodynus.

- | | |
|-----------------------------|--|
| R. Leaves of Great Mullein, | } of each,
one ounce and
a half. |
| Mallow, | |
| Linseed, | two ounces. |
| Poppy Heads, | one scruple. |
| Water, | four pounds. |

Boil for a quarter of an hour.

It is principally employed in lotions.

OIL OF GREAT MULLEIN.

Oleum Verbasci. (BRUNS.; WIRTEM.; HERBIP.)

- R. Flowers of Great Mullein, one part.
 Olive Oil, two parts.

Digest for three days over a slow fire; then boil to the consumption of the humidity, and strain with expression. (FULD.; HERBIP.)

BRUNS.; and WIRTEM. prescribe an equal part of each, and eight days' digestion.

It is soothing and anodyne, and is recommended to piles and chilblains.

VERBENA.—VERVAIN.

Two species of this genus of plants are mentioned in pharmacopœias:

1. *Verbena officinalis*, LINN.—Vervain, Holy Herb.

SYNONIMES:—Verveine commune, FR.; Eisenhart, Eisenkraut, GERM.; Wrbynka, BOHEM.; Ma-pien-tsao, CHIN.; Jernurt, DAN.; Verbena, ITAL.; SPAN.; and PORT.; Yzerhard, DUT.; Zeleznik, POL.; Schelesnik, RUSS.; Jærnært, SWED.

ANT.; HISP.; GALL.; FERR.; GENEV.; LIPP.; WIRTEM.; HERBIP.; BRUGNATELLI.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN.; ZARDA.

A plant of Europe. (*Diandria Monogynic*, LINN.; *Verbenaceæ*, JUSS.; *Fig. Flore Medic. VI. 346*)

The herb (*Herba Verbenæ*) is employed.

It consists of a tetragonal, hard stem, either simple or furnished with opposite branches, bearing opposite, petiolated, oval, oblong, rather wrinkled leaves, deeply notched into unequal, obtuse, incised lobes, the terminal one of which is much larger than the others. It is destitute of odour, but its taste is bitterish and astringent.

2. *Verbena triphylla*, LINN.—Sweet-scented Vervain.

SYNONIMES:—Verveine citronelle, FR.; Yerba luisa, SPAN.

HISP.; GALL.; FERR.; BRUGNATELLI.

A plant of Chili. (*Decandria Monogynia*, LINN.; *Verbenaceæ*, JUSS.)

The leaves (*Foliæ Aloysiæ*) are employed; they are ternate. They exhale a pleasant odour resembling that of lemon.

VERONICA.

Five species of this genus of plants are used in medicine:

1. *Veronica officinalis*, LINN.—Male Speed-well.

SYNONIMES:—Veronique officinale, Veronique mâle, Thé d'Europe, FR.; Ehrenpreiss, GERM.; Weronyka, BOHEM.; Aerenprüs, DAN.; Veronica, ITAL.; PORT.; and SPAN.; Gemeene eerenprye, DUT.; Przetacznik, POL.; Empris, SWED.

LOND.; AMER.; ANT.; AUSTR.; BATAV.; BAVAR.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FORM. HÔP. FR.; GENEV.; HANN.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GME-LIN.; PIDERIT.; SPIELMANN.; ZARDA.

A common plant of Europe. (*Diandria Monogynia*, LINN.; *Pedicularæ*, JUSS.; *Fig. Flore Medic. VI. 345*)

The flowering herb (*Herba Veronica*) is

employed. It consists of a cylindrical, hairy stem, furnished with opposite, moderately petiolated, oval, obtuse or slightly pointed leaves, narrowed at the base, rough, hairy, rather shagreened, and serrated. It is destitute of odour, but has a bitterish and feebly styptic taste.

It is anti-scorbutic, and slightly tonic.—Dose, of the juice, from one ounce to two ounces.

CONSERVE OF VERONICA.

Conserva Veronica. (WIRTEM.)

R. Fresh Herb of Veronica, one part.
Sugar, two parts.

Beat together.

PECTORAL SPECIES.

Species pectorales seu pro Thea seu Thea. (LIPP.; ROSS.; CADET DE GASSICOURT.; PIDERIT)

R. Root of Dog-grass,
one ounce and a half.

- Herb of Milfoil, two ounces.
 ——— Veronica, three ounces.
 Cut, and mix. (PIDERIT)
- R. Herb of Coltsfoot, } of each,
 ——— Veronica, } four ounces.
 Root of Liquorice, } of each,
 ——— Polypodium, } two ounces.
 ——— Marshmallows, } one ounce.
 Fennel, six drams.
 Cut, bruise, and mix. (LIPP.)
- R. Leaves of Veronica, }
 ——— Ground Ivy, } of each,
 ——— Coltsfoot, } three ounces.
 ——— Scabious, }
 ——— Balm, } of each,
 ——— Sage, } half an ounce.
 Cut, and mix. (CADET DE GASSICOURT)
- R. Flowers of Chamomile, } of each,
 Herb of Veronica, } six parts.
 Leaves of Sage, two parts.
 Cut, and mix. (ROSS.)

WATER OF VERONICA.

- Aqua Veronica.* (GALL.; PALAT.)
- R. Herb of Veronica, five parts.
 Water, twenty parts.
 Distil ten parts. (GALL.)
- PALAT. prescribes one part of herb, and three parts of water; one half to be distilled.

Dose, from two to four ounces.

SPIRIT OF VERONICA.

- Aqua Veronica vinosa.* (BRUNS.; WIRTEM.; SPIELMANN)
- R. Flowering Herb of Veronica, three pounds.
 White Wine, eight pounds.
 Macerate for several days, and distil one half.

INFUSION OF VERONICA. (FORM. HÔP. FR.; RATIER)

- R. Flowers of Veronica, half an ounce.
 Boiling Water, two pounds.
 Infuse, and strain. (RATIER)
- R. Flowers of Veronica, one hundred parts.
 Grated Liquorice, eight parts.
 Boiling Water, a sufficient quantity.
 Infuse, and strain. (FORM. HÔP. FR.)

EXTRACT OF VERONICA.

- Extractum Veronica.* (BRUNS.; SARD.; WIRTEM.)
- R. Herb of Veronica, one pound.
 Spring Water, six pounds.

Macerate for four days in a warm place, then boil a little; evaporate the expressed liquor to the consistence of an extract. (BRUNS.; WIRTEM.)

- R. Fresh Herb of Veronica, bruised, eight pounds.

Macerate for three days in an alembic; then distil off about two ounces of liquor in a water-bath: pour upon the residue,

Water, twenty-four pounds, and boil down to one third. Clarify and strain the decoction, and evaporate it to the consistence of an electuary, and add the distilled water to this when merely warm. (SARD.)

Dose, from one scruple to one dram.

SYRUP OF VERONICA.

- Syrupus de Veronica.* (WIRTEM.)
- R. Herb of Veronica, six ounces.
 Spirit of Veronica, one pound.
 Infuse in a warm place, and express. Dissolve in the liquor,
 White Sugar, sixteen ounces.
 Dose, from half an ounce to two ounces.

PECTORAL SYRUP.

- Syrupus Pectoralis.* (LIPP.)
- R. Pectoral Species, eight ounces.
 Figs, four ounces.
 Water, five pounds.
 Boil down to two-thirds, and express. Add
 White Sugar, four pounds.
 Boil for a few moments, and strain.

2. *Veronica Teucrium*, LINN.

SYNONIMES:—Véronique teucriette, FR.; Brusttheekraut, Erdbathengel, GERM.

GALL.

A plant of Europe. (*Fig. Lob. Ic. t. 473*)

The flowering herb (*Herba Veronica teucriifolia*) is employed. It consists of oval, rugose, dentated, obtuse leaves, supported on a hairy, cylindrical stem; the flowers are of a pale blue colour, and are disposed in long racemes. It is inodorous, but has a bitter and astringent taste.

3. *Veronica Virginica*, WILLD.

AMER.; COXE.

The root is employed. Its medicinal properties are not yet sufficiently known.

4. *Veronica Anagallis*, LINN.—Long-leaved Brooklime.

SYNONIMES:—Petit Beccabunga, FR.; Wasserliebendes Ehrenpreis, GERM.

HISP.; GALL.

A plant of Europe. (*Diandria Monogynia*, LINN.; *Rhinantæa*, JUSS.; *Fig. CEd. Fl. Dan.* t. 903)

The flowering herb (*Herba Anagallidis aquaticæ*) is employed. It consists of a straight stem with lanceolated, pointed, den- tated leaves, and lateral racemes of blue flowers. It is inodorous, but has a bitter and acrid taste.

It is excitant.

5. *Veronica Beccabunga*, LINN.—Brook- lime, Beccabunga.

SYNONIMES:—Grand Beccabunga, Vero- nique cressonée, FR.; Bachbungen, Bach- bohnen, Wasserbungen, GERM.; Lem- mike Ledmye, Bekbung, DAN.; Beca- bunga, ITAL.; and SPAN.; Bækebom, DUT.; Potocznik, POL.; Beccabunga; PORT.; Backabunga, SWED.

DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; WIRTEM.; BERGIUS; BRUGNATELLI; COXE; GUI- BOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN; ZARDA.

A plant of Europe, growing on the bor- ders of streams. (*Diandria Monogynia*, LINN.; *Rhinantæa*, JUSS.; *Fig. Flore Med- ic.* II. 60)

The herb (*Herba Beccabungæ*) consists of a cylindrical stem, reddish inferiorly, and green superiorly, succulent, furnished with opposite, oval, glabrous, rather fleshy, den-

ticated leaves, with short footstalks. It is only employed in the fresh state.

It is excitant and anti-scorbutic.

The two last species are given in infusion, in the proportion of one or two handfuls to a pound and a half of water.

CONSERVE OF BECCABUNGA. (WIRTEM.)

R. Flowering Herb of Beccabunga, one part.
White Sugar, two parts.
Beat together.

JUICE OF BECCABUNGA.

Succus Beccabungæ. (HERBIP.)

R. Herb of Beccabunga, any quantity.
Bruise it in a stone mortar, and express strongly; set aside the juice, then decant and preserve it in bottles, adding one dram of alcohol to each pound of the juice.

Dose, from two to four ounces.

EXTRACT OF BECCABUNGA. (SARD.)

R. Fresh Herb of Beccabunga, any quantity.

Bruise it in a mortar; then express the juice, and evaporate it in a water-bath to the consistence of an extract.

Dose, from one scruple to one dram.

WATER OF BECCABUNGA. (AUSTR.; SARD.)

R. Bruised Herb of Beccabunga, one part.
Water, two parts.

Distil two thirds. (SARD.)

R. Depurated Juice of Beccabunga, any quantity.

Distil it nearly to dryness. (AUSTR.)

Dose, from two to four ounces.

VICIA.

Two species of this genus are used in me- dicine:

1. *Vicia Ervilia*, CAND.—Tare.

SYNONIMES:—Ers, FR.; Erve, GERM.; Erve, DAN.; Erven, DUT.; Erwa, SWED. FERR.; WIRTEM.; BRUGNATELLI; MUR- RAY and GMELIN; SPIELMANN.

A plant of Europe. (*Diadelphia Decan- dria*, LINN.; *Leguminosa*, JUSS.; *Fig. Blackw. Herb.* t. 208. f. 3)

The seeds (*Semen Ervi seu Orobi*) are employed. They are flattish, brown, ino- dorous, and of a bitterish, unpleasant taste.

The meal is sometimes employed in ca- taplasms.

2. *Vicia Sativa*, LINN.

SYNONIMES:—Vesce, FR.; Futterwicke, GERM.

GALL.

A plant originally of the Levant, and cul- tivated throughout Europe. (*Fig. CEd. Flor. Dan.* t. 522)

The seeds (*Semen Viciæ*) are employed. They are black and compressed.

VINCA.—PERIWINKLE.

Two species of this genus of plants are mentioned in pharmacopœias :

1. *Vinca major*, LINN.—Great Periwinkle.

SYNONIMES :—Pervenche majeure, Grande Pervenche, FR.

GALL.

A plant of the south of Europe. (*Pentandria Monogynia*, LINN. ; *Apocynæ*, RUSS. ; *Fig. Nouv. Duh*, I. 14)

The herb is employed ; it consists of oval leaves, slightly ciliated at the borders, and heart-shaped at the base.

2. *Vinca minor*, LINN.—Small Periwinkle.

SYNONIMES :—Pervenche mineure, Petite

Pervenche, FR. ; Sinngrün, GERM. ; Vingrøn, DAN. ; Pervinca, ITAL. ; PORT. ; and SPAN. ; Vin koorde, DUT. ; Barwinek, POL.

BRUNS. ; HISP. ; GALL. ; WIRTEM. ; BERGIUS ; MURRAY and GMELIN ; PIDERIT ; SPIELMANN.

A plant common to the greater part of Europe. (*Fig. Zorn, Ic. pl. t. 67*)

The herb (*Herba Vincæ seu Pervincæ seu Clematidis seu Daphnoidis*) is employed. It consists of a slender and sarmentose stem, and of oblong-ovate, glabrous, shining leaves, supported on very short footstalks. It is inodorous, but has a bitter and astringent taste.

These two plants are slightly astringent.

VINUM.—WINE.

SYNONIMES :—*Ænos* ; *Óivos*.—Vin, FR. ; Wein, GERM. ; Khumar, ARAB. ; Winny, BOHEM. ; Scherabungury, DUK. ; Vino, ITAL. ; and SPAN. ; Drakhka mud, HIND. ; Mey, PERS. ; Wino, POL. ; Winho, PORT. ; Mada, Madira, SANS. ; Win, SWED.

LOND. ; DUBL. ; EDINB. ; AMER. ; AMST. ; ANT. ; BATAV. ; BAVAR. ; BELG. ; BRUNS. ; DAN. ; HISP. ; GALL. ; FENN. ; FERR. ; FORM. HÔP. FR. ; FULD. ; GENEV. ; HANN. ; HASS. ; LIPP. ; OLDEN. ; LUSIT. ; POL. ; CAST. BORUS. ; BORUS. ; SAX. ; SUEC. ; AINSLIE ; BERGIUS ; COXE ; MURRAY and GMELIN ; PARIS ; SAUNDERS.

A red or white liquor obtained by the fermentation of the juice or must of grapes.

Wine is a compound of water, alcohol, mucilage, acetic acid, supertartrate of potass, tartrate of lime, and colouring principles, the diversities of the respective proportions of which produce the almost innumerable varieties of this liquor which are met with in commerce.

The action of wine upon the animal economy varies considerably, according as alcohol, sugar, acetic acid, or carbonic acid may predominate in it. But in general, only those are employed medicinally which are rich in alcohol and in principles which give a more or less austere quality, such as those of Bordeaux, Burgundy, Alicante, Madeira, and Xeres, which act at the same

time as tonic and stimulant. However, it is rare that wine is prescribed alone as a medicament ; but more frequently serves as a vehicle to other more active substances.

DESICCATIVE CATAPLASM. (FERR.)

R. Lupin Meal,	eight ounces.
Rye Meal,	four ounces.
White Wine,	four pounds.

Boil till of a proper consistence.

VINOUS FOMENTATION. (FORM. HÔP. FR. ; RATIER)

R. Red Wine,	one pound.
Honey,	two ounces.

Dissolve. (RATIER)

R. Compound Tincture of Cinnamon,	one part.
Red Wine,	twenty parts.
Mix. (FORM. HÔP. FR.)	

SAMARITAN BALSAM. (CADET DE GASSICOURT ; PIERQUIN ; SPIELMANN)

R. Wine,	} of each, an equal part.
Olive Oil,	
Mix.	

VINOUS POTION.

Posca Galeni. (ELLIS ; RATIER ; SAINTE-MARIE ; SPIELMANN)

R. Wine,	three parts.
----------	--------------

Water, two parts.
Boil a little. (SPIELMANN)
R. Spanish Wine, } of each,
Spring Water, } two drams.
Syrup of Gooseberries, }
Mix. (SAINTE-MARIE)
R. White Wine, } of each, one pound.
Water, }
Alcohol, } of each, one ounce.
Tartaric Syrup, }
Mix. (RATIER)
R. Pounded Ice, a saucerful.

Madeira Wine, a spoonful.
Hordeated Cinnamon Water, a tea-spoonful.
Sugar, any quantity.
Mix. (SAINTE-MARIE)
R. Yolk of Egg, one.
Oil of Cinnamon, twenty drops.
White Wine, } of each,
Cinnamon Water, } three ounces.
Pure Water, two ounces.
Sugar, two drams.
Mix. (ELLIS)
Dose, three spoonfuls.

VIOLA.—VIOLET.

Four species of this species of plants are used in medicine:

1. *Viola pedata*, LINN.

SYNONIME:—Violette digitée, FR.

AMER.; COXE.

A plant of North America. (*Pentandria Monogynia*, LINN.; *Violaceæ*, JUSS.; *Fig. Pu. Fl. Amer.* p. 171)

The herb is employed; it consists of palmate, pectinate, stipulate leaves.

A plant growing in all parts of Europe. (*Fig. Flore Medic.* VI. 369)

The root, herb, flowers, and seeds are employed.

The root (*Radix Violariæ seu Martiæ*) is composed of a great many tufted fibres. It has an acrid and nauseous taste.

The herb presents a bunch of cordiform radical leaves, finely crenelated, green, glaucous, obtuse or a little pointed at the summit. It is inodorous, and has an herbaceous, mucous, and bitterish taste.

The flowers are bluish; they have a pleasant odour which is removed by desiccation; their taste is bitter and slightly mucilaginous.

The seeds are small, round, and whitish. The whole plant is more or less stimulant, nauseant, and emetic.

2. *Viola Canina*, LINN.—Dog-violet.

SYNONIMES:—Violette sauvage, FR.; Hundsvielfchen, GERM.; Wilde fioler, DAN.; Wilde viol, DUT.; Skogsvioler, SWED.

GALL.; MURRAY and GMELIN.

A very common plant of Europe. (*Fig. Curt. Fl. Lond.* t. 108)

The root (*Radix Viola caninæ*) is acrid and nauseous.

DISTILLED WATER OF VIOLET.

Aqua Violarum. (FERR.; PALAT.; SARD.)

R. Flowers of Violet, one part.
Water, four parts.

At the end of six hours, distil two parts. (SARD.)

PALAT. prescribes one part of flowers, and three parts of water; one part and a half to be distilled.—FERR.—three of flowers and sixteen of water; eight to be distilled.

3. *Viola odorata*, LINN.—Violet, Sweet Violet.

SYNONIMES:—Violette de Mars, FR.; Märzveilchen, GERM.; Manischar, ARM.; Fiola, wonna neh modra, BOHEM.; Keet-tuong-hoa, CHIN.; Martsfioler, DAN.; Violetta, PORT.; and SPAN.; Tamme viol, DUT.; Viola, ITAL.; Fiolki, POL.; Pachutschaja, RUSS.; Ækta fioler, SWED.

EDINB.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FERR.; FULD.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; WIRTEM.; HERBIP.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT.; MURRAY and GMELIN.; SPIELMANN.; ZARDA.

VIOLET JULEP.

Julapium Violarum. (PALAT.; WIRTEM.; SPIELMANN)

R. White Sugar, two pounds and a half.
Water of Violets, twenty ounces.

Boil for an instant, and strain.

EMULSION OF VIOLET.

Emulsio violata. (WIRTEM.; SPIELMANN)

- R. Sweet Almonds, one ounce.
 Seeds of Violet, two drams.
 ——— White Poppy, half a dram.
 Water of Violet, six ounces.
 ——— Water Lily, } of each,
 ——— Lettuce, } three ounces.

Add to the emulsion,

Syrup of Violet, one ounce.

Mix.

JUICE OF VIOLET.

Succus Violarum. (HISP.; SARD.)

- R. Flowers of Violet, freed from
 their calices, any quantity.

Bruise them in an earthen mortar; then, having expressed the juice, set it apart in a cool place, and after a short time pour off the clear part.

VIOLET LOZENGES.

Tabellæ violatæ. (SARD.)

- R. Juice of Violet, three ounces.
 White Sugar, two pounds.
 Florentine Iris, in powder, one ounce.

Make lozenges.

CONSERVE OF VIOLET.

Conserva Violarum. (HISP.; PALAT.; SARD.; WIRTEM.)

- R. Fresh Flowers of Violet, one part.
 White Sugar, two parts.

Beat together. (PALAT.; WIRTEM.)

SARD. directs one part of petals to be triturated with a little sugar, and thrown into three parts of syrup, and the mixture boiled for a few moments.—HISP.—one of petals to be bruised, added to two of thick syrup, and evaporated to the proper consistence.

INFUSION OF VIOLET. (SARD.; DISP. CHA. PARIS; RATIER)

- R. Flowers of Violet, two drams.
 Boiling Water, two pounds.
 Infuse, and strain. (RATIER)

- R. Flowers of Violet, two pounds.
 Boiling Water, eight pounds.

Digest for eight hours, and strain with slight expression. Repeat this operation nine times successively with fresh flowers. (SARD.)

- R. Flowers of Violet, } of each,
 Liquorice, } two drams.
 Boiling Water, one pound and a half.
 Strain when cold. (DISP. CHA. PARIS)

It is recommended in inflammations of the chest.

HONEY OF VIOLET.

Mel Violarum. (PALAT.; SARD.; WIRTEM.)

- R. Flowers of Violet, } of each,
 Boiling Water, } six pounds.

Macerate for twenty-four hours. Add to the expressed liquor,

Honey, six pounds.

Boil, skimming, to the consistence of syrup. (PALAT.; WIRTEM.)

- R. Depurated Juice of } of each,
 Violet, } an equal part.
 Honey, }

Boil, skim, and strain; then evaporate it to the consistence of syrup. (SARD.)

It used formerly to be employed in lavesments in the dose of from one ounce and a half to two ounces.

SYRUP OF VIOLETS.

Syrupus Violarum seu Violæ odoratæ seu Petalorum Violarum. (EDINB.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FULD.; GENEV.; HANN.; HASS.; LIPP.; OLDEN.; PALAT.; POL.; BORUS.; ROSS.; SARD.; WIRTEM.; HERBIP.; COXE; SWEDIAUR; VAN MONS)

Syrupus Violæ. (DUBL.)

- R. Fresh Petals of Violets, two pounds.
 Boiling Water, five pints.

Macerate for twenty-four hours; then express the liquor through fine linen. Add to each pint of it twenty-nine ounces of refined sugar, and form a syrup.

Syrupus Violæ odoratæ. (EDINB.; COXE)

- R. Flowers of Violet, two parts.
 Boiling Water, eight parts.
 Refined Sugar, fifteen parts.

Macerate the flowers in the water for twenty-four hours, in a covered glass or glazed earthen vessel; then strain without expression, and add the sugar.

- R. Fresh Petals of Violet, eight ounces.
 Boiling Water, sixteen ounces.

Digest for twenty-four hours at a moderate heat; then strain with very slight expression, and add to eight parts of the liquor,

White Sugar, fifteen parts.
Make a syrup. (AMST.)

The proportions of this formula varies very much;—BATAV.; and BELG. prescribe eight ounces of petals, twenty ounces of boiling water, six hours' maceration, and three pounds of sugar.—SARD.—two parts of infusion and one part of sugar.—BAVAR.—one of petals, three of water, twelve hours' infusion, and two parts of sugar to one part of liquor.—GALL.—four pounds of petals, eight of water, twelve hours' infusion, and two parts of sugar to one part of strained liquor.—SWEDIAUR.—one part of petals, two parts of water, twenty-four hours' infusion, and five parts of sugar to three of liquor.—HISP.—one pound of petals, two pounds of water, eight hours' infusion, and four pounds of sugar.—HASS.—eight ounces of petals, twenty of water, six hours' infusion, and three pounds of sugar.—HANN.; and BORUS.—the same proportions, but twelve hours' infusion.—OLDEN.; and POL.—eight ounces of petals, two pounds of water, twelve hours' maceration, and three pounds of sugar to twenty ounces of strained liquor.—HERBIP.—one pound of petals, three pounds of water, and seven of sugar.—ROSS.—one of petals, three of water, and two of sugar to one of strained liquor.—FULD.—twelve ounces of petals, eighteen of boiling water, and twelve hours' digestion; twelve other hours' maceration of the expressed liquor with six ounces of fresh petals, and the addition of twelve ounces of sugar.—FERR.—one pound of petals, one of water, two days' digestion, and two pounds of sugar.

HISP. prescribes, besides the syrup above cited, another called *Syrupus Violarum communis*, prepared with one pound of petals infused in four pounds of boiling water, and six pounds of sugar.

R. White Sugar, four pounds.
Spring Water, a sufficient quantity.

Boil to the consistence of a thick syrup, and pour it upon

Fresh Petals of Violet, two pounds.

Macerate for three days in a covered vessel, then boil a little, and express strongly. (GENEV.)

R. Fresh Petals of Violet, one pound.
Bruise in a stone mortar, adding

Boiling Water, eighteen ounces.

Digest for twenty-four hours. Add to twenty ounces of the strained liquor,

White Sugar, thirty-two ounces.
Boil slowly. (BRUNS.; PALAT.; WIRTEM.)

ANT. prescribes one pound of bruised petals, two pounds of water, twenty-four hours' digestion, and three pounds and a half of sugar.—LIPP.—one pound of petals to be bruised, and digested with heat for twelve hours, in three pounds of water, and a syrup made with the infusion and twenty ounces of sugar.

APERIENT SYRUP OF VIOLETS.

Syrupus Violarum solutivus. (WIRTEM.)

R. Petals of Violet, four ounces.
Senna, two ounces.
Boiling Spring Water, one pound.

Digest for one night in a warm place, and express in the morning. Add to nine ounces of the strained liquor.

White Sugar, sixteen ounces.
Dissolve.

OIL OF VIOLET.

Oleum Violarum. (BRUNS.; SARD.; WIRTEM.)

R. Petals of Violet, one part and a half.
Olive Oil, three parts.

Digest in a water-bath for two days; then express, and repeat the digestion with fresh petals. (SARD.)

R. Petals of Violet, } of each,
Olive Oil, } an equal part.

Boil slowly to the consumption of the humidity, and express. (BRUNS.; WIRTEM.)

4. *Viola tricolor*, LINN.—Three-coloured Violet, Heartsease.

SYNONIMES: Pensée, FR.; Dreyfaltigkeitsblume, Stiefmutterchen, Freysamkraut, GERM.; Stedmodersblomst, DAN.; Pensamientos, SPAN.; Driekleurige viol, DUT.; Fiore del pensiero, ITAL.; Fiolek potroyny, Bratzsiostra, POL.; Amor perfetto, PORT.; Wdowski, RUSS.; Styfmorsblomster, SWED.

AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GALL.; FERR.; FENN.; FULD.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; BERGIUS.; BRUGNATELLI.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN.

This plant is met with in every part of Europe. (*Fig. Zorn, Ic. pl. t. 290*)

The herb (*Herba Jaceæ seu Violæ tricoloris seu Trinitatis*) is employed. It consists of a branching, triangular stem, and of alternate, dentate, oval leaves having pinnatifid stipulæ. It has a somewhat acrid taste.

It is excitant, depurative, and diuretic, and is employed principally in herpes and crusta lactea.

EXTRACT OF HEARTSEASE. (FENN.)

- R. Herb of Heartsease, one part.
Boiling Water, six parts.

Boil for a quarter of an hour, and express; boil the residue with fresh water, and express again: then mix and evaporate the decoctions.

AQUEOUS DECOCTION OF HEARTSEASE.

Decoctum Herbae Viola tricoloris cum Aqua.
(NIEMANN)

- R. Herb of Heartsease, six drams.
Spring Water, a sufficient quantity,
to obtain six ounces of strained decoction. It is employed against crusta lactea, herpes, and tinea. — Dose, a spoonful every two hours.

LACTEAL DECOCTION OF HEARTSEASE.

Decoctum Herbae Viola tricoloris cum Lacte.
(AUGUSTIN; NIEMANN)

- R. Herb of Heartsease, half an ounce.
Cow's Milk, one pound.
Boil a little, and strain.

It is recommended in the same cases as the preceding. The dose prescribed should be repeated morning and evening.

INFUSION OF HEARTSEASE.

Infusum Herbae Jaceae. (AUGUSTIN)

- R. Herb of Heartsease, two drams.
Boiling Water, two ounces.
Infuse; add to the strained liquor,

Tartarized Antimony, one scruple.
Syrup of Marshmallows, two drams.

Mix. — Dose, one or two tea-spoonfuls every two hours.

- R. Herb of Heartsease, } of each,
Sugar of Milk, } two drams.
Precipitated Sulphur, } of each
Subcarbonate of Mag- } half a dram.
nesia,

Mix. — Dose, a scruple every two hours.

DEMULCENT POTION.

Mixtura in Hematuria chronica. (BRERA)

- R. Hemp Seed, one ounce.
Water, a sufficient quantity,
to obtain one pound of decoction. Add towards the end,

Flowers of Heartsease, half an ounce.

Infuse for some time, and add to the strained liquor,

Syrup of Marshmallows, one ounce.

Mix.

VISCUM ALBUM, LINN.—MISLETOE.

SYNONIMES:—Gui, Gui blanc, FR.; Mistel, GERM.; and SWED.; Dubowe melj, BOHEM.; Fugleljum, DAN.; Visco, SPAN.; Maréntakken, DUT.; Vischio, ITAL.; Jemiel, POL.; Visgo, PORT.

AMST.; AUSTR.; BAVAR.; BRUNS.; HISP.; GALL.; FERR.; GENEV.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; SAX.; WIRTEM.; HERBIP.; BERGIUS; BRÜGNATELLI; GUIBOURT; MURRAY and GMELIN; SPIELMANN; ZARDA.

A common small parasitical shrub of Europe. (*Diacia Tetrandria*, LINN.; *Loranthæa*, JUSS.; Fig. Zorn. Ic. pl. t. 54)

The stems (*Viscum quernum* seu *quercinum* seu *Lignum Visci*) are employed. They are ligneous, cylindrical, divided, almost from the base, into numerous dichotomous branches, of a pale green or yellowish colour. Their odour is feeble, peculiar, and disagreeable; their taste repug-

nant, rather acerb and mucilaginous: both disappear after desiccation.

It is astringent, nauseant, and emetic, and has been for a long time celebrated against epilepsy. — Dose of the powder, from half a dram to a dram, three or four times a day, in syrup. COLBATH combines it with with assafetida as HUFELAND and TISSOT did with valerian.

ANTI-SPASMODIC POWDER.

Pulvis anti-spasmodicus infantum. (AUGUSTIN; HUFELAND)

- R. Crab's Eyes, } of each,
Hartshorn Shavings, } an equal part.
Root of Valerian, }
Mistletoe,

Make a powder. — It is recommended in diarrhoea, emesis, and difficult dentition. — Dose, half a scruple. (HUFELAND)

R. Mistletoe, } of each,
 Root of Peony, } two drams and
 — Fraxineila, } a half.
 Aloe Wood, } of each, one dram.
 Crabs' Eyes, }
 Amber, half a dram.
 Charcoal, two drams.
 Mix.

AQUEOUS EXTRACT OF MISLETOE.

Extractum Visci albi aquosum. (FERR.)

R. Mistletoe, one pound.
 Water, three pounds.

Boil, and express; clarify the decoction with white of egg, and evaporate to the proper consistence.

VINOUS EXTRACT OF MISLETOE.

Extractum Visci albi vinosum. (SARD.)

R. Mistletoe, one pound.
 White Wine, five or six pounds.

Digest for three days in a sand-bath, and express strongly; then boil the residue with a sufficient quantity of fresh wine. Evaporate the mixed liquors to the consistence of an extract.

DECOCTION OF MISLETOE.

Decoction Visci albi. (NIEMANN)

R. Mistletoe, one or two ounces.
 Spring Water, three pounds.

Boil down to one half.

It is recommended in epilepsy.

VITEX AGNUS CASTUS, LINN.—AGNUS CASTUS.

SYNONIMES: — Agnus-castus, Gattilier, Petit-poivre, Poivre sauvage, FR.; Keuschbaum, Schaafmuller, Keuschlamm, GERM.; Agno casto, ITAL.; and SPAN.; Kuischboom, DUT.

BRUNS.; HISP.; GALL.; WIRTEM.; BERGIUS.; GUIBOURT.; MURRAY and GME-LIN.; SPIELMANN.

A shrub of the south of Europe. (*Didy-*

namia Angiospermia, LINN.; *Verbenaceæ*, Juss.; *Fig. Flore Medic.* I. 8)

The fruit (*Semen Agni casti*) is employed. It consists of a globular, blackish, hard berry of the size of a peppercorn or larger, enveloped at its base by the calix of the flower, and divided interiorly into four monospermal cells. It has an aromatic and somewhat stupefactive odour, and a bitter, acrid, and aromatic taste.

This plant was formerly erroneously esteemed refrigerant and aphrodisiac.

VITIS VINIFERA, LINN.—VINE.

SYNONIMES: — Vigne, FR.; Weinstock, GERM.; Inub, ARAB.; Otrik, ARM.; Kmen winny, BOHEM.; Pu-tao, CHIN.; Sur medan, CALM.; Well mudika gheddie, CYNG.; Wuntrac, DAN.; Ungur, DUK.; and PERS.; Vid, SPAN.; Dakh, HIND.; Wyngaard, DUT.; Vite, ITAL.; Buangur, MAND.; Winna macica, POL.; Videira, PORT.; Winegrad, RUSS.; Drakscha, SANS.; Vinstock, SWED.; Kodimundrie pullum, Dividatsi pullum, TAM.; Drachupundu, TEL.

A shrub supposed to be originally of Asia. (*Pentandria Monogynia* LINN.; *Vinifera*, Juss.; *Fig. Zorn, Ic. pl. t. 276*)

The leaves and fruit are employed. The leaves are large, at first hairy above and tomentose and white beneath, then glabrous; they are heart shaped at the base, divided

into from three to five toothed, incised, and acute lobes.

They are considered tonic.

The fruit, too well known to require description, is called *Grape*; *Uva*.

SYNONIMES: — Raisin, FR.; Rosinen GERM.; Zibib, ARAB.; and MAND.; Velitcha mudika gheddie, CYNG.; Kischmisch, DUK.; and HIND.; Pasas, SPAN.; Razynen, DUT.; Mewuz, PERS.; Uvas passadas, PORT.; Russin, SWED.; Dividatsi palavuttil, TAM.

HISP.; GUIBOURT.; MURRAY and GME-LIN.

Before its maturity it is called (*Verjus*, FR.; *Verjuice*; *Omphacium*, *Agresta*. It then has an acid and astringent taste. It is seldom employed, and only in cases where acids appear to be indicated.

When ripe it has a sweet and saccharine taste. Its expressed juice is called (Mout, Fr.;) *Must*; *Mustum*.

Under this form it belongs less to medicine, properly so called, than to dietetics. But when dried in the sun grapes are of very frequent use; they are then called *Raisins*. There are two kinds of them:

1. *Large Raisins*; *Passula majores*, *Uva passa*, *Zibeba*.

LOND.; EDINB.; DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HASS.; LUSIT.; ROSS.; SUEC.; SAX.; WIRTEM.; HERBIP.; BERGIUS; COXE; GUIBOURT; MURRAY and GMELIN: PIDERIT; SPIELMANN.

There are several varieties of them:

A.

Smyrna or *Damask Raisins*, nearly of the size of small plums, longish, compressed, wrinkled, and of a brownish-yellow colour.

B.

French or *Spanish Raisins*, smaller and thicker, with a brownish-colour.

C.

Calabrian Raisins, thick, of a deep colour, and viscous taste.

2. *Currants*, *Corinth Raisins*; *Passula minores* seu *Corinthiaca* seu *Corinthi-nacea*.

AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; GENEV.; FERR.; FULD.; HAMB.; HASS.; LUSIT.; ROSS.; SAX.; SUEC.; BERGIUS; GUIBOURT; MURRAY and GMELIN; PIDERIT; SPIELMANN.

These are black, of the size of small lentils, and without stones.

All these are emollient.

PULP OF RAISINS.

Pulpa passularum. (BRUNS.; PALAT.; WIRTEM.; HERBIP.)

- R. *Corinth Raisins*, any quantity.
Water, a sufficient quantity.

Soften them by boiling; then strain the pulp through a sieve, add water to it, and evaporate the whole to a proper consistence. (GALL.)

BRUNS.; PALAT.; and WIRTEM. direct one half of sugar to be added to the pulp.—HERBIP.—one third of sugar.

ROB OF GRAPES.

Roob seu Sapa Uvarum. (GALL.; SARD.; HERBIP.)

- R. *Grape Juice*, any quantity.

Boil it over a slow fire to the consistence of honey. (GALL.; SARD.)

- R. *Depurated Grape Juice*, one pound.
White Sugar, three ounces.

Boil together to the consistence of honey, (HERBIP.)

DECOCTION OF RAISINS.

Decoctum Passularum majorum. (AUGUSTIN; NIEMANN)

- R. *Raisins*, washed and freed from stones, four ounces.
Spring Water, three pounds.

Boil down to two thirds, and strain. (NIEMANN)

- R. *Raisins*, four ounces.
Pearl Barley, } of each,
Liquorice, } half an ounce.
Water, six pounds.

Boil, and add to the strained liquor when cold,

- Syrup of Lemon Juice*, two ounces.

Mix. (AUGUSTIN)

SYRUP OF GRAPE.

Syrupus Uvarum. (AUSTR.)

- R. *Recently expressed Grape Juice*, one hundred pounds.

Boil in a tinned copper vessel till no more scum rises; then remove it from the fire, and add gradually, constantly stirring,

Prepared Chalk, one pound,
or, a sufficient quantity for perfect saturation. Set apart for twelve hours, in a cool place; then pour off the clear part, and boil it, constantly stirring to the consistence of syrup.

PECTORAL SPECIES. (BRUNS.; PALAT.; WIRTEM.; SPIELMANN)

- R. *Corinth Raisins*, one ounce.
Carob Beans, } of each, six drams.
Jujubes, }
Dates, two ounces.
Figs, } of each, one ounce.
Pearl Barley, }
Liquorice, } of each,
Montpellier Maiden-hair, } half an ounce.
Hyssop, two drams.

Cut, and mix. (BRUNS.; WIRTEM.; SPIELMANN)

PALAT. gives the same formula, but substitutes sebestes for the carob beans.

JUICE OF VERJUICE.

Succus Agrestæ, Omphacium. (HISP.; GALL.; SARD.)

R. Unripe Grapes, any quantity.

Bruise them in a stone mortar, and strain the juice through flannel; then set it apart, decant, and preserve it in a long-necked bottle, under a stratum of oil.

SYRUP OF VERJUICE.

Syrupus Agrestæ seu de Agrestæ. (HISP.; GALL.; PALAT.; SARD.; WIRTEM.)

R. Juice of Verjuice, two pounds.
White Sugar, three pounds and a half.

Dissolve in a gentle heat. (GALL.)

WIRTEM. prescribes one pound and a half of juice, and two pounds of sugar.—PALAT.—five pounds of juice, and three of sugar.—HISP.—one of juice, and two of sugar.—SUEC.—a sufficient quantity of juice and of simple syrup to form a syrup by evaporation in a water-bath.

ASTRINGENT OINTMENT. (BORIES)

R. Juice of Verjuice, eight ounces.
Rose Ointment, one pound.
Yellow Wax, four ounces.

Boil to the consumption of the humidity; then set the mixture apart to solidify, separate the fæces, melt again, and flavour with essence of roses.

WRIGHTEA ANTI-DYSENTERICA, BR.—CONESSI; OVAL-LEAVED ROSE BAY.

SYNONIMES:—Codagapala, FR.; Konessrinde, Conessrinde, GERM.; Curaija, HIND.; Codaga pala, MAL.; Kutaja, SANS.; Veppalei, TAM.; Pala codija, Manupala, TEL.

BRUNS.; GALL.; FERR.; WIRTEM.; HERBIP.; AINSLIE; MURRAY and GMELIN; PARIS.

A tree of the East Indies. (*Pentandria Monogynia*, LINN.; *Apocynæ*, JUSS.; Fig. Rheed. *Malab.* l. p. 85. t. 47)

The bark of the trunk, and particularly of the root (*Cortex Codagapala seu Profluvii seu Conessi*) is employed. It is in flat or quilled pieces of the thickness of a line, blackish without, and covered with grey lichens. It has an astringent and bitter taste.

It is tonic, and was formerly used in diarrhoea and dysentery.

XANTHIUM STRUMARIUM, LINN.—LESSER BURDOCK.

SYNONIMES:—Lampourde, Petit Glouteron, Petite Bardane, FR.; Kropfklette, Kropfbetterlaus, Klissen, GERM.; Gaasekreppe, DAN.; Lampazo pequeno, SPAN.; Kleine klissen, DUT.; Lappola minore, ITAL.; Bardane menor, PORT.; Spetsborre, SWED.

BRUNS.; GALL.; LUSIT.; WIRTEM.; MURRAY and GMELIN; SPIELMANN.

A common plant of Europe. (*Monœcia Pentandria*, LINN.; *Synanthereæ*, CASS.; Fig. Zorn, *lc. pl.* t. 269)

The herb and seeds are employed.

The herb (*Herba Xanthii seu Bardane*

seu Lappæ minoris) consists of an angular, branching stem, and of alternate, petiolated, rounded leaves, cordiform, toothed, sinuated, somewhat lobed, pubescent, scabrous, trinerfed at the base.

The seeds are small, oval, pointed at both ends, flat on one side, bulged on the other, reddish, and contained in a common closed, oblong involucre truncate at the summit, monophylous, and bristled over with points externally.

The odour is disagreeable, and the taste acrid and bitter.

The juice was formerly prescribed in goitre, in the dose of six ounces.

ZANTHORRHIZA APIFOLIA, L'HER.—YELLOW ROOT, PARSLEY-LEAVED YELLOW ROOT, YELLOW ROOT SHRUB.

AMER.; COXE.

A small shrub of Virginia. (*Pentandria Monogynia*, LINN.; *Ranunculaceæ*, JUSS.; *Fig. L'Her, St. nov. p. 79. t. 38*)

The root is long, of the thickness of the little finger, and furnished with a great many fibrils. It is of a yellow colour, and has a bitter taste.

ZANTHOXYLUM CLAVATUM, LINN.—PRICKLY ASH, TOOTH-ACHE TREE.

SYNONIMES:—Clavallier à feuilles de frêne, Frêne épineux, FR.; Zahnwehholz, GERM.

AMER.; COXE.

A tree of the Antillas. (*Diœcia Pentandria*, LINN.; *Terebinthaceæ*, JUSS.)

The bark, which is brown, is employed on account of its acridity against tooth-ache. The tincture is used, in America, in chronic rheumatism.

ZEA MAYS, LINN.—INDIAN CORN, MAIZE.

SYNONIMES:—Mais, Blé d'Espagne, Blé d'Inde, Blé de Turquie, FR.; Türkischer Weizen, GERM.; Tyrkisk korn, DAN.; Maiz, SPAN.; Mays, DUT.; Gran Turco, ITAL.; Maiz, Milho de Turquia, PORT.; Türkisk korn, SWED.

GALL.

An original plant of America. (*Monœ-*

cia Triandria, LINN.; *Gramineæ*, JUSS.; *Fig. Blackw. Herb. t. 547*)

The seeds (*Semen Mays*) are employed. They are round, angulose, of the size of a pea, of a golden yellow colour externally, and white and brilliant internally.

It is farinaceous and alimentary.

ZEDOARIA.—ZEDOARY.

SYNONIMES:—Zédoaire, FR.; Zittewurzel, Zitteringwer, Gillwerwurzel, GERM.; Judwar, Bar, ARAB.; Bhuchampa, Junglie huldie, Bun huldie, BENG.; Cycwar, BOHEM.; Nagai mio, COCHIN.; Kaha, Walkaha, CYNG.; Ambie huldie, DUK.; Zedoaria, SPAN.; Nirbisi, HIND.; Bokzedoar, DUT.; Malan kua, MALAB.; Cytwar, POL.; Bhu champaka, nervischa, Vanaharidra, SANS.; Sittwer, SWED.; Casturie munjel, TAM.; Castorie pasapu, TEL.

DUBL.; AMST.; ANT.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENÈV.; HAMB.; HANN.; LIPP.; OLDEN.; POL.; BORUS.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; AINSLIE.; BERGIUS.; BRUGNATELLI.; COXE.; GUIBOURT;

MURRAY and GMELIN.; PIDERIT.; SAUNDERS.; SPIELMANN.; ZARDA.

There are two kinds of Zedoary:

1. The *long*, *Zedoaria longa*. This is several inches long, of the thickness of the finger, wrinkled, knotty, of a greyish colour without, and brownish within.

2. The *round*, *Zedoaria rotunda*. This is about an inch long, rough at the surface, rugose, and covered with fibres.

Zedoary comes to us in hard, whitish pieces, deprived of its external pellicle. It is aromatic, acrid, and pungent.

It contains much starch, and a camphorous essential oil.

The long zedoary is usually considered as the root of the *Amomum Zedoaria*, WILLD.; a plant (*Monandria Monogynia*, LINN.;

Amomeæ, JUSS.) of the East Indies (*Fig. Flore Medic. VI. 349*)

As to the round, some authors, with whom are AINSLIE and GUIBOUT, think that it belongs to the *Kampferia rotunda*, LINN.—Others think, and such is the opinion of MERAT, that they are simply varieties of the root of one plant, dependant on locality and mode of growth.

It is excitant, tonic, and stomachic.—Dose, from half a scruple to half a dram, of the powder.

EXTRACT OF ZEDOARY.

Extractum Zedoariæ. (BRUNS.; PALAT.; SARD.; WIRTEM.)

R. Root of Zedoary, one pound.
Alcohol, five or six pounds.

After sufficient extraction, express; digest the residue, and boil it a little with

Water, three pounds.

Clarify the decoction with white of egg, then add the tincture, and having distilled off the alcohol, evaporate the remainder to the proper consistence. (BRUNS.; PALAT.; WIRTEM.)

R. Zedoary, one pound.
White Wine, five or six pounds.

Digest in a sand-bath for three days, and strain; boil the residue with a sufficient quantity of new wine. Mix the liquors, and evaporate slowly to the consistence of an extract. (SARD.)

TINCTURE OF ZEDOARY.

Tinctura seu Essentia Zedoariæ. (AMST.; BRUNS.; HANN.)

R. Root of Zedoary, one part.
Alcohol (0.907), eight parts.

Digest for eight days, and filter. (AMST.)

BRUNS. prescribes one part of root, and five parts of alcohol.—HANN.—five ounces of root, and two pounds of alcohol.

It is excitant.—Dose, sixty drops, twice a day.

COMPOUND TINCTURE OF ZEDOARY.

Tinctura Zedoariæ composita, Essentia carminativa Wedelii. (BRUNS.; HANN.; PALAT.; SAX.; WIRTEM.; CADET DE GASSICOURT; NIEMANN; SPIELMANN)

R. Root of Zedoary, four ounces.
——— Carline Thistle, } of each,
——— Sweet Flag, } two ounces.
——— Galanga, }
Flowers of Chamomile, } of each,
Seeds of Anise, } one ounce.
——— Caraway, }
Cloves, } of each, six drams.
Bay Berries, }

Mace, half an ounce.
Orange Peel, one ounce.
Spirit of Lemon Peel, four pounds and a half.
Spirit of Nitric Æther, two ounces and a half.

Infuse for six days; then express, and filter. (BRUNS.; HANN.; SAX.; WIRTEM.; NIEMANN; SPIELMANN)

R. Root of Zedoary, one ounce.
——— Carline Thistle, } of each,
——— Galanga, } half
an ounce.
Orange Peel, }
Seeds of Anise, } of each,
——— Caraway, } one dram
Flowers of Chamomile, } and a half.
Cloves, }
Bay Berries, }
Sweet Flag, }
Mace, one dram.
Spirit of Lemon Peel, two pounds.
Nitric Acid, half an ounce.

After sufficient digestion, filter. (PALAT.; NIEMANN)

R. Root of Zedoary, one ounce.
——— Carline Thistle, } of each,
——— Sweet Flag, } half
——— Galanga, } an ounce.
Flowers of Chamomile, }
Seeds of Anise, } of each,
——— Caraway, } two drams.
Dried Orange Peel, }
Bay Berries, } of each,
Cloves, } one dram and a half.
Mace, one dram.
Spirit of Lemon, one pound.
Nitric Acid, five drams.

It is excitant, reputed stomachic, carminative, emmenagogue, and diuretic.—Dose, from half a dram to one dram.

BEZOARDIC TINCTURE.

Tinctura bezoardica. (PALAT.; WIRTEM.; SPIELMANN)

R. Root of Zedoary, }
——— Serpentaria, }
——— Angelica, }
——— Burnet Saxifrage, } of each,
Herb of Scordium, } six
——— Dittany of Crete, } drams.
Juniper, }
Lemon Peel, }
Myrrh, }
Rectified Spirit of Elder Flowers, one pound.

Digest for a sufficient time. Add to the expressed liquor

Pyrotartaric Acid, six ounces.
Sulphuric Acid, two ounces.
Camphor, half an ounce.

Digest, filter, and preserve. (PALAT.; WIRTEM.)

R. Root of Zedoary,	two ounces.
— Angelica,	} of each, one ounce.
— Burnet Saxifrage,	
— Valerian,	
— Swallow-wort,	
— Sweet Flag,	} of each, two ounces.
— Costus,	
Fresh Lemon Peel,	
Juniper,	
Myrrh,	three ounces.
Herb of Scordium,	one ounce and a half.
— Balm,	} of each, seventy-two ounces.
— Milfoil,	
— Rue,	
Camphor,	one ounce.
Theriac,	one pound.
Alcohol,	six pounds.

Digest for eight days, and add

Spring Water, two pounds,
and distil. Then

R. Of this Spirit,	two pounds.
Pyrotartaric Acid,	half a pound.
Sulphuric Acid,	three ounces.

Mix, colour with oneberry juice, and
prescribed in putrid fevers.—Dose, half a
dram.

ESSENTIAL OIL OF ZEDOARY.

Oleum Zedoariae athereum, Aetheroleum Zedoariae. (BRUNS.; PALAT.; WIRTEM.)

R. Zedoary,	six parts.
-------------	------------

Water,	thirty-six parts.
Muriate of Soda,	one part.

Macerate for four or five days, and distil. Collect the oil at the bottom of the receiver.

VINOUS SPIRIT OF ZEDOARY.

Aqua Zedoariae vinosa. (BRUNS.; WIRTEM.)

R. Zedoary,	one part.
Wine,	twelve parts.

Distil one half. (WIRTEM.)

R. Zedoary,	six ounces.
Wine,	three pounds.
Alcohol,	} of each, one pound and a half.
Water,	

Distil four pounds. (BRUNS.)

ANISATED SPIRIT OF ZEDOARY.

Aqua Zedoariae anisata. (PALAT.; SARD.; WIRTEM.; PIDERIT)

R. Zedoary,	six ounces.
Anise,	three ounces.
Lemon Peel,	two ounces.
Alcohol,	six pounds.

Distil one half. (PIDERIT)

PALAT.; SARD.; and WIRTEM. prescribe six ounces of zedoary, three of anise, one of lemon peel, and eight pounds of white wine; the half to be distilled after eight days' infusion.

ZIBETHA, ZIBETHUM.—CIVET.

SYNONIMES: — Civette, FR.; Zibeth, GERM.; Zibeto, SPAN.; Sivet, DUT.; Zibetto, ITAL.

BRUNS.; HISP.; GALL.; WIRTEM.; BRUGNATELLI; GUIBOUT; PIDERIT; SPIELMANN.

A semifluid, unctuous substance, of a very strong odour, and of a whitish or yel-

lowish colour, becoming very thick and brown by exposure to the air. It is a glandular secretion, and is found in a pouch situated between the anus and genitals of the civet, *Viverra Zibetha*, LINN., one of the rodentia of Asia and Africa.

It is excitant, and is considered antispasmodic.—Dose, from five to ten grains.

ZINCUM.—ZINC.

SYNONIMES: — *Stannum Indicum*.—Zinc, FR.; Zink, Spiauter, GERM.; and SWED.; Pi-yuen, CHIN.; Sungbusrie, DUK.;

Zinco, ITAL.; and PORT.; Cynk, POL.; Spiagter, RUSS.; Tutunagum, TAM. LOND.; EDINB.; DUBL.; AMER.; AMST.;

ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HAMB.; HANN.; HASS.; LIPP.; OLDEN.; LUSIT.; POL.; BORUS.; ROSS.; SAX.; SUEC.; HERBIP.; AINSLIE; COXE; PIDERIT; SPIELMANN.

A solid, bluish-white, lamellated, very ductile metal, more laminable than capable of being drawn into wire, not very hard, fusible above the red heat, volatilizable, burning with a vivid flame, and acted on very slightly by moist air. Its specific gravity is 7,1.

CHLORURET OF ZINC.

Murias seu Butyrum Zinci. (NIEMANN)

R. Pure Hydrochloric Acid, three ounces.

Throw shot-zinc gradually into it till it will dissolve no more, then add

Water, three ounces.

Filter the liquor and evaporate it to dryness. It is astringent and emetic.

ÆTHEREOUS SOLUTION OF HYDROCHLORATE OF ZINC.

Æther Zinci. (NIEMANN; VAN MONS)

R. Dried Hydro-chlorate of Zinc, half an ounce.
Pure Alcohol, one ounce.
Sulphuric Æther, two ounces.

Decant at the end of a few days.

It is anti-spasmodic.—Dose, from four to eight drops, twice a day.

CYANURET OF ZINC.

Hydrocyanas Zinci. (MAGENDIE; NIEMANN)

R. Hydrocyanate of Potass, any quantity.
Sulphate of Zinc, a sufficient quantity.

Having dissolved them, add the latter to the former till no more precipitate is formed. Dry and calcine this at a dull red heat.

The product is a mixture of cyanuret of zinc and of potassium.—It has been extolled in nervous cardialgia.—Dose, from the fourth of a grain to one grain and a half in a potion, to be taken by spoonfuls.

ANTI-SPASMODIC POWDER. (MAGENDIE)

R. Cyanuret of Zinc, half a grain.
Magnesia, four grains.
Cinnamon, three grains.

Mix.—To be taken every four hours.

PROTOXIDE OF ZINC.

Oxydum seu Oxodes Zinci seu zincicum, Zincum oxydatum.

A.

That which is met with in commerce.

1. *Nihil album.*

BATAV.; BRUNS.; FENN.; CAST. BORUS.; SAX.; SUEC.; WIRTEM.; HERBIP.; PALAT.; SPIELMANN.

In very white, light, woolly floculi, insipid, inodorous, and insoluble in water, formed by the combustion of zinc in the open air.

2. *Tutia, Tuthia, Tutia Alexandrina, Botoytis, Nihilum griseum, Oxydum zincicum griseum.*—Tutty, Impure Oxide of Zinc.

LOND.; DUBL.; EDINB.; AMER.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; HANN.; HASS.; OLDEN.; LUSIT.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; SPIELMANN.

In thick, compact, heavy scales, of an ash-grey colour, hard, difficult to be broken, shagreened and rough; it is formed in the chimneys of furnaces in which lead ores containing sulphuret of zinc are roasted.

It is the protoxide mixed with extraneous substances.

3. *Calamina, Lapis Calaminaris, Carbonas Zinci impurus, Carbonas zincicum terrestre, Oxydum Zinci nativum, Zincum terrestre.*—Calamine, Fossile Cadmia, Hydrosilicate of Zinc. (B.)

LOND.; EDINB.; DUBL.; AMER.; AMST.; ANT.; BATAV.; BELG.; BRUNS.; DAN.; HISP.; GALL.; FENN.; FERR.; FULD.; HAMB.; HASS.; LIPP.; OLDEN.; LUSIT.; SAX.; SUEC.; WIRTEM.; HERBIP.; COXE; GUIBOUT; PIDERIT; SPIELMANN.

In irregular, heavy, whitish, greyish, or reddish masses.

The name given to this substance by BEUDANT perfectly expresses its nature. It is a hydrated silicate of zinc containing accidentally oxide of iron, alumine, and carbonate of lime.

PREPARED CALAMINE.

Calamina præparata. (LOND.)

Carbonas Zinci impurus præparatus (EDINB.)

Zinci Carbonas impurum præparatum. (DUBL.)

Calcine calamine, and pulverize it;

then treat it in the manner directed for the preparation of chalk.

B.

Prepared expressly.

1. *In the dry way; Flowers of Zinc.*—*Flores Zinci, Nihil album, Pompholer, Lana philosophica, Oxydum zincicum seu Zinci album seu Zinci per se, Oxodes Zinci album, Calx Zinci, Zincum oxydatum album.*

DUBL.; EDINB.; AMER.; AMST.; ANT.; AUSTR.; BATAV.; BAVAR.; BELG.; BRUNS.; HISP.; GALL.; FENN.; FERR.; FULD.; GENEV.; HANN.; HASS.; LIPP.; LUSIT.; PALAT.; POL.; ROSS.; SAX.; SUEC.; WIRTEM.; HERBIP.; BRUGNATELLI.; COXE.; PIDERIT.; SPIELMANN.; SWEDIAUR.; VAN MONS.

Oxidum Zinci.

Place a large crucible in a furnace filled with burning coals in such a manner as to be a little inclined towards its mouth, and when the bottom of it has become of a moderate red heat, throw into it a piece of zinc of about the weight of a dram. The zinc is soon inflamed and converted into white floculi, which are from time to time to be removed from the surface of the metal, by means of an iron spatula, that the combustion may be more complete; when the inflammation has ceased remove the oxide from the crucible. Then throw in another piece of zinc, and proceed in the same manner. Finally, let the oxide be prepared in the same manner as the impure carbonate of zinc. (EDINB.)

R. Zinc, in small pieces,
any quantity.

Throw these, at intervals, into a sufficiently large crucible, heated to whiteness, and placed with its mouth inclined towards the mouth of the furnace. After each piece of zinc is thrown in, invert over the crucible another crucible, but loosely, so as not to exclude the air. Preserve the white, sublimed, light powder for use. (DUBL.)

R. Zinc, any quantity.

Put it into a large crucible placed among burning coals, and augment the heat till the metal enters into fusion, and inflames; then cover the crucible with another inverted, and afterwards separate the floculi which collect in the latter.

2. *In the moist way; Magistery of Zinc.*—*Magisterium Zinci, Zinci præcipitatum.*

AMST.; DAN.; GALL.; FERR.; HANN.; LIPP.; OLDEN.; BORUS.; SAX.; VAN MONS.

Zinci Oxydum. (LOND.)

R. Sulphate of Zinc, one pound.
Distilled Water, one pint.

Dissolve; then add

Solution of Ammonia, one part, or as much as may be sufficient to precipitate the oxide of zinc entirely. Pour off the liquor; wash the powder repeatedly with distilled water, and dry it by the heat of a sand-bath.

Dissolve sulphate of zinc in a sufficient quantity of water, and pour gradually into the solution another of subcarbonate of soda or of potass, till no more precipitate is formed. Wash and dry this.

This product is subcarbonate of zinc: GALL. speaks of it twice, under the title of oxide of zinc by precipitation and of carbonate of zinc.—The calcination for a quarter of an hour which SAX. requires the precipitate to be subjected to, renders it yellow, but does not bring it to the state of pure oxide.

Oxide of zinc is employed externally as astringent, and internally as anti-spasmodic and emetic. It has been extolled in epilepsy, hysteria, and neuralgiæ.—Dose, from six grains to half a dram, in the form of pills.

ANTI-SPASMODIC POWDER.

Pulvis Zinci oxydati albi. (FULD.; AUGUSTIN)

R. Oxide of Zinc, twelve grains.
Sugar, one dram.

Divide into six papers.—Dose, one, morning and evening, in whooping cough and the convulsive diseases of children. (AUGUSTIN)

R. Oxide of Zinc, one grain.
SYDENHAM'S Laudanum, one dram.
Sugar, one scruple.

Mix.—To be taken every three hours, in the spasms of children. (AUGUSTIN)

- R. Oxide of Zinc, one dram.
 Opium, ten grains.
 Cinchona, }
 Fennel, } of each, one ounce.

Divide into fifty parcels.—Dose, one every two hours, in hooping cough. (AUGUSTIN)

- R. Oxide of Zinc, three grains.
 Squill, of each,
 Red Sulphuret of } one grain and
 Antimony, a half.
 Subcarbonate of Magnesia, one dram.

Divide into six parcels.—Dose, one every two hours, in rheumatismal spasmodic asthma. (AUGUSTIN)

- R. Oxide of Zinc, twelve grains.
 Opium, three grains.
 Aqueous Extract of Cinchona, one dram.

Oleosaccharum of Anise, two drams.

Divide into twelve parcels. (FULD)

STOMACHIC POWDER. (BRERA)

- R. Oxide of Zinc, one grain.
 Magnesia, one scruple.

Mix.—To be taken every two hours, in dyspepsia.

- R. Oxide of Zinc, four grains.
 Calumba, thirty grains.

Divide into four parcels.—Dose, one every three hours.

DRY COLLYRIUM. (CADET DE GASSICOURT ; RATIER)

- R. Tutty, }
 Sugar Candy, } of each, half a dram.

Make a very fine powder. (RATIER)

- R. Tutty, }
 Florentine Iris, } of each, an equal part.
 Sugar Candy, }

Make a very fine powder. (CADET DE GASSICOURT)

BLACK CEPHALIC POWDER.

Pulvis cephalicus niger. (LIPP.)

- R. Oxide of Zinc, one dram.
 Powdered Mistletoe, }
 Charcoal, } of each,
 one ounce.

Mix.—Dose, from fifteen to twenty grains, to a child of one year old.

ANTI-STRUMOUS POWDER. (AUGUSTIN ; SAINTE-MARIE)

- R. Oxide of Zinc, four ounces.
 Cuttle Fish Bone, }
 Egg Shells, } of each,
 Scarlet Cloth, } two ounces,
 one ounce and a half.

Burn them together in a covered vessel, and pulverize the residue. (SAINTE-MARIE)

AUGUSTIN prescribes half an ounce of oxide, two drams of egg shells, and one dram and a half of cloth.

Dose, eighteen grains, twice a day, for an adult.—This remedy is recommended by DEHAEN.

SEDATIVE AND DIAPHORETIC BOLUSES.

Boli diaphoretici sedantes. (BRERA)

- R. Oxide of Zinc, } of each,
 Extract of Aconite, } four grains.
 Red Sulphuret of Antimony, }

Liquorice, in powder, } two grains.
 Honey, } of each,
 a sufficient quantity.

Make four boluses.—Dose, one every two hours, in sciatica, chorea, hystèria, rheumatism, and chronic gout.

SEDATIVE PILLS. (GENEV. ; BRERA ; ELLIS)

- R. Oxide of Zinc, two scruples.
 Conserve of Dog-rose,
 a sufficient quantity.

Make twenty-four pills. (GENEV. ; ELLIS)

- R. Oxide of Zinc, eight grains.
 Extract of Aconite, six grains.

Make six pills.—Dose, one every four hours. (BRERA)

These pills have been recommended in epilepsy, chorea, and chronic exanthemata.

ASTRINGENT COLLYRIUM. (SWEDIAUR)

- R. Oxide of Zinc, one dram.
 Mucilage of Quince Seeds, prepared with rose water, ten ounces.

Triturate together.

ASTRINGENT LOTION.

Lotio e Zinco et Cupro composita. (AUGUSTIN ; ELLIS ; SWEDIAUR ; SPIELMANN)

- R. Oxide of Zinc, twelve grains.
 Sulphate of Copper, four grains.
 Honey of Roses, one dram.
 Lime Water, two ounces.

Mix.—It is recommended as an application to syphilitic ulcers. (ELLIS ; SPIELMANN ; SWEDIAUR)

- R. Oxide of Zinc, one ounce.
 Elder-flower Water,
 one ounce and a half.

Mix.—THUESINK extolled this preparation in pustular erysipelas. (AUGUSTIN)

RESOLVENT COLLYRIUM. (PIERQUIN;
SPIELMANN)

- R. Tutty, half a dram.
 Aloes, one dram.
 Rose Water, } of each,
 Fennel Water, } one ounce.
 Eye-bright Water, }

Mix. (PIERQUIN)

This formula was probably framed from the following which SPIELMANN gives under the name of *Aqua ophthalmica*, and the product of which certainly does not contain an atom of zinc:

- R. Prepared Tutty, } of each, one ounce.
 Cloves, }
 Sugar Candy, }
 Camphor, } of each,
 Aloes, } one dram and a half.
 Spanish Wine, } forty-eight ounces.
 Water of Pale Roses, } six ounces.
 ——— Celandine, }
 ——— Fennel, } of each,
 ——— Eye-bright, } two ounces.
 ——— Rue, }

Macerate for fifteen days in the sun, and decant.

There is in BRUNS., under the title of *Aqua ophthalmica usualis*, the following formula, which somewhat resembles PIERQUIN'S:

- R. Rose Water, } of each, five ounces.
 Fennel Water, }
 Prepared Tutty, } six drams.
 Acetate of Lead, } one dram and a half.
 Camphor, } one dram.
 Sulphate of Zinc, } two scruples.

Mix, and preserve.

OINTMENT OF OXIDE OF ZINC.

Ceratum Calaminæ seu calaminare seu epuloticum seu e Zinco, Collyrium Nihili seu zincicum seu e Zinco, Unguentum calaminare seu Lapidis calaminaris seu e Lapide calaminari seu Carbonatis Zinci impuri seu Oxydi Zinci seu Oxydi Zinci impuri seu nativi seu de Nihilo seu Tutie seu Zinci seu griseum seu Nihili ad oculos seu ophthalmicum album seu diapompholigos. (LOND.; EDINB.; DUBL.; AMER.; AMST.; BATAV.; GALL.; FENN.; FULD.; GENEV.; HAMB.; HANN.; LIPP.; OLDEN.; LUSIT.; PALAT.; POL.; BORUS.; SAX.; AUGUSTIN.; BRUGNATELLI.; CADET DE GASSICOURT.; COXE.; ELLIS.; HUFELAND.; PIERQUIN.; SPIELMANN.; SWEDIAUR.)

Ceratum Calaminæ. (LOND.)

- R. Prepared Calamine, } of each,
 Yellow Wax, } half
 Olive Oil, } a pound.
 one pint.

Mix the oil with melted wax; then remove the mixture from the fire, and as soon as it begins to thicken, add the calamine, and stir constantly till cold.

Ceratum Carbonatis Zinci impuri.
(EDINB.)

- R. Simple Cerate, five parts.
 Prepared Impure Carbonate
 of Zinc, one part.

Mix.

Unguentum calaminare. (DUBL.)

- R. Ointment of Yellow Wax, five pounds.
 Prepared Calamine, one pound.
 Make an ointment.

AMER.; HANN.; LUSIT.; COXE.; NIEMANN.; and SWEDIAUR give the same formula as LOND.—FULD. prescribes twelve parts of oil, six of wax, and four and a half of calamine.—AMST.—two of oil, one of wax, and one and a half of calamine.—CADET DE GASSICOURT—six of oil, one of white wax, and one of calamine.

Unguentum Oxidi Zinci impuri.
(EDINB.; COXE)

- R. Simple Liniment, five parts.
 Prepared Impure Oxide of
 Zinc, one part.
 Mix.

Unguentum Tutie. (DUBL.)

- R. Ointment of White Wax, ten ounces.
 Prepared Tutty, two ounces.
 Make an ointment.

BATAV.; HANN.; OLDEN.; POL.; BORUS.; and SAX. prescribe eight parts of simple cerate, and one part of impure oxide.—GENEV.—twelve of cerate, and one of oxide.—AUGUSTIN—two drams of oxide, and one ounce of rose ointment.—LUSIT.—thirty-two of lard, one of wax, and eight of oxide.

Unguentum Zinci. (LOND.)

- R. Oxide of Zinc, one ounce.
 Prepared Lard, six ounces.
 Mix.

Unguentum Oxidi Zinci.

- R. Simple Liniment, six parts.
Oxide of Zinc, one part.

Mix. (EDINB.)

- R. Ointment of White Wax, one pound.

Oxide of Zinc,
one ounce and a half.

Make an ointment. (DUBL.)

FENN.; and COXE give the same formula as LOND.—PIERQUIN prescribes two parts of lard and one part of oxide.—PALAT.—two of goose-fat and one of oxide.—HAMB.—three of lard and one of oxide.—FULD.—three of butter and one of oxide.—BRUGNATELLI; and SWEDIAUR—four of lard and one of oxide.—PIERQUIN—four of butter and one of oxide.—AMER.; BRUGNATELLI; and COXE—five of lard and one of butter.—ANT.; and BELG.—eight of lard and one of oxide.—LIPP.—three of lard, one of oxide, and six of fresh butter.—HUFELAND—eight of lard and one of oxide.—ELLIS—six of butter and one of oxide.

- R. Oxide of Zinc, one part.
Butter, washed in }
Rose Water, } of each,
Rose Ointment, } two parts.

Mix intimately. (GALL.)

- R. Lard, twenty-four parts.
Mother of Pearl, } of each, one part.
Red Coral, }
Tutty, } of each, two parts.
Nihil Album, }

Mix. (SPIELMANN)

CAMPHORATED OINTMENT OF OXIDE OF ZINC.

Unguentum Nihili seu Nihili ad oculos seu ophthalmicum seu Zinci. (BRUNS.; DAN.; PALAT.; WIRTEM.; SWEDIAUR; VAN MONS)

- R. Oxide of Zinc, one ounce.
Camphor, one scruple.
Lard, six ounces.

Mix. (DAN.)

- R. Lard, six ounces.
Oil of Almonds, three ounces.
White Wax, two ounces.

Melt together, and add

Oxide of Zinc, three ounces.
Camphor, triturated with oil,
three drams.

Mix. (VAN MONS)

VAN MONS also prescribes one ounce of oil of almonds, half an ounce of lard, half an ounce of wax, one ounce and a half of calamine, two drams of tutty, and half a dram of camphor.

- R. Prepared Tutty, } of each, one ounce.
Nihil Album, }
Red Coral, } of each,
Mother of Pearl, } half an ounce.
Camphor, one dram.
Lard, one pound.

Mix by trituration. (BRUNS.; PALAT.; WIRTEM.)

ANTI-HÆMORRHOIDAL OINTMENT.

Unguentum exsiccans; Cerat de Hufeland, Pommade contre l'intertrigo. (AUGUSTIN; CADET DE GASSICOURT; NIEMANN; PIERQUIN)

- R. Oxide of Zinc, } of each,
Powdered Lycopodium, } one dram.
Cerate, half an ounce.

Triturate together. (AUGUSTIN; PIERQUIN)

NIEMANN prescribes one dram of oxide, one dram of lycopodium, and four ounces of rose ointment.—CADET DE GASSICOURT—thirty grains of oxide, thirty of lycopodium, and one ounce of spermaceti ointment.

- R. Oxide of Zinc, } of each,
Powdered Lycopodium, } fifteen grains.
Lard, half an ounce.
Rose Water, } of each,
Plantain Water, } asufficient quantity.

Triturate, and mix exactly. (PIERQUIN)

DESICCATIVE CERATE.

Ceratum seu Unguentum exsiccans seu ophthalmicum seu ad Ambusta. (HANN.; BORIES; CADET DE GASSICOURT; NIEMANN)

- R. Olive Oil, eight ounces
Yellow Wax, one ounce and a half.

Melt, and add

Armenian Bole, one ounce.
Oxide of Zinc, half an ounce.

Mix. (HANN.)

- R. Olive Oil, four ounces.
White Wax, } of each, one ounce.
Calamine, }
Acetate of Lead, two drams.

Make a cerate. (BORIES)

- R. Tutty, } of each,
Acetate of Lead, } two scruples.
Calamine, }
Camphor, four grains.
Rose Ointment, a sufficient quantity.

Triturate together. (NIEMANN)

RED OPHTHALMIC OINTMENT.

Unguentum ophthalmicum Sloani. (SPIELMANN)

- R. Tutty, one ounce.
Hæmatitis, two scruples.

Aloes, twelve grains.
Prepared Pearls, four grains.
Viper's Fat, two ounces.

Mix.

WHITE OPHTHALMIC OINTMENT. (VAN MONS)

R. Oxide of Zinc, three drams.
Protochloruret of Mercury, half a dram.
Camphor, one scruple.
White Wax, two drams.
Cocoa-nut Oil, } of each, one ounce.
Oil of Almonds, }

Mix by trituration,

GREEN OPHTHALMIC OINTMENT.

Unguentum oculare Keiseri. (AMST.)

R. Tutty, half an ounce.
Camphor, one dram.
Verdigris, twelve grains.
Lard, two ounces.

Mix.

CALAMINE PLASTER.

Emplastrum griseum seu de Lapide calaminari seu consolidans, Pamphilus Chirurgorum. (AMST.; BRUNS.; DAN.; MIL. COPEN.; FULD.; HANN.; LIPP.; PALAT.; SAX.; SUEC.; WIRTEM.; HERBIP.; NIEMANN.; SPIELMANN.; VAN MONS.)

R. Calamine, one pound.
Semivitreous Oxide of Lead, half a pound.
Olive Oil, one pound and a half.

Boil, constantly stirring. (FULD.; NIEMANN.)

R. Simple Plaster, one pound and a half.
Yellow Wax, } of each, nine ounces.
Olive Oil, }

Melt together, and add

Powdered Calamine, one pound and a half.

Mix. (AMST.)

R. Simple Plaster, } of each,
Powdered Calamine, } one pound and a half.
Yellow Wax, } six ounces.
Goat Suet, } of each,
Turpentine, } one ounce and a half.

Liquify, and mix perfectly. (DAN.)

SUEC.; and VAN MONS prescribe two pounds each, of simple plaster and of calamine, five ounces of wax, and one ounce and a half each, of suet and of turpentine.

R. Simple Plaster, six ounces.

Melt, and add

Powdered Calamine, two ounces.
Mastic, } of each,
Olibanum, } one dram.

Mix. (SAX.)

R. Simple Plaster, one pound.
Olive Oil, five ounces.

Melt together, and add

Powdered Calamine, four ounces.
Myrrh, } of each,
Olibanum, } two drams.

Mix. (HANN.)

LIPP. prescribes ten ounces of simple plaster, eight of olive oil, six of calamine, one ounce of mastic, and half an ounce of olibanum.

R. Oil of Rape, eighteen ounces.
Calamine, twelve ounces.
Semivitreous Oxide of Lead, six ounces.
Yellow Wax, one ounce.

Melt, and mix perfectly. (MIL. COPEN.)

R. Calamine, }
Semivitreous Oxide of Lead, } of each,
Subcarbonate of Lead, } half a pound.
Olibanum, half an ounce.
Mastic, one ounce.
Olive Oil, a sufficient quantity.

Make a plaster. (HERBIP.)

R. Wax, three ounces.
Stag's Suet, four ounces.
Turpentine, one ounce and a half.

Melt, and add

Semivitreous Oxide of Lead, } of each,
Powdered Calamine, } one ounce
Tutty, } and a half.
Mastic, } of each,
Olibanum, } half an ounce.
Camphor, triturated with a little oil, one dram.

Mix. (SPIELMANN)

BRUNS.; and WIRTEM. prescribe three ounces of wax, four of suet, one ounce and a half of turpentine, one of calamine, two ounces of semivitreous oxide, half an ounce each, of tutty, of mastic, and of olibanum, two drams of myrrh, and one dram and a half of camphor.—PALAT.—eight ounces each, of wax and of suet, six of turpentine, four of calamine, three each, of semivitreous oxide and of subcarbonate of lead, and half an ounce each, of tutty, of mastic, of olibanum, of myrrh, and of camphor.

DIAPOMPHOLIGOS PLASTER.

Emplastrum diapompholigos seu zincoplumbicum. (AMST.; BATAV.; BRUNS.; FENN.; HANN.; SAX.; SUEC.; WIRTEM.; SPIELMANN.; VAN MONS)

R. Calamine Plaster, sixteen parts.
Calcined Lead, one part.

Melt, and mix well. (SAX.)

R. Oil of Black Nightshade, thirty-four parts.
Subcarbonate of Lead, ten parts.

Boil, and add

Wax, thirty-four parts.
Calcined Lead, six parts.
Oxide of Zinc, } of each, four parts.
Olibanum, }

Mix. (SPIELMANN)

BRUNS.; and WIRTEM. prescribe two pounds and a half of oil of black nightshade, one ounce and a half of subcarbonate of lead, thirty ounces of white wax, one ounce and a half of calcined lead, and four ounces each, of oxide of zinc and of olibanum.—VAN MONS—six parts of olive oil, ten of subcarbonate of lead, eight of wax, one of resin, and two each, of calcined lead, of oxide of zinc, and of tutty.

R. Olive Oil, half a pound.
Yellow Wax, one pound.

Melt together, and add

Calcined Lead, half a pound.
Subcarbonate of Lead, four ounces.
Oxide of Zinc, three ounces.
Olibanum, one ounce and a half.

Mix intimately. (FENN.; SUEC.)

BATAV. prescribes seven parts of oil, twelve of wax, six of calcined lead, four of subcarbonate of lead, three of oxide of zinc, and one and a half of olibanum.—AMST.—thirty of oil, thirty-six of wax, twelve of calcined lead, eight of subcarbonate, six of oxide of zinc, and two and a half of olibanum.—HANN.—fifteen of oil, sixteen of wax, three each, of calcined lead and of subcarbonate, and two each, of oxide of zinc and of olibanum.

MYNSICHT'S SATURNINE PLASTER. (PALAT.; WIRTEM.)

R. Goat's Suet, } of each,
White Wax, } one ounce and a half.
Turpentine, six drams.

Melt, and add

Powdered Calcined Lead, two ounces.
Calamine, } of each,
Subcarbonate } half
of Lead, } an ounce.
Nihil Album, } of each,
Tutty, } two drams.
Myrrh, } of each, one dram.
Mastic, }

VOL. II.

Powdered Olibanum, one dram.
Aloes, } of each,
Camphor, } half a dram.
Nitro, }

Mix. (PALAT.)

WIRTEM. prescribes five ounces each, of suet and of yellow wax, two each, of turpentine and of calcined lead, half an ounce each, of calamine and of subcarbonate of lead, two drams each, of oxide of zinc and of tutty, one dram each, of myrrh, of mastic, of olibanum, of aloes, and of nitre, and half a dram of camphor.

DIAPOMPHOLIGOS OINTMENT.

Unguentum diapompholigos seu Pompholigis seu exsiccans seu ad Ambusta. (BRUNS.; OLDEN.; PALAT.; SARD.; WIRTEM.; SPIELMANN.; VAN MONS)

R. Unguentum Nutritum, thirty-two ounces.
Powdered Calamine, four ounces.

Mix. (OLDEN.)

R. Camphor, one dram.
Subcarbonate of Lead, half an ounce.
Yellow Wax, } of each, two ounces.
Calamine, }
Olive Oil, five ounces.

Triturate the subcarbonate and calamine with a little oil, add the remainder of the oil melted with the wax, then the camphor, triturated with oil, and stir constantly till cold. (VAN MONS)

R. Oil of Black Nightshade, sixteen ounces.
Wax, five ounces.

Melt slowly, and add

Powdered Subcarbonate of Lead, two ounces.
Oxide of Zinc, } of each,
Calcined Lead, } one ounce.
Olibanum, }

Mix. (SPIELMANN)

R. Rose Oil, two pounds.
Juice of Black Nightshade Berries, four ounces.

Boil to the consumption of the humidity, and add

White Wax, six ounces.
Powdered Subcarbonate of Lead, two ounces.
Calcined Lead, } of each,
Tutty, } one ounce.
Olibanum, }

Mix. (SARD.)

BRUNS.; and WIRTEM. prescribe six ounces of rose oil, four each, of black nightshade berries and of wax, and one ounce each, of calcined lead, of oxide of zinc, and of olibanum.—PALAT. gives the same pre-

portions, but reduces the wax to three ounces, and replaces the calcined lead by two ounces of subcarbonate.

COMPOUND OINTMENT OF CALAMINE.

Unguentum Calaminæ compositum seu exsiccans rubrum Acrelii. (FENN.; SUEC.)

R. Calamine,	} of each, half a pound.
Semivitreous Oxide of Lead,	
Subcarbonate of Lead,	
Red Oxide of Lead,	} of each, three ounces.
Acetate of Lead,	
Goat's Suet,	
Linseed Oil,	one pound.
	eight ounces.

Melt the suet, add the oil, then the powders, and when the mass is a little cooled,

Powdered Camphor, three drams.

Mix.

AMMONIURET OF ZINC. (VAN MONS)

R. Solution of Acetate of Zinc, any quantity.

——— Ammonia, a sufficient quantity.

Add the latter to the former till it ceases to effect a precipitate. Wash and dry this in the smallest possible quantity of solution of ammonia; then filter, evaporate, and crystallize.

SULPHATE OF ZINC.

Vitriolum album seu Goslariense seu Zinci, Sulphas Zinci seu zincicum, Zincum sulphuricum, Gilla Vitrioli seu Theophrasti, Sal vomitorium Vitrioli, Chalcantum album.—White Vitriol, White Copperas.

1. That which is found in commerce.

AMER.; AMST.; BATAV.; BELG.; BRUNS.; DAN.; MIL. COPEN.; HISP.; GALL.; FENN.; FERR.; FORM. HÔP. FR.; FULD.; GENEV.; HAMB.; HASS.; LIPP.; LUSIT.; CAST. BORUS.; ROSS.; SUEC.; WIRTEM.; HERBIP.; WURCEB.; COXE; PARIS; PIDERIT; SAUNDERS; SPIELMANN.

2. Purification of that of commerce.

R. White Vitriol, any quantity

Having calcined it, dissolve it in water, and crystallize several times successively. (GALL.)

WIRTEM. omits the calcination.—FULD.; LIPP.; and HERBIP. direct one pound of the salt to be dissolved in two pounds of boiling water, the solution to be left in digestion for six hours on six drams of shot-zinc, then to be evaporated and crystallized.

3. Prepared expressly.

LOND.; EDINB.; DUBL.; ANT.; AUSTR.;

BATAV.; BAVAR.; BELG.; HISP.; GALL.; FERR.; FENN.; HANN.; OLDEN.; POL.; BORUS.; SAX.; SUEC.; BRUGNATELLI; COXE; SWEDIAUR; VAN MONS.

Zinci Sulphas.

R. Zinc, broken into small pieces, four ounces.
Sulphuric Acid, by weight, six ounces.
Water, four pints.

Mix in a glass vessel, and when the effervescence has ceased, filter the solution through paper; then boil till a pellicle appears on the surface, and set it aside to crystallize. (LOND.)

R. Zinc, in small pieces, three parts.
Sulphuric Acid, five parts.
Water, twenty parts.

Mix, and when the effervescence has ceased, digest for a short time in a sand-bath. Filter the decanted solution through paper; then evaporate, and set aside to crystallize. (EDINB.)

R. Zinc, in small pieces, thirteen parts.
Sulphuric Acid, twenty parts.
Water, one hundred and twenty parts.

Pour the acid mixed with the water upon the zinc in a glass vessel. Digest for a short time after the effervescence has ceased; then evaporate the strained solution to the required degree, and set it apart to crystallize. (DUBL.)

R. Sulphuric Acid (20 degrees), any quantity.

Put it into a glass capsule placed in a sand-bath, and when it is moderately heated, add shot-zinc gradually to it, dilute the liquor with a little water, saturate, evaporate, and crystallize.

This salt is employed externally as astringent, and internally as tonic and astringent, in the dose of from one to five grains twice or thrice a day, and as emetic, in that of from ten to twenty grains in distilled water.

EXPECTORANT PILLS. (ELLIS; PARIS)

R. Sulphate of Zinc, half a scruple.
Conserve of Roses, a sufficient quantity.

Make twenty pills.—Dose, one, three or four times a day, in whooping cough and phthisis pulmonalis. (ELLIS)

R. Sulphate of Zinc, ten grains.
Myrrh, one dram and a half.
Conserve of Roses, a sufficient quantity.

Make twenty pills. They are prescribed in hooping cough. (ELLIS; PARIS)

SIMPLE AQUEOUS SOLUTION OF SULPHATE OF ZINC. (AMER.; COXE; NIEMANN)

1. For external use.

R. Sulphate of Zinc, two grains.
Chamomile Water, three ounces.

Dissolve.—Dose, one spoonful, two or three times a day.

WYLIE administers every two or three hours, in malignant fevers, a spoonful of a solution of nine grains of sulphate of zinc in six ounces of elder-flower water.

MOSELEY also recommends in chronic dysentery without fever, a solution of three drams of sulphate of zinc and two drams of alum in half an ounce of spirit of lavender and one pint of water, in the dose of from four to six drams in water, which he sometimes repeats at the end of three hours.

We also find in AMER.; and COXE a similar preparation, of which the following is the formula :

R. Quassia Wood, one dram.
Sulphate of Zinc, eight grains.
Water, half a pint.

Macerate for twelve hours, and strain.

2. For internal use.

A.

In the form of Gargle. (AUGUSTIN; NIEMANN)

R. Sulphate of Zinc, one dram.
Distilled Water, six ounces.
Honey, one ounce.

Dissolve. (NIEMANN)

R. Sulphate of Zinc, fifteen grains.
Infusion of Sage, six ounces.

Dissolve. (AUGUSTIN)

B.

In the form of Collyrium. — *Aqua ophthalmica seu ophthalmica adstringens seu vitriolica seu e Vitriolo Zinci, Aqua Vitrioli albi, Solutio Sulphatis Zinci, Collyrium vitriolicum seu e Zinco sulphurico seu e Vitriolo albo seu Zinci.* (EDINB.; AMER.; MIL. COPEN.; GALL.; FORM. HÔP. FR.; HAMB.; LUSIT.; CAST. BORUS.; SARD.; SUEC.; WURCEB.; AUGUSTIN; BORIES; CADET DE GASSI-

COURT; COXE; DISP. CHA. PARIS; RATIER; SAUNDERS; SPIELMANN; SWEDIAUR.

R. Sulphate of Zinc, one part.
Water, two hundred and forty parts.

Dissolve. (MIL. COPEN.; CAST. BORUS.; WURCEB.; NIEMANN)

FORM. HÔP. FR. prescribes one part of sulphate, and two hundred parts of water. — LUSIT.; SUEC.; and SPIELMANN—one of sulphate, and ninety-six of water. — MIL. COPEN.—one of sulphate, and sixty-four of water. — AMER.; and COXE—twelve grains of sulphate, and half a pint of water. — DISP. CHA. PARIS—from four to six grains of sulphate, and six ounces of water. — HAMB.—one part of sulphate, and three hundred and twenty parts of water. — RATIER—one of sulphate, and one hundred and twenty of water. — RATIER; and SWEDIAUR—one of sulphate, and ninety-six of water. — SAUNDERS; and SWEDIAUR—one of sulphate, and forty-eight of water.

Solutio Sulphatis Zinci. (EDINB.; COXE)

R. Sulphate of Zinc, sixteen grains.
Water, eight ounces.
Dilute Sulphuric Acid, sixteen drops.

Dissolve the sulphate in the water, add the acid, and filter.

R. Sulphate of Zinc, one grain.
Rose Water, one ounce.
Mucilage of Quince Seeds, one dram.

Mix.

R. Sulphate of Zinc, one scruple.
Infusion of Elder, one pound.

Dissolve. (RATIER)

RATIER prescribes elsewhere one grain of sulphate, and one ounce of plantain water. — AMER.; and COXE—two drams of sulphate, two ounces of spirit of lavender, and six ounces of water. — SARD. directs the yolk of a boiled egg to be removed and replaced by sulphate of zinc, and six ounces each, of rose water and of plantain water to be added to the liquor which drops from it. — AUGUSTIN—three grains of sulphate to be dissolved in one ounce and a half of rose water beaten with the white of an egg.

R. Sulphate of Zinc, five grains.
Plantain Water, four ounces.
Mucilage of Quince Seeds, half an ounce.

Dissolve. (CADET DE GASSICOURT)

R. Sulphate of Zinc, one dram.
Rose Water, two pounds.
Alcohol (22 degrees), one ounce.

Dissolve. (GALL.)

- R. Sulphate of Zinc, six grains.
 Camphor, four grains.
 Eye-bright Water, } of each, one ounce.
 Rose Water, }
 Plantain Water, }

Dissolve. (BORIES)

- R. Sulphate of Zinc, six grains.
 Distilled Water, six ounces.
 Mucilage of Quince, one ounce.
 Spirit of Camphor, twelve drops.

Mix. (SWEDIAUR)

C.

In the form of Lotion or Fomentation.
 (PH. SYPH.; AUGUSTIN; NIEMANN;
 SWEDIAUR)

- R. Sulphate of Zinc, two grains.
 Distilled Water, four ounces.

Dissolve. (PH. SYPH.)

NIEMANN prescribes one dram of sulphate and one ounce of water.—AUGUSTIN—three drams of sulphate, and one ounce of water.

- R. Sulphate of Zinc, } of each,
 Spirit of Camphor, } half an ounce.
 Boiling Water, two pounds.

Dissolve. (PH. SYPH.; SWEDIAUR)

D.

For Injection. (MIL. COPEN.; HISP.; PH. SYPH.; NIEMANN; SWEDIAUR)

- R. Sulphate of Zinc, one grain.
 Water, one ounce.

Dissolve. (NIEMANN)

NIEMANN also prescribes thirty-five grains of sulphate, and four ounces of water; or, from one scruple to half a dram of sulphate, and one ounce of water.

- R. Sulphate of Zinc, one dram.
 Camphorated Water, two ounces.
 Pure Water, two pounds.

Dissolve. (PH. SYPH.; SWEDIAUR)

ASTRINGENT INJECTION.

Injectio adstringens. (BRERA)

- R. Plantain Water, six ounces.
 SYDENHAM'S Laudanum, thirty drops.
 Sulphate of Zinc, one scruple.
 Mucilage of Gum Arabic, one ounce.

Mix.—For blenorrhœa. (BRERA)

- R. Sulphate of Zinc, one scruple.
 Armenian Bole, two scruples.
 Gum Arabic, two drams.
 Water, eight ounces.

Mix.—For blenorrhœa. (ELLIS)

- R. Sulphate of Zinc, ten grains.
 Gum Arabic, two drams.
 Tincture of Opium, one dram.
 Water, eight ounces.

Mix.—For gonorrhœa. (ELLIS)

ASTRINGENT GARGLE. (ELLIS)

- R. Sulphate of Zinc, one dram.
 Honey, half an ounce.
 Tincture of Myrrh, } of each,
 Proof Spirit, } one ounce.
 Rose Water, four ounces.

Mix.

ASTRINGENT FOMENTATION.

Fomentum tonico-adstringens. (BRERA)

- R. Decoction of Pomegranate Bark, four pounds.
 ————— Cinchona, two pounds.
 Sulphate of Zinc, one dram and a half.
 Tincture of Opium, one ounce.

Mix. — For external piles and vaginal blenorrhœa.

RESOLVENT COLLYRIUM. (PALAT.; CADET DE GASSICOURT; SAINTE-MARIE)

- R. Powdered Florentine Iris, three drams.
 Sulphate of Zinc, one dram and a half.
 Rose Water, } of each,
 Plantain Water, } one pound and a half.

Digest for two days, and strain. (PALAT.)

- R. Florentine Iris, } of each,
 Sulphate of Zinc, } half a dram.
 Sugar Candy, two scruples.
 Rose Water, one pound.

Digest for three days, and strain. (CADET DE GASSICOURT; SAINTE-MARIE)

- R. Fresh Yolks of Eggs, two,
 Rose Water, one pound.

Triturate together, and add

Sulphate of Zinc,
 from twelve to twenty-four grains.

Dissolve. (SAINTE-MARIE)

ASTRINGENT SEDATIVE INJECTION. (CADET DE GASSICOURT; PH. SYPH.)

- R. Sulphate of Zinc, six grains.
 Grey Oxide of Lead, twenty grains.
 Rose Water, four ounces.

Mix. (CADET DE GASSICOURT)

PH. SYPH. prescribes eighteen grains of sulphate, sixty of compound powder of sub-carbonate of lead, and twelve ounces of lime-flower or of rose water.

ASTRINGENT COLLYRIUM.

Collyrium adstringens. (AUGUSTIN; BRERA; CADET DE GASSICOURT; PIERQUIN)

- R. Sulphate of Zinc, twenty grains.
 Tutty, half a dram.

Rose Water, } of each,
 Plantain Water, } three ounces.
 Mix. (PIERQUIN)

R. Rose Water, three ounces.
 Sulphate of Zinc, three grains.
 SYDENHAM'S Laudanum, one scruple.

Mix. (BRERA)

R. Sulphate of Zinc, } of each, two ounces.
 Armenian Bole, }
 Camphor, half an ounce.

Mix.—One dram of this powder is mixed with four ounces of rose water. (AUGUSTIN; COXE)

OPHTHALMIC LIQUOR.

Aqua ophthalmic Liquor. (SPIELMANN)

R. White Sugar Candy, one ounce.
 Root of Florentine Iris, three drams.
 Prepared Tutty, two drams.
 Sarcocolla, } of each, one dram.
 Sulphate of Zinc, }
 Aloes, }
 Cloves, one scruple.
 Eye-bright Water, } of each,
 Fennel Water, } eight ounces.
 Rose Water, }
 Spanish Wine, twenty-four ounces.

Infuse in the sun for fifteen days, and decant.

This liquor very much resembles another of the same name, also indicated by SPIELMANN, given above.

STYPTIC LIQUOR.

Aqua seu Liquor seu Injectio adstringens seu Sulphatis Zinci aluminosa, Liquor Aluminis compositus, Solutio Zinci aluminosa, Injectio ad Leucorrhæam, Aqua aluminosa Bateana, Solutio vitriolica. (LOND.; MIL. COPEN.; HISP.; FERR.; LUSIT.; AUGUSTIN; BRERA; ELLIS; CADET DE GASSICOURT; PIERQUIN; SAUNDERS; VAN MONS)

Liquor Aluminis compositus. (LOND.)

R. Alum, } of each,
 Sulphate of Zinc, } half
 Boiling Water, two pints.

Dissolve, and filter.

MIL. COPEN.; COXE; and CADET DE GASSICOURT give the same formula.—AUGUSTIN; CADET DE GASSICOURT; and VAN MONS prescribe half an ounce each, of the two salts, and one pound of water.—FERR.; and LUSIT.—half an ounce each, of the salts, and two pounds and a half of boiling water.—SAUNDERS—one dram each of the two salts, and one pound of

water.—ELLIS—one dram and a half of sulphate, half a dram of alum, and half a pint of boiling water.

R. Sulphate of Zinc, } of each,
 Alum, } half an ounce.
 Rose Water, one pound.

Dissolve, and filter. (HISP.)

R. Sulphate of Zinc, } of each, one dram.
 Alum, }
 Nettle Juice, four ounces.

Dissolve. (PIERQUIN)

R. Sulphate of Zinc, half a dram.
 Sub-borate of Soda, one dram.
 Alum, four grains.
 Rose Water, six ounces.

Dissolve. (PIERQUIN)

R. Decoction of Oak Bark, ten ounces.
 Dog-rose, six ounces.
 Sulphate of Zinc, one dram.
 Alum, one dram and a half.

Add to the solution,

Alcohol, two drams.
 Aqueous Tincture of Opium, half a dram.

Mix. (BRERA)

It is astringent, and is employed externally in ecchymoses, slight hæmorrhages, and chronic mucous discharges.—Dose, a spoonful, to an adult, fasting, every morning.

TONIC COLLYRIUM. (BORIES)

R. Fresh Rue, } of each,
 Flowers of Leopard's-bane, } half
 Juniper, a handful.
 Ipecacuanha, two drams.
 White Wine, thirty grains.
 twelve ounces.

Infuse for twenty-four hours in warm ashes, and add to the strained liquor,

Sulphate of Zinc, twenty grains.
 Muriate of Ammonia, thirty grains.

Dissolve.

YVEL'S OPHTHALMIC LIQUOR. (CADET DE GASSICOURT)

R. Sulphate of Zinc, three drams.
 Copper, one dram.
 Camphor, two scruples.
 Saffron, twenty grains.
 Water, one pound and a half.

Digest for twenty-four hours, and filter. (CADET DE GASSICOURT)

R. Water, three pounds.
 Sulphate of Zinc, two ounces.
 Copper, four drams.
 Camphor, one dram.

Dissolve. (SAINTE-MARIE)

- R. Leaves of Rue, six parts.
Hot Water, fourteen parts.

Infuse for half an hour. Add to the strained liquor,

- Sulphate of Zinc, four parts.
Copper, one part.
Spirit of Camphor, one fourth part.

Dissolve. (VAN MONS)

VAN MONS gives this formula under the title of *St. Jerneron's Collyrium*.—SUEC.; and NIEMANN, under those of *Collyrium Stierncronense* seu *Decoctum Ruta zincatum* seu *Aqua ophthalmica Odhelii*, give the following proportions: two parts of rue, thirty-six of water, reduced to one half, and to twelve ounces of the strained decoction, one ounce of sulphate of zinc, three drams of sulphate of copper, and half a dram of spirit of camphor.

VULNERARY STONE.

Lapis vulnerarius externus. (FULD.)

- R. Alum, } of each,
Sulphate of Zinc, } six ounces.
Acetate of Copper, } of each,
Muriate of Ammonia, } one dram.

Triturate and fuse together; keep it over the fire till perfectly dry, adding towards the end,

Saffron, half a dram.

This preparation has been recommended to be applied to ulcers, as detergent and desiccative.

OPHTHALMIC OINTMENT.

Unguentum ophthalmicum seu ad Oculi Defusiones catarrhosas ex relaxatione. (BRUNS.; FULD.; PIERQUIN)

- R. Sulphate of Zinc, one part.
Fresh Butter, seven parts.

Triturate together. (FULD.)

BRUNS. prescribes one part of sulphate, and six parts of butter.—PIERQUIN—one of sulphate, and four of butter.

VIRGINAL OINTMENT.

Pommade de la comtesse. (CADET DE GAS-SICOURT)

- R. Sulphate of Zinc, four drams.
Leaves of Myrtle, } of each,
Sumach, } three drams.
Nut Galls, } of each,
Cypress Nuts, } two drams.
Pomegranate Bark, }
Rose Ointment, a sufficient quantity.

Mix by trituration.

POWDER AGAINST SPECKS OF THE CORNEA. (PIERQUIN)

- R. Prepared Tutty, }
Sugar Candy, } of each,
Sulphate of Zinc, } one dram.
Copper, }
Acetate of Copper, }
Proto-chloruret of Mer- }
cury, }

Make a powder.

ASTRINGENT POWDER.

Pulvis Cupressi restrictivus. (HISP.)

- R. Armenian Bole, }
Root of Bistort, } of each,
Petals of Red Roses, } one ounce.
Myrtle Berries, }
Cypress Nuts, }
Sulphate of Zinc, }
Tutty, }

Make a fine powder.

ACETATE OF ZINC.

Acetas seu Acetis Zinci, Zincum acetatum.

1. In the solid form.

VAN MONS.

- R. Oxide of Zinc, any quantity.
Distilled Vinegar, a sufficient quantity,
to dissolve the oxide with heat. Filter, concentrate, and crystallize.

2. In the liquid form.

EDINB.; DUBL.; AMER.; DAN.; PH. SYPH.; COXE; ELLIS; NIEMANN; PLANCHE; SWEDIAUR; VAN MONS.

Solutio Acetatis Zinci. (EDINB.)

- R. Sulphate of Zinc, one dram.
Acetate of Lead, four scruples.
Distilled Water, twenty ounces.

Dissolve the salts separately, each in ten ounces of the water; then mix the solutions, and filter after the precipitate has subsided.

SWEDIAUR prescribes one dram each, of the two salts and three ounces of boiling water.—PLANCHE—twenty grains, each, of the salts, and six ounces of water.—AMER.; and NIEMANN—twelve grains of sulphate of zinc, sixteen of acetate of lead, and half a pint of water.—COXE; and NIEMANN give the same formula as EDINB.

- R. Oxide of Zinc, any quantity.
Acetic Acid, a sufficient quantity,
to dissolve the oxide. Take twenty drops of the solution, and add to it four ounces of water. (PH. SYPH.)

These solutions are employed in the form of collyrium, or in that of injection in blenorrhœa.

VAN MONS proposes to triturate together twelve grains of sulphate of zinc, half a dram of acetate of lead, six ounces of rose water to be added, and the solution filtered; he gives this under the name of *Ophthalmic Liquor*.—ELLIS also gives, under the name of *Astringent Collyrium*, a solution of six grains each, of sulphate of zinc and of subacetate of lead, in four ounces of rose water.

It would be better to dissolve the dry acetate in rose water; a scruple, for instance, of the salt in eight ounces of the water.

ELLIS gives an astringent injection for gonorrhœa, of eight grains of acetate of zinc dissolved in four ounces of rose water.

AUGUSTIN quotes a similar injection, which he extols in the same case, consisting of eight or ten grains of the acetate dissolved in four or six ounces of distilled water.

PIERQUIN prescribes a collyrium prepared with one dram and a half of sulphate of zinc, two scruples of acetate of lead, and four ounces each of plantain water, and of rose water.

ANTI-OPHTHALMIC LIQUOR.

Aqua ophthalmica alba Danica. (AUGUSTIN; SPIELMANN)

R. Sulphate of Zinc, one dram.
Acetate of Lead, half a dram.
Camphor, one scruple.
Rose Water, twelve ounces.

Dissolve and filter. (SPIELMANN)

AUGUSTIN prescribes a scruple of sulphate, half a dram of acetate, two drams of spirit of camphor, and ten ounces of distilled water.

ASTRINGENT GARGLE.

Collutorium adstringens seu e Zinco compositum. (SPIELMANN; SWEDIAUR)

R. Rhazes' Trochiscs, one dram.

Sulphate of Zinc, half a dram.
Acetate of Lead, half a scruple.
Rose Water, four ounces.

Mix. (SPIELMANN)

R. Sulphate of Zinc, eight grains.
Solution of Acetate of Lead, eight drops.
Rose Water, four ounces.

Mix, and add, if required, one ounce of camphor mixture. (SWEDIAUR)

ASTRINGENT INJECTION.

Injectio ad Blenorrhœam. (BORIES PH. SYPH.)

R. Sulphate of Zinc, two drams.
Acetate of Lead, four scruples.
Camphor, one scruple and a half.
Aqueous Extract of Opium, one scruple.

Rose Water, from two to four pounds.

Dissolve. (PH. SYPH.)

R. Sulphate of Zinc, } of each, one dram.
Acetate of Lead, }
Alum, }
Camphor, ten grains,
Rose Water, } of each, eight ounces.
Pure Water, }

Dissolve. (BORIES)

The injection may be repeated twice or thrice a day.

TINCTURE OF ACETATE OF ZINC.

Tinctura Acetatis Zinci, Acetas Zinci diluta. (DUBL; COXE; NIEMANN)

Tinctura Acetatis Zinci. (DUBL.)

R. Sulphate of Zinc, } of each,
Acetate of Potass, } one part.

Triturate together, and add

Rectified Spirit, sixteen parts.

Macerate for a week, occasionally agitating, and filter through paper.

It is employed in collyrium, and in injections in blenorrhœa.

ZIRCONA.

This substance is considered to be a metallic oxide; the only compound of it employed in medicine is the following:

SILICATE OF ZIRCONA; HYACINTH.

Hyacinthus.

SYNONIMES:—Hyacinthe, FR.; Hiacinth, GERM.; Jacintho orientale, SPAN.

HISP.; WIRTEM.; GUIBOURT; SPIELMANN.

A very hard, vitreous, infusible stone, of an orange-yellow colour.

THE FIVE PRECIOUS STONES. (WIRTEM.)

R. Garnet,	} of each, an equal part.	Cornelian,	} of each, an equal part.
Hyacinth,		Emerald,	
Sapphire,			

Mix, and pulverize.—This mixture is perfectly inert.

ZIZYPHUS SATIVUS, CAND.—JUJUBE TREE.

SYNONIMES:—Jujubier, FR.; Brustbeerstrauch, GERM.; Azufeifas, SPAN.; Jotlenboom, DUT.; Ginggiolo, ITAL.; Anafegas, PORT.

ANT.; BAVAR.; HISP.; GALL.; FERR.; WIRTEM.; BRUGNATELLI; BERGIUS; GUIBOURT; MURRAY and GMELIN; SPIELMANN.

A shrub of the East Indies, but cultivated in the south of Europe. (*Pentandria Digynia*, LINN.; *Rhamnea*, JUSS.; *Fig. Nouv. Duh.* III. t. 16.)

The fruit, called *Jujubes*; *Jujuba* seu *Bacca Jujuba* seu *Zizyphi*, is employed.

It consists of drupes of the size and form of an olive of a reddish-brown colour, containing under a thick pellicle, a soft yellow-coloured, mucilaginous flesh, of a sweet and mucilaginous taste, and covering an oblong, hard, grooved stone.

AINSLIE says that the Indians prescribe the decoction of the root in certain fevers, but he thinks it does not possess much activity.

PECTORAL SPECIES.

Species pro Decocto Florum et Fructuum.
(PALAT.; WIRTEM.)

R. Root of Liquorice,	two ounces.
Herb of Montpellier	} of each,
Maidenhair,	
Endive,	two drams.
Flowers of Borage,	} of each,
Bugloss,	
Violet,	
Roses,	
Hop,	two drams.
Jujubes,	} of each,
Carob Beans	
Damask Plums,	
Tamarinds,	one ounce.

Cut and mix. (WIRTEM.)

PALAT. omits the carob beans but prescribes twenty each, of jujubes, of damask plums, and of seabastes.

PECTORAL FRUITS. (GALL.)

R. Jujubes,	} of each
Dates, stoned,	

an equal part.

Figs,	} of each,
Raisins,	

Mix.

PULP OF JUJUBES.

Pulpa Jujubarum. (GALL.)

R. Jujubes,	any quantity.
Water,	a sufficient quantity.

Boil till the fruit is softened; then pass the pulp through a hair sieve, mix the water of the decoction with, and evaporate the whole slowly to the proper consistence.

JUJUBE PASTE. (BAVAR.; GALL.; NIEMANN)

R. Jujubes,	one pound.
Gum Senegal,	six pounds.
White Sugar,	five pounds.
Filtered Water,	thirty pounds.
Tincture of Lemon Peel,	one ounce.

Boil the fruit in ten pounds of the water, strain the decoction, and dissolve the gum in the remainder of the water; mix the two strained liquors, add the sugar, and add the whites of five eggs beaten with water, and boil, skimming, till the liquor is reduced to one third; then having strained it through a linen cloth, and evaporated it to the consistence of thick syrup, add the tincture, and continue the evaporation, in a water-bath, till the whole has acquired the consistence of a soft extract; finally pour it into oiled iron-moulds and dry in a stove. (GALL.)

BAVAR.; and NIEMANN direct one part each, of jujubes, and of corinth raisins, to be boiled in thirty-six parts of water till reduced to twenty-four, twice its weight of sugar to be added to the strained decoction, and then one half part of white of egg beaten with a little water; the mixture then having been evaporated to the consistence of syrup, twenty-four parts of mucilage of gum arabic to be added, and the whole evaporated to the proper consistence; then to be poured into moulds, and dried at a gentle heat.

SYRUP OF JUJUBES.

Syrupus de Jujubis. (BRUNS.; WIRTEM.; SPIELMANN)

R. Fresh Jujubes,	three ounces.
-------------------	---------------

Liquorice,	six drams.
Montpellier Maidenhair,	of each,
Flowers of Violet,	five drams.
Seeds of Mallow,	of each,
Quince,	three drams.
White Poppy,	

Spring Water, two pounds and a half.

Boil till reduced to eighteen ounces. Dissolve in this,

Gum Tragacanth,	one dram.
White Sugar,	thirty-two ounces.

Preserve. (BRUNS.; WIRTEM.)

R. Jujubes,	three ounces.
Root of Liquorice,	of each,
Montpellier Maidenhair,	one ounce.
Pearl Barley,	
Water,	forty-four ounces.

Boil down to thirty-six ounces, adding towards the end,

Flowers of Violet,	five ounces.
--------------------	--------------

Then make an emulsion of the strained liquor with

Seeds of Mallow,	five ounces.
Quince,	of each,
White Poppy,	three drams.
Melon,	
Lettuce,	

Then dissolve in the liquor,

Sugar,	twenty-four ounces.
Gum Tragacanth,	three drams.

Boil to the consistence of syrup. (SPIELMANN)

PECTORAL DECOCTION.

Decoctum pectorale. (SARD.; CADET DE GASSICOURT)

R. Jujubes,	half an ounce.
-------------	----------------

Figs, cut,	} of each, half an ounce.
Raisins,	
Liquorice,	
Water,	six pounds.

Boil the fruits in the water till this is reduced to two thirds; then remove from the fire, add the liquorice, infuse for a short time, and strain. (SARD.)

CADET DE GASSICOURT prescribes two ounces of pectoral fruits and two pounds and a half of water, to be boiled down to two pounds.

PECTORAL BROTH. (PIERQUIN; SAINTE-MARIE)

R. Calf's Lung,	} of each, one half.
Neck of Mutton,	
Jujubes,	twenty,
Dates,	twelve.
Damask Raisins,	half an ounce.
Turnips,	two ounces.
Grits,	two drams.
Conserve of Angelica,	one ounce.
Water,	three pounds.

Boil. (PIERQUIN)

R. Calf's Lung,	} of each, one half.
Chicken,	
Jujubes,	twelve or fifteen.
Pearl Barley,	two table spoonfuls.
Water,	two pounds.

Boil down to two thirds, with skimming, and add, a few minutes before removing it from the fire, a handful of fresh leaves of lungwort.

FINIS.

INDEX.



<i>Abies balsamea</i> , i.	318	<i>Acetas plumbi dilutum al-</i>	<i>Acetum colchiticum</i> , i.	539
<i>Abies pectinata</i> , ii.	654	<i>coholisatum</i> , i.	<i>Acetum concentratum</i> , i.	41
<i>Abies picea</i> , i.	21	<i>Acetas plumbi liquidus</i> ,	<i>Acetum convallariæ ma-</i>	
<i>Abietis resina</i> , ii.	654	ii.	<i>jalis</i> , i.	542
<i>Abrotani fœminei herba</i> ,		<i>Acetas plumbi solutus</i> , ii.	<i>Acetum crudum</i> , i.	38
ii.	556	<i>Acetas plumbicum crys-</i>	<i>Acetum cupri corrosum</i> ,	
<i>Abrotani hortensis herba</i> ,		<i>tallisatum</i> , ii.	i.	590
i.	290	<i>Acetas potassæ</i> , ii.	<i>Acetum distillatum con-</i>	
<i>Abrotani maris herba</i> , i.	290	<i>Acetas potassæ aquosus</i> ,	<i>centratum</i> , i.	41
<i>Abrotani montani herba</i> ,		ii.	<i>Acetum dulcificatum</i> , i.	92
ii.	556	<i>Acetas potassæ liquidus</i> ,	<i>Acetum gelu concentra-</i>	
<i>Absinthii albi herba</i> , i.	291	ii.	<i>tum</i> , i.	38
<i>Absinthii maritimi herba</i> ,		<i>Acetas sodæ</i> , ii.	<i>Acetum glaciale</i> , i.	41
i.	24	<i>Acetas superplumbicus</i>	<i>Acetum hystericum</i> , i.	432
<i>Absinthii Romani herba</i> , i.	24	<i>aquosus</i> , ii.	<i>Acetum lavandulæ</i> , ii.	156
<i>Absinthii Pontici herba</i> , i.	24	<i>Acetas tritoxidi ferri</i> , i.	<i>Acetum lithargyri</i> , ii.	376, 386
<i>Absinthium</i> , i.	23	<i>Acetate of alumina</i> , i.	<i>Acetum Maravillianum</i> ,	
<i>Acacia Egyptinea</i> , i.	34	<i>Acetate of cinchonine</i> , i.	i.	115
<i>Acacia nostras</i> , ii.	468	<i>Acetate crystallized of</i>	<i>Acetum medicatum</i> , i.	226
<i>Acacia vera</i> , i.	34	<i>copper</i> , i.	<i>Acetum myrrhæ</i> , i.	763
<i>Acacia false</i> , ii.	509	<i>Acetate of lead</i> , ii.	<i>Acetum nicotianæ</i> , ii.	263
<i>Acacia true</i> , i.	34	<i>Acetate neutral of lead</i> , ii.	<i>Acetum odoratum</i> , ii.	231,
<i>Acanthii vulgaris herba</i> ,		<i>Acetate of morphine</i> , ii.	523	
ii.	273	<i>Aceti naphtha</i> , i.	<i>Acetum odoriferum</i> , ii.	231
<i>Acanthus mollis</i> , i.	34	<i>Acetis lixivix</i> , ii.	<i>Acetum opii</i> , ii.	275
<i>Acari</i> , i.	377	<i>Acetis plumbi</i> , ii.	<i>Acetum papaveris erratici</i> ,	
<i>Accipenser stellatus</i> , i.	697	<i>Acetosæ officinalis radix</i> ,	ii.	336
<i>Accipenser sturio</i> , i.	697	ii.	<i>Acetum plumbi</i> , ii.	376
<i>Acetabularia Mediterranea</i> ,		<i>Acetosæ pratensis radix</i> ,	<i>Acetumprimulæ veris</i> , ii.	465
ii.	2	ii.	<i>Acetum prophylacticum</i> ,	
<i>Acetas barii</i> , i.	328	<i>Acetosæ rotundifoliæ ra-</i>	i.	226
<i>Acetas ammoniæ</i> , i.	195	<i>dix</i> , ii.	<i>Acetum pseudo-narcissi</i> ,	
<i>Acetas ammoniæ aquosus</i> ,		<i>Acetosellæ herba</i> , ii.	ii.	259
i.	195	<i>Acetum acetosum</i> , i.	<i>Acetum purificatum</i> , i.	38
<i>Acetas ammoniæ dilutus</i> ,		<i>Acetum alliaceum</i> , i.	<i>Acetum purum concentra-</i>	
i.	195	<i>Acetum allii</i> , i.	<i>tum</i> , i.	38
<i>Acetas ammoniæ liqui-</i>		<i>Acetum allii compositum</i> ,	<i>Acetum quatuor latro-</i>	
<i>dum</i> , i.	195	i.	<i>num</i> , i.	115
<i>Acetas ammoniæ solutum</i> ,		<i>Acetum anti-emeticum</i> , i.	<i>Acetum radicatam</i> , i.	40
i.	195	<i>Acetum antisepticum</i> , i.	<i>Acetum rorismarini</i> , ii.	523
<i>Acetas calcis</i> , i.	363	<i>Acetum antisepticum</i> , i.	<i>Acetum rosæ</i> , ii.	516
<i>Acetas cupri</i> , i.	595	<i>Acetum aromaticum</i> , i.	<i>Acetum rubi idæi</i> , ii.	533
<i>Acetas cupri imperfectus</i> ,		ii.	<i>Acetum rutaceum</i> , ii.	537
i.	590	<i>Acetum belladonnæ</i> , i.	<i>Acetum rutæ</i> , ii.	537
<i>Acetas cupricus</i> , i.	595	<i>Acetum bezoardicum</i> , i.	<i>Acetum rutæ camphora-</i>	
<i>Acetas ferri</i> , i.	663	ii.	<i>tum</i> , ii.	537
<i>Acetas hydrargyri</i> , ii.	95	<i>Acetum camphoratum</i> , i.	<i>Acetum rutæ composi-</i>	
<i>Acetas ferri liquidus</i> , i.	663	<i>Acetum capsici spirituo-</i>	<i>tum</i> , i.	432
<i>Acetas kalicum</i> , ii.	441	<i>sum</i> , ii.	<i>Acetum salviæ</i> , ii.	546
<i>Acetas lixivix solutus</i> , ii.	442	<i>Acetum caryophylli</i> , i.	<i>Acetum sambuci</i> , ii.	553
<i>Acetas morphæi</i> , ii.	246	<i>Acetum chalybeatum</i> , i.	<i>Acetum sambucinum</i> , ii.	553
<i>Acetas morphinæ</i> , ii.	246	<i>Acetum cochleariæ armo-</i>	<i>Acetum saturni</i> , ii.	376
<i>Acetas natricum</i> , ii.	614	<i>raciæ</i> , ii.	<i>Acetum scillæ</i> , ii.	573
<i>Acetas plumbi</i> , ii.	382	<i>Acetum coffeanum extem-</i>	<i>Acetum scilliticum</i> , ii.	573
<i>Acetas plumbi acidulus</i>		<i>poraneum</i> , ii.	<i>Acetum terebinthiæ</i> , i.	40
<i>siccus</i> , ii.	382	<i>Acetum colchici</i> , i.	<i>Acetum theriacale</i> , ii.	118

Acetum tunicæ, i.	610	Acid pyrotartaric, i.	74	Acidum manganescicum,	
Acetum veratri, ii.	713	Acid rheumatic, ii.	489	ii.	217
Acetum vini camphora-		Acid sorbic, ii.	619	Acidum muriaticum, i.	53
tum, i.	373	Acid sulpho-sinapic, i.	531	Acidum muriaticum de-	
Acetum vini concentra-		ii.	588	phlogisticatum, i.	458
tum, i.	40	Acid sulphuric, i.	75	Acidum muriaticum ni-	
Acetum vini distilla-		Acid sulphurous, i.	79	troso-oxygenatum, i.	56
tum, i.	38	Acid tartaric, i.	80	Acidum muriaticum oxy-	
Acetum vini radicale, i.	40	Acid uric, i.	83	genatum, i.	458
Achillea ageratum, i.	34	Acidum acetii, i.	38	Acidum muriaticum oxy-	
Achillea atrata, i.	34	Acidum aceticum, i.	38	genatum ad contagio-	
Achillea millefolium, i.	35	Acidum aceticum aroma-		sam, i.	458
Achillea moschata, i.	34	ticum, i. 44, ii.	523	Acidum nitri, i.	67
Achillea nana, i.	34	Acidum aceticum cam-		Acidum nitri concentra-	
Achillea ptarmica, i.	37	phoratum, i.	44, 373	tum, i.	68
Achmellæ herba, ii.	621	Acidum aceticum dilu-		Acidum nitri fumans, i.	68
Achras mammosa, i.	470	tum, i.	38	Acidum nitricum, i.	67
Acid acetic, i.	38	Acidum aceticum scilliti-		Acidum nitricum alcoh-	
Acid alcoholized hydro-		cum, ii.	573	lisatum, i.	94
chloric, i.	56	Acidum aceticum tenue,		Acidum nitricum dilu-	
Acid alcoholized hydro-		i.	38	tum, i.	69
cyanic, i.	59	Acidum acetosellæ, i.	71	Acidum nitroso-nitri-	
Acid alcoholized nitric,		Acidum æreum, i.	49	cum, i.	68
i.	70	Acidum antimonium,		Acidum nitrosum, i. 68, 71	
Acid alcoholized sulphu-		i.	240	Acidum nitrosum alcoh-	
ric, i.	79	Acidum antimoniosum,		lisatum, i.	94
Acid antimoniac, i.	240	i.	236	Acidum nitrosum dilu-	
Acid arsenic, i.	286	Acidum arsenicale, i.	286	tum, i.	69
Acid arsenious, i. 281, 283		Acidum arsenicosum, i.	283	Acidum phosphori, i.	72
Acid benzoic, i.	46	Acidum azoticum, i.	67	Acidum phosphoricum,	
Acid boracic, i.	48	Acidum benzoicum, i.	46	i.	72
Acid carbonic, i.	49	Acidum benzoës, i.	46	Acidum phosphorosum,	
Acid cathartic, i.	398	Acidum boracicum, i.	48	i.	72
Acid cavadic, ii.	714	Acidum boracis, i.	48	Acidum prussicum, i.	57
Acid citric, i.	52	Acidum borussicum, i.	57	Acidum pyrotartaricum,	
Acid dilute nitric, i.	69	Acidum borussicum al-		i.	74
Acid dilute sulphuric, i.	76	cophilisatum, i.	59	Acidum sacchari, i.	71
Acid ellagic, i.	692	Acidum carbonicum, i.	49	Acidum saccharinum, i.	71
Acid of ants, i. 40, 676		Acidum citricum, i.	52	Acidum salis culinaris,	
Acid hydriodic, ii.	121	Acidum cretaceum, i.	49	i.	53
Acid hydrochloro-nitric,		Acidum hydriodicum,		Acidum salis marini, i.	53
i.	56	ii.	121	Acidum succini, i.	75
Acid hydrosulphuric i.	64	Acidum hydrocarboni-		Acidum succineum, i.	75
Acid hydrocyanic, i.	57	cum, i.	71	Acidum sulphuricosum,	
Acid hydrocyanic of Har-		Acidum hydrochlori-		i.	79
les, i.	59	cum, i.	52	Acidum sulphuricum, i.	75
Acid hydrocyanic veget-		Acidum hydrochloro-ni-		Acidum sulphuricum	
able, i.	64	tricum, i.	56	tenue, i.	76
Acid iodic, ii.	121	Acidum hydrocyanicum,		Acidum sulphuricum con-	
Acid krameric, ii.	147	i.	57	densatum, i.	76
Acid laccic, ii.	150	Acidum hydrocyanicum		Acidum sulphuricum al-	
Acid meconic, ii.	274	alcoholisatum, i.	59	cophilisatum, i.	79
Acid menispermic, i.	529	Acidum hydrosulphuri-		Acidum sulphuricum	
Acid muriatic, i.	53	cum, i.	64	aquosum, i.	76
Acid nitric flavoured, i.	69	Acidum hydrothionicum,		Acidum sulphuris vola-	
Acid nitric, i.	67	i.	64	tile, i.	79
Acid nitro-muriatic, i.	56	Acidum hydrothionicum		Acidum tartari, i.	80
Acid nitrous, i.	71	liquidum, i.	65	Acidum tartaricum, i.	80
Acid oxalic, i.	71	Acidum hydrocyanicum		Acidum tartarosum, i.	80
Acid oxymuriatic, i.	459	dilutum spirituosum		Acidum vitrioli aromati-	
Acid phosphoric, i.	72	aquosum, i.	59	cum, ii.	175
Acid phosphorous, i.	72	Acidum, iodicum, ii.	121	Acidum vitrioli deplogis-	
Acid polygalic, ii.	403	Acidum ligni pyro-oleo-		tatum, i.	76
Acid prussic, i.	57	sum, i.	39	Acidum vitrioli depura-	
Acid pyrolignic, i.	39	Acidum limonorum, i.	52	tum, i.	76

Acidum vitrioli dilutum, i.	76	Ærugo crystallisata, i.	595	Æther vitrioli aquosus, ii.	100
Acidum vitrioli purissimum, i.	76	Ærugo præparata, i.	591	Æther vitriolicus, i.	97
Acidum vitrioli rectificatum, i.	76	Æs viride crystallisatum, i.	595	Ætheroleum absinthii, i.	24
Acidum vitriolicum vinosum, i.	79	Æs ustum, i. 581, ii.	369	Ætheroleum anethi, i.	221
Acidum uricum, i.	83	Æsculus, i.	89	Ætheroleum anisi, ii.	354
Acidum zooticum, i.	57	Æsculus hippocastanum, i.	89	Ætheroleum anthos, ii.	524
Acidum zootinicum, i.	57	Æther acetic, i.	91	Ætheroleum aurantii, i.	507
Acmellæ herba, ii.	621	Æther aceticus, i.	91	Ætheroleum baccarum, juniperi, ii.	139
Aconita, i.	83	Æther aceticus martialis, i.	664	Ætheroleum calami radicis, i.	87
Aconite, i.	83	Æther alcoholicus, i.	99	Ætheroleum cardamomi, i.	395
Aconitine, i.	83	Æther alcoholized acetic, i.	92	Ætheroleum carui, i.	397
Aconiti anthora, i.	83	Æther alcoholized hydrochloric, i.	94	Ætheroleum coriandri, i.	559
Aconiti racemosi radix, i.	89	Æther alcoholized nitric, i.	96	Ætheroleum caryophyllorum aromaticorum, i.	400
Aconitinum, i.	83	Æther ammoniacal, i.	777	Ætheroleum chamæmelis, i.	448
Aconitum, camarum, i.	83	Æther anti-odontalgic, i.	374	Ætheroleum cinamomi, ii.	165
Aconitum napellus, i.	83	Æther chloruric, i.	94	Ætheroleum cubebarum, ii.	362
Aconitum neomontanum, i.	83	Æther cypriacus, i.	589	Ætheroleum cumini, i.	579
Aconitum paniculatum, i.	83	Æther hydrochloric, i.	93	Ætheroleum dictamni, ii.	322
Acor aceticus crystallinus, i.	41	Æther martialis, i.	659	Ætheroleum hyssopi, ii.	112
Acor benzoinus, i.	46	Æther mercurialis, ii.	72	Ætheroleum juniperi ligni, ii.	139
Acor boracius, i.	48	Æther muriaticus, i.	93	Ætheroleum lavandulæ, ii.	157
Acor succineus, i.	75	Æther muriaticus spurius, i.	94	Ætheroleum macidis, ii.	256
Acor sulphuris condensatus, i.	76	Æther nitric, i.	94	Ætheroleum melissæ, ii.	229
Acor sulphuris merus, i.	76	Æther nitricus, i.	94	Ætheroleum millefolii, i.	36
Acor tartaricus, i.	80	Æther nitrosus, i.	94	Ætheroleum origani, ii.	324
Acori adulterini radix, ii.	130	Æther phosphorated, ii.	349	Ætheroleum petroselinii, i.	269
Acori palustris radix, ii.	130	Æther phosphoric, i.	97	Ætheroleum pimentæ, ii.	362
Acorns torrifed, ii.	477	Æther phosphoricus, i.	97	Ætheroleum rorismarini, ii.	524
Acorus calamus, i.	85	Æther phosphoro-phosphoricus, ii.	349	Ætheroleum rosarum, ii.	517
Actæa racemosa, i.	89	Æther rectificatus, i.	97	Ætheroleum rutæ, ii.	538
Actæa spicata, i.	89	Æther rectified, i.	97	Ætheroleum sabinæ, ii.	143
Actes baccæ, ii.	551	Æther spirituosus, i.	99	Ætheroleum salviæ, ii.	546
Acus veneris radix, i.	622	Æther sulphuric, i.	97	Ætheroleum santali-citrini, ii.	196
Adderstongue, ii.	273	Æther sulphuric with alcohol, i.	100	Ætheroleum sassafra ligni, ii.	184
Adeps, i.	315	Æther sulphuricus, i.	97, 98	Ætheroleum saturejæ, ii.	558
Adeps suillus, i.	316	Æther sulphuricus acidus, i.	72	Ætheroleum serpylli, ii.	690
Adeps tartare stibii medicatus, i.	261	Æther sulphuricus cum alcohole, i.	100	Ætheroleum succini, ii.	630
Adianti albi herba, i.	299	Æther sulphuricus cum alcohole aromaticus, i.	101	Ætheroleum tanacetii, ii.	651
Adianthi aurei herba, ii.	409	Æther sulphuricus cum alcohole aromaticus, ii.	175	Ætheroleum terebinthinæ, ii.	663
Adianti Canadensis herba, i.	89	Æther sulphuricus camphoratus aquosus, i.	371	Ætheroleum thymi, ii.	689
Adiantum capillus veneris, i.	89	Æther sulphuricus ferratus, i.	659	Ætheroleum valerianæ, ii.	711
Adiantum pedatum, i.	89	Æther sulphuricus phosphoricus, ii.	349	Ætheroleum zingiberis, i.	205
Adipocera cetosa, i.	440	Æther terebinthinated, ii.	664	Æthiops alkalisatus, ii.	18
Admellæ herba, ii.	621			Æthiops antimonial, ii.	30
Ægagrophilus, i.	338				
Aer dephlogisticatus, ii.	328				
Aer fixus, i.	49				
Aer mephiticus, i.	49				
Aer purus, ii.	328				
Aer vitalis, ii.	328				
Ærugo, i.	590				

<i>Æthiops antimonialis</i> , ii.	30	<i>Alcohol ammoniacatum</i> ,	<i>Alcohol sulphuricatum</i> ,
<i>Æthiops auripigmenta-</i>		<i>aromatisatum</i> , i.	i.
<i>lis</i> , ii.	25	<i>Alcohol ammoniæ et</i>	79
<i>Æthiops graphitic</i> , ii.	26	<i>guaiaci</i> , i.	736
<i>Æthiops graphiticus</i> , ii.	26	<i>Alcohol ammoniatum</i> , i.	172
<i>Æthiops martial</i> , i.	640	<i>Alcohol ammoniatum fœ-</i>	
<i>Æthiops martialis</i> , i.	640	<i>tidum</i> , i.	304
<i>Æthiops mineral</i> , ii.	28	<i>Alcohol anisi</i> , ii.	355
<i>Æthiops mineralis</i> , ii.	28	<i>Alcohol anisi composi-</i>	
<i>Æthiops narcoticus</i> , ii.	28	<i>tum</i> , ii.	355
<i>Æthiops orpimental</i> , ii.	25	<i>Alcohol anthos compo-</i>	
<i>Æthiops saccharatus</i> , ii.	16	<i>situs</i> , ii.	528
<i>Æthiops vegetabilis</i> , i.	390	<i>Alcohol cum aromatibus</i>	
<i>Æthiops vegetable</i> , i.	679	<i>compositus</i> , ii.	173
<i>Aethusa cynapium</i> , i.	102	<i>Alcohol aromatic</i> , ii.	182
<i>Ætites</i> , i.	643	<i>Alcohol aromaticum</i> , ii.	168
<i>Agaric</i> , ii.	408	<i>Alcohol aromaticum sul-</i>	
<i>Agaric mineral</i> , i.	352	<i>phuricum</i> , ii.	175
<i>Agaric of the oak</i> , i.	340	<i>Alcohol aromaticus am-</i>	
<i>Agaricus albus</i> , ii.	408	<i>moniatus</i> , i.	174
<i>Agaricus chirurgorum</i> , i.	340	<i>Alcohol cum balsamo pe-</i>	
<i>Agaricus mineralis</i> , i.	352	<i>ruviano</i> , i.	320
<i>Agaricus muscarius</i> , i.	165	<i>Alcohol camphoræ sapo-</i>	
<i>Agaricus quernus</i> , i.	340	<i>naceum</i> , ii.	461
<i>Agaricus trochiscatus</i> ,		<i>Alcohol camphoratum</i> , i.	377
ii.	408	<i>Alcohol camphoratum</i>	
<i>Agathophyllum aromati-</i>		<i>crocatum</i> , i.	373
<i>cum</i> , i.	102	<i>Alcohol castoriatum</i> , i.	428
<i>Agave radix</i> , i.	103	<i>Alcohol cum citro au-</i>	
<i>Agave Americana</i> , i.	103	<i>rantio</i> , i.	515
<i>Ageloide</i> , i.	723	<i>Alcohol concentrated</i> , i.	108
<i>Agerati herba</i> , i.	35	<i>Alcohol cum crotone cas-</i>	
<i>Agrimonia Eupatoria</i> , i.	103	<i>carilla</i> , i.	569
<i>Agrimony hemp</i> , i.	628	<i>Alcohol diluted</i> , i.	106
<i>Agnel</i> , i.	401	<i>Alcohol dilutius</i> , i.	105
<i>Ajuga Genevensis</i> , i.	103	<i>Alcohol dilutum</i> , i.	106
<i>Ajuga pyramidalis</i> , i.	103	<i>Alcohol ferratus</i> , i.	658
<i>Ajuga reptans</i> , i.	103	<i>Alcohol formicarum dis-</i>	
<i>Alabaster</i> , i.	362	<i>tillatum</i> , i.	676
<i>Alabastrites</i> , i.	362	<i>Alcohol fortius</i> , i.	105
<i>Alabastrum</i> , i.	362	<i>Alcohol cum guaiaco of-</i>	
<i>Alami radix</i> , i.	291	<i>ficinale ammoniatus</i> , i.	736
<i>Albedo corticum citri</i> , i.	517	<i>tus</i> , ii.	528
<i>Album græcum</i> , i.	360	<i>Alcohol iodii</i> , ii.	120
<i>Albumen aluminosum</i> , i.	163	<i>Alcohol juniperi com-</i>	
<i>Alburnum sambuci</i> , ii.	551	<i>positum</i> , ii.	140
<i>Alcahest Glauberi</i> , ii.	422	<i>Alcohol nitrico-æthere-</i>	
<i>Alcannæ orientalis radix</i> ,		<i>um</i> , i.	94
ii.	185	<i>Alcohol opii dilutum</i> , ii.	284
<i>Alcannæ radix</i> , i.	216	<i>Alcohol cum opio</i> , ii.	281
<i>Alcannæ veræ radix</i> , ii.	185	<i>Alcohol rectificatus</i> , i.	109
<i>Alceæ radix</i> , ii.	216	<i>Alcohol rectified</i> , i.	108
<i>Alchemilla vulgaris</i> , i.	104	<i>Alcohol phosphorated</i> ,	
<i>Alchornea latifolia</i> , i.	112	ii.	349
<i>Alcohol absolute</i> , i.	167	<i>Alcohol potassæ</i> , ii.	414
<i>Alcohol absolutum</i> ,		<i>Alcohol cum rheo</i> , ii.	501
i.	107, 108, 109	<i>Alcohol salviæ vulnera-</i>	
<i>Alcohol aceti</i> , i.	41	<i>rius</i> , ii.	547
<i>Alcohol æthereus ferra-</i>		<i>Alcohol saponatum</i> , ii.	460
<i>tus</i> , i.	659	<i>Alcohol serpylli composi-</i>	
<i>Alcohol albuminous</i> , ii.	326	<i>tum</i> , i.	174
<i>Alcohol alcalisatum</i> , i.	107	<i>Alcohol cum succino</i> , ii.	629
<i>Alcohol alkalisatum</i> , ii.	414	<i>Alcohol cum sulphate</i>	
<i>Alcohol cum aloe perfo-</i>		<i>ferri tartarizatus</i> , i.	670
<i>liata</i> , i.	120		

Alkali vegetabile vitriola- tum, ii.	427	Aloes Socotrine, i.	117	Ammonia caustica liquida, i.	168
Alkali volatile acetatum, i.	195	Aloysiæ folia, ii.	717	Ammonia, diluted solu- tion of, i.	171
Alkali volatile aeratum, i.	182	Alpinia galanga, i.	682	Ammonia liquida, i.	168
Alkali volatile ammonia- cale, i.	182	Alsine media, i.	142	Ammonia præparata, i.	182
Alkali volatilis anisatus, i.	173	Althæa officinalis, ii.	142	Ammonia muriatica, i.	191
Alkali volatile aquosum, i.	183	Althæa rosea, i.	154	Ammonia pura, i.	168
Alkali volatile causticum, i.	168, 182	Alticis radix, i.	296	Ammonia pura liquida, i.	168
Alkali volatile nitratum, i.	190	Alum, i.	158	Ammonia, solution of, i.	168
Alkali volatile ex sale am- moniaco, i.	182	Alum compound solu- tion of i.	162	Ammonia alcoholisata, i.	172
Alkali volatile tartarisa- tum, ii.	456	Alum with dragon's blood, i.	159	Ammoniaco-mercurialisub- deutosulphate, ii.	83
Alkannæ spuria radix, i.	216	Alum dried, i.	158	Ammoniacum, i.	168
Alkanet yellow, ii.	273	Alum Roche, i.	158	Ammoniacum causticum, i.	168
Alkekenge, ii.	350	Alum Roman, i.	158	Ammoniacum succinatum, i.	180
Alkekengi bacca, ii.	350	Alum root, ii.	4	Ammoniacum volatile mite, i.	182
Alkermes liquid, ii.	712	Alum saccharated, i.	160	Ammoniaretum auri, i.	314
Alleluja herba, ii.	327	Alum, solution of, i.	162	Ammoniaretum cupri, i.	586
All-heal, ii.	107	Alum whey, i.	165	Ammonium carbonicum, i.	184
Alliariæ herba, i.	623	Alum whey with cinna- mon, i.	165	Ammonium carbonicum, i.	182
Allii alpini radix, i.	116	Alumen, i.	158	Ammonium muriaticum, i.	191
Allii montani latifolii ma- culati radix, i.	116	Alumen calcinatum, i.	158	Ammonium muriaticum depuratum, i.	191
Allium ascalonicum, i.	112	Alumen draconisatum, i.	159	Ammonium muriaticum martiale, i.	661
Allium cepa, i.	112	Alumen exsiccatum, i.	158	Ammonium muriaticum martiatum, i.	661
Allium porrum, i.	113	Alumen kinosatum, i.	159	Ammonium subcarbone- um, i.	182
Allium sativum, i.	114	Alumen plumosum, i.	295	Ammonium subcarboni- cum pyro-oleosum, i.	184
Allium scorodoprasum, i.	116	Alumen à Rocca, i.	158	Ammonium sulphuricum, i.	190
Allspice, ii.	359	Alumen rupei, i.	158	Ammonium tartaricum, ii.	456
Almonds bitter, i.	206	Alumen rupeum, i.	158	Ammoniuret of copper, i.	582
Almond common, i.	205	Alumen saccharinum, i.	160	Ammoniuret of mercury, ii.	81
Almond dwarf, i.	205	Alumen sessile, i.	362	Ammoniuretam cupri, i.	586
Almond sweet, i.	205	Alumen siccatum, i.	158	Ammoniuretam hydrar- gyri, ii.	81
Alnus glutinosa, i.	116	Alumen tinctum, i.	159	Amomi semen, ii.	359
Aloe American, i.	103	Alumen ustum, i.	158	Amomum granum para- disi, i.	203
Aloe Barbadosensis, i.	117	Alumina, i.	155	Amomum maximum, i.	203
Aloe caballina, i.	117	Alumina factitious, i.	156	Amomum racemosum, i.	394
Aloe Capensis, i.	117	Alumine, i.	155	Amomum repens, i.	394
Aloe depurata, i.	117	Aluminis spiratus, i.	75	Amomum zerumbet, i.	203
Aloe hepatica, i.	117	Aluminii oxydum, i.	155	Amomum zingiber, i.	203
Aloe linguæ-formis, i.	117	Aluminium, i.	155	Amoraci herba, ii.	322
Aloe lota, i.	117	Aluminium, oxide of, i.	155	Amygdalæ, i.	205
Aloe lucida, i.	117	Aluminium, silici-phtho- ruret of, i.	157	Amygdalæ amaræ, i.	206
Aloe pellucida, i.	117	Amadou, i.	340	Amygdalis communis, i.	205
Aloe perfoliata, i.	117	Amanila muscaria, i.	165	Amygdalæ dulces, i.	205
Aloe præparata, i.	117	Amaranthæ luteæ flores, ii.	4	Amygdalus, i.	205
Aloe purificatum, i.	118	Amaryllyis Zeylanica, ii.	563	Amygdalus nana, i.	205
Aloe rosata, i.	111	Ambarum cineritium, i.	166	Amylum, i.	213
Aloe Socotrina, i.	117	Ambarvalis radix, ii.	400	Amylum hordei, ii.	7
Aloe spicata, i.	117	Amber, ii.	629	Amyris elemifera, ii.	486
Aloe violata, i.	118	Amber black, i.	682	Amyris Gilcadensis, i.	319
Aloe vulgaris, i.	117	Ambergris, i.	166		
Aloe wood, ii.	192	Ambluti radix, ii.	368		
Aloes, i.	117	Ambra, i.	166		
Aloes Barbadoes, i.	117	Ambra ambrosiaca, i.	166		
Aloes caballine, i.	117	Ambra florida, ii.	629		
Aloes, Cape, i.	117	Ambra grisea, i.	166		
Aloes hepatic, i.	117	Ambra liquida, ii.	627		
Aloes horse, i.	117	Ambra vera, i.	166		
		Ambutæ radix, i.	505		
		Amianthus, i.	295		
		Ammios veri semen, ii.	592		
		Ammonia, i.	168		
		Ammonia caustica, i.	168		

- Amyris opobalsamum, i. 214, 319
 Anacardium longi folium, i. 215
 Anagallidis maris herba, i. 215
 Anagallis Phœnicia, i. 215
 Anarrichas lupus, i. 359
 Anatron, ii. 430
 Anchusa officinalis, i. 215
 Anchusa tinctoria, i. 216
 Anchusæ radix, i. 215, 216
 Andropogon schœnanthus, i. 216
 Anemone garden, i. 217
 Anemone meadow, i. 217
 Anemone nemorosa, i. 217
 Anemone pratensis, i. 217
 Anemone pulsatilla, i. 217
 Anemoneum, i. 217
 Anethi hortensis herba, i. 221
 Anethi ursini radix, ii. 198
 Anethum fœniculum, i. 218
 Anethum graveolens, i. 220
 Angelica archangelica, i. 221
 Angelica garden, i. 221
 Angelica preserved stalks of, i. 222
 Angelica sylvestris, i. 222
 Angelica tree, i. 270
 Angelica wild, i. 222
 Angelicæ hortensis radix, i. 222
 Angelicæ sativæ radix, i. 222
 Anguis odoratus, i. 352
 Angular-leaved physic-nut, ii. 134
 Anguriæ semen, i. 578
 Angustura bark, i. 228
 Angustura bark spurious, i. 228
 Angustura ferruginea, i. 228
 Angustura spuria, i. 228
 Angustura vera, i. 228
 Anima rhabarbari, ii. 495
 Anima rhei, ii. 495, 497
 Anise, ii. 352
 Anise Indian, ii. 113
 Anisi vulgaris semen, ii. 353
 Anisum Sinense, ii. 114
 Anisum stellatum, ii. 114
 Anserinæ radix, ii. 465
 Antennaria dioica, i. 230
 Anthemis cotula, i. 446
 Anthemis nobilis, i. 446
 Anthemis pyrethrum, i. 230
 Anthemis tinctoria, i. 446
 Anthoræ radix, i. 83
 Anthos herba, ii. 522
 Anthos sylvestris herba, ii. 185
 Anthrenus muscorum, i. 377
 Antidotus Mithridatica Damocratis, ii. 304
 Antihectic Potier's, ii. 624
 Antihecticum Poterii, ii. 624
 Antimonii clyssus, i. 79
 Antimonii minera cum sulphure mineralizata, i. 241
 Antimonii oxydum nitromuriatum, i. 254
 Antimonii oxydum vitrifactum, i. 235
 Antimonii et potassæ tartas, i. 255
 Antimonii subsulphas, i. 235
 Antimonii sulphuretum, i. 241
 Antimonii sulphuretum præcipitatum, i. 241
 Antimonii vitrum i. 235
 Antimonii vitrum hyalcynthinum, i. 235
 Antimonium, i. 232
 Antimonium calcinatum cum nitro, i. 240
 Antimonium crudum, i. 241
 Antimonium crudum præparatum, i. 241
 Antimonium diaphoreticum lotum, i. 236
 Antimonium diaphoreticum martiale, i. 236
 Antimonium diaphoreticum non ablutum, i. 240
 Antimonium diaphoreticum rubrum, i. 233
 Antimonium muriatum, i. 254
 Antimonium nativum, i. 241
 Antimonium nitratum, i. 240
 Antimonium salitum, i. 254
 Antimonium tartarissatum, i. 255
 Antimonium vitrifactum, i. 235
 Antimonium ustum, i. 240
 Antimony, i. 232
 Antimony tartarized, i. 255
 Antimony vitrified, i. 235
 Antimony Wilson's chartart, i. 235
 Antiquarium, ii. 46
 Antirrhini herba, ii. 199
 Antirrhinum majus, i. 266
 Antophylli, i. 399
 Apii montani radix, ii. 581
 Apis mellifica, i. 433, ii. 467
 Apium graveolens, i. 266
 Apium petroselinum, i. 268
 Apochylisma berberum, i. 335
 Apocynum androsæmifolium, i. 270
 Apozem alterative and diuretic, i. 297
 Apozem alterative and tonic, ii. 613
 Apozem anti-icteric, i. 496
 Apozem antiscorbutic, i. 533
 Apozem aperient, ii. 613
 Apozem aperitive, i. 297
 Apozem diuretic, i. 113, 297
 Apozem febrifuge and purgative, i. 496
 Apozem febrifuge and tonic, i. 494
 Apozem ferruginous aperitive, i. 268
 Apozem Lieutaud's refreshing, ii. 10
 Apozem pectoral and incisive, ii. 568
 Apozem purgative, i. 420
 Apozem tonic, ii. 531
 Apozem tonic and antiherpetic, i. 677
 Apozem tonic and demulcent, i. 495
 Apozema malti, ii. 214
 Apple balm, ii. 242
 Apple devil's, i. 605
 Apple of Peru, i. 605
 Apple sorb, ii. 619
 Apple tree, ii. 214
 Application anticephalgic, i. 372
 Application antispasmodic, i. 425
 Application antispasmodic and tonic, ii. 315
 Application aromatic, ii. 235
 Application detergent, ii. 18
 Application hypnotic, ii. 99
 Application purgative, i. 572
 Application resolvent, i. 168
 Application resolvent, ii. 226, 603
 Application sedative, ii. 276
 Application vermifuge, i. 572
 Aqua absinthii stillatitia, i. 24
 Aqua acetatis ammoniæ, i. 195
 Aqua acidi carbonici, i. 49
 Aqua acidulata cum acido carbonico, i. 49
 Aqua acoustica ludovici, ii. 710
 Aqua aërata, i. 49
 Aqua aërata alkalina, ii. 427
 Aqua aëris fixi, i. 49
 Aqua alkalina oxymuriatica, ii. 440
 Aqua aloetica mercurialis, ii. 35
 Aqua aluminosa, i. 162
 Aqua amara, ii. 212
 Aqua ammoniæ, i. 183
 Aqua ammoniæ carbonatis, i. 184
 Aqua amygdalarum amarum, i. 63

Aqua angelica viennensis, ii.	219	Aqua castorei vinosa, i.	426	Aqua epidemica Londinensis, i.	224
Aqua Anhaltina, ii.	666	Aqua catapultarum, ii.	547	Aqua epileptica Langii, i.	542
Aqua anisi, ii.	355	Aqua caustica ad condylomata, ii.	71, 91	Aqua foetida, ii.	305
Aqua anthos, ii.	527	Aqua celestis, i.	596	Aqua exsiccans, ii.	378
Aqua anthos composita, ii.	528	Aqua cerasorum nigrorum, i.	62	Aqua florum acaciæ, ii.	468
Aqua anti-hysterica, i.	426	Aqua cerasorum nigrorum alcoholica, i.	63	Aqua florum aurantii, i.	508
Aqua antimelancholica, i.	565	Aqua chlorinii, i.	459	Aqua florum chamomillæ vinosa, i.	448
Aqua antinephritica, ii.	606	Aqua cinnamomi, ii.	166	Aqua florum citri aurantii, i.	508
Aqua apoplectica, ii.	528	Aqua cinnamomi bu-glossata, ii.	167	Aqua florum naphæ, i.	508
Aqua armoraciæ, ii.	482	Aqua cinnamomi cordialis, ii.	167	Aqua florum sambuci, ii.	554
Aqua aromatica, ii. 168, 230, 549		Aqua cinnamomi cydoniata, ii.	168	Aqua foetida, i.	684
Aqua aromatica cum opio, ii.	280	Aqua cinnamomi hordeata, ii.	167	Aqua fortis, i.	62
Aqua atriplicis foetida, i.	458	Aqua cinnamomi simplex, ii.	166	Aqua fragariæ, i.	677
Aqua barytæ, i.	328	Aqua cinnamomi spirituosa, ii.	167	Aqua fragorum, i.	677
Aqua barytica, i.	326	Aqua cinnamomi vinosa, ii.	167	Aqua ad gingivas, i.	163
Aqua benedicta Rulandi, i.	258	Aqua citri, i.	518	Aqua Goulardi, ii.	378
Aqua benedicta serpylli, ii.	691	Aqua citri aurantii, i.	508	Aqua grisea, ii.	86
Aqua benzoica, i.	331	Aqua citri medicæ, i.	518	Aqua grysea Gohlii, ii.	86
Aqua bryoniæ composita, i.	427	Aqua cochleariæ, i.	534	Aqua hepatica, i.	65
Aqua calami radices, i.	87	Aqua Coloniensis, i.	518	Aqua hepatica acidulata, i.	65
Aqua calcariae acidula, i.	358	Aqua confortativa, i.	63	Aqua hirundinum cum castoreo, i.	426
Aqua calcariae sulphurato-stibiata, i.	254	Aqua confortiva, ii.	167	Aqua hordeata cum gummi, ii.	8
Aqua calcariae ustæ, i.	347	Aqua cornu cervi citrata, i.	702	Aqua hordei, ii.	7
Aqua calcis, i.	347	Aqua corticum aurantium, i.	508	Aqua Hungarica, ii.	528
Aqua calcis cum cera, i.	437	Aqua corticum aurantium alcoholica, i.	509	Aqua hydrargyri muratici corrosivi cum opio, ii.	76
Aqua calcis cerata, i.	437	Aqua corticum aurantium spirituosus, i.	509	Aqua hydrargyri muratici mitis nigra, ii.	34
Aqua calcis cum china, i.	488	Aqua corticum aurantium vinosa, i.	509	Aqua hydrocyanica vegetabilis, i.	64
Aqua calcis composita, ii.	183	Aqua corticum citrorum, i.	518	Aqua hydrosulphurata, i.	65
Aqua calcis muriatis, i.	363	Aqua cosmetica, i.	330	Aqua hysterica, i.	430
Aqua calcis vivæ, i.	347	Aqua cupri ammoniacalis, i.	590, 596	Aqua immortalis, i.	559
Aqua calcis vulneraria, ii.	367	Aqua cupri ammoniaci, i.	590	Aqua juniperi, ii.	139
Aqua camphoræ, i.	368	Aqua cupri ammoniati, i.	587	Aqua kali, ii.	422
Aqua camphorata Bateana, i.	584	Aqua cupri ammonio-muriatici, i.	589	Aqua kali puri, ii.	410
Aqua caponis, i.	700	Aqua cupri vitriolati composita, i.	586	Aqua kali sulphurati, ii.	418
Aqua carbonica, i.	49	Aqua cydoniorum, i.	599	Aqua kalino-ætherea, ii.	423
Aqua carbonatis acidulus calcis, i.	358	Aqua diatrion, i.	74	Aqua lauro-cerasi, i.	62
Aqua carbonatis ammoniæ, i.	483	Aqua divina Fernelii, ii.	68	Aqua lauro-cerasi cum belladonna, i.	311
Aqua carbonatis potassæ acidula, ii.	426	Aqua divina mirabilis, i.	518	Aqua laxativa, ii.	219
Aqua carbonatis sodæ acidula, ii.	607	Aqua diuretica ex nucleis, i.	64	Aqua laxativa Viennensis, i.	415
Aqua carbunculi, ii.	170	Aqua ebulli, ii.	551	Aqua liliorum convallium, i.	542
Aqua cari carui, i.	398	Aqua embryonum, ii.	549	Aqua lithargyri acetatis, ii.	375
Aqua carmelitana, ii.	230	Aqua emetica, i.	259	Aqua lithargyri composita, ii.	278
Aqua carminativa, i.	559			Aqua lithontripica, ii.	427
Aqua carui, i.	398			Aqua Lucæ, i.	180
Aqua carui seminis, i.	398			Aqua lupuli, ii.	11
				Aqua magnanimitatis, i.	676

Aqua magnanimitatis, simplex, i.	676	Aqua picis liquidæ, ii.	365	Aqua sulphureo-acida, i.	65
Aqua magnesiæ, ii.	209	Aqua pimentæ, ii.	361	Aqua sulphureti ammoniæ, i.	194
Aqua marina, ii.	603	Aqua pimentæ vinosa, ii.	362	Aqua supercarbonatis kali, ii.	427
Aqua mastichina, ii.	223	Aqua pini turionum, i.	21	Aqua supercarbonatis potassæ, ii.	426
Aqua melissæ composita, ii.	230	Aqua plumbi acetosi, ii.	378	Aqua supercarbonatis sodæ, ii.	607
Aqua melissæ spirituosacomposita, ii.	230	Aqua pontificalis, i.	37	Aqua ad synoviam, ii.	86
Aqua mellis odorata regia, i.	559	Aqua potassæ, ii.	410	Aqua tanacetii, ii.	651
Aqua menthæ alcoholica, ii.	239	Aqua Pragensis, i.	305	Aqua Thediana, i.	45
Aqua menthæ composita, ii.	240	Aqua prophylactica, i.	225	Aqua theriacalis frigida, ii.	314
Aqua menthæ spirituosacomposita, ii.	239	Aqua pruni avium alcoholica, i.	63	Aqua theriacalis Romana, ii.	314
Aqua menthæ vinosa, ii.	239	Aqua pruni lauro-cerasi, i.	62	Aqua theriacalis simplex, ii.	314
Aqua menthæ viridis, ii.	239	Aqua pruni padi, i.	63	Aqua thymi, ii.	690
Aqua mercurialis, ii. 76, 92	78, 92	Aqua pulegii, ii.	239	Aqua tilis, ii.	692
Aqua mercurialis composita, ii.	90	Aqua Rabelii, i.	79	Aqua tormentillæ, ii.	693
Aqua mercurialis nigra, ii.	34	Aqua regia, i.	56	Aqua ad tormina infantum, ii.	240
Aqua mercurii vitriolati, ii.	74	Aqua regis, i.	56	Aqua traumatica, ii.	547
Aqua mercurii vivicocta, ii.	13	Aqua rosæ, ii.	517	Aqua traumatica Thedenii, i.	45
Aqua mineralis cathartica, ii.	211	Aqua rosæ centifoliæ, ii.	517	Aqua de tribus, i.	74
Aqua mineralis ferrata, i.	651	Aqua rutæ, ii.	538	Aqua valerianæ, ii.	711
Aqua miraculosa, i.	586	Aqua rosarum, ii.	517	Aqua valerianæ composita, ii.	711
Aqua muriatico-mercurialis rubra, ii.	35	Aqua rubi idæi, ii.	533	Aqua vanillæ, ii.	712
Aqua muriatis barytæ, i.	326	Aqua cum saccharo saturni, ii.	386	Aqua vegetabilis-mineralis Goulardi, ii.	78
Aqua myrti pimentæ, ii.	361	Aqua salviæ, ii.	546	Aqua veronicæ, ii.	718
Aqua naphæ, i.	508	Aqua salviæ spirituosacomposita, ii.	547	Aqua veronicæ vinosa, ii.	718
Aqua odorifera, ii.	159	Aqua sapphirina, i.	596	Aqua violacea, ii.	133
Aqua opthalmica, ii.	377	Aqua saturni, ii.	378	Aqua violarum, ii.	721
Aqua opthalmica adstringens, i.	162	Aqua saturnina, ii. 377, 386	378, 386	Aqua viridis, i.	596
Aqua opthalmica aluminosa, i.	162	Aqua scabiosæ, ii.	559	Aqua vitæ, i.	104
Aqua opthalmica cærulea, i.	590, 596	Aqua sclopetaria, ii.	547	Aqua vitæ mulierum Bœcleri, ii.	169
Aqua opthalmica Contradi, ii.	76	Aqua sclopetaria aromatica, ii.	547	Aqua vitæ rectificata, i.	107
Aqua opthalmica mercurialis, ii.	69	Aqua sclopetaria Thedenii, i.	45	Aqua vitæ saponacea, ii.	461
Aqua opii, ii.	275	Aqua scordii, ii.	683	Aqua vitrioli cærulea, i.	590
Aqua oxygenata, i.	459	Aqua scordii composita, ii.	314	Aqua vitriolica cærulea, i.	586
Aqua oxygeno-muriatica, i.	459	Aqua serpylli, ii.	690	Aqua vulneraria, ii. 547, 682	
Aqua papaveris erratici, ii.	334	Aqua serpylli composita, ii.	691	Aqua vulneraria acida, i.	45
Aqua paralyseos, ii.	466	Aqua sine pari, i.	518	Aqua vulneraria Romana, i.	37
Aqua perlata, i.	63	Aqua sodæ, ii.	608	Aqua vulneraria spirituosacomposita, ii.	546
Aqua pestilentialis, i.	373	Aqua splenetica, i.	429	Aqua vulneraria Thedenii, i.	45
Aqua phagedenica, ii.	68	Aqua Stettensis ad usus externos, ii.	78	Aquifolii folia, ii.	113
Aqua picea, ii.	365	Aqua stibii, i.	258	Aquila alba, ii.	46
Aqua picis, ii.	365	Aqua stibiata, i.	259	Aquila mitigata, ii.	46
		Aqua stomachica, ii.	240	Aquilaria ovata, ii.	193
		Aqua styptica, i.	586	Aquilegia vulgaris, i.	270
		Aqua styptica Clementina, ii.	328, 682	Araci aromatici siliquæ, ii.	712
		Aqua succorum vegetabilium vulneraria, i.	37	Aralia naked-stalked, i.	270
		Aqua succo acetosellæ, ii.	327	Aralia nudicaulis, i.	270
		Aqua sulphatis aluminæ, i.	162	Aralia spinosa, i.	270
		Aqua sulphatis magnesiæ carbonica, ii.	212		

<i>Aranea domestica</i> , i.	271	<i>Arsenias ammoniæ</i> , i.	288	<i>Asclepias flesh-coloured</i> , i.	295
<i>Arbor vitæ American</i> , ii.	689	<i>Arsenias kali</i> , i.	288	<i>Asclepias incarnata</i> , i.	295
<i>Arbutus uva ursi</i> , i.	271	<i>Arsenias potassæ</i> , i.	288	<i>Asclepias Syriaca</i> , i.	295
<i>Arcanum Archideti</i> , i.	351	<i>Arsenis potassæ aquosus</i> , i.	286	<i>Asclepias tuberosa</i> , i.	295
<i>Arcanum bechicum Willisii</i> , ii.	418	<i>Arsenis potassæ liquidus</i> , i.	286	<i>Asclepias vincetoxicum</i> , i.	295
<i>Arcanum corallinum</i> , ii.	36	<i>Arsenias sodæ</i> , i.	288	<i>Ash mountain</i> , ii.	619
<i>Arcanum cosmeticum</i> , ii.	213	<i>Arsenis sodæ</i> , i.	287	<i>Ash prickly</i> , i.	270
<i>Arcanum duplicatum Myn- sichti</i> , ii.	427	<i>Arseniate of ammonia</i> , i.	288	<i>Ash tree</i> , i.	678
<i>Arcanum tartari</i> , ii.	441	<i>Arseniate of soda</i> , i.	288	<i>Ashes of burnt animals</i> , i.	355
<i>Archangel</i> , ii.	154	<i>Arsenidum sodæ liquidum</i> , i.	287	<i>Ashes of the hare</i> , i.	355, 391
<i>Arctium lappa</i> , i.	272	<i>Arsenic</i> , i.	281	<i>Ashes of the hedgehog</i> , i.	355, 391
<i>Argel</i> , i.	401	<i>Arsenic white</i> , i.	283	<i>Ashes of the mole</i> , i.	355, 391
<i>Argentariæ radix</i> , ii.	465	<i>Arsenici oxydum album</i> sublimatum, i.	283	<i>Ashes of silk</i> , i.	391
<i>Argentinaæ radix</i> , ii.	465	<i>Arsenicum album</i> , i.	283	<i>Ashes of the swallow</i> , i.	355, 391
<i>Argentum</i> , i.	273	<i>Arsenicum album subli- matum</i> , i.	283	<i>Ashes of the wren</i> , i.	355, 391
<i>Argentum nitratum</i> , i.	273	<i>Arsenicum antimoniatum</i> , i.	284	<i>Asparagi hortensis radix</i> , i.	296
<i>Argentum nitricum</i> , i.	273	<i>Arsenicum crudum</i> , i.	283	<i>Asparagi sativi radix</i> , i.	296
<i>Argentum nitricum crys- tallisatum</i> , i.	273	<i>Arseniate of iron</i> , i.	288	<i>Asparagine</i> , i.	296
<i>Argentum nitricum fusum</i> , i.	273	<i>Arsenicum flavum</i> , i.	282	<i>Asparaginum</i> , i.	296
<i>Argentum vivum</i> , ii.	12	<i>Arsenicum rubrum</i> , i.	282	<i>Asparagus</i> , i.	296
<i>Argilla kali-sulphurica</i> , i.	158	<i>Arsenite of potass</i> , i.	286	<i>Asparagus officinalis</i> , i.	296
<i>Argilla pura</i> , i.	156	<i>Arsenite of soda</i> , i.	287	<i>Asphaltum</i> , i.	298
<i>Argilla sulphurica alkali- sata</i> , i.	158	<i>Artanitæ radix</i> , i.	597	<i>Asphaltus</i> , i.	298
<i>Argilla sulphurica usta</i> , i.	158	<i>Artemisia abrotanum</i> , i.	290	<i>Asphodel branchy</i> , i.	298
<i>Ari radix</i> , i.	291	<i>Artemisia campestris</i> , i.	290	<i>Asphodeli albi radix</i> , i.	298
<i>Arisari radix</i> , i.	291	<i>Artemisia Chinensis</i> , i.	289	<i>Asphodeli maris radix</i> , i.	298
<i>Aristolochia clematidis</i> , i.	276	<i>Artemisia contra</i> , ii.	581	<i>Asphodeli ramosi radix</i> , i.	298
<i>Aristolochia longa</i> , i.	276	<i>Artemisia dracunculus</i> , i.	291	<i>Asphodelus ramosus</i> , i.	298
<i>Aristolochia pistolochia</i> , i.	276	<i>Artemisia glacialis</i> , i.	291	<i>Aspidium coriaceum</i> , i.	298
<i>Aristolochia rotunda</i> , i.	276	<i>Artemisia Judaica</i> , ii.	581	<i>Asplenii herba</i> , i.	443
<i>Aristolochia serpentaria</i> , i.	277	<i>Artemisia Maderapatana</i> , i.	24	<i>Asplenium ruta muraria</i> , i.	299
<i>Aristolochia trilobata</i> , i.	277	<i>Artemisia maritima</i> , i.	24	<i>Assafoetida</i> , i.	299
<i>Aristolochia cavaæ radix</i> , i.	561	<i>Artemisia Pontica</i> , i.	24	<i>Asteris conyzoides herba</i> , i.	621
<i>Aristolochiæ creticæ radix</i> , i.	276	<i>Artemisia rupestris</i> , i.	291	<i>Asteris inqualis radix</i> , i.	622
<i>Aristolochiæ fabacæ ra- dix</i> , i.	561	<i>Artemisia vulgaris</i> , i.	289	<i>Astmury</i> , i.	317
<i>Aristolochiæ longæ veræ radix</i> , i.	276	<i>Artemisiæ herba</i> , i.	289	<i>Astragalus</i> , i.	766
<i>Aristolochiæ polyrhizæ radix</i> , i.	276	<i>Artemisiæ albæ herba</i> , i.	289	<i>Astragalus exscapus</i> , i.	307
<i>Aristolochiæ rotundæ veræ radix</i> , i.	277	<i>Artemisiæ rubræ herba</i> , i.	289	<i>Astragalus gummiiferus</i> , i.	766
<i>Aristolochiæ tenuis radix</i> , i.	276	<i>Artemisiæ santonicæ se- mina</i> , ii.	581	<i>Astragalus of the hare</i> , i.	359
<i>Aristolochiæ vulgaris ra- dix</i> , i.	276	<i>Artichoke</i> , i.	601	<i>Astragalus verus</i> , i.	766
<i>Aristolochiæ vulgaris ro- tundæ radix</i> , i.	561	<i>Arum arisarum</i> , i.	291	<i>Astrantia major</i> , i.	307
<i>Armoraciæ radix</i> , ii.	481	<i>Arum common</i> , i.	291	<i>Astrantiæ majoris radix</i> , i.	307
<i>Arnica montana</i> , i.	279	<i>Arum dracunculus</i> , i.	292	<i>Astrantiæ radix</i> , ii.	114
<i>Arnica mountain</i> , i.	279	<i>Arum hooded</i> , i.	291	<i>Athanasia herba</i> , ii.	649
<i>Arnica plauensis radix</i> , i.	279	<i>Arum maculatum</i> , i.	291	<i>Athyrium filix mas</i> , i.	308
<i>Arnica spuria herba</i> , ii.	114	<i>Arum three-leaved</i> , i.	292	<i>Atriplex angustifolia</i> , i.	309
<i>Arnica Suedensis herba</i> , ii.	114	<i>Arum triphyllum</i> , i.	292	<i>Atriplicis fetidæ herba</i> , i.	458
<i>Aroph Paracelsi</i> , i.	661, 663	<i>Arundo donax</i> , i.	293	<i>Atriplex glauca</i> , i.	309
<i>Arquebusade water</i> , i.	45	<i>Arundo phragmites</i> , i.	293	<i>Atriplex hortensis</i> , i.	309
<i>Arrack</i> , i.	104	<i>Arundo saccharifera</i> , ii.	539	<i>Atriplicis odoratæ herba</i> , i.	457
<i>Arrow-root</i> , i.	632	<i>Asa dulcis</i> , i.	330	<i>Atriplicis olidæ herba</i> , i.	458
		<i>Asarabacca</i> , i.	293	<i>Atriplicis sativæ herba</i> , i.	309
		<i>Asari radix</i> , i.	293	<i>Atropa belladonna</i> , i.	309
		<i>Asarum</i> , i.	293	<i>Atropa madragora</i> , i.	312
		<i>Asarum Canadense</i> , i.	293	<i>Atropia</i> , i.	309
		<i>Asarum Canadian</i> , i.	293	<i>Atropina</i> , i.	309
		<i>Asarum Europæum</i> , i.	293	<i>Atropine</i> , i.	309
		<i>Asbestos</i> , i.	295	<i>Attmellæ herba</i> , ii.	621
		<i>Asbestus</i> , i.	295	<i>Avellanæ nuces</i> , i.	561

<i>Avens</i> , i.	718	<i>Azote</i> , ii.	265	<i>Balsam of guaiac</i> , i.	740
<i>Avens water</i> , i.	719	<i>Azure stone</i> , i.	157	<i>Balsam Hungarian</i> , i.	318
<i>Avicula margaritifera</i> , i.	354			<i>Balsam of lavender</i> , ii.	157
<i>Aurantia Curassavica</i> , i.	506	<i>Baccæ Norlandicæ</i> , ii.	531	<i>Balsam of lead</i> , ii.	375
<i>Aurantia Hispanientia</i> , i.	506	<i>Baculi liquiritiæ albi</i> , i.	726	<i>Balsam Liboschitz</i> , i.	381
<i>Aurantii folia</i> , i.	506	<i>Baculi liquiritiæ citrini</i> , i.	726	<i>Balsam Locatelli's</i> , ii.	660
<i>Aurantiorum cortices</i> , i.	506	<i>Balaustia</i> , ii.	470	<i>Balsam of marjoram</i> , ii.	323
<i>Aurate of ammonia</i> , i.	314	<i>Balls bezoardic</i> , i.	277	<i>Balsam of Mecca</i> , i.	319
<i>Auricula Judæ</i> , ii.	695	<i>Balls disinfecting</i> , i.	460	<i>Balsam Metz's</i> , i.	594
<i>Auriculæ muris herba</i> , ii.	6	<i>Balls against erysipelas</i> , ii.	373	<i>Balsam of myrrh</i> , i.	761
<i>Auriculæ muris majoris herba</i> , ii.	6	<i>Balls Molsheim's</i> , i.	667	<i>Balsam narcotic</i> , ii.	669
<i>Auridum ammoniacæ</i> , i.	314	<i>Balls musk</i> , ii.	251	<i>Balsam nervine</i> , ii.	524
<i>Auripigmentum</i> , i.	282	<i>Balm American mountain</i> , ii.	244	<i>Balsam of nutmeg</i> , ii.	250
<i>Aurum</i> , i.	312	<i>Balm apple</i> , ii.	242	<i>Balsam ophthalmic</i> , ii.	40, 527
<i>Aurum fulminans</i> , i.	314	<i>Balm bastard</i> , ii.	232	<i>Balsam opodeldoc</i> , i.	179
<i>Aurum horizontale Zuringeri</i> , i.	314	<i>Balm common</i> , ii.	227	<i>Balsam pectoral</i> , ii.	661
<i>Aurum Mosaicum</i> , ii.	625	<i>Balm field</i> , ii.	227	<i>Balsam of Peru</i> , i.	319
<i>Aurum muriaticum</i> , i.	312	<i>Balm mountain</i> , ii.	227	<i>Balsam Peruvian</i> , i.	319
<i>Aurum oxydatum</i> , i.	313	<i>Balneum incitans</i> , i.	45	<i>Balsam Plenk's mercurial</i> , ii.	21
<i>Aurum potabile</i> , i.	313	<i>Balneum sulphuratum</i> , i. 66. ii.	418	<i>Balsam rakasira</i> , i.	322
<i>Axunge</i> , i.	316	<i>Balsam acoustic</i> , i. 113, 180, 321, 633, ii. 105		<i>Balsam of rue</i> , ii.	538
<i>Axungia</i> , i.	316	<i>Balsam æthereal</i> , ii.	638	<i>Balsam Saint-Yve's ophthalmic</i> , ii.	40
<i>Axungia anatis</i> , i.	316	<i>Balsam of amber</i> , ii. 463, 631		<i>Balsam Salazar's</i> , i.	129
<i>Axungia anguillæ</i> , i.	317	<i>Balsam anisated of sulphur</i> , ii.	637	<i>Balsam saturnine</i> , ii.	385
<i>Axungia anseris</i> , i.	316	<i>Balsam anodyne</i> , ii.	462	<i>Balsam Schauer's</i> , ii.	667
<i>Axungia ardæ</i> , i.	316	<i>Balsam anti-arthritis</i> , ii.	462	<i>Balsam somniferous</i> , ii.	286
<i>Axungia aschiæ</i> , i.	317	<i>Balsam anticancerous</i> , ii.	301	<i>Balsam soothing</i> , i.	154
<i>Axungia aschii</i> , i.	317	<i>Balsam antihysteria</i> , i. 123, 190		<i>Balsam simple of sulphur</i> , ii.	637
<i>Axungia canis</i> , i.	315	<i>Balsam anti-odontalgic</i> , ii.	300	<i>Balsam succinated of sulphur</i> , ii.	638
<i>Axungia caponis</i> , i.	316	<i>Balsam antipleuritic</i> , i.	579	<i>Balsam terebinthinated of sulphur</i> , ii.	638
<i>Axungia castoris</i> , i.	315	<i>Balsam Arceus'</i> , ii.	658	<i>Balsam of Tolu</i> , i.	322
<i>Axungia cati sylvestris</i> , i.	315	<i>Balsam aromatic</i> , i. 182, 401		<i>Balsam tree</i> , ii.	409
<i>Axungia ciconiæ</i> , i.	316	<i>Balsam aromatic æthereous</i> , i.	426	<i>Balsam saponaceous of turpentine</i> , ii.	671
<i>Axungia coturnicis</i> , i.	316	<i>Balsam astrigent</i> , ii.	670	<i>Balsam vulnerary</i> , ii.	659, 665
<i>Axungia cuniculi</i> , i.	316	<i>Balsam Basville's</i> , ii.	672	<i>Balsami palustris herba</i> , ii.	233
<i>Axungia equi e collo seu juba</i> , i.	315	<i>Balsam camphorated acetie</i> , ii.	462	<i>Balsamum ad loricandum cadaver conditum</i> , ii.	628
<i>Axungia erinacei</i> , i.	315	<i>Balsam Canadian</i> , i.	318	<i>Balsamum adstringens</i> , ii.	670
<i>Axungia gallinæ</i> , i.	316	<i>Balsam Carpathian</i> , i.	318	<i>Balsamum æthereum</i> , ii.	638
<i>Axungia hominis</i> , i.	315	<i>Balsamum Carpathicum nativum</i> , i.	318	<i>Balsamum anodynum</i> , ii.	286, 461
<i>Axungia leporis</i> , i.	316	<i>Balsam chalybeate</i> , i.	656	<i>Balsamum apoplecticum</i> , ii.	524
<i>Axungia lucii piscis</i> , i.	317	<i>Balsam against chilblains</i> , ii.	670	<i>Balsamum Appariti</i> , ii.	110
<i>Axungia lupi</i> , i.	316	<i>Balsam of cinnamon</i> , ii.	165	<i>Balsamum Arcei</i> , ii.	658
<i>Axungia muris alpini</i> , i.	316, 317	<i>Balsam Condom's</i> , ii.	252	<i>Balsamum aromaticum</i> , ii.	524
<i>Axungia nitrica</i> , i.	70	<i>Balsam of copalm</i> , ii.	627	<i>Balsamum aromaticum æthereum</i> , i.	426
<i>Axungia pedum tauri</i> , i.	315	<i>Balsam Dippel's vulnerary</i> , i.	670	<i>Balsamum Basvillianum</i> , ii.	672
<i>Axungia phoci</i> , i.	316	<i>Balsam East Indian</i> , ii.	319	<i>Balsamum e benzoe</i> , i.	331
<i>Axungia porci</i> , i.	316	<i>Balsam Fioraventi's</i> , ii.	665	<i>Balsamum Brasiliense</i> , i.	554
<i>Axungia porcina</i> , i.	316	<i>Balsam Fourcroy's</i> , ii.	675		
<i>Axungia serpentum</i> , i.	316	<i>Balsam Frahm's</i> , ii.	657		
<i>Axungia sui scrofæ</i> , i.	316	<i>Balsam of gold</i> , i.	314		
<i>Axungia suilla</i> , i.	316				
<i>Axungia taxi</i> , i.	315				
<i>Axungia thymalli piscis</i> , i.	317				
<i>Axungia viperarum</i> , i.	317				
<i>Axungia ursi</i> , i.	316				
<i>Axungia vulpis</i> , i.	316				
<i>Aya-pana</i> , i.	628				
<i>Azari radix</i> , i.	293				

Balsamum calaba, i.	319	Balsamum rakasira, i.	322	Barilla, ii.	604
Balsamum Canadense, i.	318	Balsamum rutæ, ii.	538	Barium, i.	325
Balsamum catholicum, i.	331	Balsamum saponaceum, ii.	461	Bark brown, i.	470
Balsamum catholicum cum spiritu vini, ii.	108	Balsamum saponaceum terebinthinatum, ii.	671	Bark cabbage-tree, i.	716
Balsamum catholicum ex vino, ii.	108	Balsamum saponis, i.	179	Bark Carolina, i.	469
Balsamum cephalicum Saxonicum, ii.	524	Balsamum saturni, ii.	375, 376, 385	Bark grey, i.	470
Balsamum chalybeatum, i.	656	Balsamum saturninum, ii.	380	Bark greyish-brown, i.	470
Balsamum Chiron, ii.	661	Balsamus Schauerianus, ii.	667	Bark Jamaica, i.	469
Balsamum cinnamomi, ii.	165	Balsamum Scherzei, ii.	524	Bark Jesuits, i.	469
Balsamum commenda-toris, i.	331	Balsamum stomachicum, ii.	524	Bark St. Lucia, i.	471
Balsamum copaibæ, i.	554	Balsamum succini, ii.	631	Bark Peruvian, i.	469
Balsamum ad dentitionem, i.	154	Balsamum sulphuris anisatum, ii.	637	Bark red, i.	476
Balsamum embryonum, ii.	549	Balsamum sulphuris simplex, ii.	637	Bark royal yellow, i.	476
Balsamum Fioraventi, ii.	665	Balsamum sulphuris succinatum, ii.	638	Bark Spanish crown, i.	476
Balsamum Frahmii, ii.	657	Balsamum sulphuris terebinthinatum, ii.	638	Bark worm, i.	716
Balsamum guaiacinum, i.	736	Balsamum toltanum, i.	322	Bark yellow, i.	470
Balsamum Hungaricum, i.	318	Balsamum terebinthinatum, ii.	659	Barley, ii.	7
Balsamum Indicum album, i.	320	Balsamum thebaicum, ii.	286	Barley meal prepared, ii.	7
Balsamum innocentianum, ii.	108	Balsamum tranquillum, ii.	104	Barley sugar, ii.	7
Balsamum ischæmaticum, ii.	670	Balsamum traumaticum, i.	331	Barley water, ii.	7
Balsamum Italicum rubrum, ii.	660	Balsamum viride, i.	319, 661	Barm, i.	634
Balsamum Lectorense, ii.	252	Balsamum vitæ album, ii.	667	Barotes salitus, i.	325
Balsamum Locatelli, ii.	660	Balsamum vitæ externum, ii.	671	Barras, ii.	673
Balsamum mariæ, i.	319	Balsamum vitæ Rulandi, ii.	638	Baryta, i.	327
Balsamum de Mecca, i.	319	Balsamum universale, ii.	380	Baryta carbonica, i.	328
Balsamum Meccanense, i.	319	Balsamum vulnerarium, i.	331	Baryta fused, i.	328
Balsamum mercuriale, ii.	87	Balsamum vulnerarium minerale Dippelii, i.	670	Baryta muriatica, i.	325
Balsamum mercuriale Plenckii, ii.	21	Balsamum vulnerarium vegetabile, i.	37	Barytes, i.	325
Balsamum nativum pini sylvestris, ii.	654	Balsamus Ægyptiacus, i.	319	Basilica majoris herba, ii.	269
Balsamum nervinum, ii.	524	Balsamus Gileadensis, i.	319	Basilici minimi herba, ii.	269
Balsamum nucistæ, ii.	256	Balsamus Judaicus, i.	319	Basilicon black, ii.	678
Balsamum nucistæ expressum, ii.	256	Balsamus myrepsica, ii.	245	Basilicon yellow, ii.	677
Balsamita odorata, i.	317	Balsamus Orientalis, i.	319	Bath antisypilitic, ii.	69
Balsamum oleæ, i.	764	Balsamus Syriacus, i.	319	Bath aromatic, ii.	523
Balsamum opthalmicum rubrum, ii.	38	Balsamus verus, i.	319	Bath Dupuytren's sulphureous and gelatinous, ii.	418
Balsamum opthalmicum anum, ii.	40	Balzoinum, i.	330	Bath emollient, ii.	201
Balsamum Persicum, i.	331	Baneberry, i.	89	Bath resolvent, i.	192, 418
Balsamum Peruvianum, i.	319	Banillæ siliquæ, ii.	712	Bath sulphureous, i.	66, 418, 614
Balsamum Peruvianum album, i.	320	Bapti seculæ flores, i.	597	Batrachites, i.	359
		Barbæ aronis radix, i.	291	Baume de saturne de Goulard, ii.	381
		Barbæ caprinæ radix, ii.	621	Baume de vie Lelievre's, i.	140
		Barbæ hirci radix, ii.	695	Bawdmoney, ii.	197
		Bearberry, i.	355	Bay tree, ii.	178
		Bardanæ radix, i.	272	Bdellium i.	329
				Bead tree, ii.	225
				Bean French, ii.	346
				Bean garden, i.	631
				Bean picurim, ii.	680
				Bearberry bush, i.	271
				Bearsbreach, i.	34
				Bearsfoot, i.	34
				Beaumont root, i.	720
				Beccabunga, ii.	719
				Becetta, ii.	694
				Bechii radix, i.	699
				Bedeguar, i.	681
				Bedstraw ladies', i.	691

Bedstraw great ladies', i.	694	Bezoar German, i.	338	Bole white, i.	155
Bedstraw yellowladies', i.	694	Bezoar Occidental, i.	338	Boletus cervinus, ii.	577
Beech drops root, ii.	324	Bezoar Oriental, i.	338	Boletus ignarius, i.	340
Beech tree, i.	631	Bezoar of the porcupine,		Boletus salicis, i.	602
Beer, i.	440	i.	338	Boletus touchwood, i.	340
Beer alterative, ii.	590	Bezoardicum animale, i.	697	Boli anthelmintico-lax-	
Beer anti-hydropic, i.	551	Bezoardicum Curvi, ii.	578	antes, ii.	65
Beer anti-icteric, i.	409	Bezoardicum joviale, ii.	624	Boli anthelmintico-to-	
Beer anti-scorbutic, i.	534,	Bezoardicum martiale, i.	236	nici, ii.	2
ii.	483, 580	Bicarbonas ammoniæ, i.	182	Boli anti-chlorotici em-	
Beer anti-spasmodic, ii.	707	Bicarbonas sodæ, ii.	607	menagogi incitantes, i.	642
Beer aromatic, ii.	171	Bicarbonate of ammonia,		Boli anti-chlorotici em-	
Beer astrigent, ii.	477	i.	182	menagogi tonici, i.	637
Beer bitter, i.	23	Bicarbonate of soda, ii.	607	Boli antifebriles, i. 504, ii.	497
Beer cephalic, ii.	708	Bichloride of mercury, ii.	66	Boli cærulei, ii.	17
Beer chalybeate, i.	665	Bichloruret of mercury, ii.	66	Boli contrastimulantes,	
Beer of cinchona, i.	496	Bichloruretum calcis, i.	351	i.	61
Beer diuretic, ii.	590	Bidens cornua, i.	339	Boli in cystitide lenta,	
Beer prophylactic, i.	496	Bidens tripartitum, ii.	621	ii.	635
Beer purgative, i. 409, 551, 575		Bidentis herba, i.	339	Boli diaphoretici nause-	
Beer spruce, i.	23	Bilberry red, ii.	702	antes, i.	248
Beer compound spruce,		Bindweed sea, i.	366	Boli drastici, i.	695
ii.	483	Binoxalas potassæ, ii.	446	Boli emmenagogi irri-	
Beer stimulant, ii.	590	Birch, i.	338	tantes, i.	122, 135
Beer stomachic, i.	713	Bird cherry, i.	437	Boli emmenagogi resol-	
Beet common, i.	336	Birthwort round, i.	276	ventes, ii.	121
Beet white, i.	336	Biscuits purgative, i.	545	Boli ad epilepticos, ii.	705
Beetle oil, ii.	232	Biscuits purgative and ver-		Boli excitantes, i.	367
Behen albi radix, i.	433	mifuge, ii.	62	Boli e ferro compositi,	
Behen nostratis radix, ii.	587	Bislinguæ herba, ii.	536	i.	642
Behen radix, i.	433	Bismalvæ radix, i.	143	Boli incitantes necnon	
Behen rubri radix, ii.	625	Bismuthi subnitras, i.	339	diaphoretici et ex-	
Behn, ii.	244	Bismuthum nitricum, i.	339	pectorantes, i.	247
Belladonna, i.	309	Bismuthum oxydulatum		Boli incitantes emmena-	
Belliculus marinus, i.	353	album, i.	339	gogi, i.	642, ii. 479
Bellidis hortensis herba, i.	329	Bistort, ii.	406	Boli incitantes stoma-	
Bellidis majoris herba, i.	461	Bistortæ radix, ii.	406	chici, i.	642, 674
Bellidis minoris herba, i.	329	Bisulphate of copper, i.	583	Boli kermético-ammoni-	
Bellidis pratensis herba,		Bitartras potassæ, ii.	449	acales, i.	247
i.	461	Bitter apple, i.	571	Boli lithontriptici, ii.	458
Bellis perennis, i.	329	Bitter-sweet, ii.	615	Boli martiales emme-	
Ben, ii.	244	Bitumen Judaicum, i.	298	nagogi, i.	637, 642
Ben nut, ii.	245	Bitumen petroleum, ii.	345	Boli martis, i.	667
Benedictæ sylvestris herba,		Bixa orellana, ii.	352	Boli e narcotina, ii.	259
i.	719	Blackberry, ii.	532	Boli in necrosi syphili-	
Benzoë, i.	330	Black drop, ii.	275	tica, ii.	32
Benzoës flores, i.	46	Black lead, i.	639	Boli nervini, i.	302
Benzoin, i.	330	Bladder senna, i.	401, 540	Boli ad paralysim, ii.	588
Benzoinum, i.	330	Bladder wrack, i.	678	Boli in pertussi, ii.	51
Berberes, i.	335	Blancmanger, i.	704	Boli in ptyalismo mer-	
Berberides baccae, i.	335	Blatta byzantina, i.	352	curiali, ii.	419
Berberis vulgaris, i.	335	Blessed thistle, i.	524	Boli purgantes irritan-	
Berry partridge, i.	696	Blood of the goat, i.	699	tes, i.	695
Beta cycla, i.	336	Bloodroot, ii.	555	Boli in quartanam, i.	475
Beta vulgaris, i.	336	Blood-stone, i.	643	Boli resolventes, ii.	33
Betæ rubræ radix, i.	336	Blueberry, ii.	701	Boli resolventes et de-	
Betel, ii.	360	Blue-bottle, i.	597	primentes, i.	613
Betonica officinalis, i.	336	Blue-flag, ii.	131	Boli resolventes et anti-	
Betonica aquaticæ herba,		Blue Prussian, i.	674	syphilitici, ii.	33
ii.	578	Beticeæ sylvestris folia,		Boli resolventes despu-	
Betony wood, i.	336	ii.	258	mentes, ii.	121
Betula alba, i.	338	Bole Armenian, i.	155	Boli resolventes diapho-	
Betula alni cortex, i.	116	Bole artificial red, i.	156	retici, i.	247
Bezoar, i.	338	Bole dentifrice, i.	156	Boli resolventes et in-	
Bezoar Coromandel, i.	338	Bole Lemnian, i.	155	citantes, ii.	55

- Boli resolventes et pur-
 gantes, ii. 491
 Boli e rheo, ii. 491
 Boli ex rore marino, i. 637
 Boli saponacei, ii. 458
 Boli e spongia, i. 392
 Boli stomachici, i. 605
 Boli contra strumas, i. 392
 Boli tonici adstrin-
 gentes balsamici, i. 710
 Boli tonico-antiseptici, i. 302
 Boli e valeriana, ii. 705
 Bolus alba, i. 155
 Bolus anthelmintic, ii. 582
 Bolus anti-arthritis, i. 669
 Bolus anti-catarrah, ii. 246
 Bolus anti-diarrhoeic, ii. 292
 Bolus anti-lacteal, ii. 443
 Bolus anti-monial, i. 242
 Bolus anti-spasmodic, i. 340
 ii. 300
 Bolus Armenia, i. 155
 Bolus aromatic, ii. 255
 Bolus aromatic of cin-
 chona, i. 475
 Bolus astringent, ii. 491
 Bolus Chaussier's lax-
 ative, ii. 62
 Bolus of cinchona, i. 475
 Bolus opiate of cincho-
 na, i. 475
 Bolus opiate and aro-
 matic of cinchona, i. 475
 Bolus contrastimulant,
 i. 59, 61
 Bolus ad diarrhoeam, ii. 491
 Bolus ferruginous, i. 663
 Bolus ex hydrargyro,
 ii. 85
 Bolus ex hydrargyro
 oxydato rubro, ii. 38
 Bolus mercurial, ii. 85
 Bolus nervine, i. 425
 Bolus nervinus, i. 425
 Bolus of nitric oxide of
 mercury, ii. 38
 Bolus Orientalis, i. 155
 Bolus pectoral, i. 768
 Bolus of protochloruret
 of mercury, ii. 50
 Bolus purgative, i. 403
 Bolus e rheo sedativus,
 ii. 491
 Bolus rubra, i. 155
 Bolus ex sabina, ii. 50
 Bolus e stanno, ii. 623
 Bolus stimulant, i. 230
 Bolus stimulant and to-
 nic, i. 279
 Bolus of sulphur, ii. 635
 Bolus compound of sul-
 phur, ii. 635
 Bolus of tartrate of
 potash and of iron, i. 674
 Bolus tonic, i. 483
- Bolus tonic and anti-
 periodic, i. 476
 Bolus tonic and anti-
 spasmodic, i. 368
 Bolus vermifuge, ii. 66,
 623
 Boluses alterative, i. 733
 Boluses anthelmintic
 and anti-spasmodic, ii. 56
 Boluses anthelmintic
 purgative, ii. 65
 Boluses anti-asthmatic,
 ii. 635
 Boluses anti-dysenteric,
 ii. 125
 Boluses anti-epileptic,
 ii. 705
 Boluses anti-gonorrhœic,
 i. 558, ii. 663
 Boluses anti-hæmorrhoi-
 dal, ii. 635
 Boluses anti-scorbutic, i. 483
 Boluses anti-scorfulous,
 i. 392
 Boluses anti-spasmodic,
 i. 301, 367, 649, ii. 249
 Boluses anti-syphilitic, i. 313
 ii. 32
 Boluses astringent, i. 159,
 645, ii. 125
 Boluses bitter, i. 711
 Boluses camphorated, i. 367
 Boluses camphorated
 and nitrated, i. 367
 Boluses carminative, ii. 353
 Boluses chalybeate, i. 649
 Boluses of cinchona and
 rhubarb, i. 475
 Boluses for cough, ii. 438
 Boluses against cystitis,
 ii. 635
 Boluses diaphoretic, i. 248
 ii. 45, 635
 Boluses digestive, ii. 125
 Boluses of digitalis, i. 613
 Boluses diuretic and anti-
 spasmodic, i. 614
 Boluses drastic, ii. 52
 Boluses emetic, ii. 83, 125
 Boluses emmenagogue,
 i. 122, 135, 363, 637, 642
 ii. 50, 121, 142, 217, 479
 Boluses expectorant, i. 247
 Boluses febrifuge, i. 483,
 475, 504, 653, ii. 362, 479
 Boluses ferruginous, i. 642
 Boluses glycyrrhized
 mercurial, ii. 17
 Boluses against hoop-
 ing-cough, ii. 51
 Boluses lithontriptic, ii. 458
 Boluses mercurial, ii. 17, 18
 Boluses of myrrh, i. 760
 Boluses narcotic, ii. 101
 Boluses of narcotine, ii. 259
- Boluses nervine, i. 302
 Boluses nervine anthel-
 mintic, ii. 50
 Boluses purgative, i. 544,
 695, ii. 58, 62
 Boluses purgative and
 tonic, ii. 57
 Boluses for quartan
 fever, i. 475
 Boluses resolvent, i. 140, 247
 613, ii. 121, 491
 Boluses of rhubarb, ii. 491
 Boluses against saliva-
 tion, ii. 419
 Boluses saponaceous, ii. 458
 Boluses sedative, i. 367
 ii. 246, 247
 Boluses of squill, ii. 564
 Boluses stimulant, i. 367
 ii. 588
 Boluses stimulant resol-
 vent, ii. 55
 Boluses stomachic, i. 140,
 565, 637, 642, 674, ii. 209,
 605
 Boluses stomachic anti-
 syphilitic, ii. 33
 Boluses sudorific, ii. 299
 Boluses tonic, i. 475, 710
 Boluses tonic anthel-
 mintic, ii. 50
 Boluses tonic and anti-
 septic, i. 302
 Boluses tonic and dia-
 phoretic, ii. 368
 Boluses vermifuge, i. 308
 ii. 2, 64
 Bombacis semina, i. 730
 Bombax, i. 730
 Bombix mori, ii. 585
 Bone cuttle-fish, i. 354
 Bone of hart's heart, i. 360
 Bones, i. 359
 Boni Henrici herba, i. 457
 Bonifaciæ herba, ii. 536
 Bonplandia trifoliata, i. 228
 Borage, i. 341
 Boras hydrargyri, ii. 81
 Boras sodæ alkalescens,
 ii. 608
 Borassus flabelliformis,
 i. 329
 Borate of mercury, ii. 81
 Borate of soda, ii. 608
 Borax, ii. 608
 Borax Venetus, ii. 608
 Borrago officinalis, i. 341
 Borussias ferri, i. 674
 Borussias hydrargyri, ii. 80
 Botrychium lunaria, i. 342
 Botryos herba, i. 457
 Botryos Mexicanæ herba,
 i. 456
 Bougies of caustic po-
 tass, ii. 414

Bougies Falck's, ii.	42	Brunellæ herba, ii.	465	Cahuchu, i.	755
Bougies mercurial, ii. 23,	64,	Brusci radix, ii.	536	Calaguala, i.	298
	-88	Bryonia alba, i.	344	Calami radix, i.	-86
Bougies saturnine, ii.	379,	Bryony, i.	344	Calaminthæ herba, ii.	227
	382, 390	Bryony black, ii.	649	Calaminthæ humilioris	
Bougies simple, i.	437	Bubon galbanum, i.	683	herba, i.	721
Bougies soluble mercu-		Bubon Macedonicum, i.	345	Calaminthæ montanæ	
rial, ii.	77	Buckthorn, ii.	487	herba, ii.	227
Bovista lupi, ii.	204	Buckwheat, ii.	407	Calaminthæ pulegii odore	
Box tree, i.	346	Bufones exsiccati, i.	698	herba, ii.	227
Box-wood tree, i.	560	Bufo nites, i.	359	Calamus Asiaticus, i.	86
Brake, ii.	469	Bugle, i.	103	Calamus odoratus, i.	86
Bramble common, ii.	531	Bugloss, i.	215	Calamus rotang, i.	765
Bramble dwarf, ii.	531	Bugloss viper's, i.	620	Calcareus lactiformis, i.	352
Bramble mountain, ii.	531	Buglossi latifolii herba, i.	341	Calcaria carbonica, i.	352
Bran, i.	681	Buglossi radix, i.	215	Calcaria muriatica, i.	362
Brançæ ursinæ herba, i.	34	Bugulæ herba, i.	103	Calcaria phosphorica, i.	359
Brançæ ursinæ radix, ii.	5	Bulbulus thrasus, i.	601	Calcaria pura, i.	347
Brandy, i.	104	Bupthalmi herba, i.	446	Calcaria pura liquida, i.	347
Brandy purgative, i.	553	Buplevrum rotundifolium,		Calcaria sulphurata, i.	350
Brassica eruca, i.	342	i.	345	Calcaria sulphurato-sti-	
Brassica napus, i.	342	Burdock, i.	272	biata, i.	253
Brassica oleracea, i.	343	Burnet great, ii.	555	Calcaria sulphurica, i.	361
Brassicæ capitata albæ		Burnet saxifrage, ii.	356	Calcis murias, i.	362
et rubræ folia, i.	343	Burnet small, ii.	465	Calcis phosphas præcipita-	
Brassicæ marinæ herba, i.	366	Burr, i.	272	tum, i.	360
Brassicæ rubræ semen, i.	343	Bursæ pastoris herba, ii.	688	Calcitrapa stellata, i.	346
Bread, ii.	332	Bursera gummiifera, i.	346	Calcitrippæ herba, i.	608
Brodium rosarum, ii.	512	Butcher's broom, ii.	535	Calc, i.	689
Brontias, i.	359	Butonius lapis, i.	359	Calendulæ alpinæ radix, i.	279
Broom dyer's, i.	708	Butter, ii.	149	Calendula arvensis, i.	365
Broom petty, i.	708	Butter of antimony, i.	254	Calendula officinalis, i.	365
Broom-rapè Virginian, ii.	324	Butter of marjoram, ii.	323	Calendulæ herba, i.	365
Brooklime, ii.	719	Butter of wax, i.	435	Calendulæ mineralis, i.	661
Brooklime long-leaved, ii.	719	Butterbur, ii.	700	Calmontanæ herba, ii.	227
Broth alterative, i. 273, ii. 118,		Buttercup, ii.	480	Calomel, ii.	46
	530, 613	Butterfly-weed, i.	295	Calomelas, ii.	46
Broth anti-epileptic, ii.	329	Butter-nut tree, ii.	134	Calophyllum calaba, i.	319
Broth anti-herpetic, i.	677	Butuæ radix, i.	505	Calophyllum inophyllum,	
Broth anti-spasmodic, ii.	329	Butyrum, ii.	149	ii.	644
Broth aperitive, i.	297	Butyrum amygdalarum		Calthæ radix, i.	279
Broth astringent, ii.	642	dulcium, i.	206	Calthæ sativæ herba, i.	365
Broth of bones, i.	699	Butyrum antimonii, i.	254	Caltrops water, ii.	695
Broth of calf's lungs, i.	699	Butyrum cacao, ii.	687	Caluna vulgaris, i.	366
Broth of chicken, i.	700	Butyrum ceræ, i.	435	Calumba American, i.	678
Broth of crabs, i.	700	Butyrum marjoranæ, ii.	323	Calumba Marietta, i.	678
Broth demulcent, i.	703	Butyrum rutæ, ii.	538	Calumba tree, i.	527
Broth diuretic, i. 297, ii. 576		Butyrum saturni, ii.	380	Calx acetata, i.	363
Broth emollient and anti-		Butyrum stanni, ii.	625	Calx antimonii, i.	235
sceptic, ii.	460	Butyrum stibii, i.	254	Calx antimonii elota, i.	236
Broth of frogs, i.	699	Buxus semper virens, i.	346	Calx antimonii Hoffmanni,	
Broth lienteric, i.	703			i.	253
Broth of lizards, i.	699	Cabbage, i.	343	Calx antimonii nitrata, i.	236,
Broth pectoral, i. 342, 703,		Cabbage wild, i.	342		240
706, ii.	192, 338	Cabbage skunk, i.	619	Calx antimonii sulphurata,	
Broth of snails, i.	700	Cabbage swamp, i.	619	i.	253
Broth of tortoise, i.	699	Cabbagi cortex, i.	717	Calx antimonii cum sul-	
Broth of veal, i.	699	Cacao præparata, ii.	686	phure, i.	253
Broth of viper, i.	700	Cacao tabulata, ii.	686	Calx arsenici alba, i.	283
Broth of woodlice, ii.	272	Cachibou, i.	346	Calx citrata, i.	364
Brucea ferruginea, i.	228	Cactus opuntia, i.	346	Calx cinerea mercurii, ii.	84
Brucium, i.	228	Cæruleum Berolinense, i.	674	Calx cupri viridis, i.	582
Brucia, i.	228	Cæruleum Borussicum, i.	674	Calx extincta, i.	347
Brucina, i.	343	Cæruleum montanum, i.	582	Calx hydrargyri alba, ii.	89
Brucium, i.	343	Cæsalphinia echinata, ii.	193	Calx hydrargyri rubra, ii.	36

Calx mercurii, ii.	85	Caranna, i.	754	Carbonas protoxidi po-	
Calx mercurii vitriolata,		Caraway, i.	397	tassii et ammoniæ, ii.	427
ii.	82	Carbo animalis, i.	390	Carbonas sodæ, ii.	604, 607
Calx salita, i.	362	Carbo pilæ marinæ, i.	390	Carbonas sodæ alkali-	
Calx stibii cum sulphure,		Carbo e vegetabilibus,		num, ii.	604
i.	253	i.	389	Carbonas sodæ impu-	
Calx sulphurica, i.	361	Carbonas ammoniacæ,		rum, ii.	604
Calx vismuthi, i.	339	i.	182	Carbonas sodæ incom-	
Calx viva, i.	347	Carbonas ammoniaci		pletus, ii.	604
Calx usta, i.	347	cornu cervi, i.	184	Carbonas super-ammoni-	
Calyculæ glandium quercus,		Carbonas ammoniaci		acus aquosus, i.	183
ii.	474	lumbricorum crystal-		Carbonas super-ammoni-	
Calystegia soldanella, i.	366	lizatus, i.	184	acus pyro-oleosus, i.	184
Camari herba, i.	83	Carbonas ammoniaci		Carbonas super-ammoni-	
Cambogium, i.	694	viperarum, i.	184	acus pyro-oleoso-aquo-	
Camels hay, i.	216	Carbonas ammoniæ, i.	182	sus, i.	185
Camphor, i.	366	Carbonas ammoniæ al-		Carbonas super-potassi-	
Camphor artificial, ii.	670	kalinus, i.	182	cus, ii.	419
Camphorosma, i.	375	Carbonas ammoniæ al-		Carbonate of magnesia,	
Camphorosma monspe-		kalinus solutus, i.	183	ii.	206
liaca, i.	375	Carbonas ammoniæ crys-		Carbonate of mercury,	
Campion field, ii.	587	tallizatus, i.	182	ii.	86
Cancanum, i.	741	Carbonas ammoniæ in-		Carbonate native of	
Cancer pagurus, i.	354	completus, i.	182	lime, i.	352
Cancer root, ii.	324	Carbonas ammoniæ li-		Carbonate of protoxide	
Canceri herba, ii.	688	quidus, i.	183	of sodium, ii.	607
Canceris herba, ii.	2	Carbonas ammoniæ py-		Carbonate of soda, ii.	607
Candelæ cereæ, ii.	390	ro-animalis, i.	184	Carbunculus, i.	155
Candelæ chirurgicæ, ii.	379	Carbonas ammoniæ py-		Carburet of iron, i.	639
Candelæ fumales, i.	334	ro-animale liquidum,		Carburet of sulphur, ii.	633
Candelæ mercuriales, ii.	64	i.	185	Cardami majoris herba,	
Candelæ mitigantes, ii.	379	Carbonas ammoniæ so-		ii.	698
Candelæ resolventes, ii.	379	lidus, i.	182	Cardamoms, i.	394
Canella alba, i.	376	Carbonas barytæ, i.	328	Cardamomi Javanici	
Canelline, i.	376	Carbonas baryticus, i.	328	fructus, i.	394
Canis vulpes, i.	697	Carbonas calcareus, i.	352	Cardamomi longi fruc-	
Cannabinæ aquaticæ		Carbonas calcicum bitu-		tus, i.	394
radix, i.	628	minatum, i.	352	Cardamomi majoris fruc-	
Cannella Malabarica, ii.	178	Carbonas calcis, i.	352	tus, i.	394
Cannabis sativa, i.	376	Carbonas calcis nati-		Cardamomi Malabarici	
Cannella Zeylanica, ii.	161	vum pulverulatum,		fructus, i.	394
Cantharides, i.	377	i.	352	Cardamomi medii fruc-	
Cantharidine, i.	377	Carbonas calcis præci-		tus, i.	394
Cantharis spotted, i.	377	pitatum, i.	355	Cardamomi minoris fruc-	
Cantharis vesicatoria, i.	377	Carbonas ferri oxydati,		tus, i.	394
Cantharis vittata, i.	377	i.	646	Cardamomi rotundi fruc-	
Caper spurge, i.	629	Carbonas ferri et potas-		tus, i.	394
Caper tree, i.	388	sæ liquidus, i.	647	Cardamomum longum,	
Caphopierite, ii.	489	Carbonas ferri præcipi-		i.	394
Capilli veneris herba, i.	89	tatus, i.	646	Cardamomum majus, i.	394
Capituli martis radix,		Carbonas lixivie alka-		Cardamomum medium,	
i.	622	linus solutus, ii.	422	i.	394
Capparidis flores condi-		Carbonas magnesiæ, ii.	206	Cardamomum minus, i.	394
tæ, i.	389	Carbonas natriicum, ii.	604,	Cardamomum rotundum,	
Capparidis gemmæ con-			607	i.	394
ditæ, i.	389	Carbonas plumbi, ii.	371	Cardiacæ herba, ii.	185
Capparis spinosa, i.	388	Carbonas potassæ, ii.	419	Cardinal-flower blue, ii.	203
Capra ægagrus, i.	338	Carbonas potassæ alka-		Cardopatæ radix, ii.	396
Caprifolii stipites, ii.	203	lescens liquidus, ii.	422	Cardopatii radix, ii.	396
Capsicum, ii.	359	Carbonas potassæ et		Cardui benedicti herba,	
Capsicum annuum, ii.	359	ammoniæ, ii.	427	i.	524
Capsicum frutescens, ii.	359	Carbonas potassæ puris-		Cardui laceti herba, i.	395
Caput mortuum vitrioli,		simus, ii.	420	Cardui mariæ herba, i.	395
i.	644	Carbonas protoxidi po-			
Capsinium, ii.	359	tassii, ii.	426		

Cardui sancti herba, i.	524	Cassia lignea, ii.	178	Cataplasma ad singul-	
Cardui stellati radix, i.	346	Cassia cum manna, ii.	219	tum, ii.	315
Cardui tomentosi herba,		Cassia Marylandica, i.	420	Cataplasma Althææ, i.	
ii.	273	Cassia obovata, i.	401		151
Cardui veneris herba, i.	618	Cassia obtuse-leaved, i.	401	Cataplasma anti-pleuriti-	
Cardui voluntatis acu-		Cassia purging, i.	420	cum, ii.	363
lati radix, i.	622	Cassia senna, i.	401	Cataplasma ad blennor-	
Carduus benedictus, i.	524	Cassamuniar, i.	203	rhagiam, i.	213
Carduus marianus, i.	395	Cassurium occidentale,		Cataplasma ex bryonia,	
Carex arenaria, i.	396	i.	423	i.	463
Carex sea, i.	396	Castaneæ, i.	423	Cataplasma ad cance-	
Carica, i.	675	Catalpa arborea, i.	432	rum, ii.	278
Caricæ pingues, i.	675	Cataplasm ammoniacal,		Cataplasma ad cancerum,	
Carlina acaulis, i.	396	i.	189	i.	463
Carlina vulgaris, i.	396	Cataplasm anodyne, i.	566,	Cataplasma canthari-	
Carlinae humilis radix,		ii.	29, 278, 292	dum, i.	383
i.	396	Cataplasm anthelmintic,		Cataplasma e conio, i.	463
Carlinae radix, i.	396	i.	123	Cataplasma e dauco, i.	463
Carlina thistle, i.	396	Cataplasm anti-cancer-		Cataplasma emolliens,	
Carlina thistle dwarf, i.	396	ous, i.	285, ii. 278	i.	151, 566, ii. 333
Carmine, i.	530	Cataplasm antiseptic, i.	32,	Cataplasma epispasti-	
Caro et spinæ viperæ,			391, 476, ii. 475, 542	cum, ii.	588
i.	697	Cataplasm anti-spasmo-		Cataplasma fermenti pa-	
Caro et spinæ vulpis, i.	697	dic, ii.	299	nis theriacale, ii.	315
Caro testudinis, i.	697	Cataplasm aromatic, ii.	228	Cataplasma lini cum	
Carob-bean tree, i.	440	Cataplasm astringent, i.	645	galbano, i.	113
Carp-stone, i.	354	Cataplasm camphorated		Cataplasma maturans, i.	113,
Carpobalsamum, i.	215	antiseptic, ii.	475		447, 683
Carrot, i.	607	Cataplasma of cantha-		Cataplasma medullæ pa-	
Carrot narrow-leaved, ii.	683	rides, i.	383	nis, i.	151
Carthami lanati herba,		Cataplasma crustæ panis		Cataplasma medullæ pa-	
i.	433	acetosum, i.	42	nis cum cicuta, i.	463
Carthami sativi flores, i.	398	Cataplasm discutient, i.	113,	Cataplasma de mica pa-	
Carthamite, i.	398		312, 744	nis, ii.	333
Carthamus tinctorius, i.	398	Cataplasm emetic, ii.	262	Cataplasma micæ panis	
Carvi semen, i.	397	Cataplasm emollient, i.	151,	anodynum, i.	566
Carui semen, i.	397		213, ii. 200, 216, 333	Cataplasma nidi herun-	
Carum carui, i.	397	Cataplasm for ischuria,		dinis, i.	566
Caryophyllatæ aquaticæ		i.	112	Cataplasma cum opio,	
radix, i.	719	Cataplasm maturative,		ii.	292
Caryophyllatæ herba, i.	719	i.	113, ii. 458	Cataplasma pomaceum,	
Caryophylli, i.	399	Cataplasm mustard, ii.	588	i.	566
Caryophylli aromatici, i.	399	Cataplasm narcotic, ii.	102	Cataplasma resolvens, i.	447
Caryophylline, i.	399	Cataplasm Pradier's, i.	319	ii.	458
Caryophyllorum rubro-		Cataplasm resolvent, i.	151,	Cataplasma resolvens ad	
rum flores, i.	610		344, 444, 447, 463, 634,	schirrhos, i.	463
Caryophyllus aromati-		ii.	378, 379	Cataplasma saponaceum,	
cus, i.	399	Cataplasm Reuss's anti-		ii.	458
Cascarilla, i.	567	septic, i.	476	Cataplasma sinapis, ii.	588
Caseum, ii.	149	Cataplasm rubefacient,		Cataplotia ad catarrhos,	
Cashew apple, i.	423	ii.	363	ii.	301
Cashew-nut tree, i.	423	Cataplasm of sea water,		Catarixæ herba, ii.	259
Castanea equina, i.	90	i.	363	Catch-fly, ii.	587
Castanea pumila, i.	424	Cataplasm sedative, ii.	282	Catelli, i.	697
Castanea vesca, i.	423	Cataplasm against sci-		Cathartine, i.	402
Castor, i.	424	atica, i.	350	Cat mint, ii.	259
Castor fiber, i.	424	Cataplasm starch, i.	214	Catsfoot, i.	230
Castorine, i.	424	Cataplasm strengthen-		Catuputræ minoris se-	
Castranulæ herba, ii.	578	ing, i.	674	mines, i.	629
Cassia bark, ii.	177	Cataplasm suppurative,		Causticum alkalinum,	
Cassia cinnamomea, ii.	178	i.	683, ii. 534	ii.	412
Cassia fistula, i.	420	Cataplasm theriacal, ii.	315	Causticum arsenicale, i.	284
Cassia fistularis, i.	421	Cataplasm of vinegar, i.	42	Causticum commune, ii.	412
Cassia lance leaved, i.	401	Cataplasm of yeast, i.	634	Causticum commune	
Cassia lanceolata, i.	401	Cataplasma acetatum, i.	42	acerrimum, ii.	412

Causticum commune mitius, ii.	413	Cerate cosmetic, ii.	688	Ceratum plumbi super- acetatis, ii.	380
Causticum lunare, i.	273	Cerate for examination during labour, ii.	457	Ceratum refrigerans Galeni, i.	435
Causticum potentiale, ii.	412	Cerate Falck's mercu- rial, ii.	41	Ceratum resinæ, ii.	677
Ceanothus Americana, i.	432	Cerate Galen's, i.	435	Ceratum resinæ pini, ii.	675
Celandine great, i.	445	Cerate golden, i.	594	Ceratum-resinosum, ii.	675
Centaurea behen, i.	432	Cerate green, i.	593	Ceratum-sabinæ, ii.	143
Centaurea lanata, i.	433	Cerate Kirkland's desic- cative, ii.	381	Ceratum saponaceum, ii.	386
Centaurii majoris radix, i.	433	Cerate labial, i.	436, 442	Ceratum saponis, ii.	463
Centaurium officinale, i.	433	ii.	688	Ceratum saturni, ii.	380
Centaurium great, i.	433	Cerate of lard, i.	435	Ceratum saturninum, ii.	385
Centaurium lesser, i.	625	Cerate lip, i.	216	Ceratum simplex, i.	435, 441
Centinodia herba, ii.	406	Cerate mercurial, ii.	2	Ceratum spermatis ceti, i.	441
Centummorbiæ herba, ii.	205	Cerate oleo-resinous, ii.	674	Ceratum e sulphure, ii.	638
Centumnodiæ herba, ii.	406	Cerate opiate, ii.	278	Ceratum tabacinum, ii.	264
Cephælis ipecacuanha, ii.	124	Cerate Pott's, ii.	386	Ceratum viride, i.	593
Cera alba, i.	433	Cerate Rechoux's, i.	186	Ceraunius, i.	354
Cera arborea, ii.	675	Cerate red, ii.	45	Cerberus triceps, i.	239
Cera flava, i.	433	Cerate of resin, ii.	675	Cerefolii herba, i.	444
Cera inaurata, i.	594	Cerate saturnine, ii.	385	Cerefolii hispanici herba, i.	444
Cera ad labia, i.	442	Cerate of savine, ii.	143	Cerei medicati, ii.	379
Cera viridis, i.	593	Cerate of spermaceti, i.	441	Cereoli demulcentes, i.	437
Ceranium ægagrophili- um, ii.	2	Cerate simple, i.	435	Cereoli emollientes, i.	437
Ceranium albinum, ii.	2	Cerate compound of subcarbonate of lead, ii.	401	Cereoli exploratorii, i.	437
Ceranium cancellatum, ii.	2	Cerate of sulphur, ii.	638	ii.	390
Ceranium catenatum, ii.	2	Cerate of tobacco, ii.	264	Cereoli cum kali caus- tico, ii.	414
Ceranium forcipatum, ii.	2	Cerate of Virginian ju- niper, ii.	141	Cereoli mercuriales, ii.	64
Ceranium gracile, ii.	2	Ceratomia siliqua, i.	440	Cereoli plumbici, ii.	379
Ceranium incurvum, ii.	2	Ceratum æruginis, i.	593	Cereoli saturnini, ii.	379
Ceranium scoparium, ii.	2	Ceratum album, i.	441	Cereoli simplices, i.	437
Cerasi racemosi sylves- tris, i.	437	Ceratum album simplex, i.	441	Cerevisia, i.	440
Cerasus avium, i.	438	Ceratum de althæa, i.	151	Cerevisia armoraciæ, ii.	483
Cerasus lauro-cerasus, i.	439	Ceratum anodynum, ii.	391	Cerevisia cochleariæ, i.	534
Cerasus Mahaleb, i.	438	Ceratum cantharidis, i.	382, 386	Cerevisia colocynthidis, i.	575
Cerasus padus, i.	437	Ceratum de cerussa, ii.	373	Cerevisia quercina, ii.	477
Cerasus Virginiana, i.	438	Ceratum cetacei, i.	441	Cerine, i.	433
Cerasus vulgaris, i.	438	Ceratum cetacei rubrum, i.	442	Cervariæ radix, ii.	581
Cerat pour le toucher, ii.	457	Ceratum ceti, i.	441	Cervariæ albæ radix, i.	155
Cerate of acetate of lead, ii.	385	Ceratum citrinum, ii.	675	Ceruss, ii.	371
Cerate antipsoric, ii.	30	Ceratum cosmeticum, i.	441	Ceruss yellow, ii.	369
Cerate arsenical, i.	285	Ceratum detergens, ii.	41	Cerussa, ii.	371
Cerate balsamic, i.	442	Ceratum juniperi Virgini- anæ, ii.	141	Cerussa acetata, ii.	382
Cerate brown, ii.	371	Ceratum fuscum, ii.	391	Cerussa alba Norica, ii.	371
Cerate brown saturnine, ii.	391	Ceratum Galeni, i.	435	Cerussa antimonii, i.	236
Cerate camphorated, i.	375, 442	Ceratum Goulardi, ii.	385	Cerussa citrina, ii.	369
ii.	370	Ceratum ad labia, i.	436, 442	Cervus alces, i.	678
Cerate camphorated of cantharides, i.	387	Ceratum labiale, i.	436, 442	Cervus elaphus, i.	359
Cerate of cantharides, i.	386	Ceratum lithargyri, ii.	380	Ceterach officinarum, i.	443
Cerate of cinchona, i.	486	Ceratum mercuriale Falckii, ii.	41	Cetraria Islandica, ii.	187
Cerate camphorated of cinchona, i.	490	Ceratum minii, ii.	376	Chærefolii herba, i.	444
		Ceratum Noricum, ii.	399	Chærophyllyllum odora- tum, i.	444
		Ceratum plumbi, ii.	380	Chærophyllyllum sativum, i.	444
		Ceratum plumbi com- positum, ii.	382	Chærophyllyllum sylves- tre, i.	444
				Chalcitis, i.	644
				Chalk, i.	352
				Chalk French, ii.	213
				Chalk prepared, i.	355

- Chalk red, i. 155
 Chalk white, i. 352
 Chalybs, i. 634
 Chalybs præparatus, i. 634
 Chalybs tartarizatus, i. 667
 Chamæclemæ herba, i. 721
 Chamædryos herba, ii. 680
 Chamædryos aquaticæ herba, ii. 683
 Chamæleagni herba, ii. 254
 Chamæleontis albi radix, i. 396
 Chamæmeli nobilis flores, i. 446
 Chamæmeli Romani flores, i. 446
 Chamæmeli vulgaris flores, i. 446
 Chamæmori fructus, ii. 531
 Chamæmori herba, ii. 681
 Chamæpityos herba, ii. 681
 Chamæpityos Monsperliacæ herba, ii. 682
 Chameactis radix, ii. 550
 Chamomile, i. 445
 Chamomile fetid, i. 446
 Chamomile Roman, i. 446
 Chamomilla, i. 445
 Chamomilla fœtidæ flores, i. 446
 Chamomillæ nobilis flores, i. 446
 Charcoal, i. 389
 Charcoal animal, i. 390
 Charcoal of bladder-wrack, i. 390
 Charcoal of fucus, i. 390
 Charcoal of sea-pila, i. 390
 Charcoal vegetable, i. 389
 Cheirantes cheiri, i. 455
 Cheiri flores, i. 455
 Chelæ cancerorum, i. 354
 Chelapæ radix, i. 543
 Chelidonii glauci radix, i. 720
 Chelidonii minoris radix, ii. 481
 Chelidonium majus, i. 445
 Chelonites, i. 359
 Chenopodii herba, i. 457
 Chenopodium, ii. 581, 604
 Chenopodium ambrosioides, i. 456
 Chenopodium bonus Henricus, i. 457
 Chenopodium botrys, i. 457
 Chenopodium vulvaria, i. 458
 Cherry bird, i. 437
 Cherry black, i. 438
 Cherry common, i. 438
 Cherry gum, i. 755
 Cherry laurel, i. 439
 Cherry tree, i. 438
 Cherry Virginian, i. 438
 Cherry wild, i. 438
 Cherry water, ii. 350
 Chermes grana, i. 530
 Chervil garden, i. 444
 Chervil scented, i. 444
 Chervil wild, i. 444
 Chesnut, i. 423
 Chesnut chocolate, i. 424
 Chesnut dwarf, i. 424
 Chesnut water, ii. 695
 Chickweed, i. 142
 Chiendent, ii. 343
 Chimaphila corymbosa, i. 458
 China china, i. 469
 China root, ii. 592
 Chinæ veræ radix, ii. 592
 Chinina, ii. 477
 Chloras calcis, i. 351
 Chloride of calcium, i. 363
 Chlorine, i. 458
 Chlorum, i. 458
 Chloruret of antimony, i. 254
 Chloruret of barium, i. 325
 Chloruret of calcium, i. 363
 Chloruret of gold, i. 312
 Chloruret of gold and of sodium, i. 313
 Chloruret prepared of gold and of sodium, i. 313
 Chloruret of lead, ii. 369
 Chloruret of platina, ii. 368
 Chloruret of potassium, ii. 441
 Chloruret of sodium, ii. 602
 Chloruret of tin, ii. 625
 Chloruretum auri, i. 312
 Chloruretum calcis, i. 362
 Chloruretum ferri, i. 656
 Chloruretum oxidi calcii, i. 351
 Chloruretum potassæ oxydatum, ii. 440
 Chloruretum platinæ, ii. 368
 Chloruretum sodii, ii. 602
 Chocolat de santé, ii. 686
 Chocolat à la vanille, ii. 687
 Chocolata aromatica, ii. 687
 Chocolata lichenis Islandici, ii. 188
 Chocolata medica, ii. 686
 Chocolata simplex, ii. 686
 Chocolate anti-syphilitic, ii. 73
 Chocolate chesnut, i. 424
 Chocolate of health, ii. 686
 Chocolate of Iceland moss, ii. 188
 Chocolate purgative, ii. 62
 Chocolate vanilla, ii. 687
 Chocolate white, ii. 541
 Christophorianæ radix, i. 89
 Christophorianæ Americana radix, i. 89
 Chrysanthemum leucanthemum, i. 461
 Chrysolithus, i. 157
 Cicer arietinum, i. 461
 Cichorium endivia, i. 461
 Cichorium intybus, i. 462
 Cichory, i. 462
 Cicuta major, i. 463
 Cicuta terrestris herba, i. 463
 Cicutæ virosæ herba, i. 469
 Cicutæ vulgaris herba, ii. 463
 Cicutaria aquatica, i. 469
 Cicutariæ herba, i. 444
 Cicutine, i. 463
 Cider chalybeate, i. 665
 Cinchona caribæa, i. 470
 Cinchona condaminea, i. 476
 Cinchona floribunda, i. 471
 Cinchona lancifolia, i. 476
 Cinchona lanceolata, i. 470
 Cinchona longifolia, i. 476
 Cinchona nitida, i. 470
 Cinchona pulverized, i. 471
 Cinchona ovata, i. 470
 Cinchonina, i. 503
 Cinchonine, i. 503
 Cinchoninum, i. 503
 Cincus marinus, i. 698
 Cineres clavellati depurati, ii. 419
 Cinnamomum Magellanicum, i. 620
 Cinnabar, ii. 43
 Cinnabaris, ii. 43
 Cinnabaris factitia, ii. 43
 Cinnamomum canella, ii. 160
 Cinnamomum Indicum, ii. 178
 Cinnamomum Sinense, ii. 178
 Cinnamon, ii. 160
 Cinquefoil, ii. 465
 Cissampelos caepeba, i. 504
 Cissampelos pareira, i. 505
 Citrus creticus, i. 757
 Citrus calcaris animalis, i. 364
 Citras calcis, i. 364
 Citras kalicum, ii. 446
 Citras potassæ, ii. 446
 Citras potassæ aquosus, ii. 446
 Citrate of ammonia, i. 202
 Citrate of morphine, ii. 247
 Citrate of potass, ii. 426
 Citrulli semen, i. 578
 Citrus aurantium, i. 505
 Citrus medica, i. 516
 Claretum aromaticum, ii. 356
 Claretum fructuum, i. 439

Claretum ex sex semini- bus, i.	559	Coffee of acorns, ii.	477	Collyrium of henbane, ii.	101
Clarey common, ii.	544	Coffee with cinchona, i.	471	Collyrium ex hydrar- gyro muriatico corro- sivo, ii.	76
Claws crabs', i.	354	Coffea Arabica, i.	537	Collyrium hyoscyami, ii.	101
Clematis crispa, i.	523	Coffea cum cortice Peru- viano, i.	471	Collyrium Lanfranci, i.	282
Clematis erecta, i.	523	Coffea glandium, ii.	477	Collyrium Lanfranc's, i.	282
Clematis vitalba, i.	523	Coffeine, i.	537	Collyrium mercurial, ii.	69
Clematis upright, i.	523	Coix lacrymæ, i.	538	Collyrium ex muriati hydrargyri oxygenata, ii.	68
Clinopodii montani her- ba, ii.	689	Colchicum, i.	538	Collyrium narcotic, ii.	294
Clisma emolliens, ii.	202	Colchicum autumnale, i.	538	Collyrium Neumann's, i.	197
Clisma oleosum, ii.	202	Colchicum Illyricum, ii.	5	Collyrium opiate, ii.	292
Cloth Mai's waxed, i.	437	Colcothar, i.	643	Collyrium opiate satur- nine, ii.	377
Clott-burr, i.	272	Colcothar vitrioli, i.	644	Collyrium plumbatum, ii.	377
Clove-pink, i.	610	Colla, i.	696	Collyrium red, i.	121 ii. 425
Clove tree, i.	399	Colla piscium, i.	696	Collyrium resolvent, i.	192, 593, 596, ii. 387
Cloves, i.	399	Collutorium adstring- ens, i. 77, 163, ii. 516, 609		Collyrium roborant, i.	261
Club-moss, ii.	204	Collutorium alkaline, ii.	423	Collyrium of sal amoni- ac, i.	168
Clyssus antimonii, i.	79	Collutorium aluminous, i.	163	Collyrium de salibus igne fusi, i.	585
Clyisma anti-spasmodico- resolvens, ii.	603	Collutorium alumino- sum, i.	163	Collyrium sapphiricum, i.	590
Clyisma ad ascarides, ii.	603	Collutorium antiscorbu- tic, i.	352	Collyrium saturnine, ii.	379
Clyisma cum assafoetida, i.	303	Collutorium antiscorbu- ticum; i.	520	Collyrium simple, i.	547
Clyisma emolliens an- thelminticum, ii.	508	Collutorium astrigent, ii.	518	Colocynth, i.	571
Clyisma excitans, i.	492	Collutorium catechu sa- turninum, ii.	387	Colombæ radix, i.	527
Clyisma incitans, i.	371	Collutorium detergens, i.	77, ii. 516	Colophony, ii.	673
Clyisma incitans et se- dans, i.	303	Collutorium gingivale, i.	536	Colophonium, ii.	673
Clyisma nutriens, i.	700	Collutorium saturnine, ii.	377	Color Indicus, ii.	351
Clyisma oleosum, ii.	270	Collutorium sinapi, ii.	590	Coltsfoot, ii.	699
Clyisma purgans, ii.	603	Collyrium aluminosum, i.	162	Colubrinæ radix, i.	277
Clyisma resolvens, ii.	459	Collyrium aluminous, i.	162	ii.	406
Clyisma sedans et emol- liens, ii.	326	Collyrium anodyne, i.	566	Columbæ radix, i.	527
Clyisma ad soporosos af- fectus, i.	265	ii.	283, 295	Columbine, i.	270
Clyisma tanacetii, i.	303	Collyrium adstringens, i.	744	Colutea arborescens, i.	401, 540
Clyisma tonica et anti- spasmodicum, i.	303	Collyrium astrigent, i.	744	Comæ floridæ herba, i.	625
Clyisma vini antimonii irritans, i.	265	Collyrium boracicum, ii.	610	Comfits of cinchona, i.	483
Clyssus sulphuris, i.	78	Collyrium borated, ii.	610	Comfits vermifuge, ii.	51
Cobalt, i.	526	Collyrium Brown's, i.	120	Comfrey great, ii.	641
Cobaltum, i.	526	Collyrium cæruleum, i.	590	Conchæ citratæ, i.	364
Cocci cnidii grana, i.	603	Collyrium cupratum, i.	585	Concrementa astaci flu- viatilis, i.	354
Coccinella septempunc- tata, i.	526	Collyrium detersive and resolvent, i.	585	Conditum anti-epilepti- cum, ii.	296
Coccognidii baccae, i.	603	Collyrium dry, ii.	38, 57, 62	Conditum calami, i.	86
Coccoloba urifera, ii.	146	Collyrium emollient, i.	148	Conditum catechueicum, ii.	317
Cocculi semen, i.	520	Collyrium exsiccans, i.	590	Conditum corroborans, ii.	673
Cocculus Indicus, i.	529	Collyrium Fernandez, ii.	66	Conditum cydoniorum, i.	598
Cocculus palmatus, i.	527	Collyrium ferruginous, i.	652	Conditum epilepticum, ii.	705
Coccus cacti, i.	529	Collyrium Gimbernati's, ii.	414	Conditum febrifugum, ii.	669
Coccus lacca, ii.	150				
Cochineal, i.	529				
Cochlearia armoracia, ii.	481				
Cochlearia officinalis, i.	531				
Cocoa, ii.	686				
Cochleariæ hortensis herba, i.	531				
Cocos butyracea, ii.	271				
Coffee, i.	537				

Conditum hydrargyrounguinum, ii.	14	Confection anti-epileptic, ii.	296	Conserva rosarum sicca, ii.	511
Conditum iridis saccharatum, ii.	132	Confection aromatic, i. ii.	204 165	Conserva rosarum vitriolata, ii.	511
Conditum muscarium, i.	166	Confection of cassia, i.	422	Conserva sabinæ, ii.	142
Conditum nephriticum, ii.	657	Confection common, ii.	165	Conserva florum salviæ, ii.	544
Conditum pectorale, ii.	636	Confection of fly-agaric, i.	166	Conserva symphyti, ii.	642
Conditum purgativum, ii.	492	Confection Hamech, i.	572	Conserva tanacetii, ii.	649
Conditum satyrii, ii.	540	Confection of hyacinth, i.	565	Conserva florum tilia, ii.	691
Conditum tamarindaceum, i.	406	Confection of iris, ii.	132	Conserva veronicæ, ii.	717
Conditum ad vermes, i.	546	Confection kermesine, i.	530	Conserva violarum, ii.	722
Conein, i.	463	Confection laxative, i.	5.6	Conserve of angelica, i.	222
Confectio alkermes, i.	530	Confection of rue, ii.	536	Conserve anti-scorbutic, i.	531
Confectio anisi laxativa, i.	546	Confection of senna, i.	404	Conserve of arum, i.	292
Confectio aromatica, ii.	164	Confection of senna and charcoal, i.	407	Conserve of balm, ii.	228
Confectio aurantiorum, i.	506	Confection of sweet-flag, i.	86	Conserve of beccabunga, ii.	119
Confectio cassiæ, i.	422	Confection vermifuge, ii.	583	Conserve of betony, i.	337
Confectio cassiæ fistulæ, i.	422	Coni Lupuli, ii.	19	Conserve of black currant, ii.	505
Confectio chinæ anti-epileptica, i.	479	Conii maculati herba, i.	463	Conserve of blessed thistle, i.	524
Confectio chinæ antimonialis, i.	479	Conserva anthos, ii.	522	Conserve of borage, i.	341
Confectio seminis cinæ, ii.	583	Conserva rutæ, ii.	536	Conserve of bugloss, i.	215
Confectio cinnamomi regia, ii.	165	Conserva aurantii, i.	507	Conserve of cassia, i.	421
Confectio cordialis, ii.	164	Conserva corticum aurantiorum, i.	506	Conserve of chamomile, i.	446
Confectio cynosbatos, ii.	511	Conserva florum aurantiorum, i.	506	Conserve of cherry-laurel, i.	439
Confectio diascordii, ii.	317	Conserva cassiæ, i.	422	Conserve of cichory, i.	462
Confectio fistulæ, i.	422	Conserva florum chamæmeli, i.	446	Conserve of clove-pink, i.	610
Confectio hyacinthina, i.	565	Conserva citri, i.	517	Conserve of coltsfoot, ii.	699
Confectio de hyacintho, i.	565	Conserva citri aurantii, i.	566	Conserve of convallaria, i.	541
Confectio Japonica, ii.	317	Conserva cornorum, i.	460	Conserve of cornel berries, i.	560
Confectio margaritarum kermesina, i.	530	Conserva cydoniorum, i.	598	Conserve of cowslip, ii.	466
Confectio opii, ii.	306	Conserva cynosbatos, ii.	511	Conserve of daisy, i.	329
Confectio passulata, i.	403	Conserva erysimi, i.	623	Conserve of elecampane, ii.	115
Confectio de prunis, i.	404	Conserva farfaræ, ii.	699	Conserve of eringo, i.	622
Confectio de Rebecha, i.	725	Conserva lepidii, ii.	186	Conserve of fumitory, i.	679
Confectio rosæ caninæ, ii.	516	Conserva florum lillii convallium, i.	541	Conserve of great comfrey, ii.	642
Confectio rosæ Gallicæ, ii.	510	Conserva florum liliorum alborum, ii.	198	Conserve of ground-ivy, i.	721
Confectio rutæ, ii.	536	Conserva foliorum menthæ, ii.	234	Conserve of hemlock, ii.	463
Confectio santalorum cordialis, i.	565	Conserva mesembryanthemi crystallini, ii.	242	Conserve of hips, ii.	511
Confectio scammonia, i.	561	Conserva millepedum, ii.	272	Conserve of horehound, ii.	222
Confectio sennæ, i.	404	Conserva florum persicarum, ii.	344	Conserve of hyssop, ii.	111
Confectio sennæ compositum, i.	404	Conserva portulacæ, ii.	410	Conserve of iceplant, ii.	242
Confectio diaprunum, i.	404	Conserva primulæ veris, ii.	466	Conserve of Jerusalem oak, i.	457
Confectio sennæ cum pulpis, i.	404	Conserva rhododendri chrysanthi, ii.	502	Conserve of juniper, ii.	137
Confectio florum tanacetii, ii.	649	Conserva rosarum rubrarum adstringens, ii.	512	Conserve of lemon, i.	517
Confection of almonds, i.	296			Conserve of lesser centaury, i.	625
				Conserve of lily, ii.	198

Conserve of lime flowers, ii. 691	Contrayerva Germanicæ radix, i. 296	Corrector Starkeyi, ii. 671
Conserve of mallow, ii. 215	Contrayerva Virginianæ radix, i. 277	Corsican moss, ii. 2
Conserve of marigold, i. 365	Convallaria majalis, i. 541	Cortex acaciæ nostratis, ii. 467
Conserve of marjoram, ii. 322	Convolvulus jalapa, i. 542	Cortex caryophylloides, ii. 178
Conserve of mint, ii. 234	Convolvulus mechoacanna, i. 553	Cortex cascarillæ, i. 567
Conserve of orange, i. 507	Convolvulus scammonia, ii. 559	Cortex chacarillæ, i. 567
Conserve of orange flowers i. 506	Convolvulus scoparius, ii. 195	Cortex citri, i. 517
Conserve of orange peel, i. 506	Convolvulus turpethum, i. 553	Cortex cotini, ii. 503
Conserve of orchis, ii. 540	Conyza squarrosa, i. 554	Cortex culilaban, ii. 178
Conserve of peach flowers, ii. 344	Conyza cærulea herba, i. 621	Cortex Curassavenses, i. 506
Conserve of peony, ii. 329	Conyza majoris herba, i. 622	Cortex eleutheriæ, i. 567
Conserve of pepperwort, ii. 186	Conyza mediæ herba, ii. 114	Cortex gascarillæ, i. 567
Conserve of petty broom, i. 709	Conyza minoris herba, i. 621	Cortex Magellanicus, i. 620
Conserve of plantain, ii. 366	Conyza vulgaris herba, i. 554	Cortex Peruvianus, i. 469
Conserve of purslain, ii. 410	Copaiba, i. 554	Cortex Peruvianus flavus, i. 476
Conserve of quince, i. 598	Copaifera officinalis, i. 554	Cortex Peruvianus fuscus, i. 470
Conserve of red poppy, ii. 334	Copaivæ balsamum, i. 554	Cortex Peruvianus offici- nalis, i. 470
Conserve of red roses, ii. 510	Copal gum, i. 755	Cortex Peruvianus ruber, i. 476
Conserve resolvent and alterative, ii. 444	Copal resin, i. 755	Cortex pocgeregæ, ii. 402
Conserve of rhododendron ii. 502	Copalm, ii. 627	Cortex quercinus, ii. 474
Conserve of rocket, i. 342	Copper, i. 581	Cortex quercus, ii. 474
Conserve of rosemary, ii. 522	Copper Armenian, i. 582	Cortex sambuci, ii. 551
Conserve vitriolated of roses, ii. 511	Copper calcined, i. 582	Cortex simarubæ, ii. 587
Conserve of rue, ii. 536	Copperas, i. 583	Cortex Soymidæ, ii. 641
Conserve of sage, ii. 544	Coptis trifolia, i. 558	Cortex suberi, ii. 475
Conserve of savine, ii. 142	Coral red, i. 353	Cortex Winteranus, i. 620
Conserve of scordium, ii. 683	Coral white, i. 353	Cortex Winteranus spu- rius, i. 376, 561
Conserve of sloes, ii. 468	Corallina corsica, ii. 2	Corylus avellana, i. 561
Conserve of smallage, i. 266	Corallina officinarum, ii. 2	Corydalis bulbosus, i. 561
Conserve of sorrel, ii. 534	Corallina rubens, ii. 2	Costæ herba, ii. 111
Conserve spagyric, ii. 430	Corallium album, i. 353	Costi hortorum radix, i. 317
Conserve of sweet flag, i. 86	Corallium rubrum, i. 353	Costi nostratis herba, ii. 111
Conserve of tamarinds, ii. 646	Cordia myxa, i. 558	Costus amarus, i. 561
Conserve of tansey, ii. 649	Cordial Brodum's, i. 139	Costus Arabian, i. 561
Conserve of veronica, ii. 717	Coreopsis bidens, ii. 621	Costus Arabicus, i. 561
Conserve of violet, ii. 722	Coriander, i. 559	Costus corticosus, i. 376, 561
Conserve of water lily, ii. 268	Coriandrum sativum, i. 559	Costus dulcis, i. 561
Conserve of wild mustard i. 623	Cornel male, i. 560	Cotini cortex, ii. 503
Conserve of woodlice, ii. 272	Cornel round-leaved, i. 559	Cotton, i. 730
Conserve of wood sorrel, ii. 327	Cornine, i. 560	Cotton thistle, ii. 273
Conserve of wormwood, i. 31	Cornu cervi, i. 359	Cotton tree, i. 730
Consolidæ majoris radix, ii. 641	Cornu alcis, i. 359	Cotula aurea, i. 562
Consolidæ medicæ herba i. 103	Cornu cervi auratum, i. 314	Cotyledon lutea, i. 562
Consolidæ minoris herba ii. 465	Cornu cervi raspatum, i. 359	Cotyledon umbilicus, i. 562
Consolidæ regalis herba, i. 608	Cornu cervi philosophice præparatum, i. 361	Couch-grass, ii. 696
Consolidæ Saracenicæ herba, i. 618	Cornu cervi sine igne præ- paratum, i. 361	Coumarona odorata, i. 562
	Cornu cervi ustum, i. 360	Cowhage, ii. 625
	Cornu cervi ustum præ- paratum, i. 360	Cow-parsnip, ii. 5
	Cornus circinnata, i. 559	Cowslip, ii. 465
	Cornus florida, i. 560	Cow's parsley, i. 444
	Cornus mas, i. 560	Crab's claws, i. 354
	Cornus sericea, i. 560	Cranberry, ii. 702
	Coronopus ruellii, i. 560	Crane's-bill, i. 718
	Corpus pro balsamo sapo- nis, ii. 672	Cranium humanum, i. 359
	Corrallina rubra, ii. 2	Crassulæ radix, ii. 580
	Corrector opii, ii. 671	Cream of cinnabar, ii. 43
		Cream pectoral, i. 60, ii. 687
		Cream of tartar, ii. 450

Cremor, ii.	149	Cunilæ bulbulæ herba, i.	324	Cyperus edible, i.	601
Cremor tartar, ii.	450	Cunilæ sativæ herba, i.	558	Cyperus esculentus, i.	601
Cremor tartaris solubilis, ii.	454	Cupressini officinalis radix linariæ admodum similis, i.	629	Cyperus rotundus, i.	602
Crepitus lupi, ii.	204	Cupressus sempervirens, i.	580	Cyperus round, i.	602
Cress common, ii.	688	Cupri subacetæ, i.	590	Cyperus long, i.	602
Cress Indian, ii.	698	Cupridum ammoniacæ liquidum, i.	596	Cyperus longus, i.	602
Cress penny, ii.	688	Cuprum, i.	581	Cyphoides Galli, ii.	663
Cress winter, i.	623	Cuprum aceticum, i.	595	Cypress, i.	580
Cresswort thorough-stem, i.	628	Cuprum aluminatum, i.	585	Cypress nuts, i.	580
Creta, i.	352	Cuprum ammoniacale, i.	586	Cypress spurge, i.	629
Creta alba, i.	352	Cuprum ammoniaco-sulphuricum, i.	586	Cypri antiquorum radix, ii.	185
Creta barotica, i.	328	Cuprum armenus, i.	582	Cytinus hypocistis, ii.	12
Creta Gallica, ii.	213	Cuprum calcinatum, i.	582	Cytiso-genistæ herba, i.	708
Creta præcipitata, i.	355	Cuprum sulphuricum, i.	583	Dactyli, ii.	377
Creta præparata, i.	355	Cuprum vitriolatum, i.	583	Dactylus idæus, i.	354
Crithmum maritimum, i.	562	Cupulæ glandium quercus, ii.	474	Dædalea suavolens, i.	602
Crocus antimonii, i.	241	Curcuma angustifolia, i.	632	Daisy, i.	329
Crocus ferri, i.	646	Curcuma longa, i.	581	Daisy French, i.	722
Crocus martis aperitivus, i.	646	Curcuma rotunda, i.	581	Daisy greater ox-eye, i.	461
Crocus martis rore majali paratus, i.	646	Currant black, ii.	504	Dalbergia monetaria, i.	765
Crocus martis sulphuratus, i.	644	Currant red, ii.	79	Dandelion, ii.	651
Crocus martis vitriolatus, i.	644	Cuscuta epithymum, i.	596	Daphne gnidium, i.	603
Crocus metallorum, i.	241	Cuscuta Europæa, i.	597	Daphne laureola, i.	603
Crocus sativus, i.	563	Cuscutæ majoris herba, i.	597	Daphne mezereum, i.	602
Crocus solis, i.	313	Cusparia febrifuga, i.	228	Daphne thymelea, i.	603
Crocus veneris, i.	581	Cuttle-fish bone, i.	354	Date tree, ii.	347
Croton cascarilla, i.	567	Cyanuret of mercury, ii.	80	Datura fastuosa, i.	605
Croton castaneifolium, ii.	150	Cyanuret hydrated of iron, i.	674	Datura metel, i.	605
Croton eleutheria, i.	567	Cyanuretum ferri, i.	674	Datura stramonium, i.	605
Croton tiglium, i.	570	Cyanuretum hydrargyri, ii.	80	Dauci candiani semina, i.	307
Crow flower, i.	217	Cyanus segetum, i.	597	Dauci cretici semina, i.	307
Crowfoot bulbous, ii.	480	Cyclamen Europæum, i.	597	Dauci cyanopi radix, i.	356
Crowfoot marsh, ii.	480	Cyclaminis officinalis radix, i.	597	Decoctio rubi villosi, i.	532
Crowfoot upright, ii.	480	Cydonia communis, i.	597	Daucus carota, i.	607
Cruciata radix, i.	691	Cydonia cotonea, i.	598	Decoction of alcornocho, i.	112
Crystalli lunæ, i.	273	Cymini semen, i.	579	Decotion Allen's i.	495
Crystalli veneris, i.	595	Cynanchum argel, i.	401	Decotion alternative, ii.	498
Crystallum minerale, ii.	433	Cynanchum Monspelianum, i.	600	Decotion of angustura, i.	229
Crystals of tartar, ii.	450	Cynanchum Montpellier, i.	600	Decotion of annual mercury, ii.	211
Cubebs, ii.	360	Cynara cardunculus, i.	600	Decotion anthelmintic, i.	123
Cuckoo-pint, i.	291	Cynara scolymus, i.	601	Decotion anti-arthritis, ii.	598
Cucumber, i.	578	Cynoglossæ majoris radix, i.	601	Decotion anti-asthmatic, i.	432
Cucumber squirting, ii.	243	Cynoglossæ vulgaris radix, i.	601	Decotion anticachectic, i.	426
Cucumeris agrestis radix, ii.	243	Cynoglossum officinale, i.	601	Decotion antidysenteric, i.	317
Cucumeris semen, i.	578	Cynorrhodon, ii.	510	Decotion anti-hæmorrhoidal, i.	414
Cucumis colocynthis, i.	570	Cynosbati fructus, ii.	510	Decotion Rosenstien's anthelmintic, ii.	14
Cucumis melo, i.	577	Cynosbati semina, ii.	510	Decotion anti-icteric, i.	677
Cucumis sativus, i.	578	Cyperi odorati radix, i.	602	Decotion antiphthisical, i.	466
Cucurbita citrullus, i.	578	Cyperi orientalis majoris radix, i.	602	Decotion antiscorbutic, ii.	9, 486
Cucurbita lagenaria, i.	578	Cyperi Romani radix, i.	602		

Decoction antiseptic, i.	280	Decoction of lesser centaury, i.	626	Decoction emetic, i.	295
ii.	684	Decoction of dried cherries, i.	439	Decoction emmenagogue, ii.	405
Decoction antistrumous, i.	393	Decoction of china root, ii.	592	Decoction emollient, i.	147
Decoction antisyphilitic, i.	396	Decoction of cichory, i.	462	Decoction emollient and carminative, i.	147
Decoction aperient pectoral, i.	420	Decoction of cinchona, i.	489	Decoction febrifuge, i.	495
Decoction aperitive, i.	297	Decoction of cinchona and cascarilla, i.	493	ii.	424
Decoction nitrated aperitive, i.	297	Decoction of cinchona with pulverized cinchona, i.	494	Decoction saline febrifuge, i.	626
Decoction of apples, ii.	215	Decoction of cinchona and Iceland-moss, i.	493	Decoction of figs, i.	675
Decoction aqueous of hearts-ease, ii.	724	Decoction of cinchona and leopards-bane, i.	492	Decoction Forest's, ii.	10
Decoction of aralia nudicaulis, i.	270	Decoction of cinchona with opium, i.	492	Decoction of gelatine, i.	701
Decoction aromatic, ii.	545	Decoction of cinchona and rhubarb, i.	494	Decoction of Surinam Geofræa, i.	717
Decoction of asparagus, i.	296	Decoction of cinchona and seneka, i.	493	Decoction of german-der, ii.	681
Decoction of astragalus, i.	307	Decoction of cinchona with serpentaria, i.	494	Decoction of gratiola, i.	731
Decoction astringent, ii.	693	Decoction of cinchona and valerian, i.	495	Decoction of guaiac, i.	737
Decoction of avens, i.	719	Decoction acid of cinchona, i.	492	Decoction of gum arabic, i.	751
Decoction æthereous of avens, i.	719	Decoction alkaline of cinchona, i.	495	Decoction of gum arabic with hartshorn, i.	752
Decoction æthereous of avens with cascarilla, i.	720	Decoction aluminated of cinchona, i.	491	Decoction of hartshorn, i.	702
Decoction æthereous of avens with rhubarb, i.	720	Decoction factitious of cinchona, i.	90	Decoction acidulous of hartshorn, i.	702
Decoction of barley, ii.	7	Decoction nitrated of cinchona, i.	490	Decoction lacteal of hearts-ease, ii.	724
Decoction acidulated of barley, ii.	8	Decoction opiate of cinchona, i.	490	Decoction Heister's, ii.	388
Decoction of barley with gum, ii.	8	Decoction of great comfrey, ii.	642	Decoction of white helibore, ii.	715
Decoction of bead tree root, ii.	225	Decoction of cork, ii.	475	Decoction hemastatic, i.	488
Decoction of bird-cherry, i.	439	Decoction of Corsican moss, ii.	2	Decoction of horse chestnut bark, i.	91
Decoction of bistort, ii.	407	Decoction of dandelion, ii.	651	Decoction Hoffmann's of spruce fir, i.	23
Decoction æthereous bitter, i.	713	Decoction deobstruent, ii.	652	Decoction hydragogue, ii.	568
Decoction alcoholized bitter, i.	713	Decoction diaphoretic, ii.	701	Decoction bitter of Iceland moss, ii.	189
Decoction of bitter-sweet, ii.	616	Decoction of digitalis, i.	614	Decoction lacteal of Iceland moss, ii.	189
Decoction of blackberry, ii.	532	Decoction diuretic, i.	269, 405, 568	Decoction mucilaginous of Iceland moss, ii.	189
Decoction of borage, i.	341	Decoction alkaline diuretic, ii.	698	Decoction of juniper leaves, ii.	141
Decoction of bran, i.	681	Decoction of dock, ii.	535	Decoction of juniper wood, ii.	141
Decoction of bryony, i.	344	Decoction of dog-wood, i.	560	Decoction laxative, ii.	468
Decoction of burdock, i.	272	Decoction of elder bark, ii.	553	Decoction of lemon, i.	521
Decoction of burnt sponge, i.	393	Decoction of elecampane, ii.	117	Decoction of leopard's-bane, i.	280
Decoction Callac's, ii.	599	Decoction of elm bark, ii.	700	Decoction of logwood, ii.	193
Decoction of calumba, i.	528			Decoction of madder, ii.	530
Decoction of blue cardinal flower, ii.	203			Decoction alkaline of madder, ii.	531
Decoction carminative, ii.	141			Decoction of male fern, i.	308
Decoction of cassia, i.	421			Decoction of malt, ii.	214
Decoction catholic, ii.	597			Decoction of marsh-mallows, i.	147
				Decoction of mastic, ii.	223

Decoction of mercury, ii. 13	Decoction of semen con- tra, ii. 582	Decoction compound of chamomile, i. 452
Decoction of mezereon, i. 603	Decoction of seneka, ii. 404	Decoction compound of dog-grass, ii. 698
Decoction of mountain laurel, ii. 145	Decoction of serpenta- ria, i. 278	Decoction compound of elecampane, ii. 117
Decoction myrrhous sa- turnine, ii. 388	Decoction of simarouba, ii. 587	Decoction compound of gentian, i. 712
Decoction narcotic, ii. 617	Decoction of soapwort, ii. 557	Decoction compound of guaiaac, i. 737
Decoction nervine, i. 510	Decoction of spigelia, ii. 620	Decoction compound of isinglass, ii. 596
Decoction of nutgalls, i. 693	Decoction of spruce fir, i. 22	Decoction compound of mallow, ii. 216
Decoction of oak bark, ii. 476	Decoction saponaceous of spruce fir, i. 23	Decoction compound of mezeron, i. 603
Decoction of oak bark with alum, ii. 476	Decoction vinous of spruce fir, i. 23	Decoction compound of great mullein, ii. 716
Decoction of oak bark with vinegar, ii. 476	Decoction of spurred rye, ii. 579	Decoction compound of quassia, ii. 473
Decoction of orange leaves, i. 510	Decoction of squill, ii. 568	Decoction compound of sarsaparilla, ii. 595
Decoction of orange peel, i. 510	Decoction of stavesacre, i. 608	Decoction compound of tamarinds, ii. 647
Decoction of pareira brava, ii. 505	Decoction stimulant su- dorific, ii. 596	Decoction compound of tamarinds, ii. 647
Decoction of peach leaves, ii. 344	Decoction of stramon- ium, i. 605	Decoction compound of tamarinds, ii. 647
Decoction pectoral, i. 148, 729. ii. 9, 189, 325, 347, 405	Decoction sudorific and nutritive, i. 739	Decoction compound of tamarinds, ii. 647
Decoction stimulant pec- toral, ii. 190	Decoction of tamarinds, ii. 646	Decoction compound of tamarinds, ii. 647
Decoction of wall-pelli- tory, ii. 341	Decoction against tinea, i. 739	Decoction compound of tamarinds, ii. 647
Decoction of pine buds, ii. 358	Decoction of tobacco, ii. 261	Decoction compound of tamarinds, ii. 647
Decoction Pollini's, ii. 136	Decoction tonic and composing, i. 488	Decoction compound of tamarinds, ii. 647
Decoction of polygala amara, ii. 404	Decoction of tormentil, ii. 693	Decoction compound of tamarinds, ii. 647
Decoction of pomegran- ate rind, ii. 471	Decoction vermifuge, ii. 3, 582	Decoction compound of tamarinds, ii. 647
Decoction Rehmann's of pomegranate rind, ii. 471	Decoction of uva ursi, i. 271	Decoction compound of tamarinds, ii. 647
Decoction of the bark of pomegranate root, ii. 471	Decoction of walnut rind, ii. 136	Decoction compound of tamarinds, ii. 647
Decoction of poppy, ii. 338	Decoction Sydenham's white, ii. 332	Decoction compound of tamarinds, ii. 647
Decoction of quassia, ii. 473	Decoction Trommsdorf's white, i. 703	Decoction compound of tamarinds, ii. 647
Decoction of quince seeds, i. 599	Decoction of willow, ii. 543	Decoction compound of tamarinds, ii. 647
Decoction of ratany, ii. 148	Decoction of willow herb, ii. 206	Decoction compound of tamarinds, ii. 647
Decoction of rice, ii. 325	Decoction of worm bark, i. 717	Decoction compound of tamarinds, ii. 647
Decoction acidulated of rice, ii. 325	Decoction Zittmann's, ii. 599	Decoction compound of tamarinds, ii. 647
Decoction of rhubarb, ii. 495	Decoction compound of aloes, i. 134	Decoction compound of tamarinds, ii. 647
Decoction of salep, ii. 541	Decoction compound of barley, ii. 9	Decoction compound of tamarinds, ii. 647
Decoction of yellow sadd- ers, ii. 196	Decoction compound of bitter-sweet, ii. 616	Decoction compound of tamarinds, ii. 647
Decoction of sarsaparilla, ii. 594	Decoction compound of calumba, i. 528	Decoction compound of tamarinds, ii. 647
Decoction of savine, ii. 143	Decoction compound of dandelion, ii. 652	Decoction compound of tamarinds, ii. 647
Decoction bezoardic of scorzonera, ii. 578		Decoction compound of tamarinds, ii. 647

- Decoctum calumbæ, i. 528
 Decoctum carbonatis potassæ absinthii, ii. 424
 Decoctum caricarum, i. 675
 Decoctum cassiæ fistulæ, i. 421
 Decoctum cerasorum acidorum siccatorum, i. 439
 Decoctum chamædryos amarum, ii. 681
 Decoctum chamæmeli compositum, i. 452
 Decoctum chinæ acidum, i. 492
 Decoctum chinæ cum alumine, i. 491
 Decoctum chinæ cum arnica, i. 492
 Decoctum chinæ cum cascarilla, i. 493
 Decoctum chinæ cum china pulverata, i. 494
 Decoctum chinæ compositum, i. 494
 Decoctum chinæ compositum laxans, i. 496
 Decoctum chinæ cum lichen, i. 493
 Decoctum chinæ opiatum, i. 490
 Decoctum chinæ cum radice caryophyllatæ et opio, i. 492
 Decoctum chinæ radicis, ii. 592
 Decoctum chinæ cum rheo, i. 494
 Decoctum chinæ cum senega, i. 493
 Decoctum chinæ cum valeriana, i. 495
 Decoctum cinchonæ, i. 489
 Decoctum cinchonæ cardiacum, i. 494
 Decoctum cinchonæ lancifoliæ, i. 489
 Decoctum citri Mynsichti, i. 521
 Decoctum columbæ compositum, i. 528
 Decoctum corni floridæ, i. 560
 Decoctum cornu cervi, i. 703
 Decoctum cornu cervi compositum, ii. 332
 Decoctum cornu cervi gummosum, i. 752
 Decoctum cornu cervi cum mica panis, ii. 332
 Decoctum cornu cervini, i. 752
 Decoctum corticis cinchonæ, i. 489
 Decoctum corticis hippocastani, i. 91
 Decoctum corticis mezerei, i. 603
 Decoctum corticis Peruviani, i. 489
 Decoctum corticis quercus cum aceto, ii. 476
 Decoctum corticis quercus cum alumine, ii. 476
 Decoctum corticis salicis laureæ, ii. 543
 Decoctum corticis sambuci, ii. 553
 Decoctum corticis viridis nucum juglandium, ii. 136
 Decoctum crystallorum tartari, ii. 452
 Decoctum cucumerum cum extracto nucis vomicæ, ii. 267
 Decoctum cydoniæ, i. 599
 Decoctum daphnes mezerei, i. 603
 Decoctum diaphoreticum, i. 737, ii. 701
 Decoctum diaphoreticum incitans, i. 596
 Decoctum digitalis, i. 614
 Decoctum diureticum alkalinum, ii. 698
 Decoctum diureticum Hufelandi, ii. 568
 Decoctum drupæ juglandis regiæ, ii. 136
 Decoctum graminis compositum, ii. 698
 Decoctum gratiolæ, i. 731
 Decoctum guaiaci, i. 737
 Decoctum guaiaci compositum, i. 737
 Decoctum gummi arabici, i. 751
 Decoctum hæmatoxyli, ii. 193
 Decoctum helenii, ii. 117
 Decoctum helminthocorti, ii. 2
 Decoctum herbæ violæ tricoloris cum aqua, ii. 724
 Decoctum herbæ violæ tricoloris cum lacte, ii. 724
 Decoctum hordeatum amygdalinum, i. 207
 Decoctum hordei acidulatum, ii. 8
 Decoctum hordei citratum, ii. 8
 Decoctum hordei compositum, ii. 9
 Decoctum hordei decorticati, ii. 7
 Decoctum dulcamaræ, ii. 616
 Decoctum febrifugum, i. 490
 Decoctum febrifugum, laxans, i. 496
 Decoctum febrifugum magistrale, i. 495
 Decoctum febrifugum salinum, i. 626
 Decoctum febrifugum simplex, i. 91
 Decoctum foliorum aurantii, i. 510
 Decoctum fructuum juniperi carminativum, ii. 141
 Decoctum ad fungos articularum, ii. 388
 Decoctum furfuris tritici, i. 681
 Decoctum gallarum, i. 693
 Decoctum gei urbani, i. 719
 Decoctum Geoffrææ Jamaicensis, i. 717
 Decoctum Geoffrææ Surinamensis, i. 717
 Decoctum Geoffroyæ inermis, i. 717
 Decoctum graminis, ii. 698
 Decoctum hordei distichi, ii. 7
 Decoctum hydrargyri, ii. 13
 Decoctum incitans aromaticum, ii. 545
 Decoctum incrassans Fuleri, i. 751
 Decoctum ipecacuanhæ, ii. 127
 Decoctum juniperi, ii. 141
 Decoctum kalmiæ latifoliæ, ii. 145
 Decoctum lapathi, ii. 535
 Decoctum laxativum, ii. 219
 Decoctum laxativum Vieanense, i. 415
 Decoctum leontodontis, ii. 651
 Decoctum lichenis Islandici, ii. 189
 Decoctum lichenis Islandici compositum, ii. 190
 Decoctum lichenis Islandici lacteum, ii. 189
 Decoctum ligni campechiani, ii. 193
 Decoctum ligni guaiaci compositum, ii. 595
 Decoctum ligni juniperi, ii. 141
 Decoctum ligni quassiæ, ii. 473
 Decoctum ligni santali citrini, ii. 196
 Decoctum lignorum, i. 737
 Decoctum lobeliæ syphiliticæ, ii. 203
 Decoctum ad lupias synoviales Heisteri, ii. 388
 Decoctum malti, ii. 214
 Decoctum malvæ compositum, ii. 216

Decoctum mannae carminativum, ii.	219	Decoctum resolvens, ii.	652	Dentalia, i.	353
Decoctum mannae purgans, ii.	219	Decoctum resolvens et diureticum, ii.	438, 444	Dentalium dentalis, i.	353
Decoctum ad metorrhagiam, i.	510	Decoctum rhubarbari, ii.	495	Dentalium dentalis, i.	353
Decoctum mezerei, i.	603	Decoctum rubiae compositum, ii.	531	Dentaria digitata, i.	609
Decoctum mezerei compositum, i.	603	Decoctum rubiae tinctoriae, ii.	530	Dentaria pinnata, i.	609
Decoctum ex myrtillis, i.	702	Decoctum sabinæ, ii.	143	Dentariæ majoris radix, ii.	368
Decoctum nervinum, i.	510	Decoctum salicariæ, ii.	206	Dentaria radix, i.	230
Decoctum nicotianæ, ii.	261	Decoctum saponariæ officinalis, ii.	557	Dentes apri, i.	359
Decoctum nigrum Allenii, i.	495	Decoctum sarsæ, ii.	594	Dentes equi marini, i.	359
Decoctum ophthalmicum mucilaginosum, i.	147	Decoctum sarsaparillæ, ii.	594	Dentes hippopotamis, i.	359
Decoctum oryzæ, ii.	325	Decoctum sarsaparillæ compositum, ii.	595, 598	Dentis leonis radix, ii.	651
Decoctum papaveris, ii.	338	Decoctum sarsaparillæ excelsæ, ii.	599	Dentillariæ radix, ii.	368
Decoctum pareira, i.	505	Decoctum saturninum, ii.	375	Desmarestia aculeata, i.	2
Decoctum pectorale corroborans, ii.	405	Decoctum scillæ, ii.	568	Desvauxia fluitans, i.	609
Decoctum pectorale nutriens, ii.	189	Decoctum scorzonæræ bezoardicum, ii.	578	Devil-in-a-bush, ii.	264
Decoctum pectorale nutriens et incitans, ii.	190	Decoctum secalis cornuti, ii.	579	Devil's apple, i.	605
Decoctum pectorale in phthisi, ii.	347	Decoctum seminum delphinii staphisagriæ, i.	608	Devil's bit, ii.	559
Decoctum Petri Foresti, ii.	10	Decoctum senegæ, ii.	404	Deuto-acetate of copper, i.	595
Decoctum pini saponatum, i.	23	Decoctum simarubæ, ii.	587	Deuto-acetate of mercury, ii.	96
Decoctum polygalæ senegæ, ii.	404	Decoctum smilacis sarsaparillæ, ii.	594	Deutochloruret of mercury, ii.	66
Decoctum prunorum cum senna, i.	414	Decoctum spigeliæ, ii.	620	Deutohydrochloras antimonii, i.	254
Decoctum prunus padi, i.	439	Decoctum stomachicum resolvens, ii.	435	Deutohydrochloras stanni, ii.	625
Decoctum purgans antiphlogisticum, ii.	468	Decoctum sudorificum, i.	737	Deutohydrochlorate of copper, i.	558
Decoctum quassiae compositum, ii.	473	Decoctum sulphureti calcis Hoffmanni, i.	254	Deutohydrochlorate of copper and of ammonia, i.	589
Decoctum quercus, ii.	476	Decoctum syphiliticum roborans, ii.	136	Deuto-ioduret of mercury, ii.	79
Decoctum quercus roboris, ii.	476	Decoctum tabaci, ii.	261	Deutomurias stibi, ii.	254
Decoctum quinae antisepticum purgans, i.	496	Decoctum tamarindi compositum, ii.	647	Deutonnitrate of copper, i.	588
Decoctum de radice salep, ii.	541	Decoctum tamarindorum, ii.	646	Deutonnitrate of mercury, ii.	89
Decoctum radicis althææ, emolliens, i.	147	Decoctum taraxaci, ii.	651	Deutophosphate of copper, i.	583
Decoctum radicis caryophyllatæ, i.	719	Decoctum taraxaci compositum, ii.	652	Deutosulphas cupri et ammoniæ, i.	586
Decoctum radicis caryophyllatæ cum cascariilla, i.	720	Decoctum tartaricatum, ii.	452	Deutosulphuretum hydrargyri rubrum, ii.	43
Decoctum radicis caryophyllatæ cum rheo, i.	720	Decoctum turionum pini, i.	22	Deutoxycyanuret hydrated of iron, i.	674
Decoctum radicis consolida, ii.	642	Decoctum veratri, ii.	715	Deutoxide of antimony, i.	236
Decoctum radicis ratanhia, ii.	148	Decoctum verbasci andydnium, ii.	716	Deutoxide of hydrogen, ii.	328
Decoctum radicis tormen- tillæ, ii.	693	Decoctum ulmi campestris, ii.	700	Deutoxide of lead, ii.	370
		Delphia, i.	608	Deutoxydum plumbi rubrum, ii.	370
		Delphinium consolida, i.	608	Dewberry, ii.	532
		Delphinium staphisagriæ, i.	608	Diacaryon, ii.	135
		Dentalis, i.	353	Diadodium liquidum montani, ii.	339
				Diacydium, ii.	559
				Diacydium glycyrrhizatum, ii.	560
				Diacydium sulphuratum, ii.	560
				Diacydium jalapinum, i.	550

<i>Diacydinum lucidum</i> ja-		<i>Doronicum pardalian-</i>		<i>Dropwort</i> , ii.	621
<i>lapinum</i> , i.	550	<i>ches</i> , i.	618	<i>Drosera longifolia</i> , i.	619
<i>Diacydinum sollutivum</i> ,		<i>Doronicum plantagine-</i>		<i>Drosera rotundifolia</i> , i.	619
<i>pellucidum</i> , i.	550	<i>um</i> , i.	619	<i>Drymis aromatica</i> , i.	620
<i>Diacydonium aromati-</i>		<i>Douche aromatic</i> , i.	193	<i>Drymis winteri</i> , i.	620
<i>cum</i> , i.	598	<i>Dove's-foot</i> , i.	718	<i>Dryobanalops camphora</i> ,	
<i>Diacydonium lucidum</i> ,		<i>Dracæna draco</i> , i.	765	i.	366
i.	598	<i>Draco mitigatus</i> , ii.	46	<i>Dulcichinum</i> , i.	601
<i>Diacydonium simplex</i> ,		<i>Dracontia minoris radix</i> ,		<i>Dwale</i> , i.	309
i.	598	i.	291	<i>Dyers' broom</i> , i.	708
<i>Diacydonium cum spe-</i>		<i>Dracontium foetidum</i> , i.	619		
<i>ciebus</i> , i.	598	<i>Dracunculi esculenti</i>		<i>Eagle-stone</i> , i.	643
<i>Dianthus caryophyllus</i> ,		<i>herba</i> , i.	291	<i>Earth-worm</i> , i.	698
i.	610	<i>Dacunculi hortensis</i>		<i>Eau d'Anhalt</i> , ii.	666
<i>Diapensia herba</i> , ii.	556	<i>herba</i> , i.	291	<i>Eau d'Arquebusade</i> , i.	45
<i>Diaphoretic Keup's</i> , i.	236	<i>Dragon-root</i> , i.	292	ii.	547
<i>Diarrhodon cum rha-</i>		<i>Dragon's blood</i> , i.	764	<i>Eau des carmes</i> , ii.	230
<i>barbaro</i> , ii.	493	<i>Draught anti-emetic</i> , i.	50	<i>Eau cordiale de Coladon</i> ,	
<i>Diascordium</i> , ii.	317	<i>Draught antiscrofulous</i> ,		i.	519
<i>Diascordium liquidum</i> ,		i.	188	<i>Eau divine</i> , i.	518
ii.	684	<i>Draught diaphoretic</i> , i.	186	<i>Eau de Hongrie</i> , ii.	528
<i>Dictamni cretici folia</i> ,		<i>Draught sedative</i> , ii.	284	<i>Eau de Javelle</i> , ii.	602
ii.	322	<i>Draught stimulant</i> , ii.	272	<i>Eau saline fondant de</i>	
<i>Dictamnus albus</i> , i.	611	<i>Draught sudorific</i> , i.	197	<i>Lassine</i> , ii.	454
<i>Dictamnus of Candia</i> ,		<i>Drink anticatarrhal</i> , ii.	114,	<i>Eau de luce</i> , i.	180
ii.	322		482	<i>Eau medicinale de Hus-</i>	
<i>Diet-drink Lisbon</i> , ii.	598	<i>Drink antidysenteric</i> , ii.	423	<i>son</i> , i.	538
<i>Digestive Plenk's</i> , ii.	659	<i>Drink antiplagistic</i> , ii.	435	<i>Eau sans pareil</i> , i.	518
<i>Digestivum liquefactum</i> ,		<i>Drink antirachitic</i> , ii.	531	<i>Eau de vie des femmes</i> ,	
ii.	658	<i>Drink antiscrofulous</i> , i.	490	ii.	169
<i>Digitaline</i> , i.	612	<i>Drink antispasmodic</i> , i.	452	<i>Ebuli radix</i> , ii.	550
<i>Digitalis ferruginea</i> , i.	611	ii.	707	<i>Ebur</i> , i.	360
<i>Digitalis purpurea</i> , i.	611	<i>Drink astringent</i> , ii.	642	<i>Echinus</i> , i.	354
ii.	555	<i>Drink bread</i> , ii.	332	<i>Echioglossi herba</i> , ii.	274
<i>Dill</i> , i.	220	<i>Drink of cream of tartar</i> ,		<i>Echium vulgare</i> , i.	620
<i>Diospyros ebenum</i> , ii.	194	ii.	452	<i>Eclegma album</i> , i.	210
<i>Diospyros Virginiana</i> ,		<i>Drink ferruginous</i> , i.	653	<i>Eclygma diaphoretic</i> , i.	240
i.	619	<i>Drink imperial</i> , ii.	453	<i>Eel's liver</i> , i.	699
<i>Dipsacus sylvestris</i> , i.	618	<i>Drink laxative</i> , ii.	448,	<i>Egg-plant</i> , ii.	617
<i>Dipsacus fullonum</i> , i.	618	<i>Drink Milman's</i> , ii.	452	<i>Egg pullet's</i> , ii.	326
<i>Diptamni officinalis ra-</i>		<i>Drink nitrous</i> , i.	70	<i>Elæocarpus copallifera</i> ,	
<i>dix</i> , i.	611	<i>Drink purgative</i> , ii.	449	ii.	755
<i>Dittany of crete</i> , ii.	322	<i>Drink refrigerant and</i>		<i>Elais Guineensis</i> , ii.	271
<i>Dittany white</i> , i.	611	<i>diuretic</i> , ii.	438	<i>Elaterin</i> , ii.	243
<i>Dock bloody</i> , ii.	535	<i>Drink tartro-borated</i> ,		<i>Elaterinum</i> , ii.	243
<i>Dock curled</i> , ii.	534	ii.	454	<i>Elaterium</i> , ii.	243
<i>Dock red</i> , ii.	535	<i>Drop black</i> , ii.	275	<i>Elatin</i> , ii.	243
<i>Dock water</i> , ii.	534	<i>Dropax antiparalyticum</i> ,		<i>Elder</i> , ii.	551
<i>Dodder</i> , i.	597	ii.	364	<i>Elder dwarf</i> , ii.	550
<i>Dog-grass</i> , ii.	696	<i>Drops anthelmintic</i> , ii.	136	<i>Elecampane</i> , ii.	114
<i>Dogsbane</i> , i.	270	<i>Drops anti-hysteric</i> , i.	306	<i>Electuarium de acoro</i> , i.	86
<i>Dog-wood tree</i> , i.	560	<i>Drops bitter</i> , i.	267	<i>Electuarium adstringens</i>	
<i>Dog-wood swamp tree</i> ,		<i>Drops English cephalic</i> ,		<i>Fulleri</i> , ii.	512
i.	560	i.	187	<i>Electuarium alkalino-</i>	
<i>Dolichos pruriens</i> , ii.	625	<i>Drops chalybeate</i> , i.	656	<i>feratum</i> , i.	643
<i>Doronic Germanici ra-</i>		<i>Drops diuretic</i> , ii.	262	<i>Electuarium aloes com-</i>	
<i>dix</i> , i.	279	<i>Drops General la Motte's</i> ,		<i>positum</i> , i.	129
<i>Doronici latifolii radix</i> ,		i.	314	<i>Electuarium anodynum</i> ,	
i.	618	<i>Drops Jena's</i> , i.	139	<i>pro infantibus</i> , ii.	209
<i>Doronici officinalis ra-</i>		<i>Drops phosphoric</i> , i.	73	<i>Electuarium antacidum</i> ,	
<i>dix</i> , i.	618	<i>Drops sedative</i> , ii.	246	ii.	209
<i>Doronici plantaginis fo-</i>		<i>Drops stimulant</i> , ii.	270	<i>Electuarium anthelmin-</i>	
<i>lii</i> , i.	279	<i>Drops Ward's white</i> , ii.	89	<i>ticum</i> , ii.	705
<i>Doronici Romani radix</i> ,		<i>Drops Warner's cordial</i> ,		<i>Electuarium anti-epilep-</i>	
i.	618	i.	411	<i>ticum</i> , i.	479

Electuarium antispasmodicum, ii.	479	Electuarium de junipero, ii.	138	Electuarium theriacale magnum, ii.	306
Electuarium aperiens, i.	404	Electuarium laxans et emmenagogum, i.	122	Electuarium tonico-resolvens, i.	643
Electuarium aromaticopiatum, ii.	306	Electuarium laxativum, i.	422	Electuarium ad vermes, i.	546
Electuarium aromaticum, ii.	164	Electuarium lenitivum, i.	404	Electuary alternative, i.	407,
Electuarium boleti suaveolentis, i.	602	Electuarium lenitivum cum sulphure, i.	407	741, ii.	29, 32
Electuarium cardiacum, ii.	164	Electuarium de manna, ii.	218	Electuary antacid, ii.	209
Electuarium cassiæ fistulæ, i.	422	Electuarium mimosæ catechu compositum, ii.	317	Electuary anthelmintic, i.	546, ii.
Electuarium cassiæ tamarindatum, i.	422	Electuarium mithridaticum, ii.	209, 304	331, 583, 705	
Electuarium catechu compositum, ii.	318	Electuarium mundificans, i.	407, 741	Electuary antichlorotic, i.	650
Electuarium catholicum, i.	407	Electuarium opiatum, ii.	306	Electuary anti-apoplectic, i.	650
Electuarium catholicum commune, i.	404	Electuarium opii compositum, ii.	306	Electuary anti-arthritic, ii.	45, 142
Electuarium catholicum duplicatum, i.	407	Electuarium pro pecoribus, i.	761	Electuary anticardialgic, i.	477
Electuarium ceratum, i.	434	Electuarium piperinum, ii.	363	Electuary anticachectic, ii.	363
Electuarium e chamæmelo, i.	446	Electuarium polygalæ amaræ, ii.	403	Electuary antidysenteric, i.	229, ii.
Electuarium cochleariæ cum calamo aromatico, i.	531	Electuarium purgans, i.	545, ii.	125, 492	
Electuarium cordiale, i.	565	Electuarium requies Nicolai, ii.	320	Electuary anti-epileptic, i.	439, 479, ii.
Electuarium demulcens, i.	434	Electuarium resolvens antihæpeticum, ii.	26	331	
Electuarium dentifricium, ii.	451	Electuarium resolvens martiale diureticum, i.	643	Electuary antifebrile, i.	477
Electuarium deobstruens, ii.	567	Electuarium resumptivum, ii.	699	Electuary anti-gonorrhœic, i.	557
Electuarium diacatholicon, i.	407	Electuarium de rhabarbaro, i.	407	Electuary antihæmorrhoidal, i.	406, ii.
Electuarium diaphœnicum, ii.	562	Electuarium sanans, i.	434	219	
Electuarium diaprunum, i.	404	Electuarium scammonii, ii.	561	Electuary antihæmoptysic, ii.	100
Electuarium diastyron, ii.	541	Electuarium scordii, ii.	317	Electuary antihæpetic, i.	262
Electuarium diascordium, ii.	317	Electuarium scordii opiatum, ii.	317	266, 546, ii.	26, 32, 74
Electuarium diatragacanthæ, i.	767	Electuarium e scordio, ii.	317	Electuary antihydrophobic, ii.	135
Electuarium eccoprocticum, i.	404	Electuarium e senna, i.	404	Electuary antilyssic, ii.	232
Electuarium Fracastoris, ii.	317	Electuarium sennæ, i.	404	Electuary antiperiodic and tonic, i.	240
Electuarium gingivale, i.	761	Electuarium sennæ cum carbone, i.	407	Electuary antiphthisical, ii.	636
Electuarium hieræ picræ, i.	129	Electuarium sennæ compositum, i.	404	Electuary antirheumatic, ii.	636
Electuarium hydragogum, i.	545	Electuarium sennæ cum pulpis, i.	404	Electuary antiscorbutic, i.	531, ii.
Electuarium hyoscyami opiatum, ii.	320	Electuarium stomachicum, ii.	492	588	
Electuarium ipecacuanhæ moschatum, ii.	126	Electuarium stomachicum alterans, ii.	363	Electuary antiscrofulous, ii.	444, 445
Electuarium joviale ad tæniam, ii.	624	Electuarium tamarindorum, ii.	648	Electuary antisisitic, i.	622
		Electuarium terebinthinatum, ii.	668	Electuary antispasmodic, ii.	705
		Electuarium thebaicum, ii.	306	Electuary antisymphilitic, ii.	58, 76
				Electuary aperitive and purgative, i.	650
				Electuary aphrodisiac, i.	531
				Electuary Arabian, ii.	541
				Electuary aromatic, i.	593
				Electuary astringent, i.	446
				478, 584, ii.	66, 146, 148,
				206, 315, 318, 477, 480,	512
				Electuary of bayberries, ii.	182

- Electuary of carburet of iron, i. 639
- Electuary caryocostine, ii. 561
- Electuary chalybeate, i. 638, 650
- Electuary of charcoal, i. 391
- Electuary of cinchona, i. 478
- Electuary alkaline of cinchona, i. 477
- Electuary antimoniated of cinchona, i. 479
- Electuary cordial, i. 565
- Electuary dentifrice, i. 358, 361, ii. 451
- Electuary deobstruent, ii. 567
- Electuary detersive, i. 760
- Electuary diaphœnic, ii. 562
- Electuary diatartaric, i. 406
- Electuary diuretic, ii. 138, 568
- Electuary Duhaume's astringent, i. 161
- Electuary dysenteric, ii. 315
- Electuary emetic, ii. 126
- Electuary emmenagogue, i. 639, 650
- Electuary febrifuge, i. 32, 90, 278, 476, 479
- Electuary febrifuge and aperitive, ii. 445
- Electuary febrifuge and incisive, i. 479
- Electuary alkaline ferruginous, i. 643
- Electuary of fragrant boletus, i. 602
- Electuary gingival, i. 761
- Electuary Heister's anthelmintic, ii. 28
- Electuary of hiera picra, i. 129
- Electuary hydragogue, i. 550, 650, ii. 561
- Electuary incrassant, ii. 192, 337
- Electuary laxative, i. 122, 648
- Electuary laxative and pectoral, i. 550
- Electuary of leopard's-bane, i. 279
- Electuary of manna, ii. 218
- Electuary mesenteric, i. 639
- Electuary compound of myrrh, i. 761
- Electuary Neapolitan, ii. 62
- Electuary opiate, ii. 306
- Electuary Pajot-Laforest's, ii. 74
- Electuary pectoral, i. 557, ii. 204, 219, 403
- Electuary tonic pectoral, i. 638
- Electuary purgative, i. 407, 545, ii. 45, 51, 452, 636
- Electuary purgative and alterative, ii. 29
- Electuary purgative and febrifuge, i. 477
- Electuary purgative and tonic, ii. 58
- Electuary refrigerant, ii. 452
- Electuary resolvens, i. 643
- Electuary resumptive, ii. 699
- Electuary Roccamore's, i. 507
- Electuary compound of scammony, ii. 561
- Electuary sedative, ii. 388
- Electuary Senac's febrifuge, i. 478
- Electuary compound of senna, i. 407
- Electuary stimulant, i. 231, 546
- Electuary stomachic, i. 86, 204, 507, ii. 363, 492
- Electuary of sulphur, ii. 636
- Electuary of supertartrate, ii. 450
- Electuary of sweetflag, i. 86
- Electuary temperant, ii. 434
- Electuary terebinthinate, ii. 668
- Electuary tonic, i. 477, 767, 643
- Electuary traumatic, i. 332, ii. 512
- Electuary vermifuge, ii. 29, 66, 623, 626
- Electuary for ulcerations of the mouth, i. 161
- Electuary Vogler's anthelmintic, ii. 66
- Electuary of wax, i. 434
- Electuary Werlhof's alterative, i. 741
- Elemi, ii. 486
- Eleminthocorton, ii. 2
- Elichrysi montani herba, i. 230
- Elixir acido-aromaticum, ii. 175
- Elixir acidum antinephreticum, i. 79
- Elixir acidum antipodragicum, i. 79
- Elixir acidum Dippelii, i. 79
- Elixir acidum Halleri, i. 79
- Elixir aloes saponaceum, i. 133
- Elixir amarum, ii. 449
- Elixir amarum aquosum, i. 30, 510, 516
- Elixir amarum vinosum, i. 713, 715
- Elixir ammoniato-opiatum, i. 729
- Elixir anthelminticum Succorum, i. 552
- Elixir anti-arthriticum caribæum, i. 735
- Elixir anticolicum crocatum, i. 130
- Elixir antihypochondriacum, i. 502
- Elixir antiscorbuticum amarum, i. 536
- Elixir aperiens, ii. 449
- Elixir aperitivum, i. 129
- Elixir aperitivum Claudi, i. 134
- Elixir cum aqua calcarie ustæ, ii. 473
- Elixir aromaticum acidum, ii. 175
- Elixir aurantiorum compositum, i. 513, 516
- Elixir balsamicum, i. 512
- Elixir balsamicum purificans, ii. 75
- Elixir balsamicum spirituosum Hoffmanni, ii. 685
- Elixir balsamicum, stomachicum, et temperatum Hoffmanni, i. 513
- Elixir citri purgans, i. 552
- Elixir Claudi's aperient, i. 134
- Elixir cordiale et stomachicum, i. 139
- Elixir Daffy, i. 412
- Elixir extracti hippocastani, i. 91
- Elixir foetidum, i. 429
- Elixir glycyrrhizæ, i. 729
- Elixir guaiaci volatilis, i. 736
- Elixir guaiacinum, i. 736
- Elixir guaiacinum volatile, i. 736
- Elixir ad hepar, i. 133
- Elixir jalapæ compositum, i. 552
- Elixir juniperinum, ii. 138
- Elixir longæ vitæ, i. 138
- Elixir malvaticum juniperinum, i. 138
- Elixir mastichinum, ii. 224
- Elixir myrrhæ, i. 763
- Elixir papaveris, ii. 277
- Elixir paregoricum, ii. 287
- Elixir pectorale dulce, i. 729
- Elixir pectorale regis Daniæ, i. 729
- Elixir pectorale Ringlemanni, i. 729
- Elixir polychrestum Hallense, ii. 562
- Elixir polychrestum purgativum, i. 412
- Elixir proprietatis, i. 129

Elixir proprietatis sine acido, i.	131	Elixir anti-arthritic, i.	129, 290, 736	Elixir resolvent, ii.	446
Elixir proprietatis cum acido Boerhaavii, i.	132	Elixir antifebrile, i.	503	Elixir Ringlemann's, i.	729
Elixir proprietatis cum acido Paracelsi, i.	132	Elixir antihysteria, ii.	119	Elixir saponaceous aloetic, i.	133
Elixir proprietatis album Helmontii, i.	130	Elixir antiscorbutic, i.	536	Elixir stomachic, i.	27, 226, 320, 501
Elixir proprietatis aquosum, i.	134	Elixir Boerhaave's antiscorbutic, i.	535	Elixir Stoughton's, i.	139
Elixir proprietatis dulce, i.	131	Elixir Drawitz's antiscorbutic, i.	535	Elixir Lettsom's sudorific, ii.	288
Elixir proprietatis cum rheo, i.	139	Elixir Selle's antiscorbutic, i.	535	Elixir Willis' sudorific, ii.	130
Elixir proprietatis Stahl, i.	133	Elixir antiseptic, i.	503	Elixir tonic, i.	138, 661
Elixir resolvens, ii.	449, 653	Elixir antispasmodic, ii.	101, 710	Elixir tonic acid, i.	570
Elixir rhei amarum, ii.	500	Elixir antivenereal, i.	555	Elixir Regimbeau's vermifuge, i.	140
Elixir rhei compositum, ii.	500	Elixir aromatic, ii.	356	Elixir visceral, i.	516
Elixir rhei dulce, ii.	500	Elixir astringent, ii.	194	Elixir Rosenstein's visceral, ii.	498
Elixir roborans acidum, i.	570	Elixir Hoffmann's spirituous balsamic, ii.	685	Elixir Wright's, ii.	77
Elixir roborans Whyttii, ii.	502	Elixir Werlhof's balsamic, i.	499	Elk's-hoof, i.	678
Elixir sacrum, i.	138	Elixir bitter, i.	30, 133, 510, 516, 526, ii.	Elm red, ii.	700
Elixir salutis, i.	410	Elixir cholagogue, i.	133	Elm rough-leaved, ii.	700
Elixir scorbuticum, i.	536	Elixir Crollius' uterine, i.	429	Elm slippery, ii.	700
Elixir sennæ, i.	409	Elixir Daffy's, i.	412	Embrocation ammoniacal, i.	174, 187
Elixir de spina, i.	140	Elixir deobstruent, ii.	653	Embrocation for chilblains, i.	174, 321
Elixir stomachicum, i.	513, 714	Elixir Dubois', i.	715	Embrocation irritating, i.	382
Elixir stomachicus spirituosus, i.	502	Elixir expectorant, i.	320	Embrocation refrigerant, i.	102
Elixir ex succo glycyrrhizæ, i.	729	Elixir febrifuge, i.	91	Embrocation resolvent, i.	198
Elixir ex succo liquiritiæ, i.	729	Elixir Whytt's febrifuge, i.	502	Embrocation stimulant, i.	486
Elixir sudoriferum, ii.	288	Elixir alkaline of gamboge, i.	695	Emerald, i.	157
Elixir traumaticum, i.	331	Elixir Garus', i.	130	Emery, i.	155
Elixir viscerale, i.	27, 513	Elixir of juniper, ii.	138	Emetic tartar, i.	255
Elixir viscerale Hoffmanni, i.	513	Elixir of lemon, i.	519	Emetine, i.	620, ii.
Elixir viscerale Rosensteini, ii.	498	Elixir of mastic, ii.	224	Emetinum, i.	620
Elixir vitæ Matthioli, ii.	170	Elixir Matthioli's ii.	170	Empetri herba, ii.	5
Elixir vitæ mulierum, ii.	549	Elixir Mittie's, i.	205	Emplastrum adhæsivum, i.	706, ii.
Elixir vitrioli, ii.	175	Elixir of myrrh, i.	763	Emplastrum adhæsivum Woodstockii, i.	706
Elixir vitrioli aromaticum, ii.	175	Elixir compound of orange, ii.	516	Emplastrum adstringens, ii.	391
Elixir vitrioli Edinburgensium, ii.	175	Elixir Elzner's pectoral, ii.	119	Emplastrum album coc-tum, ii.	400
Elixir vitrioli Mynsichti, ii.	175	Elixir pectoral balsamic of tolu, i.	324	Emplastrum de Althæa, i.	151
Elixir vitrioli cum tinctura cinnamomi, ii.	175	Elixir simple pectoral, i.	728	Emplastrum ammoniaci, i.	747, ii.
Elixir uterinum, i.	307	Elixir Wedel's pectoral, ii.	572	Emplastrum ammoniaci cum hydrargyro, ii.	23
Elixir acid, i.	78	Elixir polychrestic, ii.	562	Emplastrum de ammoniacum mercurio, i.	23
Elixir alkermic, ii.	174	Elixir proprietatis without acid, i.	131	Emplastrum ammoniacogalbanicum, i.	685
Elixir Courcelle's American, ii.	119	Elixir proprietatis antiscorbutic, i.	133	Emplastrum Andreae a cruce, ii.	677
Elixir opiate ammoniacal, i.	729	Elixir proprietatis Boerhaave's, i.	132	Emplastrum anglicum, i.	706
Elixir antacid, i.	349, ii.	Elixir proprietatis Helmont's, i.	130	Emplastrum anticolicum, i.	690
Elixir anti-apoplectic, ii.	174	Elixir proprietatis Paracelsus', i.	132	Emplastrum antihystericum, i.	689
		Elixir purgative polychrestic, i.	412		
		Elixir Radcliff's, i.	139		
		Elixir Raulin's, i.	412		

Emplastrum antihysteri- cium, i.	747	Emplastrum conii macu- lati, i.	467	Emplastrum de galbano, i.	685
Emplastrum aromaticum, ii.	165	Emplastrum e conio, i.	467	Emplastrum de galbano crocatum, i.	687
Emplastrum assafoetidae, i.	689	Emplastrum corrosivum, ii.	72	Emplastrum ad ganglia, i.	749
Emplastrum assafoetidae compositum, i.	747	Emplastrum croci sativi compositum, i.	687	Emplastrum glutinosum, i.	706
Emplastrum attrahens, ii.	675	Emplastrum de crusta panis, ii.	644	Emplastrum gratia dei, ii.	657
Emplastrum Bechholzii, i.	690	Emplastrum cumini, i.	580	Emplastrum Guillermi servitoris, ii.	640
Emplastrum benedictum, ii.	401	Emplastrum e cumino, i.	580	Emplastrum cum gum- matibus, i.	685
Emplastrum ad bubones venereos, ii.	24	Emplastrum defensivum i.	645	Emplastrum gummatum, i.	685
Emplastrum calefaciens, i.	386	Emplastrum defensivum rubrum, i. 645, ii.	393	Emplastrum cum gummi, i.	685
Emplastrum cantharidis, i.	383	Emplastrum diachylum, ii.	389	Emplastrum gummi resi- nosum, i.	747
Emplastrum cantharidis perpetuum, i.	387	Emplastrum diachylum compositum, i.	685	Emplastrum e gummis re- sinosis, i.	685
Emplastrum cantharidis resinosum, i.	387	Emplastrum diachylum simplex, ii.	389	Emplastrum gummosum, i.	747
Emplastrum cantharidis vesicatoriae, i.	383	Emplastrum diasulphuris Rulandi, i.	761	Emplastrum ad hernias, ii.	642
Emplastrum cantharidis vesicatorium composi- tum, i.	387	Emplastrum diapalmæ, ii.	392	Emplastrum Hierneri, ii.	392
Emplastrum cantharidum, i.	383	Emplastrum diaphœni- con, ii.	392	Emplastrum hydrargyri, ii.	22
Emplastrum carbonatis plumbi ustum, ii.	401	Emplastrum diaphoretici cum Mynsichti, i.	687	Emplastrum hydrargyri cum conio, ii.	24
Emplastrum catagmati- cum, ii.	644	Emplastrum divinum, ii.	396	Emplastrum hydrargyri cum conio et opio, ii.	25
Emplastrum cephalicum, ii.	303	Emplastrum emolliens, i.	151	Emplastrum hydrargyri- cum, ii.	22
Emplastrum ceræ, ii.	675	Emplastrum epispasti- cum, i.	383	Emplastrum ex hydrar- gyro, ii.	22
Emplastrum ceræ tere- binthinatum, ii.	675	Emplastrum de euphor- bio, i.	756	Emplastrum hyoscyami, ii.	104
Emplastrum cereum, ii.	675	Emplastrum ferri, ii.	393	Emplastrum de hyoscy- amo, ii.	104
Emplastrum cerussæ, ii.	400	Emplastrum filii, i.	684	Emplastrum ichthyocollæ telæ inductum, i.	706
Emplastrum cetacei, i.	441	Emplastrum filii Zac- chariæ, i.	436	Emplastrum incognitum, ii.	196
Emplastrum de cicuta, i.	467	Emplastrum flavum, i.	151	Emplastrum ischiadicum, i.	690, 756
Emplastrum de cicuta cum ammoniaco, i.	467	Emplastrum e fœnugræco, i.	748	Emplastrum ladani, i.	758
Emplastrum de cicuta compositum, i.	467	Emplastrum fœtidum, i.	689, 747	Emplastrum lithargyri, ii.	389
Emplastrum cicutæ, i.	467	Emplastrum ad fonticu- los, i. 435, ii.	371	Emplastrum lithargyri- cum cum resina pini, ii.	679
Emplastrum citrinum, i.	151, 675	Emplastrum ad fracturas, ii.	644	Emplastrum ad lupiam, i.	595
Emplastrum ad clavos pedum, i.	594	Emplastrum frigidum Riedlini, i.	153	Emplastrum lythargyri compositum, i.	685
Emplastrum collæ ad her- niam, ii.	395	Emplastrum ad fungos articulorum, i.	757	Emplastrum lythargyri- cum resinae, ii.	679
Emplastrum comfortati- vum Vigonis, ii.	393	Emplastrum galbani, i.	685	Emplastrum lyttæ vesica- toriae, i.	383
Emplastrum commune, ii.	389	Emplastrum galbani cam- phoratum, i.	684	Emplastrum malacticum cum gummi, i.	153
Emplastrum commune ad- hæsivum, ii.	679	Emplastrum galbani com- positum, i.	685		
Emplastrum commune gummosum, i.	685	Emplastrum galbani robo- rans, i.	689		
Emplastrum commune cum resina, ii.	679	Emplastrum galbani satur- ninum, i.	685		
Emplastrum conii, i.	467	Emplastrum galbani sim- plex, i.	684		

Emplastrum malacticum sine gummi, i.	153	Emplastrum piperitum, ii.	364	Emplastrum simplex, ii.	390, 675
Emplastrum mamillare, i. 443, ii.	463	Emplastrum plumbi, ii.	389	Emplastrum de spermate ceti compositum, i.	684
Emplastrum ad mam- millas, i.	443	Emplastrum plumbi ni- grum, ii.	369	Emplastrum de spermate ranarum, ii.	401
Emplastrum matricale, i.	689	Emplastrum plumbi sub- carbonatis compositum, ii.	401	Emplastrum spermatis ceti, i.	441
Emplastrum maturans, ii.	677	Emplastrum e plumbo compositum, ii.	400	Emplastrum sticticum, ii.	393, 675
Emplastrum meliloti, ii.	226	Emplastrum polychres- tum, ii.	679	Emplastrum stomachi- cum, i. 758, ii.	315, 644
Emplastrum mercuriale, ii.	22	Emplastrum ranarum sim- plex, ii.	396	Emplastrum stypticum, i. 645, ii. 393,	680
Emplastrum mercuriale resolvens, ii.	22	Emplastrum de ranis sine mercurio, ii.	396	Emplastrum styracis sto- machicum, ii.	627
Emplastrum minii ad us- tum, ii.	399	Emplastrum e resina elas- tica, i.	755	Emplastrum subcarbo- natis plumbi, ii.	400
Emplastrum minii cam- phoratum, ii.	399	Emplastrum resinae, ii.	679	Emplastrum sulphura- tum, i.	690, 761
Emplastrum moschatum, ii.	250	Emplastrum resinarum glutinans, ii.	677	Emplastrum sulphura- tum resinosum, ii.	640
Emplastrum de mucila- ginibus, i.	748	Emplastrum resinosum, ii.	679	Emplastrum tacamah- acinum, ii.	644
Emplastrum mucilagi- nosum, i.	151	Emplastrum resinosum cantharidum, i.	383	Emplastrum ad tempo- ra, ii.	302
Emplastrum de nicoti- ana, ii.	264	Emplastrum resolvens cum sapone et cam- phora, ii.	464	Emplastrum ad tormina ventris, ii.	302
Emplastrum nigrum Bech- holzii, i.	690	Emplastrum resolvens, i.	468, 747	Emplastrum trifolii mel- iloti officinalis, ii.	226
Emplastrum nigrum sul- phuratum, i.	690	Emplastrum resolvens compositum, ii.	24	Emplastrum tripharma- cum, ii.	399
Emplastrum Noribergense, ii.	399	Emplastrum resolvens cum mercurio, ii.	24	Emplastrum ad verru- cas, i.	594
Emplastrum opiatum, ii.	303	Emplastrum resolutivum, ii.	397	Emplastrum vesicato- rium, i.	383
Emplastrum opii, ii.	297	Emplastrum rhei com- positum, i.	142	Emplastrum vesicato- rium perpetuum, i.	387
Emplastrum oxycroceum, i.	687	Emplastrum roborans, i. 645, ii. 391, 393, 675, 680		Emplastrum Vigonis sine mercurio, ii.	396
Emplastrum oxydi ferri rubri, i. 645, ii.	393	Emplastrum rubefaciens, i.	756	Emplastrum viride, i.	593
Emplastrum oxydi plum- bi diabatanum, ii.	397	Emplastrum rubefians, i.	383	Emplastrum viride com- positum, i.	593
Emplastrum oxydi plum- bi epuloticum, ii.	391	Emplastrum santalinum, ii.	196	Emulsio violata, ii.	722
Emplastrum oxydi plumbi mucilagineum, ii.	389	Emplastrum saponaceo- plumbicum, ii.	392	Emulsio acaciæ arabicæ, i.	207
Emplastrum oxydi plumbi rubrum, ii.	393	Emplastrum saponaceum, ii.	464	Emulsio amygdalina, i.	207
Emplastrum oxydi plumbi semi-vitrei, ii.	389	Emplastrum saponatum, ii.	463	Emulsio amygdalarum composita, ii.	210
Emplastrum oxydi plumbi semi-vitrei gummosum, i.	685	Emplastrum de sapone, ii.	463	Emulsio antihysterica, i.	303
Emplastrum oxydi plumbi semi-vitrei cum sulphate ferri calcinato, ii.	393	Emplastrum saponis, ii.	463	Emulsio arabica, i.	207
Emplastrum piceum, ii.	675	Emplastrum saponis cam- phoratum, ii.	464	Emulsio balsami copai- væ, i.	556
Emplastrum piceum com- positum, ii.	677	Emplastrum de scrofula- ria, i.	593	Emulsio balsamica, i.	323
Emplastrum picis com- positum, ii.	677	Emplastrum sebi anody- num, ii.	391	Emulsio cajeputi, ii.	270
Emplastrum picis nava- lis, ii.	675	Emplastrum siliquarum, ii.	393	Emulsio camphoræ, i.	370
Emplastrum picis nigrae regium, ii.	642	Emplastrum siliquarum imperiale, ii.	395	Emulsio camphorata, i.	370
Emplastrum picis, ii.	675			Emulsio cerea, i.	433
Emplastrum picis liqui- dæ, ii.	366			Emulsio cornu cervi usti, i.	752
				Emulsio externa e nuc- leis, i.	345
				Emulsio ferri composita, i.	647

- Emulsio guaiacina, i. 739
 Emulsio gummi resinæ ammoniaci, i. 742
 Emulsio gummi resinæ guaiaci, i. 739
 Emulsio hordeata amygdalina, i. 207
 Emulsio leniens, i. 208, 211
 Emulsio cum nuce vomica, ii. 267
 Emulsio phosphorata, ii. 348
 Emulsio pini, i. 23
 Emulsio sedativa, ii. 102
 Emulsio sedativa simplex, ii. 247
 Emulsio terebinthinacea, ii. 655
 Emulsio almond, i. 207
 Emulsio ammoniacal, i. 187
 Emulsion anthelmintic, ii. 322
 Emulsion anticatarrhal, ii. 346
 Emulsion astringent, i. 157, ii. 477
 Emulsion of anisated balsam of sulphur, ii. 638
 Emulsion balsamic, i. 323
 Emulsion, Barthez, ii. 435
 Emulsion Brunner's, i. 744
 Emulsion of camphor, i. 370
 Emulsion camphorated antimonial, i. 262
 Emulsion camphorated and nitrated, i. 371
 Emulsion camphorated nitrous, ii. 437
 Emulsion of cantharides, i. 382, 388
 Emulsion of cinchona, i. 487
 Emulsion cooling, i. 208
 Emulsion cosmetic, i. 320
 Emulsion demulcent, i. 212
 Emulsion of deutochloruret of mercury, ii. 72
 Emulsion diuretic, i. 378, 744
 Emulsion emetic, i. 248
 Emulsion febrifuge, i. 627
 Emulsion of fir-cone kernels, ii. 358
 Emulsion of guaiac, i. 739
 Emulsion of hempseed, i. 376
 Emulsion kermeticized, i. 248
 Emulsion of lemon pips, i. 521
 Emulsion against lum-bago, ii. 424
 Emulsion mucilaginous, i. 754
 Emulsion narcotic, ii. 102, 280
 Emulsion nitrous, ii. 435
 Emulsion oily, ii. 270
 Emulsion pectoral, i. 443, ii. 341
 Emulsion phosphorated, ii. 348
 Emulsion pistachio, ii. 364
 Emulsion purgative, i. 546, 550, ii. 220, 507, 563
 Emulsion refreshing, ii. 430
 Emulsion refrigerant, ii. 436
 Emulsion saline, ii. 448
 Emulsion sedative, i. 97, ii. 247, 337, 345
 Emulsion soothing, i. 577
 Emulsion of spruce fir, i. 23
 Emulsion stimulant, ii. 270
 Emulsion terebinthinated, ii. 668
 Emulsion of turpentine, ii. 655
 Emulsion vermifuge, i. 64, ii. 563
 Emulsion of violet, ii. 722
 Emulsion of wax, i. 433
 Emys lutaria, i. 697
 Endive, i. 461
 Endivæ herba, i. 461
 Enema cum acido sulphurico, i. 77
 Enema antispasmodicum, i. 371
 Enema antispasticum, ii. 102
 Enema ad ascarides, ii. 650
 Enema camphoratum, i. 303
 Enema catharticum, i. 118, ii. 212
 Enema chinæ cum opio, i. 484
 Enema foetidum, i. 303
 Enema hyosciami, ii. 102
 Enema mannatum, ii. 60
 Enema mercuriale, ii. 49, 60
 Enema purgans, ii. 212
 Enema sabadillæ, ii. 714
 Ens martis, i. 661
 Ens veneris Boylei, i. 661
 Entalia, i. 353
 Enulæ radix, ii. 115
 Enulæ campanæ radix, ii. 115
 Epithem antipodagric, ii. 288
 Epithem vermifuge, i. 121
 Epithymi cretici herba, i. 597
 Epithymi officinarum herba, i. 597
 Equisetum arvense, i. 621
 Equisetum fluviatile, i. 621
 Equisetum hyemale, i. 621
 Equisetum palustre, i. 621
 Erica herba, i. 366
 Erigeri herba, ii. 584
 Erigeron acre, i. 621
 Erigeron Canadense, i. 622
 Erigeron Philadelphicum, i. 622
 Erigeron viscosum, i. 622
 Erinaceus combustus, i. 391
 Erinaceus Europæus, i. 355
 Erioxylon, i. 730
 Errhine astringent, ii. 370
 Errhine for ozæna, i. 222
 Eruca semen, ii. 588
 Eruca nigræ semen, ii. 588
 Ervi semen, ii. 719
 Ervum lens, i. 631
 Eryngium campestre, i. 622
 Eryngo common, i. 622
 Eryngo sea, i. 622
 Eryngo water, i. 622
 Eryngium aquaticum, i. 622
 Eryngium maritimum, i. 622
 Erysimum alliaria, i. 623
 Erysimum barbarea, i. 623
 Erysimum officinale, i. 623
 Erythraea centaurinum, i. 625
 Erythraea chilensis, i. 628
 Erythronium flavescens, i. 628
 Esox lucius, 360
 Esprit aromatique, ii. 528
 Essence Stahl's alexipharmic, ii. 684
 Essence Stahl's temperate alexipharmic, ii. 684
 Essence anodyne, ii. 284
 Essence Ludwig's anodyne, ii. 285
 Essence anti-arthritis, i. 429
 Essence antihydrotic, i. 126
 Essence anhyseric, i. 306, 430
 Essence opiate arthritis, i. 429
 Essence compound of balm, ii. 228
 Essence balsamic, i. 520
 Essence Halle's bezoardic, i. 226
 Essence bitter, i. 626, 715
 Essence carminative, i. 516
 Essence carminative of castor, i. 431
 Essence cephalic, i. 88
 Essence Hellwig's of cinchona, i. 501
 Essence Helvetius' emetic, i. 235
 Essence of ground ivy, i. 722
 Essence dry of lemon, i. 517
 Essence of liquorice, i. 729
 Essence of malt, ii. 214
 Essence dry of orange, i. 507
 Essence vulnerary of poplar, ii. 409
 Essence of poppy, ii. 277
 Essence of rosemary, ii. 523
 Essence Keump's scillitic, ii. 445

Essence compound of scordium, ii.	684	Essentia cardamomi, i.	394	Essentia millefolii, i.	36
Essence of soot, i.	199	Essentia cardui benedicti, i.	525	Essentia moschi, ii.	252
Essence Stahl's splenetic, ii.	108	Essentia carminativa castorei, i.	431	Essentia alcholica myrrhæ, i.	762
Essence theriacal, ii.	314	Essentia carminativa Fulleri, i.	516	Essentia myrrhæ alkalisata, i.	762, 763
Essence traumatic, ii.	693	Essentia castorei, i.	428	Essentia pimpinellæ albæ, ii.	357
Essence vermifuge, ii.	650	Essentia castorei thebaica, i.	429	Essentia pini, i.	22
Essentia absinthii, i.	28	Essentia catholica purgans Rothii, i.	552	Essentia piperis, ii.	360
Essentia absinthii amara, i.	29	Essentia de cedro, i.	517	Essentia promorum aurantium immaturorum, i.	515
Essentia absinthii totius, i.	29	Essentia centaurei minoris, i.	626	Essentia oculorum populi, ii.	409
Essentia aconiti, i.	85	Essentia cephalica, i.	88	Essentia populi vulneraria, ii.	409
Essentia alexipharmaca Stahl, ii.	684	Essentia chinæ, i.	501	Essentia rorismarini, ii.	523
Essentia alexipharmaca temperata Stahl, ii.	684	Essentia cinchonæ, i.	500	Essentia sabinæ, ii.	143
Essentia aloes, i.	120	Essentia cinnamomi, ii.	172	Essentia scordii, ii.	684
Essentia aloes composita, i.	133	Essentia cochleariæ, i.	534	Essentia de scordio composita, ii.	684
Essentia amara, i.	715	Essentia corticis Peruviani alexipharmaca Huxhami, i.	501	Essentia splenetica Stahl, ii.	108
Essentia amara Halensis, i.	30	Essentia corticis Peruviani composita, i.	502	Essentia stomachica, i.	513
Essentia ambraë Hoffmanni, i.	167	Essentia corticis viridis nucum juglandis, ii.	136	Essentia stomachica polychresta Gundelsheimeri, ii.	361
Essentia ambraë liquida, i.	167	Essentia corticum aurantium, i.	515	Essentia stramonii, i.	606
Essentia ambraë sicca, i.	166	Essentia corticum aurantium vinosa, i.	511	Essentia succini alcoholica, ii.	629
Essentia Augusturæ, i.	229	Essentia croci, i.	564	Essentia succini alkalisata, ii.	630
Essentia anodyna officinalis, ii.	284	Essentia culilaban, ii.	178	Essentia terebinthinæ, ii.	655
Essentia anodyna Ludovici, ii.	285	Essentia digitalis, i.	616	Essentia theriacalis, ii.	314
Essentia antarthritica Hoffmanni cum opio, i.	429	Essentia dulcis, ii.	444	Essentia traumatica, ii.	693
Essentia arthritica Hoffmanni sine opio, i.	429	Essentia fuliginis Clauderi, i.	199	Essentia valerianæ, ii.	708
Essentia antihypropica Hoffmanni, i.	126	Essentia galbani, i.	684	Essentia valerianæ composita, ii.	708
Essentia antihysterica, i.	306	Essentia gentianæ, i.	714	Essentia valerianæ volatilis, ii.	109
Essentia antimonii, i.	258	Essentia glechomæ hederaceæ, i.	722	Essentia contra vermes, ii.	650
Essentia antiscorbutica, ii.	177	Essentia guaiaci, i.	734	Essentia vulneraria Stahl, ii.	665
Essentia antiseptica Huxhami, i.	501	Essentia gummi ammoniaci, i.	744	Esulæ minoris radix, i.	629
Essentia aromatica, ii.	173	Essentia hæmato-cathartica et alexipharmaca Clauderi, i.	199	Eucalyptus resinifera, ii.	146
Essentia assafoetida, i.	305	Essentia hyoscyami nigri, ii.	102	Euchloras potassæ, ii.	440
Essentia assafoetidæ æthereæ, i.	305	Essentia hyperici, ii.	108	Everlasting flower, ii.	4
Essentia aurantii dulcis, i.	515	Essentia lactucæ virosæ, ii.	153	Euonymus Europæus, i.	630
Essentia balsami Canadensis mundificans, ii.	75	Essentia levistici, ii.	197	Eupatorii mesues herba, i.	35
Essentia balsami Canadensis roborans, i.	318	Essentia ligni aloes, ii.	192	Eupatorii veterum herba, i.	103
Essentia balsami copaivæ kalica, i.	555	Essentia ligni sassafras, ii.	183	Eupatorium aya-pana, i.	628
Essentia balsami Peruviani, i.	320	Essentia ligni sassafras composita, ii.	184	Eupatorium cannabinum, i.	628
Essentia benzoini, i.	330	Essentia lignorum, i.	735	Eupatorium common, i.	268
Essentia calami aromatici, i.	87	Essentia liquiritiæ, i.	729	Eupatorium perfoliatum, i.	628
Essentia cantharides vesicatoriæ, i.	379	Essentia macis, ii.	255	Eupatorium purple, i.	628
		Essentia malti, ii.	214	Eupatorium purpureum, i.	682
		Essentia mastiches, ii.	224	Eupatorium teucrifolium, i.	628
		Essentia menthæ piperritæ, ii.	236	Euphorbia canariensis, i.	756
				Euphorbia corollata, i.	629

Euphorbia cyparissias, i.	629	Extract of great celandine, i.	445	Extract vinous of guaiac, i.	733
Euphorbia helioscopia, i.	629	Extract alcoholic of lesser centaury, i.	627	Extract of heart's-ease, ii.	724
Euphorbia ipecacuanha, i.	629	Extract aqueous of lesser centaury, i.	626	Extract of hemlock, i.	464
Euphorbia lathyris, i.	629	Extract of chamomile, i.	450	Extract alcoholic of henbane, ii.	102
Euphorbia officinarum, i.	756	Extract of cherry-laurel, i.	440	Extract aqueous of henbane, ii.	100
Euphorbia palustris, i.	629	Extract of chervil, i.	445	Extract alcoholic of hops, ii.	11
Euphorbia peplus, i.	629	Extract of scented chervil, i.	445	Extract aqueous of hops, ii.	16
Euphorbia sylvatica, i.	629	Extract alcoholic of cinchona, i.	482	Extract alcoholic of horehound, ii.	222
Euphorbium, i.	756	Extract aqueous of cinchona, i.	480	Extract aqueous of horehound, ii.	222
Euphrasia officinalis, i.	630	Extract emulsive of cinchona, i.	482	Extract alcoholic of horse-chesnut, i.	91
Excacaria agallocha, ii.	192	Extract hydro-alcoholic of cinchona, i.	482	Extract aqueous of horse-chesnut, i.	90
Excrement white of dogs, i.	360	Extract vinous of cinchona, i.	482	Extract alcoholic of jalap, i.	547
Extract of aconite, i.	84	Extract of cichory, i.	462	Extract aqueous of jalap, i.	547
Extract alcoholic of agaric, ii.	408	Extract of clematis, i.	523	Extract gummy of jalap, i.	547
Extract aqueous of agaric, ii.	408	Extract of cocculus indicus, i.	529	Extract resinous of jalap, i.	547
Extract of aloe wood, ii.	192	Extract of colchicum, i.	538	Extract of Indian jalap, i.	554
Extract aqueous of aloes, i.	118	Extract of colocynth, i.	573	Extract of Iceland moss, ii.	188
Extract resinous of aloes, i.	119	Extract compound of colocynth, i.	124, 574	Extract of ipecacuanha, ii.	127
Extract alcoholic of anemone, i.	218	Extract of convallaria, i.	542	Extract of florentine iris, ii.	132
Extract aqueous of anemone, i.	217	Extract of cotton thistle, ii.	273	Extract tartarized of iron, i.	667
Extract of angelica, i.	225	Extract of dandelion, ii.	652	Extract of juniper, ii.	137
Extract of angustura, i.	229	Extract of digitalis, i.	614	Extract of lead, ii.	375, 376
Extract anodyne, ii.	291	Extract of dock, ii.	535	Extract aqueous of leopard's bane, i.	280
Extract aperient, i.	409	Extract of dog-grass, ii.	697	Extract spirituous of leopard's bane, i.	281
Extract of aristolochia, i.	278	Extract of elaterium, ii.	243	Extract of garden lettuce, ii.	152
Extract of asarum, i.	295	Extract alcoholic of elecampane, ii.	119	Extract of viroous lettuce, ii.	152
Extract of avens, i.	719	Extract aqueous of elecampane, ii.	117	Extract of liquorice, i.	724
Extract of balm, ii.	228	Extract of elm, ii.	700	Extract of liquorice juice, i.	723
Extract of beccabunga, ii.	719	Extract of fumitory, i.	680	Extract of logwood, ii.	194
Extract of belladonna, i.	310	Extract of germander, ii.	681	Extract of lovage, ii.	197
Extract bezoardic, i.	225	Extract alcoholic of gentian, i.	711	Extract macrocostine, i.	126
Extract of birch, i.	338	Extract aqueous of gentian, i.	710	Extract alcoholic of madder, ii.	530
Extract of bistort, ii.	407	Extract of geoffræa, i.	717	Extract aqueous of madder, ii.	530
Extract of bitter-sweet, ii.	615	Extract of golden rod, ii.	618	Extract of marigold, i.	365
Extract of blessed thistle, i.	524	Extract Goulard's, ii.	376	Extract of may-apple, ii.	402
Extract of borage, i.	342	Extract of gratiola, i.	731	Extract of meat, i.	631
Extract of bryony, i.	344	Extract alcoholic of gratiola, i.	731	Extract alcoholic of milfoil, i.	36
Extract of burdock, i.	272	Extract of great mullein, ii.	716		
Extract of burnet saxifrage, ii.	357	Extract of ground ivy, i.	721		
Extract of calumba, i.	529	Extract of ground pine, ii.	681		
Extract of cantharides, i.	377	Extract alcoholic of guaiac, i.	733		
Extract of carob-bean, i.	440	Extract aqueous of guaiac, i.	732		
Extract of carrot, i.	608				
Extract alcoholic of cascarilla, i.	568				
Extract aqueous of cascarilla, i.	568				
Extract of cassia, i.	421				
Extract of castor, i.	428				

Extract aqueous of milfoil, i.	35	Extract alcoholic of rubarb, ii.	494	Extract mitigated of tobacco, ii.	261
Extract alcoholic of myrrh, i.	759	Extract aqueous of rubarb, ii.	493	Extract of tormentil, ii.	693
Extract aqueous of myrrh, i.	758	Extract compound of rubarb, ii.	494	Extract of toxicodendron, ii.	504
Extract compound of myrrh, i.	759	Extract of Pontic rubarb, ii.	490	Extract alcoholic of valerian, ii.	708
Extract of wild narcissus, ii.	258	Extract of rosemary, ii.	522	Extract aqueous of valerian, ii.	707
Extract of nettle, ii.	701	Extract alcoholic of rue, ii.	537	Extract of veronica, ii.	718
Extract of nutgalls, i.	693	Extract aqueous of rue, ii.	537	Extract of green walnuts, ii.	135
Extract alcoholic of nuxvomica, ii.	266	Extract alcoholic of saffron, i.	463	Extract alcoholic of willow, ii.	543
Extract aqueous of nuxvomica, ii.	266	Extract aqueous of saffron, i.	563	Extract aqueous of willow, ii.	542
Extract of oak bark, ii.	475	Extract of sage, ii.	545	Extract alcoholic of wormwood, i.	29
Extract of opium by fermentation, ii.	290	Extract of sarsaparilla, ii.	594	Extract aqueous of wormwood, i.	26
Extract acetose of opium, ii.	275	Extract fluid of sarsaparilla, ii.	594	Extractum absinthii aquosum, i.	26
Extract alcoholic of opium, ii.	282	Extract of sassafras, ii.	183	Extractum acetosellæ, ii.	327
Extract aqueous of opium, ii.	289	Extract alcoholic of savine, ii.	143	Extractum aconiti, i.	84
Extract aqueous of opium without narcotine, ii.	290	Extract aqueous of savine, ii.	142	Extractum agarici aloeticum, i.	574
Extract cinnabarine of opium, ii.	45	Extract of scordium, ii.	683	Extractum aloes, i.	118
Extract crocated of opium, ii.	291	Extract of scorzonera, ii.	578	Extractum aloes aquosum, i.	118
Extract cydoniated of opium, ii.	291	Extract of scurvy grass, i.	531	Extractum aloes gummosum, i.	118
Extract Langelot's cydoniated of opium, ii.	291	Extract of seneka, ii.	403	Extractum aloes hepaticæ, i.	118
Extract Diest's of opium, ii.	290	Extract alcoholic of seneka, i.	408	Extractum aloes purificatum, i.	118
Extract glycyrrhized of opium, ii.	291	Extract aqueous of seneka, i.	408	Extractum aloes resinosum, i.	119
Extract hard of opium, ii.	290	Extract of soapwort, ii.	557	Extractum anthemidis, i.	450
Extract vinous of opium, ii.	276	Extract of woodsorrel, ii.	327	Extractum anthemidis nobilis, i.	450
Extract of orange leaves, i.	510	Extract aqueous of spruce fir, i.	21	Extractum artemisiæ absinthii aquosum, i.	26
Extract alcoholic of orange peel, i.	515	Extract spirituous of spruce fir, i.	21	Extractum foliorum citri aurantii, i.	510
Extract aqueous of orange peel, i.	509	Extract of squill, ii.	568	Extractum belladonnæ, i.	310
Extract of ox-gall, i.	633	Extract of St. John's-wort, ii.	108	Extractum bilis, i.	632
Extract of petty broom, i.	709	Extract of stramonium, i.	606	Extractum bistortæ, ii.	407
Extract of peony, ii.	329	Extract of stramonium seeds, i.	606	Extractum calami gummo-resinosum, i.	87
Extract of plantain, ii.	367	Extract of sulphurwort, ii.	343	Extractum eardui benedicti, i.	524
Extract of polygala amara, ii.	403	Extract of swallow-wort, i.	296	Extractum cardui tomentosi, ii.	273
Extract of poppy, ii.	337	Extract alcoholic of sweet-flag, i.	87	Extractum carnis, i.	631
Extract of red poppy, ii.	334	Extract vinous of sweet-flag, i.	88	Extractum caryophyllatæ, i.	719
Extract alcoholic of quassia, ii.	472	Extract alcoholic of tansey, ii.	650	Extractum cascarillæ alcoholicum, i.	568
Extract aqueous of quassia, ii.	477	Extract aqueous of tansey, ii.	650	Extractum cascarillæ aquosum, i.	568
Extract of ratany, ii.	147	Extract theriacal, ii.	314	Extractum cascarillæ gummosum, i.	568
		Extract of tobacco, ii.	261	Extractum cascarillæ resinoso-gummosum, i.	568

Extractum cassiæ, i.	421	Extractum ferri poma-	665	Extractum opii aquosum,	289
Extractum cassiæ fistulæ,		tum, i.		ii.	
i.	421	Extractum ferri cum tar-	667	Extractum opii cinnaba-	45
Extractum castorei, i.	428	taro, i.		rinum, ii.	
Extractum catharticum,		Extractum flammulæ	523	Extractum opii cydonia-	291
i.	574	Jovis, i.		tum, ii.	
Extractum catholicum,		Extractum fructus mo-	243	Extractum opii glycyrrhi-	291
i.	574	mordicæ elaterii, ii.		zatum, ii.	
Extractum centaureæ be-		Extractum fumaricæ, i.	680	Extractum opii gummo-	289
nedictæ, i.	524	Extractum genistæ, i.	709	sum, ii.	
Extractum centaurii mi-		Extractum gentianæ, i.	710	Extractum opii vinosum,	276
noris, i.	626	Extractum gentianæ luteæ,		ii.	
Extractum chærophylli,		i.	710	Extractum panchymago-	574
i.	445	Extractum geoffrææ, i.	717	gum Crollii, i.	
Extractum chamæmeli,		Extractum glechomæ		Extractum papaveris, ii.	337
i.	450	hederacæ, i.	721	Extractum papaveris albi,	337
Extractum florum cha-		Extractum glycyrrhizæ,		ii.	
mæmeli, i.	450	i.	724	Extractum papaveris som-	337
Extractum chamomillæ,		Extractum Goulardi, ii.	376	niferi, ii.	535
i.	450	Extractum graminis, ii.	697	Extractum patientiæ, ii.	357
Extractum chelidonii ma-		Extractum gratiolæ, i.	731	Extractum pimpinellæ	
ajoris, i.	445	Extractum guaiaci resi-		albæ, ii.	21
Extractum chinæ alcoh-		nosum, i.	733	Extractum pini, i.	
olicum, i.	482	Extractum humuli, ii.	10	Extractum pini turionum,	21
Extractum cichorii, i.	462	Extractum juglandis, ii.	135	alcoholicum, i.	376
Extractum cinchonæ a-		Extractum nucum jug-		Extractum plumbi, ii.	665
quosum, i.	480	landium viridum, ii.	135	Extractum pomorum fer-	
Extractum cinchonæ gum-		Extractum juniperi, ii.	137	rarium, i.	665
mosum, i.	480	Extractum levistici, ii.	197	Extractum pomorum fer-	665
Extractum cinchonæ lan-		Extractum ligni aloes,		rarium liquidum, i.	472
cifoliæ, i.	482	ii.	192	Extractum quassiæ, ii.	472
Extractum cinchonæ resi-		Extractum ligni guaiaci		Extractum quassiæ alco-	472
nosum, i.	480	aquosum, i.	732	holicum, ii.	494
Extractum cocculi, i.	522	Extractum ligni sassafras,		Extractum quercetani, i.	490
Extractum cochleariæ,		ii.	183	Extractum ratanhæ, ii.	493, 494
i.	531	Extractum ligustici, ii.	197	Extractum rhabarbari,	
Extractum colocynthidis,		Extractum lupuli, ii.	10	alcoholicum, ii.	494
i.	573	Extractum martis, i.	663	Extractum rhapontici, ii.	494
Extractum colocynthidis		Extractum martis cydo-		Extractum rhei, ii.	494
compositum, i.	574	nium, i.	665	Extractum rhei alcoh-	494
Extractum conii, i.	464	Extractum martis poma-		licum, ii.	504
Extractum conii maculati,		tum, i.	665	Extractum rhois radican-	
i.	464	Extractum martis cum		tis, ii.	522
Extractum corticis Peru-		succo pomorum, i.	665	Extractum rorismarini,	
viani alcoholico-aquo-		Extractum martis cum	667	ii.	530
sum, i.	482	tartaro, i.		Extractum rubiæ tincto-	
Extractum corticis Peru-		Extractum nucis vomicæ	266	riæ alcoholicum, ii.	530
viani gummosum, i.	480	aquosum, ii.		Extractum rubiæ tincto-	
Extractum corticis Peru-		Extractum malatis ferri,	665	riæ aquosum, ii.	537
viani vinosum, i.	482	i.		Extractum rutæ alcoh-	
Extractum corticis quer-		Extractum myrrhæ gra-	759	licum, ii.	537
cus, ii.	475	minatum, i.		Extractum rutæ aquosum,	
Extractum croci alcoh-		Extractum myrrhæ gum-	758	ii.	537
olicum, i.	563	mosum, ii.		Extractum rutæ graveo-	
Extractum croci aquo-		Extractum myrrhæ resi-	759	lentis, ii.	542
sum, i.	463	nosum, i.		Extractum sabinæ alco-	
Extractum dauci, i.	608	Extractum narcissi pra-	258	holicum, ii.	142
Extractum dauci carottæ,		tensis, ii.		Extractum salicis aquo-	
i.	608	Extractum nicotianæ, ii.	261	sum, ii.	545
Extractum dulcamaræ,		Extractum nucis vomicæ	266	Extractum salviæ, ii.	
ii.	615	alcoholicum, ii.	135	Extractum florum sam-	552
Extractum fellis, i.	632	Extractum nucum, ii.	266	buci, ii.	
Extractum ferri, i.	663	Extract hard of nux vo-	289		
Extractum ferri cydoni-		mica, ii.			
atum, i.	665	Extractum opii, ii.			

Extractum saponariæ, ii.	557	Fat, i.	315	Ferri limatura, i.	634
Extractum sarsaparillæ, ii.	594	Fat badger's, i.	315	Ferri ramenta, i.	634
Extractum sarsaparillæ fluidum, ii.	594	Fat bear's, i.	316	Ferri rubigo, i.	646
Extractum saturni, ii.	375, 376, 386	Fat beaver's, i.	315	Ferri scobs, i.	634
Extractum scillæ, ii.	568	Fat dog's, i.	315	Ferri squamæ, i.	634
Extractum scordii, ii.	683	Fat duck's, i.	316	Ferri subcarbonas, i.	646
Extractum scorzonæræ, ii.	578	Fat fox's, i.	316	Ferri sulphas, i.	651
Extractum senegæ radicis, ii.	403	Fat hare's, i.	316	Ferrocyanate of potass solution of, i.	61
Extractum sennæ, i.	408	Fat hedge-hog's, i.	315	Ferrum, i.	634
Extractum siliquæ dulcis, i.	440	Fat hen's, i.	316	Ferrum ammoniacale, i.	661
Extractum solutivum, i.	409	Fat heron's, i.	316	Ferrum ammoniacatum, i.	661
Extractum stramonii, i.	606	Fat of the horse's neck, i.	315	Ferrum ammoniatum, i.	661
Extractum stramonii e seminibus, i.	606	Fat human, i.	315	Ferrum borussicum, i.	674
Extractum tabaci, ii.	261	Fat marmot's, i.	316	Ferrum ferri, i.	656
Extractum tanacetii alcoholicum, ii.	650	Fat quail's, i.	316	Ferrum hæmatites, i.	643
Extractum tanacetii aquosum, ii.	650	Fat rabbit's, i.	316	Ferrum muriaticum, i.	656
Extractum taraxaci, ii.	652	Fat seal's, i.	316	Ferrum muriaticum oxydatum crystallisatum, i.	656
Extractum thebaicum, ii.	289	Fat snake's, i.	316	Ferrum oxydatum carbonicum fuscum, i.	646
Extractum thebaicum ammoniacale, i.	729	Fat stork's, i.	316	Ferrum oxydulatum nigrum, i.	640
Extractum theriacale, ii.	314	Fat viper's, i.	317	Ferrum phosphoricum, i.	651
Extractum tormentillæ, ii.	693	Fat wild cat's, i.	315	Ferrum potabile, i.	667
Extractum tritici repentis, ii.	697	Fat wolf's, i.	316	Ferrum præparatum, i.	634
Extractum turpethi, i.	554	Febrifugum Cranii, i.	233	Ferrum pulveratum, i.	634
Extractum valerianæ, ii.	707	Febrifugum salsum, ii.	424	Ferrum salitum, i.	656
Extractum valerianæ alcoholicum, ii.	708	Fecula ari, i.	292	Ferrum sulphuratum, i.	639
Extractum valerianæ aquosum, ii.	707	Fecula cicutæ, i.	464	Ferrum sulphuricum, i.	651
Extractum verbasci, ii.	716	Fecula cucumeris agrestis, ii.	243	Ferrum vitriolatum, i.	651
Extractum veronicæ, ii.	718	Fecula fructuum hippocastani, i.	90	Ferrum vitriolatum exsiccatum, i.	652
Extractum virgæ aureæ, ii.	618	Fecula hyoscyami, ii.	101	Ferrum vitriolatumustum, i.	644
Exuvium serpentum, i.	678	Fecula ireos, ii.	132	Ferrum zooticum, i.	674
Eye-bright, i.	630	Fecula solani tuberosi, ii.	628	Ferula assafœtida, i.	299
Eyes crabs' i.	354	Fecula tapioka, i.	632	Ferula Persica, i.	764
Faba, febrifuga, ii.	113	Fecula of belladonna, i.	310	Fever root, ii.	296
Faba Indica, ii.	113	Fecula of bryony, i.	344	Feverfew, ii.	225
Faba marina, i.	353	Fecula of elaterium, ii.	243	Ficariæ herba, ii.	578
Faba pichurim, ii.	680	Fecula of hemlock, i.	464	Ficariæ radix, ii.	481
Faba sancti Ignatii, ii.	113	Fecula of henbane, ii.	101	Ficus carica, i.	675
Faba tunka, i.	562	Fecula of iris, ii.	132	Ficus religiosa, ii.	150
Faba vulgaris, i.	631	Fecula of marantæ, i.	632	Fig Indian, i.	346
Faburiæ radix, ii.	580	Fecula of peony, ii.	329	Fig tree, i.	675
Fagara octandra, ii.	644	Fecula of potato, ii.	618	Figwort knot-rooted, ii.	578
Fagopyri semen, ii.	407	Fecula of stramonium, i.	605	Figwort water, ii.	578
Fagus sylvatica, i.	631	Fel bovinum, i.	632	Filicis non ramosæ dentatæ radix, i.	308
Farfaræ radix, ii.	699	Fel tauri inspissatum, i.	632	Filiculæ dulcis radix, ii.	407
Farina hordei præparata, ii.	7	Fel vitri, ii.	430	Filipendulæ radix, ii.	621
Farina cypodii, ii.	204	Fellis ferræ herba, i.	625	Fish-glue, i.	696
		Feni camelorum herba, i.	217	Flammulæ herba, ii.	481
		Feniculi aquatici herba, ii.	346	Flammulæ Jovis herba, i.	523
		Feniculi porcini radix, i.	343	Flavedo corticum citri, i.	517
		Fennel, i.	218	Flax, ii.	200
		Fennel flower small garden, ii.	264	Flax purging, ii.	200
		Fenugreci semen, ii.	696		
		Fenugreek, ii.	696		
		Fermentum cerevisiæ, i.	634		
		Fern male, i.	308		
		Ferrariæ herba, ii.	578		
		Ferri cyanuretum, i.	674		

Flea-bane, i. 554, 621, ii. 114	Fomentation aromatic, i. 27, 400 ii. 524, 684	Fotus frigidum, i. 45
Flea-bane Philadelphian, i. 622	Fomentation astrigent, i. 162, 349, 653, 693	Fotus frigidus Schmuck-eri, ii. 437
Flea-bane viscous, i. 622	ii. 367, 379, 471, 476	Fotus ad gangrænam, i. 492
Fleseed, ii. 367	Fomentation Brunning-hausen's, ii. 414	Fotus ad perniones, i. 162
Flesh and spine of the fox, i. 697	Fomentation camphorated, i. 373	Fotus ad pustulas in cute favendas, i. 57
Flesh of the tortoise, i. 697	Fomentation for chil-blains, i. 460	Fotus refrigerans, ii. 437
Flesh and spine of the viper, i. 697	Fomentation for colic, i. 452	Fotus refrigerans et de-primens, i. 45
Flies Spanish, i. 377	Fomentation common, i. 44	Fowl's gizzard, i. 699
Flores æris, i. 595	Fomentation detergent, ii. 321	Foxglove purple, i. 611
Flores argentinæ anti-monii, i. 234	Fomentation diuretic, i. 193	Fragorum baccae, i. 677
Flores auri, i. 661	Fomentation emmenagogue, i. 345	Fraga, i. 677
Flores boracis, i. 48	Fomentation emollient, i. 148, ii. 201	Fragaria vesca, i. 677
Flores lupuli, ii. 10	Fomentation emollient and relaxant, i. 148	Frankincense, ii. 271
Flores macis, ii. 254	Fomentation excitant, ii. 546	Fraseria Walteri, i. 678
Flores salis ammoniaci, i. 191	Fomentation laxative and antispasmodic, ii. 459	Fraxinella, i. 611
Flores salis ammoniaci hæmatisantes, i. 661	Fomentation nitric, i. 70	Fraxini pumilæ radix, i. 611
Flores salis ammoniaci martiales, i. 661	Fomentation resolvent, i. 45, 60, 193, 197, 199, 452, ii. 379, 387, 424, 553, 672	Fraxinus argentea, ii. 218
Flores sulphuris compo-siti, i. 128	Fomentation saturnine, ii. 379	Fraxinus excelsior, i. 678
Flores viridis æris, i. 595	Fomentation sedative, i. 148, ii. 283, 338, 388	ii. 218
Flower de luce blue, ii. 130	Fomentation stimulant, i. 57, 73, 373, ii. 262, 448	Fraxinus ornus, ii. 218
Flowers of nutmeg, ii. 254	Fomentum acidi phosphorici, i. 73	Fraxinus parviflora, ii. 218
Flowers pectoral, ii. 215	Fomentum aromaticum, ii. 524, 684	Fraxinus rotundifolia, ii. 218
Flowers of sal ammo-niac, i. 191	Fomentum Belladonnæ ammoniatum, i. 197	Fraxinus sæbrufescens, ii. 218
Fly agaric, i. 165	Fomentum commune, i. 44	Frog edible, i. 697
Fly potato, i. 377	Fomentum discuteins, i. 44	Frog-spawn, i. 697
Fœniculi radix, i. 219	Fomentum nitricum, i. 70	Frumentum, ii. 579
Fœniculi dulcis radix, i. 219	Fomentum ad perniones, i. 460	Fucus, ii. 604
Fœniculi Lusitanici se-men, ii. 592	Fomentum quercinum, ii. 476	Fucus barbatus, ii. 2
Fœniculi marini folia, i. 562	Fomentum saturninum, ii. 387	Fucus cricoides, ii. 2
Fœniculi ursini radix, ii. 198	Fomentum saturninum cum opio, ii. 379	Fucus fascicla, ii. 2
Folia cochlearum, i. 354	Food antiscorbutic, i. 479	Fucus helminthocorton, ii. 2
Fomentatio discutiens, i. 60	Fool's parsley, i. 102	Fucus sodoides, ii. 2
Fomentatio refrigerans, ii. 379	Formica rufa, i. 676	Fucus vesiculosus, i. 678
Fomentation camphorated ammoniacal, i. 187	Fotus adstringens, ii. 476	Fuligo splendens, i. 679
Fomentation anodyne, ii. 338	Fotus aluminosus, i. 162	Fuller's thistle, i. 618
Fomentation anthelmin-tic, i. 278	Fotus anodynus, ii. 716	Fumaria officinalis, i. 679
Fomentation antihæ-morrhoidal, i. 114	Fotus antisepticus, ii. 476	Fumigatio muriatico-oxygenata, i. 458
Fomentation antihæmor-rhoidal, ii. 216	Fotus aromaticus, i. 27	Fumigatio Smithiana, i. 69
Fomentation antiphlo-gistic, ii. 460	Fotus communis, i. 27	Fumigation chlorine, i. 458
Fomentation antipleuri-tic, ii. 335	Fotus discutiens, i. 45	Fumigation disinfecting, i. 57, 69
Fomentation antiseptic, i. 492, 304		Fumigation resinous, ii. 675

- Gadus lota*, i. 360
Gagates, i. 682
Galanga major, i. 682
Galanga minor, i. 682
Galanga spuria, i. 682
Galanga great, i. 682
Galanga lesser, i. 682
Galarhæus piperatus, ii. 636
Galbanum, i. 682
Galbuli, i. 580
Galega officinalis, i. 691
Galega Virginiana, i. 691
Galeopsidis herba, ii. 154
Gales herba, ii. 254
Galipea cusparia, i. 228
Galium aparine, i. 691
Galium cruciatum, i. 691
Gall, i. 632
Gall nut, i. 691
Galla de Aleppo, i. 691
Galla quercina, i. 691
Galla tinctoria, i. 691
Galla turcica, i. 691
Gallato-carbonate of potass, ii. 420
Gallinæ stercus, i. 83
Gallipot, ii. 673
Gallitrichi herba, ii. 544
Galium mollugo, i. 694
Galium verum, i. 694
Galls blue, i. 692
Galls white, i. 692
Gamboge, i. 694
Gambogium, i. 694
Garcinia cambogia, i. 695
Gardenia gummifera, ii. 486
Gargarisma adstringens, i. 56
Gargarisma adstringens acidum, i. 77
Gargarisma in angina maligna, i. 46
Gargarisma antiscorbuticum, i. 537
Gargarisma ad aphthas, ii. 609
Gargarisma belladonnæ cum hydrargyro, ii. 74
Gargarisma e borace, i. 609
Gargarisma boraxatum, ii. 609
Gargarisma cochleariæ, i. 537
Gargarisma commune, ii. 516
Gargarisma mercuriale Plenckii, ii. 15
Gargarisma nitrosum, ii. 435
Gargarisma piperitum, ii. 364
Gargarisma refrigerans, ii. 435
Gargarisma salviæ cum acido muriatico, i. 56
Gargarisma saturninum, ii. 387
Gargarisma e scilia, ii. 575
Gargarisma viride, i. 592
Gargarisma in ulceribus venereis, ii. 69
Gargle, ii. 351
Gargle acetose, i. 193
Gargle acidulous, i. 56
Gargle ammoniacal, i. 171, 186
Gargle anti-odontalgic, i. 231
Gargle antiphlogistic, i. 97
Gargle antiscorbutic, i. 520, 536, 537, 609, ii. 483
Gargle antiseptic, i. 172, 460, 491, 494
Gargle antisiphilitic, ii. 49, 71
Gargle astringent, i. 56, 77, 163, 197, 371, 693, ii. 148, 378, 387, 471, 590, 693
Gargle borated, ii. 609
Gargle chloric, i. 460
Gargle common, ii. 434
Gargle demulcent, ii. 216, 341
Gargle detergent, ii. 516, 553
Gargle detersive, i. 70, 77, 763
Gargle emollient, i. 148, ii. 642
Gargle gingival, i. 536
Gargle humectant, i. 521
Gargle mercurial, ii. 69, 74, 77
Gargle Plenck's mercurial, ii. 15
Gargle opiate, ii. 283
Gargle oxymellated, i. 44
Gargle refrigerant, ii. 435
Gargle resolvent, i. 192
Gargle scillitic, ii. 575
Gargle sedative, ii. 293, 338, 387, 590
Gargle sialagogue, i. 231
Gargle stimulant, i. 46, 231, 603, ii. 363, 364, 545, 590
Gargle styptic, ii. 378
Garlic, i. 114
Garnet, i. 157
Gas hepaticum, i. 64
Gas hydrocarbonatum, ii. 98
Gas hydrogenium carbonatum, ii. 98
Gas hydrogenium sulphuratum, i. 64
Gas inflammable, ii. 98
Gas percarbonated hydrogen, ii. 98
Gaultheria procumbens, i. 696
Gaz azoticum, ii. 265
Gaz azoticum oxygenatum, ii. 265
Gelatina, i. 696
Gelatina ex amylo solani tuberosi, ii. 618
Gelatina animalis, i. 704
Gelatina cornu cervi, i. 704
Gelatina cornu cervi acida, i. 705
Gelatina cornu cervi amygdalata, i. 704
Gelatina cornu cervi citrata, i. 705
Gelatina cydoniorum, i. 598
Gelatina glycyrrhizæ, i. 724
Gelatina lichenis Islandici, ii. 190
Gelatina liquiritiæ, i. 724
Gelatina ribesiorum nigrorum, ii. 505
Gelatina ribium, ii. 506
Gelatina radicis saleb, ii. 540
Gelatine, i. 696
Gelapii radix, i. 543
Genepi albi herba, i. 291
Genicellæ radix, ii. 405
Geniculatæ radix, ii. 405
Genipi spuri herba, i. 34
Genipi veri herba, i. 34
Genista Canariensis, ii. 195
Genista juncea, i. 708
Genista sagittalis, i. 708
Genista scoparia, i. 708
Genista tinctoria, i. 708
Genistæ angulosæ herba, i. 708
Genistellæ herba, i. 708
Gentia, i. 716
Gentian, i. 709
Gentian blue, i. 709
Gentiana Catesbæi, i. 709
Gentiana chirayita, i. 86
Gentiana lutea, i. 709
Gentianæ albæ radix, ii. 155
Gentianæ cruciatæ radix, i. 691
Gentianæ majoris radix, i. 709
Gentianæ minoris radix, i. 691
Gentianæ nigrae radix, ii. 581
Gentianæ rubrae radix, i. 709
Gentianæ veterum radix, i. 709
Gentiana, i. 716
Gentianina, i. 716
Gentianine, i. 709, 716
Geoffræa Jamaicensis cortex, i. 717
Geoffræa inermis, i. 716
Geoffræa Surinamensis, i. 717
Geranii batrachoides herba, i. 718
Geranium maculatum, i. 718
Geranium pratense, i. 718

Geranium Robertianum, i.	718	Gold mosaic, ii.	625	Gum lac, ii.	150
Geranium rotundifolium, i.	718	Golden-rod, ii.	618	Gum olive, i.	764
Geranium sanguineum, i.	718	Gold thread trefoil, i.	558	Gum senegal, i.	766
Geranium spotted, i.	718	Good king Henry, i.	457	Gum tragacanth, i.	766
Germander common, ii.	680	Goose-foot fetid, i.	458	Gummi adstringens Fothergilli, ii.	145
Germander yellow-flowered, ii.	681	Goose-foot Mexican, i.	456	Gummi aloes, i.	117
Germander water, ii.	682	Gooseberry, ii.	506	Gummi ammoniacum, i.	742
Geum rivale, i.	719	Gossypina, i.	730	Gummi ammoniacum al-kalisatum, i.	749
Geum urbanum, i.	718	Gossipium herbaceum, i.	730	Gummi anime, i.	741
Ghitta jemoco, i.	694	Gossypii semina, i.	730	Gummi arabicum, i.	749
Gialappæ radix, i.	543	Gourd, i.	578	Gummi assafoetida, i.	299
Gigartina helminthocorton, ii.	2	Gouttes uterines de la reine d'Espagne, ii.	267	Gummi benzoës, i.	330
Gigartina plicata, ii.	2	Grains of paradise, i.	203	Gummi carannæ, i.	754
Gilla, i.	644	Graminis albi radix, ii.	697	Gummi ceraci, i.	755
Gillenia trifoliata, i.	720	Graminis canini radix, ii.	697	Gummi cerasorum, i.	755
Gin, i.	104	Graminis majoris radix, i.	396	Gummi copal, i.	755
Ginger, i.	203	Graminis mannæ semen, i.	609	Gummi elasticum, i.	755
Ginger broad-leaved, i.	203	Graminis orientalis herba i.	217	Gummi euphorbii, i.	756
Ginger wild, i.	293	Graminis radix, ii.	697	Gummi gamandraræ, i.	694
Ginseng, ii.	331	Graminis rubri radix, i.	396	Gummi gambiense, ii.	145
Glacies mariæ vulgaris, i.	362	Grana actes, ii.	551	Gummi de Goa, i.	694
Gladioli lutei radix, ii.	130	Grana cnidia, i.	603	Gummi guaiaci, i.	732
Gladioli cærulei radix, ii.	130	Grana kermes, i.	530	Gummi guttæ, i.	694
Gladiolus communis, i.	720	Grana manguinetta, i.	203	Gummi guttæ Americana-num, i.	695
Glandes quercûs, ii.	474	Grana meliguetta, i.	203	Gummi guttæ sianense, i.	695
Glandes quercûs tostæ, ii.	477	Grana molucca, i.	570	Gummi guttæ spurium, i.	695
Glans unguentaria, ii.	245	Grana paradisi, i.	203	Gummi guttæ verum, i.	695
Glass of antimony, i.	235	Grana tiglia, i.	570	Gummi guttæ Zeylanicum, i.	695
Glass cerated of antimony, i.	235	Granatum, i.	157	Gummi hederæ, i.	757
Glasti herba, ii.	133	Graphites, i.	639	Gummi de jemu, i.	694
Glaucium corniculatum, i.	720	Graphites hydrargyratus, ii.	26	Gummi juniperi, ii.	555
Glechoma hederacea, i.	720	Grass creeping panic, ii.	342	Gummi kino, ii.	145
Globularia vulgaris, i.	722	Grass five-leaved, ii.	465	Gummi laccæ, ii.	150
Globuli carminativi, i.	633	Gratiæ dei radix, i.	730	Gummi ladanum, i.	757
Globuli ad erysipelas, ii.	373	Gratiola officinalis, i.	730	Gummi ligni sancti, i.	732
Globuli ferri tartarisati, i.	667	Gravel root, i.	628	Gummi mimosæ, i.	749
Globuli martiales soluti, i.	667	Grease capon's, i.	316	Gummi mimosæ, i.	749
Globuli moschati, ii.	251	Grease capon's, i.	316	Gummi myrrhæ, i.	758
Globuli tartratis ferri et lixivii, i.	667	Grease goose, i.	316	Gummi nostras, i.	755
Globuli tartari martiales, i.	667	Great centauri, i.	433	Gummi oleæ, i.	764
Glycyrrhiza ecchinata, i.	723	Gromwell, ii.	203	Gummi olibanum, ii.	271
Glycyrrhiza glabra, i.	722	Ground-ivy, i.	720	Gummi panacis, ii.	321
Glycyrrhiza lucida, i.	724	Ground-pine, ii.	681	Gummi sagapenum, i.	764
Glycyrrhizine, i.	722	Groundsel, ii.	584	Gummi sanguis draconis, i.	764
Glue, i.	696	Guaiac, i.	731	Gummi sarcocollæ, i.	765
Glue fish, i.	696	Guaiaci lignum, i.	732	Gummi Senegalense, i.	766
Gluten pectorale, i.	724	Guaiacina, i.	732	Gummi serapinum, i.	764
Gnaphalii dioici herba, i.	230	Guaiacum officinale, i.	731	Gummi serapionis, i.	749
Goat's blood, i.	699	Guilandina moringa, ii.	195	Gummi tragacantha, i.	766
Goat's rue, i.	691	Gum ammoniac, i.	742	Gutta gamandra, i.	694
Gold, i.	312	Gum arabic, i.	749	Gutta gamba, i.	694
Gold divided, i.	312	Gum anime, i.	741	Guttulæ albæ Wardii, ii.	89
		Gum bdellium, i.	329	Guttæ acidæ tonicæ, i.	79
		Gum benzoin, i.	330	Guttæ anglicanæ cephalicæ, i.	187
		Gum caranna, i.	754	Guttæ ad blennorrhœam, i.	556
		Gum cherry, i.	755	Guttæ e cupro, i.	582
		Gum copal, i.	755	Guttulæ diureticæ, ii.	262
		Gum elastic, i.	755	Guttæ febrifugæ, i.	74
		Gum guaiac, i.	732		
		Gum hederæ, i.	757		
		Gum juniper, ii.	555		

Guttæ ferratæ, i.	656	Heart's-ease, ii.	723	Herbæ Brittanicæ radix,	
Guttæ nervinæ, i.	659	Heath, i.	366	ii.	534
Guttulæ phosphoratæ,		Hedera helix, i. 757, ii.	1	Hermodactyles, ii.	5
i.	73	Hederæ arboreæ folia,		Hermodactyles spurious,	
		ii.	1	ii.	5
Hæmatites lapis, i.	643	Hederæ terrestus herba,		Hermodactyli, ii.	5
Hæmatinum, ii.	193	i.	721	Herniaria glabra, ii.	5
Hæmatoxylinum, ii.	193	Hedge hyssop, i.	730	Herniaria hirsula, ii.	6
Hæmatoxylum campe-		Hedysarum alhagi, ii.	218	Hibisci radix, i.	143
chianum, ii.	193	Helenii radix, ii.	115	Hieracium murorum, ii.	6
Halicacabi baccae, ii.	350	Helianthus tuberosus,		Hieracium pilosella, ii.	6
Haloïdum oxygenatum,		ii.	1	Hippocastani cortex, i.	90
ii.	440	Heliotropium Europæum,		Hippocras, ii.	171
Hamamelis Virginiana,		ii.	1	Hippolapathi radix, ii.	535
ii.	1	Helix pomatia, i.	354, 698	Hips, ii.	519
Hare, i.	697	Hellebore white, ii.	714	Hirundinariæ radix, i.	295
Hartshorn, i.	359	Hellebore albi radix, ii.	714	Hirundines combustæ, i.	391
Hartshorn burnt, i.	360	Helminthocorton, ii.	2	Hirundo officialis, ii.	6
Hartshorn calcined, i.	360	Helminthocortos, ii.	2	Hirundo rustica, i.	355
Hartshorn golden, i.	314	Helxines herba, ii.	341	Hispidulæ herba, i.	230
Hart's tongue, ii.	577	Helychrysum arenarium,		Histrix cristatus, i.	338
Hart's truffle, ii.	577	ii.	4	Holly knee, ii.	535
Hastuli regis radix, i.	298	Helychrysum stœchas,		Holly sea, i.	622
Haustus anti-emeticus,		ii.	4	Hollyhock, i.	154
i.	50	Hemlock, i.	463	Holy herb, ii.	717
Haustus astringens, ii.	588	Hemlock water, i.	469	Honey of annual mer-	
Haustus balsamicus, i.	323	ii.	346	cury, ii.	241
Haustus ad blennor-		Hemp, i.	376	Honey compound of an-	
hagiam, i.	556	Hemp agrimony, i.	628	annual mercury, ii.	241
Haustus catharticus pro		Henbane, ii.	99	Honey of celandine, i.	456
infantibus, ii.	493	Henbane white, ii.	99	Honey of colchicum,	538
Haustus catharticus oleo-		Henchera Americana, ii.	4	Honey of elder flowers,	
sus, ii.	508	Hepar anguillæ, i.	699	ii.	552
Haustus catharticus sa-		Hepar antimonii, i. 241,	254	Honey glycyrrhized, i.	728
linus, ii.	211	Hepar canis rabidi, i.	697	Honey of horseradish,	
Haustus collæ animalis,		Hepar lupi, i.	697	ii.	482
i.	701	Hepar sulphuris, ii.	416	Honey mercurial, ii. 17, 38,	
Haustus diureticus Mi-		Hepar sulphuris calca-		85, 248	
manni, ii.	445	reum, i.	350	Honey with protochlo-	
Haustus diaphoreticus		Hepar sulphuris salinum,		ruret of mercury, ii.	49
resolvens, i.	84	ii.	416	Honey of raspberries,	
Haustus incitans nervi-		Hepar sulphuris volatile,		ii.	532
nus, i.	372	i.	194	Honey of rosemary, ii.	522
Haustus incitans et re-		Hepar sulphuris vulgare,		Honey of roses, ii.	515
solvens, ii.	495	ii.	416	Honey solutive of roses,	
Haustus ex olibano, ii.	272	Hepaticæ nobiles herba,		ii.	516
Haustus pectoralis inci-		ii.	4	Honey saponaceous, ii.	459
tans, ii.	405	Hepatica triloba, ii.	4	Honey of squill, ii.	569
Haustus pectoralis resol-		Hepaticæ albæ herba,		Honey compound of	
vens, ii.	617	ii.	342	squill, ii.	569
Haustus sedans narco-		Hepaticæ fontanæ herba,		Honey terebinthinated,	
ticus, ii.	102	ii.	221	ii.	669
Haustus stomachicus, ii.	473	Hepatorii herba, i.	103	Honey of violet, ii.	722
Haustus terebinthinatus,		Heptaphylli radix, ii.	692	Honey water, i.	559
i.	556	Herocanthæ radix, i.	396	Honeysuckle, ii.	203
Haustus tonicus adstrin-		Heracleum spondilium,		Hop, ii.	10
gens, ii.	280	ii.	5	Hordeum præparatum,	7
Haustus tonicus martia-		Heracium gummifera,		Hordeum vulgare, ii.	7
lis, i.	666	i.	724	Horehound white, ii.	221
Hawk-weed Hungarian,		Herb Bennet, i.	718	Horehound wild, i.	628
ii.	111	Herb Christopher, i.	89	Hormini herba, ii.	544
Hawthorn, ii.	242	Herb Robert, i.	718	Horn of the rhinoceros,	
Hazel, i.	561	Herba Ægyptiaca, ii.	225	i.	699
Hazel white, ii.	1	Herba sardoa, ii.	480	Horse chesnut, i.	89
Heal-weed, i.	596	Herbæ aromaticæ, ii.	545	Horseradish, ii.	481

Horse-tail, i.	621	Hydrargyrum stibiato-		Hydrochlorate of mer-	
Hound's tongue, i.	601	sulphuratum, ii.	30	cury and of ammo-	
Houseleek great, ii.	584	Hydrargyrum sulphura-		nia, ii.	89
Humulus lupulus, ii.	10	tum nigrum, ii.	28	Hydrochlorate of pro-	
Hyacinth musk, ii.	254	Hydrargyrus, ii.	12	toxide of sodium,	
Hydragogum Boylei, i.	273	Hydrargyrus acetatus, ii.	95	ii.	602
Hydrargyri acetas, ii.	95	Hydrargyrus causticus		Hydrochlorate of silver	
Hydrargyri calcinatus,		flavus, ii.	82	and of ammonia, i.	276
ii.	36	Hydrargyrus causticus		Hydrochlorate of soda,	
Hydrargyri cyanuretum,		luteus, ii.	82	ii.	602
ii.	80	Hydrargyrus corrosivus		Hydrochlorinas ferri, i.	656
Hydrargyri nitrico-oxy-		albus, ii.	66	Hydrochlorinas natricus,	
dum, ii.	36	Hydrargyrus corrosivus		ii.	602
Hydrargyri oxydum cine-		flavus, ii.	82	Hydrocyanas ferri, i.	674
reum, ii.	32	Hydrargyrus emeticus		Hydrogale, ii.	150
Hydrargyri oxydum ni-		flavus, ii.	82	Hydrogen, ii.	98
grum, ii.	32	Hydrargyrus flavus, ii.	82	Hydrogenium, ii.	98
Hydrargyri oxydum ru-		Hydrargyrus muriaticus		Hydrogeno-sulphuretum	
brum, ii.	36	dulcis, ii.	46	ammoniacæ liquidum,	
Hydrargyri oxymurias,		Hydrargyrus muriatus,		i.	194
ii.	67	ii.	66	Hydrolapathi radix, ii.	534
Hydrargyri persulphas,		Hydrargyrus muriatus		Hydromel anti-asthmatic,	
ii.	82	ammoniacalis fortior,		i.	615
Hydrargyri submurias,		ii.	89	Hydromel anti-catarrhal,	
ii.	46	Hydrargyrus muriatus		ii.	192
Hydrargyri submuriatus		corrosivus, ii.	66	Hydromel asthmaticum,	
ammoniacus, ii.	90	Hydrargyrus nitratus		i.	615
Hydrargyri sulphuretum,		ruber, ii.	36	Hydromel vinous, i.	110
nigrum, ii.	28, 42	Hydrargyrus præcipita-		Hydropiperis herba, ii.	406
Hydrargyri sulphuretum		tus flavus, ii.	82	Hydrosellini radix, i.	266
rubrum, ii.	43	Hydrargyrus sulphura-		Hydrosulphas ammo-	
Hydrargyrum, ii.	12	tus niger, ii.	28	niæ, i.	194
Hydrargyrum acetatum,		Hydrargyrus cum sul-		Hydrosulphas calcis, i.	362
ii.	95	phure, ii.	28	Hydrosulphas protoxidi	
Hydrargyrum ammoniato-		Hydrargyrus tartarisa-		calcii, i.	362
muriaticum, ii.	89	tum, ii.	97	Hydrosulphate of ammo-	
Hydras calcariae, i.	347	Hydrargyrus vitriolatus,		nia, i.	194
Hydrargyrum cum creta,		ii.	82	Hydrosulphate sulphuret-	
ii.	18	Hydras kalicum, ii.	412	ted of ammonia, i.	194
Hydrargyrum cum mag-		Hydrastes Canadensis,		Hydrosulphate sulphuret-	
nesia, ii.	18	ii.	98	ted of antimony, i.	249
Hydrargyrum muriati-		Hydrate of lime, i.	347	Hydrosulphate of potass,	
cum mite, ii.	46	Hydrate of sulphur, ii.	633	ii.	441
Hydrargyrum muriati-		Hydriodas potassæ, ii.	121	Hydrosulphate of soda,	
cum præcipitatum, ii.	89	Hydriodate of iron, ii.	123	ii.	614
Hydrargyrum muriatum		Hydrocarbonate of cop-		Hydrosulphuratum oxydi	
corrosivum, ii.	66	per, i.	582	stibii rubrum, i.	243
Hydrargyrum nitricum,		Hydrochloras ammoniæ,		Hydrosulphuretum am-	
ii.	85	i.	191	monia, i.	194
Hydrargyrum oxydatum		Hydrochloras auri et so-		Hydrosulphuretum am-	
nigrum, ii.	84	dii, i.	313	moniacale aquosum,	
Hydrargyrum oxydatum		Hydrochloras deutoxidi		i.	194
rubrum, ii.	36	cupri, i.	558	Hydrosulphuretum anti-	
Hydrargyrum oxydula-		Hydrochloras potassæ,		monii rubrum, i.	243
tum nigrum, ii.	32	ii.	440	Hydrosulphuretum liqui-	
Hydrargyrum phospho-		Hydrochlorate of ammo-		dum, i.	65
ratum, ii.	81	nia, i.	191	Hydrosulphuratum oxydi	
Hydrargyrum phospho-		Hydrochlorate of ammo-		stibii fuscum, i.	243
reum, ii.	81	nia and of iron, i.	661	Hydrosulphuretum lu-	
Hydrargyrum præcipita-		Hydrochlorate of gold,		teum oxydi stibii sul-	
tum album, ii.	89	i.	313	phuratum, i.	249
Hydrargyrum purifica-		Hydrochlorate of iron,		Hydrosulphuretum oxy-	
tum, ii.	12	i.	656	duli stibii, i.	243
Hydrargyrum sacchara-		Hydrochlorate of man-		Hpdrosulphuretum ru-	
tum, ii.	16	ganese, ii.	217	brum stibii, i.	243

Hydrosulphuretum stibiosum, i.	243	Jacea nigra, ii.	112	Illicium anisatum, ii.	113
Hygra, i.	592	Jaceæ herba, ii.	723	Imbricaria parietina, ii.	187
Hymenæa courbaril, i.	742	Jaceæ nigra radix, ii.	112	Imperatoria ostruthium, ii.	114
Hyoscyama, ii.	99	Jacobææ herba, ii.	584	Imperatoria radix, ii.	114
Hyoscyaminum, ii.	99	Jalap, i.	542	Indian physic, i.	720
Hyoscyamus albus, ii.	99	Jalap Indian, i.	553	Indian pink, ii.	620
Hyoscyamus niger, ii.	99	Jalapæ albæ radix, i.	553	Indigo, ii.	351
Hypercarbonas ammoniæ, i.	182	Jalapæ radix, i.	543	Indigofera anil, ii.	351
Hypercarbonas potassæ, ii.	419	Jamaicine, i.	717	Indigofera argentea, ii.	351
Hypercarbonas sodæ, ii.	604	Jasminum officinale, ii.	134	Indigofera tinctoria, ii.	351
Hyperici perforati summitates, ii.	107	Jatropha curcas, ii.	134	Indigotine, ii.	133, 351
Hyperici summitates, ii.	107	Jatropha manihot, i.	632	Inhalation balsamic, i.	323
Hypericum androsæmum, ii.	107	Jaw-bone of the pike, i.	360	Infusio anthelmintico-incitans, ii.	683
Hypericum bacciferum, i.	695	Iberidis herba, ii.	186	Infusio anthelmintico-tonica, i.	528
Hypericum Cayenense, i.	695	Iceland, ii.	241	Infusio laxativa sedans, ii.	445
Hypericum quadrangulare, ii.	107	Icica caranna, i.	754	Infusio nervina ætherea, i.	281
Hypericum perforatum, ii.	107	Icica icicriba, ii.	486	Infusio nervina martialis, i.	281
Hyperoxodes ferri nigrum, i.	640	Ichthyocola, i.	696	Infusion of acorns, ii.	477
Hyperoxodes hydrargyri, ii.	36	Jelly analeptic, i.	705	Infusion of American winter-green, i.	458
Hypersulphas aluminæ et potassæ, i.	188	Jelly of bread, ii.	333	Infusion of anemone, i.	218
Hypertartras potassæ, ii.	449	Jelly of red cabbage, i.	343	Insusio of angelica, i.	225
Hypoboras sodæ, ii.	608	Jelly calf's foot, i.	704	Infusion of angustura, i.	229
Hypocarbonas ammoniæ pyro-animale, i.	184	Jelly of cinchona, i.	492	Infusion of anise, ii.	356
Hypocarbonas ammoniæ pyro-animale liquidum, i.	185	Jelly of corsican moss, ii.	3	Infusion anthelmintic, i.	528
Hypocarbonas potassæ liquidum, ii.	422	Jelly of red currants, ii.	506	Infusion antihydronic, ii.	486
Hypochaeris maculata, ii.	111	Jelly demulcent, i.	704	Infusion antiparalytic, ii.	485
Hypocist, ii.	12	Jelly of glue, i.	703	Infusion antiscorbutic, i.	533
Hypocistis, ii.	12	Jelly of hartshorn, i.	704	Infusion of astragalus, i.	307
Hypoglossi herba, ii.	536	Jelly acidulous hartshorn, i.	705	Infusion astrigent, ii.	513
Hypophosphate of potass, ii.	427	Jelly amygdaline hartshorn, i.	704	Infusion of aya-pana, i.	628
Hyposulphite of mercury and of ammonia, ii.	84	Jelly of Iceland moss, ii.	190	Infusion of balm, ii.	228
Hyposulphite of mercury and of potass, ii.	84	Jelly of Iceland moss with cinchona, ii.	191	Infusion of belladonna, i.	311
Hyposulphite of mercury and of soda, ii.	84	Jelly of Iceland moss with opium, ii.	191	Infusion bitter, i.	525
Hyssop, ii.	111	Jelly isinglass, i.	704	Infusion alkalized bitter, i.	712
Hyssop hedge, i.	730	Jelly laxative, ii.	218	Infusion of bitter-sweet, ii.	616
Hyssopi herba, ii.	111	Jelly purgative, i.	550	Infusion of blessed thistle, i.	525
Hyssopus officinalis, ii.	111	Jelly aromatic of quince, i.	598	Infusion of bloodroot, ii.	555
		Jelly of salep, ii.	540	Infusion of borage, i.	341
		Jelly starch, i.	214	Infusion of bugloss, i.	216
		Jelly of potatoe-starch, ii.	618	Infusion of burnet saxifrage, ii.	357
		Jelly of tortoise, i.	704	Infusion calcareous of cinchona, i.	488
		Jerusalem artichoke, ii.	1	Infusion of calumba, i.	528
		Jerusalem oak, i.	457		
		Jessamine, ii.	134		
		Jet, i.	682		
		Jew's ear, ii.	695		
		Ignatia amara, ii.	113		
		Ignis herba, ii.	136		
		Ilex aquifolium, ii.	113		
		Ilex vomitoria, ii.	113		
		Illecebr majoris radix, ii.	580		
		Illecebr vermicularis herba, ii.	580		

- | | | | | | |
|--|----------|--|---------------|--|----------|
| Infusion of cantharides, i. | 378 | Infusion of hemp seed, i. | 376 | Infusion of quassia with sulphate of zinc, ii. | 473 |
| Infusion of capsicum, ii. | 364 | Infusion of hops, ii. | 11 | Infusion of rhododendron, i. | 503 |
| Infusion cardiac, i. | 278 | Infusion of horseradish, ii. | 485 | Infusion of rhubarb, ii. | 494 |
| Infusion cardiac with vinegar, i. | 278 | Infusion of hyssop, ii. | 111 | Infusion of rosemary, ii. | 522 |
| Infusion carminative, i. | 452 | Infusion of Iceland moss, ii. | 188 | Infusion acidulous of roses, ii. | 512 |
| Infusion of cascarrilla, i. | 569 | Infusion of ipecacuanha, ii. | 127 | Infusion of roses, ii. | 512 |
| Infusion of cassia, i. | 421 | Infusion of juniper, ii. | 140 | Infusion of saffron, i. | 513 |
| Infusion cathartic and anodyne, i. | 452 | Infusion compound of juniper, ii. | 141 | Infusion of sage, ii. | 545 |
| Infusion of chamomile, i. | 451 | Infusion of yellow ladies'-bedstraw, i. | 694 | Infusion acidulated of sage, ii. | 545 |
| Infusion compound of chamomile, i. | 452 | Infusion of lavender, ii. | 156 | Infusion vinous of sage, ii. | 545 |
| Infusion of cherry-laurel, i. | 439 | Infusion laxative, i. | 415 | Infusion of sassafras, ii. | 183 |
| Infusion of cherry-stalks, i. | 439 | Infusion of leopard's-bane, i. | 280 | Infusion of savine, ii. | 143 |
| Infusion of chick peas, i. | 461 | Infusion æthereous of leopard's-bane, i. | 281 | Infusion acidulated of savine, i. | 73 |
| Infusion of cinchona, i. | 487 | Infusion compound of leopard's-bane, i. | 280 | Infusion of semen contra, ii. | 582 |
| Infusion acidulated of cinchona, i. | 489 | Infusion ferruginous of leopard's-bane, i. | 281 | Infusion of senna, i. | 413 |
| Infusion compound of cinchona, i. | 488, 489 | Infusion of marsh ledum, ii. | 185 | Infusion of American senna, i. | 420 |
| Infusion magnesiated of cinchona, i. | 487 | Infusum vinous of leopard's bane, i. | 281 | Infusion of senna with lemon juice, i. | 414 |
| Infusion potassated of cinchona, i. | 488 | Infusion of lesser centaury, i. | 626 | Infusion of senna with tamarinds, i. | 415 |
| Infusion of cinchona prepared with heat, i. | 487 | Infusion of lime flowers, ii. | 691 | Infusion of serpentaria, i. | 278 |
| Infusion of cinchona prepared without heat, i. | 486 | Infusion of linseed, ii. | 201 | Infusion of simarouba, ii. | 587 |
| Infusion of cinnamon, ii. | 171 | Infusion of liquorice, i. | 728 | Infusion of anthelmintic spigelia, ii. | 620 |
| Infusion of clematis, i. | 523 | Infusion of mallow, ii. | 216 | Infusion of common spleenwort, i. | 443 |
| Infusion of cloves, i. | 399 | Infusion of malt, ii. | 213 | Infusion of spruce fir, i. | 22 |
| Infusion of corsican moss, ii. | 2 | Infusion of wild marjorum, ii. | 324 | Infusion of spurred rye, ii. | 579 |
| Infusion of thoroughstem cresswort, i. | 629 | Infusion of Mexican tea, i. | 457 | Infusion stomachic, i. | 488 |
| Infusion of digitalis, i. | 614 | Infusion of mint, ii. | 235 | Infusion of sweet-flag, i. | 87 |
| Infusion diuretic, i. | 269, 615 | Infusion compound of mint, ii. | 235 | Infusion of sweet-flag with chlorine, i. | 460 |
| Infusion of elder flowers, ii. | 553 | Infusion nervine, ii. | 706 | Infusion of tamarinds, ii. | 646 |
| Infusion of elder with tartarized antimony, i. | 260 | Infusion of nutgalls, i. | 693 | Infusion compound of tamarinds, ii. | 646 |
| Infusion of elm, ii. | 700 | Infusion aqueous of opium, ii. | 288 | Infusion of tansey, ii. | 650 |
| Infusion emmenagogue, i. | 712 | Infusion of orange leaves, i. | 510 | Infusion of thyme, ii. | 689 |
| Infusion of purging flax, ii. | 201 | Infusion compound of orange peel, i. | 510 | Infusion of tobacco, ii. | 262 |
| Infusion compound of gentian, i. | 711 | Infusion of Pareira brava, i. | 505 | Infusion tonic, i. | 712, 488 |
| Infusion of germander, ii. | 681 | Infusion of peach flowers, ii. | 344 | ii. | 127 |
| Infusion of gratiola, i. | 731 | Infusion pectoral, i. | 147, 721, 336 | Infusion of valerian, ii. | 706 |
| Infusion of ground ivy, i. | 721 | ii. | 329, 186 | Infusion compound of valerian, ii. | 706 |
| Infusion of gum arabic paste, i. | 753 | Infusion of peony, ii. | 329, 186 | Infusion vermifuge, ii. | 620, 683 |
| Infusion of heart's-ease, ii. | 724 | Infusion of pepperwort, ii. | 186 | Infusion of veronica, ii. | 718 |
| Infusion of hemlock, i. | 466 | Infusion of polygala amara, ii. | 404 | Infusion of violet, ii. | 722 |
| | | Infusion of red poppy, ii. | 335 | Infusion of water-lily, ii. | 268 |
| | | Infusion of quassia, ii. | 473 | Infusion of wood-lice, ii. | 272 |
| | | | | Infusion of wormwood, i. | 25 |

Infusion of wormwood with nitre, i.	25	Infusum cinchonæ frigidum, i.	486	Infusum turionum pini, i.	22
Infuso-decoction Bremer's of willow, ii.	543	Infusum cinchonæ lancifoliae, i.	486	Infusum quassiae, ii.	473
Infusum absinthii nitrata, i.	25	Infusum cinchonæ cum magnesia usta, i.	487	Infusum quassiae excelsae, ii.	473
Infusum alexiterium, i.	278	Infusum cinchonæ cum succo limorum, i.	489	Infusum quassiae cum natro, ii.	607
Infusum amarum, i.	711	Infusum colombæ, i.	528	Infusum quassiae cum sulphate zinci, ii.	473
Infusum amarum cum nitro, i.	712	Infusum conii maculati, i.	466	Infusum raphani, ii.	485
Infusum amarum et rorismarini, i.	712	Infusum corticis Peruviani vinosum, i.	496	Infusum rhabarbari, ii.	494
Infusum amarum cum natro, ii.	606	Infusum cuspariæ, i.	229	Infusum rhei, ii.	494
Infusum amarum vinosum, i.	713	Infusum digitalis, i.	614	Infusum rhei alkalisatum, ii.	497
Infusum angusturæ, i.	229	Infusum digitalis purpureæ, i.	614	Infusum rhei amarum, ii.	495
Infusum anthelminticum, ii.	2, 620	Infusum dulcamaræ, ii.	616	Infusum rhei boraxatum, ii.	498
Infusum anthemidis, i.	451	Infusum eupatorii perforiati, i.	629	Infusum rhei salinum, ii.	498
Infusum anthemidis nobilis, i.	451	Infusum flammulæ Jovis, i.	523	Infusum rhei cum subcarbonate potassæ, ii.	497
Infusum armoraciæ, ii.	485	Infusum gallarum, i.	693	Infusum rhododendri chrysanthi, ii.	503
Infusum armoraciæ compositum, ii.	485	Infusum gentianæ compositum, i.	711	Infusum reborans, i.	712
Infusum aurantii compositum, i.	510	Infusum glandium quercus, tostorum, ii.	477	Infusum rosæ acidulum, ii.	512
Infusum bruceæ, i.	229	Infusum radicis glycyrrhizæ, i.	728	Infusum rosæ compositum, ii.	512
Infusum bynes, ii.	213	Infusum gratiolæ, i.	731	Infusum rosæ Gallicæ, ii.	513
Infusum calami aromatici, i.	87	Infusum herbæ jaceæ, ii.	724	Infusum rosarum, ii.	512
Infusum calami aromatici cum acido muriatico oxygenato, i.	460	Infusum ipecacuanhæ vinosum, ii.	128	Infusum sabinæ, ii.	143
Infusum calcis cinchonatum, i.	488	Infusum juglandis compositum, ii.	136	Infusum sabinæ cum acido phosphorico, i.	75
Infusum calumbæ, i.	528	Infusum baccarum juniperi, ii.	140	Infusum salep cum adstringentibus, ii.	609
Infusum cardiacum, i.	278	Infusum kali carbonici cum scilla vinosum, ii.	571	Infusum salviæ, ii.	545
Infusum cardiacum acetatum, i.	278	Infusum laxativum simplex, i.	415	Infusum salviæ vinosum, ii.	545
Infusum cardui benedicti, i.	525	Infusum laxativum Vienne, i.	415	Infusum sanguinariæ canadensis, ii.	555
Infusum caryophyllorum, i.	399	Infusum lepidii, ii.	486	Infusum seminis santonici, ii.	582
Infusum cascarillæ, i.	569	Infusum lini, ii.	201	Infusum scilliticum vinosum compositum, ii.	570
Infusum cassiæ marylandicæ, i.	420	Infusum lini cathartici, ii.	201	Infusum secalis cornuti, ii.	579
Infusum cassiæ sennæ, i.	413	Infusum lini usitatissimi, ii.	201	Infusum florum sambuci nigrae, ii.	553
Infusum cathartico-anodynum, i.	452	Infusum liquiritæ, i.	728	Infusum sennæ, i.	413
Infusum centaureæ bedictæ, i.	525	Infusum malti, ii.	213	Infusum sennæ compositum sine aromaticis, i.	415
Infusum centaurii minoris, i.	626	Infusum menthæ, ii.	235	Infusum sennæ citratum, i.	414
Infusum foliorum ceterach, i.	443	Infusum millepedum, ii.	292	Infusum sennæ compositum, i.	413, 415
Infusum chamæmeli, i.	451	Infusum nicotianæ tabaci, ii.	262	Infusum foliorum cassiæ sennæ compositum, i.	415
Infusum chamomillæ, i.	451	Infusum florum nymphaeæ, ii.	268	Infusum sennæ cum manna, i.	415
Infusum chenopodii ambrosioides, i.	457	Infusum orchidis, ii.	541	Infusum sennæ tamarindis, i.	415
Infusum chinæ, i.	487	Infusum ad paralysim, ii.	485	Infusum sennæ tartarissatum, ii.	415
Infusum cinchonæ, i.	487	Infusum paralyticum, ii.	485	Infusum simarubæ, ii.	587
Infusum cinchonæ cum aqua calcis, i.	488	Infusum pectorale, i.	721		
Infusum cinchonæ sine calore, i.	486	Infusum persicarum, ii.	344		
		Infusum picis empyreumaticæ liquidæ, ii.	365		
		Infusum picis liquidæ, ii.	365		

Infusum seminum lini, i. 201	Injection antigonorrhœic, i. 64	Iodate of lime, ii. 121
Infusum seminum sina- peos, ii. 485	Injection antiseptic, ii. 543	Iodate of potass, ii. 121
Infusum tabaci, ii. 262	Injection astringent, i. 162	Iodine, ii. 120
Infusum tamarindorum, ii. 646	492, 588, 653, 693, ii. 16, 146, 377, 387, 388, 476	Iodinium, ii. 120
Infusum tamarindorum, compositum, ii. 646	Injectio balsamic, i. 555	Ioduretted hydriodate of potass, ii. 122
Infusum tanacetii, ii. 650	Injection against carnos- ities of the urethra, ii. 49	Ioduret of potassium, ii. 122
Infusum tonico nervo- sum, ii. 706	Injection contrastimu- lant, i. 62	Iodurets of mercury, ii. 79
Infusum valerianæ, ii. 706	Injection detergent, ii. 537	Ipecacuanha, ii. 124
Infusum valerianæ com- positum, ii. 706	Injection detersive, i. 122, 134, 592, 763, ii. 69	Ipecacuanha bastard, ii. 696
Infusum valerianæ vino- sum, ii. 708	Injection diuretic, i. 614	Ipecacuanha spurge, i. 629
Infusum ulmi, ii. 700	Injection emmenagogue, i. 171	Ireos nostratis radix, ii. 130
Injectio anodyna, ii. 293	Injection emollient, ii. 338	Ireos palustris radix, ii. 130
Injectio cum aqua lauro- cerasi, i. 62	Injection Girtanner's, ii. 414	Ireos radix, ii. 131
Injectio auricularis deter- siva, ii. 537	Injection of isinglass and myrrh, i. 701	Iridis Florentinæ radix, ii. 131
Injectio chinæ, i. 492	Injection lithontriptic, ii. 434, 606	Iris Florentina, ii. 131
Injectio chinæ antiputrida, i. 491	Injection mercurial, ii. 34, 69	Iris fœtidissima, ii. 131
Injectio contrastimulans, i. 62	Injection simple mercu- rial, ii. 15	Iris Germanica, ii. 130
Injectio cuprata, i. 587	Injection opiate mercu- rial, ii. 76	Iris pseudacorus, ii. 130
Injectio detergens, i. 763	Injection mercurio-satur- nine, ii. 77	Iris tuberosa, ii. 5
Injectio detergens et ro- borans, i. 134	Injection of mercury and turpentine, ii. 18	Iris versicolor, i. 131
Injectio fellea, i. 633	Injection mucilaginous, ii. 201	Iris Florentine, ii. 131
Injectio ad fistulas, i. 349, 380	Injection narcotic, ii. 102, 259	Iris yellow, ii. 130
Injectio ex gallis, i. 693	Injection opiate, ii. 293	Iron, i. 634
Injectio hydrargyri mu- riatis camphorata, ii. 66	Injection against phy- mosis, i. 585	Iron filings prepared, i. 634
Injectio hydrargyro-sa- turnina, ii. 77	Injection of protochlo- ruret of mercury, ii. 49	Isatine, ii. 133
Injectio hydrargyro-te- rebinthinata, ii. 18	Injection saturnine, ii. 370	Isatis herba, ii. 133
Injectio lactis opiatæ, ii. 293	Injection resolvent, ii. 386	Isatis tinctoria, ii. 133, 351
Injectio ad leucorrhœam, ii. 377	Injection sedative, ii. 246, 279, 282, 293, 294, 379	Isinglass, i. 696
Injectio leniens, ii. 246	Injection soothing, ii. 10,	Isolusine, ii. 403
Injectio mercurialis Plen- kii, ii. 15	Injection stimulant, i. 380, 349, 633	Ivæ arthriticæ herba, ii. 681
Injectio ex muriate hy- drargyri, ii. 49	Injectio terebinthinata, ii. 656	Ivæ herba, ii. 682
Injectio in necrosi syphi- litica, ii. 69	Injection tonic, i. 491	Juglans cinerea, ii. 134
Injectio de plumbo et mercurio, ii. 77	Injection Whately's, ii. 71	Juglans regia, ii. 134
Injectio terebinthinacea, ii. 656	Ink-powder, i. 653	Juice of apples, ii. 214
Injectio vaginalis emme- nagoga, i. 171	Intybi angusti herba, ii. 152	Juice of barberries, i. 335
Injectio vaginalis cum mercurio, ii. 34	Intybi hortensis herba, i. 461	Juice of beccabunga, ii. 719
Injection acoustic, i. 321	Inspissamentum cucume- ris agrestis, ii. 243	Juice of blueberries, ii. 702
Injection compound alu- minous, ii. 388	Inspissamentum urtica- rum, ii. 701	Juice of borage, i. 341
Injection of ammoniacal copper, i. 582	Inula dysenterica, ii. 114	Juice of bryony, i. 344
	Inula helenium, ii. 114	Juice of buckthorn, ii. 487
	Inuline, ii. 115	Juice of bugloss, i. 215
	Job's tears, i. 538	Juice of carrot, i. 607
		Juice of cherries, i. 438
		Juice of chervil, i. 444
		Juice of cichory, i. 462
		Juice of cranberries, ii. 702
		Juice of cucumber, i. 578
		Juice of black currants, ii. 505
		Juice of red currants, ii. 79
		Juice diuretic, ii. 482
		Juice diuretic and cooling, i. 445
		Juice of dandelion, ii. 651
		Juice of dog-grass, ii. 697
		Juice of dwarf elder ber- ries, ii. 551
		Juice of elaterium, ii. 243
		Juice of elder berries, ii. 551
		Juice of fumitory, i. 680
		Juice of hemlock, i. 464
		Juice of horehound, ii. 222

Juice of kermes, i.	530	Julep of mithridate, ii.	306	Kermes, i.	530
Juice of lemon, i.	520	Julep refreshing, i.	521	Kermes grains, i.	530
Juice of liquorice, i.	723	Julep of rose, ii.	518	Kermes minerale, i.	243
Juice of liquorice anisated, i.	723	Julep scillitic, ii.	576	Keyrini flores, i.	455
Juice of liquorice with gum, i.	723	Julep sedative, ii.	387	Kininum, ii.	477
Juice of liquorice purified, i.	723	Julep sedative and purgative, ii.	436	Kino, ii.	145
Juice of lungwort, ii.	469	Julep succinated, ii.	630	Knapweed common, ii.	112
Juice of milfoil, i.	35	Julep temperant, ii.	436	Knotgrass, ii.	406
Juice of mulberries, ii.	248	Julep violet, ii.	721	Krameria triandria, ii.	147
Juice of orange, i.	507	Junci odorati herba, i.	217	Labdanum, i.	757
Juice of peach flowers, ii.	344	Juniper, ii.	137	Lac, ii.	150
Juice of plantain, ii.	366	Juniper Virginian, ii.	141	Lac ammoniacale, i.	742
Juice of pomegranate, ii.	471	Juniperus communis, ii.	137, 555	Lac ammoniaci, i.	742
Juice of quince, i.	599	Juniperus lycia, ii.	271, 555	Lac amygdalæ, i.	207
Juice of raspberries, ii.	532	Juniperus oxycedrus, ii.	555	Lac asininum, ii.	149
Juice of pale roses, ii.	510	Juniperus phœnicea, ii.	271	Lac asininum artificiale, i.	701
Juice of scurvy-grass, i.	531	Juniperus sabina, ii.	142	Lac assafoetidæ, i.	303
Juice compound of scurvy-grass, i.	533	Juniperus thurifera, ii.	271	Lac bubulum, ii.	149
Juice of self-heal, ii.	466	Juniperus Virginiana, ii.	141	Lac caprinum, ii.	149
Juice of soapwort, ii.	557	Ivory, i.	360	Lac ebutyratum, ii.	149
Juice of sorrel, ii.	534	Ivory fossile, i.	360	Lac guaiacinum, i.	739
Juice of wood sorrel, ii.	327	Jupiter, ii.	623	Lac gummi guaiaci, i.	739
Juice of strawberries, i.	677	Jus millepedum, ii.	272	Lac lunæ, i.	352
Juice of violet, ii.	722	Jus ossium, i.	699	Lac sulphuris, ii.	633
Julapium acidulatum, i.	97	Jusculum helecinum, i.	700	Lac terebinthinaceum, ii.	655
Julapium acidum, i.	77	Jusculum pulli gallinacii lientericum, i.	703	Lac terræ, ii.	206
Julapium æthereum, i.	101	Jusculum siccum, i.	705	Lac vaccinum, ii.	149
Julapium e camphora, i.	369	Jusculum viperæ, i.	700	Lac Virginis, i.	330
Julapium e camphora cum aceto, i.	370	Jusculum viperinum, i.	700	Lacca, ii.	150
Julapium camphoratum, i.	369	Jusquiami radix, ii.	99	Lacca cærulea, ii.	694
Julapium mithridaticum, ii.	306	Justicia adhatoda, ii.	144	Lacca musica, ii.	694
Julapium moschatum, ii.	252	Justicia pectoralis, ii.	144	Laccin, ii.	150
Julapium moschi, ii.	252	Ivy ground, i.	720	Lauro-cerasi folia, i.	439
Julapium regium, ii.	518	Izarinum, ii.	529	Lacrymæ Job semen, i.	538
Julapium rosarum, ii.	518	Kadmium, ii.	145	Lacerta agilis, i.	697
Julapium rosatum, ii.	518	Kæmpferia Galanga, i.	682	Lactuca sativa, ii.	152
Julapium sambucinum, ii.	554	Kali acetatum, ii.	441	Lactuca scariola, ii.	152
Julapium sistens, ii.	319	Kali carbonicum, ii.	419	Lactuca virosa, ii.	152
Julapium stomachicum, ii.	425	Kali causticum siccum, ii.	412	Lactucarium, ii.	154
Julapium succinatum, ii.	630	Kali citratum, ii.	446	Ladies' bedstraw, i.	691
Julapium violarum, ii.	721	Kali nitricum, ii.	432	Ladies' hair, i.	89
Julep antihysterical, ii.	538	Kali oxalicum acidulum, ii.	446	Ladies' mantle, i.	104
Julep antiphthisical, i.	584	Kali præparatum, ii.	419	Ladies' thistle, i.	395
Julep balsamic, i.	324	Kali subcarbonicum, ii.	410	Lady-bird, i.	526
Julep diuretic, ii.	668	Kali subcarbonicum purissimum, ii.	420	Lady-cow, i.	526
Julep of elder, ii.	554	Kali sulphurato-stibiatum, i.	254	Lagophthalmi herba, i.	719
Julep emmenagogue, i.	556	Kali sulphuratum, ii.	416	Lamii montani herba, ii.	232
Julep fetid, i.	303	Kali sulphuricum, ii.	427	Lamium album, ii.	154
Julep hydragogue, i.	265	Kali tartarisatum, ii.	447	Lapathi aquatici radix, ii.	534
		Kali vegetabile causticum, ii.	410	Lapathi crispi herba, ii.	535
		Kali vegetabile purum, ii.	410	Lapathi unctuosius herba, i.	457
		Kali vitriolatum, ii.	427	Laphæ radix, i.	291
		Kalmia latifolia, ii.	145	Lapides cancerorum, i.	354
		Kekekunemalo, ii.	147	Lapides cancerorum acetati, i.	363
				Lapides cancerorum citrati, i.	364
				Lapides percarum, i.	354

Lapides spongiarum, i.	354	Laudanum opiatum cydoniatum, ii.	291	Lavement sedative, ii.	102
Lapilli cancrorum, i.	354	Laudanum opiatum Rousseau, ii.	282	278, 292, 293, 338,	656
Lapis armenia, i.	582	Laudanum Rousseau's, ii.	282	Lavement starch, i.	214
Lapis auris ceti, i.	360	Laudanum simplex, ii.	289	Lavement stimulant, i.	492
Lapis carptionis, i.	354	Laudanum Sydenham's, ii.	277	Lavement terebinthinate, ii.	656
Lapis causticus, ii.	412	Laudanum urinarium, ii.	291	Lavement terebinthinated, ii.	667
Lapis divinus, i.	585	Laudanum Warner's, ii.	286	Lavement tonic, i.	484, 492
Lapis infernalis, i.	273	Lavement acetose, i.	42	Lavement vermifuge, i.	375
Lapis judaicus, i.	354	Lavement acid, i.	73	ii.	508, 603, 650
Lapis infernalis alkalinus, ii.	412	Lavement aloetic, i.	118	Lavender, ii.	155
Lapis lazuli, i.	157	Lavement anodyne, ii.	283, 338	Lavender French, ii.	159
Lapis lycis, i.	354	Lavement anthelmintic, ii.	706	Lavender sea, ii.	625
Lapis manati, i.	360	Lavement antiperiodic, i.	490	Laurel mountain, ii.	145
Lapis medicamentosus, i.	160	Lavement antispasmodic, i.	303, 452, ii. 252, 283,	Laurel tree, ii.	178
Lapis miraculosus, i.	586	Lavement antisyphilitic, ii.	69	Lauri Alexandrini herba, ii.	536
Lapis nephreticus, ii.	260	Lavement against ascari-des, ii.	714	Laurus cassia, ii.	177
Lapis ophthalmicus, i.	585	Lavement astringent, i.	77	Laurus cinnamomum, ii.	160
Lapis porcinus, i.	338, 352	693, ii. 319, 407, 471,	656	Laurus culilawan, ii.	178
Lapis prunellæ, ii.	433	Lavement astringent and tonic, i.	280	Laurus nobilis, ii.	178
Lapis pumicis, ii.	470	Lavement of belladonna, i.	311	Laurus sassafras, ii.	183
Lapis septicus, ii.	412	Lavement butyraceous, i.	317	Lawsonia inermis, ii.	184
Lapis specularis, i.	362	Lavement camphorated, i.	371, 373	Lawsonia spinosa, ii.	185
Lapis vulnerarius externus, i.	586	Lavement carminative, i.	179, 356	Lazulite, i.	157
Lappæ majoris radix, i.	272	Lavement of chamomile, i.	452	Leach medicinal, ii.	6
Lappulæ hepaticæ herba, i.	103	Lavement of cinchona, i.	489	Lead, ii.	368
Lard, i.	315, 316	Lavement demulcent laxative, i.	421	Lead black, i.	639
Lard hog's, i.	316	Lavement emollient, i.	148	Lead red, ii.	370
Larix Europæa, ii.	218, 654	681, ii.	202, 216, 326	Lead white, ii.	371
Larkspur, i.	608	Lavement irritant, ii.	262	Leadwort, ii.	368
Laserpitii Germanici, ii.	197	Lavement mercurial, ii.	49, 60, 68	Ledum marsh, ii.	185
Laserpitium latifolium, ii.	155	Lavement nutritive, i.	700	Ledum palustre, ii.	185
Laserpitium siler, ii.	155	Lavement oily, ii.	270	Leek, i.	113
Lasserwort broad-leaved, ii.	155	Lavement opiate, ii.	293	Leminthocorton, ii.	2
Lathyridis majoris semines, i.	629	Lavement oxymellated, i.	44	Lemon grass, i.	216
Lavandula spica, ii.	155	Lavement of poppy, ii.	338	Lemon tree, i.	516
Lavandula stœchas, ii.	159	Lavement purgative, i.	259, 265, 419, 731, ii. 212, 241,	Lemonade alcoholic, i.	111
Laudanum antihysteri-c, ii.	291	603, 612		Lemonade antimonial, i.	260
Laudanum cinnabarinum, ii.	45	Lavement refreshing, i.	579	Lemonade dry, i.	52
Laudanum compositum, ii.	277	Lavement refrigerant, ii.	435	Lemonade laxative, ii.	454
Laudanum crocatum, ii.	277	Lavement resolvent, i.	452	Lemonade mineral, i.	55
Laudanum diureticum, ii.	291	Lavement saponaceous, ii.	459	Lemonade nitric, i.	69
Laudanum Lalouette's, ii.	276			Lemonade phosphoric, i.	73
Laudanum liquidum, ii.	277, 281			Lemonade powder, i.	82
Laudanum liquidum Sydenhami, ii.	277			Lemonade vegetable, i.	520
Laudanum liquidum War-neri, ii.	286			Linimentum ceræ sim-plex, i.	435
Laudanum opiatum, ii.	276, 289, 291			Lentil, i.	631
				Leontopodii radix, i.	104
				Leonurus cardiaca, ii.	185
				Leopard's bane, i.	279, 618
				Lepidium iberis, ii.	186
				Lepidium latifolium, ii.	185
				Lepus combustus, i.	391
				Lepus timidus, i.	355, 697
				Lettuce garden, ii.	152
				Lettuce hemlock, ii.	152
				Lettuce-opium, ii.	154

Lettuce strong-scented, ii.	152	Lily of the valley, i.	541	Linguae bovis radix, i.	215
Lettuce virous, ii.	152	Lily white, ii.	198	Linguae cervinae herba, ii.	577
Leucanthemi odorati flores, i.	446	Lily white water, ii.	267	Linguae sepentariae herba, ii.	274
Leucosi lutei, i.	455	Lily yellow water, ii.	268	Liniment æthereous, ii.	179, 462
Levistici radix, ii.	197	Limonada sicca, i.	52	Liniment sedative æthereal, ii.	288
Ley juniper, ii.	423	Limonii radix, ii.	625	Liniment stimulant æthereous, i.	381
Ley soap-boilers, ii.	601	Limatura chalybis, i.	634	Liniment alkaline, ii.	424
Libanus thurifera, ii.	271	Limatura ferri, i.	634	Liniment aluminous, i.	163
Lichen, ii.	186, 187	Limatura ferri alcoholi-sata, i.	634	Liniment of ammonia and of tartarized antimony, i.	262
Lichen ash-coloured ground, ii.	186	Limatura ferri præparata, i.	634	Liniment ammoniacal, i.	177
Lichen cinereus terrestris, ii.	186	Limatura martis, i.	634	Liniment camphorated ammoniacal, i.	174, 189
Lichen cocciferous, ii.	186	Limatura martis depurata, i.	634	Liniment compound ammoniacal, i.	179
Lichen common yellow, ii.	187	Limatura stanni, ii.	623	Liniment fetid ammoniacal, i.	179
Lichen funnel-shaped, ii.	187	Lime quick, i.	347	Liniment saponaceous ammoniacal, i.	179
Lichen hairy oak, ii.	187	Lime slaked, i.	347	Liniment ammoniacal æthereal, i.	182
Lichen Iceland, ii.	187	Lime tree, ii.	691	Liniment amygdaline, i.	206
Lichen Islandicus, ii.	187	Lime water, i.	347	Liniment of ammoniuret of mercury, ii.	81
Lichen slow tree, ii.	188	Lime water with cinchona, i.	349	Liniment anodyne, i.	179
Lichen snow-white, ii.	186	Lin, ii.	200	ii. 101, 279, 285, 295	
Lichen terrestris cinereus, ii.	186	Linaria cymbalaria, ii.	199	Liniment anthelmintic, i.	572
Lichenis petræi latifolii herba, ii.	221	Linaria spuria, ii.	199	Liniment anti-arthritic, i.	190
Lichenis stellati herba, ii.	221	Linaria vulgaris, ii.	199	ii.	672
Lignum agallochi veri, ii.	192	Linctus acidus, i.	76, 600	Liniment antihæmorrhoidal, i.	440, ii. 107, 278, 658
Lignum agallochum, ii.	192	Linctus amygdalinus, i.	210	Liniment antihysterical, ii.	382
Lignum aquilæ, ii.	193	Linctus aperiens, ii.	219	Liniment anti-ophthalmic, i.	645, ii. 373
Lignum aspalathi, ii.	193	Linctus albus, i.	210	Liniment antipleuritic, ii.	463
Lignum benedictum, i.	732	Linctus ad aphthas, ii.	609	Liniment antipsoric, i.	350
Lignum Brasilianum rubrum, ii.	193	Linctus de borace, i.	609	Liniment antirheumatic, ii.	676
Lignum Brasilense, ii.	193	Linctus cetacei, i.	443	Liniment antispasmodic, ii.	279
Lignum cæruleum, ii.	193	Linctus communis, i.	216, 512	Liniment aphrodisiac, i.	321
Lignum Campescanum, ii.	193	Linctus demulcens, i.	211	Liniment stimulant aromatic, i.	381
Lignum Campechianum, ii.	193	Linctus emolliens, i.	211	Liniment arsenical, i.	285
Lignum cedrinum, ii.	137	Linctus expectorans, ii.	575	Liniment barytic, i.	328
Lignum colubrinum, ii.	626	Linctus gelatinosus nutritiens, i.	704	Liniment black, i.	690
Lignum ebenum, ii.	194	Linctus ad gingivas, i.	763	Liniment of camphor, i.	374
Lignum Fernambuci, ii.	193	Linctus laxans, i.	422	Liniment camphorated ammoniacal, i.	178
Lignum Indicum, i.	732	Linctus mannatus, ii.	219	Liniment camphorated saponaceous, i.	374
Lignum moluccanum, i.	570	Linctus olei terebinthinæ, ii.	669	Liniment stimulant camphorated, i.	381
Lignum nephriticum, ii.	195	Linctus oleosus, i.	211	Liniment of cantharides, i.	382
Lignum Rhodii, ii.	195	Linctus oxymuriatis calci, i.	352	Liniment ammoniacal of cantharides, i.	881
Lignum sanctum, i.	732	Linctus saponatus, ii.	459		
Lignum santali, ii.	195	Linctus ad stomacacem, i.	352, 460		
Lignum santali albi, ii.	196	Linctus borated, ii.	609		
Lignum santali citrini, ii.	196	Linctus camphorated, i.	368		
Lignum santali rubri, ii.	195	Linctus commen, i.	443		
Lignum Sappan, ii.	193	Linctus for cough, ii.	280		
Lignum visci, ii.	724	Linctus demulcent, i.	443		
Ligusticum levisticum, ii.	197	Linctus detersive, i.	763		
Ligusticum meum, ii.	197	Linctus emetic, i.	262		
Ligustri Ægyptiaci radix, ii.	185	Linctus emollient, i.	211		
Ligustrum vulgare, ii.	198	Linctus expectorant, i.	251		
Lilium candidum, ii.	198	Linctus refrigerant, ii.	434		
		Linctus saponaceous, ii.	459		
		Linctus vermifuge, ii.	705		
		Linden tree, ii.	691		
		Linguae avis semen, i.	678		

- Liniment composing, i. 374
 Liniment for cynanche, i. 204
 Liniment demulcent, ii. 584
 Liniment depilatory, i. 282
 Liniment diuretic, ii. 140, 568, 668
 Liniment emollient, i. 213
 Liniment excitant, ii. 283
 Liniment gingival, i. 761
 Liniment of gum ammoniac, i. 743
 Liniment hydrargyroacetose, ii. 96
 Liniment irritating, i. 182
 Liniment of juniper, ii. 140
 Liniment of lime water, i. 349
 Liniment of lime water with opium, i. 349
 Liniment mammillary, i. 321
 Liniment ammoniacal mercurial, ii. 25
 Liniment camphorated ammoniacal mercurial, ii. 25
 Liniment opiate mercurial, ii. 26
 Liniment narcotic, ii. 278
 Liniment opiate, ii. 278, 283, 292
 Liniment camphorated opiate, i. 375, ii. 279, 285
 Liniment saponaceous opiate, ii. 279
 Liniment of opium with calf's gall, ii. 297
 Liniment of petroleum, ii. 345
 Liniment phosphorated, ii. 350
 Liniment phosphorated of camphor, ii. 350
 Liniment purgative, i. 572
 Liniment quercico-saturnine, ii. 389
 Liniment resolvent, i. 178, 189, 322, 352, 368, 386, 633, 679
 Liniment rheumatic, i. 180
 Liniment irritating saponaceous, i. 381, ii. 461
 Liniment sedative, ii. 106, 292
 Liniment camphorated sedative, ii. 285
 Liniment spirituous, i. 188
 Liniment stimulant, i. 178, 372, 378, 386, ii. 144, 256
 Liniment of subacetate of copper, i. 592
 Liniment of subcarbonate of ammonia, i. 189
 Liniment of sulphuret of lime, i. 351
 Liniment sulphuro-saponaceous, ii. 602
 Liniment of tar, ii. 365
 Liniment stimulant terebinthine, i. 381
 Liniment for tinea, i. 179
 Liniment of tobacco, ii. 263
 Liniment of turpentine, ii. 669
 Liniment æthereal of turpentine, ii. 664
 Liniment vermifuge, ii. 650
 Linimentum acetatis cuprici, i. 591
 Linimentum æruginis, i. 591
 Linimentum æthereum, ii. 179, 462
 Linimentum album, i. 441
 Linimentum aluminosum, i. 162
 Linimentum ad ambustiones cum opio, i. 349
 Linimentum ammoniæ, i. 177
 Linimentum ammoniæ fortius, i. 177
 Linimentum ammoniato-æthereum, i. 381
 Linimentum ammoniato-aromaticum, i. 381
 Linimentum ammoniatum, i. 177
 Linimentum ammonii opiatum, i. 179
 Linimentum anglicanum, i. 177
 Linimentum anodynum, i. 179, ii. 286, 461
 Linimentum anti-arthritic, i. 178
 Linimentum anti-arthriticum, i. 320
 Linimentum antihæmorrhoidale, ii. 199
 Linimentum antiherpeticum, i. 639
 Linimentum antipleuriticum, ii. 463
 Linimentum antispasticum, i. 179
 Linimentum ad aphthas, ii. 609
 Linimentum ex aqua calcis, i. 349
 Linimentum aquæ calcis, i. 349
 Linimentum aromaticum, ii. 157
 Linimentum aromaticum stimulans, ii. 524
 Linimentum ad arthrodyniam, i. 386
 Linimentum balsami Peruviani, i. 320
 Linimentum de borate, ii. 609
 Linimentum calcaris sulphuratæ, i. 351
 Linimentum calcicum, i. 349
 Linimentum calcis, i. 349
 Linimentum camphoræ, i. 374
 Linimentum camphoræ compositum, i. 174
 Linimentum camphoratum, i. 374
 Linimentum phosphoratum camphoratum, ii. 350
 Linimentum volatile camphoratum, i. 178
 Linimentum cepæ, i. 113
 Linimentum ad combustiones, i. 348
 Linimentum diureticum gastro-scliticum, ii. 568
 Linimentum farinaceum cum pice, ii. 365
 Linimentum ex felle tauri, i. 633
 Linimentum gingivale, i. 56
 Linimentum gingivale mundificans, i. 761
 Linimentum gummoso-balsamicum, i. 321
 Linimentum hydrargyri, ii. 25
 Linimentum irritans, i. 381
 Linimentum juniperinum, ii. 140
 Linimentum opiatum, ii. 461
 Linimentum petrolei ammoniatum, i. 178
 Linimentum plumbatum, ii. 380
 Linimentum psoricum, ii. 639
 Linimentum resolvens, ii. 670
 Linimentum resolvens et diureticum, ii. 140
 Linimentum rubefaciens, ii. 588
 Linimentum saponaceum i. 179, ii. 460, 461
 Linimentum saponaceum camphoratum, ii. 462
 Linimentum saponaceum hydrosulphuretum, ii. 417
 Linimentum saponato-camphoratum, i. 179
 Linimentum saponis, ii. 460, 461, 463
 Linimentum saponis camphoratum, ii. 286, 461
 Linimentum saponis compositum, i. 179, ii. 286, 461
 Linimentum saponis et opii, ii. 286
 Linimentum saponis cum opio, ii. 461
 Linimentum saturnino-quercicum, ii. 389

Linimentum stimulans, ii.	671	Liquor ammonii hydrothiodis, i.	194	Liquor digestivus Boerhaavii, ii.	442
Linimentum e sulphure, ii.	637	Liquor ammonii pyrooleosi, i.	185	Liquor discutiens, i.	44
Linimentum tabaci, ii.	263	Liquor ammonii pyrooleosi succinici, i.	200	Liquor enixus, i.	190
Linimentum terebinthinæ, ii.	669	Liquor ammonii sulphureti, i.	194	Liquor excitans, i.	180
Linimentum volatile, i.	177	Liquor ammonii tartarici, i.	202	Liquor excitorius, i.	382
Linimentum volatile cum camphora, i.	178	Liquor ammonii vinosus, i.	172	Liquor ferri alkalini, i.	647
Linseed, ii.	200	Liquor anodynus animalis, i.	97	Liquor fumans Libavii, i.	625
Linum catharticum, ii.	200	Liquor antinephritic, ii.	338	Liquor hydrargyri camphorati, ii.	70
Linum usitatissimum, ii.	200	Liquor anodynus martialis, i.	659	Liquor hydrargyri muratici corrosivi, ii.	68
Liquamen martis, i.	657	Liquor anodynus mineralis aquosus, i.	100	Liquor hydrargyri nitrici, ii.	86
Liquamen myrrhæ, i.	762	Liquor anodynus mineralis Hoffmanni, i.	99	Liquor hydrargyri oxy-muriati, ii.	70
Liquamen salis tartari, ii.	422	Liquor anodynus mineralis martialis, i.	659	Liquor hydrogenii sulphuratus, i.	65
Liquidambar Orientale, ii.	628	Liquor anodynus vegetabilis, i.	92	Liquor hydrosulphuretico-acidulus, i.	65
Liquidambar styraciflua, ii.	627	Liquor anti-arthritis, i.	736	Liquor hydro-sulphureticus pro balneo, i.	66
Liquiritia cocta citrina, i.	726	Liquor antimonii tartarisati, i.	288	Liquor kali acetici, ii.	442
Liquiritiæ radix, i.	722	Liquor aperitivus, ii.	453	Liquor kali acetici cum rheo, ii.	445
Liquor acetatis ferri æthereus, i.	664	Liquor ad aphthas, ii.	609	Liquor kali carbonici, ii.	422
Liquor acidi muriatici oxygenati, i.	459	Liquor argenti muriatico-ammoniaci, i.	276	Liquor kali caustici, ii.	410
Liquor acidus Halleri, i.	79	Liquor aromaticus saponato-ammoniacus, i.	188	Liquor kali subcarbonici, ii.	422
Liquor æris cinnamomatus, i.	585	Liquor arsenicalis, i.	286, 287	Liquor martiales, i.	659
Liquor æruginis, i.	585	Liquor arthriticus Pottii, ii.	670	Liquor mercurialis, ii.	68
Liquor æthereus oleosus, i.	100	Liquor antimonii tartarisati, ii.	259	Liquor Mindereri, i.	195
Liquor æthereus sulphuricus, i.	99	Liquor baryticus, i.	327	Liquor muriatis deutoxydohydrargyro-calcarei, ii.	35
Liquor alexiterius acetatus, i.	42	Liquor Bellosti, ii.	86	Liquor muriatis ferri, i.	658
Liquor alexiterius oxygenatus, i.	459	Liquor caleis, i.	347	Liquor muriatis hydrargyri, ii.	68
Liquor alexiterius extemporaneus, i.	460	Liquor calcis muriatis, i.	363	Liquor muriatis hydrargyro-ammoniacalis, ii.	90
Liquor alkali volatilil compositus, i.	201	Liquor carbonatis ammoniæ, i.	183	Liquor myrrhæ, i.	762
Liquor aluminis composita, i.	162	Liquor cornu cervi acetatus, i.	195	Liquor ad nephritidem calculosam, ii.	338
Liquor ex ammonia et oleo succini, i.	180	Liquor cornu cervi alkalinus, i.	185	Liquor nervinus Bangii, i.	371
Liquor ammoniacalis cum aceto, i.	195	Liquor cornu cervi succinatus albus, i.	207	Liquor nitratis hydrargyri, ii.	86
Liquor ammoniæ, i.	168	Liquor cornu cervi tartarizatus, i.	202	Liquor olei animalis et cajeputi, ii.	270
Liquor ammoniæ acetatis, i.	195	Liquor cornu cervi terebinthinatus, i.	196	Liquor oleosus Sylvii, i.	174
Liquor ammoniæ anisatus, i.	173	Liquor ad condylomata, ii.	70, 78	Liquor ophthalmicus roborans, i.	135
Liquor ammoniæ puræ, i.	168	Liquor cupri ammoniaci, i.	587	Liquor ad parvum vel unquem oculorum consumendum, i.	595
Liquor ammoniæ spirituosus, i.	172	Liquor cupri ammonio-muriatici, i.	589	Liquor ad perniones, ii.	370
Liquor ammoniæ subcarbonatis, i.	184	Liquor cupri Kœchlini, i.	589	Liquor plumbi subacetatis dilutus, ii.	378
Liquor ammonii aceteci, i.	195			Liquor potassæ, ii.	410
Liquor ammonii caustici, i.	168				

Liquor potassæ arseniatis, ii.	286	Liquor oxymuriated alkaline, ii.	440	Liquor temperate cordial, ii.	169
Liquor potassæ causticæ, ii.	410	Liquor muriatico-hydrargyro-cupric ammoniacal, ii.	92	Liquor cosmetic i.	209, 331, 633, ii. 36, 610, 613
Liquor potassæ subcarbonatis, ii.	422	Liquor Fuller's anodyne, i.	188	Liquor dentifrice, ii.	231
Liquor probatorius Wirtembergicus, i.	283	Liquor succinated anodyne, ii.	632	Liquor digestive, ii.	658
Liquor pyrotartareus, i.	74	Liquor Vicat's anodyne, i.	174	Liquor Labarague's disinfecting, i.	351
Liquor salinus laxativus, ii.	211, 453	Liquor Eller's anti-arthritic, i.	202	Liquor diuretic, ii.	273
Liquor salis alkali vegetabilis acetati, ii.	442	Liquor anti-asthmatic, ii.	116, 288	Liquor camphorated diuretic, ii.	436
Liquor salis ammoniaci anisatus, i.	173	Liquor anti-gangrenous, i.	193	Liquor emetic, ii.	483
Liquor salis tartari, ii.	422	Liquor antigonorrhœic, ii.	667	Liquor epidemic, i.	224
Liquor saponis stibiati, i.	252	Liquor cardinal de Luynes antiherpetic, ii.	78	Liquor febrifuge, i.	653
Liquor splenicus ex tartaro, ii.	447	Liquor antihysterical, ii.	229	Liquor gummy ferruginous, i.	653
Liquor stibiatus cum acenito, i.	84	Liquor anti-melancholic, i.	565	Liquor fetid, i.	684
Liquor stypticus, i.	157, 657	Liquor ammoniated antimonial, i.	202	Liquor distilled of frog's spawn, i.	708
Liquor subacetatis lythargyri, ii.	375	Liquor anti-odontalgic, ii.	224	Liquor Hahnemann's i.	65
Liquor succinicus anodynus, ii.	632	Liquor antiphlogistic, i.	45	Liquor hydrargyro-acetic, ii.	96
Liquor syphiliticus Turneri, ii.	70	Liquor antipsoric, i.	362	Liquor hysteric, i.	450
Liquor tartari solubilis, ii.	456	Liquor antimiasmatic, i.	589, 590	Liquor laxative, ii.	453
Liquor tartari solubilis chalybeati, i.	673	Liquor antinephritic, ii.	606	Liquor Sanders' lithontriptic, ii.	414
Liquor tartari stibiati, i.	259	Liquor antiscorbutic, ii.	177	Liquor mercurial, ii.	86, 92
Liquor tartari tartarisati, ii.	448	Liquor antispasmodic, i.	60, 201	Liquor aloetic mercurial, ii.	35
Liquor tartari vegetabilis, ii.	442	Liquor Grimaud's antispasmodic, i.	202	Liquor aluminous mercurial, ii.	74
Liquor tartratis ammoniaci, i.	202	Liquor antisiphilitic, ii.	70, 71	Liquor Plenk's balsamic mercurial, ii.	63
Liquor terræ foliatæ, ii.	442	Liquor Chaussier's antisiphilitic, ii.	81	Liquor camphorated mercurial, ii.	70
Liquor terræ ponderosæ salitæ, i.	326	Liquor antisiphilitic of Rouen, ii.	595	Liquor Gohl's grey mercurial, ii.	86
Liquor vini probatorius Hahnemanni, i.	65	Liquor apoplectic, ii.	528	Liquor mercurio-cuprous, i.	590
Liquor ad ulcera atonica, ii.	440	Liquor compound aromatic, ii.	682	Liquor Pereboom's nerve vine, ii.	672
Liquor ad ulcera faucium, i.	763	Liquor spirituous aromatic, ii.	255	Liquor odoriferous, ii.	159
Liquor ad ulcera oris, i.	592	Liquor Pott's arthritic, ii.	670	Liquor ophthalmic, i.	135, 596, ii. 375, 377, 387
Liquor volatilis cornu cervi, i.	185	Liquor astringent, i.	157	Liquor anodyne ophthalmic, i.	767
Liquor vulnerarius, ii.	665	Liquor astringent aromatic, ii.	667	Liquor Conrad's ophthalmic, ii.	76
Liquor vulnerarius mineralis, i.	670	Liquor catarrhal, ii.	169	Liquor mercurial ophthalmic, ii.	68
Liquor vulnerarius vegetabilis Dippelii, i.	37	Liquor Plenk's caustic, ii.	78	Liquor resolvent ophthalmic, ii.	414
Liquor acoustical, ii.	710	Liquor Charles the fifth's cephalic, ii.	169	Liquor black phagedenic, ii.	34
Liquor oily æthereal, i.	100	Liquor chalybeate, i.	673	Liquor compound phagedenic, ii.	69
Liquor sulphureous æthereal, i.	98	Liquor nitric æthereal chalybeate, i.	656	Liquor Grindel's phagedenic, ii.	71
		Liquor against chilblains, ii.	379	Liquor Piderit's phagedenic, ii.	69
				Liquor white phagedenic, ii.	92
				Liquor compound white phagedenic, ii.	92

Liquor yellow phagedenic, ii.	35	Lixivium ammoniacale vinosum, i.	172	Lohoch sedative, ii.	280
Liquor physagogue, ii.	355	Lixivium causticum, ii.	601	Lohoch terebinthine, ii.	655
Liquor Mœgling's probatory, i.	283	Lixivium causticum vegetabile, ii.	410	Lohoch terebinthinated, ii.	668
Liquor prophylactic, i.	225	Lixivium magistrale, ii.	410	Lohoch vermifuge, ii.	3
Liquor against pterygium, i.	595	Lixivium martis, i.	657	Lonchitidis herba, ii.	577
Liquor purgative and emetic, ii.	613	Lixivium mater salis marini, i.	362	Long-rooted gargle, i.	116
Liquor resolvent, ii. 18,	445, 613	Lixivium saponarium, ii.	410	Loniceræ periclymenum, ii.	203
Liquor saline, ii.	449	Lixivium sodæ, ii.	601	Loose-strife, ii.	205
Liquor scillitic, ii.	445	Lixivium tartari, ii.	422	Lopez root, ii.	479
Liquor Spangenberg's, ii.	76	Lizard, i.	697	Lopezia racemosa, ii.	479
Liquor Stettin's, ii.	78	Load-stone, i.	640	Loti odoratæ herba, ii.	225
Liquor stimulant and vulnerary, i.	762	Lobaria pulmonaria, ii.	188	Loti urbanæ herba, ii.	225
Liquor stomachic, ii. 240,	607	Lobelia inflata, ii.	203	Lotio acida, i.	70
Liquor styptic, i.	445	Lobelia syphilitica, ii.	203	Lotio alcoholisata, i.	349
Liquor Loof's styptic, i.	657	Logwood, ii.	193	Lotio antipsorica, ii.	49
Liquor tonic, i.	167	Lohoch album, i.	210	Lotio hydrargyri, ii.	34, 68
Liquor vegeto-mercurial, ii.	97	Lohoch anthelmintic, ii.	583	Lotio hydrargyri lutea, ii.	35
Liquor visceral i. 666, ii.	110	Lohoch antiasthmatic, i.	744, 567	Lotio nicotianæ, ii.	262
Liquor vulnerary, ii.	367	Lohoch anticatarrhal, i.	94	Lotio plumbata, ii.	379
Liquor camphorated vulnerary, ii.	388	Lohoch antisyphilitic, ii.	92	Lotio plumbi composita, ii.	78
Liquor Plenk's vulnerary, ii.	388	Lohoch ad asthma, ii.	567	Lotio quercus saturnina, ii.	389
Liquor spirituous vulnerary, ii.	546	Lohoch balsamic, i. 318,	322	Lotio saturnina, ii.	379
Liquorice, i.	722	Lohoch leniens balsamicum, ii.	655	Lotio e staphisagria, i.	609
Liquorice spiny, i.	723	Lohoch of red cabbage, i.	343	Lotio tartari stibiati, i.	260
Liriodendron tulipifera, i. 470, ii.	202	Lohoch of coltsfoot, ii.	699	Lotio ad tineam, ii.	613
Lithagrostis semen, i.	538	Lohoch compound, ii.	295	Lotio veratri mercurialis, ii.	49
Litharge, ii.	369	Lohoch demulcent, i.	767	Lotio e veratro, ii.	715
Lithargyrum, ii.	369	Lohoch expectorant, i.	248, 358, 753, ii.	Lotion anodyne, ii.	338
Lithomarge, i.	155	Lohoch de farfara, ii.	699	Lotion antipsoric, i.	66, ii. 49
Lithontriptic Tulp's, i.	380	Lohoch of fox's lung, i.	728	Lotion Bonhomme's antirachitic, ii.	422
Lithospermum officinale, ii.	203	Lohoch of gum arabic, i.	754	Lotion antiseptic, i. 490, ii.	387
Litmus, ii.	509	Lohoch gummoso-oleosum, i.	210	Lotion astringent, ii.	378
Litus acidus, i.	56	Lohoch of iceplant, ii.	242	Lotion against chilblains, ii.	423
Litus ad hæmorrhoides, ii.	107	Lohoch laxative, i.	463	Lotion corroborant, ii.	379
Liver of sulphur, ii.	416	Lohoch leniens, i.	767	Lotion desiccative, ii.	78
Liverwort, i. 103, ii.	4	Lohoch of manna, ii.	219	Lotion excitant, i.	609
Liverwort ash-coloured, ii.	186	Lohoch pectoral, i. 36, 600, 752, ii.	354, 569, 575	Lotus ad hæmorrhoides, ii.	107
Liverwort Iceland, ii.	187	Lohoch pectoralis leniens, i.	211, 752	Lotion of hellebore, ii.	715
Lixivia acetata, ii.	441	Lohoch pulmonale, i. 728, ii.	354	Lotion hydrocyanic, i.	59
Lixivia caustica, ii.	410	Lohoch de pulmone vulpis, i. 728, ii.	354	Lotion mercurial, ii.	68
Lixivia purificata, ii.	419	Lohoch sanum et expertum, i.	727	Lotion Alibert's mercurial, ii.	69
Lixivia tartarisata, ii.	447	Lohoch saponaceous, ii.	459	Lotion black mercurial, ii.	34
Lixivia vitriolata, ii.	427	Lohoch saponaceum, ii.	459	Lotion Manry's mercurial, ii.	86
Lixivium alkali vegetabilis, ii.	422	Lohoch saponatum, ii.	459	Lotion yellow mercurial, ii.	35
Lixivium ammoniacale, i. 168		Lohoch de scilla, ii.	567	Lotion ophthalmic, i.	192
Lixivium ammoniacale anisatum, i.	173			Lotion phagedenic, i.	763
Lixivium ammoniacale aromaticum, i.	174			Lotion resolvent ophthalmic, i.	264
				Lotion Schmucker's refrigerant, ii.	437

Lotion sedative, ii.	338	Lupine white, ii.	204	Magnesia carbonas, ii.	206
Lotion stimulant, i.	260	Lupinus albus, ii.	204	Magnesia subcarbonas,	
Lotion sulphuro-saponaceous, ii.	639	Lupuline, ii.	10	ii.	206
Lotion terebinthinated, ii.	665	Lybisticæ radix, ii.	197	Magnesia sulphas, ii.	210
Lotion against tinea, ii.	602	Lycoperdon bovista, ii.	204	Magnesium, ii.	206
Lotion for tinea, i.	349	Lycopodium, ii.	204	Magnesium oxydatum nativum, ii.	217
Lotus ad rheumatismum, ii.	670	Lycopodium clavatum, ii.	204	Magnet, i.	640
Lovage, ii.	197	Lynx-stone, i.	354	Magnet arsenical, i.	283
Love-apple, ii.	617	Lysimachia nummularia, ii.	205	Magnolia glauca, ii.	213
Lozenges Tronchin's anti-catarrhal, i.	247	Lysimachia vulgaris, ii.	205	Magnolia sweet-scented, ii.	213
Lozenges Kunkel's anti-monial, i.	242	Lysimachia lutea herba, ii.	205	Mahogany, ii.	641
Lozenges antiperiodic, i.	484	Lysimachia purpurea radix, ii.	205	Maidenhair Canadian, i.	89
Lozenges antistruumal, i.	393	Lythargyrus, ii.	369	Maidenhair great golden, ii.	409
Lozenges aphrodisiac, ii.	251	Lythrum salicaria, ii.	205	Mala aurantia Hispanensia, i.	506
Lozenges aromatic, ii.	163	Lytta, i.	377	Mala aurantia matura, i.	506
Lozenges of citric acid, i.	53	Mace, ii.	254	Mala citrea, i.	517
Lozenges Wistar's cough, i.	727	Macis, ii.	254	Mala insana, ii.	617
Lozenges of crab's eyes, i.	356	Madder, ii.	529	Mala lycopersica, ii.	617
Lozenges emetic, i.	283	Madrepore oculata, i.	353	Mala punica, ii.	470
Lozenges ferruginous, i.	643	Magisterium bismuthi, i.	339	Malabathrum, ii.	178
Lozenges of guaiac, i.	733	Magisterium jalapæ, i.	549	Malacca bean-tree, i.	215
Lozenges of iron, i.	638	Magisterium marcasitæ, i.	339	Malacca nut-tree, i.	215
Lozenges kermesine, i.	245	Magisterium saturni, ii.	369	Malacorum, ii.	470
Lozenges of liquorice, i.	725	Magisterium sulphuris, ii.	633	Malas ferri, i.	665
Lozenges of marshmallows, i.	146	Magisterium tartari purgans, ii.	441	Malas protoxidi ferri, i.	665
Lozenges nitrated, ii.	438	Magisterium vitriolis martis, i.	644	Mallow, ii.	215
Lozenges of opium, ii.	292	Magistery of jalap, i.	549	Malt, ii.	213
Lozenges of oxalic acid, i.	72	Magistery of sulphate of iron, i.	644	Maltum, ii.	213
Lozenges pectoral, i.	322	Magistery of sulphur, ii.	633	Malva alcea, ii.	216
Lozenges of hydriodate of potass, ii.	122	Magistrantia radix, ii.	114	Malva rotundifolia, ii.	215
Lozenges purgative, i.	550, 561	Magma scytodepsicum ex quercu, ii.	389	Malva sylvestris, ii.	215
Lozenges Spitzlay's, ii.	354	Magnes, i.	640	Malvavisci radix, i.	143
Lozenges of squill, ii.	567	Magnes arsenicalis, i.	283	Malvæ hortensis flores, i.	155
Lozenges of sweetflag, i.	86	Magnesia, ii.	206	Malvæ roseæ flores, i.	155
Lozenges against thirst, ii.	430	Magnesia ærata, ii.	206	Malvæ arboreæ flores, i.	155
Lozenges for allaying thirst, i.	82	Magnesia alba, ii.	206	Malus communis, ii.	214
Lozenges of violet, ii.	722	Magnesia alba calcinata, ii.	206	Mandibulæ lucii piscis, i.	360
Lozenges Ching's worm, ii.	51	Magnesia Edinburgensis, ii.	206	Mandragoræ radix, i.	312
Lujulæ herba, ii.	327	Magnesia nigra, ii.	217	Mandrake, i.	312
Lumbricus terrestris, i.	698	Magnesia opulina, i.	233	Manganese, ii.	217
Lunaria botrytidos herba, i.	342	Magnesia pura, ii.	206	Manganesium, ii.	217
Lunaria herba, i.	342	Magnesia subcarbonica, ii.	206	Manna, ii.	218
Lungwort, ii.	88, 469	Magnesia sulphurica, ii.	210	Manna di fronde, ii.	218
		Magnesia vitrariorum, ii.	217	Manna metallorum, ii.	46
		Magnesia vitriolica, ii.	210	Manganum oxydatum nativum, ii.	217
		Magnesia usta, ii.	206	Manna tabulata, ii.	451
				Manna tartarisata, ii.	451
				Maranta arundinacea, i.	632
				Marathri radix, i.	219
				Marble white, i.	352
				Marcasita, i.	339
				Marchantia polymorpha, ii.	221
				Margaritæ, i.	354
				Mari syriaci herba, ii.	682

Mari vulgaris herba, ii.	689	Meadow saffron, i.	538	Melilotas cæruleus, ii.	225
Marigold, i.	365	Meal prepared barley, ii.	7	Melilotus officinalis, ii.	226
Marigold diamond fig, ii.	241	Meals emollient, ii.	200	Melissa calamintha, ii.	227
Marjoram of candia, ii.	324	Meals resolvent, ii.	579, 696	Melissæ cedronellæ herba,	
Marjoram common, ii.	323	Mechoacan, i.	553	ii.	228
Marjoram cretan, ii.	324	Mechoacannæ radix, i.	553	Melissæ citrinæ herba,	
Marjoram sweet, ii.	322	Mechoacannæ albæ radix,		ii.	229
Marjoram wild, ii.	323	i.	543	Melissæ hortensis herba,	
Marjoranæ herba, ii.	322	Mechoacannæ nigræ		ii.	228
Marmalade antinephritic,		radix, i.	543	Melissæ humilis herba,	
ii.	657	Meconate of soda, ii.	615	ii.	232
Marmalade astringent,		Meconium, ii.	276	Melissa nepeta, ii.	227
ii.	673	Medlar Dutch, ii.	242	Melissa officinalis, ii.	227
Marmalade Dehaen's, i.	422	Medulla bovillæ, i.	317	Melissophylli herba, ii.	232
Marmalade expectorant,		Medulla saxorum, i.	155	Melithocorton, ii.	2
i.	48, 248	Mei radix, ii.	198	Mellago graminis, ii.	697
Marmalade febrifuge, i.	447	Mei athamantici radix,		Mellago leontodontis,	
ii.	669	ii.	198	ii.	652
Marmalade of iris, ii.	132	Mel acetati colchitici, i.	540	Meloë majalis, ii.	232
Marmalade mercurial, ii.	14	Mel aceti, i.	43	Meloë proscarabeus, ii.	232
Marmalade pectoral, ii.	125,	Mel aceti scillæ, ii.	574	Melon, i.	577
	636	Mel anthosatum, ii.	522	Melon water, i.	578
Marmalade of tamarinds,		Mel armoraciæ, ii.	482	Melonum semen, i.	577
i.	106	Mel boracis, ii.	609	Menstruum auri, i.	56
Marmalade Tronchin's,		Mel boraxatum, ii.	609	Mentha aquatica, ii.	233
i.	422	Mel chelidonii, i.	456	Mentha arvensis, ii.	233
Marmalade purgative, i.	605	Mel cupruratum, i.	591	Mentha crispa, ii.	233
ii.	492	Mel glycyrrhizatum, i.	728	Mentha gentilis, ii.	233
Marmor album, i.	352	Mel hydrargyrum, ii.	17	Mentha piperita, ii.	234
Marrow beef, i.	315	Mel hydrargyri, ii.	49	Mentha pulegium, ii.	234
Marrubium vulgare, ii.	221	Mel liquiritiæ, i.	728	Mentha rotundifolia, ii.	233
Marrubii albi herba, ii.	222	Mel mercuriale, ii.	241	Mentha sativa, ii.	233
Mars, i.	634	Mel mercurialis annuæ,		Mentha sylvestris, ii.	234
Mars solubilis, i.	667	ii.	241	Mentha viridis, ii.	234
Mars vitriolatus, i.	651	Mel fructuum mororum,		Menthæ balsaminæ herba,	
Marsh spurge, i.	629	ii.	248	ii.	233
Marshmallows, i.	142	Mel e muriate hydrargyri,		Menthæ catariciæ herba,	
Martis radix, ii.	721	ii.	49	ii.	259
Marum, ii.	682	Mel rosaceum, ii.	515	Menthæ equinæ herba, ii.	233
Mass odontalgic, ii.	223	Mel rosæ, ii.	515	Menthæ longifoliæ herba,	
Mass purgative, i.	403	Mel rosæ gallicæ, ii.	515	ii.	234
Massa ad fornacem, i.	334	Mel rosatum, ii.	515	Menthæ Romanæ radix,	
Massa pro fornace odori-		Mel rorismarini, ii.	522	i.	317
fera, i.	334	Mel rosatum solutivum,		Menthæ rubræ herba, ii.	233
Massa de gummo arabica,		ii.	516	Menthæ saracenicæ radix,	
i.	752	Mel fructuum sambuci,		i.	317
Masterwort, ii.	114	ii.	552	Menthæ sylvestris folio	
Masterwort great black,		Mel scillæ acetatum, ii.	574	rotundiore herba, ii.	233
i.	307	Mel scillæ compositum,		Menthæ sylvestris herba,	
Masticatory stimulant,		ii.	569	ii.	233
i.	231	Mel scilliticum, ii.	569	Menthæ verticillatæ herba,	
Mastic, ii.	223	Mel sub boratis sodæ, ii.	609	ii.	233
Mastiche, ii.	223	Mel superacetatis cupri,		Menthætri herba, ii.	234
Mastichina, ii.	218	i.	591	Mephatidis hepatica, i.	64
Mastichinæ gallorum		Mel vermium majalium,		Mercurialis annua, ii.	241
herba, ii.	680	ii.	232	Mercurius, ii.	12
Mater margaritarum, i.	353	Mel violarum, ii.	722	Mercurius aceticus, ii.	95
Mater perlarum, i.	353	Melaleuca cajaputi, ii.	270	Mercurius alkalizatus,	
Matonia cardamomum,		Melaleuca leucodendron,		ii.	18
i.	394	ii.	270	Mercurius boraxatus, ii.	81
Matricaria chamomilla,		Melanthii semen, ii.	264	Mercurius corallinus, ii.	36
i.	445	Melia azederach, ii.	225	Mercurius corrosivus	
Matricaria parthenium,		Melilot blue, ii.	225	ruber, ii.	36
ii.	225	Melilot common, ii.	226	Mercurius cosmeticus, ii.	89
May apple, ii.	432	Melilot trefoil, ii.	225	Mercurius dulcis, ii.	46

Mercurius dulcis martia- tus, ii.	58	Milk of sulphur, ii.	633	Mixtura antiherpetico-di- aphoretica, i.	84
Mercurius gummosus Plenkii, ii.	14	Milk-thistle, i.	395	Mixtura antimiasmatica, i.	589
Mercurius liquidus, ii.	92	Milk-vetch hairy-podded, i.	307	Mixtura antiphlogistica, ii.	437
Mercurius muriaticus, ii.	46	Milk virgin, ii.	388	Mixtura antiscabiosa, i.	639
Mercurius muriatus, ii.	66	Milk virgin's, ii.	375	Mixtura simplex anti- scorbutica, i.	535
Mercurius niger Mos- chati, ii.	52	Milkwort, ii.	402	Mixtura antispasmodica et diuretica, i.	102
Mercurius nitratus ruber, ii.	30	Millefolii herba, i.	35	Mixtura antispasmodica simplex, i.	92
Mercurius nitrosus, ii.	85	Millepede, ii.	272	Mixtura boraxata, ii.	609
Mercurius præcipitatus albus, ii.	46, 89	Millet, ii.	332	Mixtura e camphora cum aceto, i.	370
Mercurius præcipitatus fuscus Wurtzii, ii.	81	Millii semen, ii.	332	Mixtura camphoræ, i.	369
Mercurius præcipitatus viridis, ii.	38	Milligranæ herba, ii.	5	Mixtura camphoræ cum magnesia, i.	370
Mercurius ruber, ii.	36	Millietum de cupro ace- tatum, i.	591	Mixtura camphorata ætherea, i.	372
Mercurius saccharatus, ii.	16	Mimosa cinerea, ii.	150	Mixtura carbonatis potas- sæ absinthii stomachica, ii.	425
Mercurius solubilis Hah- nemanni, ii.	84	Mimosa corinda, ii.	150	Mixtura castorei ætherea, i.	431
Mercurius sublimatus cor- rosivus, ii.	66	Mimosa nilotica, i.	34, 750	Mixtura castorei ammo- niato spirituousa, i.	431
Mercurius sublimatus dulcis, ii.	46	Mimosa senegalensis, i.	766	Mixtura cathartica, i.	282
Mercurius cum sulphure, ii.	28	Mimosa unguis cati, ii.	195	Mixtura cerea, i.	433
Mercurius terrestris, ii.	406	Mineral turbith, ii.	82	Mixtura chelidonii, i.	456
Mercurius vitæ, i.	254	Mineral black turbith, ii.	84	Mixtura chinæ phospho- rata, ii.	349
Mercury, ii.	12	Minium, ii.	370	Mixtura cicutæ, i.	465
Mercury annual, ii.	241	Mint cat, ii.	259	Mixtura citre-muriatica, ii.	603
Mercury with chalk, ii.	18	Mint corn, ii.	233	Mixtura e conio, i.	465
Mercury with magnesia, ii.	18	Mint curled, ii.	233	Mixtura cornu cervi, i.	752
Mercury saccharated, ii.	16	Mint green, ii.	234	Mixtura cretacea, i.	358
Mercury with sulphur, ii.	28	Mint pennyroyal, ii.	234	Mixtura demulcens, ii.	541
Mercury with sulphur and sulphuret of anti- mony, ii.	30, 42	Mint pepper, ii.	234	Mixtura deobstruens, i.	449
Mesembryanthemum crys- tallinum, ii.	241	Mint round leaved, ii.	233	Mixtura depressans, i.	615
Mespilus arbutifolius, i.	470	Mint spear, ii.	233	Mixtura depressans et valde diuretica, i.	614
Mespilus Germanica, ii.	242	Mint water, ii.	233	Mixtura diatrion, i.	74
Mespilus oxyacantha, ii.	242	Mirospermum toluife- rum, i.	323	Mixtura digitalis campho- rata, i.	617
Metellæ nucis semines, i.	605	Misletoe, ii.	724	Mixtura ad dyslochiam, i.	102
Mezerei cortex, i.	603	Mistura ammoniaci, i.	742	Mixtura extracti rhois radicantis, ii.	504
Mezereon, i.	602	Mistura assafoetidæ, i.	303	Mixtura ferrea, i.	659
Milli solis semines, ii.	203	Mistura camphoræ, i.	369	Mixtura ferri aromatica, i.	635
Militis melissophyllum, ii.	232	Mistura cornu usti, i.	752	Mixtura ferri composita, i.	647
Milk anodyne, i.	209	Mistura cretæ, i.	358	Mixtura frigorifica, ii.	437
Milk artificial, i.	209	Mistura guaiaci, i.	739	Mixtura guaiaci gummosa, i.	739
Milk artificial asses, i.	701	Mistura millefolii, i.	36	Mixtura guaiacina, i.	734
Milk asses, ii.	149	Mistura mucilaginoso- balsamica, i.	22		
Milk astrigent, ii.	476	Mithridate, ii.	304		
Milk cow's, ii.	149	Miva cydoniorum, i.	598, 599		
Milk goat's, ii.	149	Miva cydoniorum aro- matica, i.	598		
Milk of lead, ii.	369	Mixtura acidi muriatici, i.	56		
Milk Plenk's mercurial, ii.	15	Mixtura acidi nitrici, i.	70		
		Mixtura acidi phosphorici, i.	73		
		Mixtura ætherea, i.	101		
		Mixtura alterans, i.	260		
		Mixtura amara cum rheo, ii.	493		
		Mixtura amygdalarum, i.	207, 208		
		Mixtura assafoetidæ pro clysmate, i.	303		
		Mixtura anodyna, i.	101		
		Mixtura antacida, ii.	210, 606		
		Mixtura anthelmintico-in- citans, ii.	683		
		Mixtura antidysentericus, i.	260		

Mixtura gummosa ex hydrargyro muriatico corrosivo, ii.	91	Mixtura ad partum, ii.	609	Mixture æthereous of castor, i.	431
Mixtura in hæmaturia chronica, ii.	724	Mixtura piperita, ii.	177	Mixture camphorated æthereous, i.	372
Mixtura Hulmiensis, i.	51	Mixtura pyrotartarica, i.	74	Mixture alkaline, ii.	424
Mixtura incitans mar- tialis, i.	666	Mixtura refrigerans, ii.	34	Mixture aloetic, i.	134
Mixtura incitans ner- vina, i.	281	Mixtura resinæ copaibæ, i.	556	Mixture alterative, i.	260
Mixtura incitans et se- dans, ii.	259	Mixtura resinæ guaiaci aquosa, i.	739	Mixture ammoniated spirituous of castor, i.	431
Mixtura e kali nitrico, ii.	434	Mixtura resolvens, ii.	123, 357	Mixture analeptic, ii.	326
Mixtura kali tartarici, ii.	450	Mixtura resolvens et ner- vina, i.	281	Mixture anodyne, ii.	279
Mixtura laxans, ii.	219, 647	Mixtura resolvens robo- rans, ii.	449	Mixture anodyne acid, ii.	288
Mixtura leniens, i.	211	Mixtura resolvens spiri- tuosa, ii.	416	Mixture antacid saline, ii.	212
Mixtura liquoris ammo- niæ causticæ, i.	171	Mixtura resolvent, i.	327	Mixture anthelmintic, i.	557, 64, 345
Mixtura e lupulo, ii.	11	Mixtura roborans, ii.	146	Mixture anti-aphthous, ii.	326
Mixtura lycopodii, ii.	205	Mixtura salina, ii.	446	Mixture anti-arthritis, i.	37, 84
Mixtura Macbridi, ii.	447	Mixtura salina compo- sita, ii.	447	Mixture anti-asthmatic, i. 117, ii.	552
Mixtura e magnesia sul- phurica, ii.	211	Mixtura salina paupe- rum, ii.	442	Mixture Van Swieten's anti-asthmatic, i.	187
Mixtura marrubii, ii.	222	Mixtura salina volatilis, i.	195	Mixture antiblennor- rhagic, i.	342
Mixtura martiata amara, i.	666	Mixtura saponacea, ii.	428	Mixture anticatarrhal, ii.	571
Mixtura mercurialis, ii.	90	Mixtura sedativo-narco- tica pectoralis, ii.	280	Mixture anticathartic, ii.	284
Mixtura mercurii gum- mosa, ii.	15	Mixtura sedativo-refrige- rans, ii.	434	Mixture antichlorotic, i.	79
Mixtura moschata, ii.	252	Mixtura simplex, i.	45, 74	Mixture anti-diarrhæic, ii.	256
Mixtura moschi, ii.	252	Mixtura simplex Ludo- vici, i.	74	Mixture antidysenteric, i. 70, ii.	206, 294
Mixtura mucilaginoso- aromatica, i.	568	Mixtura sistens, ii.	319	Mixture anti-emetic, i.	25
Mixtura myrrhæ alkali- sata, i.	763	Mixtura sulphurico acida, i.	79	Mixture anti-epileptic, i.	188
Mixtura myrrhæ Griffi- thii, i.	647	Mixtura sulphuris præ- cipitati, ii.	637	Mixture antifebrile, i.	56, 485
Mixtura nicotianæ diure- tica, ii.	263	Mixtura tonico-adstrin- gens, i.	494	Mixture antifebrile aci- dulous, i.	78
Mixtura nitrata, ii.	434	Mixtura tonico-antisepti- ca, i.	304	Mixture antigonorrhæic, ii.	267
Mixtura nitrata cam- phorica, ii.	437	Mixtura tonico-balsami- ca, i.	666	Mixture anti-hæmorrhagic, i.	701
Mixtura nitri phospho- rici, ii.	611	Mixtura tonico-balsami- ca diaphoretica, i.	493	Mixture antihydrotic, i.	388, 646
Mixtura nitrosa, ii.	434	Mixtura tonico-nervina Stahl, ii.	415	Mixture anti-icteric, ii.	443
Mixtura nucum juglan- dium, ii.	136	Mixtura tonico-stoma- chica, i.	528	Mixture antilethargic, ii.	281
Mixtura olei ricini cum jalapa, i.	553	Mixtura ad vomitum san- guinis, ii.	319	Mixture anti-odontalgic, ii.	299
Mixtura oleo-acida, i.	53	Mixtura vulneraria acida, i.	45	Mixture anti-otodynamic, ii.	299
Mixtura oleosa, i.	211	Mixture absorbent, i.	349, 364, ii.	Mixture antiperiodic, i.	485, 987
Mixtura oleosa cum aceto, i.	44	Mixture acid, i.	55, 57, 70, 73, 77, ii.	Mixture antiphthisical, ii.	616
Mixtura oleoso-balsamica, ii.	166	Mixture acidulated, i.	97	Mixture antipsoric, i.	639
Mixtura opiato-æthereæ, ii.	281	Mixture acidulated of camphor, i.	370		
Mixtura opiato-antidy- senterica, ii.	281	Mixture acidulous, i.	42		
Mixtura opii adstringens, ii.	319	Mixture acidulous pur- gative, ii.	453		
Mixtura oxygeno-muria- tica, i.	460				

- Mixture antirheumatical, i. 754
- Mixture antispasmodic, i. 70, 92, 102, 187, 201
- Mixture antispasmodic diuretic, i. 102
- Mixture antispasmodic resolvent, i. 102
- Mixture antisyphilitic, i. 592, ii. 63
- Mixture Dehaen's antisyphilitic, ii. 76
- Mixture aperitive, i. 263
- Mixture aromatic, ii. 177
- Mixture aromatic of iron, i. 635
- Mixture aromatic sedative, ii. 284
- Mixture arthritic, i. 56
- Mixture of assafoetida, i. 303
- Mixture astringent, i. 79, 486, ii. 194, 319, 512, 657
- Mixture balsamic, i. 22, 324, ii. 655
- Mixture bitter, ii. 493
- Mixture of camphor, i. 369
- Mixture of camphor and magnesia, i. 370
- Mixture spirituous camphorated, i. 373
- Mixture camphoric, i. 374
- Mixture cardiac, ii. 166
- Mixture carminative, ii. 356, 499
- Mixture cathartique Arabe, ii. 563
- Mixture comforting, i. 63
- Mixture composing, i. 371
- Mixture for convulsive cough, i. 485
- Mixture cordial, ii. 173
- Mixture cretaceous, i. 358
- Mixture demulcent, i. 209, 211, 600
- Mixture deobstruent, ii. 445
- Mixture depressing, i. 62
- Mixture detergent, ii. 518
- Mixture gummy of deutochloruret of mercury, ii. 91
- Mixture diaphoretic, i. 188, 566
- Mixture diuretic, i. 70, 540, 614, ii. 205, 263, 437, 438, 571, 655, 665
- Mixture for dysmenorrhoea, i. 102
- Mixture emetic, i. 621, ii. 127
- Mixture emeto-anodyne, i. 248
- Mixture excitant, ii. 603
- Mixture expectorant, i. 70, 615
- Mixture expectorant and tonic, ii. 128
- Mixture febrifuge, i. 485
- Mixture foetid, i. 305
- Mixture Griffiths' tonic, i. 647
- Mixture of gum ammoniac, i. 742
- Mixture compound of gum ammoniac, i. 744
- Mixture hydragogue, ii. 244
- Mixture kermeticized, i. 248
- Mixture laxative, ii. 453
- Mixture lithontriptic, i. 56, 101
- Mixture Ludwig's simple, i. 74
- Mixture mercurial, ii. 15
- Mixture balsamic mercurial, ii. 15
- Mixture mineral, ii. 71
- Mixture mucilaginous, i. 754
- Mixture Mutzel's resolvent, ii. 448
- Mixture of musk, ii. 252
- Mixture nervine, i. 304
- Mixture stimulant nervine, i. 281
- Mixture nitrous, ii. 434
- Mixture camphorated nitrous, ii. 437
- Mixture odontalgic, i. 400
- Mixture oleo-acid i. 53
- Mixture oleo-balsamic, ii. 166
- Mixture oleose and acetose, i. 44
- Mixture oleoso-acid, i. 521
- Mixture oleoso-saline, ii. 212
- Mixture opiate, ii. 279, 284
- Mixture oxymellated, i. 44
- Mixture pectoral, i. 60, 61, 192, 204, ii. 117, 341, 576
- Mixture with protochloruret of mercury, ii. 49
- Mixture purgative, i. 552
- Mixture saponaceous purgative, i. 549
- Mixture refreshing, i. 82
- Mixture refrigerant, i. 754, ii. 434, 435, 438
- Mixture resolvent, i. 193, 265, ii. 445, 446, 456, 616, 665
- Mixture resolvent and nervine, i. 281
- Mixture spirituous resolvent, ii. 416
- Mixture saline, ii. 443, 445
- Mixture compound saline, ii. 447
- Mixture simple saline, ii. 438, 416
- Mixture sedative, i. 48, 614, 617, ii. 247, 259, 281, 288, 295
- Mixture simple, i. 45, 74
- Mixture stimulant, i. 177
- Mixture stomachic, i. 711, 516, ii. 11, 425
- Mixture sudorific, i. 197
- Mixture Sylvius', ii. 276
- Mixture tartareous, ii. 450
- Mixture tartaric, i. 419
- Mixture theriacal, ii. 316
- Mixture tonic and antiseptic, i. 304
- Mixture tonic, i. 503, ii. 146, 447
- Mixture tonic and balsamic, i. 666
- Mixture Stahl's tonic-nervine, ii. 415
- Mixture vermifuge and purgative, i. 547
- Mixture vinous, ii. 430
- Mixture volatile, i. 202
- Molybdænum magnesiæ, ii. 217
- Momordica balsamina, ii. 242
- Momordica elaterium, ii. 243
- Momordica elaterii radix, ii. 243
- Monarda punctata, ii. 244
- Moneywort, ii. 205
- Monoceros, i. 359
- Monodon monoceros, i. 359
- Monkshood, i. 83
- Moon-wort, i. 342
- Mori rubri baccæ, ii. 531
- Moringa oleifera, ii. 244
- Morpheum, ii. 245
- Morphina, ii. 245
- Morphine, ii. 245, 274
- Morsuli aceti, i. 42
- Morsuli anthelmintici, ii. 583
- Morsuli antimoniales, i. 242
- Morsuli aromatici, ii. 163
- Moschus artificialis, ii. 631
- Morsuli chinati, i. 476
- Morsuli citri, i. 520
- Morsuli menthæ piperitæ, ii. 237
- Morsuli purgantes, i. 550
- Morsuli restaurantes Kunkelii, i. 242
- Morsuli ad scabiem, ii. 636
- Morsuli stibii, i. 242
- Morsuli contra vermes, ii. 583

Morsus diaboli radix, ii.	559	Mucilago lichenis Islan- dici, ii.	190	Murias hydrargyri oxy- genatus sublimatione paratus, ii.	66
Morsus gallinæ herba, i.	142	Mucilago liquiritiæ, i.	724	Murias hydrargyrisuboxy- genatus, ii.	46
Morus Indica, ii.	479	Mucilago mercurialis Plenkii, ii.	14	Murias hydrargyricum corrosivum, ii.	66
Morus nigra, ii.	248	Mucilago oleosa, i.	211	Murias hydrargyro-am- moniacalis salitus, ii.	90
Moschus, ii.	249	Mucilago psyllii, ii.	367	Murias hydrargyro-am- moniacale acerrimum, ii.	89
Moschus moschiferus, ii.	249	Mucilago radidis saleb, ii.	540	Murias hydrargyrosus mitis, ii.	46
Moss Corsican, ii.	2	Mucilago tragacanthæ, i.	766	Murias manganesi, ii.	217
Moss Iceland, ii.	187	Mugwort, i.	289	Murias oxydi hydrargyri ammoniacalis, ii.	89
Motacilla troglodytes, i.	335	Mulberry, ii.	248	Murias oxyduli hydrar- gyri mitis, ii.	46
Mother of pearl, i.	353	Mullein great, ii.	716	Murias oxyduli stibii, i.	254
Mother of thyme, ii.	690	Mungos radix, ii.	274	Murias oxygenatus, i.	254
Mountain ash, ii.	619	Muria, ii.	602	Murias platinae, ii.	368
Mountain blue, i.	582	Murias ammoniæ, i.	191	Murias potassæ, ii.	440
Mountain bugle, i.	103	Murias ammoniacæ æru- ginatus, i.	589	Murias potassæ oxyge- natum, ii.	440
Mountain green, i.	582	Murias ammoniæ et ferri, i.	661	Murias sodæ, ii.	602
Mountain-tea, i.	696	Murias ammoniacæ ferri- cus, i.	661	Murias stanni, ii.	625
Mouse-ear, ii.	6	Murias ammoniæ mar- tius, i.	661	Murias stibii, i.	254
Mucharum rosarum, ii.	512	Murias ammoniæ super- hydrargyricus, ii.	90	Murias stibiosus, i.	254
Mucilage camphorated, i.	368	Murias ammonio-ferri- cum, i.	661	Muriate of ammonia, i.	191
Mucilage of cantharides, i.	380	Murias antimonii, i.	254	Muriate of antimony, i.	254
Mucilage of great com- frey, ii.	642	Murias auri, i.	312	Muriate of barium, i.	325
Mucilage of fenugreek, ii.	696	Murias auri liquidus, i.	313	Muriate of iron, i.	656
Mucilage of fleaseed, ii.	367	Murias auri et sodii, i.	313	Muriate liquid of iron, i.	657
Mucilage of gum arabic, i.	751	Murias aurico-nitricum, i.	313	Muriate sublimed of iron, i.	656
Mucilage mercurial of gum arabic, ii.	14	Murias baryticum, i.	325	Muriate mild of mercury, ii.	46
Mucilage of linseed, ii.	200	Murias calcicum, i.	362	Muriate of soda, ii.	602
Mucilage of marshmal- lows root, i.	146	Murias calcis, i.	362	Muscæ Hispanicæ, i.	377
Mucilage of quince seeds, i.	600	Murias cupri, i.	558	Muscari ambrosiacum, ii.	254
Mucilage acidulated of quince seeds, i.	600	Murias ferri, i.	656	Musci canini herba, i.	186
Mucilage of starch, i.	214	Murias ferri ammonia- calis, i.	661	Musci cathartici herba, ii.	187
Mucilage of tragacanth, i.	766	Murias ferri liquidus, i.	657	Musci clavati herba, ii.	204
Mucilago acaciæ, i.	751	Murias ferricus, i.	656	Musci pyxidati herba, ii.	186
Mucilago althææ, i.	146	Murias ferro-ammonia- cale, i.	661	Musci terrestres herba, ii.	204
Mucilago radidis althææ, i.	146	Murias hydrargyri cor- rosivus, ii.	66	Muscus albus quernus, ii.	187
Mucilago arabica, i.	751	Murias hydrargyri basi oxydi imperfecti, ii.	66	Muscus arboreus, ii.	187
Mucilago astragali tra- gacanthæ, ii.	766	Murias hydrargyri ferra- tus, ii.	58	Muscus capillaceus ma- jor, ii.	409
Mucilago cantharidum, i.	380	Murias hydrargyri mitis, ii.	46	Muscus corallinus, ii.	2
Mucilago consolidæ ma- joris, ii.	642	Murias hydrargyri oxy- dulati, ii.	46	Muscus helminthocor- tos, ii.	2
Mucilago cretica, i.	358	Murias hydrargyri oxy- genatus præcipitatione paratus, ii.	66	Muscus inarinus, ii.	2
Mucilago seminum cydo- norum acidulata, i.	600	Murias hydrargyri spiri- tuosus liquidus, ii.	70	Muscus pulmonarius, ii.	188
Mucilago glycyrrhiæ, i.	724	Murias oxygenatus hy- drargyri sublimatus, ii.	66	Mushroom scarlet, i.	601
Mucilago gummi ara- bici, i.	751				
Mucilago gummi traga- canthæ, i.	766				

Musk, ii.	249	Nasturtii sylvestris herba,	614	Nitras natricum, ii.	614
Musk artificial, ii.	631	i.	560	Nitras potassæ, ii.	432
Musk yarrow, i.	34	Nasturtium, ii.	698	Nitras potassæ cum sul-	
Mustard bastard, ii.	402	Natron, ii.	604	phure fusus, ii.	433
Mustard black, ii.	588	Natrum, ii.	604	Nitras protoxidi potassii,	
Mustard white, ii.	588	Natrum aceticum, ii.	614	ii.	432
Mustard wild, i.	623	Natrum carbonicum, ii.	604	Nitras sodæ, ii.	614
Mustum hordei, ii.	214	Natrum glaciale, i.	362	Nitras sub-bismuthicum,	
Myrica gale, ii.	254	Natrum muriaticum, ii.	602	i.	339
Myricine, i.	433	Natrum oxyphosphoro-		Nitrate of ammonia, i.	190
Myristica aromatica,		des, ii.	610	Nitratis argenti crystalli,	
ii.	254	Natrum phosphoricum,		ii.	73
Myrobolanus bellerica,		ii.	610	Nitrate of mercury, ii.	85
ii.	257	Natrum sulphuratum,		Nitrate of silver, i.	273
Myrobolanus chebula,		ii.	602	Nitrate ammoniacal of	
ii.	257	Natrum sulphuricum, ii.	611	silver, i.	275
Myrobolanus citrina, ii.	257	Natrum tartarisatum,		Nitrate of soda, ii.	614
Myroxylum peruiferum,		ii.	455	Nitre, ii.	432
i.	320	Nauclea longiflora, ii.	145	Nitre cubic, ii.	614
Myrrh, i.	758	Navel-stone, i.	353	Nitre fixed, ii.	421
Myrrha, i.	758	Navel wort, i.	562	Nitre fixed by tartar,	
Myrrha liquida, i.	762	Nenupharis radix, ii.	268	ii.	421
Myrrha rubra, i.	758	Nepeta cataria, ii.	259	Nitre pearled, ii.	433
Myrrhidis annuæ semina,		Nepetæ herba, ii.	227	Nitre regenerated, ii.	432
i.	307	Nerium oleander, ii.	260	Nitre rhomboidal, ii.	614
Myrtacanthæ radix, ii.	536	Nettle common, ii.	701	Nitri naphtha, i.	94
Myrti brabanticæ herba,		Nettle stinging, ii.	701	Nitrogen, ii.	265
ii.	254	Nettle white dead, ii.	154	Nitrogen oxygenated, ii.	265
Myrtle, ii.	257	Nezetta, ii.	694	Nitrogenium, ii.	265
Myrtle Dutch, ii.	254	Nicotiana rustica, ii.	260	Nitrum, ii.	432
Myrtus communis, ii.	257	Nicotiana tabacum, ii.	260	Nitrum antimoniatum, i.	241
Myrtus pimenta, ii.	359	Nicotianæ minoris folia,		Nitrum antimoniatum	
Mytilus margarati ferus,		ii.	260	Stahlî, ii.	241
i.	353	Nicotine, ii.	260	Nitrum fixum, ii.	421
Myxæ fructus, i. 558, ii.	99	Nicotianine, ii.	260	Nitrum flammans, i.	190
		Nicotianum, ii.	260	Nitrum lunæ, i.	273
Nacra perlarum, i.	353	Nidi hirundinum, i.	355	Nitrum papaveratum,	
Nadæ celticæ alterius		Nigella arvensis, ii.	264	ii.	433
radix, i.	279	Nigella damascena, ii.	264	Nitrum perlatum, ii.	433
Naphæ flores, i.	506	Nigella sativa, ii.	264	Nitrum regeneratum, ii.	432
Naphtha, ii.	258	Nightshade American,		Nitrum rosatum, ii.	433
Naphtha aceti, i.	91	ii.	351	Nitrum saturninum, ii.	384
Naphtha nitri, i.	94	Nightshade deadly, i.	309	Nitrum tabulatum, ii.	433
Naphtha petrolei, ii.	345	Nightshade garden, ii.	617	Nitrum vitriolatum Schœ-	
Naphtha vitrioli, i.	97	Nightshade woody, ii.	615	deri, ii.	477
Naphtha vitrioli campho-		Ninsi radix, ii.	592	Nodulus purgans, i.	403
rata, i.	371	Ninsing radix, ii.	592	Nolæ culinaris radix,	
Naphtha vitrioli diluta,		Nitras ammoniæ, i.	190	i.	217
i.	100	Nitras ammoniacæ, i.	190	Nostochium commune,	
Naphtha vitrioli martia-		Nitras argenti, i.	273	ii.	265
lis, i.	659	Nitras argenti crystalli-		Nuces avellanæ, i.	561
Naphtha vitrioli phos-		nus, i.	273	Nuces cupressi, i.	580
phorata, ii.	349	Nitras argenti fusas,		Nuces quercus, ii.	474
Narcissus pseudo-narcis-		i.	273	Nuces quercus tostæ, ii.	477
sus, ii.	258	Nitras barii, i.	328	Nucis juglandis cortex,	
Narcissus wild, ii.	258	Nitras bismuthi, i.	339	ii.	135
Narcotina, ii.	259	Nitras cubicum, ii.	614	Nucis juglandis nucleus,	
Narcotine, ii.	259, 274	Nitras cupri, i.	588	ii.	135
Nard celtic, ii.	703	Nitras deutoxidi cupri,		Nuclei juglandis epider-	
Nardi rusticæ radix,		i.	588	mis, ii.	135
i.	293	Nitras hydrargyri, ii.	85	Nummulariæ herba, ii.	205
Nardi sylvestris radix,		Nitras hydrargyri liqui-		Nut-gall, i.	691
i.	293	dus, ii.	86	Nut Malabar, ii.	144
Nardus celtica, ii.	703	Nitras kalicum, ii.	432	Nutmeg, ii.	254
Narwhal, i.	359	Nitras lixivix, ii.	432	Nuts cyprus, i.	580

Nux Barbadosensis, i.	134	Oil of cocoa-nut, ii.	687	Oil olive, ii.	270
Nux ben, ii.	245	Oil of colocynth, i.	576	Oil of orange peel, i.	507
Nux cathartica Americana, ii.	134	Oil of copalm, ii.	627	Oil of palm, ii.	271
Nux juglandis, ii.	135	Oil of croton, i.	570	Oil paregoric, ii.	100
Nux moschata, ii.	254	Oil of cucumber, i.	578	Oil of petroleum, ii.	345
Nux myristicæ, ii.	254	Oil compound of cyclamen Europæum, i.	576	Oil phosphorated, ii.	350
Nux vomica, ii.	265	Oil of dill, ii.	221	Oil aromatic phosphorated, ii.	350
Nymphæa alba, ii.	267	Oil of dittany of crete, ii.	322	Oil of physic nut, ii.	134
Nymphæa lutea, ii.	268	Oil of earthworms, i.	705	Oil of pistachio, ii.	364
Oak, i.	474	Oil eel's, i.	317	Oil of poppy, ii.	337
Oak black, i.	474	Oil of egg, ii.	326	Oil of puppies, ii.	110
Oak Jerusalem, i.	457	Oil of elaterium, ii.	244	Oil purgative, ii.	562
Oak quercitron, ii.	474	Oil compound of elaterium, ii.	244	Oil of Rhodium wood, ii.	195
Oak poison, ii.	503	Oil of elder, ii.	554	Oil of ricinus, ii.	507
Oak sea, i.	678	Oil compound of elecampane, ii.	120	Oil of rose, ii.	517
Oak white, ii.	474	Oil of euphorbium, i.	756	Oil infused of rue, ii.	538
Ochra lutea, i.	643	Oil expressed of fennel, i.	219	Oil of sage, ii.	546
Ochra rubra, i.	643	Oil of figwort, ii.	578	Oil of saffron, i.	564
Ochra veneris, i.	582	Oil of flowers, ii.	109	Oil of yellow sanders, ii.	196
Ocre red, i.	643	Oil of fox, ii.	110	Oil of sassafras, ii.	184
Ocre yellow, i.	643	Oil of frog's spawn, i.	706	Oil of savine, ii.	143
Oculi cancrorum, i.	354	Oil of great mullein, ii.	716	Oil of scorpions, i.	705
Ocymum basilicum, ii.	269	Oil of gum ammoniac, i.	744	Oil compound of scorpions, i.	706
Ocymum minimum, ii.	269	Oil of guaiac, i.	732	Oil of sesame, ii.	586
Œsypus, ii.	187	Oil compound of guaiac, i.	732	Oil of seven flowers, i.	454
Offa alba, i.	113	Oil of hartshorn, i.	707	Oil of soap, ii.	464
Oil acoustic, i.	99, 100	Oil rectified of hartshorn, i.	707	Oil of St. John's-wort, ii.	108
Oil æthereal, i.	206	Oil succinated of hartshorn, i.	707	Oil compound of St. John's-wort, ii.	110
Oil of almonds, i.	118	Oil of hemlock, i.	466	Oil of St. John's-wort by infusion, ii.	108
Oil Batavian aloetic, i.	630	Oil of hempseed, i.	376	Oil stomachic, i.	31
Oil of amber, ii.	244	Oil of henbane, ii.	103	Oil of stramonium, i.	607
Oil of American mountain balm, ii.	317	Oil green of henbane, ii.	103	Oil of sulphur, i.	79
Oil animal, i.	707	Oil of henbane seeds, ii.	100	Oil of sulphur wort, ii.	344
Oil Dippel's animal, i.	354	Oil of hyssop, ii.	112	Oil of sweet basil, ii.	269
Oil of anise, ii.	669	Oil of iris, ii.	133	Oil of tacamahac, ii.	644
Oil anthelmintic, ii.	737	Oil of juniper, ii.	139	Oil deliquesced of tartar, ii.	422
Oil arthritic, i.	685	Oil of juniper wood, ii.	139	Oil of turpentine, ii.	663
Oil antirheumatic, ii.	676	Oil of lily, ii.	198	Oil of violet, ii.	723
Oil of ants, i.	229	Oil of linseed, ii.	202	Oil of wall flower, i.	455
Oil of balm, ii.	178	Oil of lizards, i.	706	Oil of walnuts, ii.	135
Oil of bay berries, ii.	245	Oil of marjoram, ii.	323, 324	Oil of wax, i.	434
Oil of ben, ii.	375	Oil of marjoram by infusion, ii.	323	Oil of wormwood, i.	24, 30
Oil Wedel's bezoardic, i.	270	Oil of marshmallows, i.	151	Oil essential of aloes, i.	118
Oil of brick, ii.	576	Oil of mastic, ii.	224	Oil essential of angelica, i.	222
Oil compound of bryony, ii.	269	Oil of mellilot, ii.	226	Oil essential of asphaltum, i.	298
Oil of cajeput, ii.	382	Oil of mint by infusion, ii.	236	Oil essential of benzoin, i.	332
Oil of cantharides, i.	389	Oil mucilaginous, ii.	201, 696	Oil nitric of camphor, i.	374
Oil caper, i.	629	Oil infused of mustard, ii.	589	Oil essential of caraway, i.	397
Oil expressed of caper spurge, i.	397	Oil of myrrh, i.	763	Oil essential of cardamoms, i.	395
Oil expressed of caraway, i.	448, 454	Oil of myrtle, ii.	258	Oil essential of cascarilla, i.	568
Oil æthereous carminative, i.	507	Oil neat's foot, i.	315		
Oil castor, i.	550, 570	Oil of garden nightshade, ii.	617		
Oil artificial castor, i.	453	Oil expressed of nutmeg, ii.	256		
Oil of chamomile, i.	448				
Oil compound of chamomile, i.	460				
Oil with chlorine, i.					

Oil essential of chamomile, i.	447	Ointment anthelmintic, i.	31	Ointment cosmetic, i.	441
Oil essential of chervil, i.	445	Ointment anti-hæmorrhoidal, i.	39, 350, 352, 566, 692, ii. 45, 49, 83, 93, 294, 374, 381, 382, 385, 386, 391, 624, 640	Ointment of cucumbers, i.	578
Oil essential of cinnamon, ii.	165	Ointment antiherpetic, i.	248	Ointment of cyanuret of mercury, ii.	81
Oil essential of coriander, i.	559	Ointment Cullerier's antiherpetic, ii.	83	Ointment of cyclamen Europeanum, i.	576
Oil essential of cubebs, ii.	362	Ointment anti-ophthalmic, ii.	279	Ointment Cyrillo's, ii.	72
Oil essential of cumin, i.	579	Ointment antiparalytic, ii.	463	Ointment of the resin of daphne gnidium, i.	604
Oil essential of dill, i.	221	Ointment antipsoric, i.	391	Ointment defensive, i.	156
Oil essential of fennel, i.	219	ii. 25, 30, 34, 38, 72, 93, 264, 424, 426, 602		Ointment demulcent and balsamic, ii.	658
Oil essential of galbanum, i.	683	Ointment antiscorbutic, ii.	628	Ointment deobstruent, i.	469
Oil compound essential of galbanum, i.	684	Ointment antiscrofulous, ii.	123	Ointment depilatory, ii.	657
Oil essential of ginger, i.	205	Ointment antispasmodic, i.	426	Ointment Desault's, ii.	41
Oil essential of lavender, ii.	157	Ointment astringent, ii.	375	Ointment desiccative, ii.	78, 374
Oil essential of lemon, i.	517	Ointment antistrumous, ii.	39, 326	Ointment Stahl's detergent, ii.	25
Oil essential of mace, ii.	256	Ointment antisymphilitic, i. 312, ii.	63	Ointment detergent, ii.	30, 263
Oil essential of milfoil, i.	36	Ointment of apples, i.	436	Ointment of deutoxide of lead, ii.	370, 398
Oil essential of mint, ii.	236	Ointment aromatic, ii.	527	Ointment camphorated of deutoxide of lead, ii.	370
Oil essential of mountain tea, i.	696	Ointment camphorated aromatic, ii.	181	Ointment digestive, ii.	628
Oil essential of mugwort, i.	289	Ointment arsenical, i.	285	Ointment opiate digestive, ii.	660
Oil essential of orange, i.	507	Ointment astringent, i.	162	Ointment simple digestive, ii.	659
Oil essential of parsley, i.	269	Ointment basilicon, ii.	677	Ointment of digitalis, i.	618
Oil essential of pimenta, ii.	362	Ointment of bay, ii.	179	Ointment discutient, ii.	588
Oil essential of rosemary, ii.	524	Ointment of belladonna, i.	311	Ointment diuretic, ii.	568
Oil essential of rue, ii.	538	Ointment Bell's, ii.	41	Ointment Dubois', ii.	88
Oil essential of savory, ii.	558	Ointment black, ii.	45	Ointment of elder, ii.	554
Oil essential of sweet flag, i.	87	Ointment of black pepper, ii.	363	Ointment of elecampane, ii.	115
Oil essential of tansey, ii.	651	Ointment against burns, ii.	104, 374, 381	Ointment of elemi, ii.	486
Oil essential of thyme, ii.	689	Ointment of cantharides, i.	383, 384	Ointment compound of elemi, ii.	487, 658
Oil essential of valerian, ii.	711	Ointment compound of cantharides, i.	382	Ointment emollient, i.	151
Oil essential of wild thyme, ii.	690	Ointment opiate of cantharides, i.	378	ii.	657
Ointment alabastrine, i.	154	Ointment of carburet of iron, i.	639	Ointment epispastic, i.	382, 384, 387, 757
Ointment alkaline, ii.	606	Ointment cardiac, ii.	617	Ointment epispastic ammoniacal, i.	177
Ointment of American nightshade, ii.	351	Ointment carminative, ii.	181, 454	Ointment camphorated epispastic, i.	378, 387
Ointment ammoniacal, i.	190	Ointment catheretic, i.	586	Ointment green epispastic, i.	387
Ointment saponaceous ammoniacal, i.	177	Ointment of charcoal, i.	391	Ointment escharotic, ii.	39
Ointment anodyne, ii.	104, 294, 296, 299	Ointment against chilblains, ii.	674	Ointment Fabre's, i.	262
Ointment antiarthritic, ii.	21	Ointment for chilblains, i.	375	Ointment Falck's, ii.	42
		Ointment of chlorine, i.	460	Ointment hæmorrhoidal, ii.	100
		Ointment citric, i.	53	Ointment of white hellebore, ii.	715
		Ointment clysmatic, i.	154	Ointment Hellmund's, i.	284
				Ointment of hemlock, i.	466
				Ointment of henbane, ii.	103
				Ointment of hops, ii.	11
				Ointment of hydriodate of potass, ii.	122
				Ointment hypnotic, ii.	303

- Ointment golden, ii. 38
 Ointment green, i. 594, ii. 110
 Ointment of green precipitate ii. 98
 Ointment Jadelot's, ii. 93
 Ointment Janin's, ii. 93
 Ointment Jenner's, i. 262
 Ointment of infusion of cantharides, i. 377
 Ointment of iodine, ii. 121
 Ointment of ioduretted hydriodate of potass, ii. 123
 Ointment of ioduret of mercury, ii. 80
 Ointment Joerden's, ii. 41
 Ointment Juch's, ii. 22
 Ointment of juniper, ii. 141
 Ointment against lice, i. 220, 231, 269, 609, ii. 22
 Ointment compound of liquorice, ii. 374
 Ointment of turpentine, ii. 12
 Ointment Lyon's, ii. 38
 Ointment of mallow, ii. 216
 Ointment of marigold, i. 365
 Ointment mercurial, ii. 19, 40
 Ointment Bœcking's mercurial, ii. 39
 Ointment camphorated mercurial, ii. 25
 Ointment camphorated grey mercurial, ii. 35
 Ointment digestive mercurial, ii. 21
 Ointment grey mercurial, ii. 34
 Ointment nitrous mercurial, ii. 21
 Ointment opiate mercurial, ii. 26
 Ointment Sanchez' mercurial, ii. 21
 Ointment saponaceous mercurial, ii. 25
 Ointment sulphuretted mercurial, ii. 30
 Ointment of mezereon, i. 604
 Ointment of myrrh, i. 761
 Ointment of muriate of soda, ii. 603
 Ointment narcotic, ii. 296
 Ointment nervine, i. 182, ii. 179
 Ointment of garden nightshade, ii. 618
 Ointment of nitrate of mercury, ii. 87
 Ointment of nitric oxide of mercury, ii. 38
 Ointment nitrous, i. 70
 Ointment of nutgalls, i. 692
 Ointment opiate of nutgalls, i. 692
 Ointment ophthalmic, i. 594
 633, ii. 41
 Ointment Dupuytren's ophthalmic, ii. 40
 Ointment Pilletan's ophthalmic, ii. 38
 Ointment Richter's ophthalmic, ii. 39
 Ointment opiate, ii. 278, 296
 Ointment pectoral, ii. 133
 Ointment phosphorated, ii. 350
 Ointment of phosphoric acid, i. 73
 Ointment of poplar, ii. 106, 410
 Ointment of propolis, ii. 467
 Ointment of protochloruret of mercury, ii. 49
 Ointment of protoxide of cobalt, i. 526
 Ointment red, i. 216, ii. 41
 Ointment refrigerant, i. 375
 Ointment Regent's, ii. 41
 Ointment renal, ii. 301
 Ointment of resin, ii. 675
 Ointment brown resinous, ii. 679
 Ointment resolvent, i. 378
 ii. 65, 385, 463
 Ointment Ricour's, ii. 401
 Ointment of rose, ii. 518
 Ointment compound of rosemary, ii. 527
 Ointment of rue, ii. 538
 Ointment Rust's, ii. 41
 Ointment santaline, ii. 196
 Ointment saponaceous, ii. 459, 462
 Ointment saturnine, ii. 385
 Ointment compound saturnine, ii. 374
 Ointment saturno-mercurial, ii. 88
 Ointment for scrofulous ulcers, ii. 40
 Ointment sedative, i. 375
 ii. 283, 386
 Ointment soothing, i. 317
 ii. 296
 Ointment spermaceti, i. 441
 Ointment splenic, i. 749
 Ointment of squill, ii. 567
 Ointment stimulant, ii. 40
 Ointment stomachic, i. 31, 399, 454
 Ointment of storax, ii. 627
 Ointment of stramonium, i. 607
 Ointment of subacetate of copper, i. 593
 Ointment compound of subacetate of copper, i. 593
 Ointment of subacetate of lead, ii. 380
 Ointment camphorated of subacetate of lead, ii. 381
 Ointment opiate of subacetate of lead, ii. 381
 Ointment saponaceous of subacetate of lead, ii. 382
 Ointment of subcarbonate of lead, ii. 373, 401
 Ointment camphorated of subcarbonate of lead, ii. 373
 Ointment of sub-deuto-sulphate of mercury, ii. 83
 Ointment of sulphate of copper, i. 584
 Ointment compound of sulphur, ii. 639
 Ointment simple of sulphur, ii. 638
 Ointment of red sulphuret of mercury, ii. 45
 Ointment sulphuric, i. 78
 Ointment sulphuro-alkaline, ii. 410
 Ointment sulphuro-saponaceous, ii. 639
 Ointment sultana's, i. 442
 Ointment suppurative, i. 113
 ii. 40
 Ointment for syphilitic ulcers, ii. 40
 Ointment of tar, ii. 365
 Ointment of tartarized antimony, i. 261
 Ointment against tic d'oreux, ii. 373
 Ointment against tinea, i. 391, 679, ii. 89, 93, 640, 675
 Ointment of toadflax, ii. 199
 Ointment of tritoxide of lead, ii. 371
 Ointment of turpentine, ii. 657
 Ointment camphorated of turpentine, ii. 659, 661
 Ointment of the twelve apostles, ii. 395
 Ointment of veratrine, ii. 713
 Ointment vermifuge, i. 122
 Ointment of vinegar, i. 45
 Ointment of willow, ii. 542
 Ointment Zeller's, ii. 92
 Olea Europœa, ii. 270
 Oleo-margarate of mercury, ii. 98
 Oleo-resina copabii, i. 554
 Oleo-resina de Mecca, i. 319
 Oleosaccharum anisi, ii. 353
 Oleosaccharum aurantii, i. 507
 Oleosaccharum calami aromatici, i. 87
 Oleosaccharum cephalicum, ii. 252

Oleosaccharum cinna- momi, ii.	165	Oleum amygdalarum dul- cium, i.	206	Oleum cataputræ majoris, i.	629
Oleosaccharum citri, i.	517	Oleum anethi æthereum, i.	221	Oleum de cedro, i.	517
Oleosaccharum myrrhæ, i.	759	Oleum animale æthe- reum, i.	707	Oleum ceræ, i.	434
Oleosaccharum succini, ii.	631	Oleum æthereum ani- male empyreumaticum, i.	707	Oleum chamæmeli æthe- reum, i.	447
Oleosaccharum of amber, ii.	631	Oleum animale Dippelii, i.	707	Oleum cæreuleum chamæ- meli compositum, i.	448
Oleosaccharum of anise, ii.	353	Oleum animale fœtidum, i.	707	Oleum cicutæ, i.	466
Oleosaccharum of caje- put, ii.	270	Oleum volatile animale, i.	707	Oleum citri corticum, i.	517
Oleosaccharum cephalic, ii.	252	Oleum anisi, ii.	354	Oleum cinnamomi æthe- reum, ii.	165
Oleosaccharum of chamo- mile, i.	448	Oleum anisi æthereum, ii.	354	Oleum chamæmelinum, i.	453
Oleosaccharum of cinna- mon, ii.	165	Oleum anthemidis, i.	447	Oleum colocynthidis, i.	576
Oleosaccharum of cloves, i.	400	Oleum volatile anthemi- dis nobilis, i.	448	Oleum conyzæ compo- situm, ii.	244
Oleosaccharum of croton tigilium, i.	570	Oleum anthos æthereum, ii.	524	Oleum coriandri æthe- reum, i.	559
Oleosaccharum of fennel, i.	219	Oleum apparitii, ii.	110	Oleum cornu cervi fœti- dum, i.	707
Oleosaccharum of juni- per, ii.	139	Oleum arthanitæ com- positum, i.	576	Oleum cornu cervi recti- ficatum, i.	707
Oleosaccharum of lemon, i.	517	Oleum asphalti, i.	298	Oleum expressum corti- cis flavi citri medicæ, i.	517
Oleosaccharum of mace, ii.	256	Oleum aurantii æthereum, i.	507	Oleum expressum corti- cis flavi fructus citri aurantii, i.	507
Oleosaccharum of myrrh, i.	759	Oleum baccarum juniperi æthereum, ii.	139	Oleum corylinum, i.	561
Oleosaccharum of orange, i.	507	Oleum baccarum lauri, ii.	178	Oleum cubebarum æthe- reum, ii.	362
Oleosaccharum of pepper- mint, ii.	237	Oleum baccarum pi- mentæ, ii.	362	Oleum seminis cucu- meris, i.	578
Oleosaccharum of sassa- fras, ii.	184	Oleum balatinum, ii.	245	Oleum de cucurbita, i.	579
Oleosaccharum of sweet- flag, i.	87	Oleum been, ii.	245	Oleum cucurbitæ pleuri- ticum, i.	579
Oleosaccharum of tur- pentine, ii.	668	Oleum bryoniæ compo- situm, ii.	576	Oleum cumini æthe- reum, i.	579
Oleosaccharum of vale- rian, ii.	711	Oleum bufonum, ii.	104	Oleum dictamni æthe- reum, ii.	322
Oleum, i.	445	Oleum cacao, ii.	687	Oleum Dippelii, i.	707
Oleum absinthii æthe- reum, i.	24	Oleum fructus cacao, ii.	687	Oleum empyreumaticum animale, i.	707
Oleum absinthii distilla- tum, i.	24	Oleum cacao spissatum, ii.	687	Oleum empyreumaticum cornu cervi rectifica- tum, i.	707
Oleum absinthii coctum, i.	30	Oleum cardamomi se- minis æthereum, i.	395	Oleum enulæ compo- situm, ii.	120
Oleum absinthii infu- sum, i.	30	Oleum carminativum, i.	454	Oleum formicarum, i.	676
Oleum æthereo-pingvæ nucleorum myristicæ moschatæ, ii.	256	Oleum carpathicum, i.	318	Oleum distillatum Gauth- eriæ, i.	696
Oleum æthereum, i.	100	Oleum carui æthereum, i.	397	Oleum guaiaci compo- situm, i.	732
Oleum ammoniatum, i.	177	Oleum carui essentielle, i.	397	Oleum hieraclinum, i.	561
Oleum amygdalæ com- munis, i.	206	Oleum caryophyllorum essentiale, i.	400	Oleum hyoscyami ammo- niato-camphoratum, i.	179
Oleum amygdalarum, i.	206	Oleum castorei, i.	426	Oleum foliorum hyoscy- ami coctum, ii.	103
		Oleum catellorum, ii.	110	Oleum herbæ hyoscy- ami coctum, ii.	103

Oleum hyoscyami infusum, ii.	103	Oleum myrrhæ, i.	763	Oleum sambuci, ii.	534
Oleum hyoscyami viride, ii.	103	Oleum myrrhæ per deliquium, i.	762	Oleum santali citrini æthereum, ii.	196
Oleum hyperici per infusum, ii.	108	Oleum myrtinum, ii.	258	Oleum saponis, ii.	464
Oleum hyssopi æthereum, ii.	112	Oleum naphæ, i.	507	Oleum sassafras ligni æthereum, ii.	184
Oleum ireos, ii.	133	Oleum narcoticorum, ii.	104	Oleum saturejæ æthereum, ii.	558
Oleum iridis, ii.	133	Oleum nucis moschatæ, ii.	256	Oleum saturni, ii.	385
Oleum irinum, ii.	133	Oleum nucistæ, ii.	256	Oleum scarabeorum, ii.	232
Oleum juglandis regiæ, ii.	135	Oleum nucistæ expressum, ii.	256	Oleum serpylli æthereum, ii.	690
Oleum lacertarum viridium, i.	706	Oleum nucum juglandium, ii.	135	Oleum scorpionum, i.	705
Oleum æthereum laticis, ii.	664	Oleum olivarum, ii.	270	Oleum scorpionum compositum, i.	706
Oleum laterum, ii.	270	Oleum olivarum empyreumaticum, ii.	270	Oleum septem florum, i.	454
Oleum lavandulæ æthereum, ii.	157	Oleum olivarum oxynatum, i.	460	Oleum seminum sinapi infusum, ii.	589
Oleum lauri expressum, ii.	178	Oleum origani æthereum, ii.	324	Oleum florum Slotani, ii.	109
Oleum laurinum, ii.	179	Oleum ovorum expressum, ii.	326	Oleum solani nigri, ii.	617
Oleum laurinum infusum, ii.	179	Oleum oxydi cupri viride, i.	594	Oleum spermatis ranæ, i.	706
Oleum laurinum, ii.	178	Oleum palmæ, ii.	271	Oleum stramonii, i.	607
Oleum ligni juniperini æthereum, ii.	139	Oleum papaveris albi, ii.	337	Oleum stramonii compositum, ii.	104
Oleum liliorum alborum, ii.	198	Oleum volatile petrolei, ii.	345	Oleum succini, ii.	630
Oleum lini cum calce, i.	349	Oleum petræ, ii.	345	Oleum succini æthereum, ii.	630
Oleum lini expressum, ii.	202	Oleum petroselinæ æthereum, i.	269	Oleum succini oxydatum, ii.	631
Oleum lumbricorum, i.	705	Oleum philosophorum, ii.	270	Oleum succini purissimum, ii.	631
Oleum macis æthereum, ii.	256	Oleum phosphoratum, ii.	350	Oleum succini rectificatum, ii.	631
Oleum mariæ, i.	319	Oleum pimentæ, ii.	362	Oleum sulphuratum, ii.	637
Oleum marjoranæ infusum, ii.	323	Oleum pimentæ æthereum, ii.	362	Oleum tacamahacæ, ii.	644
Oleum martis per deliquium, i.	657	Oleum pinguidinis, i.	317	Oleum tanacetæ æthereum, ii.	651
Oleum mastiches, ii.	224	Oleum volatile pini, ii.	664	Oleum tartari per deliquium, ii.	422
Oleum mastichinum, ii.	224	Oleum pistaciorum, ii.	364	Oleum terebinthinæ, ii.	663
Oleum melissæ æthereum, ii.	229	Oleum pulegii, ii.	237	Oleum terebinthinæ sulphuratum, ii.	638
Oleum menthæ crispæ, ii.	237	Oleum purgans, ii.	562	Oleum thapsiæ anti-rheumaticum, ii.	685
Oleum menthæ piperitæ, ii.	237	Oleum pyro-animale depuratum, i.	707	Oleum theobromæ cacao expressum, ii.	687
Oleum menthæ sativæ, ii.	237	Oleum ricini, ii.	507	Oleum thymi æthereum, ii.	689
Oleum menthæ viridis, ii.	237	Oleum ricini Americani, ii.	134	Oleum ad tormina ventris, i.	454
Oleum millefolii æthereum, i.	36	Oleum rorismarini, ii.	524	Oleum de tribus, ii.	669
Oleum momordicæ, ii.	244	Oleum volatile rorismarini officinalis, ii.	524	Oleum valerianæ æthereum, ii.	711
Oleum monardæ æthereum, ii.	244	Oleum rosarum æthereum, ii.	517	Oleum verbasci, ii.	716
Oleum mucilagineum, ii.	201	Oleum rosatum, ii.	517	Oleum vini, i.	100
Oleum de mucilagibus, ii.	201	Oleum rutæ æthereum, ii.	538	Oleum violarum, ii.	723
		Oleum rutæ infusum, ii.	538	Oleum e vitellis ovorum, ii.	326
		Oleum sabinæ æthereum, ii.	143	Oleum vitrioli, i.	75
		Oleum salviæ æthereum, ii.	546	Oleum vitrioli depuratum, i.	76

Oleum vitrioli dulce, i.	99	Orpine, ii.	580	Oxodes hydrargyri ru- brum, ii.	36
Oleum vitrioli purifica- tum, i.	76	Orris, ii.	131	Oxodes mangani nativa, ii.	217
Oleum vulpinum, ii.	110	Orritanum, ii.	316	Oxodes stibii album, i.	236
Oleum zingiberis æthe- reum, i.	205	Oryza sativa, ii.	325	Oxodes stibii sulphura- tum, i.	249
Olibanum, ii.	271	Os sepia, i.	354	Ox tongue, i.	215
Olive gum, i.	764	Osmazome, i.	631	Oxyacanthæ flores, ii.	242
Olivile, i.	764	Ossa, i.	359	Oxycocci baccæ, ii.	702
Olsnitii radix, ii.	581	Ossa de cordibus cervi, i.	360	Oxycrat, i.	42
Oneberry, ii.	342	Ossa usta præparata, i.	360	Oxycratum, i.	42
Ongula alces, i.	678	Osteocolla, i.	352	Oxyduli stibii sulphura- tum, i.	249
Onion, i.	112	Ostrea edulis, i.	353	Oxydulum antimonii hy- dro-sulphuratum au- rantiacum, i.	249
Onion sea, ii.	563	Ostruthii radix, ii.	114	Oxydulum ferri nigrum, i.	640
Oniscus asellus, ii.	272	Ovum gallinaceum, ii.	326	Oxydulum hydrargyri Hahnemanni, ii.	84
Onopordon acanthium, ii.	273	Oxalas acidulum potassæ, ii.	446	Oxydulum stanni, ii.	623
Onosma echioides, ii.	273	Oxalas ammoniæ, i.	200	Oxydulum stibii per aci- dum muriaticum, i.	234
Opercula cochlearum, i.	354	Oxalas hydrargyri, ii.	95	Oxydulum stibii griseum, i.	234
Opercula of snail-shells, i.	354	Oxalas potassæ acidulus, ii.	446	Oxydulum stibii sulphu- ratum semi-vitreum, i.	241
Ophioglossum vulgatum, ii.	273	Oxalas subpotassicus, ii.	446	Oxydulum stibii vitria- tum, i.	235
Ophiorrhiza mungos, ii.	274	Oxalate of ammonia, i.	200	Oxydulum stibii vitreum, i.	235
Opiate anti-odontalgic, i.	401	Oxalate of mercury, ii.	95	Oxydulum stibiosum cum sulphure, i.	249
Opium, ii.	274	Oxalidis radix, ii.	534	Oxydum antimonii al- bum, i.	254
Opium colatum, ii.	289	Oxalis acetosella, ii.	327	Oxydum antimonii nitro- muriaticum, i.	254
Opium depuratum, ii.	289	Oxalis corniculata, ii.	327	Oxydum antimonii cum phosphate calcis, i.	236
Opium thebaicum, ii.	274	Oxalium, ii.	446	Oxydum antimonii sul- phuratum rubrum, i.	243
Opobalsam, i.	214	Ox-eye, i.	446	Oxydum antimonii cum sulphure vitrificatum, i.	235
Opobalsamum verum, i.	319	Ox-eye greater daisy, i.	461	Oxydum antimonii sul- phuretum griseum,	235
Opodeldoc balsam, i.	179	Ox-gall, i.	632	Oxydum arsenici album, ii.	283
Opoponax, ii.	321	Ox-gall inspissated, i.	632	Oxydum arsenici album cum sulphureto stibii, i.	284
Opuntia folia, i.	346	Oxi-chloruretum calcis, i.	351	Oxydum auri ammonia- tum, i.	314
Orange-tree, i.	505	Oxide grey sulphuretted of antimony, i.	235	Oxydum calcii, i.	347
Orcella, ii.	509	Oxide white of arsenic, i.	283	Oxydum cupri per aci- dum vegetabile, i.	590
Orchilla, ii.	509	Oxide black of iron, i.	640	Oxydum ferri ferricum, i.	643
Oreoselinii radix, ii.	581	Oxide red of iron, i.	643	Oxydum ferri nigrum, i.	640
Origani aquatici radix, i.	628	Oxide of kadmium, ii.	145	Oxydum ferri nigrum purificatum, i.	640
Origani vulgaris herba, ii.	324	Oxide red of lead, ii.	370		
Origanum, ii.	323	Oxide semivitreous of lead, ii.	369		
Origanum creticum, ii.	324	Oxide yellow of lead, ii.	369		
Origanum marjorana, ii.	322	Oxide of magnesium, ii.	206		
Origanum dictamnus, ii.	322	Oxide black of man- ganese, ii.	217		
Origanum vulgare, ii.	323	Oxide grey of mercury, ii.	32		
Orleana, ii.	352	Oxide Moretti's grey of mercury, ii.	34		
Ormin, ii.	544	Oxide nitric of mercury, ii.	36		
Ornithoglossæ semen, i.	678	Oxide red of mercury, ii.	36		
Orobanchæ Virginiana, ii.	324	Oxide of platina, ii.	368		
Orobi semen, ii.	719	Oxodes ferri, i.	640		
Orontii majoris herba, i.	266	Oxodes hydrargyri ni- grum, ii.	84		
Orpiment, i.	282				
Orpiment red, i.	282				
Orpiment yellow, i.	282				

Oxydum ferri rubrum, i.	643	Oxydum stibii acido muriatico oxygenato paratum, i.	254	Oxytartarus, ii.	441
Oxydum ferricum, i.	640	Oxydum stibii hydrosulphuratum rubro-fusum, i.	243	Oxysaccharum emeticum, i.	261
Oxydum ferricum crystallisatum nativum, i.	643	Oxydum stibii sulphuratum rubrum, i.	243	Oxysaccharum scilliticum, ii.	574
Oxydum ferroso-ferricum, i.	640	Oxydum stibii potassicum, i.	240	Oxytriphylly herba, ii.	327
Oxydum ferrosus, i.	640	Oxydum stibii cum potassa, i.	240	Padi cortex, i.	437
Oxydum hydrargyri ammoniacale, ii.	84	Oxydum stibii sulphuratum vitreum, i.	235	Pæonia officinalis, ii.	328
Oxydum hydrargyri cinereum, ii.	32	Oxydum stibiosum, i.	236	Palampissa, ii.	673
Oxydum hydrargyri completum, ii.	36	Oxygen, ii.	328	Palma christi, ii.	507
Oxydum hydrargyri nigrum, ii.	32, 84	Oxygenium, ii.	328	Palmulæ, ii.	347
Oxydum hydrargyri nitratum, ii.	36	Oxygeno-chloruretum potassii, ii.	440	Paludii radix, i.	266
Oxydum hydrargyri nitratum rubrum, ii.	36	Oxymel æruginis, i.	591	Panacea duplicata, ii.	427
Oxydum hydrargyri nitrico-ammoniacale, ii.	84	Oxymel chalybeatum, i.	664	Panacea hydrargyri rubra, ii.	36
Oxydum hydrargyri nitricum, ii.	36	Oxymel colchici, i.	540	Panacea Kornmanni, i.	314
Oxydum hydrargyri rubrum, ii.	36	Oxymel narcissi pratensis, ii.	259	Panacea Kornmann's, i.	314
Oxydum hydrargyri rubrum per acidum nitricum, ii.	36	Oxymel oxyduli cupri, i.	591	Panacea lapsorum, i.	279
Oxydum hydrargyri saccharatum, ii.	16	Oxymel primulæ veris, ii.	465	Panada anglorum, ii.	332
Oxydum hydrargyri salinum, ii.	84	Oxymel pectoral, ii.	118	Panado, ii.	332
Oxydum hydrargyri sulphuricum, ii.	82	Oxymel saccharine, i.	43	Panavæ lignum, i.	570
Oxydum hydrargyricum, ii.	36	Oxymel of squill, ii.	574	Panax quinquefolium, ii.	331
Oxydum hydrargyricum præcipitatum, ii.	32	Oxymel of subacetate of copper, i.	591	Panchymagogum minerale, ii.	46
Oxydum hydrargyrosus, ii.	32	Oxymel simple of subacetate of copper, i.	592	Panchymagogus quercetanus, ii.	46
Oxydum kadmii, ii.	145	Oxymel tartareous, ii.	450	Pancratii veri radix, ii.	563
Oxydum mangani nativum, ii.	217	Oxymel of tobacco, ii.	263	Panes saccharati purgantes, i.	545
Oxydum magnesi nigrum, ii.	217	Oxymurias calcis, i.	351	Panicum miliaceum, ii.	332
Oxydum platinæ, ii.	368	Oxymurias hydrargyri, ii.	66	Panis, ii.	332
Oxydum plumbi album, ii.	371	Oxymuriate of mercury, ii.	66	Panis cassaviensis, i.	632
Oxydum plumbi carbonatum per acidum acetosum, ii.	371	Oxynitras hydrargyri et ammoniæ, ii.	84	Panis cydoniorum, i.	598
Oxydum plumbi fusum, ii.	369	Oxyphosphas calcis, i.	361	Panis cydoniorum cum aromatibus, i.	598
Oxydum plumbi purum, ii.	371	Oxysaccharum, i.	43	Panis cydoniorum compositus, i.	598
Oxydum plumbi rubrum, ii.	370	Oxysaccharum digitalis, i.	615	Panis porcini radix, ii.	597
Oxydum plumbi semivitreum, ii.	369	Oxymel, i.	43	Papaveris albi semen, ii.	336
Oxydum stibii ablutum, i.	240	Oxymel sambuci, ii.	553	Papaveris corniculati radix, i.	720
Oxydum stibii non ablutum, i.	240	Oxymel scillæ, ii.	574	Papaver rhæas, ii.	334
		Oxymel scilliticum, ii.	574	Papaver somniferum, ii.	336
		Oxymel simplex, i.	43	Paper caustic, ii.	657
		Oxymel tabaci, ii.	263	Paper waxed, i.	437
		Oxymel antispasmodic, i.	304	Paralyseos radix, ii.	465
		Oxymel of belladonna, i.	311	Pareira brava, i.	505
		Oxymel of colchicum, i.	540	Pareira bravæ radix, i.	505
		Oxymel of cowslip, ii.	465	Paridis radix, ii.	342
		Oxymel of elder, ii.	553	Parietaria officinalis, ii.	341
		Oxymel of garlic, i.	115	Parigline, ii.	593
		Oxymel of iron, i.	664	Paris quadrifolia, ii.	342
		Oxymel of wild narcissus, ii.	259	Parnassia palustris, ii.	342
				Paronychia argentea, ii.	342
				Paronychia herba, i.	299
				Parsley, i.	268
				Parsley cow's, i.	444
				Parsley Macedonian, i.	345
				Parsnip, ii.	343

Parsnip cow, ii.	51	Pastilles emmenagogue,		Pepper Jamaica, ii.	359
Parthenii herba, ii.	225	i.	638	Pepper long, ii.	359
Partridge-berry, i.	696	Pastilles ferruginous,		Pepper white, ii.	359
Paspalum dactylum, ii.	342	i.	638	Peppermint, ii.	234
Pasque flower, i.	217	Pastilles of ginseng, ii.	332	Pepperwort, ii.	185
Passæ ficus, i.	675	Pastilles of Iceland moss,		Perca fluviatilis, i.	354
Passula anthelminticæ,		ii.	191	Perch stones, i.	354
ii.	492	Pastilles kermesine, i.	246	Perchloruret of mercury,	
Passulæ laxativæ, i.	403	Pastilles Lagneau's, ii.	16	ii.	66
Pasta althææ, i.	752	Pastilles of lemon, i.	520	Perfoliata herba, i.	345
Pasta amygdalina, i.	206	Pastilles mercurial, ii.	45	Periploca indica, ii.	593
Pasta cacaotina, ii.	686	Pastilles mucilaginous, i.	768	Periploca scammonia, ii.	560
Pasta cacaotina licheni-		Pastilles odoriferous, i.	334	Perlæ, i.	354
fera, ii.	188	Pastilles pectoral, i.	247, ii.	Peroxide of copper, i.	581
Pasta cacaotina saccha-			292	Peroxide of iron, i.	643
rata, ii.	686	Pastilles of red poppy,		Peroxide of manganese,	
Pasta emulsiva, i.	206	ii.	336	ii.	217
Pasta glycyrrhizæ, i.	724	Pastilles stimulant, i.	204,	Peroxydum ferri, i.	643
Pasta de gummo ara-		230, ii.	362	Peroxydum maganesiini-	
bico, i.	752	Pastilles of stramonium,		grum nativum, ii.	217
Pasta gummosa saccha-		i.	606	Peroxydum plumbi, ii.	371
rata, i.	752	Pastilles vermifuge, ii.	29,	Persica vulgaris, ii.	344
Pasta lichenifera, ii.	191		51, 60	Persicaria spotted, ii.	406
Pasta limacina, i.	705	Pastilles Barthez' ver-		Persicariæ minoris herba,	
Pasta liquiritiæ, i.	724	mifuge, ii.	51	ii.	406
Pasta liquiritiæ fusca,				Persicariæ urentis herba,	
i.	725	Pastilles of vinegar, i.	42	ii.	406
Pasta ad nævos, i.	350	Pastinaca opoponax, ii.	321	Persimmon, i.	619
Pasta regia, i.	206	Pastinaca sativa, ii.	343	Personatæ radix, i.	172
Paste almond, i.	206	Patæ lupinæ herba, ii.	185	Persulphas hydrargyri,	
Paste amygdaline cos-		Pavonæ lignum, i.	570	ii.	82
metic, i.	207	Pavonæ stercus, i.	83	Persulphate of mercury,	
Paste arsenical, i.	286	Peach tree, ii.	344	ii.	82
Paste astringent, i.	156	Pearl mother of, i.	353	Persulphuret of tin, ii.	625
Paste Else's caustic,		Pearls, i.	354	Petroleum, ii.	345
ii.	414	Pearlash, ii.	419	Petroselinî radix, i.	268
Paste for chilblains, i.	161	Pearlash prepared, ii.	419	Petty broom, i.	708
Paste cocoa, ii.	686	Peas caustic, ii.	657	Phaseolus communis,	
Paste of dates, ii.	347	Peas chick, i.	461	ii.	346
Paste depilatory, i.	282	Peas grey, i.	461	Phellandrium aquaticum,	
Paste of florentine iris,		Peau de Goulard, ii.	382	ii.	346
ii.	133	Pediluvium alkaline, ii.	423	Philonium Romanum,	
Paste of gum arabic,		Pediluvium ex armoracia,		ii.	320
i.	752	ii.	481	Peucedanum officinale,	
Paste liquorice, i.	724	Pediluvium irritant, ii.	481	ii.	343
Paste anisated liquorice,		Pediluvium sedative, ii.	153	Peucedanum silaus,	
i.	725	Pedis cati herba, i.	230	ii.	343
Paste opiate liquorice,		Pedis columbi herba, i.	718	Phœnicites, i.	354
i.	725	Pedis leonis radix, i.	104	Phœnix dactylifera, ii.	347
Paste of Iceland moss,		Pelias berus, i.	697	Phosphas ammoniæ, i.	190
ii.	191	Pellitory of Spain, i.	230	Phosphas calcicum sti-	
Paste against nævi, i.	350	Pellitory of the wall, ii.	341	biatum, i.	236
Paste pectoral, i.	751	Peltigera aphthosa, ii.	186	Phosphas calcis, i.	359
Paste rubefacient and		Peltigera canina, i.	186	Phosphas calcis stibia-	
febrifuge, ii.	537	Peltigera horizontalis,		tus, i.	236
Paste of snails, i.	705	ii.	186	Phosphas cupri, i.	583
Pastilles antihysterical, i.	565	Peltigera polydactyla, ii.	186	Phosphas ferri, i.	651
Pastilles of balsam of		Peltigera venosa, ii.	186	Phosphas hydrargyri,	
tolu, i.	323	Penis of the bull, i.	698	ii.	81
Pastilles of berberries,		Penis of the stag, i.	698	Phosphas magnesiæ, ii.	210
i.	335	Penis of the whale, i.	698	Phosphas natricum, ii.	610
Pastilles Chaussier's, ii.	300	Pencea sarcocolla, i.	765	Phosphas sodæ, ii.	610
Pastilles of elecampane,		Pentaphylli radix, ii.	465	Phosphate of ammonia,	
ii.	116	Peony, ii.	328	i.	190
Pastilles emetic, i.	620	Pepper black, ii.	358	Phosphate of soda, ii.	610
		Pepper Guinea, ii.	359		

- Phosphate precipitated of lime, i. 360
- Phosphate of mercury, ii. 81
- Phosphate of soda and of ammonia, ii. 611
- Phosphorus, ii. 347
- Phosphorus pulverized, ii. 348
- Phosphuretum calcis, i. 350
- Plirasi herba, ii. 222
- Phyllanthus emblica, ii. 257
- Phyllitidis herba, ii. 577
- Physalis alkekengi, ii. 350
- Physcia nivalis, ii. 186
- Physcia prunastri, ii. 188
- Physeter macrocephalus, i. 441
- Phytolacca decandra, ii. 351
- Pichurim, ii. 680
- Picrotoxa, i. 529
- Picrotoxinum, i. 529
- Pierre divine, i. 585
- Pigmentum indicum, ii. 351
- Pigmentum urucu, ii. 352
- Pila marina, ii. 352
- Pilæ damarum, i. 338
- Pilewort, ii. 481
- Pills absorbent, i. 633, ii. 492, 606
- Pills of acetate of lead, ii. 384
- Pills of acetate of lead and of ipecacuanha, ii. 385
- Pills mercurial of aconite, ii. 73
- Pills saponaceous aloetic, i. 121
- Pills ferruginous alterative, i. 649
- Pills anodyne, i. 250
- Pills anthelmintic, ii. 52
- Pills anti-asthmatic, i. 393
- Pills antiherpetic, i. 327, ii. 30, 54, 59, 74
- Pills antihydrotic, i. 747, ii. 63, 65
- Pills antihysteria, i. 425, ii. 101
- Pills anti-icteric, i. 141, 310, ii. 50
- Pills antilacteal, i. 367, ii. 58, 614
- Pills compound antimonial, i. 239, ii. 55
- Pills of aloes, i. 121
- Pills of aloes and agaric, i. 122
- Pills of aloes and assafoetida, i. 122
- Pills of aloes and gum ammoniac, i. 127
- Pills of aloes and of myrrh, i. 135
- Pills saponaceous of aloes and assafoetida, i. 122
- Pills of aloes and tartarized antimony, i. 122
- Pills aloetic, i. 121
- Pills compound aloetic, i. 121
- Pills ferruginous aloetic, i. 125
- Pills alterative, i. 250, 734, ii. 31, 50, 63
- Pills compound alterative, ii. 53
- Pills simple alterative, ii. 52
- Pills against amaurosis, i. 302
- Pills of amber, i. 128
- Pills of ambergris, i. 166
- Pills mercurial ammoniacal, ii. 94
- Pills Anderson's, i. 127
- Pills angelic, i. 140
- Pills anodyne and sudorific, i. 264
- Pills anthelmintic, i. 41, ii. 51, 54, 56, 267, 583, 714
- Pills anti-arthritis, i. 186, 243, 250, 252, 265, 734, ii. 33, 59
- Pills antidiarrhœic, ii. 146
- Pills antidysenteric, i. 475, ii. 303
- Pills anti-emetic, ii. 297
- Pills anti-epileptic, i. 275, ii. 708
- Pills antifebrile, i. 483
- Pills antigonorrhœic, i. 558, ii. 29
- Pills Klein's antimonial, i. 243
- Pills antiphlogistic, ii. 384
- Pills antirheumatic, ii. 59
- Pills antiscrofulous, i. 327
- Pills antiseptic, i. 367
- Pills antispasmodic, i. 527, ii. 300
- Pills antistruumic, ii. 29
- Pills antisiphilitic, ii. 33, 50, 51, 55, 58, 59, 74, 76, 79, 94
- Pills Kœmpf's anti-siphilitic, ii. 55
- Pills antiveneereal, ii. 26
- Pills aperient, i. 121, 125
- Pills ferruginous aperient, i. 637
- Pills aromatic aperient, i. 122
- Pills aperitive, ii. 433
- Pills of arseniate of iron, i. 289
- Pills arsenical, i. 285
- Pills Asiatic, i. 285
- Pills of assafoetida, i. 299
- Pills compound of assafoetida, i. 300
- Pills of assafoetida and of iron, i. 302
- Pills of assafoetida and of squill, i. 302
- Pills astringent, i. 160, 319, 638, 760, ii. 57, 146, 148, 663, 674, 693
- Pills of aurate of ammonia, i. 314
- Pills Baldinger's, ii. 32
- Pills balsamic, i. 47, 141, 318, 760, ii. 271, 663, 674
- Pills ferruginous balsamic, i. 760
- Pills opiate balsamic, ii. 292
- Pills of belladonna, i. 310
- Pills Belloste's, ii. 26
- Pills benzoic, i. 47
- Pills bitter, i. 760, ii. 492
- Pills aloetic bitter, i. 137
- Pills cathartic bitter, i. 136
- Pills of bitter-sweet, ii. 615
- Pills of brucina, i. 343
- Pills camphorated, i. 367
- Pills of cantharides, i. 388
- Pills mercurial of cantharides, i. 388
- Pills opiate of cantharides, i. 388
- Pills of carburet of iron, i. 639
- Pills carminative, i. 633
- Pills cathartic, ii. 61
- Pills of celandine, i. 456
- Pills cephalic, i. 549
- Pills chalybeate, i. 637, 642, 648
- Pills aperient chalybeate, i. 674
- Pills alkaline of charcoal, ii. 417
- Pills of chloruret of gold, i. 313
- Pills of cinchona, i. 475
- Pills camphorated of cinchona, i. 483
- Pills of cocoa nut oil, ii. 687
- Pills compound of colocyath, i. 124
- Pills corroborant, i. 584, 760
- Pills demulcent, i. 751
- Pills deobstruent, i. 643
- Pills of deutochloruret of mercury and of opium, ii. 76
- Pills diaphoretic, i. 239, ii. 32, 53
- Pills diaphoretic and sedative, i. 247

Pills of digitalis, i.	613	Pills mercurial, ii. 19, 26, 58,	Pills compound saponaceous of protochloruret of mercury and of jalap, ii.	61
Pills dinner, i.	128	59, 78		
Pills diuretic, ii.	65, 567	Pills of mercurial ointment, ii.		
Pills mercurial diuretic, ii.	85	21		
Pills eccoprotic, i.	125	Pills Moscati's mercurial, ii.		
Pills compound of elaterium, ii.	243	33		
Pills emetic, i.	584	Pills Nolte's mercurial, ii.		
Pills emmenagogue, i.	126, 137, 302, 637, 642, 648, 747, ii.	94		
57, 142, 247		Pills Plenck's mercurial, ii.		
Pills English, i.	122	16		
Pills excitant, ii.	267	Pills Plenck's compound mercurial, ii.		
Pills expectorant, i. 250, ii.	56, 125, 567	16		
Pills English febrifuge, i.	475	Pills purgative mercurial, ii.		
Pills ferruginous, i.	636	27		
Pills fetid, i.	572	Pills of mercury, ii.		
Pills compound of galbanum, i.	683	Pills glycyrrhized of mercury, ii.		
Pills compound gamboge, i.	127, 695	17		
Pills golden, i.	142	Pills mellated of mercury, ii.		
Pills for gonorrhœa, i.	368	17		
Pills to terminate gonorrhœa, ii.	57	Pills of mercury and of turpentine, ii.		
Pills Griffiths', i.	648	18		
Pills guaiacine, i.	740	Pills of musk, ii.		
Pills of gum ammoniac, i.	745	250		
Pills laxative gum, i.	122	Pills narcotic, i. 465, ii.		
Pills Hahnemann's, ii.	85	279		
Pills of hemlock, i.	465	Pills nauseant, ii.		
Pills compound of hound's tongue, ii.	301	125		
Pills hydragogue, i. 613, 570, ii.	56, 61, 243, 455, 566	Pills Neapolitan, ii.		
Pills Bontius' hydragogue, i.	127	26		
Pills Janin's hydragogue, ii.	56	Pills of nitric oxide of mercury, ii.		
Pills of hydrochlorate of manganese, ii.	217	38		
Pills of hydriodate of potass, ii.	122	Pills opiate, ii.		
Pills James', i.	239	295		
Pills incisive, i. 114, ii.	567	Pills saponaceous of opium, ii.		
Pills of iodine, ii.	121	299		
Pills of ioduret of mercury, ii.	80	Pills optic, i.		
Pills kermesine, i.	247	125		
Pills camphorated of kino, ii.	146	Pills oriental, ii.		
Pills laxative, i.	549	301		
Pills ferruginous laxative, i.	642	Pills pectoral, i. 565, ii.		
Pills Leslie's, ii.	63	566		
Pills lithontriptic, ii.	605	Pills opiate pectoral, ii.		
Pills Hoffmann's major, ii.	72	279		
Pills Machiavel's, i.	136	Pills physagogue, ii.		
Pills of mastic, i.	128	566		
		Pills Plummer's, ii.		
		52		
		Pills compound Plummer's, ii.		
		53		
		Pills polychrest, ii.		
		714		
		Pills Pringle's, ii.		
		63		
		Pills prophylactic for smallpox, ii.		
		52		
		Pills Meglin's, ii.		
		101		
		Pills of protochloruret of mercury, ii.		
		50		
		Pills of protochloruret of mercury and of camphor, ii.		
		57		
		Pills of protochloruret of mercury and of guaiac, ii.		
		59		
		Pills of protochloruret of mercury and of opium, ii.		
		63		
		Pills compound of protochloruret of mercury and of jalap, ii.		
		61		
		Pills compound of protochloruret of mercury and of opium, ii.		
		56		
		Pills saponaceous of protochloruret of mercury and of jalap, ii.		
		61		

Pills styptic, ii.	385	Pilulæ aureæ, i.	142	Pilulæ ferri compositæ, i.	648
Pills of subcarbonate of copper and of ammonia, i.	582	Pilulæ balsamicæ Junkeri, ii.	674	Pilulæ e ferro laxantes, i.	642
Pills of subsulphate of copper and of ammonia, i.	587	Pilulæ balsamicæ mor-tonianæ, i.	47	Pilulæ ferulacæ, cum ammonio, i.	300
Pills sudorific, i.	734	Pilulæ balsamicæ cum opio, ii.	292	Pilulæ fetidæ, i.	300
Pills of sulphate of iron, i.	652	Pilulæ benedictæ, i.	126	Pilulæ galbani composi-tæ, i.	683
Pills of sulphuret of anti-mony, i.	242	Pilulæ benzoicæ, i.	47	Pilulæ e galbano mar-tiatæ, i.	636
Pills of sulphuret of anti-mony and of lime, i.	254	Pilulæ bilis, i.	633	Pilulæ gambogiæ com-positæ, i.	695
Pills compound of sul-phuret of mercury and of antimony, ii.	31	Pilulæ in bronchitide lenta, ii.	347	Pilulæ e gentiana croca-tæ, i.	137
Pills of toxicodendron, ii.	504	Pilulæ cachecticæ, i.	637	Pilulæ Griffithii, i.	648
Pills traumatic, ii.	663	Pilulæ cæruleæ, i.	587	Pilulæ guaiaci, i.	740
Pills of turpentine, ii.	661	Pilulæ cambogiæ com-positæ, i.	127	Pilulæ guaiaci cum hy-drargyro stibio et ex-tractis narcoticis, i.	243
Pills of veratrine, ii.	713	Pilulæ camphoratæ, i.	367	Pilulæ guaiaci cum sti-bio, i.	252
Pills Vigarou's, ii.	50	Pilulæ carbonum kali-næ, ii.	417	Pilulæ guaiacinæ, i.	740
Pills Vogler's, ii.	73	Pilulæ catarrhales, ii.	61	Pilulæ de gummatibus, i.	683
Pills Werlhof's, ii.	64	Pilulæ e chelidonio, i.	456	Pilulæ gummi guttæ aloeticæ, i.	127
Pills Zeller's, ii.	87	Pilulæ chinæ alumina-tæ, i.	160	Pilulæ gummosæ, i.	300, 683
Pilosellæ herba, ii.	6	Pilulæ conii cinnaba-rinæ, ii.	45	Pilulæ gummosæ laxati-væ, i.	683
Pilulæ aloes amaræ, i.	137	Pilulæ cocchii, i.	124	Pilulæ de hiera, i.	122
Pilulæ aloes et cincho-næ, i.	137	Pilulæ coccinæ, i.	124	Pilulæ hieræ cum agarico, i.	122
Pilulæ aloes cum colo-cynthide, i.	124	Pilulæ cochæ, i.	124	Pilulæ hydrargyri, ii.	17, 85, 455
Pilulæ aloes crocatæ, i.	135	Pilulæ cocciæ, i.	124	Pilulæ hydrargyri gum-mosæ, ii.	16
Pilulæ aloeticæ martia-les, i.	125	Pilulæ colocynthidis com-positæ, i.	124	Pilulæ hydrargyri jala-pinæ, ii.	61
Pilulæ aloes cum myrrha, i.	135	Pilula colocynthidis ex-tracti compositi, i.	575	Pilulæ hydrargyri oxy-muriatis, ii.	94
Pilulæ aluminis, i.	160	Pilulæ communes, i.	135	Pilulæ ex hydrargyro catharticæ, ii.	60
Pilulæ amaræ, i.	711	Pilulæ conii, i.	465	Pilulæ ex hydrargyro diureticæ, ii.	85
Pilulæ amaro-catharticæ, i.	136	Pilulæ e conio, i.	465	Pilulæ ex hydrargyro muriatico, ii.	52
Pilulæ ammoniuretæ cu-pri, i.	587	Pilulæ cupri ammoniaci, i.	587	Pilulæ ex hydrargyro muriatico miti, ii.	50
Pilulæ analepticæ, i.	239	Pilulæ e cupro ammo-niaco, i.	587	Pilulæ ex hydrargyro nitrico, ii.	87
Pilulæ anethinæ, i.	125	Pilulæ de cynoglossa, ii.	301	Pilulæ ex hydrargyro oxydulato nigro, ii.	85
Pilulæ anglicæ, i.	122	Pilulæ deobstruentes, i.	745, 566	Pilulæ hydropicæ, i.	127
Pilulæ ante cibum, i.	137	Pilulæ deprimentes, ii.	246	Pilulæ hypnoticæ, ii.	292
Pilulæ anthelminticæ pur-gantes, ii.	52	Pilulæ depurantes, ii.	53	Pilulæ hystericæ, i.	683
Pilulæ antihystericæ, i.	300, 250	Pilulæ diaphoreticæ sti-biatæ, i.	239	Pilulæ iodatæ, ii.	121
Pilulæ antispasmodicæ, i.	275	Pilulæ diureticæ resol-ventes, ii.	566	Pilulæ e kino et campho-ra, ii.	146
Pilulæ antispasticæ, i.	300	Pilulæ diureticæ robo-rantes, ii.	673	Pilulæ laxantes, ii.	61
Pilulæ antisiphiliticæ et resolventes, ii.	50	Pilulæ de duobus, i.	142	Pilulæ laxantes cum mer-curio, ii.	61
Pilulæ antiveneræ, ii.	38	Pilulæ ecphracticæ, i.	136, 492	Pilulæ lucis majores, i.	125
Pilulæ aperientes lax-an-tes, i.	141	Pilulæ Emanuelis, i.	136	Pilulæ lunares, i.	275
Pilulæ aperientes Stahl-ii, i.	637	Pilulæ emmenagogæ, ii.	250		
Pilulæ aperitivæ, ii.	433	Pilulæ leviter emmena-gogæ deprimentes, ii.	247		
Pilulæ assafoetidæ com-positæ, i.	300	Pilulæ excitantes, i.	275		
		Pilulæ extracti catholici, ii.	561		
		Pilulæ fellis tauri amaræ, i.	633		

<i>Pilulæ manganensii muriatici</i> , ii.	217	<i>Pilulæ roborantes ad blennorrhœam</i> , ii.	492	<i>Pimpinellæ hircinæ radix</i> , ii.	356
<i>Pilulæ menagogæ catharticæ</i> , i.	142	<i>Pilulæ Rufi</i> , i.	135	<i>Pimpinellæ minoris radix</i> , ii.	465
<i>Pilulæ ad menstrua</i> , i.	126	<i>Pilulæ rupicaprarum</i> , i.	338	<i>Pimpinellæ nalicæ radix</i> , ii.	556
<i>Pilulæ mercuriales</i> , ii.	61	<i>Pilulæ saponacæ</i> , ii.	458	<i>Pimpinellæ nigræ radix</i> , ii.	356
<i>Pilulæ mercuriales Plenckii</i> , ii.	16	<i>Pilulæ saponis cum opio</i> , ii.	299	<i>Pincneya pubescens</i> , i.	469
<i>Pilulæ ex mercurio dulci</i> , ii.	50	<i>Pilulæ scilliticæ</i> , ii.	565	<i>Pine ground</i> , ii.	681
<i>Pilulæ ex mercurio solubili Hahnemanni</i> , ii.	65	<i>Pilulæ scillæ compositæ</i> , ii.	565	<i>Pine tree</i> , ii.	358
<i>Pilulæ ad morbos cutaneos</i> , i.	250	<i>Pilulæ sedativæ</i> , ii.	292	<i>Pine wild</i> , ii.	358
<i>Pilulæ ad morbos deploratos</i> , i.	239	<i>Pilulæ solares</i> , i.	315	<i>Pinus canadensis</i> , i.	318
<i>Pilulæ muriatis hydrargyri camphoratæ</i> , ii.	57	<i>Pilulæ specificæ</i> , ii.	94	<i>Pinus maritima</i> , ii.	654
<i>Pilulæ e myrrha compositæ</i> , i.	648	<i>Pilulæ stibiata cum arnica et camphora</i> , i.	250	<i>Pinus mugho</i> , ii.	664
<i>Pilulæ myrrhæ amaræ cum ammoniaco</i> , i.	760	<i>Pilulæ stibiata cum cicuta</i> , i.	250	<i>Pinus mughos</i> , i.	318
<i>Pilulæ myrrhæ cum arnica</i> , i.	760	<i>Pilulæ stibiata cum opio et laxantibus resinosis</i> , i.	243, 252	<i>Pinus pinea</i> , ii.	358
<i>Pilulæ myrrhæ compositæ</i> , i.	300	<i>Pilulæ ex stibio et hydrargyro</i> , ii.	52	<i>Pinus sylvestris</i> , ii.	358, 654
<i>Pilulæ myrrhæ ferrulacæ</i> , i.	300	<i>Pilulæ stimulantés ad gonorrhœam</i> , ii.	146, 673	<i>Piper album</i> , ii.	359
<i>Pilulæ nauseam cientes</i> , ii.	125	<i>Pilulæ stomachicæ</i> , i.	137	<i>Piper betel</i> , ii.	360
<i>Pilulæ opiatæ</i> , ii.	295	<i>Pilulæ e storace</i> , ii.	298, 628	<i>Piper caudatum</i> , ii.	360
<i>Pilulæ opii compositæ</i> , ii.	301	<i>Pilulæ strychnini</i> , ii.	626	<i>Piper cubeba</i> , ii.	360
<i>Pilulæ ex opio</i> , ii.	295	<i>Pilulæ submuriatis hydrargyri compositæ</i> , ii.	53	<i>Piper Hispanicum</i> , ii.	359
<i>Pilulæ opticæ</i> , i.	125	<i>Pilulæ subsulphatis ammonico-cuprici</i> , i.	587	<i>Piper Jamaicense</i> , ii.	359
<i>Pilulæ pectorales</i> , i.	745, ii. 566	<i>Pilulæ de succino</i> , i.	128	<i>Piperis indicî fructus</i> , ii.	359
<i>Pilulæ perpetuæ</i> , i.	234	<i>Pilulæ sulphatis potassæ aloeticæ</i> , i.	127	<i>Piper indicum</i> , ii.	359
<i>Pilulæ physagogæ</i> , ii.	566	<i>Pilulæ syphiliticæ</i> , ii.	38	<i>Piper longum</i> , ii.	359
<i>Pilulæ picis</i> , ii.	366	<i>Pilulæ systema nervorum summopere velligantes</i> , i.	275	<i>Piper nigrum</i> , ii.	358
<i>Pilulæ Plummeri</i> , ii.	52, 53	<i>Pilulæ nervosum et musculare systema velligantes</i> , ii.	266	<i>Piper tabasci</i> , ii.	359
<i>Pilulæ Polychrestæ</i> , ii.	38	<i>Pilulæ tartaræ Bontii</i> , i.	127	<i>Piper turcicum</i> , ii.	359
<i>Pilulæ polychrestæ balsamicæ</i> , i.	136	<i>Pilulæ e tartaro boraxato</i> , ii.	455	<i>Pipera</i> , ii.	359
<i>Pilulæ polychrestæ Starkei</i> , ii.	714	<i>Pilulæ e terebinthina</i> , ii.	673	<i>Piperinum</i> , ii.	359
<i>Pilulæ purgantes</i> , ii.	61	<i>Pilulæ torpentes et ressolventes</i> , ii.	154	<i>Pisa pro fonticulis</i> , ii.	657
<i>Pilulæ purgantes cum hydrargyro muriatico miti</i> , ii.	60	<i>Pilulæ toxicodendri</i> , ii.	504	<i>Pissa cedria</i> , ii.	364
<i>Pilulæ ex resina jalapæ</i> , ii.	6	<i>Pilulæ de tribus</i> , i.	141	<i>Pistachio tree</i> , ii.	364
<i>Pilulæ resolventes</i> , i.	745, ii. 458	<i>Pilulæ valerianæ ferulacæ</i> , i.	300	<i>Pistacia lentiscus</i> , ii.	223
<i>Pilulæ resolventes cum mercurio</i> , ii.	53	<i>Pilulæ ad variolam</i> , ii.	52	<i>Pistacia terebinthus</i> , ii.	654
<i>Pilulæ resolventes cum rheo</i> , ii.	492	<i>Pilulæ veneræ</i> , i.	587	<i>Pistacia vera</i> , ii.	364
<i>Pilulæ rhabarbarinæ antacidæ</i> , ii.	492	<i>Pimenta</i> , ii.	359	<i>Pistolochiæ radix</i> , i.	276
<i>Pilulæ rhei</i> , ii.	491	<i>Pimento</i> , ii.	359	<i>Pitch</i> , ii.	673
<i>Pilulæ e rheo aceticæ</i> , ii.	492	<i>Pimpernel</i> , ii.	465	<i>Pitch Burgundy</i> , ii.	673
<i>Pilulæ e rheo cum natro acetico</i> , ii.	614	<i>Pimpernel red</i> , i.	215	<i>Pitch Jew's</i> , i.	298
		<i>Piminel saxifrage</i> , ii.	356	<i>Pix atra</i> , ii.	673
		<i>Pimpinella anisum</i> , ii.	352	<i>Pix Burgundica</i> , ii.	673
		<i>Pimpinella magna</i> , ii.	356	<i>Pix cedria</i> , ii.	364
		<i>Pimpinella saxifraga</i> , ii.	356	<i>Pix Græca</i> , ii.	673
		<i>Pimpinellæ albæ radix</i> , ii.	356	<i>Pix liquida</i> , ii.	364
				<i>Pix navalis</i> , ii.	673
				<i>Pix solida</i> , ii.	673
				<i>Pix vegetabilis</i> , ii.	673
				<i>Plantago cynops</i> , ii.	367
				<i>Plantago lanceolata</i> , ii.	366
				<i>Plantago major</i> , ii.	366
				<i>Plantago media</i> , ii.	366
				<i>Plantago psyllium</i> , ii.	367
				<i>Plantain clammy</i> , ii.	367
				<i>Plantain great</i> , ii.	366
				<i>Plantain hoary</i> , ii.	366
				<i>Plantain small</i> , ii.	366
				<i>Platina</i> , ii.	368
				<i>Plaster adhesive</i> , ii.	679
				<i>Plaster ferruginous adhesive</i> , ii.	680

- Plaster saponaceous ammoniacal, i. 193
- Plaster André de la Croix, ii. 677
- Plaster anglo-saxon, ii. 399
- Plaster anodyne, ii. 301, 302, 391
- Plaster anti-apoplectic, ii. 661
- Plaster anti-arthritis, ii. 400
- Plaster anticolical, ii. 302
- Plaster antihæmorrhoidal, ii. 315
- Plaster antihernial, ii. 642
- Plaster antihysterical, i. 689
- Plaster antimonial, i. 262
- Plaster anti-odontalgic, ii. 224
- Plaster antirachitic, ii. 677
- Plaster antispasmodic, ii. 300
- Plaster aromatic, i. 399, 749, ii. 165
- Plaster astringent, ii. 391
- Plaster balsamic, i. 321, ii. 400, 464
- Plaster basilic, ii. 303
- Plaster of bayberries, ii. 182
- Plaster of belladonna, i. 311
- Plaster of betony, i. 337
- Plaster black of lead, ii. 369
- Plaster Bechholz's black, i. 690
- Plaster Mejean's blistering, i. 387
- Plaster for the breasts, i. 443
- Plaster against buboes, ii. 24
- Plaster Stahl's camphorated, ii. 399
- Plaster of cantharides, i. 383
- Plaster camphorated of cantharides, i. 386
- Plaster of cantharides and euphorbium, i. 387
- Plaster capucin's, i. 757
- Plaster of carburet of iron, i. 639
- Plaster against colic, i. 690
- Plaster common, ii. 390
- Plaster for corns, i. 594
- Plaster Rajot Laforet's corn, i. 182
- Plaster corroborant, ii. 680
- Plaster corrosive, ii. 72
- Plaster court, i. 706
- Plaster of cumin, i. 580
- Plaster compound of cumin, i. 580
- Plaster saponaceous of cumin, ii. 463
- Plaster Green's defensive, ii. 677
- Plaster red defensive, ii. 371
- Plaster simple diachylon, ii. 389
- Plaster diabatanum, ii. 397
- Plaster diapalm, ii. 392
- Plaster diaphoretic, i. 687
- Plaster of digitalis, i. 618
- Plaster discutient, i. 749
- Plaster divine, ii. 396
- Plaster Doyen's, ii. 400
- Plaster of elastic gum, i. 755
- Plaster camphorated epispastic, i. 383
- Plaster epulotic, ii. 391
- Plaster of euphorbium, i. 756
- Plaster compound of euphorbium, i. 756
- Plaster against fractures, ii. 644
- Plaster of frog's spawn, ii. 401
- Plaster of galbanum, i. 684
- Plaster camphorated of galbanum, i. 684
- Plaster compound of galbanum, i. 685
- Plaster crocated of galbanum, i. 687
- Plaster for goitre, i. 262
- Plaster of gum ammoniac, i. 747
- Plaster compound of gum ammoniac, i. 747
- Plaster compound of gums, i. 687
- Plaster of hemlock, i. 467
- Plaster compound of hemlock, i. 467
- Plaster of henbane, ii. 104
- Plaster Hjerne's, ii. 392
- Plaster ischiatic, i. 690
- Plaster Juck's, ii. 22
- Plaster of labdanum, i. 758
- Plaster of leek, ii. 399
- Plaster magnetic, i. 283
- Plaster mammellary, ii. 399
- Plaster of melilot, ii. 226
- Plaster camphorated melilot, ii. 227
- Plaster mercurial, ii. 22
- Plaster mercurial with hemlock and opium, ii. 25
- Plaster ammoniacal mercurial, ii. 23
- Plaster of mercury with hemlock, ii. 24
- Plaster miraculous, ii. 400
- Plaster mucilaginous, i. 748
- Plaster of musk, ii. 250
- Plaster Nuremberg, ii. 399
- Plaster odontalgic, i. 386, ii. 302
- Plaster opiate, ii. 297
- Plaster opodeldoch, ii. 395
- Plaster of red oxide of iron, i. 645
- Plaster of red oxide of lead, ii. 370
- Plaster compound of pitch, ii. 677
- Plaster of pyrethrum, i. 231
- Plaster resolvent, i. 190, 468, 595, ii. 14, 25, 371, 386, 398, 679
- Plaster resolvent and anti-cancerous, ii. 418
- Plaster of the four solvents, ii. 24
- Plaster compound of rhubarb, i. 142
- Plaster rubefacient, i. 386
- Plaster saponaceous, ii. 463
- Plaster Barbette's saponaceous, ii. 400
- Plaster camphorated saponaceous, ii. 464
- Plaster simple, ii. 389, 390
- Plaster compound of spermaceti, i. 684
- Plaster splenetic, i. 749
- Plaster black sticking, i. 706
- Plaster stimulant, ii. 364
- Plaster stomachic, ii. 315, 644
- Plaster storax, ii. 627
- Plaster strengthening, i. 645
- Plaster red strengthening, ii. 393
- Plaster subcarbonate of lead, ii. 400
- Plaster of sulphur, ii. 640
- Plaster myrrhine of sulphur, i. 761, ii. 677
- Plaster suppurative, i. 345
- Plaster of tar, ii. 366
- Plaster of tobacco, ii. 264
- Plaster Vigo's, ii. 396
- Plaster volatile, i. 190
- Plaster of wax, i. 436
- Plaster yellow, ii. 326
- Plum tree, ii. 467
- Plumbago, i. 639
- Plumbago Europæa, ii. 368
- Plumbum, ii. 368
- Plumbum acetatum, ii. 382
- Plumbum oxydatum rubrum, ii. 370
- Pocgeregæ cortex, ii. 402

<i>Podophyllum peltatum</i> , ii.	402	<i>Potass alcoholated</i> , ii.	413	<i>Potio anthelmintica pa- rum laxans</i> , ii.	582
<i>Poison-oak</i> , ii.	503	<i>Potass caustic</i> , ii.	412	<i>Potio anticolica</i> , ii.	219
<i>Polanisia dodecandria</i> , ii.	402	<i>Potass caustic with lime</i> , ii.	413	<i>Potio antiphlogistico- laxans</i> , i.	265
<i>Polii cretici herba</i> , ii.	682	<i>Potas of commerce</i> , ii.	419	<i>Potio ad atrabilem</i> , ii.	448
<i>Pollen coryllorum</i> , i.	561	<i>Potass ferrocyanate of</i> , i.	61	<i>Potio carbonatis calcis</i> , i.	358
<i>Polychroïte</i> , i.	563	<i>Potass fused</i> , ii.	412	<i>Potio cathartica</i> , ii.	211
<i>Polygala amara</i> , ii.	402	<i>Potass medicinal hydro- cyanate of</i> , i.	60	<i>Potio cathartico-diapho- reticus</i> , i.	265
<i>Polygala rubelia</i> , ii.	403	<i>Potass liquid</i> , ii.	410	<i>Potio comfortans acidu- lata</i> , i.	567
<i>Polygala seneka</i> , ii.	403	<i>Potass purified</i> , ii.	419	<i>Potio cretacea</i> , i.	358
<i>Polygala vulgaris</i> , ii.	403	<i>Potassa cum calce</i> , ii.	413	<i>Potio demulcens</i> , i.	753
<i>Polygalæ Virginianæ</i> radix, ii.	403	<i>Potassa caustica</i> , ii.	412	<i>Potio emetico-laxans</i> , i.	265
<i>Polygaline</i> , ii.	403	<i>Potassa caustica cum calce</i> , ii.	413	<i>Potio frigida martialis</i> , i.	674
<i>Polygonati vulgaris radix</i> , ii.	405	<i>Potassa caustica fusa</i> , ii.	412	<i>Potio juniperina</i> , ii.	139
<i>Polygoni herba</i> , ii.	406	<i>Potassa fusa</i> , ii.	412	<i>Potio laxans</i> , i. 417, ii. 211, 219, 647	
<i>Polygonatum angulosum</i> , ii.	405	<i>Potassa impura</i> , ii.	419	<i>Potio laxativa Viennensis</i> , i.	415
<i>Polygonum aviculare</i> , ii.	406	<i>Potassa pura</i> , ii.	410	<i>Potio leniens</i> , i.	211, 752
<i>Polygonum historta</i> , ii.	406	<i>Potassa venalis</i> , ii.	419	<i>Potio mannæ tamarindi- nata</i> , ii.	219
<i>Polygonum fagopyrum</i> , ii.	407	<i>Potassæ acetæ</i> , ii.	441	<i>Potio mercurii vivi</i> , ii.	13
<i>Polygonum hydropiper</i> , ii.	406	<i>Potassæ carbonas</i> , ii.	419	<i>Potio narcotica</i> , ii.	245
<i>Polygonum persicaria</i> , ii.	406	<i>Potassæ carbonas e lixive cinere</i> , ii.	420	<i>Potio nutriens et sedans</i> , ii.	154, 189
<i>Polypodium vulgare</i> , ii.	407	<i>Potassæ carbonas e tar- tari crystallis</i> , ii.	421	<i>Potio oleosa drastica</i> , i.	570
<i>Polypody</i> , i.	308	<i>Potassæ carbonatis aqua</i> , ii.	422	<i>Potio picea</i> , ii.	365
<i>Polypody common</i> , ii.	407	<i>Potassæ nitras</i> , ii.	432	<i>Potio purgans</i> , i.	417
<i>Polyporus officinalis</i> , ii.	408	<i>Potassæ et ferri prus- sias</i> , i.	61	<i>Potio resolvens ex iodio</i> , ii.	122
<i>Polytrichum commune</i> , ii.	409	<i>Potassæ ferrocyanas</i> , i.	61	<i>Potio resolvens et laxans</i> , ii.	495
<i>Poma renetia</i> , ii.	214	<i>Potassæ hydriodas</i> , ii.	121	<i>Potio salina mannata</i> , ii.	219
<i>Poma sorbi aucupariæ</i> , ii.	619	<i>Potassæ hydrocyanas ferruginosus</i> , i.	61	<i>Potio sistens</i> , i.	566
<i>Pomatum albissimum</i> , ii.	441	<i>Potassæ subcarbonas</i> , ii.	419	<i>Potio stibiata</i> , i.	259
<i>Pomatum florum naphæ</i> , i.	509	<i>Potassæ supertartras</i> , ii.	449	<i>Potio stomachica visce- rale</i> , i.	27
<i>Pomatum orange-flower</i> , i.	509	<i>Potassæ sulphas</i> , ii.	428	<i>Potio terebinthinata</i> , ii.	655
<i>Pomatum saturni</i> , ii.	380	<i>Potassæ tartras</i> , ii.	447	<i>Potio tonico-anodyna</i> , i.	527
<i>Pomegranate tree</i> , ii.	470	<i>Potasse à l'alcohol</i> , ii.	413	<i>Potion absorbent</i> , i.	171,
<i>Pommade antihyperpetique de Goulard</i> , ii.	381	<i>Potassidum myrrhæ</i> , i.	761	ii. 283, 423	
<i>Pommade de Goulard</i> , ii.	382	<i>Potassium</i> , ii.	410	<i>Potion acidulated</i> , i.	82
<i>Pomum hierosolymita- num</i> , ii.	243	<i>Potassium cyanuret of</i> , i.	60	<i>Potion acidulous</i> , ii.	506
<i>Pomum melongenæ</i> , ii.	617	<i>Potassium proto-hydro- cyanate of</i> , i.	60	<i>Potion æthereous</i> , i.	101
<i>Poplar tree</i> , ii.	409	<i>Potassium proto-hydro- ferro-cyanate of</i> , i.	61	<i>Potion alexipharmic</i> , i.	201
<i>Poplar trembling</i> , ii.	405	<i>Potassium proto-hydro- sulpho-cyanate of</i> , i.	62	<i>Potion alkaline</i> , ii.	427
<i>Poppy</i> , ii.	336	<i>Potatoes</i> , ii.	613	<i>Potion aluminous</i> , i.	164
<i>Poppy corn</i> , ii.	334	<i>Potentilla anserina</i> , ii.	465	<i>Potion Peyrilhe's ammo- niacal</i> , i.	186
<i>Poppy red</i> , ii.	334	<i>Potentilla reptans</i> , ii.	465	<i>Potion analeptic</i> , i.	110
<i>Poppy white</i> , ii.	336	<i>Poterium sanguisorba</i> , ii.	465	<i>Potion angelica</i> , i.	225
<i>Poppy yellow horned</i> , i.	720	<i>Potio amara antifebrilis</i> , i.	91	<i>Potion Andry's anodyne</i> , ii.	17
<i>Populus balsamifera</i> , ii.	409,	<i>Potio alkalina</i> , ii.	427	<i>Potion antacid</i> , ii.	210
	644	<i>Potio angelica</i> , ii.	219	<i>Potion antihelmintic</i> , ii.	508,
<i>Populus nigra</i> , ii.	409	<i>Potio anthelmintica oleo- so-aromatica</i> , ii.	508		582
<i>Populus tremula</i> , ii.	409			<i>Potion anti-arthritic</i> , i.	734
<i>Portulaca oleracea</i> , ii.	410			<i>Potion anti-asthmatic</i> , i.	198
				<i>Potion anticatarrhal</i> , ii.	575
				<i>Potion anticroupal</i> , i.	263

- Potion antidiarrhœal, i. 200
 Potion anti-epileptic, i. 694, ii. 705
 Potion antidysenteric, ii. 281
 Potion antigonorrhœic, i. 556, ii. 360
 Potion antihæmorrhagic, i. 485, ii. 280
 Potion antiherpetic, ii. 418
 Potion antihydropic, ii. 220
 Potion antihysterical, i. 201, 425
 Potion anti-icteric, ii. 530
 Potion antilactæal, ii. 424
 Potion antilyssic, i. 198, ii. 232
 Potion antinephritic, i. 754, ii. 438
 Potion antiperiodic, ii. 281
 Potion antiphlogistic, ii. 435
 Potion antipleuritic, i. 240
 Potion antiscorbutic, i. 484
 Potion antiscrofulous, i. 490, ii. 424, 530, 606
 Potion antiseptic, i. 51, 198, 486
 Potion antispasmodic, i. 171, 201, 240, 430, 457, ii. 127, 240, 268, 316, 423
 Potion antispasmodic and tonic, i. 485
 Potion antistrumous, ii. 423
 Potion antisiphilitic, ii. 69
 Potion antitetanic, i. 182
 Potion antitoxic, ii. 127
 Potion aperitive, ii. 434, 443
 Potion for aphonia, i. 111
 Potion aromatic, i. 566
 Potion asthmatic, i. 171
 Potion astringent, i. 157, 199, 484, 494, 693, 765, ii. 146, 149, 194, 283, 407, 512, 588, 609
 Potion astringent tonic, ii. 280
 Potion astringent and tonic, i. 491
 Potion Bourquenod's, ii. 71
 Potion cardiac, i. 188
 Potion carminative, i. 171, ii. 228, 280
 Potion cathartic, ii. 211, 613
 Potion bitter chalybeate, i. 666
 Potion cold chalybeate, i. 674
 Potion chalybeate tonic, i. 666
 Potion chloric, i. 460
 Potion compound of cinchona, i. 487
 Potion comforting, i. 567
 Potion cordial, i. 567, ii. 316
 Potion demulcent, i. 212, 443, 752, 753, ii. 201, 338, 541, 724
 Potion diaphoretic, i. 202, 264
 Potion diaphoretic and pectoral, i. 240
 Potion digestive, ii. 606
 Potion diuretic, i. 97, 198, 322, 380, 445, 520, 539, 614, 617, ii. 408, 424, 434, 435, 438, 443, 444, 453, 454, 569, 572, 575, 668
 Potion diuretic and resolvent, ii. 438
 Potion diuretic and sedative, ii. 446
 Potion cordial diuretic, ii. 118
 Potion Milmann's diuretic, ii. 445
 Potion emetic, i. 248, 259, 261, 263, 294, ii. 126, 128
 Potion emeto-cathartic, i. 265
 Potion emmenagogue, i. 370, 380, 661, ii. 143, 144, 537
 Potion excitant, ii. 349, 504, 707
 Potion excitant and antispasmodic, ii. 705
 Potion ferruginous excitant, i. 666
 Potion excitant and resolvent, ii. 495
 Potion exciting, i. 495
 Potion expectorant, i. 197, 198, 260, 264, 304, 466, 617, 701, ii. 423, 574, 607, 632
 Potion expectorant and incisive, ii. 128
 Potion febrifuge, i. 91, 287, 288, 484, 537, ii. 128, 294, 434, 478, 608
 Potion febrifuge and laxative, i. 485, ii. 453
 Potion bitter febrifuge, i. 480
 Potion purgative febrifuge, i. 420
 Potion ferruginous, i. 659
 Potion against the gripes of infants, i. 499
 Potion hæmastatic, i. 36
 Potion against whooping-cough, ii. 387
 Potion hydragogue, i. 264, 456, 615, ii. 139, 280, 452
 Potion incisive, ii. 425, 576
 Potion against inebriation, i. 171
 Potion kermeticized, i. 248
 Potion laxative, ii. 430, 611, 647
 Potion laxative and resolvent, ii. 495
 Potion laxative and sedative, ii. 495
 Potion oleose laxative, i. 411
 Potion lithontriptic, ii. 210, 423, 427, 607
 Potion mercurial, ii. 87
 Potion of mercury, ii. 13
 Potion mucilaginous, i. 766
 Potion narcotic, ii. 425, 293
 Potion nitrated, ii. 434
 Potion nitrous, ii. 434
 Potion ocytic, ii. 579
 Potion oily, i. 767, ii. 270
 Potion opiate, ii. 294
 Potion opio-antimonial, i. 264
 Potion oxygenated, ii. 440
 Potion paregoric, ii. 280
 Potion pectoral, i. 322, 751, ii. 216, 222, 280, 405, 541, 617
 Potion bitter pectoral, i. 487, ii. 403
 Potion phosphorated, ii. 348
 Potion purgative, i. 248, 266, 417, 547, 551, 553, 570, ii. 210, 219, 453, 488, 493, 499, 508, 563, 612
 Potion purgative for children, ii. 449
 Potion camphorated purgative, i. 419
 Potion purgative and febrifuge, ii. 424
 Potion major purgative, i. 547
 Potion purgative and tonic, ii. 499
 Potion with pulverized cinchona, i. 479
 Potion purgative and vermifuge, ii. 493
 Potion of manna and of rhubarb, ii. 499
 Potion refrigerant, ii. 436
 Potion resolvent, i. 263, ii. 123, 430
 Potion resolvent stomachic, ii. 435
 Potion resolvent and diuretic, ii. 444
 Potion saline, ii. 211
 Potion saponaceous, ii. 423, 459
 Potion sedative, i. 311, 465, 754, ii. 102, 154, 179, 247, 279, 280, 281, 283, 341, 424, 436
 Potion with decoction of cinchona, i. 490
 Potion stimulant, i. 36, 87, 110, 187, 198, 344, 460, ii. 122, 626, 668
 Potion stimulant and tonic, i. 486

- Potion stomachic, i. 27, ii. 240, 407, 473, 528
- Potion stomachic purgative, i. 122
- Potion sudorific, i. 84, 192, 199, 260
- Potion sulphureous, ii. 418
- Potion sulphurico-saline, ii. 431
- Potion sulphurous, ii. 637
- Potion against tæniæ, ii. 668
- Potion compound tartaric, ii. 453
- Potion temperant, ii. 44, 434
- Potion against tinea, i. 557
- Potion tonic, i. 490, 229, 484, 486, 491, 492, 493, 494, 527, 566, 653, ii. 173, 316, 495
- Potion tonic and antihyperpetic, ii. 530
- Potion tonic and antispasmodic, i. 485, 567
- Potion camphorated tonic, i. 485
- Potion of turpentine, ii. 655
- Potion vermifuge, i. 120, 177, 419, 695, ii. 3, 168, 436, 583, 606, 669, 683
- Potus acidus, i. 44, 76, ii. 506
- Potus analepticus, i. 110
- Potus antiphlogisticus diureticus, ii. 438
- Potus antiphlogisticus diaphoreticus et leviter laxans, i. 260
- Potus antiphlogisticus leviter laxans, i. 260
- Potus ad biennorrhæam, i. 556
- Potus carminativus, ii. 141
- Potus cerasorum, i. 439
- Potus citri stibiatus, i. 260
- Potus communis incitans, i. 110
- Potus diureticus, ii. 141
- Potus diureticus simplex, ii. 444
- Potus hydrocyanicus, i. 60
- Potus mercurialis Belleti, ii. 87
- Potus muriatico-acidus, i. 55
- Potus oryzæ citratus, ii. 325
- Potus oxygenata, ii. 440
- Potus oxymellicus, i. 44
- Potus ex pomis, ii. 215
- Potus sacchari lactis, ii. 150
- Potus spirituosus, i. 110
- Poultice carrot, i. 607
- Poudre de la princesse de Carignan, ii. 330
- Powder against abortion, ii. 520
- Powder absorbent, i. 238, ii. 208, 490, 579
- Powder of aconite, i. 84
- Powder nitrated of aconite, i. 84
- Powder aerophoric, i. 49
- Powder alexipharmic, i. 156, 277, ii. 436
- Powder Rollwag's alexipharmic, i. 156
- Powder alexiteric, ii. 125
- Powder alkaline gum, i. 750
- Powder of aloes with cannella, i. 128
- Powder compound of aloes, i. 126, 128
- Powder alterative, ii. 31
- Powder compound alterative, ii. 53
- Powder alterative and resolvent, ii. 429
- Powder simple alterative, ii. 52
- Powder of alum, i. 159
- Powder compound of alum, i. 159
- Powder ammoniacal, i. 186
- Powder camphorated ammoniacal, i. 186
- Powder analeptic, ii. 43
- Powder anodyne, i. 340, ii. 292, 297, 299
- Powder anonymous, i. 768
- Powder antacid, i. 356
- Powder anthelmintic, i. 544, 653, ii. 43, 704
- Powder anti-arthritic, i. 84, 710, 733, ii. 561
- Powder bitter anti-arthritic, i. 625
- Pulvis arthriticus, i. 84
- Powder anticachectic, i. 635, 645, 649
- Powder anticalculus, i. 271, 505
- Powder anticatarrhal, ii. 330, 346, 433, 634
- Powder antichlorotic, i. 636
- Powder antidiarrhæic, ii. 124
- Powder antidysenteric, ii. 297, 634
- Powder anti-epileptic, ii. 44, 330
- Powder anti-erysipelatus, ii. 32
- Powder antifebrile, i. 472, 663
- Powder anti hæmoptysic, i. 765
- Powder anti hæmorrhagic, ii. 642
- Powder antihyperpetic, ii. 31
- Powder antihydropsic, i. 613
- Powder antihysteria, i. 434
- Powder anti-icteric, i. 402
- Powder antileucorrhæic, ii. 223, 680
- Powder antilyssic, ii. 443, 115, 188, 35.
- Powder count d'Issembourg's antilyssic, i. 358
- Powder antimonial, i. 236
- Powder antimonial febrifuge, i. 471
- Powder antimoniated, i. 261
- Powder Stahl's antinephritic, i. 364
- Powder antiperiodic, ii. 478
- Powder antiphlogistic, ii. 429
- Powder antiphthitical, i. 351, 483, 584, ii. 384
- Powder antipleuritic, i. 361, ii. 634
- Powder antipsoric, ii. 384, 633
- Powder antirachitic, ii. 530
- Powder antirheumatic, i. 250
- Powder antiscrofulous, ii. 255
- Powder antiseptic, i. 472, 474
- Powder antispasmodic, i. 246, 424, 474, 612, ii. 55, 249, 300, 704, 724
- Powder antistrumous, i. 357, 392, ii. 29
- Powder aperitive, ii. 433
- Powder aphrodisiac, ii. 249
- Powder aromatic, ii. 161, 522
- Powder aromatic of elecampane, ii. 115
- Powder aromatic ferruginous, i. 652
- Powder aromatic nutritive, ii. 161
- Powder arsenical, i. 284
- Powder Baumann's arsenical, i. 285
- Powder Justamond's arsenical, i. 284
- Powder Pleneiz' arsenical, i. 285
- Powder Plunket's arsenical, i. 284
- Powder Rousselot's arsenical, i. 284
- Powder compound of asarum, i. 294
- Powder astringent, i. 156, 159, 472, 583, 645, 649, 765, ii. 297, 492
- Powder astringent antifebrile, i. 471
- Powder astringent and demulcent, i. 750
- Powder of avens, i. 719
- Powder of avens and rubarb, i. 719

- Powder basilic, ii. 53
 Powder of belladonna, i. 309
 Powder benzoic, i. 47
 Powder English bezoardic, i. 238
 Powder Halle's bezoardic, i. 238
 Powder cachectic, i. 641
 Powder Camerarius', i. 361
 Powder camphorated, i. 367
 Powder camphorated temperant, ii. 436
 Powder capucin's, ii. 714
 Powder of carburet of iron, i. 639
 Powder against caries, i. 299
 Powder carminative, ii. 125, 353
 Powder cathartic, i. 118
 ii. 64
 Powder Plenk's caustic, ii. 37
 Powder Kruger's caustic, ii. 75
 Powder compound of celtic nard, ii. 703
 Powder cephalic, i. 337, ii. 44, 261, 520
 Powder compound of chalk, i. 356
 Powder of chloruret of gold, ii. 313
 Powder camphorated of cinchona, i. 471
 Powder of chocolate with cinchona, i. 471
 Powder compound of cinchona, i. 473
 Powder of cinchona with muriate of ammonia, i. 474
 Powder compound of cinamon, ii. 161
 Powder citrine, ii. 131
 Powder against colic, ii. 422
 Powder of colocynth, i. 572
 Powder corroborant, i. 472
 Powder cosmetic, i. 90, 207
 ii. 458
 Powder demulcent, i. 146, 750
 Powder nitrous demulcent, i. 750
 Powder dentifrice, i. 357, 361, 391, 474, ii. 450, 633
 Powder Pelletier's dentifrice, ii. 478
 Powder white dentifrice, ii. 470
 Powder deobstruent, ii. 208, 403
 Powder Jasser's depurative, i. 243
 Powder desicative, i. 367
 Powder for diabetes, i. 157
 Powder diaphoretic, i. 362, 733, ii. 55, 62, 439, 633
 Powder diaphoretic and sedative, ii. 299
 Powder diarrhodon, ii. 520
 Powder digestive, i. 265
 ii. 64, 207, 211, 353, 422, 429
 Powder compound digestive, ii. 429
 Powder Klein's digestive, ii. 448
 Powder of digitalis, i. 612
 Powder diuretic, i. 271, 388, 612, ii. 137, 438, 613, 614
 Powder ferruginous diuretic, ii. 437
 Powder eccoprotic, ii. 452
 Powder ephratic, ii. 452
 Powder of elder, ii. 552
 Powder embalming, i. 135
 Powder emetic, i. 263, 584
 ii. 125
 Powder compound emetic, i. 261
 Powder Helvetius' emetic, i. 263
 Powder Marryat's emetic, i. 262
 Powder emeto-cathartic, ii. 124
 Powder emmenagogue, i. 388, 636, ii. 142
 Powder errhine, i. 541, 756
 ii. 58, 261, 360, 714
 Powder against erysipelas, ii. 370, 551
 Powder escharotic, i. 160, 583, 595, ii. 37, 94, 384
 Powder expectorant, i. 245, 250, 330, ii. 63, 125
 Powder of extract of anemone, i. 218
 Powder factitious of cinchona, i. 90
 Powder febrifuge, i. 192, 238, 250, 284, 483, 587, 709, 719, ii. 125, 362
 Powder febrifuge and anticachectic, i. 472
 Powder febrifuge and incisive, i. 472
 Powder cathartic febrifuge, i. 472
 Powder Fizes' febrifuge, i. 473
 Powder Fuller's febrifuge, i. 472
 Powder Gaubius' febrifuge, i. 473
 Powder Hoffmann's febrifuge, i. 472
 Powder Jadelot's febrifuge, i. 473
 Powder Leschude's febrifuge, i. 474
 Powder nervine febrifuge, i. 473
 Powder saline febrifuge, ii. 439
 Powder compound of fennel, i. 220
 Powder ferruginous, i. 642
 Powder ferruginous tonic, i. 472
 Powder Fontaneilles', i. 285
 Powder Fordyce's, ii. 456
 Powder fumigatory, ii. 629
 Powder Grimaldi's, i. 649
 Powder of gum and sulphur, i. 750
 Powder for gutta serena, i. 218
 Powder Haly's, i. 213
 Powder red hepatic, ii. 450
 Powder hippiatric, i. 243
 Powder for hooping cough, i. 246
 Powder Hungary, i. 157
 Powder hydragogue, i. 549
 ii. 64
 Powder compound of jalap, i. 543
 Powder incisive, i. 246, 261, 364, ii. 684
 Powder for infants, i. 357
 Powder for ink, i. 653
 Powder for intertrigo, i. 357
 Powder compound of ipecacuanha, ii. 298
 Powder mercurial of ipecacuanha, ii. 60
 Powder kermesine, i. 245
 Powder camphorated kermesine, i. 246
 Powder compound of kino, ii. 146
 Powder Knox's, i. 351
 Powder laxative, i. 402
 ii. 634
 Powder laxative for children, ii. 60
 Powder Tronchin's laxative and temperant, ii. 452
 Powder of iron filings, i. 635
 Powder for lemonade, i. 82
 Powder compound of leopard's-bane, i. 279
 Powder against lice, i. 609
 Powder compound of liquorice, i. 402
 Powder lithontriptic, i. 271
 Powder Ludwig's, i. 361
 Powder of lupuline, ii. 11
 Powder compound of madder, ii. 530
 Powder English martial, i. 635
 Powder masticatory, ii. 114

- Powder laxative mercurial, ii. 55
 Powder of musk, ii. 249
 Powder compound of musk, ii. 249
 Powder compound of myrrh, i. 759
 Powder nauseant, i. 612
 ii. 124
 Powder nitro-camphorated, ii. 436
 Powder nutritive, i. 699
 Powder of nux vomica, ii. 265
 Powder compound of oak bark, ii. 475
 Powder ocytic, ii. 363
 Powder odoriferous, ii. 520
 Powder compound of olibanum, ii. 271
 Powder ophthalmic, i. 630
 ii. 353
 Powder Baldinger's ophthalmic, ii. 451
 Powder opiate, ii. 295
 Powder opiate febrifuge, i. 473
 Powder opio-antimonial, i. 264
 Powder of factitious orpiment, i. 282
 Powder of grey oxide of mercury, ii. 33
 Powder of nitric oxide of mercury, ii. 37
 Powder paretic, ii. 208
 Powder pectoral, i. 192, 239, 246, 402, ii. 150, 337, 635
 Powder Plummer's, ii. 52
 Powder compound Plummer's, ii. 53
 Powder of protochloruret of mercury, ii. 48
 Powder of protochloruret of mercury and of opium, ii. 62
 Powder purgative, i. 264, 633, 730, ii. 207, 211
 Powder purgative and diuretic, i. 612
 Powder absorbent purgative, ii. 634
 Powder English purple, i. 358
 Powder Struensee's pyric, ii. 440
 Powder Quarin's, i. 246
 Powder refrigerant, ii. 437
 Powder resolvent, i. 264
 ii. 64, 346, 448
 Powder pectoral resolvent, ii. 634
 Powder compound resolvent, i. 239
 Powder Stahl's resolvent, i. 239
 Powder restorative, ii. 686
 Powder of rhubarb and of magnesia, ii. 491
 Powder saponaceous of rhubarb, ii. 471
 Powder of rhubarb and of sulphate of potass, ii. 490
 Powder of rhubarb and of supertartrate of potass, ii. 490
 Powder Ruding's, i. 235
 Powder of salep, ii. 540
 Powder saline, ii. 452
 Powder compound saline, ii. 211
 Powder against salivation, ii. 417
 Powder compound saturnine, ii. 372
 Powder compound of scammony, ii. 560
 Powder scidlitz, i. 50
 Powder sedative, i. 612
 ii. 297
 Powder compound of senna, i. 402
 Powder sialagogue, i. 192
 Powder of nitrate of silver, i. 275
 Powder soothing, i. 48
 Powder compound of squill, ii. 564
 Powder sternutatory, i. 230, 336
 Powder stimulant, ii. 57, 139, 178, 490
 Powder stomachic, i. 292, 472, 473, 527, 709, ii. 353
 Powder of strychnine, ii. 626
 Powder styptic, ii. 384
 Powder sudorific and tonic, ii. 593
 Powder of sulphuret of antimony, i. 242
 Powder of sulphuret of mercury and of antimony, ii. 31
 Powder of superoxalate of potass, ii. 446
 Powder of sweet flag, i. 86
 Powder compound of tansey, ii. 649
 Powder tartarized nitrous, ii. 438
 Powder temperant, ii. 384, 439
 Powder Bonillon Lagrange's temperant, ii. 438
 Powder Rosenstein's temperant, ii. 439
 Powder Stahl's temperant, ii. 44
 Powder Unzer's temperant, ii. 439
 Powder ferruginous of tin, ii. 623
 Powder against tinea, i. 327
 Powder tonic, i. 472, 559, 567, 583, 642, 649, 652, 674, 675
 Powder tonic and anti-spasmodic, i. 635
 Powder tonic and cathartic, i. 473
 Powder ferruginous tonic, i. 636
 Powder tonic and incisive, i. 246
 Powder tonic and stomachic, i. 636
 Powder compound of tragacanth, i. 767
 Powder of vanilla, ii. 712
 Powder vermifuge, i. 239, 308, 730, ii. 2, 17, 29, 48, 56, 59, 60, 65, 142, 582, 603, 620, 623, 625
 Powder mercurial vermifuge, ii. 29
 Powder Vernix's, i. 584
 Precipitatus ruber, ii. 36
 Precipitatus solubilis niger, ii. 64
 Prassii herba, ii. 222
 Precipitate black, ii. 84
 Precipitate green, ii. 38
 Precipitate perse, ii. 36
 Precipitate red, ii. 36
 Precipitate rose, ii. 82
 Precipitate white, ii. 47, 90
 Precipitate Wurtz', ii. 81
 Precipitate yellow, ii. 82
 Priapus cervi, i. 698
 Priapus ceti, i. 698
 Priapus tauri, i. 698
 Prickly ash, i. 270
 Primula officinalis, ii. 465
 Prinos verticillatus, ii. 467
 Privet, ii. 198
 Propolis, ii. 467
 Proto-acetas barii, i. 328
 Proto-acetate of barium, i. 328
 Proto-acetate of calcium, i. 363
 Proto-acetate of lead, ii. 382
 Proto-acetate of mercury, ii. 95
 Proto-acetate of potassium, ii. 441
 Proto-acetate of sodium, ii. 614
 Proto-carbonate of potassium, ii. 426
 Proto-carbonate of potassium and of ammonia, ii. 427
 Proto-carbonate of sodium, ii. 607

Proto-chlorate of potassium, ii.	440	Proto-oleomargarate of potassium, ii.	456	Protoxi-sulphuret of potassium, ii.	416
Proto-chloruret of antimony, i.	254	Proto-oleomargarate of sodium, ii.	615	Protoxi-sulphuret of sodium, ii.	602
Proto-chloruret of iron, i.	656	Proto-phosphate of magnesium, ii.	210	Protoxi-sulphuretum calcis, i.	350
Proto-chloruret of mercury, ii.	46	Proto-sulphas barii, i.	328	Protoxydum antimonii, i.	234
Proto-chloruret ferrated of mercury, ii.	58	Proto-sulphate of barium, i.	328	Protoxydum hydrargyri nitrico-ammoniacale, ii.	84
Proto-chloruretum hydrargyri, ii.	46	Proto-sulphate of calcium, i.	361	Protoxydum stibii griseum, i.	234
Proto-citrate of calcium, i.	364	Proto-sulphate of iron, i.	651	Prune, ii.	467
Proto-citrate of iron, i.	666	Proto-sulphate of magnesium, ii.	210	Prunella vulgaris, ii.	465
Proto-citrate of potassium, ii.	446	Proto-sulphate of potassium and of ammonia, ii.	431	Pruneoli sylvestris cortex, ii.	467
Proto-gallate of lead, ii.	389	Proto-sulphate of potassium, ii.	427	Prunus domestica, ii.	467
Proto-hydriodate of potassium, ii.	121	Proto-sulphate of potassium and of magnesium, ii.	197	Prunus spinosa, ii.	467
Proto-hydrochloras antimonii, i.	254	Proto-sulphate of sodium, ii.	611	Prussias lixivie et ferri, i.	61
Proto-hydrochlorate of calcium, i.	362	Proto-sulphite of potassium, ii.	432	Prussian blue, i.	674
Proto-hydrochlorate of iron, i.	656	Proto-sulphite of sodium, ii.	614	Prussiate of potass and of iron, i.	674
Proto-hydrochlorate of iron solid or crystallized, i.	656	Proto-sulphuret of copper, i.	581	Pseudacori radix, ii.	130
Proto-hydrochlorate of magnesium, ii.	213	Proto-sulphuretum antimonii, i.	241	Pseudo-angustura, i.	228
Proto-hydrochlorate of potassium, ii.	440	Proto-tartrate of antimony, i.	255	Psoralea glandulosa, ii.	468
Proto-hydrosulphate of antimony, i.	243	Proto-tartrate of antimony and of potass, i.	255	Psoralea stripe-flowered, ii.	468
Proto-hydrosulphate of calcium, i.	362	Proto-tartrate of potassium, ii.	447	Psychotria emetica, ii.	124
Proto-hyposulphate of sodium, ii.	614	Proto-tartrate of potassium and of ammonia, ii.	456	Psyllii semines, ii.	367
Proto-hydrosulpho-cyanate of iron, i.	675	Proto-tartrate of potassium and of iron, i.	667	Ptarmicæ radix, i.	37
Proto-hyposulphate of potassium, ii.	432	Proto-tartrate of potassium and of sodium, ii.	455	Ptarmicæ montanæ radix, i.	279
Proto-iodate of mercury, ii.	89	Protoxi-chloruret of lime, i.	351	Pteris aquilina, ii.	468
Proto-ioduret of mercury, ii.	79	Protoxi-chloruret of sodium, ii.	602	Pterocarpus draco, i.	765
Proto-malate of iron, i.	665	Protioxide of antimony, i.	234	Pterocarpus santalinus, i.	765
Proto-meconate of sodium, ii.	615	Protioxide of barium, i.	327	ii.	191
Proto-nitras barii, i.	328	Protioxide of calcium, i.	347	Ptisan acid, i.	520
Proto-nitrate of barium, i.	328	Protioxide of cobalt, i.	526	Ptisan Mascagni's alkaline, ii.	423
Proto-nitrate of lead, ii.	375	Protioxide of lead, ii.	369	Ptisan alterative, ii.	617
Proto-nitrate of mercury, ii.	85	Protioxide of potassium, ii.	410	Ptisan Vigarou's alterative, ii.	597
Proto-nitrate of mercury and of ammonia, ii.	89	Protioxide of sodium, ii.	601	Ptisan antilacteal, i.	444
Proto-nitrate of potassium, ii.	432	Protioxide of tin, ii.	624	Ptisan antiscorbutic, i.	536
Proto-nitrate of sodium, ii.	614	Protoxidum antimonii sulphuratum, i.	241	ii.	482
Proto-oleomargarate of lead, ii.	389	Protoxi-phosphuret of calcium, i.	360	Ptisan antiscrofulous, i.	493
				Ptisan antisyphilitic, ii.	599
				Ptisan Arabian, ii.	595
				Ptisan astringent, ii.	642, 693
				Ptisan camphorated, i.	371
				Ptisan of cherry juice, i.	438
				Ptisan of cichory, i.	462
				Ptisan demulcent, i.	148
				Ptisan diuretic, i.	396, ii. 141, 424, 536
				Ptisan of elecampane, ii.	117
				Ptisan febrifuge, i.	537
				Ptisan Feltz', ii.	596
				Ptisan Jussierand's, ii.	598
				Ptisan compound of lemon, i.	521
				Ptisan Lisbon, ii.	598
				Ptisan pectoral, i.	148, ii. 150

Ptisan purgative anti-syphilitic, ii.	597	Pulvis absorbens, ii.	208	Pulvis ad arenulas, i.	505
Ptisan purgative and sudorific, ii.	599	Pulvis absorbens pro usu externo, ii.	579	Pulvis ari alkalinus, i.	292
Ptisan Sainte-Marie's, ii.	595	Pulvis ex acido tartarico, i.	82	Pulvis ari compositus, i.	292
Ptisan sudorific, i. 273, ii.	596	Pulvis acidulus, i.	82	Pulvis aromatico-ferratus, i.	641
Ptisan Astruc's sudorific, ii.	598	Pulvis aerophorus, i.	49	Pulvis aromaticus, ii.	161
Ptisan vermifuge, i.	308	Pulvis ex Æthiope antimoniali, ii.	31	Pulvis aromaticus ferratus, i.	652
Ptisan Vinache's, ii.	598	Pulvis adstringens, i. 765, ii.	297	Pulvis asari compositus, i.	294
Ptisan vulnerary, i.	280	Pulvis alexiterius, ii.	298	Pulvis atramentarius, i.	653
Ptisana Callaci, ii.	599	Pulvis aluminosus Helvetii, i.	159	Pulvis ex baccis lauri compositus, ii.	255
Ptisana Datisbii, i.	521	Pulvis alexipharmicus sinicus, ii.	250	Pulvis basilicus, ii.	53
Ptisana Hippocratica, ii.	7	Pulvis algarothi, i.	254	Pulvis Bernhardi, i.	284
Ptisana laxativa, i.	415	Pulvis aloes cum canella, i.	128	Pulvis bezoardicus albus Ludovici, i.	361
Ptisana oryzæ, ii.	325	Pulvis aloes cum guaiaco, i.	128	Pulvis bezoardicus anglicanus, i.	236
Ptisana ex pane, ii.	332	Pulvis aluminis compositus, i.	159	Pulvis bezoardicus Curri, i.	474
Ptisana passulata, i.	415	Pulvis aluminosus, i.	159	Pulvis bezoardicus dia-phoreticus, i.	238
Puff-ball, ii.	204	Pulvis aluminoso-mercurialis, ii.	37	Pulvis bezoardicus Halensis, i.	238
Pukeweed American, ii.	351	Pulvis amarus ferratus, i.	653	Pulvis bezoardicus Kentianus, i.	238
Pulegii herba, ii.	234	Pulvis analepticus, ii.	43	Pulvis bezoardicus Sennertii, i.	356
Pulicariæ semines, ii.	367	Pulvis angelicus, i.	254	Pulvis bezoardicus stibiatus, i.	238
Pulmonaria officinalis, ii.	469	Pulvis anodynus, ii.	298	Pulvis cachecticus Ludovici, i.	236
Pulmonariæ gallicæ herba, ii.	6	Pulvis antacida, ii.	208	Pulvis ad calculum, i.	505
Pulp of cassia, i.	421	Pulvis infantum antacidus, ii.	208	Pulvis camphoratus, i.	367
Pulp of cassia for lavements, i.	421	Pulvis anthelminticus, i. 653, ii.	17, 582	Pulvis capitalis sancti-angeli, i.	294
Pulp of dates, ii.	347	Pulvis anti-arthritis purgans, ii.	561	Pulvis carbonatis calcis compositus, i.	356
Pulp emollient, i.	146	Pulvis anti-arthritis purgativus, i.	402	Pulvis cardiacus, ii.	43
Pulp of figs, i.	675	Pulvis anticolicus cemerarii, ii.	422	Pulvis carthusianorum, i.	243
Pulp of hips, ii.	511	Pulvis antihectico-scrofulous, ii.	255	Pulvis caryocostinus, ii.	560
Pulp of lily, i.	198	Pulvis antilyssicus Tun-chinensis, ii.	250	Pulvis castoreatus, i.	424
Pulp of onion, i.	112	Pulvis antilyssicus Ormskirkianus, ii.	115	Pulvis contra casum, i.	765
Pulp of prunes, ii.	468	Pulvis antilyssus, ii. 188, 537		Pulvis catarrhalis, ii.	634
Pulp of squill, ii.	563	Pulvis antimonialis, i.	236, 242	Pulvis catharticus, ii.	490
Pulp of tamarinds, ii.	645	Pulvis antimonialis cum scammonio, i.	239	Pulvis catharticus absorbens, ii.	491, 634
Pulpa cassiæ, i.	421	Pulvis antisepticus, i.	474	Pulvis catholicus, ii.	560
Pulpa cassiæ fistulæ, i.	421	Pulvis antispasmodicus, ii.	44	Pulvis ad botium, i.	392
Pulpa cassiæ pro clysteribus, i.	421	Pulvis antispasmodicus infantum, ii.	724	Pulvis causticus, ii.	37
Pulpa cynorrhodontos, ii.	511	Pulvis antispasmodicus et anticonvulsivus stomachicus, i.	424	Pulvis cellensis aureus, ii.	43
Pulpa dactylorum, ii.	347	Pulvis antispasticus, ii.	704	Pulvis cephalicus, ii.	330
Pulpa ebuli, ii.	551	Pulvis aperitivus, ii.	433	Pulvis cervinus Camerarii, i.	361
Pulpa ficarum, i.	675	Pulmonariæ arboreæ herba, ii.	188	Pulvis e cerussa compositus, ii.	372
Pulpa lili, ii.	198			Pulvis chamomillæ cum antimonio, i.	238
Pulpa prunorum, ii.	468			Pulvis Cheltenhamensis, ii.	211
Pulpa scillæ, ii.	563			Pulvis chinæ, i.	474
Pulpa tamarindorum, ii.	645				
Puls arsenicalis, i.	285				
Pulsatillæ cæruleæ radix, i.	217				
Pulsatillæ nigricantis herba, ii.	217				
Pulsatillæ vulgaris radix, i.	217				
Pulticula de manna, ii.	218				
Pulticula saponacea, ii.	492				
Pulvis contra abortum, ii.	520				

Pulvis chinæ antisepticus, i.	474	Pulvis evacuans puero- rum, ii.	60	Pulvis ex hydrargyro sti- biato-sulphurato, ii.	31
Pulvis chocolatæ china- tæ, i.	471	Pulvis in excoriatione ægri, i.	357	Pulvis hydrargyri stibiato- sulphurato cum cicuta, ii.	31
Pulvis e cinchona cathar- ticus, i.	473	Pulvis ad excrescentias, i.	160	Pulvis hydrargyri stibia- tus cum hyosciamo, ii.	55
Pulvis colocynthidis, i.	572	Pulvis expectorans, i.	245, 246	Pulvis hypnoticus, ii.	28
Pulvis comitis Warwick, i.	239	Pulvis exsiccans et inci- tans, i.	367	Pulvis Jacobi, i.	236
Pulvis ad condienda in- tranea et cavitates im- plendas, i.	135	Pulvis febrifugus anti- monialis, i.	236	Pulvis jalapæ compositus, i.	543
Pulvis ad condiendas et infarciendas carnes, i.	135	Pulvis febrifugus Mor- toni, i.	238	Pulvis jalapæ hydrargy- ratus, i.	543
Pulvis ad condiendum cor, i.	135	Pulvis febrifugus et ner- vinus, i.	473	Pulvis jalapæ cum hy- dragyro muriatico miti, i.	543
Pulvis Constantini, ii.	97	Pulvis febrifugus Unzeri, i.	250	Pulvis jalapæ cum kali sulphurico, i.	543
Pulvis cordialis, ii.	43	Pulvis ferratus, i.	642	Pulvis jalapæ salinus, i.	543
Pulvis Cornachini, i.	239	Pulvis ferri alcoholisatus, i.	634	Pulvis jalapæ tartaricus, i.	543
Pulvis cornu usti cum opio, ii.	295	Pulvis febrifugus pau- perum, i.	719	Pulvis Jamesii, i.	236, 238
Pulvis Cosmii, i.	284	Pulvis ferri limati, i.	635	Pulvis imperatoris, ii.	161
Pulvis cretæ compositus, i.	356	Pulvis ad flatum, ii.	353	Pulvis incidens, ii.	448
Pulvis cretæ compositus cum opio, ii.	295	Pulvis contra fluorem album, ii.	680	Pulvis incrassans Fulleri, i.	750
Pulvis dentrificus, ii.	450	Pulvis fœniculi compo- situs, i.	220	Pulvis pro infantibus, ii.	208
Pulvis deprimens et nau- seam ciens, i.	612	Pulvis fumalis, i. 332, ii	629	Pulvis infantum cum rheo, ii.	491
Pulvis diacarthami, ii.	560	Pulvis fumalis nobilis, i.	333	Pulvis Johannis, ii.	36
Pulvis diaphoreticus, i.	362, 298	Pulvis ad fumigationes muriaticas, i.	458	Pulvis Johannis de Vigo, ii.	36
Pulvis diapnoicus, ii.	298	Pulvis galactopoicus, ii.	208	Pulvis ipecacuanhæ com- positus, ii.	298
Pulvis diaromaton, ii.	161	Pulvis galactopoieticus, i.	220	Pulvis ipecacuanhæ opia- tus, ii.	298
Pulvis diarrhodon, ii.	520	Pulvis gratiolæ cum hy- drargyro, i. 730, ii.	56	Pulvis ipecacuanhæ et opii, ii.	298
Pulvis diascordii, ii.	317	Pulvis gnaiaei cum senna, i.	402	Pulvis ipecacuanhæ cum opio, ii.	298
Pulvis diasennæ, i.	402	Pulvis gummosus, i. 750, 767	767	Pulvis iridis cum nitro, ii.	433
Pulvis digestivus, ii. 429, 471	471	Pulvis gummosus cum liquiritia et sulphure stibiato, i.	250	Pulvis pro juseulo extem- poraneo, i.	699
Pulvis digestivus cum rheo, ii.	429	Pulvis gummosus stibia- tus, i.	245	Pulvis Kermesianus, i.	245
Pulvis digitalis purpureæ, i.	612	Pulvis de gutteta, ii.	330	Pulvis Kermesinus, i.	245
Pulvis diureticus, ii.	438	Pulvis ad guttetam, ii. 44, 330	330	Pulvis Kermesinus cum camphora, i.	246
Pulvis Doveri, ii.	298	Pulvis hepaticus ruber, ii.	450	Pulvis kino compositus, ii.	146
Pulvis ad dysuriam, i.	146	Pulvis ad herpetem, ii.	31	Pulvis kino gummosus, ii.	146
Pulvis ecphracticus Sellii, ii.	208	Pulvis hieræ picræ, i.	128	Pulvis kinosatus, i.	159
Pulvis emmenagogus, i.	950	Pulvis hydrargyri cum camphora, ii.	57	Pulvis lauri carbonicus, ii.	178
Pulvis epilepticus, ii. 330, 704	330	Pulvis cinereus hydrar- gyri, ii.	32	Pulvis laxans, i. 402, 543, ii.	491
Pulvis epilepticus Hol- satis, ii.	331	Pulvis hydrargyri cina- reas, ii.	84	Pulvis laxans cum mer- curio, i.	543
Pulvis epilepticus mar- chionum, ii.	208, 330	Pulvis hydrargyri cum opio, ii.	62	Pulvis laxans cum sale, i.	543
Pulvis epilepticus ruber, ii.	44	Pulvis hydrargyri oxydati rubri, ii.	37	Pulvis laxativus vegeta- bilis, i.	543
Pulvis equorum, i.	243	Pulvis ex hydrargyro ru- bro cum alumine, ii.	37		
Pulvis errhinus, i.	294				
Pulvis ad erysipelam, ii.	551				
Pulvis externus ad erysi- pelas, i.	32				
Pulvis erysipelatodes, ii.	370				
Pulvis erysipelatodes ex- ternus, i.	32				

Pulvis leniens antihypochondriacus Kleinii, ii.	448	Pulvis opiatius, ii.	295	Pulvis roborans antispasmodicus, ii.	704
Pulvis lenitivus, ii.	490	Pulvis oxydi Stibii compositus, i.	239	Pulvis sacchari acidulus, i.	82
Pulvis lenitivus tartarizatus, i. 402, ii.	490	Pulvis pæoniæ compositus, ii.	330	Pulvis salino-stibiatus, i.	264
Pulvis pro limonada, i.	82	Pulvis pannonicus ruber, i.	157	Pulvis salinus, ii.	452
Pulvis liquiritiæ compositus, i.	402	Pulvis ad partum promovendum, ii.	609	Pulvis salinus compositus, ii.	44, 211
Pulvis ad lumbricos, ii.	60	Pulvis pectoralis, i. 246, 402	402	Pulvis saponis compositus, i.	633
Pulvis lycopodii, ii.	204	Pulvis pectoralis balsamicus, ii.	635	Pulvis contra scabiem, ii.	633
Pulvis ad maculas corneæ, ii.	451	Pulvis pectoralis resolvens, i. 250, ii.	634	Pulvis scammonæ compositus, ii.	560
Pulvis magnesiæ compositus, ii.	469	Pulvis Peruvianus Banjii, i.	473	Pulvis scammonii compositus, ii.	560
Pulvis magnesiæ tartaricus, i.	49	Pulvis ex phellandro aquatico gummosus, ii.	346	Pulvis scillæ compositus, ii.	564
Pulvis manualis, i.	90	Pulvis pleuriticus, i.	361	Pulvis sedativus, ii.	297
Pulvis marchionis, ii.	330	Pulvis pleuriticus Mynsichti, ii.	634	Pulvis sennæ compositus, i.	402
Pulvis martialis, i.	635	Pulvis e plumbo, ii.	372	Pulvis sennæ cum sulphure, i.	402
Pulvis mercurialis argenteus, ii.	97	Pulvis precipitans Stahlian, i.	238	Pulvis sodæ tartaricus, i.	49
Pulvis e mercurio, ii.	55	Pulvis principis, ii.	36	Pulvis spongæ compositus, i.	392
Pulvis e mercurio dulci, ii.	48	Pulvis ad procidentiam ani, i.	156	Pulvis stanni, ii.	623, 624
Pulvis ad mictum involuntarium, i.	649	Pulvis ad prolapsum ani, i.	156	Pulvis stanni martiatus, ii.	623
Pulvis mitigans Wepferi, i.	146	Pulvis puerorum citrinus, ii.	131, 208	Pulvis sternutatorius, i.	294, 714
Pulvis ad morsum canis rabidi, ii.	537	Pulvis puerorum epilepticus cum rheo, ii.	491	Pulvis sternutatorius albus, ii.	252
Pulvis moschi compositus, ii.	250	Pulvis purgans, i. 543, ii.	490	Pulvis stibiatus, i.	236, 261
Pulvis moschi opiatius, ii.	249	Pulvis purgans anthelmintica, ii.	64	Pulvis stomachicus Birkmanni, i.	292
Pulvis motum peristalticum invertens, ii.	124	Pulvis purgans et irritans, i.	730	Pulvis ad strumas, i.	392
Pulvis muriatis hydrargyri thebaicus, ii.	62	Pulvis purificans, ii.	31	Pulvis contra strumas, i.	392
Pulvis myrrhæ compositus, i.	759	Pulvis purificans Kämpferi, i.	242	Pulvis stypticus, i. 159, ii.	146
Pulvis nasalis odoratus, ii.	520	Pulvis purpureus anglicus, i.	358	Pulvis sudorificus, ii.	298
Pulvis natri carbonici cum acido tartarico, i.	49	Pulvis pyricus, ii.	440	Pulvis sudorificus Doveri, ii.	298
Pulvis nervinus, ii.	704	Pulvis contra quartanam, i.	472	Pulvis pro suffimigio, ii.	629
Pulvis nitri oxalicus, ii.	439	Pulvis quercûs marinæ, i.	390	Pulvis ad suffiandum, ii.	629
Pulvis nitri tartaricus, ii.	438	Pulvis radicis jalapæ cum natro, i.	543	Pulvis ex sulphure stibiato, i.	250
Pulvis nitri thebaicus, ii.	297	Pulvis refrigerans, i. 82, ii.	452	Pulvis ex sulphure stibiato rubro cum camphora, i.	246
Pulvis nitroso-camphoratus, ii.	436	Pulvis relaxans, i.	264	Pulvis ex sulphure stibiato rubro, i.	246
Pulvis nitrosus, ii. 438, 439	439	Pulvis resolvens antiphlogisticus, ii.	403	Pulvis sulphuris compositus, i.	128
Pulvis nitrosus cinnabarinus, ii.	44	Pulvis rhei antihæcticus, ii.	490	Pulvis sulphureosus, ii.	634
Pulvis nobilis Cellensis, ii.	43	Pulvis rhei compositus, ii.	490	Pulvis tanacetii compositus, ii.	649
Pulvis nucis vomicæ, ii.	265	Pulvis rhei cum magnesia, ii.	471	Pulvis tartari argillatus, ii.	451
Pulvis nucum moschatarum compositus, ii.	255	Pulvis rhei salinus, ii.	469	Pulvis tartari compositus, ii.	450, 452
Pulvis nutricum, i. 220, ii.	208	Pulvis rhei cum tartaro, ii.	490	Pulvis temperans, ii.	452
Pulvis ophthalmicus, ii.	451				

Pulvis temperans albus, ii.	438	Quassia amara, ii.	472	Regulus antimonii medicinalis, i.	233
Pulvis temperans camphoratus, ii.	436	Quassia excelsa, ii.	472	Regulus antimonii simplex, i.	233
Pulvis temperans diureticus, ii.	438	Queen of the meadows, ii.	621	Regulus antimonii stellatus, i.	233
Pulvis temperans ruber, ii.	44	Querculæ minoris herba, ii.	681	Regulus antimonii martialis, i.	233
Pulvis terebinthinatus, ii.	668	Quercus alba, ii.	474	Regulus of antimony, i.	233
Pulvis thebaicus, ii.	295	Quercus cerris, i.	692	Regulus martial of antimony, i.	233
Pulvis tonico-adstringens, i.	472	Quercus ilex, ii.	474	Regulus medicinal of antimony, i.	233
Pulvis tonico-antispasmodica martialis, i.	635	Quercus infectoria i.	692	Regulus simple of antimony, i.	233
Pulvis tonico-martialis, i.	636	Quercus marina, i.	678	Regulus stibii, i.	233
Pulvis tonicus Banjii, i.	473	Quercus pedunculata, ii.	474	Remedium antiphthisicum Adairi, i.	584
Pulvis tormentillæ compositus, ii.	692	Quercus robur, ii.	474	Remedium antitæniolum, ii.	48
Pulvis ad tormina infantum, ii.	471	Quercus suber, ii.	475	Remedy Mittie's antisyphilitic, ii.	135
Pulvis ad tormina infantum antacidus, ii.	208	Quercus tinctoria, ii.	474	Remedy Redfearn's antidiabetic, ii.	425
Pulvis tragacanthæ compositus, i.	767	Quickgrass, ii.	696	Rennet apple, ii.	214
Pulvis de tribus, i.	239	Quicksilver, ii.	12	Requies Nicolai, ii.	320
Pulvis contra tussim infantum, ii.	330	Quina, ii.	477	Resin of amber, ii.	630
Pulvis valerianæ aromaticus, ii.	704	Quince tree, i.	597	Resin of benzoin, i.	331
Pulvis ad vermes, ii.	60	Quinine, ii.	477	Resin of jalap, i.	548
Pulvis contra vermes, ii.	29, 582	Quinquefolii majoris radix, ii.	465	Resin prepared of jalap, i.	549
Pulvis vermifugus, ii.	582	Quinquina, i.	470	Resin of scammony, ii.	560
Pulvis Viennensis albus virgineus, i.	362	Radish turnip, ii.	481	Resin yellow, ii.	673
Pulvis Vigonis, ii.	36	Radix Brasiliensis, ii.	124	Resina alba, ii.	673
Pulvis vitæ imperatoris, ii.	161	Radix Lopeziana, ii.	479	Resina anime, i.	741
Pulvis ad ulcera tendinum, i.	759	Ragwort, ii.	584	Resina fluida canadensis, i.	318
Pumex, ii.	470	Raisins purgative, i.	403	Resina carannæ, i.	754
Pumpion i.	579	Raisins vermifuge, ii.	492	Resina cayennensis, i.	755
Pumpkin, i.	579	Ramenta ferri, i.	634	Resina copallina, i.	755
Punch, i.	111	Rana esculenta, i.	697	Resina corticis Peruviani, i.	482
Punica granatum, ii.	470	Ranunculi albi herba, i.	217	Resina elastica, i.	755
Puppies i.	697	Ranunculi nemorosi herba, i.	217	Resina elemi, ii.	486
Purgative Grant's, ii.	448	Ranunculus acris, ii.	480	Resina flava, ii.	673
Purgative Lefroy's, i.	551	Ranunculus aquatilis, ii.	480	Resina galbani, i.	682
Purple of cassius, i.	314	Ranunculus bulbosus, ii.	480	Resina gialappæ, i.	548
Purslain, ii.	410	Ranunculus ficaria, ii.	481	Resina guaiaci, i.	733
Pyrethri officinalis radix, i.	230	Ranunculus flammula, ii.	481	Resina guaiaci nativa, i.	732
Pyrethrum, i.	230	Ranunculus sceleratus, ii.	480	Resina hederæ, i.	757
Pyri martiales, i.	667	Rape, i.	342	Resina jalapæ, i.	548
Pyrites arsenicalis micaceus, i.	282	Raphanus rusticus, ii.	481	Resina juniperi, ii.	555
Pyrites ferri artificialis, i.	639	Raphanus sativus, ii.	481	Resina lentisci, ii.	223
Pyrola rotundifolia, ii.	471	Rapi rubri radix, i.	336	Resina mastichis, ii.	223
Pyroleum ossium rectificatum, i.	707	Raspatura cornu cervi, i.	359	Resina meccanense fluida, i.	319
Pyroleum succini, ii.	630	Raspatura ligni guaiaci, i.	732	Resinæ oleæ, i.	764
Quadrichloruret of sodium, ii.	602	Raspberry, ii.	532	Resina liquida pini bal-samei, i.	318
Quassia, ii.	472	Rassura cornu cervi, i.	359	Resina pini empyreumatica liquida, ii.	364
		Rasura ligni guaiaci, i.	732	Resina liquida pini lenticis, ii.	654
		Ratafia antiscorbutic, i.	439	Resina sanguis draconis, i.	764
		Ratany, ii.	147		
		Rattlesnake root, ii.	403		
		Realgar, i.	282		
		Recrementa vitri, ii.	430		
		Reed corn flag, i.	720		
		Reed-grass, i.	293		
		Reed-grass mannard, i.	293		
		Regina pratensis radix, ii.	621		
		Reguli usti, ii.	391		
		Regulus antimonii, i.	233		

Resina scammonii, ii.	560	Rob corticis nucum jug-	Rose mineral, ii.	82
Resina succini, ii.	630	landis regiæ, ii.	Rose hundred leaved, ii.	509
Resina succini balsamica,		Rob of cranberries, ii.	Rose musk ii.	509
ii.	631	Rob of red currants, ii.	Rose Provence, ii.	509
Resolvent Richard's, ii.	670	Rob cydoniorum, i.	Rose red, ii.	509
Rhabbarbari monachorum		Rob cynosbatos, ii.	Rose white, ii.	509
radix, ii.	535	Rob dauci, i.	Rosebay common, ii.	260
Rhabbarbari nigri radix,		Robdauci carottæ, i.	Rosemary, ii.	521
i.	543	Rob diacarii Galeni, ii.	Rosewort, ii.	580
Rhabbarbarine, ii.	489	Rob of dwarf elder, ii.	Rosmarinus officinalis,	
Rhabbarbarum, ii.	489	Rob of elder, ii.	ii.	521
Rhabbarbarum tostum, ii.	490	Rob of fragariæ, i.	Rossolis of cinchona, i.	501
Rhamnus cathartica, ii.	487	Rob of juniper, ii.	Rotulæ anthelminticæ, ii.	65
Rhamnus frangula, ii.	487	Rob juniperi, ii.	Rotulæ berberum, i.	335
Rhamnus jujuba, ii.	150	Rob of lemon, i.	Rotulæ diaireos, ii.	132
Rhapontici radix, ii.	534	Rob mororum, ii.	Rotulæ manus christi,	
Rhaponticine, ii.	490	Rob of the fruit of moun-	ii.	540
Rhatania, ii.	147	tain ash, ii.	Rotulæ menthæ piperitæ,	
Rheum, ii.	489	Rob of mulberries, ii.	ii.	237
Rheumine, ii.	489	Rob myrtillorum, ii.	Rotulæ pectorales, ii.	635
Rhinoceros cornu, i.	359, 699	Rob nucum, ii.	Rotulæ contra vermes, ii.	29
Rhinoceros indicus, i.	359	Rob oxycoccos, ii.	Rotuli citri, i.	520
Rhodiæ radix, ii.	580	Rob prunorum sylves-	Rubetæ terrestres majores,	
Rhododaphnis folio, ii.	260	trium, ii.	i.	698
Rhododendron yellow-		Rob of quince, i.	Rubia manjista, ii.	529
flowered, ii.	502	Rob of raspberries, ii.	Rubigo, i.	646
Rhododendrum chrysan-		Rob ribesiorum nigrorum,	Rubigo ferri, i.	646
thum, ii.	502	ii.	Rubinus, i.	155
Rhodomel, ii.	515	Rob rubi idæi, ii.	Rubrica fabrilis, i.	155
Rhodosaccharum, ii.	513	Rob of sloes, ii.	Rubus arcticus, ii.	531
Rhubarb, ii.	489	Rob spinæ cervinæ, ii.	Rubus chamæmorus, ii.	531
Rhubarb bastard, ii.	535	Rob of strawberries, i.	Rubus fruticosus, ii.	531
Rhubarb monk's, ii.	535	Rob vaccinii myrtilli, ii.	Rubus idæus, ii.	532
Rhubarb torrified, ii.	490	Rob vaccinii oxycocci, ii.	Rubus trivialis, ii.	532
Rhus copallinus, i.	755	Rob vitis idææ, ii.	Rubus villosus, ii.	532
Rhus coriaria, ii.	503	Rob of walnuts, ii.	Ruby, i.	155
Rhus cotinus, ii.	503	Robinia pseudo-acacia,	Rue goat's, i.	691
Rhus toxicodendrum et		ii.	Rumex acetosa, ii.	533
radicans, i.	503	Roambole, i.	Rumex alpina, ii.	535
Ribes grossularia et uva		Roccella tinctoria, ii.	Rumex aquatica, ii.	534
crispa, ii.	506	Rocella, ii.	Rumex crispa, ii.	534
Ribes nigrum, ii.	504	Rocket, i.	Rumex sanguinea, ii.	535
Ribes rubrum, ii.	79	Roob ebuli, ii.	Rumex scutata, ii.	534
Ribwort, ii.	366	Roob ribium rubrorum,	Ruperti herba, i.	718
Rice, ii.	325	ii.	Ruptorium commune, ii.	412
Richardsonia Brasiliensis,		Roob sambuci, ii.	Rupture wort, ii.	5
ii.	124	Roob sorbi ancupariæ, ii.	Ruscus aculeatus, ii.	535
Richweed, i.	89	Root yellow, ii.	Ruscus hypoglossum, ii.	536
Ricini majoris semines, ii.	134	Rorillæ herba, i.	Rust of iron, i.	646
Ricinus, ii.	507	Roris solis herba, i.	Ruta graveolens, ii.	536
Ricinus communis, ii.	507	Rorismarini sylvestris	Rutæ capariæ herba, i.	691
Rob fructuum acaciæ,		herba, ii.	Rutæ murariæ herba, i.	299
Germanicæ, ii.	468	Ros sulphuris, i.	Rye, ii.	579
Rob antisypilitic, ii.	597	Ros vitrioli, i.	Rye spurred, ii.	579
Rob baccharum juniperi,		Rosa alba, ii.		
ii.	138	Rosa canina, ii.	Sabadilla, ii.	713
Rob of barberries, i.	335	Rosa centifolia, ii.	Sabadilline, ii.	714
Rob berberum, i.	335	Rosa Gallica, ii.	Sabinæ herba, ii.	142
Rob of red bilberries,		Rosa moschata, ii.	Sacchar, ii.	539
ii.	703	Rosæ benedictæ, ii.	Saccharum, ii.	539
Rob of black currants, ii.	505	Rosæ regię, ii.	Saccharum clarificatum,	
Rob of blueberries, ii.	702	Rosaginis folia, ii.	ii.	539
Rob of buckthorn, ii.	487	Rosariæ radix, ii.	Saccharum glycyrrhisa-	
Rob of cherries, i.	438	Rose damask, ii.	tum, ii.	724
Rob citri medicæ, i.	520	Rose dog, ii.	Saccharum hordeatum, ii.	7

Saccharum lactis, ii.	6	Sal cornu cervi succina-		Salab, ii.	540
Saccharum mercuriale		tus, i.	200	Salep, ii.	540
compositum, ii.	26	Sal cornu cervi volatile, i.	184	Salicaria radix, ii.	205
Saccharum mercurii ver-		Sal culinaris, ii.	602	Salicine, ii.	542
nifugum, ii.	16	Sal digestivus, ii.	440	Salivalis radix, i.	230
Saccharum myrrhæ, i.	759	Sal diureticum vegetabile,		Salix alba, ii.	542
Saccharum myrrhatum, i.	759	ii.	447	Salix capræa, ii.	542
Saccharum perlatum, ii.	511,	Sal diureticus, ii.	441	Salix fragilis, ii.	542
	540	Sal de duobus, ii.	427	Salix pentandra, ii.	542
Saccharum rosatum, ii.	511	Sal Ebsharnense, ii.	210	Salix vitellina, ii.	542
Saccharum saturni, ii.	382	Sal Epsoniense, ii.	210	Salsaparilla germanicæ	
Sacco-gummit, i.	722	Sal essentielle, i.	46	radix, i.	396
Safflower, i.	398	Sal febrifugus Sylvii, ii.	440	Salsefi, ii.	695
Saffron, i.	563	Sal gemme, ii.	602	Salsilago ammoniacalis	
Saffron astringent of		Salvia horminum, ii.	544	i.	195
mars, i.	643	Sal marinus, ii.	602	Salsilago ammoniacalis	
Saffron of mars Stahl's		Sal margaritarum, i.	363	acetosa, i.	195
aperitive antimoniated,		Sal matris perlarum, i.	363	Salsilago cornu cervi suc-	
i.	645	Sal martis, i.	651	cinatus, i.	200
Saffron of mars by sul-		Sal martis muriaticum		Salsilago succinatus am-	
phur, i.	644	ammoniacale, i.	661	moniaci, i.	200
Saffron meadow, i.	538	Sal martis muriaticus, i.	656	Salsilago succinatus am-	
Sagapenum, i.	764	Sal mercuriatus acetosus,		moniac, i.	200
Sage common, ii.	544	ii.	95	Salsola, ii.	604
Sage purple-topped, ii.	544	Sal mercurii acetosus, ii.	95	Salsola fragus, ii.	543
Sal absinthii citratum, ii.	446	Sal mirabile Glauberi, ii.	611	Salsola kali, ii.	544
Sal acetosellæ, ii.	446	Sal muriaticum baroticum,		Salsola soda, ii.	543
Sal acetosus ammoniaca-		i.	325	Salt, ii.	602
lis, i.	195	Sal oculorum cancrorum,		Salt essential of cinchona,	
Sal acidum, i.	46	i.	363	i.	482
Sal alembroth, ii.	89	Sal panchrestum, ii.	447	Salt epsom, ii.	210
Sal amarum, ii.	210	Sal polychrest soluble, ii.	455	Salt Lagresie's ferrugi-	
Sal ammoniacum, i.	191	Sal polychrestum glaseri,		nous, i.	652
Sal ammoniacum depu-		ii.	427	Salt of glass, ii.	430
ratum, i.	191	Sal polychrestus Seignetti,		Salt Glauber's, ii.	611
Sal ammoniacum fixum,		ii.	455	Salt Guindre's, ii.	613
i.	360	Sal prunella, ii.	433	Salt of hartshorn, i.	184
Sal ammoniacum mar-		Sal prunellæ, ii.	433	Salt Lagaraye's, i.	482
tile, i.	661	Sal rupellensis, ii.	455	Salt of lead, ii.	383
Sal ammoniacum nitro-		Sal sapientiæ, ii.	89, 428	Salt Tachenius' lixivial,	
sum, i.	190	Sal sardlitzense, ii.	210	ii.	421
Sal ammoniacum secre-		Sal scientiæ, ii.	89	Salt essential of opium,	
tum Glauberi, i.	190	Sal sedativum mercuriale,		ii.	290
Sal ammoniacum tarta-		ii.	81	Salt petre, ii.	432
reum, ii.	456	Sal sedativus Hombergii,		Salt purging, i.	42
Sal volatile oleosum an-		i.	48	Salt Rochelle, ii.	455
gelicæ Boerhaavii, i.	223	Sal sedlitzense, ii.	210	Salt essential of rhubarb,	
Sal anglicanum cathar-		Sal scydschutzensense, ii.	210	ii.	494
ticum, ii.	210	Sal succini, i.	75	Salt sedative, ii.	608
Sal armoniacum, i.	191	Sal volatile succini, i.	75	Salt essential of senna,	
Sal astracanensis artifi-		Sal tachenianus, ii.	421	i.	488
cialis, ii.	613	Sal tartari, ii.	420	Salt of sorrel, ii.	446
Sal volatile benzoës, i.	46	Sal tartari essentielle, i.	80	Salt of tartar, ii.	420
Sal acidum boracis, i.	48	Sal tartari per se liquidus,		Salt essential of tartar,	
Sal volatile boracis, i.	48	ii.	422	ii.	449
Sal catharticum, ii.	64	Sal tartari sulphuratum,		Salt English volatile, i.	183
Sal catharticum amarum,		ii.	428	Salt of vinegar, i.	42
ii.	210	Sal vegetabile, ii.	447	Salt aromatic volatile, i.	186
Sal chalybis, i.	651	Sal vitæ, ii.	89	Salt of urine, ii.	429
Sal cinerum clavellato-		Sal vitri, ii.	430	Salt Westendorf's, i.	42
rum, ii.	419	Sal volatile oleosum sylvii,		Salt of wormwood, ii.	421
Sal corallinorum, i.	363	i.	174	Saltwort common, ii.	543
Sal communis, ii.	602	Sal vitrioli narcoticum, i.	48	Salvia officinalis, ii.	544
Sal cornu cervi depura-		Sal urinosus purissimus,		Salvia pratensis, ii.	544
tum, i.	184	i.	182	Salvia sclarea, ii.	544

<i>Salviæ sylvestris herba</i> , ii.	683	<i>Sapohydrosulphureti oxy-</i> <i>duli stibii sulphurati</i> , i.	251	<i>Scilla maritima</i> , ii.	563
<i>Sambucus Canadensis</i> , ii.	551	<i>Sapo jalapæ</i> , i.	549	<i>Scillitinum</i> , ii.	563
<i>Sambucus ebulus</i> , ii.	550	<i>Sapo jalapinus</i> , i.	549	<i>Scincus marinus</i> , i.	698
<i>Sambucus nigra</i> , ii.	551	<i>Sapo kalicus</i> , ii.	457	<i>Scincus maximus</i> , i.	698
<i>Samphire</i> , i.	562	<i>Sapo medicatus</i> , ii.	457	<i>Scincus officinalis</i> , i.	698
<i>Sampsuchi herba</i> , ii.	322	<i>Sapo medicinalis</i> , ii.	457	<i>Sclæræ herba</i> , ii.	544
<i>Sanamundæ herba</i> , i.	719	<i>Sapo mollis</i> , ii.	457	<i>Scleroderma cervinum</i> , ii.	577
<i>Sandarac</i> , ii.	555	<i>Sapo niger</i> , ii.	457	<i>Sclerotium clarum</i> , ii.	579
<i>Sandaraca</i> , i. 282, ii.	555	<i>Sapo potassæ terebinthi-</i> <i>naceus</i> , ii.	671	<i>Scobs ferri</i> , i.	634
<i>Sanders red</i> , ii.	195	<i>Sapo Starkeyanus</i> , ii.	671	<i>Scolopendrii veri herba</i> , i.	443
<i>Sanders white</i> , ii.	196	<i>Sapo stibiatus</i> , i.	251	<i>Scolopendrium officinale</i> , ii.	577
<i>Sanders yellow</i> , ii.	196	<i>Sapo sulphurato-croca-</i> <i>tus</i> , ii.	417	<i>Scordii herba</i> , ii.	683
<i>Sanguinalis herba</i> , ii.	406	<i>Sapo tartareus</i> , ii.	671	<i>Scordium</i> , ii.	682
<i>Sanguinaria Candensis</i> , ii.	555	<i>Sapo tartareus Starkeyi</i> , ii.	671	<i>Scorodonæ herba</i> , ii.	683
<i>Sanguinariæ herba</i> , ii.	406, 688	<i>Sapo terebinthinatus</i> , ii.	671	<i>Scorpio Europæus</i> , i.	698
<i>Sanguinariæ minoris her-</i> <i>ba</i> , ii.	342	<i>Sapo Venetus</i> , ii.	456	<i>Scorpion European</i> , i.	698
<i>Sanguinariæ radix</i> , i.	718	<i>Sapo viridis</i> , ii.	457	<i>Scorzoneræ garden</i> , ii.	577
<i>Sanguis capræ ibicis</i> , i.	699	<i>Sapo vomitoria Sylvii</i> , i.	259	<i>Scorzoneræ Hispanica</i> , ii.	577
<i>Sanguis draconis</i> , i.	764	<i>Sapo Wologdensis</i> , ii.	457	<i>Scorzoneræ humilis</i> , ii.	577
<i>Sanguis hirci</i> , i.	699	<i>Saponaria officinalis</i> , ii.	556	<i>Scrofularia</i> , ii.	578
<i>Sanguisorba officinalis</i> , ii.	555	<i>Sapota dissecta</i> , i.	470	<i>Scrophularia aquatica</i> , ii.	578
<i>Sanicle</i> , ii.	556	<i>Sapota salicifolia</i> , i.	470	<i>Scrophularia nodosa</i> , ii.	578
<i>Sanicle American</i> , ii.	4	<i>Sapphirus</i> , i.	155	<i>Scurvy grass</i> , i.	531
<i>Sanicula Europæa</i> , ii.	556	<i>Saracen's wound-wort</i> , ii.	618	<i>Scurvy grass wild</i> , i.	560
<i>Saniculæ albæ radix</i> , i.	609	<i>Sarcocolla</i> , i.	765	<i>Scutum stomachicum</i> , ii.	315
<i>Saniculi radix</i> , i.	86	<i>Sarcocollina</i> , i.	765	<i>Scutum pro umbilico</i> , i.	689
<i>Santalina</i> , ii.	196	<i>Sarsa</i> , ii.	593	<i>Scyphophorus cocciferus</i> , ii.	186
<i>Santalina chamæcy pa-</i> <i>rissias</i> , ii.	556	<i>Sarsaparilla</i> , ii.	593	<i>Scyphophorus pyxidatus</i> , ii.	187
<i>Santalum album</i> , ii.	196	<i>Sassafras</i> , ii.	183	<i>Sea-bindweed</i> , i.	366
<i>Santolinæ herba</i> , ii.	556	<i>Satureja capitata</i> , ii.	558	<i>Sea-oak</i> , i.	678
<i>Santonici semen</i> , ii.	581	<i>Satureja hortensis</i> , ii.	558	<i>Seabaceum palmæ</i> , ii.	271
<i>Saphire</i> , i.	155	<i>Satureja thymbra</i> , ii.	558	<i>Sebestenæ</i> , i.	558
<i>Sapo albus</i> , ii.	456	<i>Saturnus</i> , ii.	368	<i>Sebestes</i> , i.	558
<i>Sapo alcoholico-ammo-</i> <i>niatus</i> , i.	179	<i>Sauce-alone</i> , i.	623	<i>Secale cereale</i> , ii.	579
<i>Sapo alcoholico-campho-</i> <i>ratus</i> , i.	179	<i>Savory cretan</i> , ii.	558	<i>Secale cornutum</i> , ii.	579
<i>Sapo Alicantinus</i> , ii.	456	<i>Savory garden</i> , ii.	558	<i>Sedi majoris herba</i> , ii.	584
<i>Sapo Aloniensis</i> , ii.	456	<i>Saxifraga granulata</i> , ii.	558	<i>Sedum acre</i> , ii.	580
<i>Sapo ammoniaco-cam-</i> <i>phoratus</i> , i.	179	<i>Saxifragæ anglicæ radix</i> , ii.	343	<i>Sedum Rhodiola</i> , ii.	580
<i>Sapo ammoniæ</i> , i.	177	<i>Saxifragæ rubræ radix</i> , ii.	621	<i>Sedum telephium</i> , ii.	580
<i>Sapo amygdalinus</i> , ii.	457	<i>Saxifrage burnet</i> , ii.	356	<i>Seeds major cold</i> , i.	578
<i>Sapo antimonialis</i> , i.	251	<i>Saxifrage common</i> , ii.	558	<i>Serum lactis mannatum</i> , ii.	220
<i>Sapo antimonii</i> , i.	251	<i>Saxifrage pimpinel</i> , ii.	356	<i>Segestria cellaria</i> , i.	271
<i>Sapo arthriticus Pottii</i> , ii.	670	<i>Scabiosa arvensis</i> , ii.	559	<i>Self-heal</i> , ii.	465
<i>Sapo aromaticus</i> , i.	179	<i>Scabiosa succisa</i> , ii.	559	<i>Selinum cervaria</i> , ii.	580
<i>Sapo cocautinus</i> , ii.	457	<i>Scabiosa field</i> , ii.	559	<i>Selinum oreoselinum</i> , ii.	581
<i>Sapo chemicus pro casu</i> <i>externo</i> , ii.	671	<i>Scammonea</i> , ii.	559	<i>Selinum palustre</i> , ii.	581
<i>Sapo domesticus</i> , ii.	456	<i>Scammonium</i> , ii.	559	<i>Semen cinæ</i> , ii.	581
<i>Sapo guaiacinus</i> , i.	740	<i>Scammony</i> , ii.	559	<i>Semen contra</i> , ii.	581
<i>Sapo gummi ammoniaci</i> , i.	749	<i>Scammony glycyrrhized</i> , ii.	560	<i>Semen lumbricorum</i> , ii.	581
<i>Sapo Hispanicus albus</i> , ii.	456	<i>Scammony sulphuretted</i> , ii.	560	<i>Semines lycopodii</i> , ii.	204
<i>Sapo hydrargyri</i> , ii.	98	<i>Scarabeus unctuarii</i> , ii.	232	<i>Semen sanctum</i> , ii.	581
		<i>Scariolæ herba</i> , i.	461	<i>Semen sinæ</i> , ii.	581
		<i>Scarlet mushroom</i> , i.	601	<i>Sempervivum tectorum</i> , ii.	584
		<i>Schenanthi herba</i> , i.	217	<i>Semen contra vermes</i> , ii.	581
				<i>Semen zedoariæ</i> , ii.	581

Sementina, ii.	581	Serum lactis tartarisatum, ii.	585	Sison amomum, ii.	591
Senecio Jacobæa, ii.	584	Serum lactis temperans, ii.	647	Sisymbrium sophia, ii.	591
Senecio vulgaris, ii.	584	Serum lactis vinosum, ii.	585	Sium ammi, ii.	592
Senecionis cærulei herba, i.	621	Sesame, ii.	586	Sium angustifolium, ii.	591
Senegæ radix, ii.	403	Sesamum orientale, ii.	586	Sium latifolium, ii.	591
Senegine, ii.	403	Seseli Cretici radix, ii.	692	Sium ninsi, ii.	592
Seneka, ii.	403	Seseli tortuosum, ii.	586	Sium nodiflorum, ii.	591
Senekæ radix, ii.	401	Seslios pratensis radix, ii.	343	Sium sisarum, ii.	591
Senna, i.	401	Sevum, i.	315	Skins of snakes, i.	678
Senna of Aleppo, i.	401	Sevum bovillum, i.	315	Skull human, i.	359
Senna Alexandrian, i.	420	Sevum bubalum, i.	315	Skunk-cabbage, i.	619
Senna American, i.	401	Sevum cervinum, i.	315	Slater, ii.	272
Senna of Barbary, i.	401	Sevum hircinum, i.	316	Sloe, ii.	467
Senna Bicharian, i.	540	Sevum ovillus, i.	316	Smallage, i.	266
Senna bladder, i.	401	Sevum vervecinum, i.	316	Smallage stalks preserved, i.	266
Senna East Indian, i.	401	Shalot, i.	112	Smalt i.	526
Senna Egyptian, i.	401	Shells egg, i.	353	Smaltum, i.	526
Senna Italian, i.	401	Shells oyster, i.	353	Smaragdus, i.	157
Sennæ folii, i.	402	Shells snail, i.	353	Smilax china, ii.	592
Sennæ follicula, i.	401	Shepherd's purse, ii.	688	Smilax sarsaparilla, ii.	593
Senna Nubian, i.	401	Shield-fern male, i.	308	Smiris, i.	155
Senna of Sayd, i.	401	Sicilianæ herba, ii.	107	Smyris, i.	155
Senna Spanish, i.	692	Sideritis hirsutæ herba, ii.	622	Snail orchard, i.	698
Septifolii radix, ii.	585	Sief album, ii.	372	Snake-root, ii.	620
Serici folliculi, ii.	706	Sief cum opio, ii.	372	Snake-root Canadian, i.	293
Sericum adhæsivum, i.	585	Sigesbechia orientalis, ii.	621	Snake-root Indian, ii.	274
Sericum crudum, ii.	391	Sigilli Salomonis radix, ii.	405	Snake-root Virginian, i.	277
Sericum tostum, i.	277	Silene behen, ii.	587	Snakeweed, ii.	406
Serpentaria, i.	291	Silene otites, ii.	587	Snakewood, ii.	626
Serpentariæ minoris radix, i.	406	Silene Virginica, ii.	587	Snap dragon, i.	266
Serpentariæ rubræ radix, ii.	277	Sileris montani semines, ii.	155	Sneezewort, i.	37
Serpentariæ Virginicæ radix, i.	406	Silicate of alumina and of iron, i.	157	Soap almond, ii.	457
Serpentariæ vulgaris radix, ii.	274	Silicate of alumina and of soda, i.	157	Soap aromatic animal, ii.	457
Serpentum radix, ii.	690	Silicate of magnesium, ii.	213	Soap antimonial, i.	251
Serpylli herba, ii.	619	Siliqua dulcis, i.	440	Soap Singer's antisialagogue, ii.	417
Service tree, ii.	149	Siliqua hirsuta, ii.	626	Soap aromatic, ii.	459
Serum, ii.	585	Silk Guilbert's epispastic, i.	378	Soap beef-marrow, ii.	457
Serum lactis, ii.	586	Silk raw, ii.	585	Soap Dippel's chemical, ii.	416
Serum lactis acetosum, ii.	585	Silk vesicatory, i.	295	Soap cocoa, ii.	457
Serum lactis acidum, ii.	165	Silk-weed, i.	273	Soap common, ii.	456
Serum lactis alumina-tum, i.	165	Silver, i.	273	Soap of croton oil, i.	570
Serum lactis aluminosum, i.	165	Silver assayed, i.	273	Soap gamboge, i.	696
Serum lactis aluminosum cum cinnamomo, i.	533	Silver pure, i.	465	Soap guaiac, i.	737
Serum lactis antiscorbuticum, i.	585	Silverweed, ii.	697	Soap guaiacine, i.	740
Serum lactis chalybeatum, ii.	585	Silurus glanis, i.	587	Soap of gum ammoniac, i.	749
Serum lactis compositum, ii.	586	Simarouba amara, ii.	588	Soap jalapine, i.	549
Serum lactis dulci Hoffmanni, ii.	585	Sinapis nigra, ii.	588	Soap ley, ii.	410
Serum lactis sincerum, ii.	590	Sinapism, ii.	588	Soap medicinal, ii.	457
Serum lactis sinapinum, ii.	647	Sinapismus, ii.	755	Soap Chaussier's mercurial, ii.	98
Serum lactis tamarindatum, ii.		Siphonia cahuchu, i.	562	Soap sulphuretted of soda, ii.	602
		Sirop anti-arthritisque, ii.	562	Soap soft, ii.	457
		Sirop antihydrique, ii.	209	Soap Spanish, ii.	456
		Sirop d'orgeat, i.		Soap Starkey's, ii.	671
				Soap vegetable, ii.	422
				Soap Venice, ii.	456
				Soap of wax, ii.	601
				Soap Wologd, ii.	457
				Soap-wort, ii.	556
				Soda, ii.	604

Soda acētata, ii.	614	Solutio arsenitis kalicæ,		Solutio nitratis hydrar-	
Soda acetosa, ii.	614	i.	286	gyri ammoniacalis, ii.	89
Soda caustic, ii.	601	Solutio balsamica alkali-		Solutio nitratis sodæ,	
Soda caustica, ii.	601	sata, ii.	606	ii.	614
Soda ley, ii.	601	Solutio barytæ muriati-		Solutio opii alcoholi-	
Soda phosphorata, ii.	610	cæ, i.	326	sata, ii.	282
Soda pura, ii.	601	Solutio calcaris sulphura-		Solutio oxymuriatis cal-	
Soda tartarisata, ii.	455	to-stibiata, i.	254	cici, i.	351
Soda tartarized, ii.	455	Solutio calcis, i.	347	Solutio plumbi in acido	
Soda vitriolata, ii.	611	Solutio camphoræ æthe-		acetoso, ii.	375
Soda water, ii.	608	rea, i.	371	Solutio rob juniperi, ii.	138
Sodæ acetas, ii.	614	Solutio camphoræ aquo-		Solutio styptica Loeffii,	
Sodæ bicarbonas, ii.	607	sa, i.	368	i.	657
Sodæ carbonas, ii.	604	Solutio camphoræ oleosa,		Solutio subacetatis plum-	
Sodæ carbonas siccatum,		i.	374	bici, ii.	378
ii.	604	Solutio catheretica, i.	282	Solutio subcarbonatis	
Sodæ carbonatis aqua,		Solutio cerata, i.	433	ammonia, i.	184
ii.	605	Solutio citratis kalici,		Solutio subcarbonatis	
Sodæ murias, ii.	602	ii.	446	ammoniaci spirituosas,	
Sodæ phosphas, ii.	610	Solutio citro-muriatica,		i.	172
Sodæ et potassæ tartras,		ii.	603	Solutio sulphatis cupri	
ii.	455	Solutio cupri ammonia-		composita, i.	586
Sodæ sub-boras, ii.	608	calis, i.	596	Solutio tartari depurati,	
Sodæ subcarbonas, ii.	604	Solutio elaterii, ii.	244	ii.	452
Sodæ subcarbonas ex-		Solutio ferrata, i.	673	Solutio tartari solubilis,	
siccata, ii.	604	Solutio ferri muriatici		ii.	448
Sodæ sulphas, ii.	611	cum baryta muriatica,		Solution of acetate of	
Sodium, ii.	601	i.	327	ammonia, i.	195
Solamen hypochondria-		Solutio fuliginis fœtida,		Solution aqueous of ace-	
corum, ii.	448	i.	306	tate of lead, ii.	386
Solamen scabiosorum, i.	679	Solutio gummi guttæ		Solution opiate aqueous	
Solani furiosi radix, i.	309	alkalina, i.	695	of acetate of lead, ii.	386
Solani lethalis radix, i.	309	Solutio hydrargyrata,		Solution of acetate of	
Solani maniaci herba,		ii.	77	mercury, ii.	96
i.	605	Solutio hydrargyri gum-		Solution alcoholic of ace-	
Solani quadrifolii radix,		mosi, ii.	15	tate of potass, ii.	444
ii.	342	Solutio mercurialis, ii.	90	Solution of alum, i.	162
Solani racemosi radix,		Solutio mercurii corro-		Solution compound of	
ii.	351	sovi, ii.	70	alum, i.	162
Solani scandentis stipites,		Solutio mineralis Fowl-		Solution of ammonia,	
ii.	615	eri, i.	287	i.	168
Solani vescarii baccæ,		Solutio mucilaginosas,		Solution diluted of am-	
ii.	350	i.	751	monia, i.	171
Solanine, ii.	615, 617	Solutio muriatis barytæ,		Solution antiscorbutic,	
Solanum dulcamara, ii.	615	i.	326	i.	327
Solanum lycopersicum,		Solutio muriatis calcis,		Solution antiscrofulous,	
ii.	617	i.	363	i.	327
Solanum melongena, ii.	617	Solutio muriatis calcis		Solution Niemann's anti-	
Solanum nigrum, ii.	617	hydrargyrata, ii.	35	scrofulous, i.	363
Solanum tuberosum, ii.	618	Solutio muriatis ferri,		Solution antistrumous,	
Soldanellæ herba, i.	366	i.	658	ii.	123
Solidaginis Saracenicae		Solutio muriatis ferri		Solution of arseniate of	
herba, ii.	618	composita, i.	657	ammonia, i.	288
Solidago virga aurea, ii.	618	Solutio muriatis ferrici		Solution bitter arsenical,	
Solomon's seal, ii.	405	ætherea, i.	659	i.	287
Solutio ætherea, i.	100	Solutio muriatis ferrici		Solution Brera's arseni-	
Solutio alexiteria, i.	55	spirituosa, i.	658	cal, i.	286
Solutio alexiteria Gau-		Solutio muriatis hydrar-		Solution Fowler's arse-	
biana, i.	57	gyri oxygenati, ii.	70	nical, i.	287
Solutio alexiteria oxy-		Solutio muriatis potassæ		Solution Hein's arseni-	
genata, i.	459	oxygenati, ii.	440	cal, i.	286
Solutio arsenicalis, i.	287	Solutio myrrhæ alkali-		Solution Jacob's arseni-	
Solutio arsenicalis Borus-		sata, i.	762, 763	cal, i.	286
sica, i.	287	Solutio alcoholica myrr-		Solution of arsenious	
Solutio arsenicata, i.	286	hæ, i.	762	acid, i.	284

- | | | | | | |
|---|-----|--|----------|--|----------|
| Solution compound of arsenite of soda, i. | 287 | Solution Dellalena's lithontriptic, ii. | 444 | Solution of subsulphate of copper and of ammonia, i. | 587 |
| Solution alkaline balsamic, ii. | 606 | Solution of liver of sulphur, ii. | 418 | Solution aqueous of sulphate of copper, i. | 584 |
| Solution of baryta, i. | 328 | Solution of malate of iron, i. | 665 | Solution aromatic of sulphate of copper, i. | 585 |
| Solution barytic, i. | 327 | Solution Marryat's, ii. | 71 | Solution camphorated of sulphate of copper, i. | 584 |
| Solution of belladonna, i. | 310 | Solution of muriate of iron, i. | 657 | Solution compound of sulphate of copper, i. | 586 |
| Solution of artificial camphor, ii. | 670 | Solution of muriate of lime, i. | 363 | Solution carbonic of sulphate of magnesia, ii. | 212 |
| Solution cathartic, i. | 585 | Solution of myrrh, i. | 762 | Solution aqueous of sulphate of potass, ii. | 430 |
| Solution of caustic potass, ii. | 410 | Solution alkaline of myrrh, i. | 762 | Solution of sulphate of silver, i. | 276 |
| Solution of chlorate of potass, ii. | 440 | Solution of nitrate of silver, i. | 274 | Solution of sulphuret of antimony and of lime, i. | 254 |
| Solution of chlorine, i. | 459 | Solution of nitrate of soda, ii. | 614 | Solution alcoholic of sulphuret of potass, ii. | 419 |
| Solution extemporaneous of chlorine, i. | 460 | Solution nitrous, i. | 70 | Solution of sulphuret of potassium, ii. | 417 |
| Solution Cohen's, ii. | 414 | Solution ophthalmic, i. | 586 | Solution of supercarbonate of potass, ii. | 426 |
| Solution æthereal of deutochloruret of mercury, ii. | 72 | Solution of potass, ii. | 410 | Solution of supertartrate of potass, ii. | 452 |
| Solution alcoholic of deutochloruret of mercury, ii. | 70 | Solution alcohol of potass, ii. | 414 | Solution of tartarized antimony, i. | 258 |
| Solution opiate alcoholic of deutochloruret of mercury, ii. | 76 | Solution prophylactic against scarlatina, i. | 311 | Solution of tartrate of ammonia, i. | 202 |
| Solution aqueous of deutochloruret of mercury, ii. | 68 | Solution Niemann's solvent, i. | 193 | Solution alcoholic of tartrate of potass, ii. | 449 |
| Solution æthereal of deuto-ioduret of mercury, ii. | 79 | Solution Vogler's solvent, i. | 193 | Solution aqueous of tartrate of potass, ii. | 448 |
| Solution alcoholic of deuto-ioduret of mercury, ii. | 79 | Solution of rob of juniper, ii. | 138 | Solution Tracher's, i. | 186 |
| Solution epispastic, i. | 259 | Solution salino-gelatinous, i. | 703 | Solution of veratrine, ii. | 713 |
| Solution evaporatory, i. | 102 | Solution Saviard's, ii. | 414 | Solution Weikard's, ii. | 92 |
| Solution of extract of nux vomica, ii. | 267 | Solution sedative, ii. | 283, 293 | Sonchus oleraceus, ii. | 619 |
| Solution Fallopius', i. | 162 | Solution of soda, ii. | 601 | Soot, i. | 679 |
| Solution of ferrocyanate of potass, i. | 61 | Solution æthereal of soap, ii. | 462 | Sophiæ chirurgorum herba, ii. | 591 |
| Solution gelatinous, i. | 701 | Solution alcoholic of soap, ii. | 460 | Sorb apple, ii. | 619 |
| Solution saline gelatinous, ii. | 613 | Solution aromatic and camphorated of soap, ii. | 461 | Sorb tree, ii. | 619 |
| Solution of gum arabic, i. | 751 | Solution camphorated alcoholic of soap, ii. | 460 | Sorbus aucuparia, ii. | 619 |
| Solution Heineke's, i. | 238 | Solution camphorated opiate alcoholic of soap, ii. | 461 | Sorbus domestica, ii. | 619 |
| Solution of hydriodate of potass, ii. | 122 | Solution of antimonial soap, i. | 252 | Sorrel common, ii. | 533 |
| Solution of ioduretted hydriodate of potass, ii. | 123 | Solution aromatic alcoholic of soap, ii. | 460 | Sorrel small, ii. | 534 |
| Solution of hydrochlorate of ammonia and of iron, i. | 663 | Solution of subacetate of lead, ii. | 375 | Sorrel wood, ii. | 327 |
| Solution alkaline of iron, i. | 647 | Solution diluted of subacetate of lead, ii. | 378 | Sorrel yellow wood, ii. | 327 |
| Solution of lime, i. | 347 | Solution of subcarbonate of ammonia, i. | 183 | Sotirella parva, ii. | 300 |
| Solution oxygenated of lime, i. | 351 | Solution alcoholic of subcarbonate of ammonia, i. | 187 | Soup portable, i. | 705 |
| Solution lithontriptic, ii. | 422 | Solution aqueous of subcarbonate of potass, ii. | 422 | Southernwood garden, i. | 290 |
| | | Solution of subcarbonate of soda, ii. | 605 | Southernwood wild, ii. | 290 |
| | | | | Sparadrap adhesive, ii. | 675 |
| | | | | Sparadrap aromatic, ii. | 256 |
| | | | | Sparadrap common, i. | 437, 657 |
| | | | | Sparadrap desiccative, ii. | 374 |
| | | | | Sparadrap epulotic, ii. | 391 |
| | | | | Sparadrap Gauthier's, ii. | 393 |
| | | | | Sparadrap Goulard's, ii. | 382 |

Sparadrap green, i.	593	Species pro cataplasmate,		Species diatron santalon,	
Sparadrap red, ii.	45, 371	i.	143	ii.	197
Sparadrap resinous, ii.	677	Species cephalic, i.	191, ii.	Species diaturbith with	
Sparadrap ad corpus con-		132, 156, 234		rhubarb, ii.	490
ditum involvendum, ii.	256	Species cephalicæ, i.	191,	Species discutientes, i.	33
Sparadrap resolvent, ii.	386	ii.	234	Species diuretic, i.	266, 268,
Sparadrap of plasters, ii.	661	Species cephalicæ pro		296, ii.	406
Sparadrap spermaceti,		cucuphis, ii.	132	Species emmenagogue,	
i.	443	Species cephalicæ pro		i.	345
Sparadrap vesicant, i.	386	fomentatione, ii.	156	Species emollient, i.	143, 447,
Sparadrap white, ii.	374	Species cold pearl, ii.	268	ii.	215
Sparadrapum album, ii.	393	Species cordial, ii.	192	Species emollient for ca-	
Sparadrapus pallidus, ii.	391	Species cordialis tempera-		taplasm, ii.	200
Spathulæ fetidæ radix,		tæ, ii.	192	Species emollientes, i.	143
ii.	131	Species decocti antine-		Species pro enemate	
Spathum ponderosum, i.	328	phretici, i.	268	emolliente, i.	143
Spearwort lesser, ii.	481	Species decocti antipoda-		Species ad fomentum	
Species adstringentes pro		grici Viennensis, i.	741	emollientem, i.	143
gargarismate, ii.	470	Species decocti pro gar-		Species for epithem, ii.	521
Species alterative, i.	21	garmate adstringente,		Species pro epithemate	
Species amaræ, i.	32	ii.	470	cordis et pulsuum, ii.	521
Species amaricantes, i.	32	Species decocti lignorum,		Species pro fomento, i.	143
Species anodyne, i.	389	i.	740	Species pro fotu ad coli-	
ii.	337	Species for decoction of		cam, i.	447
Species anthelmintic, i.	277	the woods, i.	740	Species pro fotu emme-	
Species anti-apoplectic,		Species pro decocto an-		nagogo, i.	345
ii.	329	tipsorico, i.	731	Species pro fotu ad gan-	
Species antigangrenous,		Species pro decocto car-		grenam, i.	33
i.	33	minativo, i.	397	Species pro fotu laterali,	
Species anti hæmorrhoi-		Species pro decocto de-		ii.	337
dal, i. 114, ii.	471	mulcente, i.	750	Species pro fotu ad hæ-	
Species antinephritic, i.	268	Species pro decocto ad		morrhoidum fluxum,	
Species antipsoric, i.	731	gangrænam, ii.	521	ii.	471
Species antiscorbutic, ii.	482	Species pro decocto in-		Species fotus discutientes	
Species antispasmodic,		fantum, i.	699	et nervini, i.	191
ii.	704	Species pro decocto lig-		Species fumales, i.	332
Species aperitivæ majores,		norum, i.	740	Species fumales pro suffi-	
i.	268	Species pro decocto ni-		mento, ii.	271
Species aperitivæ mino-		gro, i.	692	Species for fumigation, i.	42,
res, i.	269	Species pro decocto pro		332	
Species greater aperitive,		potu infantum, i.	723	Species ad fumigationem	
i.	269	Species pro decocto vul-		anticontagiosam, i.	80
Species lesser aperitive,		nerario, ii.	357	Species fumigatory, i.	55, ii.
i.	269	Species ad decoctum lig-		271	
Species aperitive with rhu-		norum, i.	740	Species pro fumo, i.	332
barb, i.	269	Species demulcent, i.	750	Species against gangrene,	
Species contra apoplex-		Species deterrent, i.	741	ii.	521
iam, ii.	329	Species diaboracis, ii.	608	Species pro gargarismate,	
Species aromatic, ii.	155, 228, 522, 545	Species diacinnamomi,		i.	143
Species aromaticæ, ii.	161, 545	ii.	161	Species de gemmis frigi-	
Species aromaticæ rosa-		Species diacretæ, i.	356	dæ, ii.	268
tæ, ii.	161	Species diajalapæ, i.	543	Species herbarum pro	
Species for arthritic de-		Species diaireos, ii.	131	infuso analapctico, i.	279
coction, i.	741	Species diamargaritæ,		Species of hiera picra, i.	128
Species astrigent, i.	35, 470	i.	204	Species hieræ picræ, i.	128
Species bitter, i.	32, 625	Species diamargaritæ frigi-		Species of hyacinth, ii.	520
Species for black decoc-		gidæ, ii.	268	Species imperatoris, ii.	161
tion, i.	692	Species diambæræ, ii.	161	Species pro infuso diure-	
Species of bryony, i.	345	Species dianisi, ii.	353	tico, ii.	406
Species carminativæ, i.	397	Species dianthos, ii.	522	Species pro infuso pecto-	
Species carminative, i.	397, 447	Species diarrhoden, ii.	520	rale, i.	145
		Species diascordii, ii.	317	Species ad infusum pec-	
		Species of diascordium, ii.	317	torale, i.	145
		Species diatragacanthæ		Species lignorum, i.	740
		frigidæ, i.	767	Species mundificantes, i.	21

- Species nephro-cathartici Foresti, i.* 268
Species nervinæ pro cuphis, ii. 649
Species nervinæ externæ, ii. 234
Species nervine, i. 506, *ii.* 649
Species pectoral, i. 145, 444, 675, 699, 721, 723, *ii.* 188, 403, 699, 717
Species pro odoramento, ii. 131
Species pectorales, i. 145, *ii.* 717
Species for perfume, ii. 131
Species against pleurisy, ii. 337
Species for ptisan, ii. 697
Species purgative, i. 402
Species resolvent, i. 230, *ii.* 530, 557
Species resolventes, ii. 530
Species resolventes externæ, i. 33
Species resolutive, i. 33
Species saluterous, i. 403
Species of the three sanders, ii. 197
Species contra scorbuticum, ii. 482
Species contra sputum sanguinis, i. 35
Species cold of precious stones, ii. 268
Species sudorific, i. 272, 396, *ii.* 551
Species sudorific and cathartic, i. 264
Species ad suffiendum, i. 332
Species pro suffitu, i. 332
Species pro thea, ii. 717, 228
Species pro vaporibus acidi acetici, i. 42
Species pro vaporibus acidi muriatici, i. 55
Species pro vaporibus acidi nitrici, i. 69
Species pro vaporibus superoxydi muriatici, i. 458
Species vermifuge, ii. 32
Species vulnerary, i. 626, *ii.* 357
Specific anti-atrophic, i. 679
Specific Croll's antifebrile, i. 347
Specific Zobel's, i. 161
Specificatum jalapinum, i. 543
Specificum antifebrile Crollii, i. 347
Specificum anti-atrophicum Wepteri, i. 670
Specificum anti-epilepticum Weismanni, i. 586
Specificum Cranii, i. 233
Specificum Paracelsi, ii. 428
Specificum stomachicum Poterii, i. 236
Specificum tartari pharyngicum, i. 161
Specilla cerea, ii. 390
Speedwell male, ii. 717
Sperma ranæ, i. 697
Spermaceti, i. 440
Sperniola Crollii, i. 760
Spicæ alpinæ radix, ii. 703
Spicæ celticæ radix, ii. 703
Spider cellar, i. 271
Spider house, i. 271
Spigelia anthelmia, ii. 619
Spigelia anthelmintic, ii. 619
Spigelia Maryland, ii. 620
Spigelia Marylandica, ii. 620
Spiguel broad-leaved, ii. 580
Spiguel divaricated, ii. 581
Spiguel marsh, ii. 581
Spilanthus acmella, ii. 620
Spilanthus common, ii. 620
Spinacia oleracea, ii. 621
Spinæ albæ flores, ii. 242
Spinæ albæ herba, i. 395, *ii.* 273
Spinæ cervinæ bacca, ii. 487
Spinæ dorsi mustelæ piscis, i. 360
Spinage, ii. 621
Spindle tree, i. 630
Spiræa filipendula, ii. 621
Spiræa ulmaria, ii. 621
Spirit acoustic, i. 189
Spirit alkaline of fennel, i. 220
Spirit of ammonia, i. 172
Spirit anisated of ammonia, i. 173
Spirit aromatic of ammonia, i. 174
Spirit fetid of ammonia i. 176
Spirit succinated of ammonia, i. 180
Spirit of ammonia and cantharides, i. 174
Spirit of ammonia and turpentine, i. 174
Spirit ammoniacous, i. 193
Spirit of angelica, i. 222
Spirit compound of angelica, i. 223
Spirit theriacal of angelica, i. 223
Spirit of anise, ii. 355
Spirit compound of anise, ii. 355
Spirit anodyne, i. 174
Spirit anti-epileptic, i. 542
Spirit antihysterical, i. 304
Spirit anti-icteric, ii. 665
Spirit of ants, i. 40, 676
Spirit aromatic, ii. 168, 594, 690
Spirit ammoniated of assafoetida, i. 304
Spirit of balm, ii. 229
Spirit of Peruvian balsam, i. 320
Spirit Bussius' bezoardic, i. 201
Spirit of bread, ii. 334
Spirit compound of bryony, i. 427
Spirit of brucina, i. 344
Spirit of camphor, i. 372
Spirit of camphor with saffron, i. 373
Spirit of caraway, i. 398
Spirit carminative, i. 559, *ii.* 528
Spirit Sylvius' carminative, i. 224
Spirit compound of castor, i. 426
Spirit of cherry-laurel, i. 62
Spirit of black cherries, i. 63
Spirit of cinnamon, ii. 167
Spirit of cloves, i. 400
Spirit ammoniated of colchicum, i. 539
Spirit vinous of convallaria, i. 542
Spirit of black currants, ii. 79
Spirit of copaiba, i. 555
Spirit of elder, ii. 554
Spirit of fumitory, i. 681
Spirit of grain, i. 104
Spirit of guaiac wood, i. 732
Spirit of hartshorn, i. 185
Spirit of horseradish, ii. 482
Spirit compound of horseradish, ii. 482
Spirit of jessamine, ii. 134
Spirit of juniper, ii. 140
Spirit compound of juniper, ii. 140
Spirit of lavender, ii. 157
Spirit compound of lavender, ii. 158
Spirit of lead, i. 41
Spirit of lemon peel, i. 518
Spirit of mastic, ii. 323
Spirit compound of mastic, ii. 224
Spirit compound of millfoil, i. 37
Spirit of mint, ii. 239
Spirit saccharine of mint, ii. 240
Spirit Riemer's nervine, i. 188
Spirit of nitric æther, i. 95

Spirit nitrous æthereus, i.	95	Spiritus ætheris sulphurici, i.	99	Spiritus cochleariæ, i.	534
Spirit of nutmeg, ii.	255	Spiritus ætheris sulphuricus aromaticus, i.	100	Spiritus cochleariæ compositus, i.	535
Spirit compound of nutmeg, ii.	255	Spiritus ætheris vitriolici, i.	99	Spiritus colchici ammoniatus, i.	539
Spirit odoriferous, i.	518	Spiritus ætheris vitriolici compositus, i.	99	Spiritus concentratissimus, i.	107, 109
Spirit ophthalmic, i.	321	Spiritus alcoholisatus, i.	108	Spiritus concentratus, i.	109
Spirit of orange peel, i.	569	Spiritus alexiterius, ii.	168	Spiritus cornu cervi, i.	185
Spirit of pimenta, ii.	362	Spiritus aluminis, i.	75	Spiritus croci, i.	565
Spirit proof, i.	106	Spiritus ammoniæ, i.	172	Spiritus croci compositus, i.	130
Spirit of pyrethrum, i.	232	Spiritus ammoniæ anisatus, i.	173	Spiritus diphlegmatissimus, i.	108
Spirit compound of pyrethrum, i.	232	Spiritus ammoniæ aromaticus, i.	174	Spiritus diatrion, i.	74
Spirit rectified, i.	104, 106	Spiritus ammoniæ compositus, i.	174	Spiritus dilutus, i.	109
Spirit Schmucker's solvent, i.	193	Spiritus ammoniæ succinatus, i.	181	Spiritus eboris, i.	185
Spirit of rosemary, ii.	527	Spiritus ammoniæ dulcis, i.	172	Spiritus formicarum, i.	40, 676
Spirit compound of rosemary, ii.	528	Spiritus ammoniæ fætidus, i.	176, 304	Spiritus frumenti rectificatus, i.	109
Spirit of roses, ii.	518	Spiritus ammoniæ cum succino, i.	180	Spiritus vinosus frumenti, i.	104
Spirit of saffron, i.	565	Spiritus ammoniato-anisatus, i.	173	Spiritus fuliginis, i.	40
Spirit of sage, ii.	547	Spiritus angelicæ aromaticus camphoratus, i.	223	Spiritus fumans, i.	53
Spirit of salt, i.	53	Spiritus angelicæ compositus, i.	223	Spiritus fumans Beguini, i.	194
Spirit of savine, ii.	144	Spiritus Anhaltinus, ii.	666	Spiritus juniperi, ii.	140
Spirit of scurvy grass, i.	534	Spiritus anisi, ii.	355	Spiritus juniperi compositus, ii.	140
Spirit compound of scurvy grass, i.	535	Spiritus anisi compositus, ii.	355	Spiritus ligni guaiaci, i.	732
Spirit of soot, i.	40, 679	Spiritus anthos, ii.	327	Spiritus liliorum convallium, i.	542
Spirit of spruce fir, i.	22	Spiritus anti-ictericus, ii.	665	Spiritus lumbricorum, i.	185
Spirit stimulant for headache, i.	174	Spiritus antiscorbuticus Drawitzii, i.	535	Spiritus lumbricorum vinosus, i.	173
Spirit of sulphur, i.	79	Spiritus armoraciæ compositus, ii.	482	Spiritus lavandulæ, ii.	157
Spirit of sulphuric æther, i.	100	Spiritus aromaticus, i.	37	Spiritus mastiches compositus, ii.	224
Spirit aromatic of sulphuric æther, i.	100	Spiritus aromaticus acidus, i.	37	Spiritus matricalis, ii.	224
Spirit compound of sulphuric æther, i.	100	Spiritus arthriticus Pottii, ii.	670	Spiritus menthæ, ii.	239
Spirit supercamphorated, i.	372	Spiritus aurantiorum abstractitius, i.	509	Spiritus menthæ crispæ, ii.	240
Spirit theriacal, ii.	314	Spiritus balsamicus, ii.	667	Spiritus menthæ piperitæ, ii.	240
Spirit of thyme, ii.	690	Spiritus Beguini, i.	194	Spiritus menthæ viridis, ii.	240
Spirit of wild thyme, ii.	690	Spiritus bezoardicus Bassii, i.	201	Spiritus millefolii acetatus, i.	37
Spirit compound of wild thyme, ii.	691	Spiritus bryoniæ albæ compositus, i.	427	Spiritus millefolii compositus, i.	37
Spirit of turpentine, ii.	663	Spiritus cæruleus, i.	596	Spiritus Mindereri, i.	195
Spirit of vanilla, ii.	713	Spiritus camphoræ, i.	372	Spiritus muriatico-æthereus, i.	93
Spirit compound of valerian, ii.	711	Spiritus camphoratus, i.	372	Spiritus myristicæ aromatica, ii.	255
Spirit of veronica, ii.	718	Spiritus cari carui, i.	398	Spiritus myrti pimentæ, ii.	362
Spirit vulneraria, ii.	547	Spiritus carminativus, ii.	528	Spiritus nitri acidus, i.	67
Spirit of wine, i.	104	Spiritus carui, i.	398	Spiritus nitri dulcis, i.	94
Spirit of wormwood, i.	24	Spiritus castorei compositus, i.	426	Spiritus nitri fumans Glauberi, i.	68
Spiritus abietis, i.	22	Spiritus cinnamomi, ii.	167	Spiritus nitri tenuior, i.	69
Spiritus absinthii, i.	24			Spiritus nitri tenuis, i.	69
Spiritus aceti, i.	38				
Spiritus acetico-æthereus, i.	92				
Spiritus æruginis, i.	40				
Spiritus æthereus vitriolatus, i.	99				
Spiritus ætheris ferratus, i.	659				
Spiritus ætheris nitratus, i.	94				

Spiritus nitrico-æthereus, i.	94	Spiritus salis marini dephlogisticatus, i.	458	Spiritus volatilis causticus, i.	168
Spiritus nucis moschatæ, ii.	255	Spiritus salis terebinthinatus, i.	670	Spiritus volatilis foetidus, i.	304
Spiritus olei terebinthinæ æthereus, ii.	664	Spiritus sambuci, ii.	554	Spiritus volatilis oleosus, i.	174
Spiritus oleosus aromaticus Sylvi, i.	174	Spiritus saponaceus, ii.	460	Spiritus vulnerarius, i.	37
Spiritus ophthalmicus Himly, i.	321	Spiritus saponatus, ii.	460	Spleenwort, i.	443
Spiritus ophthalmicus Mindereri, i.	195	Spiritus, saponis compositus, ii.	671	Spodylli radix, ii.	5
Spiritus pimentæ, ii.	362	Spiritus saturni, i.	41	Sponge, ii.	622
Spiritus potassatus, ii.	414	Spiritus serpylli, ii.	690	Sponge burnt, ii.	622
Spiritus raphani compositus, i.	535	Spiritus suavolens pro suffitu, i.	518	Sponge prepared, ii.	622
Spiritus raphani rusticani, ii.	482	Spiritus sulphurico-æthereus, i.	99	Sponge stones, i.	354
Spiritus rectificatissimus, i.	108, 109	Spiritus sulphurico-æthereus camphoratus, i.	371	Spongia calcinata, i.	390
Spiritus rectificatus, i.	104, 105, 106, 107, 108, 109	Spiritus sulphurico-æthereus martiatus, i.	659	Spongia cerata, ii.	622
Spiritus aperitivus Renotti, i.	74	Spiritus sulphuricus per campanum, i.	79	Spongia cynosbati, i.	681
Spiritus resolvens, ii.	460	Spiritus sulphuris, i.	79	Spongia lacinulosa, ii.	622
Spiritus rorismarini, ii.	527	Spiritus sulphuris volatilis, i.	194	Spongia marina, ii.	622
Spiritus rorismarini officinalis, ii.	527	Spiritus tartari, i.	74	Spongia præparata, ii.	622
Spiritus rosarum, ii.	518	Spiritus tartari vitriolatus, i.	74	Spongia usta, i.	390, ii. 622
Spiritus rosmarini acetatus, i.	37	Spiritus tenuior, i.	105, 106, 108	Spruce beer, i.	23
Spiritus rosmarini compositus, i.	37	Spiritus terebinthinæ, ii.	663	Spruce fir, i.	21
Spiritus sabinæ, ii.	144	Spiritus thymi, ii.	690	Spurge caper, i.	629
Spiritus salinus aromaticus, i.	174	Spiritus de tribus, i.	74	Spurge cypress, i.	629
Spiritus salis, i.	53	Spiritus vanillæ, ii.	713	Spurge ipecacuanha, i.	629
Spiritus salis ammoniaci, i.	194	Spiritus vini, i.	104, 109	Spurge large-flowering, i.	629
Spiritus salis ammoniaci anisatus, i.	173	Spiritus vini communis, i.	107	Spurge marsh, i.	629
Spiritus salis ammoniaci aquosus, i.	168, 183	Spiritus vini camphoratus, i.	373	Spurge olive, i.	602
Spiritus salis ammoniaci cum calce paratus, i.	168	Spiritus vini dephlegmatissimus, i.	108	Squamariæ radix, ii.	368
Spiritus salis ammoniaci cum cineribus clavellatis, i.	183	Spiritus vini depuratus, i.	107	Squill, ii.	563
Spiritus salis ammoniaci dulcificatus, i.	172	Spiritus vini fortior, i.	105	Squillæ radix, ii.	563
Spiritus salis ammoniaci lactescens, i.	180	Spiritus vini rectificatissimus, i.	105, 107, 108, 109	Squinanthi herba, i.	217
Spiritus salis ammoniaci lacteus, i.	180	Spiritus vini rectificatus, i.	107, 108, 109	Stalagmitis cambogioides, i.	695
Spiritus salis ammoniaci simplex, i.	183	Spiritus vini tartarizatus, i.	108	Stannate of gold, i.	314
Spiritus salis ammoniaci tartarizatus, i.	183	Spiritus vini tenuis, i.	108	Stanni butyrum, ii.	625
Spiritus salis ammoniaci vinosus, i.	172	Spiritus vinosus, i.	99	Stannum, ii.	623
Spiritus salis dulcis, i.	93	Spiritus vinosus rectificatus, i.	105	Stannum granulatum, ii.	623
Spiritus salis marini coagulatus, ii.	440	Spiritus, vinosus tenuior, i.	105	Stannum præparatum, ii.	623
		Spiritus vitrioli, i.	75	Stannum pulveratum, ii.	623, 624
		Spiritus vitrioli dulcis, i.	99	Stannum raspatum, ii.	623
		Spiritus vitrioli phlogisticatus, i.	79	Stannum suboxydatum, ii.	624
		Spiritus viperarum, i.	185	Stachys recta, ii.	622
		Spiritus vitrioli coagulatus Mynsichti, ii.	428	Stachys white, ii.	622
		Spiritus unguæ alcis, i.	185	Staphidis agriæ semines, i.	608
		Spiritus volatilis aromaticus, i.	174	Staphisagriæ semines, i.	608
				Star thistle, i.	346
				Starch, i.	213
				Starch of arum, i.	292
				Starch blue, i.	526
				Starch of horse chesnut, i.	99
				Starch of potato, ii.	618
				Stargrass, i.	112
				Starwort, i.	112
				Statice Caroliniana, ii.	625
				Statice limonicum, ii.	625
				Stella terræ, ii.	213
				Stercus diaboli, i.	299

<i>Stercus caninum album</i>		<i>Strychnina</i> , ii.	626	Subcarbonate of ammo-	
i.	360	<i>Strychnine</i> , ii.	626	nia, i.	182
<i>Stibias potassæ</i> , i.	236	<i>Strychnos colubrina</i> , ii.	626	Subcarbonate of copper, i.	582
<i>Stibium</i> , i.	232	<i>Strychnos nux vomica</i> ,		Subcarbonate of copper	
<i>Stibium oxydatum album</i>		ii.	265	and of ammonia, i.	582
ablutum, i.	236	<i>Styracine</i> , ii.	627	Subcarbonate of iron, i.	646
<i>Stibium oxydatum album</i>		<i>Styrax benzoin</i> , i.	330	Subcarbonate of lead, ii.	371
non ablutum, i.	240	<i>Styrax liquidambar</i> , ii.	627	Subcarbonate of lime pre-	
<i>Stibium oxydatum fus-</i>		<i>Styrax officinale</i> , ii.	628	pared, i.	355
<i>cum</i> , i.	241	<i>Styrax purificata</i> , ii.	628	Subcarbonate of magne-	
<i>Stibium oxydatum sul-</i>		<i>Styrax solidus</i> , ii.	628	sia, ii.	206
<i>phuratum vitrifactum</i> ,		<i>Subacetas cupri</i> , i.	590	Subcarbonate of protox-	
i.	235	<i>Subacetas cupri impu-</i>		ide of sodium, ii.	604
<i>Stibium oxydulatum</i> , i.	234	<i>rus</i> , i.	590, 595	Subcarbonate of soda, ii.	604
<i>Stibium purum</i> , i.	233	<i>Subacetas cupri præpa-</i>		Subcarbonate dried of so-	
<i>Stibium striatum</i> , i.	241	<i>ratum</i> , i.	591	da, ii.	604
<i>Stibium sulphuratum le-</i>		<i>Subacetas cupricus</i> , i.	590,	<i>Subchloras calcis</i> , i.	351
<i>vigatum</i> , i.	241		595	<i>Sub-deuto-acetate of cop-</i>	
<i>Stibium sulphuratum ni-</i>		<i>Subacetas plumbi</i> , ii.	371	<i>per</i> , i.	590
<i>grum</i> , i.	241	<i>Subacetas plumbi dilu-</i>		<i>Sub-deuto-acetate of cop-</i>	
<i>Stibium sulphuratum ni-</i>		<i>tus</i> , ii.	378	<i>per</i> and of ammonia, i.	596
<i>grum alcoholizatum</i> ,		<i>Subacetas plumbi liqui-</i>		<i>Sub-deuto carbonate of</i>	
i.	241	<i>dus</i> , ii.	375	<i>copper</i> , i.	582
<i>Sticks white liquorice</i> ,		<i>Subacetate of copper</i> , i.	590	<i>Sub-deuto-nitrate of cop-</i>	
i.	726	<i>Subacetate of lead</i> , ii.	371	<i>per</i> and of ammonia, i.	558
<i>Sticks yellow liquorice</i> ,		<i>Sub-boras natricum</i> , ii.	608	<i>Sub-deuto-sulphate of</i>	
i.	726	<i>Sub-boras sodæ</i> , ii.	608	<i>copper</i> and of ammo-	
<i>Stillicidium aromaticum</i> ,		<i>Sub-borate of soda</i> , ii.	608	<i>nia</i> , i.	566
i.	193	<i>Subcarbonas ammoniæ</i> ,		<i>Sub-deuto-sulphate of</i>	
<i>Stinking gladwyn</i> , ii.	131	i.	182	<i>mercury</i> , ii.	82
<i>St. John's-wort</i> , ii.	107	<i>Subcarbonas ammoniæ</i>		<i>Sublimate corrosive</i> , ii.	66
<i>Stizolobium pruriens</i> ,		<i>animale</i> , i.	184	<i>Submurias antimonii</i> , i.	254
ii.	625	<i>Subcarbonas ammonia-</i>		<i>Submurias hydrargyri</i> , ii.	46
<i>Stœchadis citrinæ flo-</i>		<i>cum empyreumaticum</i> ,		<i>Submurias hydrargyri</i>	
<i>res</i> , ii.	4	i.	184	<i>ammoniatum</i> , ii.	90
<i>Stœchadis Germanicæ</i>		<i>Subcarbonas ammoniæ</i>		<i>Submurias hydrargyri</i>	
<i>flores</i> , ii.	4	<i>pyro-animale liquidum</i> ,		<i>mitis</i> , ii.	46
<i>Stœchadis Neapolitanæ</i> ,		i.	185	<i>Submurias hydrargyri</i>	
<i>flores</i> , ii.	4	<i>Subcarbonas ammoniæ</i>		<i>præcipitatus</i> , ii.	46, 90
<i>Stœchadis purpuræ Ara-</i>		<i>pyro-oleosum</i> , i.	184	<i>Submurias hydrargyri</i>	
<i>bicæ flores</i> , ii.	160	<i>Subcarbonas bismuthi-</i>		<i>sublimatus</i> , ii.	46
<i>Stone Armenian</i> , i.	582	<i>cum</i> , i.	339	<i>Subnitras bismuthi</i> , i.	339
<i>Stone blood</i> , i.	643	<i>Subcarbonas calcaricæ cre-</i>		<i>Subnitras bismuthicum</i> ,	
<i>Stone carp</i> , i.	354	<i>taceus</i> , i.	352	i.	339
<i>Stone eagle</i> , i.	643	<i>Subcarbonas calcareus</i> , i.	352	<i>Sub-phosphas calcaricæ</i> ,	
<i>Stone Goa</i> , ii.	251	<i>Subcarbonas calcis</i> , i.	352	i.	359
<i>Stone judaic</i> , i.	354	<i>Subcarbonas ferri</i> , i.	646	<i>Sub-phosphate of soda</i> , ii.	610
<i>Stone looking-glass</i> , i.	362	<i>Subcarbonas ferri præ-</i>		<i>Sub-phosphuretum calcis</i> ,	
<i>Stone lynx</i> , i.	354	<i>paratus</i> , i.	646	i.	350
<i>Stone medicinal</i> , i.	160	<i>Subcarbonas ferroso-fer-</i>		<i>Sub-proto-acetate of lead</i> ,	
<i>Stone navel</i> , i.	353	<i>ricum</i> , i.	646	ii.	375
<i>Stone nephretic</i> , ii.	260	<i>Subcarbonas kalicum de-</i>		<i>Sub-proto-borate of so-</i>	
<i>Stone savine</i> , i.	352	<i>puratum</i> , ii.	419	<i>dium</i> , ii.	608
<i>Stones perch</i> , i.	354	<i>Subcarbonas magnesiæ</i> ,		<i>Sub-proto-carbonate of</i>	
<i>Stones sponge</i> , i.	354	ii.	206	<i>barium</i> , i.	328
<i>Stonewort field</i> , ii.	591	<i>Subcarbonas natricum</i> ,		<i>Sub-proto-carbonate of</i>	
<i>Stonewort small</i> , ii.	592	ii.	604	<i>bismuth</i> , i.	339
<i>Storax</i> , ii.	628	<i>Subcarbonas plumbi</i> , ii.	371	<i>Sub-proto-carbonate of</i>	
<i>Storax liquid</i> , ii.	627	<i>Subcarbonas potassæ</i> ,		<i>calcium</i> , i.	352
<i>Storax purified</i> , ii.	628	<i>purissimum</i> , ii.	420	<i>Sub-proto-carbonate of</i>	
<i>Stramonium</i> , i.	605	<i>Subcarbonas protoxidi</i>		<i>lead</i> , ii.	371
<i>Strawberry</i> , i.	677	<i>barii</i> , i.	328	<i>Sub-proto-carbonate of</i>	
<i>Strobili lupuli</i> , ii.	10	<i>Subcarbonas sodæ</i> ii.	604	<i>sodium</i> , ii.	604
<i>Strombus lentiginosus</i> , i.	353	<i>Subcarbonas tritoxidifer-</i>		<i>Sub-proto-carbonate of</i>	
<i>Strychnia</i> , ii.	626	<i>ri</i> , i.	646	<i>potassium</i> , ii.	419

Sub-proto-nitrate of bis- muth, i. 339	Succus dauci inspissatus, i. 608	Sulphas ammoniacæ, i. 190
Sub-proto-nitrate of mer- cury, ii. 84	Succus ebuli, ii. 551	Sulphas ammoniacæ cu- pratus, i. 586
Sub-proto-phosphate of calcium, i. 359	Succus elaterii, ii. 243	Sulphas ammoniacæ, i. 190
Sub-proto-phosphate of of sodium, ii. 610	Succus ex floribus persi- carum, ii. 344	Sulphas argenti, i. 276
Sub-proto-sulphate of lead, ii. 375	Succus fumaricæ, i. 680	Sulphas argenti liquidus, i. 276
Sub-sulphas ammonio-cu- pricus, i. 586	Succus granatorum, ii. 471	Sulphas barytæ, i. 328
Sub-sulphas hydrargyri flavus, ii. 82	Succus juniperi inspissa- tus, ii. 138	Sulphas cadmii, ii. 145
Sub-trito-carbonate of iron, i. 646	Succus granorum kermes, i. 530	Sulphas calceis, i. 361
Succinas ammoniacæ, i. 200	Succus liquiritiæ, i. 723	Sulphas calcicum, i. 361
Succinas ammoniacæ pyro- animale liquidum, i. 200	Succus liquiritiæ depu- ratus, i. 723	Sulphas cinchonini, i. 504
Succinas ammoniacæ pyro- oleoso-aquosum, i. 200	Succus millefolii, i. 35	Sulphas cupri, i. 583
Succinate of ammonia, i. 200	Succus mororum, ii. 248	Sulphas cupri alumin- osus camphoratus, i. 585
Succinate liquid of am- monia, i. 200	Succus mororum inspis- satus, ii. 248	Sulphas cupri ammoni- acalis, i. 586
Succinic acid, i. 75	Succus plantaginis, ii. 366	Sulphas ferri, i. 651
Succinum, ii. 629	Succus pomorum, ii. 214	Sulphas ferri calcinatum, i. 644
Succinum oxygenatum, ii. 631	Succus prunellæ, ii. 466	Sulphas ferri exsiccatus, i. 652
Succisæ radix, ii. 559	Succus pulmonariæ, ii. 469	Sulphas hydrargyri cum excessu oxydi, ii. 82
Succus acaciæ Germa- nicæ inspissatus, i. 34	Succus rhamni cathartici, ii. 487	Sulphas hydrargyri lu- teum, ii. 82
Succus acetosæ, ii. 534	Succus ribesiorum nigro- rum, ii. 505	Sulphas kalico-alumini- cum, i. 158
Succus spissatus aconiti napelli, i. 84	Succus ribium rubrorum, ii. 79	Sulphas magnesiæ, ii. 210
Succus antiscorbuticus, i. 533	Succus rosarum palida- rum, ii. 510	Sulphas natricus, ii. 611
Succus armoraciæ com- positus, ii. 482	Succus rubi idæi, ii. 532	Sulphas oxyduli ferri, i. 651
Succus aurantiorum, i. 507	Succus sambuci, ii. 551	Sulphas potassæ, ii. 428
Succus e baccis myrtil- lorum, ii. 702	Succus sambuci inspissa- tus, ii. 552	Sulphas potassæ et am- moniacæ, ii. 431
Succus beccabunga, ii. 719	Succus saponariæ, ii. 557	Sulphas potassæ et mag- nesiæ, ii. 197
Succus spissatus bella- donnæ, i. 310	Succus spinæ cervinæ, ii. 487	Sulphas potassæ cum sul- phure, ii. 429
Succus berberum, i. 335	Succus taraxaci, ii. 651	Sulphas protoxidi barii, i. 328
Succus cerefolii, i. 444	Succus tritici, ii. 697	Sulphas protoxidi ferri, i. 651
Succus inspissatus aco- niti, i. 84	Succus violarum, ii. 722	Sulphas sodæ, ii. 611
Succus carottæ, i. 607	Suet, i. 315	Sulphas sodæ acidulus, ii. 611
Succus cerasorum, i. 438	Suet beef, i. 315	Sulphas sodæ et magne- siæ, ii. 613
Succus cichorii, i. 462	Suet goat, i. 315	Sulphas super-ammoni- aco-cupreus, i. 586
Succus cicutæ, i. 464	Suet mutton, i. 316	Sulphate of ammonia, i. 190
Succus spissatus cicutæ, i. 464	Suet stag's, i. 315	Sulphate of cinchonine, i. 504
Succus citri, i. 520	Suffitus chlorini, i. 458	Sulphate of copper, i. 583
Succus citrorum, i. 520	Suffitus oxymuriaticus, i. 458	Sulphate of deutoxide of copper, i. 583
Succus cochleariæ, i. 531	Sugar, ii. 539	Sulphate liquid of indigo, ii. 352
Succus cochleariæ com- positus, i. 533	Sugar of cinchona, i. 487	Sulphate of iron, i. 651
Succus spissatus conii, i. 465	Sugar of lead, ii. 383	Sulphate of iron dried, i. 652
Succus spissatus conii maculati, i. 465	Sugar compound mercu- rial, ii. 26	Sulphate of kadmium, ii. 145
Succus cucumeris, i. 578	Sugar of milk, ii. 150	Sulphate of magnesia, ii. 210
Succus cydoniorum, i. 599	Sulphas aluminæ, i. 158	Sulphate of mercury, ii. 83
	Sulphas aluminæ aceta- tus, i. 160	Sulphate of morphine, ii. 246
	Sulphas acidulum alu- minæ, i. 158	Sulphate of potass, ii. 428
	Sulphas aluminæ et po- tassæ acidulus, i. 158	Sulphate of quinine, ii. 478
	Sulphas acidus aluminæ et potassæ, i. 158	
	Sulphas aluminæ fusus, i. 158	

Sulphate of silver, i.	276	Sulphuret simple of potass, ii.	417	Sulphuretum vini, i.	101
Sulphate of soda, ii.	611	Sulphuret of protoxide of potassium, ii.	416	Sulphuris clyssus, i.	79
Sulphate of soda and of magnesia, ii.	613	Sulphuretted sulphate of potass and of mercury, ii.	84	Sulphuris ros, i.	75
Sulphur, ii.	632	Sulphuretum ammoniæ, i.	194	Sulphurwort common, ii.	343
Sulphur antimoniated, i.	261	Sulphuretum antimonii, i.	241	Sulphurwort meadow, ii.	343
Sulphur antimoniatum fuscum, i.	252	Sulphuretum antimonii præcipitatum, i.	252	Sumac folia, ii.	503
Sulphur antimomi præcipitatum, i.	243	Sulphuretum arsenici flavum, i.	282	Sumach curriers, ii.	503
Sulphur antimoniatum tartarizatum, i.	243	Sulphuretum arsenici rubrum, i.	282	Sumach common upright, ii.	503
Sulphur auratum liquidum, i.	252	Sulphuretum calcaris et stibii, i.	253	Sundew long-leaved, i.	619
Sulphur auratum saponatum, i.	251	Sulphuretum calcis et arsenici, i.	283	Sundew round-leaved, i.	619
Sulphur cupri, i.	582	Sulphuretum calcis stibiatus, i.	253	Superacetates cupri impurus, i.	590
Sulphur compound flow-ers of, i.	135	Sulphuretum ferri, i.	639	Superacetates plumbi, ii.	382
Sulphur lotum, ii.	632	Sulphuretum hydrargyri nigrum, ii.	28	Supercarburetum ferri, i.	639
Sulphur lycopodii, ii.	204	Sulphuretum hydrargyri rubrum, ii.	43	Super-deuto-sulphate of copper, i.	583
Sulphur oil of, i.	79	Sulphuretum hydrargyri simplex, ii.	28	Supermuriates hydrargyri, ii.	66
Sulphur præcipitatum, ii.	633	Sulphuretum hydrargyri stibiatum, ii.	30	Superoxalates calicum, ii.	446
Sulphur precipitated, ii.	633	Sulphuretum hydrargyri et stibii, ii.	30	Superoxalates potassæ, ii.	446
Sulphur rubrum, i.	243	Sulphuretum hydrargyri sublimatum, ii.	143	Superoxalate of potass, ii.	446
Sulphur spirit of, i.	79	Sulphuretum hydrosulphas antimonii, i.	249	Superoxydum manganicum, ii.	217
Sulphur stibiatum rubrum, i.	243	Sulphuretum kalicum, ii.	416	Superoxydum plumbi, ii.	371
Sulphur sublimatum lotum, ii.	632	Sulphuretum lixivis, ii.	416	Superoxydum plumbosum, ii.	370
Sulphur vegetabile, ii.	204	Sulphuretum lixivis stibi-atum, i.	240	Superoxygenated muriate of mercury, ii.	66
Sulphur washed, ii.	632	Sulphuretum oxyduli stibii hydrogenatum, i.	249	Superphosphates calcis, i.	361
Sulphuret of antimony, i.	241	Sulphuretum plumbi, ii.	369	Super-proto-carbonate of calcium, i.	358
Sulphuret of antimony and of lime, i.	252	Sulphuretum potassæ, ii.	416	Super-proto-carbonate of iron, i.	650
Sulphuret of antimony and of potass, i.	254	Sulphuretum potassæ et ferri, i.	640	Super-proto-phosphate of calcium, i.	361
Sulphuret golden of antimony, i.	249	Sulphuretum sodæ, ii.	602	Super-proto-sulphate of potassium, ii.	431
Sulphuret precipitated of antimony, i.	252	Sulphuretum sodii protoxidi, ii.	602	Super-proto-sulphate of sodium, ii.	611
Sulphuret prepared of antimony, i.	241	Sulphuretum stanni, ii.	625	Super-proto-tartrate of potassium, ii.	449
Sulphuret pulverized of antimony, i.	241	Sulphuretum stibii arsenicatum, i.	284	Super-proto-tartrate borated of potassium, ii.	454
Sulphuret of arsenic, i.	282	Sulphuretum stibii hydro-genatum, i.	249	Super-protoxalate of potassium, ii.	446
Sulphuret red of arsenic, i.	282	Sulphuretum stibii nativum, i.	241	Superstibias kalica, i.	236
Sulphuret yellow of arsenic, i.	282	Sulphuretum stibii oxydulati, i.	249	Supersulphates aluminæ et potassæ, i.	158
Sulphuret of barium, i.	325	Sulphuretum stibii præparatum, i.	241	Supersulphates argillæ alkalisatum, i.	158
Sulphuret of iron, i.	639	Sulphuretum stibii venale, i.	241	Supersulphates potassæ, ii.	431
Sulphuret of lead, ii.	369			Supersulphate of alumina and of potassa, i.	158
Sulphuret liquid of lime, i.	351			Supertartrate of potass, ii.	449
Sulphuret of mercury, ii.	42			Supertartras potassæ, ii.	449
Sulphuret of mercury and of antimony, ii.	30			Supertartras kalicum, ii.	449
Sulphuret black of mercury, ii.	28			Supersulphuret of potass, ii.	417
Sulphuret of potass, ii.	416			Suppository anthelmintic, i.	122, 126
Sulphuret of potass and of iron, i.	640			Suppository antihæmorrhoidal, i.	391
Sulphuret martial of potass, ii.	418				

Suppository corroborant, ii.	475	Syrup of birch, i.	338	Syrup of digitalis, i.	615
Suppository emollient, ii.	688	Syrup of blessed thistle, i.	525	Syrup of dittany, ii.	322
Suppository irritant, i.	572	Syrup of blueberries, ii.	702	Syrup drastic, ii.	562
Suppository laxative, ii.	458	Syrup of borage, i.	341	Syrup of elder, ii.	552
Suppository sedative, ii.	297	Syrup Boullay's, i.	489	Syrup of elecampane, ii.	117
Suriumamine, i.	717	Syrup Boyle's, ii.	642	Syrup emetic, i.	621
Sus scrofa, i.	359	Syrup of dwarf bramble, ii.	532	Syrup Ange Sala's emetic, i.	261
Swamp-cabbage, i.	619	Syrup of bryony, i.	344	Syrup of emetine, i.	621
Swallow-wort, i.	295	Syrup of buckthorn, ii.	487	Syrup expectorant, ii.	128
Sweet basil, ii.	269	Syrup of bugloss, i.	216	Syrup of French lavender, ii.	160
Sweet flag, i.	85	Syrup byzantine, i.	267	Syrup compound of French lavender, ii.	160
Sweet yarrow, i.	35	Syrup of red cabbage, i.	343	Syrup of fumitory, i.	680
Swietenia mahogoni, i.	470	Syrup of calf's lung, i.	702	Syrup of garlic, i.	114
ii.	641	Syrup cardiac, ii.	172	Syrup of gentianine, i.	716
Suppositorium mellis irritans, i.	572	Syrup of cascarrilla, i.	569	Syrup of ginger, i.	204
Suppositorium ad alvum sollicitandum, ii.	458	Syrup of lesser centaury, i.	626	Syrup of gold, i.	312
Symphyti dentarii radix, i.	609	Syrup aromatic chalybeate, i.	672	Syrup of ground ivy, i.	721
Symphyti maculosi herba, ii.	469	Syrup of chamomile, i.	452	Syrup of guaiac, i.	737
Symphyti minimi herba, i.	329	Syrup of cherries, i.	438	Syrup aqueous of gum ammoniac, i.	743
Symphytum officinale, ii.	641	Syrup compound of china root, ii.	593	Syrup vinous of gum ammoniac, i.	744
Symphytum tuberosum, ii.	641	Syrup of chloruret of gold and of sodium, i.	313	Syrup of gum arabic, i.	752
Synopsis, i.	155	Syrup of cichory, i.	463	ii.	188
Syringii, i.	622	Syrup compound of cichory, i.	463	Syrup of hemlock, i.	464
Syringites, i.	353	Syrup ferruginous of cichory, i.	671	Syrup of henbane, ii.	101
Syrup of acetate of morphine, ii.	447	Syrup vinous of cinchona, i.	497	Syrup of honeysuckle, ii.	204
Syrup alkaline, ii.	423	Syrup aqueous of cinchona, i.	491	Syrup against whooping cough, ii.	642
Syrup of almonds, i.	209	Syrup of cinchonine, i.	504	Syrup of horehound, ii.	222
Syrup bitter alterative, ii.	485	Syrup alcoholic of cinnamon, ii.	172	Syrup of horseradish, ii.	485
Syrup Larrey's alterative and antisyphilitic, ii.	92	Syrup aqueous of cinnamon, ii.	171	Syrup of houseleek, ii.	584
Syrup compound of ammonia, i.	172	Syrup vinous of cinnamon, ii.	172	Syrup of hydrochloric acid, i.	55
Syrup antimoniated of vinegar, i.	261	Syrup of citric acid, i.	53	Syrup of hydrocyanate of potass, i.	61
Syrup antiseptic, ii.	484	Syrup of clove-pink, i.	610	Syrup of hydrocyanic acid, i.	60
Syrup Portal's antiseptic, ii.	74	Syrup of colchicum, i.	540	Syrup of hyssop, ii.	111
Syrup antiscrofulous, i.	313	Syrup of coltsfoot, ii.	699	Syrup compound of hyssop, ii.	112
Syrup St. Idéphont's antisyphilitic, ii.	71	Syrup of columbine, i.	270	Syrup scillitic of hyssop, ii.	112
Syrup of apples, ii.	214	Syrup compound of comfrey, ii.	642	Syrup of jalap, i.	551
Syrup compound of apples, i.	414	Syrup of great comfrey, ii.	642	Syrup of Iceland moss, ii.	188
Syrup astrigent, ii.	480	Syrup of copaiba, i.	555	Syrup of iceplant, ii.	242
Syrup Joubert's astrigent, ii.	702	Syrup of coral, i.	364	Syrup compound of Jerusalem oak, i.	458
Syrup of balm, ii.	228	Syrup of corsican moss, ii.	5	Syrup of iodine, ii.	120
Syrup of Peruvian balsam, i.	320	Syrup of cowslip, ii.	465	Syrup of ipecacuanha, ii.	130
Syrup of balsam of tolu, i.	323	Syrup of cranberries, ii.	702	Syrup aqueous of ipecacuanha, ii.	127
Syrup of barberries, i.	335	Syrup Cuisinier's, ii.	600	Syrup compound of ipecacuanha, ii.	129
Syrup of belladonna, i.	311	Syrup of black currants, ii.	505	Syrup of kermes, i.	530
Syrup Bellet's, ii.	87	Syrup of red currants, ii.	506	Syrup of kino, ii.	146
Syrup of benzoïn, i.	330	Syrup of daisy, i.	329	Syrup of lactucarium, ii.	154
Syrup of betony, i.	337	Syrup demulcent pectoral, i.	449	Syrup of lemon, i.	521
		Syrup Dessessart's, ii.	129	Syrup compound of lemon, i.	522
				Syrup of lemon juice, i.	522

Syrup aqueous of lemon-peel, i.	521	Syrup of plantain, ii.	367	Syrup aqueous of squill, ii.	569
Syrup vinous of lemon-peel, i.	522	Syrup glycyrrhized of polygala, ii.	404	Syrup compound of squill, ii.	569
Syrup Lescure's, i.	493	Syrup of pomegranate, ii.	471	Syrup of stramonium, i.	606
Syrup of liquorice, i.	728	Syrup of poppy, ii.	339	Syrup of strawberries, i.	677
Syrup of lupuline, ii.	12	Syrup anisated of poppy, ii.	340	Syrup Steven's, ii.	600
Syrup of manna, ii.	220	Syrup crocated of poppy, ii.	340	Syrup of St. John's-wort, ii.	108
Syrup of marshmallows, i.	149	Syrup of red poppy, ii.	335	Syrup of sulphate of morphine, ii.	246
Syrup compound of marshmallows, i.	150	Syrup of protonitrate of mercury, ii.	85	Syrup of sulphate of quinine, ii.	479
Syrup mercurial, ii.	96	Syrup of pumpkin, i.	579	Syrup compound of sulphate of quinine, ii.	479
Syrup Cheron's mercurial, ii.	72	Syrup of quince, i.	599	Syrup of sulphuret of potass, ii.	418
Syrup of mercury and of gum, ii.	14	Syrup of raspberries, ii.	532	Syrup sulphuretic, i.	195
Syrup Lagneau's of mercury, ii.	14	Syrup of raspberry vinegar, i.	43	Syrup of sulphuric acid, i.	76
Syrup of Mexican tea, i.	457	Syrup resolvent, i.	263	Syrup of sulphuric æther, i.	101
Syrup of milfoil, i.	36	Syrup of rhubarb, ii.	495	Syrup of sulphurwort, ii.	343
Syrup of mint, ii.	235	Syrup alkaline of rhubarb, ii.	499	Syrup of sweetflag, i.	87
Syrup of mugwort, i.	289	Syrup aromatic of rhubarb, ii.	496	Syrup of tartaric acid, i.	82
Syrup compound of mugwort, i.	289	Syrup compound of rhubarb, ii.	496	Syrup of tartrate of mercury, ii.	98
Syrup of mulberries, ii.	248	Syrup compound of roses, ii.	514	Syrup of tobacco, ii.	262
Syrup of myrtle, ii.	258	Syrup of pale roses, ii.	514	Syrup of tolu, i.	323
Syrup of nettle, ii.	701	Syrup of red roses, ii.	513	Syrup of valerian, ii.	706
Syrup of nutmeg, ii.	255	Syrup of safflower, i.	399	Syrup of vanilla, ii.	712
Syrup of opium, ii.	293	Syrup aqueous of saffron, i.	563	Syrup vermifuge, ii.	3
Syrup alcoholic of opium, ii.	283	Syrup vinous of saffron, i.	564	Syrup purgative vermifuge, i.	419
Syrup emulsive of opium, ii.	293	Syrup of sarsaparilla, ii.	595	Syrup of vinegar, i.	43
Syrup succinated of opium, ii.	294	Syrup of sarsaparilla and guaiac, ii.	600	Syrup of violets, ii.	716
Syrup vinous of opium, ii.	276	Syrup vinous of sarsaparilla, ii.	601	Syrup aperient of violets, ii.	723
Syrup of orange flowers, i.	508	Syrup of sassaparilla, ii.	183	Syrup of viper, i.	701
Syrup of orange juice, i.	507	Syrup of scammony, ii.	562	Syrup of water lily, ii.	268
Syrup alcoholic of orange peel, i.	515	Syrup of scurvy-grass, i.	533	Syrup of wild mustard, i.	623
Syrup aqueous of orange peel, i.	510	Syrup compound of scurvy-grass, i.	533	Syrup compound of wild mustard, i.	624
Syrup vinous of orange peel, i.	511	Syrup of scordium, ii.	683	Syrup of wood-lice, ii.	272
Syrup of peach-flowers, ii.	344	Syrup of selfheal, ii.	466	Syrup of wood sorrel, ii.	327
Syrup of peach leaves, ii.	344	Syrup of senna, i.	413	Syrup of wormwood, i.	25
Syrup pectoral, i.	324	Syrup compound of sen- na, i.	414	Syrup of wormwood and of quince, i.	25
ii.	294, 340, 576, 718	Syrup of seneka, ii.	404	Syrupus absinthii, i.	25
Syrup balsamic pectoral, i.	325	Syrup moschated of se- neka, ii.	405	Syrupus de absinthio, i.	25
Syrup Courti's pectoral, ii.	128	Syrup simple, ii.	539	Syrupus absinthii com- positus, i.	25
Syrup Maloet's pectoral, ii.	294	Syrup of sloe flowers, ii.	468	Syrupus aciaciarum flo- rum, ii.	468
Syrup of peony, ii.	329	Syrup of smallage, i.	267	Syrupus aceti, i.	43
Syrup of peppermint, ii.	236	Syrup compound of smal- lage, i.	267	Syrupus acetosæ, ii.	534
Syrup of phosphoric acid, i.	73	Syrup Boudet's of snails, i.	701	Syrupus acetosellæ, ii.	327
		Syrup of sorrel, ii.	534	Syrupus acetositis ci- tri, i.	522
		Syrup of spigelia, ii.	620		
		Syrup acetose of squill, ii.	574		

Syrupus acidi acetici, i.	43	Syrupus cerasorum, i.	438	Syrupus digitalis acetici, i.	615
Syrupus acidi citrici, i.	522	Syrupus cerasorum acidorum, i.	438	Syrupus emulsivus, i.	209
Syrupus acidi phosphorici, i.	73	Syrupus chalybeatus, i.	672	Syrupus enulæ, ii.	117
Syrupus acidi sulphurici, i.	76	Syrupus chamæmeli, i.	452	Syrupus erysimi, i.	623
Syrupus de acido tartarico, i.	82	Syrupus chamomillæ, i.	452	Syrupus de erysimo, i.	623
Syrupus cum æthere sulphurico paratus, i.	101	Syrupus chenopodii, i.	457	Syrupus de erysimo, compositus, i.	624
Syrupus Alexandrinus, ii.	518	Syrupus chinæ chinæ, i.	491	Syrupus de erysimo Lobellii, i.	624
Syrupus allii, i.	114	Syrupus chinæ compositus, ii.	593	Syrupus fragorum, i.	677
Syrupus de alio, i.	114	Syrupus de cichorio, i.	463	Syrupus de frambæsiis, ii.	532
Syrupus de althæa, i.	149	Syrupus de cichorio compositus, ii.	496	Syrupus de floribus aurantiorum, i.	508
Syrupus althææ, i.	149	Syrupus cicutæ, i.	464	Syrupus de floribus dianthi caryophylli, i.	610
Syrupus althææ officinalis, i.	149	Syrupus cinchonæ, i.	491	Syrupus florum aurantii, i.	508
Syrupus amomi zingiberis, i.	204	Syrupus citri aurantii, i.	510	Syrupus florum citri aurantii, i.	508
Syrupus amygdalarum, i.	209	Syrupus citri medicæ, i.	522	Syrupus florum paralyseos, ii.	465
Syrupus amygdali persicæ, ii.	344	Syrupus citri e toto, i.	521	Syrupus de fumaria, i.	680
Syrupus de amygdali persicæ floribus, ii.	344	Syrupus de citrorum succo, i.	522	Syrupus fumarie, i.	680
Syrupus de amygdalis, i.	209	Syrupus de cochlearia officinali, i.	533	Syrupus glycyrrhizæ, i.	728
Syrupus amygdalium, i.	209	Syrupus cochleariæ compositus, i.	484, 533	Syrupus de granatis, ii.	471
Syrupus anethi fœniculi compositum, i.	267	Syrupus colchici, i.	540	Syrupus granatorum, ii.	471
Syrupus anodynus, i.	606	Syrupus colchici acetici, i.	540	Syrupus granorum kermes, i.	530
Syrupus anthemidis, i.	452	Syrupus colchici autumnalis, i.	540	Syrupus guaiacini, i.	737
Syrupus antiscorbuticus, ii.	484	Syrupus communis, ii.	539	Syrupus gummi acaciæ, i.	752
Syrupus aperitivus, i.	267	Syrupus copaivicus, i.	555	Syrupus hederæ terrestis, i.	721
Syrupus de armoracia, compositus, ii.	484	Syrupus coralli, ii.	540	Syrupus hordeatus, i.	209
Syrupus armoracæ, ii.	485	Syrupus coralli rubri, i.	364	Syrupus hydrargyri, ii.	14
Syrupus de arundine, i.	87	Syrupus corallinorum, i.	364	Syrupus hyperici, ii.	108
Syrupus atriplicis Mexicanæ, i.	457	Syrupus de cortice citri, i.	521	Syrupus hypnoticus, ii.	293
Syrupus aurantii, i.	510	Syrupus e corticibus ligni sassafras, ii.	183	Syrupus hyssopi, ii.	111
Syrupus de aurantiis, i.	507	Syrupus corticis citri aurantii, i.	510	Syrupus hyssopi cum scilla, ii.	112
Syrupus aurantiorum, i.	510	Syrupus corticum citri, i.	521	Syrupus hyssopi scilliticus, ii.	112
Syrupus balsami tolutani, i.	323	Syrupus corticis peruviani, i.	491	Syrupus de hyssopo, ii.	111
Syrupus balsamicus, i.	323	Syrupus corticum citri amarus, i.	522	Syrupus de jalapa, i.	551
Syrupus balsamicus de tolu, i.	323	Syrupus corticum citri vinosus, i.	522	Syrupus jalapinus, i.	551
Syrupus de berberide, i.	335	Syrupus corticum aurantiorum, i.	510	Syrupus ipecacuanhæ, ii.	130
Syrupus berberum, i.	335	Syrupus croci, i.	563, 564	Syrupus kali sulphurati, ii.	418
Syrupus Boyleanus, ii.	642	Syrupus Cuisnieri, ii.	600	Syrupus kermesinus, i.	530
Syrupus byzantinus Mesues, i.	267	Syrupus de cydoniis, i.	599	Syrupus de kinkina, i.	491
Syrupus cardui benedicti, i.	525	Syrupus cydoniorum, i.	599	Syrupus limonis, i.	522
Syrupus de capitibus papaveris, ii.	339	Syrupus diacodii, ii.	339	Syrupus limonum, i.	522
Syrupus carthami solutior, i.	399	Syrupus diacodion, ii.	339	Syrupus longæ vitæ, ii.	241
Syrupus caryophylli rubri, i.	610	Syrupus diacodion crocatum, ii.	340	Syrupus liquiritiæ, i.	728
Syrupus cascarillæ, i.	569	Syrupus de dictamno, ii.	332	Syrupus magistralis, i.	691
Syrupus cassiæ sennæ, i.	413	Syrupus digitalis, i.	615	Syrupus e manna, ii.	220
				Syrupus de manna laxativus, ii.	220
				Syrupus mannæ cum sen- na, ii.	220
				Syrupus de marrubio, ii.	222
				Syrupus marrubii,	222
				Syrupus de meconio, ii.	339

Syrupus menthæ, ii.	235	Syrupus e quinque radicibus, i.	267	Syrupus scillæ maritimæ, ii.	574
Syrupus menthæ crispæ, ii.	235	Syrupus quinque radicum aperientum, i.	267	Syrupus scilliticus, ii.	569
Syrupus menthæ piperitidis, ii.	236	Syrupus raphani rusticani, ii.	485	Syrupus scordii, ii.	683
Syrupus mercurialis, ii.	241	Syrupus derhabarbaro, ii.	495	Syrupus sedativus, ii.	293
Syrupus Belleti mercurialis, ii.	87	Syrupus rhamni, ii.	487	Syrupus sedi majoris, ii.	584
Syrupus de millefoliis, i.	36	Syrupus rhamni cathartici, ii.	487	Syrupus sempervivi, ii.	584
Syrupus mineralis, i.	76	Syrupus regius, ii.	518	Syrupus senegæ, ii.	404
Syrupus mori, ii.	248	Syrupus rhei, ii.	495	Syrupus senegæ cum liquiritia, ii.	404
Syrupus mori nigræ, ii.	248	Syrupus rhei alkalisatus, ii.	499	Syrupus senegæ moschatatus, ii.	405
Syrupus mororum, ii.	248	Syrupus rhei aromaticus, ii.	496	Syrupus de senna rosaceus, ii.	514
Syrupus muriaticus, i.	55	Syrupus rhei cum senna, i.	414	Syrupus sennæ, i.	413
Syrupus de musco quercino, ii.	188	Syrupus de rheo, ii.	496	Syrupus sennæ compositum, i.	413
Syrupus myrtilli, ii.	702	Syrupus rhœados, ii.	335	Syrupus simplex, ii.	539
Syrupus de myrto, ii.	258	Syrupus ribesiæ nigræ, ii.	79	Syrupus similacis sarsaparillæ, ii.	600
Syrupus naphæ, i.	508	Syrupus ribis nigri, ii.	79	Syrupus spleneticus, i.	267
Syrupus de nicotiana, ii.	262	Syrupus ribium, ii.	506	Syrupus de stœchade, ii.	160
Syrupus nucistæ, ii.	255	Syrupus rosæ, ii.	514	Syrupus destœchade compositus, ii.	160
Syrupus nymphææ, ii.	268	Syrupus rosæ centifoliæ, ii.	514	Syrupus succi citri, i.	522
Syrupus opiatius, ii.	293	Syrupus rosæ Gallicæ, ii.	513	Syrupus succi citri medicæ, i.	522
Syrupus opii, ii.	293	Syrupus rosarum pallidarum, ii.	514	Syrupus e succo aurantiorum, i.	507
Syrupus opii succinatus, ii.	294	Syrupus rosarum rubrarum, ii.	513	Syrupus e succo citri, i.	522
Syrupus ex oxydulo hydrargyri Hahnemannii, ii.	85	Syrupus rosarum solutivarum, ii.	514	Syrupus sudorificus, ii.	600
Syrupus papaveris erratici, ii.	335	Syrupus rosarum solutivus cum senna, ii.	514	Syrupus sulphureticus, i.	195
Syrupus papaveris albi, ii.	339	Syrupus rosatus simplex, ii.	514	Syrupus de sulphureto potassæ, ii.	418
Syrupus papaveris rhœados, ii.	335	Syrupus rosatus solutivus, ii.	514	Syrupus sulphuris, ii.	418
Syrupus papaveris rubri, ii.	335	Syrupus de rosis pallidis, ii.	514	Syrupus symphyti, ii.	642
Syrupus papaveris somniferi, ii.	339	Syrupus e rosis siccis, ii.	513	Syrupus e symphyto Fernellii, ii.	642
Syrupus pectoralis, ii.	718	Syrupus rubi arctici, ii.	532	Syrupus thebaicus, ii.	293
Syrupus persicarum florum, ii.	344	Syrupus rubi idæi, ii.	532	Syrupus toluiferæ balsami, i.	323
Syrupus ad pertussim, i.	489	Syrupus rubi idæi acetatus, i.	43	Syrupus toluianus, i.	323
Syrupus ad pertussim, ii.	642	Syrupus sacchari, ii.	539	Syrupus tunicæ, i.	610
Syrupus petalorum violarum, ii.	716	Syrupus saccharinus cum aqua naphæ, i.	509	Syrupus tussilaginis, ii.	699
Syrupus plantaginis, ii.	367	Syrupus sambuci, ii.	552	Syrupus valerianæ, ii.	706
Syrupus de pomis compositus, i.	414	Syrupus de sarsaparilla compositus, ii.	600	Syrupus de veronica, ii.	718
Syrupus de pomis solutivus, i.	414	Syrupus de sarsaparilla et senna compositus, ii.	600	Syrupus violæ odoratæ, ii.	716
Syrupus pomorum redolentium, ii.	214	Syrupus sarsaparillæ, ii.	595	Syrupus violarum, ii.	716
Syrupus prassii albi, ii.	222	Syrupus de scammonio, ii.	562	Syrupus violarum solutivus, ii.	723
Syrupus primulæ veris, ii.	465	Syrupus de scilla, ii.	569	Syrupus viperarum, i.	701
Syrupus prophylacticus, ii.	241	Syrupus scillæ, ii.	569	Syrupus vitriolatus, i.	76
Syrupus prunellæ, ii.	466	Syrupus scillæ compositus, ii.	569	Syrupus de urtica, ii.	701
Syrupus psychotriæ emeticæ, ii.	130			Syrupus zingiberis, i.	204
Syrupus e pulmonibus vitulinis, i.	702			Tabaci herba, ii.	260
				Tabellæ absorbentes, ii.	100
				Tabellæ althææ, i.	146
				Tabellæ anisi compositæ, i.	726

Tabellæ ad ardorem ventriculi, i.	357	Tartarum emeticum, i.	256	Tartris stibii cum potassa, i.	255
Tabellæ diacarthami, ii.	561	Tartarus ammoniacalis, ii.	456	Tea green, ii.	686
Tabellæ diarrhodon, ii.	511	Tartarus ammoniatus, ii.	456	Tea hæmastic, ii.	513
Tabellæ diatragacanthæ, i.	768	Tartarus antimonialis, i.	255	Tea Mexican, i.	456
Tabellæ glychrrhizæ glabræ compositæ, i.	726	Tartarus chalybeatus, i.	667	Tea mountain, i.	606
Tabellæ magnesiæ, ii.	209	Tartarus chalybeatus solubilis, i.	667	Tea New Jersey, i.	432
Tabellæ manus christi, ii.	511	Tartarus crudus, ii.	449	Teazle, i.	618
Tabellæ de rheo, ii.	493	Tartarus diureticus, ii.	441	Teazle wild,	618
Tabellæ rosatæ, ii.	511	Tartarus emeticus, i.	255	Tecolithes, i.	354
Tabellæ iridis florentinæ compositæ, ii.	132	Tartarus mercurialis, ii.	97	Teeth of the hippopotamus, i.	359
Tabellæ de semine santónico, ii.	583	Tartarus natronatus, ii.	455	Teeth of the unicorn, i.	359
Tabellæ ad sodam, i.	387	Tartarus solubilis, ii.	447	Teeth of the wild boar, i.	359
Tabellæ de tribus, ii.	438	Tatarus solubilis ammoniacalis, ii.	456	Tela emplastica epulotica, ii.	391
Tabellæ contra vermes, ii.	583	Tartarus tartarisatus, ii.	447	Tela Gaulthieri, ii.	393
Tabellæ violatæ, ii.	722	Tartarus vitriolatus, ii.	428	Tela ichthyocollæ glutinans, i.	706
Tabulæ alimentariæ, i.	705	Tartras ammoniacæ pyro-oleoso-aquosus, i.	202	Telephii vulgaris, ii.	580
Tabulæ althææ, i.	146	Tartras antimonii, i.	255	Terebinthina, ii.	653
Tabulæ ferri compositæ, i.	643	Tartras antimonii, et potassæ, i.	255	Terebinthina balsamea, i.	318
Tabulæ instantes, i.	638	Tartras kalico-ferricum, i.	667	Terebinthina Canadensis, i.	318
Tabulæ martiales, i.	638, 643	Tartras kalico-natricum, ii.	455	Terebinthina cocta, ii.	673
Tacamahaca, ii.	644	Tartras kalinus, ii.	447	Terebinthina empyreumatica, ii.	364
Tafia, i.	104	Tartras lixivie, ii.	447	Terra aluminis, i.	156
Talc, ii.	213	Tartras potassæ, ii.	447	Terra absorbens mineralis, ii.	206
Talcum venetæ, ii.	213	Tartras acidulus potassæ, ii.	449	Terra amara ærata, ii.	206
Talpa Europæa, i.	355	Tartras potassæ ferratus, i.	667	Terra amara sulphurica, ii.	210
Talpæ combustæ, i.	391	Tartras potassæ et ferri, i.	667	Terra argillacea pura, i.	156
Talus leporum, i.	359	Tartras potassæ et ferri liquidus, i.	669	Terra bolaris, i.	156
Tamarind tree, ii.	645	Tartras potassæ ferrosus, i.	667	Terra calcaria pura, i.	347
Tamarindus Indica, ii.	645	Tartras potassæ ferruginosus, i.	667	Terra calcarata, i.	352
Tamarisk French, ii.	648	Tartras potassæ et sodæ, ii.	455	Terra foliata tartari, ii.	441
Tamarisk German, ii.	648	Tartras potassæ sodicus, ii.	455	Terra foliata mercurii, ii.	95
Tamarix Gallica, ii.	648	Tartras potassæ stibialis, i.	255	Terra foliata mineralis, ii.	614
Tamarix Germanica, ii.	648	Tartras potassæ stibiosus, i.	255	Terra lemnia, i.	155
Tamnus, ii.	649	Tartras sodæ et kali, ii.	455	Terra lemniana, i.	155
Tamnus communis, ii.	649	Tartras stibii, i.	255	Terra orleana, ii.	352
Tanaceti hortensis radix, i.	317	Tartras subpotassicus, ii.	449	Terra ossium, i.	359
Tanacetum vulgare, ii.	649	Tartras superpotassicus, ii.	449	Terra ponderosa acetata, i.	328
Tansey, ii.	649	Tartrate of mercury, ii.	97	Terra ponderosa ærata, i.	328
Tapioca, i.	632	Tartrate of mercury and of potass, ii.	97	Terra ponderosa muriata, i.	325
Tar, ii.	364	Tartrate of potass, ii.	447	Terra ponderosa pura, i.	327
Tar Barbadoes, ii.	345	Tartrate of potass and of soda, ii.	455	Terra ponderosa salita, i.	325
Tar mineral, ii.	345	Tartris lixivie stibiatus, i.	255	Terra sigellata, i.	155
Taraxacum dens leonis, ii.	651	Tartris potassæ acidulus stibiatus, i.	255	Terra tripolitiana, ii.	686
Tare, ii.	719			Terra talcosa oxyanthracodes, ii.	206
Tarragon, i.	291			Terra tartari crystallibilis, ii.	614
Tartar borated, ii.	454			Terra viridis, i.	582
Tartar crude, ii.	449			Terra vitrioli dulcis, i.	644
Tartar emetic, i.	255			Testæ cochleæ, i.	353
Tartar purified, ii.	450			Testæ ostræarum, i.	353
Tartar soluble, ii.	447			Testæ ovorum, i.	353
Tartar vitriolated, ii.	428				
Tartari crystalli, ii.	450				
Tartarized antimony, i.	255				
Tartarum antimoniatum, i.	255				

Testes equi, i.	698	Tinctura absinthii compo-		Tinctura antimonii alka-	
Testes of the horse, i.	678	sita, i.	29	lina, ii.	414
Tetranthera pichurim, ii.	680	Tinctura acetatis ferri,		Tinctura antimonii acris,	
Tetroxide of manganese,		i.	664	ii.	414
ii.	217	Tinctura acetatis ferri cum		Tinctura antimonii bal-	
Teucrium chamædrys, ii.	680	alcoholæ, i.	664	samica, ii.	414
Teucrium chamæpitys, ii.	681	Tinctura acida aromatica,		Tinctura antimonii nigra,	
Teucrium creticum, ii.	682	ii.	175	ii.	414
Teucrium flavum, ii.	681	Tinctura acidi sulphurici,		Tinctura balsamica com-	
Teucrium marum, ii.	682	ii.	175	posita, i.	318
Teucrium montanum, ii.	682	Tinctura aconiti, i.	85	Tinctura antimonii tarta-	
Teucrium scordium, ii.	682	Tinctura aconiti ætherea,		rizata, ii.	414
Teucrium scorodonia, ii.	683	i.	85	Tinctura antimonii sopo-	
Thalictrum flavum, ii.	685	Tinctura ætherea, i.	99	nata, i.	252
Thapsia asclepium, ii.	685	Tinctura ætherea campho-		Tinctura antirheumatis-	
Thea bohea, ii.	686	rata, i.	371	mal, i.	607
Thea viridis, ii.	686	Tinctura ætherea ferrata.		Tinctura aromatica, ii.	158,
Theobroma cacao, ii.	686	i.	659	170, 173	
Theriac, ii.	306	Tinctura ætherea guaiaci,		Tinctura aristolochiæ ser-	
Theriac officinal, ii.	116	i.	734	pentariæ, i.	278
Theriaca, ii.	306	Tinctura ætherea acetatis		Tinctura artemisiæ absin-	
Theriaca celestis Hoff-		ferri, i.	664	thii, i.	28
manni, ii.	319	Tinctura alceæ, ii.	217	Tinctura arthritica, ii.	143
Theriaca diatesseron, i.	761	Tinctura alcoholica aloes		Tinctura assafœtida, i.	305
Theriaca officialis, i.	761	composita, i.	129	Tinctura assafœtidæ æthe-	
Theriaca pauperum, i.	761	Tinctura alexipharmaca		rea, i.	305
Thistle blessed, i.	524	Huxhami, i.	501	Tinctura assafœtidæ am-	
Thistle carline, i.	396	Tinctura alexipharmaca		moniata, i.	177, 304
Thistle cotton, ii.	273	corticis Peruviani Hux-		Tinctura aurea ferrata, i.	659
Thistle dwarf carline, i.	396	hami, i.	501	Tinctura aurantii, i.	515
Thistle Fuller's, i.	618	Tinctura alkalina, ii.	414	Tinctura aurantii dulcis, i.	515
Thistle ladies', i.	395	Tinctura alkalina acris,		Tinctura aurantium	
Thistle milk, i.	395	ii.	414	ætherea, i.	511
Thistle sow, ii.	619	Tinctura aloes, i.	120	Tinctura auri, i.	313
Thistle woolly, i.	433	Tinctura aloes ætherea,		Tinctura balsami cana-	
Thlaspi arvense, ii.	688	i.	132	densis roborans, i.	318
Thlaspi bursa pastoris, ii.	688	Tinctura aloes aquosa, i.	120	Tinctura balsami Peru-	
Thlaspi campestre, ii.	688	Tinctura aloes composita,		viani, i.	320
Thlaspi sativum, ii.	688	i.	129	Tinctura balsami tolutani,	
Thorn apple, i.	605	Tinctura aloes crocata, i.	129	i.	324
Thorough-stem cresswort,		Tinctura aloes-cum myrrha,		Tinctura balsamica, i.	324,
i.	628	i.	130	ii.	629
Thorough-wax, i.	345	Tinctura aloes cum myrrha		Tinctura benzoës, i.	331
Thus, ii.	271	crocata, i.	129	Tinctura benzoini, i.	330
Thuya articulata, ii.	555	Tinctura aloetica alkalina,		Tinctura benzoini balsa-	
Thuya occidentalis, ii.	689	i.	131	mica, i.	331
Thyme alpine, ii.	689	Tinctura amara, i.	513, 714	Tinctura benzoini compo-	
Thyme cretan, ii.	558	Tinctura amara Halensis,		sita, i.	331
Thyme garden, ii.	689	i.	30	Tinctura bonplandiæ tri-	
Thyme mastic, ii.	689	Tinctura ambraë liquida,		foliatæ, i.	229
Thyme wild, ii.	690	i.	167	Tinctura botryos, i.	457
Thymi cretici herba, ii.	558	Tinctura ambraë moschata		Tinctura bruceæ, i.	229
Thymus alpinus, ii.	689	ætherea, i.	167	Tinctura Burgundica, i.	499
Thymus mastichina, ii.	689	Tinctura ammoniata aro-		Tinctura calami aroma-	
Thymus serpyllum, ii.	690	matica, i.	174	tici, i.	87
Thymus vulgaris, ii.	689	Tinctura amomi carda-		Tinctura calumbæ, i.	528
Tigllii grana, i.	570	moni, i.	394	Tinctura camphoræ, i.	372
Tigline, i.	570	Tinctura amomi repentis,		Tinctura cantharidis, i.	379
Tilii Europæa, ii.	691	i.	394	Tinctura cantharidum, i.	379
Tillii grana, i.	570	Tinctura amomi zingi-		Tinctura cantharidum	
Tin, ii.	623	beris, i.	205	ætherea, i.	382
Tin powdered, ii.	623	Tinctura angusturæ, i.	229	Tinctura cantharidum	
Tinctura absinthii, i.	28	Tinctura anodyna, ii.	277	composita, i.	280
Tinctura absinthii alka-		Tinctura volatilis anodyna,		Tinctura cantharidum	
lina, ii.	29	ii.	285	opiata, i.	381

Tinctura calami compo- sita, i.	88	Tinctura cinchonæ mar- tiatæ, i.	666	Tinctura digitalis semi- spirituosa, i.	617
Tinctura alcoholica can- tharidum camphorata, i.	380	Tinctura cinnamomi phosphorico acida, i.	73	Tinctura digitalis spiri- tuosa, i.	617
Tinctura cantharidum terebinthinata, i.	381	Tinctura coccinellæ sep- tempunctatæ, i.	526	Tinctura elastica, i.	647
Tinctura capsici, ii.	360	Tinctura coccionellæ, i.	530	Tinctura ellebori albi, ii.	715
Tinctura capsici annui, ii.	360	Tinctura cochleariæ, i.	534	Tinctura euphorbii, i.	756
Tinctura capsici et can- tharidum, ii.	361	Tinctura colchici autum- nalis, i.	539	Tinctura febrifuga acida, i.	756
Tinctura cascarillæ, i.	569	Tinctura colocynthidis anisata, i.	575	Tinctura ferrata, i.	671
Tinctura cascarillæ aquo- sa, i.	569	Tinctura colocynthidis Dalhbergi, i.	575	Tinctura ferri, i.	657
Tinctura cassiæ lignæ, ii.	172	Tinctura colocynthidis stibiata, i.	575	Tinctura ferri acetici æ- therea, i.	664
Tinctura castorei, i.	428	Tinctura colombar, i.	528	Tinctura ætherea ferri, i.	659
Tinctura castorei æthe- rea, i.	431	Tinctura colombo, i.	528	Tinctura ferri ammoni- ati, i.	663
Tinctura castorei compo- sita, i.	429	Tinctura conii maculati composita, i.	466	Tinctura ferri aurantia- ca, i.	667
Tinctura castorei fœti- da, i.	429	Tinctura convulvuli ja- lapæ, i.	552	Tinctura ferri muriati, i.	657
Tinctura castorei fœti- da ammoniacata, i.	429	Tinctura copaivæ balsa- mi, i.	555	Tinctura ferri muriatis, i.	657
Tinctura castorei kali- na, i.	428	Tinctura corallii, ii.	540	Tinctura ferri tartarisa- ta, i.	667
Tinctura castorei thebai- ca, i.	429	Tinctura corallinorum, i.	364	Tinctura vinosa ferri tar- tarisata, i.	669
Tinctura cardamomi, i.	394	Tinctura alcoholica cor- ticum aurantium	502	Tinctura ferulæ assafœ- tidæ, i.	306
Tinctura cardamomi composita, i.	395	Whytii, i.	502	Tinctura florum aquili- giæ, i.	270
Tinctura cardui benedic- ti, i.	525	Tinctura aquosa corti- cis, i.	488	Tinctura florum rosarum, ii.	518
Tinctura caryophyllatæ, i.	719	Tinctura corticis auran- tiorum, i.	515	Tinctura floræ salis am- moniaci martialium, i.	663
Tinctura centaurii mi- noris, i.	626	Tinctura corticis chinæ composita, i.	502	Tinctura fœtida pyro- oleosa, ii.	288
Tinctura chamæmeli, i.	451	Tinctura corticis Peru- viani, i.	500	Tinctura formicarum composita, i.	676
Tinctura chamomillæ, i.	451	Tinctura corticis Peru- viani composita, i.	502	Tinctura fuliginis alka- lina, i.	199
Tinctura chenopodii æ- therea, i.	457	Tinctura corticis Peru- viani volatilis, i.	500	Tinctura fuliginis Clau- deri, i.	199
Tinctura chenopodii am- brosioides, i.	457	Tinctura corticis quer- cus, ii.	475	Tinctura fuliginis fœti- da, i.	306
Tinctura chinæ, i.	500	Tinctura corticum winter- anorum, i.	620	Tinctura galbani, i.	684
Tinctura chinæ compo- sita, i.	502	Tinctura croci, i.	564	Tinctura galangæ, i.	682
Tinctura chinæ gingiva- li, i.	503	Tinctura crotonis cas- carillæ, i.	569	Tinctura gallarum, i.	693
Tinctura alcoholica chi- næ composita, i.	501	Tinctura crotonis eleu- theriæ, i.	569	Tinctura gambogiæ am- moniata, i.	696
Tinctura cicutæ ætherea, i.	466	Tinctura culilaban, ii.	178	Tinctura gei urbani, i.	719
Tinctura cinchonæ, i.	500	Tinctura cupri sulphuri- ci, i.	585	Tinctura gentianæ, i.	714
Tinctura cinchonæ æthe- rea composita, i.	503	Tinctura curcumæ, i.	581	Tinctura gentiana com- posita, i.	714
Tinctura cinchonæ ama- ra, i.	502	Tinctura Daffyi, i.	412	Tinctura gentianæ rubræ, i.	714
Tinctura cinchonæ am- moniata, i.	500	Tinctura delphinii, i.	608	Tinctura guaici, i.	734, 735, 736
Tinctura cinchonæ compo- sita, i.	501	Tinctura digitalis, i.	616	Tinctura guaiaci æthe- rea, i.	734
Tinctura cinchonæ lan- cifoliæ, i.	500	Tinctura digitalis æthe- rea, i.	617	Tinctura guaiaci ammo- niata, i.	736
		Tinctura digitalis aquo- so ætherea, i.	617	Tinctura guaiaci fœni- culata, i.	736
		Tinctura digitalis com- posita, i.	615		
		Tinctura digitalis pur- pureæ, i.	616		

<i>Tinctura guaiaci officinalis</i> , i.	735	<i>Tinctura malatis ferri</i> , i.	665	<i>Tinctura alcoholica myrrhæ uterina</i> , i.	306
<i>Tinctura guaiaci volatilis</i> , i.	736	<i>Tinctura mari veri æthereæ</i> , ii.	682	<i>Tinctura nervina Bestucheffii</i> , i.	659
<i>Tinctura gummi ammoniaci</i> , i.	744	<i>Tinctura matris perlarum</i> , i.	364	<i>Tinctura nervina Halensis</i> , i.	659
<i>Tinctura gummi guttæ ammoniata</i> , i.	696	<i>Tinctura martis</i> , i.	657, 665	<i>Tinctura nervino-tonica Lamotte</i> , i.	659
<i>Tinctura gummi guaiaci</i> , i.	735	<i>Tinctura martis aperiens</i> , i.	667, 669	<i>Tinctura nervino-tonica martialis</i> , i.	659
<i>Tinctura hæmatitis</i> , i.	663	<i>Tinctura martis aperitiva</i> , i.	670	<i>Tinctura mucis vomicæ</i> , ii.	267
<i>Tinctura Hassfieldi</i> , i.	737	<i>Tinctura martis aperitiva Zwelferi</i> , i.	664	<i>Tinctura oculorum cancrorum</i> , i.	364
<i>Tinctura helenii</i> , ii.	118	<i>Tinctura martis aurea</i> , i.	657	<i>Tinctura opii</i> , ii.	281
<i>Tinctura Hjærnæi contra tæniæ</i> , i.	123	<i>Tinctura martis cydoniata</i> , i.	665	<i>Tinctura alcoholica opii</i> , ii.	285
<i>Tinctura humuli</i> , ii.	11	<i>Tinctura martis helleborata</i> , i.	671	<i>Tinctura opii alcoholica saponacea</i> , ii.	286
<i>Tinctura humuli lupuli</i> , ii.	11	<i>Tinctura martis pomata</i> , i.	665	<i>Tinctura opii ammoniacalis</i> , ii.	285
<i>Tinctura hydragoga</i> , i.	659	<i>Tinctura martis salita</i> , i.	657	<i>Tinctura opii ammoniacata</i> , ii.	286
<i>Tinctura hyoscyami æthereæ</i> , i.	103	<i>Tinctura martis cum spiritu salis</i> , i.	657	<i>Tinctura opii ammoniata</i> , ii.	287
<i>Tinctura hyperici</i> , ii.	108	<i>Tinctura martis cum succo cydoniorum</i> , i.	665	<i>Tinctura opii benzoica</i> , ii.	287
<i>Tinctura alcoholica hyperici vulneraria</i> , ii.	108	<i>Tinctura martis cum succo pomorum</i> , i.	665	<i>Tinctura opii benzoica composita</i> , ii.	288
<i>Tinctura hypnotica</i> , ii.	284	<i>Tinctura martis tartariata</i> , i.	667, 669	<i>Tinctura opii camphorata</i> , ii.	287
<i>Tinctura jalapæ</i> , i.	552	<i>Tinctura martis tartariata Glauberi</i> , i.	670	<i>Tinctura opii crocata</i> , ii.	277
<i>Tinctura jalapæ confortans</i> , i.	552	<i>Tinctura martis tartariata Ludovici</i> , i.	670	<i>Tinctura opii Eccardi</i> , ii.	284
<i>Tinctura jalapæ composita</i> , i.	552	<i>Tinctura martis cum vino malvatico et pomis auratis</i> , i.	667	<i>Tinctura opii simplex</i> , ii.	276, 281, 284
<i>Tinctura jalapæ resinæ</i> , i.	552	<i>Tinctura martis vinosa</i> , i.	671	<i>Tinctura opii spirituosa</i> , ii.	281
<i>Tinctura jalappæ</i> , i.	552	<i>Tinctura mastiches</i> , ii.	224	<i>Tinctura opii vinosa</i> , ii.	276
<i>Tinctura iodii</i> , ii.	120	<i>Tinctura mastichina</i> , ii.	224	<i>Tinctura opii vinosa Sydenhami</i> , ii.	277
<i>Tinctura iodii æthereæ</i> , ii.	121	<i>Tinctura menthæ piperitæ</i> , ii.	236	<i>Tinctura oreosolini</i> , ii.	581
<i>Tinctura ipecacuanhæ</i> , ii.	128	<i>Tinctura menthæ viridis</i> , ii.	236	<i>Tinctura papaveris rhæados</i> , ii.	336
<i>Tinctura kalina</i> , ii.	414	<i>Tinctura metallorum</i> , ii.	414	<i>Tinctura phellandri aquatici</i> , i.	469
<i>Tinctura kinkinæ amara</i> , i.	502	<i>Tinctura millefolii</i> , i.	36	<i>Tinctura alcoholica picis græcæ composita</i> , i.	129
<i>Tinctura laccæ aquosa</i> , ii.	150	<i>Tinctura moschata</i> , ii.	252	<i>Tinctura pimentæ sulphurica</i> , ii.	361
<i>Tinctura laccæ kalica</i> , ii.	151	<i>Tinctura moschi</i> , ii.	252	<i>Tinctura pimpinellæ albæ</i> , ii.	357
<i>Tinctura lactucæ verosæ</i> , ii.	153	<i>Tinctura moschi artificialis</i> , ii.	631	<i>Tinctura pimpinellæ composita</i> , ii.	357
<i>Tinctura lactucarii</i> , ii.	154	<i>Tinctura moschi vinosæ</i> , ii.	252	<i>Tinctura pini</i> , i.	22
<i>Tinctura lavandulæ composita</i> , ii.	158	<i>Tinctura muriatis ferri cum oxydo rubro</i> , i.	658	<i>Tinctura pini composita</i> , i.	22
<i>Tinctura laudanum</i> , ii.	277	<i>Tinctura æthereæ alcoholica de muriate ferri</i> , i.	659	<i>Tinctura plumbosa</i> , ii.	378
<i>Tinctura laxativa</i> , i.	120	<i>Tinctura myrrhæ</i> , i.	762	<i>Tinctura pomorum aurantium immaturorum</i> , i.	515
<i>Tinctura ligni aloes</i> , ii.	192	<i>Tinctura alcoholica myrrhæ</i> , i.	762	<i>Tinctura populi</i> , ii.	409
<i>Tinctura ligni sassafras</i> , ii.	183	<i>Tinctura myrrhæ alkalisata</i> , i.	763	<i>Tinctura proprietatis Mynsichti</i> , ii.	356
<i>Tinctura lignorum</i> , i.	22, 735	<i>Tinctura myrrhæ aquosa</i> , i.	762	<i>Tinctura quassia</i> , ii.	473
<i>Tinctura lobeliæ inflatæ</i> , ii.	203				
<i>Tinctura longæ vitæ</i> , i.	138				
<i>Tinctura lupuli</i> , ii.	11				
<i>Tinctura lupuli strobilorum</i> , ii.	11				
<i>Tinctura lyttæ vesicatoriæ</i> , i.	379, 382				
<i>Tinctura macidis</i> , ii.	255				

<i>Tinctura quassiae excelsae</i> , ii. 474	<i>Tinctura sedativa</i> , ii. 281, 284	<i>Tinctura vinosa opii an-</i> <i>ticolica</i> , ii. 285
<i>Tinctura raphani compo-</i> <i>sita</i> , ii. 483	<i>Tinctura sedativa campho-</i> <i>rata</i> , ii. 287	<i>Tinctura pro vitam pro-</i> <i>ducenda</i> , i. 138
<i>Tinctura ratanhiæ</i> , ii. 148	<i>Tinctura seminum col-</i> <i>chici</i> , i. 539	<i>Tinctura vulneraria sim-</i> <i>plex</i> , ii. 224
<i>Tinctura ratanhiæ compo-</i> <i>sita</i> , ii. 148	<i>Tinctura seminum col-</i> <i>chici vinosa Williamsi</i> , i. 539	<i>Tinctura zingiberis</i> , i. 205
<i>Tinctura regia</i> , i. 167, ii. 170	<i>Tinctura senegæ</i> , ii. 404	<i>Tincture of acetate of iron</i> , i. 664
<i>Tinctura resinæ guaiaci</i> , i. 735	<i>Tinctura sennæ</i> , i. 410	<i>Tincture æthereal</i> , i. 371
<i>Tinctura resinæ succini</i> , ii. 631	<i>Tinctura sennæ aroma-</i> <i>tica</i> , i. 410	<i>Tincture æthereal of aco-</i> <i>nite</i> , i. 85
<i>Tinctura rhabbarbari</i> , ii. 497, 501	<i>Tinctura sennæ compo-</i> <i>sita</i> , i. 410	<i>Tincture alcoholic of æo-</i> <i>nite</i> , i. 85
<i>Tinctura rhabbarbari vi-</i> <i>nosa</i> , ii. 500	<i>Tinctura serpentariæ</i> , i. 278	<i>Tincture of alcea</i> , ii. 217
<i>Tinctura rhei</i> , ii. 497	<i>Tinctura stibiata</i> , ii. 288	<i>Tincture of aloe wood</i> , ii. 192
<i>Tinctura rhei et aloes</i> , i. 138, ii. 502	<i>Tinctura stibii alkalizata</i> , ii. 414	<i>Tincture of aloes</i> , i. 120
<i>Tinctura rhei amara</i> , ii. 502	<i>Tinctura stomachica</i> , i. 513, 673	<i>Tincture of aloes and</i> <i>myrrh</i> , i. 129
<i>Tinctura rhei aquosa</i> , ii. 494	<i>Tinctura stomachica acu-</i> <i>ta Ernstringii</i> , ii. 361	<i>Tincture of aloes myrrh</i> <i>and saffron</i> , i. 129
<i>Tinctura rhei composita</i> , ii. 501, 502	<i>Tinctura stomachica Lip-</i> <i>siensis</i> , i. 174	<i>Tincture æthereal of aloes</i> , i. 132
<i>Tinctura rhei Darelli</i> , ii. 500	<i>Tinctura stomachica mar-</i> <i>tialis</i> , i. 673	<i>Tincture æthereal of am-</i> <i>ber</i> , ii. 629
<i>Tinctura rhei dulcis</i> , ii. 500	<i>Tinctura stramonii</i> , i. 606	<i>Tincture alcoholic of am-</i> <i>ber</i> , ii. 629
<i>Tinctura rhei et gentianæ</i> , ii. 502	<i>Tinctura stramonii æthe-</i> <i>rea</i> , i. 607	<i>Tincture alkaline of am-</i> <i>ber</i> , ii. 630
<i>Tinctura rhei spirituosa</i> , ii. 501	<i>Tinctura strychnos</i> , ii. 267	<i>Tincture fetid of amber</i> , ii. 632
<i>Tinctura rhei vinosa</i> , ii. 500	<i>Tinctura succini ætherea</i> , ii. 629	<i>Tincture of ambergris</i> , i. 167
<i>Tinctura rhododendri chry-</i> <i>santhi</i> , ii. 503	<i>Tinctura succini balsa-</i> <i>mica</i> , ii. 631	<i>Tincture æthereal of am-</i> <i>bergris</i> , i. 167
<i>Tinctura rhois radicantis</i> , ii. 504	<i>Tinctura succini alcoho-</i> <i>lica</i> , ii. 629	<i>Tincture alkaline of am-</i> <i>bergris</i> , i. 167
<i>Tinctura roborans</i> , i. 501, 715, ii. 173	<i>Tinctura succini alkali-</i> <i>sata</i> , ii. 630	<i>Tincture balsamic of am-</i> <i>bergris</i> , i. 167
<i>Tinctura rosarum acidula</i> , ii. 518	<i>Tinctura tanacetii</i> , ii. 650	<i>Tincture balsamic of am-</i> <i>bergris Dippel's</i> , i. 167
<i>Tinctura sabinæ</i> , ii. 143	<i>Tinctura tartari</i> , ii. 414	<i>Tincture of ambergris</i> <i>Mindererus'</i> , i. 167
<i>Tinctura sabinæ kalina</i> , ii. 143	<i>Tinctura tartratis potassæ</i> <i>ferrici</i> , i. 669	<i>Tincture of angelica</i> , i. 226
<i>Tinctura sacra</i> , i. 119	<i>Tinctura terebinthinæ</i> , ii. 655	<i>Tincture ammoniated of</i> <i>angelica</i> , i. 223
<i>Tinctura salina compo-</i> <i>sita</i> , i. 715	<i>Tinctura thebaica</i> , ii. 284	<i>Tincture of angustura</i> , i. 229
<i>Tinctura salis ammoniaci</i> <i>cuprifera</i> , i. 589	<i>Tinctura thebaica genuina</i> , ii. 276	<i>Tincture anodyne</i> , ii. 710
<i>Tinctura salis tartari</i> , ii. 414	<i>Tinctura theriacalis</i> , ii. 314	<i>Tincture anti-arthritis</i> , ii. 425
<i>Tinctura sanguinaris can-</i> <i>adensis</i> , ii. 555	<i>Tinctura toluiferæ bal-</i> <i>sami</i> , i. 324	<i>Tincture anti-epileptic</i> , i. 542
<i>Tinctura sanguis draconis</i> , i. 765	<i>Tinctura tolutana</i> , i. 324	<i>Tincture antigonorrhœic</i> , ii. 437
<i>Tinctura saponata</i> , ii. 460	<i>Tinctura tonica martialis</i> <i>lutea</i> , i. 658	<i>Tincture antihysterical</i> , i. 430
<i>Tinctura saponis campho-</i> <i>rata</i> , i. 179, ii. 461	<i>Tinctura tormentillæ</i> , ii. 693	<i>Tincture anti-odontalgic</i> , i. 231
<i>Tinctura saponis et opii</i> , ii. 286	<i>Tinctura valerianæ</i> , ii. 708	<i>Tincture Gramann's anti-</i> <i>phthisical</i> , i. 666, ii. 388
<i>Tinctura saturnino-mar-</i> <i>tialis</i> , ii. 387	<i>Tinctura valerianæ æthe-</i> <i>rea</i> , ii. 710	<i>Tincture antispasmodic</i> , i. 306, ii. 710
<i>Tinctura sarcollæ</i> , i. 765	<i>Tinctura valerianæ ammo-</i> <i>niata</i> , ii. 709	<i>Tincture Chrestien's an-</i> <i>ti-spasmodic</i> , ii. 282, 285
<i>Tinctura scammonii</i> , ii. 562	<i>Tinctura valerianæ cam-</i> <i>phorata</i> , ii. 710	<i>Tincture Dumas' antispas-</i> <i>modic</i> , ii. 285
<i>Tinctura scillæ</i> , ii. 571	<i>Tinctura valerianæ com-</i> <i>posita</i> , ii. 708	<i>Tincture antispasmodic</i> <i>and emmenagogue</i> , i. 129
<i>Tinctura scillæ kalina</i> , ii. 572	<i>Tinctura vanigliæ</i> , ii. 712	
<i>Tinctura scordii</i> , ii. 684	<i>Tinctura veratri albi</i> , ii. 715	
<i>Tinctura scordii compo-</i> <i>sita</i> , ii. 684		

- Tincture antispasmodic and tonic, ii. 709
 Tincture antisyphilitic, ii. 71
 Tincture Besnard's antisyphilitic, ii. 425
 Tincture Falk's antisyphilitic, ii. 75
 Tincture Talbot's antisyphilitic, ii. 425
 Tincture compound of ants, i. 676
 Tincture aperitive, i. 56, ii. 499
 Tincture aromatic, ii. 170
 Tincture compound aromatic, ii. 156
 Tincture of artificial musk, ii. 631
 Tincture of asarum, i. 295
 Tincture of assafoetida, i. 305
 Tincture æthereal of assafoetida, i. 305
 Tincture alkaline of assafoetida, i. 306
 Tincture antimoniated of assafoetida, i. 252
 Tincture fuliginous of assafoetida, i. 306
 Tincture astringent, i. 73, ii. 387, 480
 Tincture of avens, i. 719
 Tincture of balm, ii. 229
 Tincture compound balsamic, i. 555
 Tincture of Canadian balsam, i. 318
 Tincture of balsam of Mecca, i. 319
 Tincture of Peruvian balsam, i. 320
 Tincture of balsam of tolu, i. 324
 Tincture æthereal of balsam of tolu, i. 325
 Tincture of belladonna, i. 311
 Tincture of benzoin, i. 330
 Tincture compound of benzoin, i. 331
 Tincture Ludwig's bezoardic, i. 226
 Tincture bitter, i. 30
 Tincture of black pepper, ii. 360
 Tincture of blessed thistle, i. 525
 Tincture of bloodroot, ii. 555
 Tincture Burgundic, i. 499
 Tincture of burnet saxifrage, ii. 357
 Tincture alkaline of burnet saxifrage, ii. 357
 Tincture compound of burnet saxifrage, ii. 357
 Tincture of calumba, i. 528
 Tincture of cantharides, i. 379, 381
 Tincture æthereal of cantharides, i. 382
 Tincture camphorated of cantharides, i. 380
 Tincture æthereous camphorated of cantharides, i. 382
 Tincture compound of cantharides, i. 380
 Tincture compound æthereous of cantharides, i. 382
 Tincture of capsicum, ii. 360
 Tincture of capsicum and of cantharides, ii. 361
 Tincture of cardamoms, i. 394
 Tincture compound of cardamoms, i. 395
 Tincture of cascarilla, i. 569
 Tincture alcoholic and alkaline of cascarilla, i. 569
 Tincture aqueous of cascarilla, i. 569
 Tincture of castor, i. 428
 Tincture æthereal of castor, i. 431
 Tincture alkaline of castor, i. 428
 Tincture compound of castor, i. 429
 Tincture opiate of castor, i. 429
 Tincture of lesser centaury, i. 626
 Tincture Ludwig's chalybeate, i. 670
 Tincture of chamomile, i. 451
 Tincture Cheston's, i. 440
 Tincture ammoniated of cinchona, i. 500
 Tincture compound of cinchona, i. 501
 Tincture ferruginous of cinchona, i. 666
 Tincture of grey cinchona, i. 500
 Tincture of red cinchona, i. 500
 Tincture potassated of cinchona, i. 501
 Tincture of yellow cinchona, i. 500
 Tincture of cinchonine, i. 504
 Tincture of cinnamon, ii. 172
 Tincture æthereous of cinnamon, ii. 175
 Tincture compound of cinnamon, ii. 173
 Tincture of cloves, i. 400
 Tincture of coccinella, i. 526
 Tincture of cochineal, i. 530
 Tincture of colchicum bulb, i. 539
 Tincture of colchicum seeds, i. 539
 Tincture of colocynth, i. 575
 Tincture anisated of colocynth, i. 575
 Tincture antimoniated of colocynth, i. 575
 Tincture crocated of colocynth, i. 576
 Tincture of columbine, i. 270
 Tincture alcoholic of copaiba, i. 555
 Tincture alkaline of copaiba, i. 555
 Tincture of copper, i. 587
 Tincture corroborant, i. 715
 Tincture cosmetic, i. 331
 Tincture of culilawan, ii. 178
 Tincture of daisy, i. 329
 Tincture of the resin of daphne gnidium, i. 604
 Tincture Darel's, ii. 500
 Tincture of digitalis, i. 616
 Tincture æthereal of digitalis, i. 617
 Tincture compound of digitalis, i. 615
 Tincture Flittner's of digitalis, i. 617
 Tincture Fowler's aqueous of digitalis, i. 614
 Tincture Harless' of digitalis, i. 617
 Tincture Maclean's of digitalis, i. 616
 Tincture Planche's of digitalis, i. 616
 Tincture Remer's of digitalis, i. 615
 Tincture Trommsdorff's of digitalis, i. 617
 Tincture diuretic, i. 617
 Tincture Eccard's, ii. 284
 Tincture of dragon's blood, i. 765
 Tincture of elecampane, ii. 118
 Tincture febrifuge, i. 288
 Tincture of euphorbium, i. 756
 Tincture Clutton's febrifuge, i. 226
 Tincture Mutis' febrifuge, i. 502
 Tincture of fecula of henbane, ii. 101
 Tincture ferruginous, i. 671
 Tincture fetid, i. 306
 Tincture of fraxinella, i. 611
 Tincture of galanga, i. 682
 Tincture of galbanum, i. 684
 Tincture compound of galbanum, i. 684

Tincture of gamboge, i.	696	Tincture aromatic of jalap, i.	552	Tincture of myrrh, i.	762
Tincture alkaline of gamboge, i.	696	Tincture compound of jalap, i.	552	Tincture alkaline of myrrh, i.	763
Tincture ammoniacal of gamboge, i.	696	Tincture of iodine, ii.	120	Tincture of myrrh phosphoric, i.	73
Tincture of gentian, i.	714	Tincture æthereal of iodine, ii.	121	Tincture nervine and tonic, ii.	709
Tincture alkaline of gentian, i.	715	Tincture of ipecacuanha, ii.	129	Tincture of nitric oil of camphor, i.	374
Tincture compound of gentian, i.	714	Tincture anisated of ipecacuanha, ii.	130	Tincture of nutgalls, i.	693
Tincture compound alkaline of gentian, i.	715	Tincture aperitive of iron, i.	667	Tincture of nux vomica, ii.	267
Tincture of gentianine, i.	716	Tincture of iron and of orange, i.	667	Tincture of oak bark, ii.	475
Tincture alkaline of gentianine, i.	716	Tincture of kermes, i.	530	Tincture ophthalmic, i.	264
Tincture of Geoffræa, i.	717	Tincture of kino, ii.	147	Tincture acetose of opium, ii.	275
Tincture of ginger, i.	205	Tincture alcoholic of lac, ii.	151	Tincture alcoholic of opium, ii.	281
Tincture of gold, i.	313	Tincture alkaline of lac, ii.	151	Tincture aromatic alcoholic of opium, ii.	284
Tincture of guaiac and fennel, i.	736	Tincture aluminated of lac, ii.	151	Tincture ammonical and camphorated of opium, ii.	285
Tincture ammoniacal of guaiac, i.	736	Tincture aqueous of lac, ii.	150	Tincture compound alcoholic of opium, ii.	285
Tincture saponaceous of guaiac, i.	737	Tincture compound of lac, ii.	151	Tincture crocated alcoholic of opium, ii.	285
Tincture Dewee's volatile of guaiac, i.	737	Tincture of lactucarium, ii.	154	Tincture crocated vinous of opium, ii.	277
Tincture of guaiac resin, i.	735	Tincture of larkspur, i.	608	Tincture vinous of opium, ii.	276
Tincture compound of guaiac resin, i.	736	Tincture of leopard's bane, i.	281	Tincture aromatic vinous of opium, ii.	277
Tincture of guaiac wood, i.	734	Tincture æthereal of leopard's bane, i.	281	Tincture alcoholic of opium and of benzoate of ammonia, ii.	287
Tincture compound of guaiac wood, i.	735	Tincture of viroous lettuce, ii.	153	Tincture camphorated alcoholic of opium and of benzoic acid, ii.	287
Tincture of gum ammoniac, i.	744	Tincture of lovage, ii.	197	Tincture of orange peel, i.	515
Tincture alkaline of gum ammoniac, i.	744	Tincture of lupuline, ii.	12	Tincture of sweet orange peel, i.	515
Tincture for the gums, i.	503	Tincture of mace, ii.	255	Tincture of unripe orange, i.	515
Tincture of white hellebore, ii.	715	Tincture of malate of iron, i.	665	Tincture of pareira brava, i.	505
Tincture ferruginous of hellebore, i.	671	Tincture æthereous of marum, ii.	682	Tincture physagogue, i.	511
Tincture æthereal of hemlock, i.	466	Tincture of mastic, ii.	224	Tincture æthereal of pimenta, ii.	361
Tincture alcoholic of hemlock, i.	466	Tincture compound of mastic, ii.	224	Tincture of peppermint, ii.	236
Tincture compound of hemlock, i.	466	Tincture of mentha crispay, ii.	236	Tincture of poplar, ii.	409
Tincture of water hemlock, i.	469, ii. 346	Tincture of Mexican tea, i.	457	Tincture aqueous of poppy flowers, ii.	336
Tincture of henbane seeds, ii.	103	Tincture of Mexican tea, ii.	457	Tincture aqueous of red poppy, ii.	336
Tincture æthereal of henbane, ii.	103	Tincture of milfoil, i.	36	Tincture of proto-hydrochlorate of iron, i.	658
Tincture alcoholic of henbane, ii.	103	Tincture of green mint, ii.	236	Tincture æthereous of pyrethrum, ii.	232
Tincture compound of horse radish, ii.	483	Tincture of muriate of iron, i.	657	Tincture alcoholic of quassia, ii.	473
Tincture hydragogue, ii.	140	Tincture æthereal of muriate of iron, i.	659	Tincture vinous of quassia, ii.	473
Tincture of hydrochlorate of ammonia and iron, i.	663	Tincture of musk, ii.	252		
Tincture of jalap, i.	552	Tincture æthereal of musk, ii.	252		
		Tincture vinous of musk, ii.	252		
		Tincture Mynsicht's, ii.	356		

Tincture of ratany, ii.	148	Tincture of St. John's-wort, ii.	108	Tincture camphorated æthereal of valerian, ii.	710
Tincture acetose of ratany, ii.	148	Tincture compound of St. John's-wort, ii.	108	Tincture compound of valerian, ii.	708
Tincture aqueous of ratany, ii.	148	Tincture compound vinous of St. John's-wort, ii.	108	Tincture compound ammoniated of valerian, ii.	709
Tincture compound of ratany, ii.	148	Tincture stomachic, i.	174, 516, ii.	Tincture of vanilla, ii.	712
Tincture of resin of jalap, i.	552	Tincture aromatic stomachic, ii.	174	Tincture of veratrine, ii.	713
Tincture of resin of indigo, ii.	352	Tincture Ernsting's stomachic, ii.	361	Tincture vermifuge, ii.	288
Tincture of rhododendron, ii.	503	Tincture ferruginous stomachic, i.	671	Tincture of walnut rind, ii.	136
Tincture of rhubarb and aloes, i.	138, ii. 502	Tincture Lentin's stomachic, i.	88	Tincture of Winter's bark, i.	620
Tincture alcoholic of rhubarb, ii.	501	Tincture æthereal of stramonium, i.	607	Tincture of woodlice, ii.	273
Tincture alkaline aqueous of rhubarb, ii.	497	Tincture alcoholic of stramonium, i.	606	Tincture of wormwood, i.	28
Tincture aromatic alcoholic of rhubarb, ii.	501	Tincture vinous of stramonium, i.	606	Tincture alkaline of wormwood, i.	29
Tincture bitter of rhubarb, ii.	502	Tincture strengthening, ii.	173	Tincture compound of wormwood, i.	29
Tincture saline aqueous, of rhubarb, ii.	498	Tincture of strychnine, ii.	626	Tinea flavi frontella, ii.	377
Tincture of roses, ii.	516	Tincture phosphorated of subcarbonate of potass, ii.	349	Tinkal, ii.	608
Tincture acid of roses, ii.	518	Tincture of sulphate of copper, i.	585	Tinker-weed, ii.	696
Tincture of saffron, i.	564	Tincture of sulphate of quinine, ii.	479	Tithymali cyparissiae radix, i.	629
Tincture saline, ii.	415	Tincture of sulphur, ii.	637	Tithymali latifolii semines, i.	629
Tincture of sarcocolla, i.	765	Tincture of sulphuro-prussiate of iron, i.	675	Toad-flax, ii.	199
Tincture of sassafras, ii.	183	Tincture of swallow-wort, i.	296	Toads dried, i.	698
Tincture compound of sassafras, ii.	184	Tincture of sweet flag, i.	87	Tobacco, ii.	200
Tincture of savine, ii.	143	Tincture compound of sweet flag, i.	88	Tobacco Indian, ii.	203
Tincture alkaline of savine, ii.	143	Tincture against tænia, i.	123	Tomata, ii.	617
Tincture of scammony, ii.	562	Tincture of tansey, ii.	650	Tonka-bean tree, i.	562
Tincture of scordium, ii.	684	Tincture of tartrate of potass and of iron, i.	667	Toothache tree, i.	270
Tincture of scurvy-grass, i.	534	Tincture theriacal, ii.	314	Topasius, i.	157
Tincture antimoniated sedative, ii.	288	Tincture of tobacco, ii.	263	Topaz, i.	157
Tincture of seneka, ii.	404	Tincture of Indian tobacco, ii.	203	Tordylium officinale, ii.	692
Tincture of senna, i.	410	Tincture of tolu, i.	324	Tormentil, ii.	692
Tincture aromatic of senna, i.	412	Tincture tonic, i.	488, 501	Tormentilla erecta, ii.	692
Tincture compound of senna, i.	410	Tincture of tormentil, ii.	693	Tornasolis cærulea, ii.	693
Tincture of serpentaria, i.	278	Tincture of toxicodendron, ii.	504	Totæ sanæ herba, ii.	107
Tincture of soap and of opium, ii.	286	Tincture of trito-hydrochlorate of iron, i.	658	Touchwood boletus, i.	340
Tincture of spignel, ii.	581	Tincture of turmeric, i.	581	Toxicodendron, ii.	503
Tincture of spruce fir, i.	22	Tincture of turpentine, ii.	655	Tragacanth, i.	766
Tincture compound of spruce fir, i.	22	Tincture of valerian, ii.	70	Tragacantha, i.	766
Tincture of squill, ii.	571	Tincture æthereal of valerian, ii.	710	Tragea alba, i.	750
Tincture alkaline of squill, ii.	572	Tincture ammoniated of valerian, ii.	709	Tragea aromatica, ii.	161
Tincture aromatic of squill, ii.	571	Tincture camphorated of valerian, ii.	708	Tragea granorum actes, ii.	552
Tincture compound of squill, ii.	571			Tragea ophthalmica, i.	630, ii. 353
Tincture stimulant, i.	380			Tragenata de semine santonico, ii.	583

- Trichicus manatus*, i. 338
Trifolii acetosi herba, ii. 327
Trifolii aurei herba, ii. 4
Trifolii cervini radix, i. 628
Trifolium pratense, ii. 695
Trigonella fœnum græcum, ii. 696
Trinitatis herba, ii. 723
Tritestum perfoliatum, ii. 696
Tripela, ii. 696
Tripoli, ii. 696
Trissaginis herba, ii. 681
Tritici fagini semen, ii. 407
Triticum repens, ii. 696
Trito-acetate of iron, ii. 663
Trito-hydrochlorate of iron, i. 657
Trito-hydroferrocyanate of iron, i. 674
*Trito-hydrocyanate ferru-
retted of iron*, i. 674
Trito-phosphate of iron, i. 651
Tritoxide of antimony, i. 240
Tritoxide of iron, i. 643
Tritoxide of lead, ii. 371
Tritoxidum antimonii, i. 240
Trochisci de agarico, ii. 408
Trochisci albi, ii. 372
Trochisci alhandali, i. 571
Trochisci aliptæ moschata, ii. 251
Trochisci alkekengi, ii. 350
Trochisci althææ, i. 146
*Trochisci althææ compo-
siti*, i. 146
Trochisci amyli, i. 213
Trochisci antacidi, ii. 209
Trochisci anthelmintici, ii. 65, 583
*Trochisci anthelmintici
doctoris ching*, ii. 51
Trochisci aromatici, ii. 163
*Trochisci aromatico sto-
machici*, ii. 163
Trochisci bechici albi, i. 146, 213, 725, 726, 750, ii. 132
Trochisci bechici nigri, i. 726
*Trochisci calami aroma-
tici*, i. 86
Trochisci calcis carbonatis, i. 357
*Trochisci carbonatis mag-
nesiæ*, ii. 209
*Trochisci carbonatis plum-
bi camphorati*, ii. 372
Trochisci citratis ferrici, i. 667
*Trochisci fructuum colo-
cynthidis*, i. 571
*Trochisci colocynthidis
præparatæ*, i. 571
*Trochisci de cremore tar-
tari*, ii. 451
*Trochisci fructuum cu-
pressi*, i. 580
Trochisci cypheos, ii. 663
Trochisci fumales, i. 334
Trochisci galliæ moschata, ii. 251
Trochisci glycyrrhizæ, i. 725, 726
*Trochisci glycyrrhizæ
glabræ*, i. 726
*Trochisci glycyrrhizæ cum
opio*, i. 727, ii. 297
*Trochisci glycyrrhizæ the-
baici*, i. 727
Trochisci gummi acaciæ, i. 750
Trochisci gummosi, i. 750
Trochisci hedychroi, i. 86
Trochisci hermetici, ii. 60
*Trochisci hermetici se-
cretæ*, ii. 65
Trochisci hydrargyri, ii. 85
Trochisci imperatoris, ii. 163
Trochisci liquiritiæ, i. 726
*Trochisci liquiritiæ cum
gummi*, i. 726
Trochisci magnesiæ, ii. 209
Trochisci de minio, ii. 77
*Trochisci menthæ pipe-
ritæ*, ii. 237
*Trochisci muriatis oxy-
genati hydrargyri*, ii. 73
Trochisci nitratis potassæ, ii. 434
Trochisci de nitro, ii. 434
Trochisci pectorales, i. 750
*Trochisci pectorales regis
Danorum*, i. 727
Trochisci refrigerantes, ii. 439
Trochisci Rhazis, ii. 372
*Trochisci de scilla Damo-
cratis*, ii. 564
Trochisci scillitici, ii. 564
*Trochisci sedativo-balsa-
mici*, i. 727
Trochisci stibiati, i. 242
Trochisci stomachici, ii. 163
Trochisci e sulphure, ii. 635
Trochisci temperantes, ii. 439
Trochisci contra vermes, ii. 60
Trochisci viperarum, i. 699
Trochiscs absorbent, ii. 109
Trochiscs of agaric, ii. 408
Trochiscs alhandal, i. 571
Trochiscs of amber, ii. 303
Trochiscs of ambergris, i. 166
Trochiscs anthelmintic, i. 545, 653, ii. 65, 583
Trochiscs antipsoric, ii. 636
*Trochiscs of carbonate
of lime*, i. 357
Trochiscs of castor, i. 425
Trochiscs of cinchona, i. 476
Trochiscs cypheos, ii. 663
Trochiscs of cypress, i. 580
Trochiscs diarrhodon, ii. 511
Trochiscs digestive, ii. 608
Trochiscs escharotic, ii. 73, 77
Trochiscs Gordon's, i. 751
Trochiscs of gum arabic, i. 750
Trochiscs of ipecacuanha, ii. 126
*Trochiscs compound of
ipecacuanha*, ii. 126
*Trochiscs opiate of ipeca-
cuanha*, ii. 126
Trochiscs of iris, ii. 132
*Trochiscs citronated of
of iron*, i. 667
Trochiscs kermesine, i. 531
Trochiscs Keyser's, ii. 96
Trochiscs of manna, ii. 218
Trochiscs manus christi, ii. 540
Trochiscs masticatory, ii. 223
*Trochiscs compound of
myrrh*, i. 302
Trochiscs mercurial, ii. 85
Trochiscs of mercury, ii. 16
Trochiscs mogul, ii. 251
Trochiscs musk, ii. 251
Trochiscs black pectoral, i. 726
Trochiscs opiate pectoral, i. 727, ii. 297
Trochiscs red pectoral, i. 768
Trochiscs white pectoral, i. 725
Trochiscs yellow pectoral, i. 726, 750
Trochiscs of peppermint, ii. 237
Trochiscs refrigerant, ii. 439
Trochiscs of rhubarb, ii. 493
Trochiscs of squill, ii. 564
Trochiscs of starch, i. 213
Trochiscs of sulphur, ii. 635
*Trochiscs of supertartrate
of potass*, ii. 451
Trochiscs temperant, ii. 446
Trochiscs of viper, i. 699
Trochiscs Rhazes' white, ii. 372
Trochiscs of winter cherry, ii. 350
Troæolum majus, ii. 698
Tryphera magna, ii. 516
Tubera cervina, ii. 577
Tulip-tree Virginian, ii. 202
Tunicæ flores, i. 610
Turmeric, i. 581
Turnerie round, i. 581
Turnip Indian, i. 292
Turnsol, ii. 1, 693
Turpethi albi radix, i. 554
*Turpethi vegetabilis ra-
dix*, i. 554
Turpethum minerale, ii. 82
Turpethum nigrum, ii. 84

Turpethum rubrum, ii.	81	Vincetoxici radix, i.	295	Vinum aloetico-alkalinum, i.	133
Turpentine, ii.	653	Vinegar compound of angelica, i.	226	Vinum aloes, i.	119
Tussilago farfara, ii.	699	Vinegar anti-emetic, i.	115	Vinum aloes socotorinæ, i.	119
Tussilago petasites, ii.	700	Vinegar antihysteria, i.	432	Vinum aloeticum, i.	119
Tyllelini radix, ii.	581	Vinegar aperient, i.	412	Vinum amaro-chalybeatum, i.	672
Vaccinium myrtillus, ii.	701	Vinegar aromatic, i.	44, 133, 231, 523	Vinum amarum, i.	713
Vaccinium oxycoccos, ii.	702	Vinegar aromatic with garlic, i.	115	Vinum amarum alkalisatum, i.	513
Vaccinium vitis idææ, ii.	702	Vinegar camphorated, i.	373	Vinum amarum diureticum, ii.	570
Valantia aparine, i.	691	Vinegar of capsicum, ii.	361	Vinum amarum scilliticum compositum, ii.	570
Valerian garden, ii.	703	Vinegar of belladonna, i.	311	Vinum amarum cum spiritu vini, i.	713
Valerian wild, ii.	703	Vinegar of clove-pink, i.	610	Vinum anthelminticum, i.	27
Valeriana celtica, ii.	703	Vinegar of coffee, i.	537	Vinum antimoniale, i.	258
Valeriana phu, ii.	703	Vinegar of colchicum, i.	539	Vinum antimoniatum, i.	258
Valerianæ officinalis, ii.	703	Vinegar concentrated by freezing, i.	38	Vinum antimonii, i.	258
Valerianella olitoria, ii.	711	Vinegar of convallaria, i.	542	Vinum armoraciæ, ii.	483
Vanigliæ siliquæ, ii.	712	Vinegar of cowslip, ii.	465	Vinum aromo-chalybeatum, i.	672
Vanilla aromatica, ii.	712	Vinegar crude, i.	38	Vinum aurantiorum compositum, i.	512
Vapor aceticus, i.	42	Vinegar of elder, ii.	553	Vinum calami aromatici, i.	88
Vapor hepaticus, i.	65	Vinegar compound of the four thieves, i.	116	Vinum calidum, ii.	171
Vapour hydro-sulphureous, i.	65	Vinegar of garlic, i.	115	Vinum chalybeate, i.	671
Variolaria orcina, ii.	509	Vinegar glacial, i.	41	Vinum chalybeatum compositum, i.	673
Ventriculus gallinæ, i.	699	Vinegar of white hellebore, ii.	715	Vinum chinæ amarum, i.	498
Veratrina, ii.	713	Vinegar of horseradish, ii.	483	Vinum chinæ martiatum, i.	497
Veratrine, ii.	713	Vinegar of lavender, ii.	156	Vinum chinæ compositum, i.	498
Veratrinum, ii.	713	Vinegar compound of lavender, ii.	156	Vinum chinatum, i.	496
Veratrum album, ii.	714	Vinegar lythargyrate, ii.	376	Vinum cinnamomi, ii.	171
Veratrum sabadilla, ii.	713	Vinegar of marigold, i.	365	Vinum cochleariæ compositum, i.	483
Verbasci albi herba, ii.	716	Vinegar of myrrh, i.	763	Vinum colchici, i.	538
Verbasculi pratensis radix, ii.	465	Vinegar of wild narcissus, ii.	259	Vinum colocynthidis, i.	575
Verbascum nigrum, ii.	716	Vinegar of opium, ii.	275	Vinum corticis Peruviani, i.	496
Verbascum phlomoides, ii.	716	Vinegar of pimenta, ii.	361	Vinum crocato-opiatum, ii.	277
Verbascum thapsus, ii.	716	Vinegar of red poppy, ii.	336	Vinum crocatum, i.	564
Verberna officinalis, ii.	717	Vinegar radical, i.	40, 41	Vinum fructuum cupressi compositum, i.	580
Verberna triphylla, ii.	717	Vinegar raspberry, ii.	533	Vinum dictamni martiatum, i.	673
Verbesinæ herba, i.	339	Vinegar of rosemary, ii.	523	Vinum emeticum, i.	258, 129
Verdigris, i.	590	Vinegar of roses, ii.	516	Vinum enulæ, ii.	118
Vermis majalis, ii.	232	Vinegar compound of roses, ii.	517	Vinum enulatum, ii.	118
Vernix sicca, ii.	555	Vinegar of rue, ii.	537	Vinum ferratum, i.	671
Veronica anagallis, ii.	719	Vinegar camphorated of rue, ii.	537	Vinum ferri, i.	671
Veronica beccabunga, ii.	719	Vinegar of sage, ii.	546	Vinum genistæ Sydenhami, ii.	453
Veronica officinalis, ii.	717	Vinegar saturnine, ii.	376	Vinum gentianæ, i.	713
Veronica teucrium, ii.	178	Vinegar of squill, ii.	573	Vinum gentianæ compositum, i.	713
Veronica virginica, ii.	178	Vinegar tarragon, i.	291	Vinum Hippocraticum, ii.	171
Verrucariæ herba, i.	365, ii. 2	Vinegar theriacal, ii.	118	Vinum ipecacuanhæ, ii.	128
Verrucarii herba, i.	560	Vinegar thieves', i.	115		
Vertebræ mustelæ piscis, i.	360	Vinegar of tobacco, ii.	263		
Vervain, ii.	717	Vinegar of turpentine, i.	40		
Vervain sweet-scented, ii.	717	Vinegar of wine, i.	38		
Vesicatorium magistræ, i.	383	Vinegar of wood, i.	39		
Vicia ervilla, ii.	719	Vini oleum, i.	100		
Vicia sativa, ii.	719	Vini sulphuretum, i.	161		
Victoralis femineæ radix, i.	720	Vinum absinthii, i.	27		
Victoralis longæ radix, i.	116	Vinum adstringens, ii.	531		
Victoralis rotundæ radix, i.	720	Vinum adustum, i.	104		
Vin opiacé préparé par la fermentation, ii.	282				

Vinum juniperinum, ii.	141	Viride montanum, i.	582	Unguentum aceti, i.	45
Vinum Lusitanicum ar- tefactum, i.	110	Virgin's bower, i.	523	Unguentum acidi nitrici, i.	70
Vinum martiatum, i.	671	Viscaginis herba, ii.	587	Unguentum acidi nitrosi, i.	70
Vinum martis, i.	671	Viscum album, ii.	724	Unguentum Ægyptiacum, i.	591, 592, ii. 660
Vinum nicotianæ, ii.	263	Viscum quercinum, ii.	724	Unguentum æruginis, i.	591, 593
Vinum opii, ii.	277	Vitis idææ baccæ, ii.	703	Unguentum adipoceræ cetorum, i.	441
Vinum opii aromaticum, ii.	277	Vitri herba, ii.	543	Unguentum Agrippæ, ii.	576
Vinum opii compositum, ii.	277	Vitriol blue, i.	583	Unguentum album, i.	435, 441, 373
Vinum opii crocatum, ii.	277	Vitriol green, i.	651	Unguentum album cam- phoratum, ii.	373
Vinum opii paregoricum, ii.	277	Vitriol of potass, ii.	428	Unguentum album resol- vens, i.	177
Vinum oxydi antimonii, i.	258	Vitrioli naphtha, i.	97	Unguentum album sim- plex, ii.	373
Vinum psychotriæ, ii.	128	Vitrioli oleum, i.	75	Unguentum aloes cum pe- troleo, i.	122
Vinum psychotriæ eme- ticæ, ii.	128	Vitrioli ros, i.	75	Unguentum althææ, i.	151
Vinum raphani martia- tum, i.	673	Vitroli spiritus, i.	75	Unguentum amarum, i.	576
Vinum rhei, ii.	500	Vitrioli spiritus phlogis- ticatus, i.	79	Unguentum ad ambusta, ii.	104
Vinum rhei amarum, ii.	590	Vitriolicum magnesæ, ii.	210	Unguentum amygdali- num, i.	435
Vinum rhei compositum, ii.	500	Vitriolum anglicum, i.	651	Unguentum amyridis, ii.	136
Vinum rhei palmati aro- maticum, ii.	500	Vitrum antimonii, i.	235	Unguentum anodynum, ii.	104
Vinum roborans, i.	28, 497, 498	Vitriolum cæruleum, i.	583	Unguentum anthelminti- cum, i.	31, 122, 576
Vinum salviæ, ii.	546	Vitriolum camphoratum, i.	585	Unguentum antipsoricum, ii.	639
Vinum scillæ, ii.	569	Vitriolum cupri, i.	583	Unguentum apostolorum, ii.	395
Vinum scilliticum, ii.	569	Vitriolum cyprinum, i.	583	Unguentum aquæ rosæ, ii.	518
Vinum sennæ, i.	409	Vitriolum cyprium, i.	583	Unguentum arcæi, ii.	658
Vinum sinapeos, ii.	590	Vitriolum ferreum, i.	651	Unguentum aromaticum, ii.	179, 524, 527
Vinum stibiatum cum ex- tractis narcoticis, i.	85, 260	Vitriolum ferri, i.	651	Unguentum de arthani- ta, i.	576
Vinum stramonii, i.	606	Vitriolum hungaricum, i.	651	Unguentum aureum, ii.	677
Vinum stypticum, i.	580	Vitriolum Londinense vi- ride cubicum, i.	651	Unguentum balsami et olei laricis, ii.	657
Vinum tabaci, ii.	263	Vitriolum martis, i.	651	Unguentum balsamicum, ii.	658
Vinum tartari emetici, i.	258	Vitriolum martis calcina- tum, i.	652	Unguentum basilicum, ii.	677
Vinum ex tartrate anti- monii, i.	258	Vitriolum martis ad albe- dinem calcinatum, i.	652	Unguentum basilicum ni- grum, ii.	678
Vinum tartratis antimo- nii, i.	258	Vitriolum martis siccatum, i.	652	Unguentum ad blenorr- hagiam cordatum, ii.	25
Vinum tonico-stimulans, i.	498	Vitriolum romanum, i.	651	Unguentum de bolo, i.	156
Vinum veratri albi, ii.	715	Vitriolum veneris, i.	583	Unguentum bryoniæ, ii.	576
Vinum vincetoxici ama- rum, i.	627	Vitriolum viride, i.	651	Unguentum Bardenii, ii.	426
Vinum viperarum, i.	699	Vitrum antimonii cera- tum, i.	235	Ungulæ caballinæ radix, ii.	699
Viola canina, ii.	721	Vitrum stibii, i.	235	Ungulæ cancrorum, i.	354
Viola odorata, ii.	721	Volkameria aculeata, i.	470	Unguentum cantharidis, i.	377
Viola pedata, ii.	721	Vulnerary Swiss, i.	279	Unguentum cantharidis cereum, i.	382
Viola tricolor, ii.	723	Vulvaræ herba, i.	458	Unguentum cantharidum colatum, i.	387
Violæ dentariæ radix, i.	609	Ulmariæ radix, ii.	621		
Violæ tricoloris herba, ii.	723	Ulmus campestris, ii.	700		
Violaræ radix, ii.	721	Ulmus fulva, ii.	700		
Violet, ii.	721	Ulma lactuca, ii.	2		
Viperinæ radix, i.	277, 620	Ulva pavonia, ii.	2		
Viper's bugloss, i.	620	Ulva squammaria, ii.	2		
Virgæ aureæ herba, i.	618	Umbel, i.	644		
Virideæ ris preparatum, i.	591	Umbilici folia, i.	562		
		Umbra, i.	644		
		Unguentum acetatis cu- pri, i.	593		
		Unguentum acetatis cu- pri cummelle, i.	591		
		Unguentum acetatis plum- bi, ii.	380		
		Unguentum acetatis plum- bi carbonatum, ii.	324		

Unguentum carbonatis plumbici, ii.	373	Unguentum gallarum, i.	692	Unguentum martiatum, ii.	182
Unguentum ceræ albæ, i.	435	Unguentum ex gallis opiatum, i.	692	Unguentum matris, ii.	324
Unguentum ceratum, i.	435	Unguentum gemmarum populi compositum, ii.	106	Unguentum medullæ mulagineum, i.	436
Unguentum cereum, i.	435	Unguentum hæmorrhoidale, i.	566	Unguentum mercuriale, ii.	19
Unguentum ceræ flavæ, i.	435	Unguentum ad hæmorrhoides, ii.	391	Unguentum mercuriale album, ii.	92
Unguentum cerussæ, ii.	373	Unguentum helenii, ii.	115	Unguentum mercuriale aptalizomenon, ii.	72
Unguentum ceti, i.	441	Unguentum hydrargyri, ii.	19	Unguentum mercuriale camphoratum, ii.	25
Unguentum cetacei album, i.	441	Unguentum hydrargyri album, ii.	92	Unguentum mercuriale citrinum, ii.	87
Unguentum citricum, i.	53	Unguentum hydrargyri citrinum, ii.	87	Unguentum mercuriale corrosivum, ii.	72
Unguentum citrinum, i.	151, 87, 677	Unguentum hydrargyri compositum, ii.	87	Unguentum mercurii rubrum, ii.	38
Unguentum comitissæ, i.	692	Unguentum hydrargyri fortius, ii.	20	Unguentum mezerei, i.	604
Unguentum commune, i.	435	Unguentum hydargyri griseum, ii.	34	Unguentum minii, ii.	370
Unguentum conyzæ, i.	757	Unguentum hydrargyri mitius, ii.	20	Unguentum de minio, ii.	398
Unguentum cupratum, i.	587, 593	Unguentum hydrargyri nitrati, ii.	38, 87	Unguentum mundificans Poterii, ii.	263
Unguentum Grilli, ii.	72	Unguentum hydrargyri nitrico-oxydi, ii.	38	Unguentum mundificans Stahlî, ii.	25
Unguentum defensivum, i.	156	Unguentum hydrargyri rubri, ii.	38	Unguentum mundificans Zelleri, ii.	92
Unguentum defensivum cæruleum, i.	526	Unguentum ex hydrargyro oxydato rubro, ii.	38	Unguentum muriatis hydrargyri ammoniacalis, ii.	92
Unguentum depurans, ii.	324	Unguentum hydrargyri rubrum, ii.	40	Unguentum muriatis hydrargyri oxygenati, ii.	72
Unguentum deutoxidi hydrargyri nitrati, ii.	38	Unguentum hyoscyami, ii.	103	Unguentum myrrhæ, i.	761
Unguentum digestivum, ii.	659	Unguentum infusi cantharidis vesicatoriæ, i.	377	Unguentum myrrhatum, ii.	659
Unguentum digestivum aloeticum, ii.	659	Unguentum inulæ, ii.	115	Unguentum myrtinum compositum, i.	692
Unguentum digestivum flavum, ii.	658	Unguentum iodatum, ii.	121	Unguentum Neapolitanum, ii.	19
Unguentum diureticum resolvens, ii.	568	Unguentum irritans, i.	385	Unguentum nervinum, ii.	179, 524
Unguentum dodecapharmacum, ii.	395	Unguentum juglandicum cum felle tauri, i.	633	Unguentum nitratis hydrargyri, ii.	87
Unguentum elemi, ii.	486, 658	Unguentum lacteum, ii.	659	Unguentum nitricatum, i.	70
Unguentum elemi cum alumine, ii.	40	Unguentum laurino-camphoratum, ii.	181	Unguentum nutritum, ii.	380
Unguentum elemi compositum, i.	487	Unguentum laurinum, ii.	179	Unguentum oculare, ii.	386
Unguentum elemi et terebinthinæ, ii.	658	Unguentum lenitivum, i.	151, 657	Unguentum ophthalmicum Richteri, ii.	39
Unguentum emolliens, i.	113, 435, 677	Unguentum de liquiritia, ii.	374	Unguentum ophthalmicum rubrum, ii.	38
Unguentum emolliens et sedans, ii.	283	Unguentum lithargyri, ii.	380	Unguentum opiatum, ii.	297
Unguentum empiricum Flaki in tumoribus hæmorrhoidalibus, ii.	624	Unguentum lithargyri resinosum martiale, ii.	680	Unguentum opodeldoch, i.	179
Unguentum enulatum, ii.	115	Unguentum lupuli, ii.	11	Unguentum oxydi hydrargyri rubri, ii.	38
Unguentum epispasticum, i.	377, 384, 604	Unguentum ad maculas corneæ, i.	633	Unguentum oxydi cobaltii, i.	526
Unguentum fellis bovini compositum, i.	633	Unguentum magistrale, ii.	398	Unguentum oxydi cupri melleum, i.	591
Unguentum filii, i.	436	Unguentum malorum insanorum, i.	593	Unguentum oxydi plumbi acetatum, ii.	380
Unguentum filii Zachariæ, i.	436, 151, 675	Unguentum marjoranæ, ii.	323	Unguentum oxydi plumbi carbonati, ii.	373
Unguentum ad fonticulos Bergii, i.	387				
Unguentum fuscum, ii.	391, 679				
Unguentum gallæ, i.	692				

Unguentum oxydi plum- bi fuscum, ii. 391	Unguentum rosmarini compositum, ii. 179	Unguentum sulphuris, ii. 638
Unguentum oxydi plum- bi rubrum, ii. 398	Unguentum rubefaciens, i. 604	Unguentum sulphuris compositum, ii. 639
Unguentum oxyduli plum- bi semivitrei adustum, ii. 391	Unguentum rubrum ad labia oris, i. 216	Unguentum supernitra- tis hydrargyri, ii. 87
Unguentum oxygena- tum extempore paran- dum, i. 460	Unguentum rubrum po- tabile, i. 216	Unguentum suppurans, i. 113
Unguentum oxynome- non, i. 70	Unguentum rutæ, ii. 538	Unguentum suppurati- vum, ii. 40
Unguentum paralyticum, ii. 463	Unguentum sabadillæ cum mercurio, ii. 22	Unguentum ad tabida membra, i. 757
Unguentum pectorale, ii. 133	Unguentum sabinæ, ii. 143	Unguentum tartratis po- tassæ stibiati, i. 261
Unguentum pediculorum, ii. 22	Unguentum salicis, ii. 542	Unguentum terebinthi- næ, ii. 659
Unguentum contra pedi- culos-capitis, i. 609	Unguentum sambuci, ii. 554	Unguentum terebinthi- natum ii. 657
Unguentum piceum regi- um, ii. 677	Unguentum sambuci- num, ii. 554	Unguentum terebinthi- næ resinosum, ii. 677
Unguentum picis, ii. 677	Unguentum santalinum, ii. 146	Unguentum tetraphar- macum, ii. 677
Unguentum picis empy- reumaticum, ii. 677	Unguentum saponaceum, ii. 462	Unguentum tripharma- cum, ii. 380
Unguentum picis liqui- dæ, ii. 365	Unguentum saturni sa- ponaceum, ii. 382	Unguentum turpethi mi- neralis, ii. 83
Unguentum piperis nigri, ii. 363	Unguentum saturninum, ii. 380	Unguentum ad varices ani, i. 492, ii. 381, 391
Unguentum pomadinum, ii. 518	Unguentum ad scabiem, ii. 34, 87, 638	Unguentum veratri, ii. 715
Unguentum populeum, ii. 106, 410	Unguentum ad scabiem Zilleri, ii. 92	Unguentum ad vermes, i. 31
Unguentum propolicum, ii. 467	Unguentum contra sca- biem, ii. 639	Unguentum contra ver- mes, i. 122, 576
Unguentum pulveris can- tharidis vesicatoriæ, i. 385	Unguentum scillæ, ii. 567	Unguentum ad vesicata, i. 385
Unguentum purpureum Wurzii, i. 592	Unguentum sebi arcæi, ii. 658	Unguentum vesicatori- um, i. 385
Unguentum radices phy- tolacæ, ii. 351	Unguentum sedativo- narcoticum, ii. 296	Unguentum viride, i. 593
Unguentum refrigerans, ii. 518	Unguentum siccativum, ii. 78	Unguentum viride pota- bile, ii. 110
Unguentum resinæ albæ, ii. 675	Unguentum simplex, i. 435	Unguentum ad ulcera lymphatica, ii. 40
Unguentum resinæ ele- mi, ii. 658	Unguentum sinapis, ii. 588	Unguentum ad ulcera venerea, ii. 40
Unguentum resinæ flavæ, ii. 658, 679	Unguentum solani, ii. 618	Unguentum de uvis, i. 436
Unguentum resinæ ni- græ, ii. 678	Unguentum spermatis ceti, i. 441	Unicorn, i. 359
Unguentum resinæ pini sylvestris compositum, i. 151	Unguentum staphisa- græ, ii. 22	Unicornu, i. 359, 569
Unguentum resinosum, ii. 675	Unguentum stramonii, i. 607	Unicornu fossile, i. 360
Unguentum resolvens, ii. 25	Unguentum e styrace compositum, ii. 627	Unicornu marinum, i. 359
Unguentum resumpti- vum, i. 151	Unguentum styracis re- sinosum, ii. 627	Unicornu verum, i. 359
Unguentum rorismari- ni compositum, ii. 527	Unguentum subacetatis, cupri cum melle, i. 591	Uniones, i. 354
Unguentum rosarum, ii. 518	Unguentum subacetatis plumbici, ii. 380	Ursini veri herba, i. 34
Unguentum rosatum, ii. 518	Unguentum subcarbona- tis plumbi, ii. 373	Urtica dioica, ii. 701
Unguentum rosatum compositum, ii. 92	Unguentum subnitrat- is hydrargyrici, ii. 38	Urtica pilulifera, ii. 701
	Unguentum sulphatis hy- drargyri, ii. 83	Urtica urens, ii. 701
	Unguentum sulphura- tum, ii. 639	Urticæ hispanicæ semen, ii. 701
	Unguentum sulphura- tum simplex, ii. 638	Urticæ mortuæ herba, ii. 154
	Unguentum e sulphure hydrargyrato, ii. 30	Urticæ romanæ semen, ii. 701
		Uruçu, ii. 352
		Usnea plicata, ii. 187
		Uvæ versæ, ii. 342
		Uvæ ursi folia, i. 271
		Uvæ vulpinæ, ii. 342
		Uvulariæ herba, ii. 536

Wakaka Indian, ii.	712	Water of cichory, i.	462	Water of mentha crispa, ii.	238
Wake-robin, i.	291	Water buglossated cinna- mon, ii.	1	Water alkaline mephitic, ii.	427
Wallflower, i.	455	Water cordial cinnamon, ii.	167	Water simple mercurial, ii.	13
Wall pepper, ii.	580	Water hordeated cinna- mon, ii.	167	Water of mugwort, i.	289
Wallnut tree, ii.	134	Water simple cinnamon, ii.	166	Water of mustard, ii.	589
Wash black, ii.	34	Water vinous cinnamon, ii.	167	Water of myrrh, i.	764
Water camphorated æthe- reous, i.	371	Water of cloves, i.	400	Water of Naples, i.	67
Water of Aix la Chapelle, i.	67	Water of coltsfoot, ii.	699	Water of garden night- shade, ii.	618
Water acetic alexipharmic, i.	223	Water comforting, ii.	167	Water of opium, ii.	275
Water simple alexiphar- mic, i.	222	Water of convallaria, i.	542	Water of orange leaves, i.	508
Water spirituous alexi- pharmic, i.	223	Water cold cordial, ii.	167	Water of orange flowers, i.	508
Water gaseous alkaline, ii.	608	Water of coriander, i.	559	Water of orange peel, i.	508
Water of bitter almonds, i.	63	Water cosmetic, i.	64	Water oxygenated, ii.	328
Water of anemone, i.	218	Water of cowslip, ii.	466	Water of parsley, i.	269
Water of angelica, i.	222	Water parsnip creeping, ii.	591	Water of wall pellitory, ii.	341
Water of anise, ii.	355	Water of dill, i.	221	Water of pennyroyal mint, ii.	239
Water antiscorbutic, i.	534	Water diuretic, i. 64, ii.	482	Water of peony, ii.	329
Water compound of assa- foetida, i.	305	Water of elderberries, ii.	554	Water of black pepper, ii.	361
Water Balaruc, ii.	604	Water of dwarf elder, ii.	551	Water of peppermint, ii.	238
Water of balm, ii.	229	Water of elder flowers, ii.	554	Water of plantain, ii.	367
Water of Barèges, i.	66	Water of elecampane, ii.	116	Water of red poppy, ii.	334
Water barley, ii.	7	Water eryngo, i.	622	Water of potass, ii.	410
Water of beccabunga, ii.	719	Water of eye-bright, i.	630	Water of privet, ii.	198
Water artificial of birch, i.	110	Water of fennel, i. 219, ii.	219	Water of purslain, ii.	410
Water of blessed thistle, i.	525	Water of fetid goose-foot, i.	458	Water Pymont, ii.	604
Water of blue-bottle, i.	597	Water ferruginous, i.	647	Water Pymont factitious, ii.	209
Water of Bonnes, i.	67	Water saline ferruginous, i.	651	Water of quince, i.	599
Water of borage, i.	341	Water of fumitory, i.	681	Water Rabel's, i.	79
Water Bourbonne-les- bains, ii.	604	Water compound of harts- horn, i.	188	Water of raspberries, ii.	533
Water spirituous of bread, ii.	334	Water of hemlock, i. 466, ii.	466	Water of rhodium wood, ii.	195
Water of bugloss, i.	216	Water honey, i.	559	Water of rose, ii.	517
Wine camphorated, i.	368, 373	Water of hops, ii.	11	Water of rosemary, ii.	527
Water caraway, i.	398	Water of horehound, ii.	223	Water of rue, ii.	538
Water carminative, i.	509	Water of horseradish, ii.	482	Water of sage, ii.	546
Water simple carmina- tive, i.	449	Water Hungary, ii.	528	Water compound of sage, ii.	547
Water spirituous carmi- tive, i.	449	Water of hyssop, ii.	112	Water of sassafras, ii.	184
Water of cascarrilla, i.	567	Water juniper, ii.	139	Water compound of sas- safras, ii.	184
Water spirituous of castor, i.	426	Water of lavender, ii.	157	Water of savine, ii.	144
Water of celandine, i.	456	Water distilled of lemon, i.	518	Water of scabious, ii.	559
Water of lesser centaury, i.	625	Water of garden lettuce, ii.	153	Water of scordium, ii.	683
Water chalybeate, i.	650	Water of lily, ii.	198	Water of scurvy-grass, i.	534
Water of chamomile, i.	448	Water of water lily, ii.	268	Water artificial sea, ii.	603
Water anisated chamo- mille, i.	450	Water of lime, i.	347	Water artificial Seidlitz, ii.	212
Water vinous of chamo- mille, i.	448	Water compound of lime, ii.	183	Water Selter, ii.	604
Water of black cherries, i.	62	Water of lime flowers, ii.	692	Water Seltz, ii.	604
Water of bird cherry, i.	63	Water of lime with wax, i.	467	Water of sloe flowers, ii.	468
Water of cherry laurel, i.	62	Water of lohoch sanum et expertum, ii.	167	Water soda, ii.	608
Water of chervil, i.	445	Water of magnesia, ii.	209	Water of sorrel, ii.	534
Water of chlorine, i.	459	Water of mallow, ii.	216	Water of wood sorrel, ii.	327
		Water of marjoram, ii.	322	Water of spearmint, ii.	239
		Water melon, i.	578	Water of spruce fir, i.	21

Water of strawberries, i.	677	Wine antileucorrhœic, i.	499	Wine compound of orange peel, i.	512
Water sulphureous, i.	66	Wine antimonial, i.	258	Wine artificial port, i.	110
Water compound sulphureous, ii.	553	Wine antiperiodic, i.	499	Wine purgative and tonic, i.	409
Water of swallows, i.	188	Wine antirheumatic, i.	260	Wine of quince, i.	110
Water of sweet flag, i.	87	Wine antiscorbutic, i.	499, 483	Wine of rhubarb, ii.	500
Water of tansey, ii.	651	Wine bitter, i.	627	Wine bitter of rhubarb, ii.	500
Water tar, ii.	365	Wine bitter and diuretic, i.	409	Wine of saffron, i.	564
Water compound theriacal, ii.	314	Wine aromatic chalybeate, ii.	672	Wine of sage, ii.	548
Water Roman theriacal, ii.	314	Wine bitter chalybeate, i.	672	Wine of senna, i.	409
Water simple theriacal, ii.	314	Wine of cinchona, i.	496	Wine spirit of, i.	104
Water of thyme, ii.	690	Wine aromatic of cinchona, i.	498	Wine sulphate of quinine, ii.	479
Water of wild thyme, ii.	690	Wine compound of cinchona, ii.	498	Wine of squill, ii.	569
Water of tormentil, ii.	693	Wine ferruginous of cinchona, i.	497	Wine alkaline of squill, ii.	571
Water of vanilla, ii.	712	Wine of cinchonine, i.	504	Wine stimulant, i.	378
Water of valerian, ii.	711	Wine of colchicum, i.	538	Wine stomachic, i.	499, 525
Water Bell's vegeto-mineral, ii.	388	Wine of colocynth, i.	575	Wine of sweetflag, i.	88
Water of veronica, ii.	718	Wine compound of cypress, i.	580	Wine of tartarized antimony, i.	258
Water violet, ii.	133, 721	Wine of digitalis, i.	616	Wine of tobacco, ii.	263
Water vulnerary, i.	79	Wine diuretic, ii.	141, 273, 454, 484, 665	Wine Barthez' tonic and aperitive, i.	389
Water compound of wormwood, i.	25	Wine bitter diuretic, ii.	570	Wine of valerian, ii.	708
Water-avens, i.	719	Wine Sydenham's diuretic, ii.	453	Wine of vipers, i.	699
Water parsnip narrow-leaved, ii.	591	Wine of elecampane, ii.	118	Wine of wormwood, i.	27
Wax virgin, i.	433	Wine excitant, ii.	453	Wintera aromatica, i.	620
Wax white, i.	433	Wine chalybeate of fraxinella, i.	673	Winterberry deciduous, ii.	467
Wax yellow, i.	433	Wine of gentian, i.	713	Winter-cherry, ii.	350
Whey, ii.	585	Wine compound of gentian, i.	713	Winter-cress, i.	623
Whey acetose, ii.	586	Wine of white hellebore, ii.	715	Wintergreen, ii.	471
Whey acidulous, ii.	585	Wine hippocratic, ii.	171	Wintergreen American, i.	458
Whey antiscorbutic, i.	533, 647	Wine of horseradish, ii.	483	Wismuthum, i.	339
Whey aperient, ii.	213	Wine chalybeate of horseradish, i.	673	Woad, ii.	133
Whey chalybeate, ii.	585	Wine compound of horseradish, ii.	483	Wood aloe, ii.	192
Whey compound, ii.	586	Wine hydragogue, ii.	453	Wood Brazil, ii.	193
Whey Hagenot's, i.	722	Wine of ipecacuanha, ii.	128	Wood Campechy, ii.	103
Whey Hoffmann's, ii.	585	Wine compound of ipecacuanha, ii.	129	Wood ebony, ii.	194
Whey mustard, ii.	590	Wine of iron, i.	671	Wood nephritic, ii.	195
Whey purgative, ii.	220	Wine compound of iron, i.	673	Wood Rhodium, ii.	195
Whey sweet, ii.	585	Wine alkalized of juniper, ii.	453	Wood sanders, ii.	195
Whey tamarind, ii.	647	Wine of mustard, ii.	590	Wood-louse, i.	272
Whey Van Swieten's, ii.	456	Wine ophthalmic, i.	122	Woolly thistle, i.	433
Whey vinous, ii.	585	Wine opiate, ii.	276	Worm bark, i.	716
Whey Weiss's, i.	412	Wine astringent opiate, ii.	319	Worm earth, i.	698
Whiskey, i.	104	Wine of orange peel, i.	511	Wormwood, i.	23
White dittany, i.	611	Wine alkaline of orange peel, i.	513	Wormwood alpine, i.	291
White walnut-tree, ii.	134			Wormwood beer, i.	27
Willow, ii.	542			Wormwood Chinese, i.	289
Willow-herb, ii.	205			Wormwood common, i.	23
Wine alkaline of aloes, i.	133			Wormwood creeping, i.	291
Wine of aloes, i.	119			Wormwood Roman, i.	24
Wine anti-apoplectic, i.	551			Wormwood sea, i.	24
Wine anti-icteric, i.	409, 512			Wrightea antidysenterica, i.	470
Wine anti-arthritic, i.	85				
Wine aromatic, ii.	546				
Wine bitter aromatic, i.	28				
Wine camphorated aromatic, i.	373				
Wine astringent, i.	555, 531				
ii.					

INDEX.

863

Yarrow, i.	34	Zarsaparilla, ii.	593	Zingiber vulgare, i.	203
Yeast, i.	634	Zerumbet, i.	203	Zingiberis radix, i.	203
Yelk of wool, ii.	269	Zest, i.	517	Zostera gorgonia, ii.	2
Yellow root, ii.	98	Zincum silesiacum, ii.	145	Zostera marina, ii.	2
		Zingiber album, i.	203	Zostera sertularia, ii.	2
		Zingiber commune, i.	205	Zostera tubularia, ii.	2
Zarsa, ii.	593	Zingiber nigrum, i.	203		

END OF THE SECOND VOLUME.

